Soil Monitoring System

Throughput:

3" Deep .25' max depth of materi
24" Wide 2.0' (active belt width)
1.5 ft per sec (0.45 m/sec) max travel speed for rec. sensitivity (~1.5 pc:/s

.25' x 2.0' x 1.5 ft/sec x 60 sec/min x 60 min/hr
= 2700 CF/hr
= 100 CY/hr MAX

Correction factors allocated:

1. .8 3" is max depth at bottom of concave conveyor belt only. Depth diminishes in both directions toward the sides.

2. .8 Throughput interruptions attributed to positive alarms and subsequent materials marking, sorting, and restarts.

3. .8 System start-up/shut-down operati QC - achieve 8 hrs productive effort over 10 hr work day

4. .5 Weather related delays, system obstruction, equipment failure.

100 cy/hr x 0.8 x 0.8 x 0.8 x 0.5 = ~25 cy

MicroShield v5.01 (5.01-01003)
Allied Technology Group, Inc.
Results With Buildup
FILE: C:\JOEL\MS5\DATA\CO2VEYOR.MS5
Case Title: Cesium point source
This case was run on Monday, November 23, 1998 at 12:51:49 PM
Dose Point # 3 - (6,6.57e+00,0) in

Group #	Energy (MeV)	Activity photons/sec	Fluence Rate photons/cm²/sec	Energy Fluence MeV/cm²/sec	Exposure Rate mR/hr
1	0.0318	3.623e+003	2.385e-008	7.588e-010	6.321e-012
2	0.0322	6.685e+003	8.197e-008	2.639e-009	2.124e-011
3	0.0364	2.433e+003	6.391e-006	2.326e-007	1.322e-009
4	0.6616	1.575e+005	1.698e+001	1.123e+001	2.178e-002
	TOTALS:	1.702e+005	1.698e+001	1.123e+001	2.178e-002

MicroShield v5.01 (5.01-01003)
Allied Technology Group, Inc.
Results With Buildup
FILE: C:\JOEL\MS5\DATA\CO2VEYOR.MS5
Case Title: Cesium point source
This case was run on Monday, November 23, 1998 at 12:51:49 PM
Dose Point # 4 - (6,0,0) in

Group #	Energy (MeV)	Activity photons/sec	Fluence Rate photons/cm²/sec	Energy Fluence MeV/cm²/sec	Exposure Rate mR/hr
1	0.0318	3.623e+003	2.497e-005	7.945e-007	6.618e-009
2	0.0322	6.685e+003	7.046e-005	2.268e-006	1.826e-008
3	0.0364	2.433e+003	1.003e-003	3.652e-005	2.075e-007
4	0.6616	1.575e+005	4.436e+001	2.935e+001	5.690e-002
	TOTALS:	1.702e+005	4.436e+001	2.935e+001	5.690e-002

MicroShield v5.01 (5.01-01003)
Allied Technology Group, Inc.
Results With Buildup
FILE: C:\JOEL\MS5\DATA\CO2VEYOR.MS5
Case Title: Cesium point source
This case was run on Monday, November 23, 1998 at 12:51:49 PM
Dose Point # 1 - (6.5,1.97e+01,0) in

Group #	Energy (MeV)	Activity photons/sec	Fluence Rate photons/cm²/sec	Energy Fluence MeV/cm²/sec	Exposure Rate mR/hr
1	0.0318	3.623e+003	1.092e-018	3.475e-020	2.895e-022
2	0.0322	6.685e+003	7.388e-018	2.378e-019	1.914e-021
3	0.0364	2.433e+003	1.979e-013	7.204e-015	4.093e-017
4	0.6616	1.575e+005	1.240e+000	8.203e-001	1.590e-003
	TOTALS:	1.702e+005	1.240e+000	8.203e-001	1.590e-003

MicroShield v5.01 (5.01-01003)
Allied Technology Group, Inc.
Results With Buildup
FILE: C:\JOEL\MS5\DATA\CO2VEYOR.MS5
Case Title: Cesium point source
This case was run on Monday, November 23, 1998 at 12:51:49 PM
Dose Point # 2 - (6,1.30e+01,0) in

oup #	Energy (MeV)	Activity photons/sec	Fluence Rate photons/cm²/sec	Energy Fluence MeV/cm²/sec	Exposure Rate mR/hr
1	0.0318	3.623e+003	8.047e-014	2.560e-015	2.133e-017
2	0.0322	6.685e+003	3.958e-013	1.274e-014	1.026e-016
3	0.0364	2.433e+003	6.913e-010	2.516e-011	1.430e-013
4	0.6616	1.575e+005	4.232e+000	2.800e+000	5.428e-003
	TOTALS:	1.702e+005	4.232e+000	2.800e+000	5.428e-003

CASE 4

MICROSHIELD OUTPUT

5 uCi, Cs-137 point source

MicroShield v5.01 (5.01-01003)
Allied Technology Group, Inc.
Results With Buildup
FILE: C:\JOEL\MS5\DATA\CO2VEYOR.MS5
Case Title: Cobalt point source
This case was run on Monday, November 23, 1998 at 11:41:43 AM
Dose Point # 3 - (6,6.57e+00,0) in

Group #	Energy (MeV)	Activity photons/sec	Fluence Rate photons/cm²/sec	Energy Fluence MeV/cm²/sec	Exposure Rate mR/hr
1	0.6938	3.018e+001	3.240e-003	2.248e-003	4.340e-006
2	1.1732	1.850e+005	1.971e+001	2.312e+001	4.131e-002
3	1.3325	1.850e+005	1.976e+001	2.633e+001	4.567e-002
	TOTALS:	3.700e+005	3.947e+001	4.945e+001	8.699e-002

MicroShield v5.01 (5.01-01003)
Allied Technology Group, Inc.
Results With Buildup
FILE: C:\JOEL\MS5\DATA\CO2VEYOR.MS5
Case Title: Cobalt point source
This case was run on Monday, November 23, 1998 at 11:41:43 AM
Dose Point # 4 - (6,0,0) in

Group #	Energy (MeV)	Activity photons/sec	Fluence Rate photons/cm²/sec	Energy Fluence MeV/cm²/sec	Exposure Rate mR/hr
1	0.6938	3.018e+001	8.446e-003	5.860e-003	1.131e-005
2	1.1732	1.850e+005	5.041e+001	5.914e+001	1.057e-001
3	1.3325	1.850e+005	5.034e+001	6.707e+001	1.164e-001
	TOTALS:	3.700e+005	1.008e+002	1.262e+002	2.221e-001

MicroShield v5.01 (5.01-01003)
Allied Technology Group, Inc.
Results With Buildup
FILE: C:\JOEL\MS5\DATA\CO2VEYOR.MS5
Case Title: Cobalt point source
This case was run on Monday, November 23, 1998 at 11:41:43 AM
Dose Point # 1 - (6.5,1.97e+01,0) in

Group #	Energy (MeV)	Activity photons/sec	Fluence Rate photons/cm²/sec	Energy Fluence MeV/cm²/sec	Exposure Rat mR/hr
1	0.6938	3.018e+001	2.408e-004	1.671e-004	3.226e-007
2	1.1732	1.850e+005	1.734e+000	2.035e+000	3.636e-003
3	1.3325	1.850e+005	1.809e+000	2.411e+000	4.183e-003
	TOTALS:	3.700e+005	3.544e+000	4.446e+000	7.819e-003

MicroShield v5.01 (5.01-01003)
Allied Technology Group, Inc.
Results With Buildup
FILE: C:\JOEL\MS5\DATA\CO2VEYOR.MS5
Case Title: Cobalt point source
This case was run on Monday, November 23, 1998 at 11:41:43 AM
Dose Point # 2 - (6,1.30e+01,0) in

Group #	Energy (MeV)	Activity photons/sec	Fluence Rate photons/cm²/sec	Energy Fluence MeV/cm²/sec	Exposure Rate mR/hr
1	0.6938	3.018e+001	8.136e-004	5.645e-004	1.090e-006
2	1.1732	1.850e+005	5.315e+000	6.236e+000	1.114e-002
3	1.3325	1.850e+005	5.430e+000	7.235e+000	1.255e-002
	TOTALS:	3.700e+005	1.075e+001	1.347e+001	2.370e-002

CASE 3

MICROSHIELD OUTPUT

5 uCi, Co-60 point source

MicroShield v5.01 (5.01-01003)
Allied Technology Group, Inc.
Results With Buildup
FILE: C:\JOEL\MS5\DATA\CO_VEYOR.MS5
Case Title: Conveyor - Cesium
This case was run on Monday, November 23, 1998 at 11:31:07 AM
Dose Point # 3 - (6,19.5,6) in

Group #	Energy (MeV)	Activity photons/sec	Fluence Rate photons/cm²/sec	Energy Fluence MeV/cm²/sec	Exposure Rate mR/hr
1	0.0318	3.001e+002	1.714e-003	5.453e-005	4.542e-007
2	0.0322	5.537e+002	3.278e-003	1.055e-004	8.492e-007
3	0.0364	2.015e+002	1.724e-003	6.277e-005	3.566e-007
4	0.6616	1.304e+004	1.035e+000	6.846e-001	1.327e-003
	TOTALS:	1.410e+004	1.041e+000	6.848e-001	1.329e-003

MicroShield v5.01 (5.01-01003)
Allied Technology Group, Inc.
Results With Buildup
FILE: C:\JOEL\MS5\DATA\CO_VEYOR.MS5
Case Title: Conveyor - Cesium
This case was run on Monday, November 23, 1998 at 11:31:07 AM
Dose Point # 4 - (6,19.5,3) in

Group #	Energy (MeV)	Activity photons/sec	Fluence Rate photons/cm²/sec	Energy Fluence MeV/cm²/sec	Exposure Rate mR/hr
1	0.0318	3.001e+002	1.497e-003	4.763e-005	3.968e-007
2	0.0322	5.537e+002	2.863e-003	9.217e-005	7.418e-007
3	0.0364	2.015e+002	1.504e-003	5.475e-005	3.111e-007
4	0.6616	1.304e+004	8.913e-001	5.897e-001	1.143e-003
	TOTALS:	1.410e+004	8.972e-001	5.899e-001	1.145e-003

MicroShield v5.01 (5.01-01003)
Allied Technology Group, Inc.
Results With Buildup
FILE: C:\JOEL\MS5\DATA\CO_VEYOR.MS5
Case Title: Conveyor - Cesium
This case was run on Monday, November 23, 1998 at 11:31:07 AM
Dose Point # 1 - (6,19.5,12) in

Group #	Energy (MeV)	Activity photons/sec	Fluence Rate photons/cm²/sec	Energy Fluence MeV/cm²/sec	Exposure Rate mR/hr
1	0.0318	3.001e+002	1.816e-003	5.776e-005	4.812e-007
2	0.0322	5.537e+002	3.472e-003	1.118e-004	8.996e-007
3	0.0364	2.015e+002	1.828e-003	6.653e-005	3.780e-007
4	0.6616	1.304e+004	1.127e+000	7.458e-001	1.446e-003
	TOTALS:	1.410e+004	1.134e+000	7.461e-001	1.448e-003

MicroShield v5.01 (5.01-01003)
Allied Technology Group, Inc.
Results With Buildup
FILE: C:\JOEL\MS5\DATA\CO_VEYOR.MS5
Case Title: Conveyor - Cesium
This case was run on Monday, November 23, 1998 at 11:31:07 AM
Dose Point # 2 - (6,19.5,9) in

Group #	Energy (MeV)	Activity photons/sec	Fluence Rate photons/cm²/sec	Energy Fluence MeV/cm²/sec	Exposure Rate mR/hr
1	0.0318	3.001e+002	1.794e-003	5.709e-005	4.755e-007
2	0.0322	5.537e+002	3.432e-003	1.105e-004	8.891e-007
3	0.0364	2.015e+002	1.806e-003	6.574e-005	3.735e-007
4	0.6616	1.304e+004	1.106e+000	7.317e-001	1.418e-003
	TOTALS:	1.410e+004	1.113e+000	7.319e-001	1.420e-003

CASE 2

MICROSHIELD OUTPUT

Soil with 5 pCi Cs-137 per gram.

MicroShield v5.01 (5.01-01003)
Allied Technology Group, Inc.
Results With Buildup
FILE: Case 1
Case Title: Co-veyor
This case was run on Monday, November 23, 1998 at 11:19:41 AM
Dose Point # 3 - (6,19.5,6) in

Group #	Energy (MeV)	Activity photons/sec	Fluence Rate photons/cm²/sec	Energy Fluence MeV/cm²/sec	Exposure Rate mR/hr
1	0.6938	4.999e-001	3.964e-005	2.751e-005	5.311e-008
2	1.1732	3.065e+003	2.455e-001	2.880e-001	5.146e-004
3	1.3325	3.065e+003	2.464e-001	3.283e-001	5.696e-004
	TOTALS:	6.130e+003	4.919e-001	6.163e-001	1.084e-003

MicroShield v5.01 (5.01-01003)
Allied Technology Group, Inc.
Results With Buildup
FILE: Case 1
Case Title: Co-veyor
This case was run on Monday, November 23, 1998 at 11:19:41 AM
Dose Point # 4 - (6,19.5,3) in

Group #	Energy (MeV)	Activity photons/sec	Fluence Rate photons/cm²/sec	Energy Fluence MeV/cm²/sec	Exposure Rate mR/hr
1	0.6938	4.999e-001	3.416e-005	2.370e-005	4.576e-008
2	1.1732	3.065e+003	2.120e-001	2.487e-001	4.445e-004
3	1.3325	3.065e+003	2.130e-001	2.838e-001	4.923e-004
	TOTALS:	6.130e+003	4.250e-001	5.325e-001	9.369e-004

MicroShield v5.01 (5.01-01003)
Allied Technology Group, Inc.
Results With Buildup
FILE: Case 1
Case Title: Co-veyor - I
This case was run on Monday, November 23, 1998 at 11:19:41 AM
Dose Point # 1 - (6,19.5,12) in

Group #	Energy (MeV)	Activity photons/sec	Fluence Rate photons/cm²/sec	Energy Fluence MeV/cm²/sec	Exposure Rate mR/hr
1	0.6938	4.999e-001	4.319e-005	2.997e-005	5.785e-008
2	1.1732	3.065e+003	2.672e-001	3.135e-001	5.602e-004
3	1.3325	3.065e+003	2.681e-001	3.573e-001	6.199e-004
	TOTALS:	6.130e+003	5.354e-001	6.708e-001	1.180e-003

MicroShield v5.01 (5.01-01003)
Allied Technology Group, Inc.
Results With Buildup
FILE: Case 1
Case Title: Co-veyor
This case was run on Monday, November 23, 1998 at 11:19:41 AM
Dose Point # 2 - (6,19.5,9) in

Group #	Energy (MeV)	Activity photons/sec	Fluence Rate photons/cm²/sec	Energy Fluence MeV/cm²/sec	Exposure Rate mR/hr
1	0.6938	4.999e-001	4.237e-005	2.940e-005	5.676e-008
2	1.1732	3.065e+003	2.622e-001	3.076e-001	5.496e-004
3	1.3325	3.065e+003	2.631e-001	3.506e-001	6.082e-004
	TOTALS:	6.130e+003	5.253e-001	6.582e-001	1.158e-003

CASE 1

MICROSHIELD OUTPUT

Soil with 1 pCi Co-60 per gram.

Calculation MCAB-01

DETECTABILITY - MINIMUM DETECTABLE COUNT RATE (MDCR)

The scanning MDCR applies both to a detector moving along the ground, as well as soil on a conveyor belt moving under a detector. This MDCR is a function of background count rate, observation interval and confidence level (d'). As shown on the hand calculation sheet, MDCR is 676 cpm (net) for a background of 8,000 cpm and 2sec observation interval.

SUMMARY

Case	Calc'ed Count Rate (net)	MDCR (net)	Detectable Amount*
1	464 cpm	676	1.5 pCi/g
2	1188	676	2.8 pCi/g
3	24,983	676	135 nCi
4	14,760	676	229 nCi

* Concentration or source strength, ratioed to MDCR.

MDCR
(MARSSIM eqn. 6-9)

$MDCR = S_i \times 60/i$

$S_i = d' \sqrt{b_i}$

i = 2sec obs. interval

$d' = 1.38$ (per MARSSIM)

b_i = 8000 cpm × 2/60 = 267

$S_i = \sqrt{b_i} \times 1.38 = 22.5$

$MDCR = 22.5 \times 60/2 = 676$ cpm

CASE 2:

Same detector geometry.

Microshield results show dose rate fairly constant across width of belt. Average dose rate 1.32 uR/hr, from soil at 5 pCi-Cs-137/gram. Using a dose rate to count rate conversion of 900 cpm/uR/hr[2], gives 1188 cpm (net.)

Results are scalable to other concentrations of Cs-137.

CASE 3:

In this case, a point source of Co-60 travels towards and under the detectors. Width of the belt is not considered, because the particle will pass directly under one of the detectors. It is assumed to be under the 3" of soil. Since Ludlum ratemeters integrate for 2 seconds, the particle will travel 1m (39") in that time.

Seven particle positions are modeled, from 19.7" in front of the detectors to 19.7" after the detectors. The average dose rate of the seven is 58.1 uR/hr, as shown; yielding a count rate of 24,983 cpm.

Distance from Detectors	Dose Rate uR/hr
±19.7"	7.8
±13.0	23.7
±6.6	87.0
0	222.1
Average	58.1

CASE 4:

The geometry is the same as Case 3. The point source in this case is 5 uCi Cs-137.

The average dose rate is 16.4 uR/hr, as shown; yielding a count rate of 14,760 cpm.

Distance from Detectors	Dose Rate uR/hr
±19.7"	1.9
±13.0	5.4
±6.6	21.8
0	56.9
Average	16.4

[2] MARSSIM Table 6-7.

Calculation MCAB-01



CONVEYOR-MOUNTED DETECTOR MODEL

DIMENSIONS:

24" wide
belt moves at 0.5 m/s; covers 1m (39") in 2sec. counting interval
soil 3" high on belt
detectors mounted 6" above belt
10 2x2 NaI detectors cover width of belt

ASSUMPTIONS:

Soil is uniformly contaminated at 1 pCi Co-60/gram soil (Case 1.)
Soil is uniformly contaminated at 5 pCi Cs-137/gram soil (Case 2.)
Soil covers a 5 uCi point source of Co-60 (Case 3.)
Soil covers a 5 uCi point source of Cs-137 (Case 4.)

Summary of four Cases shown below, followed by discussion of detectability.

CASE 1:

Four detector positions modeled with Microshield Ver. 5. 1) center of belt (12" from edge); 2) 9" from edge; 3) 6" from edge; and 4) 3" from edge.

Mirror-image positions are the same for distances from the other side of the belt. Ignore soil on belt, which is further than 19.5" from detectors.

Microshield results show dose rate fairly constant across width of belt. Average dose rate 1.08 uR/hr, from soil at 1 pCi-Co-60/gram. Using a dose rate to count rate conversion of 430 cpm/uR/hr[1], gives 464 cpm (net.)

[1] MARSSIM Table 6-7.

ATG INC.

CALCULATION COVER SHEET

CALC. NO. MCAB-01
No. of Sheets: 1 of 76

PROJECT: Fort McClellon Burial Mound Decommissioning

PURPOSE: Calculate the performance of a conveyor-mounted detector system, scanning soil with potential cobalt-60 and cesium-137 contamination.

SOURCES OF DATA: Preliminary design of conveyor system - Lake City Project Manager, Frank Whitaker

SOURCES OF FORMULAE & REFERENCES: Multi-Agency Radiation Survey and Site Investigation Manual, MARSSIM (NUREG-1575) December, 1997.

ATTACHMENT: Summary of hand calculations and output from Microshield™ runs.

CONCLUSION: System minimum detectable count rate (MDCR) is approx. 676 cpm (net). This will allow detection of Co-60 at 1.5 pCi/g or 135 nCi discrete source; or Cs-137 at 2.8 pCi/g or 80 nCi discrete source.

PRELIMINARY CALC. ☐ FINAL CALC. ☐ SUPERSEDES CALC. NO. _____

REV. NO.	REVISION	BY:	DATE	CHECKED	DATE	APPROVED	DATE

APPENDIX 5

PRELIMINARY DESIGN OF CONVEYOR-MOUNTED DETECTION SYSTEM

APPENDIX 6

DEVELOPMENT OF DERIVED CLEANUP GUIDELINES FOR THE PELHAM RANGE 'BURIAL MOUND' FORT McCLELLAN

Development of Derived Cleanup Guidelines for the Pelham Range Burial Mound, Fort McClellan, Revision 1

June 29, 1999

Prepared for:
Allied Technology Group
669 Emory Valley Road,
Oak Ridge, TN 37830

Prepared by:
Auxier & Associates, Inc.
10317 Technology Drive, Suite 1
Knoxville, Tennessee 37932

TABLE OF CONTENTS

1.0 Introduction 1
1.1 Background 1
1.2 Objective 1
1.3 Organization of the Report 1
2.0 Description of Approach 2
3.0 Exposure Assessment 3
3.1 Characterization of the Site 3
3.1.1 Physical Setting 3
3.1.2 Site History 3
3.1.3 Land Use 4
3.1.4 Conceptual Model of the Site 4
3.1.5 Sources and Release Mechanisms 5
3.1.6 Potential Transport Pathways 5
3.1.7 Potential Exposure Routes 5
3.1.8 Receptor Scenarios Considered 8
3.1.9 Receptors Selected 10
3.2 Application of RESRAD 12
3.3 Summary of Receptor Exposures 13
3.3.1 Guardsman 13
3.3.2 Resident Adult with Garden 13
3.3.3 Resident Child with Garden 13
4.0 Derived Cleanup Guidelines 15
4.1 Role of Dose Based Criteria 15
4.2 Dose Based Cleanup Guidelines 15
4.3 Influence of Multiple Contaminants on Cleanup Guidelines 15
4.4 Derived Cleanup Guidelines Based on Expected Land Use 18
4.5 Impact of DCGLs and Soil Sorting on Final Status of Burial Mound Soil 18
4.6 Health Protectiveness of Expected Residual Soil Contamination Levels 18
5.0 References 20
Attachment A RESRAD Output 21

1.0 Introduction

1.1 Background

Fort McClellan is an army base sited on 45,679 acres adjacent to Anniston, AL. A part of the base, known as Rideout Field, served as a radiological training area from the mid 1950's through May of 1972. The radiological training area was decommissioned and some contaminated soil was excavated and moved to a burial mound located in the northwest corner of the Pelham Range. This mound, called the Pelham Burial Mound in this report, contains elevated concentrations of Co-60 and Cs-137.

1.2 Objective

The Army Base Closure and Realignment Committee has identified Fort McCellan as an installation for closure. Fort McCellan will therefore be closed, and the Pelham Range will be licensed to the Alabama Army National Guard (USACE, 1998). As part of this closure, the Fort's NRC radioactive materials licenses 01-02861-05 will be terminated. Termination of these licenses requires submission of a formal decontamination and decommissioning (D&D) plan. Cleanup levels form an integral part of this D&D plan. This report is intended to present the site-specific derived cleanup guidelines proposed for the Pelham Range Burial Mound and to document the method used to derive them.

1.3 Organization of the Report

The remainder of this report is organized into four sections. Section 2, contains a description of the approach used to derive the cleanup goals. Section 3 contains the results of the exposure assessment performed on selected receptors. Section 4 presents the DCGL's and summarizes the report's recommendations. Appendix A contains the modeling parameters and results that form the basis for the DCGL's developed in this report.

2.0 Description of Approach

In order to terminate a NRC radioactive materials license and release a site, a licensee must demonstrate that the site is suitable for release in accordance with the criteria for decommissioning in Subpart E, "Radiological Criteria for License Termination," of 10 CFR 20, "Standards of Protection Against Radiation." This report uses site-specific information to derive the release criteria that must be met before the license can be terminated and the site licensed to the Alabama Army National Guard. These criteria are based on an annual dose limit of 25 mrem/y to the group of individuals reasonably expected to receive the greatest exposure to residual radioactivity, assuming reasonable and expected use of the land in the future.

Information on the site was assembled and analyzed to develop a conceptual model of the site. This model provided a basis to identify the important sources of contamination, principle means of release and environmental transport, and likely receptors, both now and in the future.

Once these were identified, site-specific derived cleanup guidelines (DCGLs) were calculated using RESRAD version 5.82 (ANL, 1995). RESRAD was developed partially under contract with the NRC and is widely used in the decommissioning industry for calculating doses and soil cleanup criteria.

After the site-specific DCGL's were determined, the philosophy of ALARA was then applied. The DCGLs were compared to the levels detectable by the survey method and the soil sorter equipment. The lowest of these limits was identified as the lowest practical site-specific soil cleanup criteria. These lowest levels are the cleanup criteria proposed for Burial Mound soils.

Because of the bimodal distribution of the radionuclide concentrations in the pile, it is expected that the average concentration of residual contamination in the pile will be much lower than the cleanup criteria. Using measured data, the expected residual concentrations that should result from the application of these cleanup criteria were then determined. These residual concentrations were then compared to the concentrations predicted to produce 25 mrem/y to the critical member of a resident family, assuming immediate occupancy of the site. This commonly applied standard default exposure scenario was selected to determine compliance with unconditional release of the property.

3.0 Exposure Assessment

The purpose of an exposure assessment is to estimate the nature and magnitude of exposures from a site under current and plausible future conditions. This is accomplished by following a phased approach that involves the following tasks:

- Characterizing the exposure setting on and around the site,
- Identifying potentially complete human exposure pathways, and
- Quantifying the magnitude of plausible contaminant intakes by hypothetical receptors.

This section presents a description of the methods used to evaluate exposures from Pelham Mound, and the results of that assessment. The setting and physical characteristics of the Site are summarized below in Section 3.1. Section 3.2 presents the conceptual model describing the sources, contaminant migration, receptors, and exposure routes evaluated for the Mound. The methods used to quantify potential intakes by plausible receptors and the estimated intakes are presented in Section 3.3.

3.1 Characterization of the Site

The following sections summarize information on the physical setting of the site, its history, and its current and projected uses. For more detailed information on these and related subjects, see the Industrial Radiation Study No. 27-MH-0987-R2-97 (USCHPPM, Jan 1996).

3.1.1 Physical Setting

Fort McClellan is an army based sited on 45,679 acres adjacent to Anniston, AL. It is divided into three areas: the Main Post, the Choccolocco Corridor, and the Pelham Range. The Pelham Range Burial Mound is located at UMT coordinates 593300 E, 3732500 N, which is near the northwest corner of Pelham Range, on the northern end of the Battle Drill Area of Range 24C. The mound is oblong in shape and is approximately 25 meters long by 15 meters wide. It extends to three to four meters below grade, and is piled up to approximately two meters above grade in places.

3.1.2 Site History

Part of the area known as Rideout Field served as a radiological training area from the mid 1950's through May of 1972. The area was also used as an active radiological material burial

site as late as 1959. Best information indicates radioactive low-level laboratory waste and contaminated dirt were collected and placed in one burial mound located in the northwest corner of the Pelham Range.

3.1.3 Land Use

The area surrounding the burial mound has historically been used as a maneuver training area for students at the U.S. Army Chemical and Military Police Schools, and by Active Duty, Reserve, and Alabama National Guard units. The area will continue to be used as a maneuver training area for the foreseeable future. The potential for unexploded munitions also exists in some areas of the Pelham Range[1].

Because of its future use and the potential danger to the public from unexploded munitions, access will generally be restricted to authorized personnel only. This restriction will be enforced by limited access along controlled roads, and physical barriers like fences and gates.

3.1.4 Conceptual Model of the Site

The conceptual model for the Pelham Range Burial Mound has been developed to provide the basis for identifying and evaluating the potential risks to human health from radioactive materials at this site. This model presents the relationships between the following elements necessary to construct a complete exposure pathway:

- Sources and contaminants
- Release mechanisms
- Transport pathways
- Exposure routes
- Receptors

Figure 3-1 presents the conceptual model for potential human exposure to the contaminants detected in the Burial Mound soil. The objective of this conceptual model development and the concurrent analysis of potential exposure routes and receptors is to focus subsequent efforts on those pathways and sources that drive the potential impacts on human health risk.

---

[1] The Decommissioning Plan includes an OE Avoidance Plan to elminate an unexpected encounter (with odnance and explosives).

The conceptual model illustrated in Figure 3-1 of this appendix traces the exposure pathways and receptors for the Mound from the source through likely release mechanisms and exposure routes to potential receptors. The conceptual model also indicates which exposure routes are carried through the quantitative dose assessment for each receptor under current and far-future land-use scenarios.

3.1.5 Sources and Release Mechanisms

The radionuclides of concern at this site are Co-60 and Cs-137. This material is mixed with soil and the soil has been collected into a discrete pile that extends both above and below grade level. This burial mound is the principle source of radioactive material at this site, and it is a relatively small, discrete area. A previous investigation (USCHPPM, 1996) collected data that indicate the majority of the pile does not contain measurable amounts of Co-60 or Cs-137 (Figures 3-2 and 3-3 of this appendix.)

The exposed surface of the pile is subject to limited water and wind erosion. In addition, rainwater percolating through the pile may mobilize one or both of the contaminants. The impact of these releases are expected to be minimal during the study duration, but they are included the conceptual model for completeness.

3.1.6 Potential Transport Pathways

If released, these contaminants could be transported by wind, surface water, or groundwater to the vicinity of the receptor. In addition, plants may absorb some contaminants via root uptake. Each of these pathways has been included in the exposure assessment calculations made in this report.

3.1.7 Potential Exposure Routes

A receptor can come into contact with the Co-60 and Cs-137 in the Mound in a variety of ways, generally as the result of a receptor's behavior or lifestyle that brings him/her into contact with a contaminated exposure medium. This assessment defines a route mechanism as a stylized description of the behavior that brings a receptor into contact with a contaminated medium.

An exposure route describes how a radionuclide may enter or affect humans. Exposures are divided into two types: internal exposures and external exposures. Internal exposures occur when contaminants enter the human body through inhalation and ingestion. External exposures do not require physical contact and occur when a receptor is close to a source of radiation. Proximity to such a source can result in the irradiation of an individual by penetrating radiation.



Figure 3-1 Conceptual Model of Exposures for Pelham Range Burial Ground





Figure 3.2 Frequency Distributions of Co-60 and Cs-137 Concentrations in Burial Mound Soil (Data from USCHPPM, 1996)



Figure 3-3 Cumulative Probability Plot of Co-60 and Cs-137 Concentrations in Soil

The remainder of this section describes the exposure routes evaluated in this assessment. The receptors evaluated for these exposure routes are described in Sections 3.2.4 and Section 3.2.5.

3.1.7.1 Exposures from Inhalation

This route assumes a receptor is immersed in air containing suspended particulates from the Mound. Subsequent exposures occur via inhalation.

3.1.7.2 Exposures from Ingestion

This route assumes a receptor eats or drinks contaminated soil, food, or water.

3.1.7.3 Direct Exposure to Radiation

Direct exposures to radiation from radioactive material can occur when a receptor is near a radioactive source. Physical contact with a contaminated exposure medium is not necessary for exposure to external radiation to occur. The magnitude of exposure is directly related to the distance of the receptor from the source, the activity of the radionuclides present, and the amount of shielding between the source and the postulated receptor.

3.1.8 Receptor Scenarios Considered

3.1.8.1 Role of Land Use in Receptor Selection

The release criteria for the Pelham Range "Burial Mound" and its surrounding area will be based, in part, on the projected land use of the site. This is because the land use of this site will dictate the types of receptors that may be exposed at the site. The land use will also influence their behavior, and their behavior will influence the types and durations of possible exposures.

This site is currently part of a military base. At decommissioning, it will be licensed to the Alabama Army National Guard for use as a battle drill training area. This land use is expected to continue for the foreseeable future in order to comply with the recommendations of the 1995 Defense Base Closure and Realignment Committee. Therefore public access to the site will be restricted by administrative controls and base security. This land use limits plausible human receptors to military personnel and transient members of the general public.

At some point in the future, the Pelham Range may be released to the general public for unrestricted use. It is not anticipated that this will occur in the near future, but to assess the potential doses from this possibility, hypothetical exposures are assessed from materials in the

Pelham Mound to a family living in a home built on the site after 30 years of use as a National Guard training facility.

3.1.8.2 Receptor Scenarios Considered Under Planned Use Conditions

The possible receptors envisioned for the planned land use at Fort McClellan are military personnel and transient members of the general public. Two receptors have been selected to represent these two classes of exposure, a National Guard soldier on maneuvers and a hunter.

3.1.8.2.1 National Guard Scenario

It is assumed that members of the National Guard spend 14 days a year training at Fort McCellan. This 14 days includes 1 day for mobilization, 1 day for demobilization, and 12 days for maneuvers and training. It was assumed an individual would spend no more than 12 hours a day at the site itself. It is felt that a daily exposure of 12 hours is a very conservative, health protective estimate, because it is unlikely that training activities would allow an individual to spend all their time at one location. These assumptions produce an annual exposure time of 144 hours, which is equivalent to 18 eight hour days. The guardsman is assumed to remain in the Guard for 20 years and receive annual exposures during that period.

3.1.8.2.2 Hunter Scenario

It is assumed that transient exposures could occur to individuals who hunt on Ft. McCellan. A hunter could conceivably build a hunting blind on the site and spend part of each day waiting for game. This site is not as attractive as some surrounding terrain, and it is unlikely that these individuals would spend all of their time on this site, but overestimating the exposure duration in this manner is judged to be health protective.

It was assumed that a hunter will take 1 week of vacation to hunt. Assuming he spends one day building his blind during the weekend before hunting season, he could be present on the site for a total of ten days. If the hunter spends 10 hours a day on this site, the hunter will be in close proximity of the site for 100 hours each year. The hunter is assumed to hunt the site for 30 years.

3.1.8.3 Receptor Scenarios Considered Under Far-Future Use Conditions

This scenario assumes governmental control of the site ceases at 30 years in the future. If administrative controls cease, many types of land use become plausible. In this case, it was judged that residential types of land use would produce the greatest exposure potential.

Therefore, a typical residential scenario has been used to evaluate the exposures from these conditions.

3.1.8.3.1 Resident Adult Scenario

A typical residential adult was selected to represent exposures. This hypothetical resident lives on the property but works elsewhere. This adult spends most of the time indoors, but spends some time engaged in outdoor activities such as light gardening. The resident is assumed to grow his own vegetables and resides at the site for 30 years.

3.1.8.3.2 Resident Child Scenario

A typical residential family consists of both adults and children. A hypothetical child, aged 1 to 6 years old, was selected for evaluation because children are a critical subpopulation that may ingest higher amounts of soil than an adult. The child's behavior pattern is similar to the adult described above, except that less time is spent off-site.

3.1.9 Receptors Selected

Four receptor scenarios were initially considered to be plausible representations of potential uses of this site. Three of these receptors were selected for quantitative evaluation in this study. These potential receptors are described in the Section 3.2.4. Because the radionuclides of potential concern are strong gamma emitters, their most important exposure pathway is external radiation from the ground surface. The total exposure via this pathway is directly proportional to the time exposed to the contaminated ground surface, which means the calculated doses attributed to the receptors will be roughly proportional to the their assumed exposure times.

3.1.9.1 Receptor Selected for Evaluation of Planned Land Use

Two receptors are described in the section introducing potential receptors that were judged to be plausible under the planned use of the site. Table 3-1 presents the parameters describing the behavior of the two receptors. Because the radionuclides of potential concern are strong gamma emitters, their most important exposure pathway is external radiation from the ground surface. The total exposure via this pathway is directly proportional to the time exposed to the contaminated ground surface, which means the calculated doses attributed to the two receptors will be proportional to the their assumed exposure times.

TABLE 3-1 PARAMETERS USED TO ESTIMATE POTENTIAL EXPOSURES FOR THE FORT MCCELLAN PELHAM BURIAL MOUND

	Current Hypothetical Receptor				Future Hypothetical Receptors			
Pathway	National Guardsman		Civilian Hunter		Resident Adult		Resident Child	
Parameter (units)	Age 19+		Age 19+		Age 19+		Age 1-6	
Time and duration of exposure								
ET indoors (hr/d)	0	a	0	b	16.4	c	20.2	d
ET outdoors (hr/d)	12	a	10	b	2	c	3.3	d
EF (d/yr)	12	a	10	b	350	e	350	f
ED (yr)	20	a	30	g	30	e	6	f
Inhalation of dust								
IR (m3/hr)	0.63	h	0.63	h	0.63	h	0.31	i
Incidental ingestion of soil								
IR (kg/d)	0.00005	j	0.00005	j	0.00005	j	0.00020	k
Ingestion of food and water								
IR_{veg} (kg/d)	na	l	na	l	0.309	m	0.129	n
IR_{fruit} (kg/d)	na	l	na	l	0.244	m	0.102	n
IR_{water} (L/d)	na	l	na	l	2.3	o	1.3	p

a - The guardsman is assumed to spend 12 hours a day training outdoors in equipment or facilities located over the burial ground. The training lasts 2 weeks each year, with two days subtracted for mobilization and demobilization activities. The guardsman is assumed to remain in the Guard until eligible for a pension at 20 years.

b - The civilian hunter is assumed to spend 10 hours a day in a blind located on the burial ground site. The hunter is assumed to spend one day prior to hunting season building a blind, and spend 1 week of vacation (five days plus two weekends) each year actively hunting.

c - Recommended values for indoor residential and outdoor exposures from Table 5-176 in Exposure Factors Handbook, EPA 1997.

d - Time weighted average of mean values for indoor residential and outdoor exposures for 1-4 year old from Table 5-131 and Table 15-132 in Exposure Factors Handbook, EPA 1997. (Indoor = 1211 min/d, and Outdoor = 195 min/d).

e - The resident is assumed to spend 30 years living in a house built on the site, and take two weeks of vacation away from the home each year (EPA's 1991 OSWER Directive 9285.6-03.).

f - Reflects six years of life between birth and age 6.

g - The hunter is assumed to hunt the site each year he is in the area. Based on a 30 years residence time (Exposure Factors Handbook, EPA 1997).

h - Inhalation rate of adult. From Table 5-23 of Exposure Factors Handbook, EPA 1997 (15.2 m3/d x d/24 h).

i - Time weighted average of recommended values inhalation rate of children, ages 1-6. From Table 5-23 of Exposure Factors Handbook, EPA 1997. (1 y x 4.5 m3/d+2 y x 6.8 m3/d + 2y x 8.3m3/d + 1y x 10 m3/d) / 6y x d/24 h.

j - Rate of incidental soil ingestion by adult (Exposure Factors Handbook, EPA 1997, Table 4-23).

k - Soil ingestion rate (Exposure Factors Handbook, EPA 1997, pg. 4-24).

l - These receptors are expected to bring their own packaged food and water.

m - 95%tile combined home produced fruit and vegetable ingestion rate by a 71.8 kg adult (Exposure Factors Handbook, EPA 1997, pg. 9-44).

n - 95%tile combined home produced fruit and vegetable ingestion rate by a 30.1 kg child (Exposure Factors Handbook, EPA 1997, pg. 9-44).

o - 90%tile Tap water ingestion rate by an adult (Exposure Factors Handbook, EPA 1997, pg. 3-26).

p - 90%tile Tap water ingestion rate by a 1-10 year old child (Exposure Factors Handbook, EPA 1997, pg. 3-26).

Since the Guardsman Scenario assumes receptor behavior that produces 144 hours of exposure each year, while the Hunter Scenario produces 100 hours a year, the calculated dose from the Hunter Scenario will be less than 70% of the calculated dose from the Guardsman Scenario. Remediating the site to levels which are safe for the Guardsman will therefore insure the safety of the Hunter as well. Therefore, the Guardsman was selected to evaluate the highest plausible exposures from the site's planned land use.

3.1.9.2 Receptors Selected for Evaluation of Far Future Conditions

Two potential receptors, the resident adult and the resident child, are described in the section introducing potential receptors that were judged to be plausible in the unlikely event that the site is released to the public after 30 years. Therefore potential exposures to the resident adult and the resident child were both quantitatively evaluated in this study. Table 3-1 presents the parameters used to quantify the behavior of the two receptors.

3.2 Application of RESRAD

The guardsman and resident scenarios involve activities performed while on top of the burial mound soil. The computer code RESRAD 5.82 (ANL,1993) was used to assess the potential long-term doses from these activities. Default input parameters were used, except where otherwise noted in Table 3-1. Summary input files presented in Appendix A contain a detailed list of parameter values used in the RESRAD calculations for each receptor assessed.

Doses to a guardsman training on a uniform layer of the mound material during 30 years of institutional control were calculated for the following pathways using RESRAD default values:

- Soil ingestion
- Inhalation of resuspended soil, and
- Direct exposure to external radiation.

Doses to a resident living on a uniform layer of the mound material after 30 years of institutional control ends were calculated for the following pathways using RESRAD default values:

- Drinking water
- Food ingestion (home-grown vegetables)
- Soil ingestion
- Inhalation of resuspended soil, and
- Direct exposure to external radiation.

3.3 Summary of Receptor Exposures

The results of the RESRAD runs described in Section 3.3 are summarized in Table 3-2. The residual soil concentrations for the guardsman scenario yield a calculated dose of 25 mrem/y. The soil concentrations at for the residential scenarios will decay to concentrations that yield a calculated dose of 25 mrem/y after the site is used by the National Guard for 30 years.

Table 3-2 Allowable Concentrations in Current Soil For Current and Far-Future Land Use

Guardsman Scenario		Resident Adult Scenario		Resident Child Scenario	
Co-60	Cs-137	Co-60	Cs-137	Co-60	Cs-137
100	478	154	26	123	21

[a] If a resident adult moves onto the site in 30 years, the Co-60 concentration listed here will have decayed to 3 pCi/g, and the Cs-137 concentration will have decayed to 13 pCi/g.

[b] If a resident child moves onto the site in 30 years, the Co-60 concentration listed here will have decayed to 2 pCi/g, and the Cs-137 concentration will have decayed to 11 pCi/g.

3.3.1 Guardsman

Exposures to the hypothetical guardsman evaluated in this study would be dominated by direct gamma radiation emitted directly by the soil. The soil concentrations are based on occupation of the Pelham Mound by a trainee for 144 hours per year. This exposure time is considered excessive, so the resulting derived guidelines calculated for this receptor are judged to be very conservative.

3.3.2 Resident Adult with Garden

Exposures to a hypothetical adult with a garden evaluated in this study would be dominated by direct gamma radiation emitted directly by the soil (over 99% for Co-60 and almost 90% of the Cs-137) . About 10% of the dose from Cs-137 is associated with consumption of food from a garden.

3.3.3 Resident Child with Garden

Exposures to a hypothetical child with a garden evaluated in this study are very similar to those of the adult discussed earlier. They would be dominated by direct gamma radiation emitted directly by the soil (over 99% for Co-60 and almost 90% of the Cs-137) . About 10% of the dose from Cs-137 would be attributable to consumption of food from a garden. The dose based

soil guidelines for the child are lower than for the parent because the child is assumed to spend more time per year on site exposed to the underlying soil.

4.0 Derived Cleanup Guidelines

4.1 Role of Dose Based Criteria

Potential radiation exposures from the Pelham Mound and surrounding areas must be limited to safe levels before the NRC license can be terminated. The NRC has determined that a dose limit of 25 mrem/y is health protective at NRC licensed sites (10CFR20). The Derived Cleanup Guidelines (DCGLs) developed in this document are judged to be meet this requirement under the planned and far future land use proposed for the site.

Application of these DCGLs to the soil pile will result in removal of soil with concentrations exceeding these guidelines. Because only a small percentage of soil samples contained Co-60 or Cs-137 concentrations exceeding the guidelines, it is anticipated that the average concentrations of these two radionuclides in the soil remaining on site will be much less than those required for unconditional release of the site.

4.2 Dose Based Cleanup Guidelines

The dose based cleanup level for a nuclide is the smallest of the soil concentrations calculated for the three receptors (Table 3-2). This assures that final cleanup goals for individual radionuclides are health protective for all the receptors evaluated. The individual dose based Derived Cleanup Guidelines (DCGLs) are 100 pCi/g for Co-60 and 21 pCi/g Cs-137.

4.3 Influence of Multiple Contaminants on Cleanup Guidelines

A multi pathway analysis was used to develop individual dose based DCGL's of 100 pCi/g Co-60 and 21 pCi/g Cs-137. Because Co-60 and Cs-137 are both present at the site, it is necessary to assure that their combined activities do not result in exposures that exceed 25 mrem/y. A simple way of achieving this is to set a combined cleanup standard to be equal to the smaller of the two cleanup criteria in soil. This approach protects human health and the environment at the site and provides one clear, reproducible soil cleanup objective for the proposed remedial activities.

The smaller of the two dose-based cleanup guidelines calculated in this report is the 21 pCi/g limit for Cs-137. This approach would yield a combined cleanup criteria of 21 pCi/g for soil containing both radionuclides.

4.4 Derived Cleanup Guidelines Based on Expected Land Use

The approach used in this report sets 21 pCi/g as the cleanup concentration for site soils containing a mixture of Cs-137 and C-60 at the site. The final set of DCGLs for this site is presented in Table 4-1. The limits for individual nuclides will be applied if it can be demonstrated that only one of these radionuclides is present in the soil. If the presence of both is suspected, the value of 21 pCi/g will be compared to the sum of their activities.

4.5 Impact of DCGLs and Soil Sorting on Final Status of Burial Mound Soil

The previous section presents DCGL's for the Burial Mound soil. For the purpose of this project these become the maximum residual soil concentrations allowed to remain, averaged over a 100 m^2 throughout the survey area. However, the philosophy of ALARA (As Low As Reasonably Achievable) acts as an incentive to reduce the residual concentrations to lower levels, if practical.

The soil sorter is capable of discerning Co-60 and Cs-137 at 9 pCi/g and 16 pCi/g, respectively. These levels are lower than the DCGLs and have been selected as the defacto delineation between soil that will remain on site and soil that will be packaged and transported to an off-site disposal facility.

Analysis of the soil data, collected from various locations and depths of the pile, indicates the bulk of the material in the pile does not contain appreciable levels of either Co-60 or Cs-137. In fact, less than 10% of the samples contain these radionuclides in concentrations exceeding the contract MDA (Figures 3-2 and 3-3 in this appendix). This implies that the pile is essentially clean dirt, with some limited areas of elevated concentration.

The intent of the remedial action is to remove soil containing more than 9 pCi/g Co-60 or 16 pCi/g Cs-137 from the site. If this is done, current sampling results indicate the average concentrations of each isotope in the remaining soil will be at or near 0.1 pCi/g (Table 4-2).

4.6 Health Protectiveness of Expected Residual Soil Contamination Levels.

Once the average expected residual concentration was determined, RESRAD was used to confirm the health-protectiveness of the remedy. This was done by calculating the soil concentrations predicted to produce 25 mrem/y to the critical member of a hypothetical residential family living on the site immediately after remediation.

Based on the results of the residential scenario at 30 years, it was determined that the small child was the critical member of the proposed receptor group. Attachment A of this Appendix

contains the results of these calculations. In summary, these calculations determined that the DCGL at t(0) for Co-60, would be 2.4 pCi/g, while for Cs-137 it would be 10.6 pCi/g. The average concentrations projected for the soil after remediation are well below these levels (Table 4-3).

After remediation, the health protectiveness of the site will be verified. Dose rates from external exposures will be measured throughout the site. The contribution to dose from soil, water, and on-property food sources will be investigated using the measured values of Co-60 and Cs-137 in soil and the RESRAD scenario for immediate occupation by a hypothetical farm family. These results will be used to determine the annual dose to the hypothetical receptor from the remediated site. The post-remediation dose (above background) meet the 25 mrem/y release limit to allow for unconditional release.

Table 4-1 DCGL's of Individual and Mixed Radionuclides at the Pelham Range Burial Mound for Expected Use

Radionuclide	Soil Limit
Co-60	100 pCi/g
Cs-137	21 pCi/g
Co-60 + Cs-137	21 pCi/g

Table 4-2 Summary Statistics for Pelham Field Burial Mound Assuming Various Soil Cleanup Levels

Summary Stats for Cs-137					Summary Stats for Co-60				
Statistic	**No Action**	**DCGL 21 pCi/g**	**Scan Limit 21 pCi/g**	**Sorter Limit 16 pCi/g**	**Statistic**	**No Action**	**DCGL 100 pCi/g**	**Scan Limit 21 pCi/g**	**Sorter Limit 9 pCi/g**
n	585	583	583	581	n	585	582	578	575
mean	0.46	0.11	0.11	0.06	mean	1.83	0.40	0.15	0.08
SD	7.53	0.96	0.96	0.48	SD	20.72	3.32	1.09	0.60
$UCL_{95\%\ mean}$*	**0.97**	**0.18**	**0.18**	**0.09**	$UCL_{95\%\ mean}$*	**3.24**	**0.63**	**0.22**	**0.13**
Max. value	179.0	17.0	17.0	9.5	Max. value	330.0	49.0	14.0	8.0
95th Prcntl	0.30	0.29	0.29	0.20	95th Prcntl	0.20	0.20	0.10	0.10
iterated t	1.647508	1.647518	1.647518	1.647522	*iterated t*	1.647508	1.647522	1.647541	1.647555
t calc	*0.0499958*	*0.0499957*	*0.0499957*	*0.0499962*	*t calc*	*0.0499958*	*0.0499957*	*0.0499957*	*0.0499956*

*The 95% UCL on the arithmetic mean (= ArithMean + (t_score * StDev / SQRT(n))

TABLE 4-3 Summary Comparison of Predicted Soil Concentrations After Remediation, and Concentrations of Soil Required to Produce 25 mrem/y to the Critical Member of a Hypothetical Resident Family

Radionuclide	Predicted Soil Concentrations Required to Produce 25 mrem/y to a Resident Child	Projected $95\%UCL_{mean}$ for Soil Concentration In Pile after Sorting
Co-60	2.4 pCi/g	0.13 pCi/g
Cs-137	10.6 pCi/g	0.09 pCi/g

5.0 References

ANL, 1993 "Manual for Implementing Residual Radioactive Material Guidelines Using RESRAD, Version 5.0., September 1993." ANL/EAD/LD-2. USDOE, Environmental Assessment Division, Argonne National Laboratory.

ANL, 1998 RESRAD for Windows, 5.82. Environmental Assessment Division, Argonne National Laboratory.

USACE, 1998 "Final Environmental Impact Statement, Disposal and Reuse of Fort McClellan, Alabama, August 1998." US Army Corps of Engineers, Mobile District.

USCHPPM, 1996 "Industrial Radiation Study No 27-MH-0987-R2-97. Pelham Range Burial Mound, Fort McClellan, Alabama. 29 August – 15 September 1995 and 14-28 January 1996. US Army Center for Health Promotion and Preventive Medicine.

USEPA, 1997 "Exposure Factors Handbook, Volumes 1, 2, and 3" August 1997, EPA/600/P-95/002Fa, b, & c. Us Environmental Protection Agency.

Attachment A RESRAD Output

Attachment A RESRAD Output

A.1 Results of RESRAD Calculations for Guardsman Scenario A-1

A.1.1 Co-60: Guardsman Summary Output File A-1

A.1.2 Cs-137: Guardsman Summary Output File A-19

A.2 Results of RESRAD Calculations for Resident Adult Scenario A-37

A.2.1 Co-60: Resident Adult Summary Output File A-37

A.2.2 Cs-137: Resident Adult Summary Output File A-55

A.3 Results of RESRAD Calculations for Resident Child Scenario A-73

A.3.1 Co-60: Resident Child Summary Output File A-73

A.3.2 Cs-137: Resident Child Summary Output File A-92

A.1 Results of RESRAD Calculations for Guardsman Scenario

A.1.1 Co-60: Guardsman Summary Output File

RESRAD, Version 5.82 T½ Limit = 0.5 year 12/15/98 09:15 Page 1

Summary : Co 60, guardsman, 10000m2 File: C060G10K.RAD

Table of Contents

Part I: Mixture Sums and Single Radionuclide Guidelines

Dose Conversion Factor (and Related) Parameter Summary ... 2
Site-Specific Parameter Summary 3
Summary of Pathway Selections 6
Contaminated Zone and Total Dose Summary 7
Total Dose Components
Time = 0.000E+00 8
Time = 1.000E+00 9
Time = 3.000E+00 10
Time = 1.000E+01 11
Time = 3.000E+01 12
Time = 5.000E+01 13
Time = 1.000E+02 14
Time = 3.000E+02 15
Time = 1.000E+03 16
Dose/Source Ratios Summed Over All Pathways 17
Single Radionuclide Soil Guidelines 17
Dose Per Nuclide Summed Over All Pathways 18
Soil Concentration Per Nuclide 18

Dose Conversion Factor (and Related) Parameter Summary
File: DOSFAC.BIN

Menu	Parameter	Current Value	Default	Parameter Name
B-1	Dose conversion factors for inhalation, mrem/pCi:			
B-1	Co-60	2.190E-04	2.190E-04	DCF2(1)
D-1	Dose conversion factors for ingestion, mrem/pCi:			
D-1	Co-60	2.690E-05	2.690E-05	DCF3(1)
D-34	Food transfer factors:			
D-34	Co-60 , plant/soil concentration ratio, dimensionless	8.000E-02	8.000E-02	RTF(1,1)
D-34	Co-60 , beef/livestock-intake ratio, (pCi/kg)/(pCi/d)	2.000E-02	2.000E-02	RTF(1,2)
D-34	Co-60 , milk/livestock-intake ratio, (pCi/L)/(pCi/d)	2.000E-03	2.000E-03	RTF(1,3)
D-5	Bioaccumulation factors, fresh water, L/kg:			
D-5	Co-60 , fish	3.000E+02	3.000E+02	BIOFAC(1,1)
D-5	Co-60 , crustacea and mollusks	2.000E+02	2.000E+02	BIOFAC(1,2)

Site-Specific Parameter Summary

Menu	Parameter	User Input	Default	Used by RESRAD (If different from user input)	Parameter Name
R011	Area of contaminated zone (m**2)	1.000E+04	1.000E+04	---	AREA
R011	Thickness of contaminated zone (m)	2.000E+00	2.000E+00	---	THICK0
R011	Length parallel to aquifer flow (m)	not used	1.000E+02	---	LCZPAQ
R011	Basic radiation dose limit (mrem/yr)	2.500E+01	3.000E+01	---	BRDL
R011	Time since placement of material (yr)	0.000E+00	0.000E+00	---	TI
R011	Times for calculations (yr)	1.000E+00	1.000E+00	---	T(2)
R011	Times for calculations (yr)	3.000E+00	3.000E+00	---	T(3)
R011	Times for calculations (yr)	1.000E+01	1.000E+01	---	T(4)
R011	Times for calculations (yr)	3.000E+01	3.000E+01	---	T(5)
R011	Times for calculations (yr)	5.000E+01	1.000E+02	---	T(6)
R011	Times for calculations (yr)	1.000E+02	3.000E+02	---	T(7)
R011	Times for calculations (yr)	3.000E+02	1.000E+03	---	T(8)
R011	Times for calculations (yr)	1.000E+03	0.000E+00	---	T(9)
R011	Times for calculations (yr)	not used	0.000E+00	---	T(10)
R012	Initial principal radionuclide (pCi/g): Co-60	1.000E+02	0.000E+00	---	S1(1)
R012	Concentration in groundwater (pCi/L): Co-60	not used	0.000E+00	---	W1(1)
R013	Cover depth (m)	0.000E+00	0.000E+00	---	COVER0
R013	Density of cover material (g/cm**3)	not used	1.500E+00	---	DENSCV
R013	Cover depth erosion rate (m/yr)	not used	1.000E-03	---	VCV
R013	Density of contaminated zone (g/cm**3)	1.500E+00	1.500E+00	---	DENSCZ
R013	Contaminated zone erosion rate (m/yr)	0.000E+00	1.000E-03	---	VCZ
R013	Contaminated zone total porosity	4.000E-01	4.000E-01	---	TPCZ
R013	Contaminated zone effective porosity	2.000E-01	2.000E-01	---	EPCZ
R013	Contaminated zone hydraulic conductivity (m/yr)	1.000E+01	1.000E+01	---	HCCZ
R013	Contaminated zone b parameter	5.300E+00	5.300E+00	---	BCZ
R013	Average annual wind speed (m/sec)	2.000E+00	2.000E+00	---	WIND
R013	Humidity in air (g/m**3)	not used	8.000E+00	---	HUMID
R013	Evapotranspiration coefficient	5.000E-01	5.000E-01	---	EVAPTR
R013	Precipitation (m/yr)	1.000E+00	1.000E+00	---	PRECIP
R013	Irrigation (m/yr)	2.000E-01	2.000E-01	---	RI
R013	Irrigation mode	overhead	overhead	---	IDITCH
R013	Runoff coefficient	2.000E-01	2.000E-01	---	RUNOFF
R013	Watershed area for nearby stream or pond (m**2)	not used	1.000E+06	---	WAREA
R013	Accuracy for water/soil computations	not used	1.000E-03	---	EPS
R014	Density of saturated zone (g/cm**3)	1.500E+00	1.500E+00	---	DENSAQ
R014	Saturated zone total porosity	4.000E-01	4.000E-01	---	TPSZ
R014	Saturated zone effective porosity	2.000E-01	2.000E-01	---	EPSZ
R014	Saturated zone hydraulic conductivity (m/yr)	1.000E+02	1.000E+02	---	HCSZ
R014	Saturated zone hydraulic gradient	2.000E-02	2.000E-02	---	HGWT
R014	Saturated zone b parameter	5.300E+00	5.300E+00	---	BSZ
R014	Water table drop rate (m/yr)	1.000E-03	1.000E-03	---	VWT
R014	Well pump intake depth (m below water table)	1.000E+01	1.000E+01	---	DWIBWT
R014	Model: Nondispersion (ND) or Mass-Balance (MB)	ND	ND	---	MODEL
R014	Well pumping rate (m**3/yr)	2.500E+02	2.500E+02	---	UW
R015	Number of unsaturated zone strata	not used	1	---	NS

Site-Specific Parameter Summary (continued)

Menu	Parameter	User Input	Default	Used by RESRAD (If different from user input)	Parameter Name
R015	Unsat. zone 1, thickness (m)	not used	4.000E+00	---	H(1)
R015	Unsat. zone 1, soil density (g/cm**3)	not used	1.500E+00	---	DENSUZ(1)
R015	Unsat. zone 1, total porosity	not used	4.000E-01	---	TPUZ(1)
R015	Unsat. zone 1, effective porosity	not used	2.000E-01	---	EPUZ(1)
R015	Unsat. zone 1, soil-specific b parameter	not used	5.300E+00	---	BUZ(1)
R015	Unsat. zone 1, hydraulic conductivity (m/yr)	not used	1.000E+01	---	HCUZ(1)
R016	Distribution coefficients for Co-60				
R016	Contaminated zone (cm**3/g)	1.000E+03	1.000E+03	---	DCUCC(1)
R016	Unsaturated zone 1 (cm**3/g)	not used	1.000E+03	---	DCUCU(1,1)
R016	Saturated zone (cm**3/g)	not used	1.000E+03	---	DCUCS(1)
R016	Leach rate (/yr)	0.000E+00	0.000E+00	1.666E-04	ALEACH(1)
R016	Solubility constant	0.000E+00	0.000E+00	not used	SCLUBK(1)
R017	Inhalation rate (m**3/yr)	9.072E+01	8.400E+03	---	INHALR
R017	Mass loading for inhalation (g/m**3)	1.000E-04	1.000E-04	---	MLINH
R017	Exposure duration	2.000E+01	3.000E+01	---	ED
R017	Shielding factor, inhalation	4.000E-01	4.000E-01	---	SHF3
R017	Shielding factor, external gamma	7.000E-01	7.000E-01	---	SHF1
R017	Fraction of time spent indoors	0.000E+00	5.000E-01	---	FIND
R017	Fraction of time spent outdoors (on site)	1.640E-02	2.500E-01	---	FOTD
R017	Shape factor flag, external gamma	1.000E+00	1.000E+00	>0 shows circular AREA.	FS
R017	Radii of shape factor array (used if FS = -1):				
R017	Outer annular radius (m), ring 1:	not used	5.000E+01	---	RAD_SHAPE(1)
R017	Outer annular radius (m), ring 2:	not used	7.071E+01	---	RAD_SHAPE(2)
R017	Outer annular radius (m), ring 3:	not used	0.000E+00	---	RAD_SHAPE(3)
R017	Outer annular radius (m), ring 4:	not used	0.000E+00	---	RAD_SHAPE(4)
R017	Outer annular radius (m), ring 5:	not used	0.000E+00	---	RAD_SHAPE(5)
R017	Outer annular radius (m), ring 6:	not used	0.000E+00	---	RAD_SHAPE(6)
R017	Outer annular radius (m), ring 7:	not used	0.000E+00	---	RAD_SHAPE(7)
R017	Outer annular radius (m), ring 8:	not used	0.000E+00	---	RAD_SHAPE(8)
R017	Outer annular radius (m), ring 9:	not used	0.000E+00	---	RAD_SHAPE(9)
R017	Outer annular radius (m), ring 10:	not used	0.000E+00	---	RAD_SHAPE(10)
R017	Outer annular radius (m), ring 11:	not used	0.000E+00	---	RAD_SHAPE(11)
R017	Outer annular radius (m), ring 12:	not used	0.000E+00	---	RAD_SHAPE(12)
R017	Fractions of annular areas within AREA:				
R017	Ring 1	not used	1.000E+00	---	FRACA(1)
R017	Ring 2	not used	2.732E-01	---	FRACA(2)
R017	Ring 3	not used	0.000E+00	---	FRACA(3)
R017	Ring 4	not used	0.000E+00	---	FRACA(4)
R017	Ring 5	not used	0.000E+00	---	FRACA(5)
R017	Ring 6	not used	0.000E+00	---	FRACA(6)
R017	Ring 7	not used	0.000E+00	---	FRACA(7)
R017	Ring 8	not used	0.000E+00	---	FRACA(8)
R017	Ring 9	not used	0.000E+00	---	FRACA(9)
R017	Ring 10	not used	0.000E+00	---	FRACA(10)
R017	Ring 11	not used	0.000E+00	---	FRACA(11)
R017	Ring 12	not used	0.000E+00	---	FRACA(12)
R018	Fruits, vegetables and grain consumption (kg/yr)	not used	1.600E+02	---	DIET(1)

Site-Specific Parameter Summary (continued)

Menu	Parameter	User Input	Default	Used by RESRAD (If different from user input)	Parameter Name
R018	Leafy vegetable consumption (kg/yr)	not used	1.40E+01	---	DIET(2)
R018	Milk consumption (L/yr)	not used	9.20E+01	---	DIET(3)
R018	Meat and poultry consumption (kg/yr)	not used	6.30E+01	---	DIET(4)
R018	Fish consumption (kg/yr)	not used	5.40E+00	---	DIET(5)
R018	Other seafood consumption (kg/yr)	not used	9.00E-01	---	DIET(6)
R018	Soil ingestion rate (g/yr)	6.00E-01	3.65E+01	---	SOIL
R018	Drinking water intake (L/yr)	not used	5.10E+02	---	DWI
R018	Contamination fraction of drinking water	not used	1.00E+00	---	FDW
R018	Contamination fraction of household water	not used	1.00E+00	---	FHHW
R018	Contamination fraction of livestock water	not used	1.00E+00	---	FLW
R018	Contamination fraction of irrigation water	not used	1.00E+00	---	FIRW
R018	Contamination fraction of aquatic food	not used	5.00E-01	---	FR9
R018	Contamination fraction of plant food	not used	-1	---	FPLANT
R018	Contamination fraction of meat	not used	-1	---	FMEAT
R018	Contamination fraction of milk	not used	-1	---	FMILK
R019	Livestock fodder intake for meat (kg/day)	not used	6.80E+01	---	LFI5
R019	Livestock fodder intake for milk (kg/day)	not used	5.50E+01	---	LFI6
R019	Livestock water intake for meat (L/day)	not used	5.00E+01	---	LWI5
R019	Livestock water intake for milk (L/day)	not used	1.60E+02	---	LWI6
R019	Livestock soil intake (kg/day)	not used	5.00E-01	---	LSI
R019	Mass loading for foliar deposition (g/m**3)	not used	1.00E-04	---	MLFD
R019	Depth of soil mixing layer (m)	1.50E-01	1.50E-01	---	DM
R019	Depth of roots (m)	not used	9.00E-01	---	DROOT
R019	Drinking water fraction from ground water	1.00E+00	1.00E+00	---	FGWDW
R019	Household water fraction from ground water	not used	1.00E+00	---	FGWHH
R019	Livestock water fraction from ground water	not used	1.00E+00	---	FGWLW
R019	Irrigation fraction from ground water	not used	1.00E+00	---	FGWIR
R19B	Wet weight crop yield for Non-Leafy (kg/m**2)	not used	7.00E-01	---	YV(1)
R19B	Wet weight crop yield for Leafy (kg/m**2)	not used	1.50E+00	---	YV(2)
R19B	Wet weight crop yield for Fodder (kg/m**2)	not used	1.10E+00	---	YV(3)
R19B	Growing Season for Non-Leafy (years)	not used	1.70E-01	---	TE(1)
R19B	Growing Season for Leafy (years)	not used	2.50E-01	---	TE(2)
R19B	Growing Season for Fodder (years)	not used	8.00E-02	---	TE(3)
R19B	Translocation Factor for Non-Leafy	not used	1.00E-01	---	TIV(1)
R19B	Translocation Factor for Leafy	not used	1.00E+00	---	TIV(2)
R19B	Translocation Factor for Fodder	not used	1.00E+00	---	TIV(3)
R19B	Dry Foliar Interception Fraction for Non-Leafy	not used	2.50E-01	---	RDRY(1)
R19B	Dry Foliar Interception Fraction for Leafy	not used	2.50E-01	---	RDRY(2)
R19B	Dry Foliar Interception Fraction for Fodder	not used	2.50E-01	---	RDRY(3)
R19B	Wet Foliar Interception Fraction for Non-Leafy	not used	2.50E-01	---	RWET(1)
R19B	Wet Foliar Interception Fraction for Leafy	not used	2.50E-01	---	RWET(2)
R19B	Wet Foliar Interception Fraction for Fodder	not used	2.50E-01	---	RWET(3)
R19B	Weathering Removal Constant for Vegetation	not used	2.00E+01	---	WLAM
C14	C-12 concentration in water (g/cm**3)	not used	2.00E-05	---	C12WTR
C14	C-12 concentration in contaminated soil (g/g)	not used	3.00E-02	---	C12CZ
C14	Fraction of vegetation carbon from soil	not used	2.00E-02	---	CSOIL
C14	Fraction of vegetation carbon from air	not used	9.80E-01	---	CAIR
C14	C-14 evasion layer thickness in soil (m)	not used	3.00E-01	---	DMC

Site-Specific Parameter Summary (continued)

Menu	Parameter	User Input	Default	Used by RESRAD (If different from user input)	Parameter Name
C14	C-14 evasion flux rate from soil (1/sec)	not used	7.000E-07	---	EVSN
C14	C-12 evasion flux rate from soil (1/sec)	not used	1.000E-10	---	REVSN
C14	Fraction of grain in beef cattle feed	not used	8.000E-01	---	AVFG4
C14	Fraction of grain in milk cow feed	not used	2.000E-01	---	AVFG5
STOR	Storage times of contaminated foodstuffs (days):				
STOR	Fruits, non-leafy vegetables, and grain	1.400E+01	1.400E+01	---	STOR_T(1)
STOR	Leafy vegetables	1.000E+00	1.000E+00	---	STOR_T(2)
STOR	Milk	1.000E+00	1.000E+00	---	STOR_T(3)
STOR	Meat and poultry	2.000E+01	2.000E+01	---	STOR_T(4)
STOR	Fish	7.000E+00	7.000E+00	---	STOR_T(5)
STOR	Crustacea and mollusks	7.000E+00	7.000E+00	---	STOR_T(6)
STOR	Well water	1.000E+00	1.000E+00	---	STOR_T(7)
STOR	Surface water	1.000E+00	1.000E+00	---	STOR_T(8)
STOR	Livestock fodder	4.500E+01	4.500E+01	---	STOR_T(9)
R021	Thickness of building foundation (m)	not used	1.500E-01	---	FLOOR
R021	Bulk density of building foundation (g/cm**3)	not used	2.400E+00	---	DENSFL
R021	Total porosity of the cover material	not used	4.000E-01	---	TPCV
R021	Total porosity of the building foundation	not used	1.000E-01	---	TPFL
R021	Volumetric water content of the cover material	not used	5.000E-02	---	PH2OCV
R021	Volumetric water content of the foundation	not used	3.000E-02	---	PH2OFL
R021	Diffusion coefficient for radon gas (m/sec):				
R021	in cover material	not used	2.000E-06	---	DIFCV
R021	in foundation material	not used	3.000E-07	---	DIFFL
R021	in contaminated zone soil	not used	2.000E-06	---	DIFCZ
R021	Radon vertical dimension of mixing (m)	not used	2.000E+00	---	HMIX
R021	Average building air exchange rate (1/hr)	not used	5.000E-01	---	REXG
R021	Height of the building (room) (m)	not used	2.500E+00	---	HRM
R021	Building interior area factor	not used	0.000E+00	---	FAI
R021	Building depth below ground surface (m)	not used	-1.000E+00	---	DMFL
R021	Emanating power of Rn-222 gas	not used	2.500E-01	---	EMANA(1)
R021	Emanating power of Rn-220 gas	not used	1.500E-01	---	EMANA(2)

Summary of Pathway Selections

Pathway	User Selection
1 -- external gamma	active
2 -- inhalation (w/o radon)	active
3 -- plant ingestion	suppressed
4 -- meat ingestion	suppressed
5 -- milk ingestion	suppressed
6 -- aquatic foods	suppressed
7 -- drinking water	suppressed
8 -- soil ingestion	active
9 -- radon	suppressed
Find peak pathway doses	suppressed

Contaminated Zone Dimensions		Initial Soil Concentrations, pCi/g	
Area:	10000.00 square meters	Co-60	1.000E+02
Thickness:	2.00 meters		
Cover Depth:	0.00 meters		

Total Dose TDOSE(t), mrem/yr
Basic Radiation Dose Limit = 25 mrem/yr
Total Mixture Sum M(t) = Fraction of Basic Dose Limit Received at Time (t)

t (years):	0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	5.000E+01	1.000E+02	3.000E+02	1.000E+03
TDOSE(t):	2.500E+01	2.192E+01	1.684E+01	6.701E+00	4.813E-01	3.458E-02	4.782E-05	1.750E-16	0.000E+00
M(t):	1.000E+00	8.766E-01	6.737E-01	2.680E-01	1.925E-02	1.383E-03	1.913E-06	7.000E-18	0.000E+00

Maximum TDOSE(t): 2.500E+01 mrem/yr at t = 0.000E+00 years

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 0.000E+00 years
Water Independent Pathways (Inhalation excludes radon)

Radio-	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
Nuclide	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	2.500E+01	1.0000	5.516E-07	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.647E-05	0.0000
Total	2.500E+01	1.0000	5.516E-07	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.647E-05	0.0000

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)

As mrem/yr and Fraction of Total Dose At t = 0.000E+00 years
Water Dependent Pathways

Radio-	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
Nuclide	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.500E+01	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.500E+01	1.0000

*Sum of all water independent and dependent pathways.

Summary : Co 60, guardsman, 10000m2 File: C060G10K.RAD

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+00 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	2.192E+01	1.0000	4.836E-07	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.320E-05	0.0000
Total	2.192E+01	1.0000	4.836E-07	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.320E-05	0.0000

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)

As mrem/yr and Fraction of Total Dose At t = 1.000E+00 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.192E+01	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.192E+01	1.0000

*Sum of all water independent and dependent pathways.

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 3.000E+00 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	1.684E+01	1.0000	3.716E-07	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.783E-05	0.0000
Total	1.684E+01	1.0000	3.716E-07	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.783E-05	0.0000

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 3.000E+00 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.684E+01	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.684E+01	1.0000

*Sum of all water independent and dependent pathways.

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+01 years
Water Independent Pathways (Inhalation excludes radon)

Radio-	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
Nuclide	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	6.701E+00	1.0000	1.478E-07	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	7.094E-06	0.0000
Total	6.701E+00	1.0000	1.478E-07	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	7.094E-06	0.0000

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+01 years
Water Dependent Pathways

Radio-	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
Nuclide	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	6.701E+00	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	6.701E+00	1.0000

*Sum of all water independent and dependent pathways.

Summary : Co 60, guardsman, 10000m2 File: C060G10K.RAD

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 3.000E+01 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	4.813E-01	1.0000	1.062E-08	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	5.096E-07	0.0000
Total	4.813E-01	1.0000	1.062E-08	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	5.096E-07	0.0000

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)

As mrem/yr and Fraction of Total Dose At t = 3.000E+01 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	4.813E-01	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	4.813E-01	1.0000

*Sum of all water independent and dependent pathways.

Summary : Co 60, guardsman, 10000m2 File: C060G10K.RAD

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 5.000E+01 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	3.458E-02	1.0000	7.630E-10	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	3.661E-08	0.0000
Total	3.458E-02	1.0000	7.630E-10	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	3.661E-08	0.0000

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 5.000E+01 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	3.458E-02	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	3.458E-02	1.0000

*Sum of all water independent and dependent pathways.

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+02 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	4.782E-05	1.0000	1.055E-12	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	5.063E-11	0.0000
Total	4.782E-05	1.0000	1.055E-12	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	5.063E-11	0.0000

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+02 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	4.782E-05	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	4.782E-05	1.0000

*Sum of all water independent and dependent pathways.

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 3.000E+02 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	1.750E-16	1.0000	3.861E-24	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.853E-22	0.0000
Total	1.750E-16	1.0000	3.861E-24	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.853E-22	0.0000

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 3.000E+02 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.750E-16	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.750E-16	1.0000

*Sum of all water independent and dependent pathways.

Summary : Co 60, guardsman, 10000m2 File: C060G10K.RAD

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+03 years
Water Independent Pathways (Inhalation excludes radon)

Radio-	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
Nuclide	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)

As mrem/yr and Fraction of Total Dose At t = 1.000E+03 years
Water Dependent Pathways

Radio-	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
Nuclide	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000

*Sum of all water independent and dependent pathways.

Summary : Co 60, guardsmen, 10000m2 File: CO60G10K.RAD

Dose/Source Ratios Summed Over All Pathways
Parent and Progeny Principal Radionuclide Contributions Indicated

Parent (i)	Product (j)	Branch Fraction*	DSR(j,t) (mrem/yr)/(pCi/g) t= 0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	5.000E+01	1.000E+02	3.000E+02	1.000E+03
Co-60	Co-60	1.000E+00	2.500E-01	2.192E-01	1.684E-01	6.701E-02	4.813E-03	3.458E-04	4.782E-07	1.750E-18	0.000E+00

*Branch Fraction is the cumulative factor for the j't principal radionuclide daughter: CUMBRF(j) = BRF(1)*BRF(2)* ... BRF(j).
The DSR includes contributions from associated (half-life <= 0.5 yr) daughters.

Single Radionuclide Soil Guidelines G(i,t) in pCi/g

Basic Radiation Dose Limit = 25 mrem/yr

Nuclide (i)	t= 0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	5.000E+01	1.000E+02	3.000E+02	1.000E+03
Co-60	1.000E+02	1.141E+02	1.484E+02	3.731E+02	5.194E+03	7.230E+04	5.227E+07	*1.131E+15	*1.131E+15

*At specific activity limit

Summed Dose/Source Ratios DSR(i,t) in (mrem/yr)/(pCi/g)

and Single Radionuclide Soil Guidelines G(i,t) in pCi/g
at tmin = time of minimum single radionuclide soil guideline
and at tmax = time of maximum total dose = 0.000E+00 years

Nuclide (i)	Initial pCi/g	tmin (years)	DSR(i,tmin)	G(i,tmin) (pCi/g)	DSR(i,tmax)	G(i,tmax) (pCi/g)
Co-60	1.000E+02	0.000E+00	2.500E-01	1.000E+02	2.500E-01	1.000E+02

Individual Nuclide Dose Summed Over All Pathways
Parent Nuclide and Branch Fraction Indicated

Nuclide (j)	Parent (i)	BRF(i)	DOSE(j,t), mrem/yr								
			t= 0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	5.000E+01	1.000E+02	3.000E+02	1.000E+03
Co-60	Co-60	1.000E+00	2.500E+01	2.192E+01	1.684E+01	6.701E+00	4.813E-01	3.458E-02	4.782E-05	1.750E-16	0.000E+00

BRF(i) is the branch fraction of the parent nuclide.

Individual Nuclide Soil Concentration
Parent Nuclide and Branch Fraction Indicated

Nuclide (j)	Parent (i)	BRF(i)	S(j,t), pCi/g								
			t= 0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	5.000E+01	1.000E+02	3.000E+02	1.000E+03
Co-60	Co-60	1.000E+00	1.000E+02	8.766E+01	6.737E+01	2.680E+01	1.925E+00	1.383E-01	1.913E-04	7.000E-16	0.000E+00

BRF(i) is the branch fraction of the parent nuclide.

A.1.2 Cs-137: Guardsman Summary Output File

Summary : Cs137, guardsmen, 10000m2 File: C137G10K.RAD

Table of Contents

Part I: Mixture Sums and Single Radionuclide Guidelines

Dose Conversion Factor (and Related) Parameter Summary ... 2
Site-Specific Parameter Summary 3
Summary of Pathway Selections 6
Contaminated Zone and Total Dose Summary 7
Total Dose Components
Time = 0.000E+00 8
Time = 1.000E+00 9
Time = 3.000E+00 10
Time = 1.000E+01 11
Time = 3.000E+01 12
Time = 5.000E+01 13
Time = 1.000E+02 14
Time = 3.000E+02 15
Time = 1.000E+03 16
Dose/Source Ratios Summed Over All Pathways 17
Single Radionuclide Soil Guidelines 17
Dose Per Nuclide Summed Over All Pathways 18
Soil Concentration Per Nuclide 18

Dose Conversion Factor (and Related) Parameter Summary
File: DOSFAC.BIN

Menu	Parameter	Current Value	Default	Parameter Name
B-1	Dose conversion factors for inhalation, mrem/pCi:			
B-1	Cs-137+D	3.190E-05	3.190E-05	DCF2(1)
D-1	Dose conversion factors for ingestion, mrem/pCi:			
D-1	Cs-137+D	5.000E-05	5.000E-05	DCF3(1)
D-34	Food transfer factors:			
D-34	Cs-137+D , plant/soil concentration ratio, dimensionless	4.000E-02	4.000E-02	RTF(1,1)
D-34	Cs-137+D , beef/livestock-intake ratio, (pCi/kg)/(pCi/d)	3.000E-02	3.000E-02	RTF(1,2)
D-34	Cs-137+D , milk/livestock-intake ratio, (pCi/L)/(pCi/d)	8.000E-03	8.000E-03	RTF(1,3)
D-5	Bioaccumulation factors, fresh water, L/kg:			
D-5	Cs-137+D , fish	2.000E+03	2.000E+03	BIOFAC(1,1)
D-5	Cs-137+D , crustacea and mollusks	1.000E+02	1.000E+02	BIOFAC(1,2)

Site-Specific Parameter Summary

Menu	Parameter	User Input	Default	Used by RESRAD (If different from user input)	Parameter Name
R011	Area of contaminated zone (m**2)	1.000E+04	1.000E+04	---	AREA
R011	Thickness of contaminated zone (m)	2.000E+00	2.000E+00	---	THICKD
R011	Length parallel to aquifer flow (m)	not used	1.000E+02	---	LCZPAQ
R011	Basic radiation dose limit (mrem/yr)	2.500E+01	3.000E+01	---	BRDL
R011	Time since placement of material (yr)	0.000E+00	0.000E+00	---	TI
R011	Times for calculations (yr)	1.000E+00	1.000E+00	---	T(2)
R011	Times for calculations (yr)	3.000E+00	3.000E+00	---	T(3)
R011	Times for calculations (yr)	1.000E+01	1.000E+01	---	T(4)
R011	Times for calculations (yr)	3.000E+01	3.000E+01	---	T(5)
R011	Times for calculations (yr)	5.000E+01	1.000E+02	---	T(6)
R011	Times for calculations (yr)	1.000E+02	3.000E+02	---	T(7)
R011	Times for calculations (yr)	3.000E+02	1.000E+03	---	T(8)
R011	Times for calculations (yr)	1.000E+03	0.000E+00	---	T(9)
R011	Times for calculations (yr)	not used	0.000E+00	---	T(10)
R012	Initial principal radionuclide (pCi/g): Cs-137	4.775E+02	0.000E+00	---	S1(1)
R012	Concentration in groundwater (pCi/L): Cs-137	not used	0.000E+00	---	W1(1)
R013	Cover depth (m)	0.000E+00	0.000E+00	---	COVER0
R013	Density of cover material (g/cm**3)	not used	1.500E+00	---	DENSCV
R013	Cover depth erosion rate (m/yr)	not used	1.000E-03	---	VCV
R013	Density of contaminated zone (g/cm**3)	1.500E+00	1.500E+00	---	DENSCZ
R013	Contaminated zone erosion rate (m/yr)	0.000E+00	1.000E-03	---	VCZ
R013	Contaminated zone total porosity	4.000E-01	4.000E-01	---	TPCZ
R013	Contaminated zone effective porosity	2.000E-01	2.000E-01	---	EPCZ
R013	Contaminated zone hydraulic conductivity (m/yr)	1.000E+01	1.000E+01	---	HCCZ
R013	Contaminated zone b parameter	5.300E+00	5.300E+00	---	BCZ
R013	Average annual wind speed (m/sec)	2.000E+00	2.000E+00	---	WIND
R013	Humidity in air (g/m**3)	not used	8.000E+00	---	HUMID
R013	Evapotranspiration coefficient	5.000E-01	5.000E-01	---	EVAPTR
R013	Precipitation (m/yr)	1.000E+00	1.000E+00	---	PRECIP
R013	Irrigation (m/yr)	2.000E-01	2.000E-01	---	RI
R013	Irrigation mode	overhead	overhead	---	IDITCH
R013	Runoff coefficient	2.000E-01	2.000E-01	---	RUNOFF
R013	Watershed area for nearby stream or pond (m**2)	not used	1.000E+06	---	WAREA
R013	Accuracy for water/soil computations	not used	1.000E-03	---	EPS
R014	Density of saturated zone (g/cm**3)	1.500E+00	1.500E+00	---	DENSAQ
R014	Saturated zone total porosity	4.000E-01	4.000E-01	---	TPSZ
R014	Saturated zone effective porosity	2.000E-01	2.000E-01	---	EPSZ
R014	Saturated zone hydraulic conductivity (m/yr)	1.000E+02	1.000E+02	---	HCSZ
R014	Saturated zone hydraulic gradient	2.000E-02	2.000E-02	---	HGWT
R014	Saturated zone b parameter	5.300E+00	5.300E+00	---	BSZ
R014	Water table drop rate (m/yr)	1.000E-03	1.000E-03	---	VWT
R014	Well pump intake depth (m below water table)	1.000E+01	1.000E+01	---	DWIBWT
R014	Model: Nondispersion (ND) or Mass-Balance (MB)	ND	ND	---	MODEL
R014	Well pumping rate (m**3/yr)	2.500E+02	2.500E+02	---	UW
R015	Number of unsaturated zone strata	not used	1	---	NS

Site-Specific Parameter Summary (continued)

Menu	Parameter	User Input	Default	Used by RESRAD (If different from user input)	Parameter Name
R015	Unsat. zone 1, thickness (m)	not used	4.000E+00	---	H(1)
R015	Unsat. zone 1, soil density (g/cm**3)	not used	1.500E+00	---	DENSUZ(1)
R015	Unsat. zone 1, total porosity	not used	4.000E-01	---	TPUZ(1)
R015	Unsat. zone 1, effective porosity	not used	2.000E-01	---	EPUZ(1)
R015	Unsat. zone 1, soil-specific b parameter	not used	5.300E+00	---	BUZ(1)
R015	Unsat. zone 1, hydraulic conductivity (m/yr)	not used	1.000E+01	---	HCUZ(1)
R016	Distribution coefficients for Cs-137				
R016	Contaminated zone (cm**3/g)	1.000E+03	1.000E+03	---	DCNUCC(1)
R016	Unsaturated zone 1 (cm**3/g)	not used	1.000E+03	---	DCNUCU(1,1)
R016	Saturated zone (cm**3/g)	not used	1.000E+03	---	DCNUCS(1)
R016	Leach rate (/yr)	0.000E+00	0.000E+00	1.666E-04	ALEACH(1)
R016	Solubility constant	0.000E+00	0.000E+00	not used	SOLUBK(1)
R017	Inhalation rate (m**3/yr)	9.072E+01	8.400E+03	---	INHALR
R017	Mass loading for inhalation (g/m**3)	1.000E-04	1.000E-04	---	MLINH
R017	Exposure duration	2.000E+01	3.000E+01	---	ED
R017	Shielding factor, inhalation	4.000E-01	4.000E-01	---	SHF3
R017	Shielding factor, external gamma	7.000E-01	7.000E-01	---	SHF1
R017	Fraction of time spent indoors	0.000E+00	5.000E-01	---	FIND
R017	Fraction of time spent outdoors (on site)	1.640E-02	2.500E-01	---	FOTD
R017	Shape factor flag, external gamma	1.000E+00	1.000E+00	>0 shows circular AREA.	FS
R017	Radii of shape factor array (used if FS = -1):				
R017	Outer annular radius (m), ring 1:	not used	5.000E+01	---	RAD_SHAPE(1)
R017	Outer annular radius (m), ring 2:	not used	7.071E+01	---	RAD_SHAPE(2)
R017	Outer annular radius (m), ring 3:	not used	0.000E+00	---	RAD_SHAPE(3)
R017	Outer annular radius (m), ring 4:	not used	0.000E+00	---	RAD_SHAPE(4)
R017	Outer annular radius (m), ring 5:	not used	0.000E+00	---	RAD_SHAPE(5)
R017	Outer annular radius (m), ring 6:	not used	0.000E+00	---	RAD_SHAPE(6)
R017	Outer annular radius (m), ring 7:	not used	0.000E+00	---	RAD_SHAPE(7)
R017	Outer annular radius (m), ring 8:	not used	0.000E+00	---	RAD_SHAPE(8)
R017	Outer annular radius (m), ring 9:	not used	0.000E+00	---	RAD_SHAPE(9)
R017	Outer annular radius (m), ring 10:	not used	0.000E+00	---	RAD_SHAPE(10)
R017	Outer annular radius (m), ring 11:	not used	0.000E+00	---	RAD_SHAPE(11)
R017	Outer annular radius (m), ring 12:	not used	0.000E+00	---	RAD_SHAPE(12)
R017	Fractions of annular areas within AREA:				
R017	Ring 1	not used	1.000E+00	---	FRACA(1)
R017	Ring 2	not used	2.732E-01	---	FRACA(2)
R017	Ring 3	not used	0.000E+00	---	FRACA(3)
R017	Ring 4	not used	0.000E+00	---	FRACA(4)
R017	Ring 5	not used	0.000E+00	---	FRACA(5)
R017	Ring 6	not used	0.000E+00	---	FRACA(6)
R017	Ring 7	not used	0.000E+00	---	FRACA(7)
R017	Ring 8	not used	0.000E+00	---	FRACA(8)
R017	Ring 9	not used	0.000E+00	---	FRACA(9)
R017	Ring 10	not used	0.000E+00	---	FRACA(10)
R017	Ring 11	not used	0.000E+00	---	FRACA(11)
R017	Ring 12	not used	0.000E+00	---	FRACA(12)
R018	Fruits, vegetables and grain consumption (kg/yr)	not used	1.600E+02	---	DIET(1)

Site-Specific Parameter Summary (continued)

Menu	Parameter	User Input	Default	Used by RESRAD (If different from user input)	Parameter Name
R018	Leafy vegetable consumption (kg/yr)	not used	1.400E+01	---	DIET(2)
R018	Milk consumption (L/yr)	not used	9.200E+01	---	DIET(3)
R018	Meat and poultry consumption (kg/yr)	not used	6.300E+01	---	DIET(4)
R018	Fish consumption (kg/yr)	not used	5.400E+00	---	DIET(5)
R018	Other seafood consumption (kg/yr)	not used	9.000E-01	---	DIET(6)
R018	Soil ingestion rate (g/yr)	6.000E-01	3.650E+01	---	SOIL
R018	Drinking water intake (L/yr)	not used	5.100E+02	---	DWI
R018	Contamination fraction of drinking water	not used	1.000E+00	---	FDW
R018	Contamination fraction of household water	not used	1.000E+00	---	FHHW
R018	Contamination fraction of livestock water	not used	1.000E+00	---	FLW
R018	Contamination fraction of irrigation water	not used	1.000E+00	---	FIRW
R018	Contamination fraction of aquatic food	not used	5.000E-01	---	FR9
R018	Contamination fraction of plant food	not used	-1	---	FPLANT
R018	Contamination fraction of meat	not used	-1	---	FMEAT
R018	Contamination fraction of milk	not used	-1	---	FMILK
R019	Livestock fodder intake for meat (kg/day)	not used	6.800E+01	---	LFI5
R019	Livestock fodder intake for milk (kg/day)	not used	5.500E+01	---	LFI6
R019	Livestock water intake for meat (L/day)	not used	5.000E+01	---	LWI5
R019	Livestock water intake for milk (L/day)	not used	1.600E+02	---	LWI6
R019	Livestock soil intake (kg/day)	not used	5.000E-01	---	LSI
R019	Mass loading for foliar deposition (g/m**3)	not used	1.000E-04	---	MLFD
R019	Depth of soil mixing layer (m)	1.500E-01	1.500E-01	---	DM
R019	Depth of roots (m)	not used	9.000E-01	---	DROOT
R019	Drinking water fraction from ground water	1.000E+00	1.000E+00	---	FGWDW
R019	Household water fraction from ground water	not used	1.000E+00	---	FGWHH
R019	Livestock water fraction from ground water	not used	1.000E+00	---	FGWLW
R019	Irrigation fraction from ground water	not used	1.000E+00	---	FGWIR
R19B	Wet weight crop yield for Non-Leafy (kg/m**2)	not used	7.000E-01	---	YV(1)
R19B	Wet weight crop yield for Leafy (kg/m**2)	not used	1.500E+00	---	YV(2)
R19B	Wet weight crop yield for Fodder (kg/m**2)	not used	1.100E+00	---	YV(3)
R19B	Growing Season for Non-Leafy (years)	not used	1.700E-01	---	TE(1)
R19B	Growing Season for Leafy (years)	not used	2.500E-01	---	TE(2)
R19B	Growing Season for Fodder (years)	not used	8.000E-02	---	TE(3)
R19B	Translocation Factor for Non-Leafy	not used	1.000E-01	---	TIV(1)
R19B	Translocation Factor for Leafy	not used	1.000E+00	---	TIV(2)
R19B	Translocation Factor for Fodder	not used	1.000E+00	---	TIV(3)
R19B	Dry Foliar Interception Fraction for Non-Leafy	not used	2.500E-01	---	RDRY(1)
R19B	Dry Foliar Interception Fraction for Leafy	not used	2.500E-01	---	RDRY(2)
R19B	Dry Foliar Interception Fraction for Fodder	not used	2.500E-01	---	RDRY(3)
R19B	Wet Foliar Interception Fraction for Non-Leafy	not used	2.500E-01	---	RWET(1)
R19B	Wet Foliar Interception Fraction for Leafy	not used	2.500E-01	---	RWET(2)
R19B	Wet Foliar Interception Fraction for Fodder	not used	2.500E-01	---	RWET(3)
R19B	Weathering Removal Constant for Vegetation	not used	2.000E+01	---	WLAM
C14	C-12 concentration in water (g/cm**3)	not used	2.000E-05	---	C12WTR
C14	C-12 concentration in contaminated soil (g/g)	not used	3.000E-02	---	C12CZ
C14	Fraction of vegetation carbon from soil	not used	2.000E-02	---	CSOIL
C14	Fraction of vegetation carbon from air	not used	9.800E-01	---	CAIR
C14	C-14 evasion layer thickness in soil (m)	not used	3.000E-01	---	DMC

Site-Specific Parameter Summary (continued)

Menu	Parameter	User Input	Default	Used by RESRAD (if different from user input)	Parameter Name
C14	C-14 evasion flux rate from soil (1/sec)	not used	7.000E-07	---	EVSN
C14	C-12 evasion flux rate from soil (1/sec)	not used	1.000E-10	---	REVSN
C14	Fraction of grain in beef cattle feed	not used	8.000E-01	---	AVFG4
C14	Fraction of grain in milk cow feed	not used	2.000E-01	---	AVFG5
STOR	Storage times of contaminated foodstuffs (days):				
STOR	Fruits, non-leafy vegetables, and grain	1.400E+01	1.400E+01	---	STOR_T(1)
STOR	Leafy vegetables	1.000E+00	1.000E+00	---	STOR_T(2)
STOR	Milk	1.000E+00	1.000E+00	---	STOR_T(3)
STOR	Meat and poultry	2.000E+01	2.000E+01	---	STOR_T(4)
STOR	Fish	7.000E+00	7.000E+00	---	STOR_T(5)
STOR	Crustacea and mollusks	7.000E+00	7.000E+00	---	STOR_T(6)
STOR	Well water	1.000E+00	1.000E+00	---	STOR_T(7)
STOR	Surface water	1.000E+00	1.000E+00	---	STOR_T(8)
STOR	Livestock fodder	4.500E+01	4.500E+01	---	STOR_T(9)
R021	Thickness of building foundation (m)	not used	1.500E-01	---	FLOOR
R021	Bulk density of building foundation (g/cm**3)	not used	2.400E+00	---	DENSFL
R021	Total porosity of the cover material	not used	4.000E-01	---	TPCV
R021	Total porosity of the building foundation	not used	1.000E-01	---	TPFL
R021	Volumetric water content of the cover material	not used	5.000E-02	---	PH2OCV
R021	Volumetric water content of the foundation	not used	3.000E-02	---	PH2OFL
R021	Diffusion coefficient for radon gas (m/sec):				
R021	in cover material	not used	2.000E-06	---	DIFCV
R021	in foundation material	not used	3.000E-07	---	DIFFL
R021	in contaminated zone soil	not used	2.000E-06	---	DIFCZ
R021	Radon vertical dimension of mixing (m)	not used	2.000E+00	---	HMIX
R021	Average building air exchange rate (1/hr)	not used	5.000E-01	---	REXG
R021	Height of the building (room) (m)	not used	2.500E+00	---	HRM
R021	Building interior area factor	not used	0.000E+00	---	FAI
R021	Building depth below ground surface (m)	not used	-1.000E+00	---	DMFL
R021	Emanating power of Rn-222 gas	not used	2.500E-01	---	EMANA(1)
R021	Emanating power of Rn-220 gas	not used	1.500E-01	---	EMANA(2)

Summary of Pathway Selections

Pathway	User Selection
1 -- external gamma	active
2 -- inhalation (w/o radon)	active
3 -- plant ingestion	suppressed
4 -- meat ingestion	suppressed
5 -- milk ingestion	suppressed
6 -- aquatic foods	suppressed
7 -- drinking water	suppressed
8 -- soil ingestion	active
9 -- radon	suppressed
Find peak pathway doses	suppressed

Contaminated Zone Dimensions		Initial Soil Concentrations, pCi/g	
Area:	10000.00 square meters	Cs-137	4.77E+02
Thickness:	2.00 meters		
Cover Depth:	0.00 meters		

Total Dose TDOSE(t), mrem/yr
Basic Radiation Dose Limit = 25 mrem/yr
Total Mixture Sum M(t) = Fraction of Basic Dose Limit Received at Time (t)

t (years):	0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	5.000E+01	1.000E+02	3.000E+02	1.000E+03
TDOSE(t):	2.500E+01	2.442E+01	2.331E+01	1.981E+01	1.244E+01	7.808E+00	2.439E+00	2.322E-02	1.955E-09
M(t):	9.999E-01	9.769E-01	9.325E-01	7.923E-01	4.975E-01	3.123E-01	9.756E-02	9.289E-04	7.821E-11

Maximum TDOSE(t): 2.500E+01 mrem/yr at t = 0.000E+00 years

Summary : Cs137, guardsman, 10000m2 File: C137G10K.RAD

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 0.000E+00 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	2.500E+01	1.0000	3.837E-07	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.350E-04	0.0000
Total	2.500E+01	1.0000	3.837E-07	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.350E-04	0.0000

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)

As mrem/yr and Fraction of Total Dose At t = 0.000E+00 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.500E+01	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.500E+01	1.0000

*Sum of all water independent and dependent pathways.

Summary : Cs137, guardsman, 10000m2 File: C137G10K.RAD

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+00 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	2.442E+01	1.0000	3.749E-07	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.295E-04	0.0000
Total	2.442E+01	1.0000	3.749E-07	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.295E-04	0.0000

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+00 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.442E+01	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.442E+01	1.0000

*Sum of all water independent and dependent pathways.

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 3.000E+00 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	2.331E+01	1.0000	3.578E-07	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.191E-04	0.0000
Total	2.331E+01	1.0000	3.578E-07	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.191E-04	0.0000

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 3.000E+00 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.331E+01	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.331E+01	1.0000

*Sum of all water independent and dependent pathways.

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+01 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	1.981E+01	1.0000	3.040E-07	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.862E-04	0.0000
Total	1.981E+01	1.0000	3.040E-07	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.862E-04	0.0000

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+01 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.981E+01	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.981E+01	1.0000

*Sum of all water independent and dependent pathways.

Summary : Cs137, guardsman, 10000m2 File: C137G10K.RAD

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 3.000E+01 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	1.244E+01	1.0000	1.909E-07	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.169E-04	0.0000
Total	1.244E+01	1.0000	1.909E-07	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.169E-04	0.0000

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)

As mrem/yr and Fraction of Total Dose At t = 3.000E+01 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.244E+01	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.244E+01	1.0000

*Sum of all water independent and dependent pathways.

Summary : Cs137, guardsman, 10000m2 File: C137G10K.RAD

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 5.000E+01 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	7.808E+00	1.0000	1.199E-07	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	7.339E-05	0.0000
Total	7.808E+00	1.0000	1.199E-07	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	7.339E-05	0.0000

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 5.000E+01 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	7.808E+00	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	7.808E+00	1.0000

*Sum of all water independent and dependent pathways.

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+02 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	2.439E+00	1.0000	3.744E-08	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.293E-05	0.0000
Total	2.439E+00	1.0000	3.744E-08	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.293E-05	0.0000

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+02 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.439E+00	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.439E+00	1.0000

*Sum of all water independent and dependent pathways.

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 3.000E+02 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	2.322E-02	1.0000	3.565E-10	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.183E-07	0.0000
Total	2.322E-02	1.0000	3.565E-10	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.183E-07	0.0000

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)

As mrem/yr and Fraction of Total Dose At t = 3.000E+02 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.322E-02	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.322E-02	1.0000

*Sum of all water independent and dependent pathways.

Summary : Cs137, guardsman, 10000m2 File: C137G10K.RAD

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+03 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	1.955E-09	1.0000	3.001E-17	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.838E-14	0.0000
Total	1.955E-09	1.0000	3.001E-17	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.838E-14	0.0000

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)

As mrem/yr and Fraction of Total Dose At t = 1.000E+03 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.955E-09	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.955E-09	1.0000

*Sum of all water independent and dependent pathways.

Dose/Source Ratios Summed Over All Pathways
Parent and Progeny Principal Radionuclide Contributions Indicated

Parent (i)	Product (j)	Branch Fraction*	DSR(j,t) (mrem/yr)/(pCi/g) t= 0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	5.000E+01	1.000E+02	3.000E+02	1.000E+03
Cs-137	Cs-137	1.000E+00	5.235E-02	5.114E-02	4.882E-02	4.148E-02	2.604E-02	1.635E-02	5.108E-03	4.863E-05	4.094E-12

*Branch Fraction is the cumulative factor for the j't principal radionuclide daughter: CUMBRF(j) = BRF(1)*BRF(2)* ... BRF(j).
The DSR includes contributions from associated (half-life <= 0.5 yr) daughters.

Single Radionuclide Soil Guidelines G(i,t) in pCi/g

Basic Radiation Dose Limit = 25 mrem/yr

Nuclide (i)	t= 0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	5.000E+01	1.000E+02	3.000E+02	1.000E+03
Cs-137	4.776E+02	4.888E+02	5.121E+02	6.027E+02	9.600E+02	1.529E+03	4.895E+03	5.141E+05	6.106E+12

Summed Dose/Source Ratios DSR(i,t) in (mrem/yr)/(pCi/g)

and Single Radionuclide Soil Guidelines G(i,t) in pCi/g

at tmin = time of minimum single radionuclide soil guideline
and at tmax = time of maximum total dose = 0.000E+00 years

Nuclide (i)	Initial pCi/g	tmin (years)	DSR(i,tmin) (pCi/g)	G(i,tmin) (pCi/g)	DSR(i,tmax)	G(i,tmax) (pCi/g)
Cs-137	4.775E+02	0.000E+00	5.235E-02	4.776E+02	5.235E-02	4.776E+02

Individual Nuclide Dose Summed Over All Pathways
Parent Nuclide and Branch Fraction Indicated

Nuclide (j)	Parent (i)	BRF(i)	DOSE(j,t), mrem/yr								
		t=	0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	5.000E+01	1.000E+02	3.000E+02	1.000E+03
Cs-137	Cs-137	1.000E+00	2.500E+01	2.442E+01	2.331E+01	1.981E+01	1.244E+01	7.808E+00	2.439E+00	2.322E-02	1.955E-09

BRF(i) is the branch fraction of the parent nuclide.

Individual Nuclide Soil Concentration
Parent Nuclide and Branch Fraction Indicated

Nuclide (j)	Parent (i)	BRF(i)	S(j,t), pCi/g								
		t=	0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	5.000E+01	1.000E+02	3.000E+02	1.000E+03
Cs-137	Cs-137	1.000E+00	4.775E+02	4.666E+02	4.453E+02	3.784E+02	2.376E+02	1.492E+02	4.660E+01	4.436E-01	3.735E-08

BRF(i) is the branch fraction of the parent nuclide.

A.2 Results of RESRAD Calculations for Resident Adult Scenario

A.2.1 Co-60: Resident Adult Summary Output File

RESRAD, Version 5.82 T½ Limit = 0.5 year 12/14/98 17:38 Page 1

Summary : Co60, resident adult, 10000m2 File: CO60R10K.RAD

Table of Contents

Part I: Mixture Sums and Single Radionuclide Guidelines

Dose Conversion Factor (and Related) Parameter Summary ... 2
Site-Specific Parameter Summary ... 3
Summary of Pathway Selections ... 6
Contaminated Zone and Total Dose Summary ... 7
Total Dose Components
Time = 0.000E+00 ... 8
Time = 1.000E+00 ... 9
Time = 3.000E+00 ... 10
Time = 1.000E+01 ... 11
Time = 3.000E+01 ... 12
Time = 5.000E+01 ... 13
Time = 1.000E+02 ... 14
Time = 3.000E+02 ... 15
Time = 1.000E+03 ... 16
Dose/Source Ratios Summed Over All Pathways ... 17
Single Radionuclide Soil Guidelines ... 17
Dose Per Nuclide Summed Over All Pathways ... 18
Soil Concentration Per Nuclide ... 18

Dose Conversion Factor (and Related) Parameter Summary
File: DOSFAC.BIN

Menu	Parameter	Current Value	Default	Parameter Name
B-1	Dose conversion factors for inhalation, mrem/pCi:			
B-1	Co-60	2.190E-04	2.190E-04	DCF2(1)
D-1	Dose conversion factors for ingestion, mrem/pCi:			
D-1	Co-60	2.690E-05	2.690E-05	DCF3(1)
D-34	Food transfer factors:			
D-34	Co-60 , plant/soil concentration ratio, dimensionless	8.000E-02	8.000E-02	RTF(1,1)
D-34	Co-60 , beef/livestock-intake ratio, (pCi/kg)/(pCi/d)	2.000E-02	2.000E-02	RTF(1,2)
D-34	Co-60 , milk/livestock-intake ratio, (pCi/L)/(pCi/d)	2.000E-03	2.000E-03	RTF(1,3)
D-5	Bioaccumulation factors, fresh water, L/kg:			
D-5	Co-60 , fish	3.000E+02	3.000E+02	BIOFAC(1,1)
D-5	Co-60 , crustacea and mollusks	2.000E+02	2.000E+02	BIOFAC(1,2)

Site-Specific Parameter Summary

Menu	Parameter	User Input	Default	Used by RESRAD (If different from user input)	Parameter Name
R011	Area of contaminated zone (m**2)	1.000E+04	1.000E+04	---	AREA
R011	Thickness of contaminated zone (m)	2.000E+00	2.000E+00	---	THICKD
R011	Length parallel to aquifer flow (m)	1.000E+02	1.000E+02	---	LCZPAQ
R011	Basic radiation dose limit (mrem/yr)	2.500E+01	3.000E+01	---	BRDL
R011	Time since placement of material (yr)	0.000E+00	0.000E+00	---	TI
R011	Times for calculations (yr)	1.000E+00	1.000E+00	---	T(2)
R011	Times for calculations (yr)	3.000E+00	3.000E+00	---	T(3)
R011	Times for calculations (yr)	1.000E+01	1.000E+01	---	T(4)
R011	Times for calculations (yr)	3.000E+01	3.000E+01	---	T(5)
R011	Times for calculations (yr)	5.000E+01	1.000E+02	---	T(6)
R011	Times for calculations (yr)	1.000E+02	3.000E+02	---	T(7)
R011	Times for calculations (yr)	3.000E+02	1.000E+03	---	T(8)
R011	Times for calculations (yr)	1.000E+03	0.000E+00	---	T(9)
R011	Times for calculations (yr)	not used	0.000E+00	---	T(10)
R012	Initial principal radionuclide (pCi/g): Co-60	1.542E+02	0.000E+00	---	S1(1)
R012	Concentration in groundwater (pCi/L): Co-60	not used	0.000E+00	---	W1(1)
R013	Cover depth (m)	0.000E+00	0.000E+00	---	COVER0
R013	Density of cover material (g/cm**3)	not used	1.500E+00	---	DENSCV
R013	Cover depth erosion rate (m/yr)	not used	1.000E-03	---	VCV
R013	Density of contaminated zone (g/cm**3)	1.500E+00	1.500E+00	---	DENSCZ
R013	Contaminated zone erosion rate (m/yr)	0.000E+00	1.000E-03	---	VCZ
R013	Contaminated zone total porosity	4.000E-01	4.000E-01	---	TPCZ
R013	Contaminated zone effective porosity	2.000E-01	2.000E-01	---	EPCZ
R013	Contaminated zone hydraulic conductivity (m/yr)	1.000E+01	1.000E+01	---	HCCZ
R013	Contaminated zone b parameter	5.300E+00	5.300E+00	---	BCZ
R013	Average annual wind speed (m/sec)	2.000E+00	2.000E+00	---	WIND
R013	Humidity in air (g/m**3)	not used	8.000E+00	---	HUMID
R013	Evapotranspiration coefficient	5.000E-01	5.000E-01	---	EVAPTR
R013	Precipitation (m/yr)	1.000E+00	1.000E+00	---	PRECIP
R013	Irrigation (m/yr)	2.000E-01	2.000E-01	---	RI
R013	Irrigation mode	overhead	overhead	---	IDITCH
R013	Runoff coefficient	2.000E-01	2.000E-01	---	RUNOFF
R013	Watershed area for nearby stream or pond (m**2)	1.000E+06	1.000E+06	---	WAREA
R013	Accuracy for water/soil computations	1.000E-03	1.000E-03	---	EPS
R014	Density of saturated zone (g/cm**3)	1.500E+00	1.500E+00	---	DENSAQ
R014	Saturated zone total porosity	4.000E-01	4.000E-01	---	TPSZ
R014	Saturated zone effective porosity	2.000E-01	2.000E-01	---	EPSZ
R014	Saturated zone hydraulic conductivity (m/yr)	1.000E+02	1.000E+02	---	HCSZ
R014	Saturated zone hydraulic gradient	2.000E-02	2.000E-02	---	HGWT
R014	Saturated zone b parameter	5.300E+00	5.300E+00	---	BSZ
R014	Water table drop rate (m/yr)	1.000E-03	1.000E-03	---	VWT
R014	Well pump intake depth (m below water table)	1.000E+01	1.000E+01	---	DWIBWT
R014	Model: Nondispersion (ND) or Mass-Balance (MB)	ND	ND	---	MODEL
R014	Well pumping rate (m**3/yr)	2.500E+02	2.500E+02	---	UW
R015	Number of unsaturated zone strata	1	1	---	NS

Site-Specific Parameter Summary (continued)

Menu	Parameter	User Input	Default	Used by RESRAD (If different from user input)	Parameter Name
R015	Unsat. zone 1, thickness (m)	4.000E+00	4.000E+00	---	H(1)
R015	Unsat. zone 1, soil density (g/cm**3)	1.500E+00	1.500E+00	---	DENSUZ(1)
R015	Unsat. zone 1, total porosity	4.000E-01	4.000E-01	---	TPUZ(1)
R015	Unsat. zone 1, effective porosity	2.000E-01	2.000E-01	---	EPUZ(1)
R015	Unsat. zone 1, soil-specific b parameter	5.300E+00	5.300E+00	---	BUZ(1)
R015	Unsat. zone 1, hydraulic conductivity (m/yr)	1.000E+01	1.000E+01	---	HCUZ(1)
R016	Distribution coefficients for Co-60				
R016	Contaminated zone (cm**3/g)	1.000E+03	1.000E+03	---	DCNUCC(1)
R016	Unsaturated zone 1 (cm**3/g)	1.000E+03	1.000E+03	---	DCNUCU(1,1)
R016	Saturated zone (cm**3/g)	1.000E+03	1.000E+03	---	DCNUCS(1)
R016	Leach rate (/yr)	0.000E+00	0.000E+00	1.666E-04	ALEACH(1)
R016	Solubility constant	0.000E+00	0.000E+00	not used	SOLUBK(1)
R017	Inhalation rate (m**3/yr)	4.057E+03	8.400E+03	---	INHALR
R017	Mass loading for inhalation (g/m**3)	1.000E-04	1.000E-04	---	MLINH
R017	Exposure duration	3.000E+01	3.000E+01	---	ED
R017	Shielding factor, inhalation	4.000E-01	4.000E-01	---	SHF3
R017	Shielding factor, external gamma	7.000E-01	7.000E-01	---	SHF1
R017	Fraction of time spent indoors	6.550E-01	5.000E-01	---	FIND
R017	Fraction of time spent outdoors (on site)	8.000E-02	2.500E-01	---	FOTD
R017	Shape factor flag, external gamma	1.000E+00	1.000E+00	>0 shows circular AREA.	FS
R017	Radii of shape factor array (used if FS = -1):				
R017	Outer annular radius (m), ring 1:	not used	5.000E+01	---	RAD_SHAPE(1)
R017	Outer annular radius (m), ring 2:	not used	7.071E+01	---	RAD_SHAPE(2)
R017	Outer annular radius (m), ring 3:	not used	0.000E+00	---	RAD_SHAPE(3)
R017	Outer annular radius (m), ring 4:	not used	0.000E+00	---	RAD_SHAPE(4)
R017	Outer annular radius (m), ring 5:	not used	0.000E+00	---	RAD_SHAPE(5)
R017	Outer annular radius (m), ring 6:	not used	0.000E+00	---	RAD_SHAPE(6)
R017	Outer annular radius (m), ring 7:	not used	0.000E+00	---	RAD_SHAPE(7)
R017	Outer annular radius (m), ring 8:	not used	0.000E+00	---	RAD_SHAPE(8)
R017	Outer annular radius (m), ring 9:	not used	0.000E+00	---	RAD_SHAPE(9)
R017	Outer annular radius (m), ring 10:	not used	0.000E+00	---	RAD_SHAPE(10)
R017	Outer annular radius (m), ring 11:	not used	0.000E+00	---	RAD_SHAPE(11)
R017	Outer annular radius (m), ring 12:	not used	0.000E+00	---	RAD_SHAPE(12)
R017	Fractions of annular areas within AREA:				
R017	Ring 1	not used	1.000E+00	---	FRACA(1)
R017	Ring 2	not used	2.732E-01	---	FRACA(2)
R017	Ring 3	not used	0.000E+00	---	FRACA(3)
R017	Ring 4	not used	0.000E+00	---	FRACA(4)
R017	Ring 5	not used	0.000E+00	---	FRACA(5)
R017	Ring 6	not used	0.000E+00	---	FRACA(6)
R017	Ring 7	not used	0.000E+00	---	FRACA(7)
R017	Ring 8	not used	0.000E+00	---	FRACA(8)
R017	Ring 9	not used	0.000E+00	---	FRACA(9)
R017	Ring 10	not used	0.000E+00	---	FRACA(10)
R017	Ring 11	not used	0.000E+00	---	FRACA(11)
R017	Ring 12	not used	0.000E+00	---	FRACA(12)
R018	Fruits, vegetables and grain consumption (kg/yr)	1.936E+02	1.600E+02	---	DIET(1)

Site-Specific Parameter Summary (continued)

Menu	Parameter	User Input	Default	Used by RESRAD (If different from user input)	Parameter Name
R018	Leafy vegetable consumption (kg/yr)	0.000E+00	1.400E+01	---	DIET(2)
R018	Milk consumption (L/yr)	not used	9.200E+01	---	DIET(3)
R018	Meat and poultry consumption (kg/yr)	not used	6.300E+01	---	DIET(4)
R018	Fish consumption (kg/yr)	not used	5.400E+00	---	DIET(5)
R018	Other seafood consumption (kg/yr)	not used	9.000E-01	---	DIET(6)
R018	Soil ingestion rate (g/yr)	1.750E+01	3.650E+01	---	SOIL
R018	Drinking water intake (L/yr)	8.400E+02	5.100E+02	---	DWI
R018	Contamination fraction of drinking water	1.000E+00	1.000E+00	---	FDW
R018	Contamination fraction of household water	not used	1.000E+00	---	FHHW
R018	Contamination fraction of livestock water	not used	1.000E+00	---	FLW
R018	Contamination fraction of irrigation water	1.000E+00	1.000E+00	---	FIRW
R018	Contamination fraction of aquatic food	not used	5.000E-01	---	FR9
R018	Contamination fraction of plant food	-1	-1	0.500E+00	FPLANT
R018	Contamination fraction of meat	not used	-1	---	FMEAT
R018	Contamination fraction of milk	not used	-1	---	FMILK
R019	Livestock fodder intake for meat (kg/day)	not used	6.800E+01	---	LFI5
R019	Livestock fodder intake for milk (kg/day)	not used	5.500E+01	---	LFI6
R019	Livestock water intake for meat (L/day)	not used	5.000E+01	---	LWI5
R019	Livestock water intake for milk (L/day)	not used	1.600E+02	---	LWI6
R019	Livestock soil intake (kg/day)	not used	5.000E-01	---	LSI
R019	Mass loading for foliar deposition (g/m**3)	1.000E-04	1.000E-04	---	MLFD
R019	Depth of soil mixing layer (m)	1.500E-01	1.500E-01	---	DM
R019	Depth of roots (m)	9.000E-01	9.000E-01	---	DROOT
R019	Drinking water fraction from ground water	1.000E+00	1.000E+00	---	FGWDW
R019	Household water fraction from ground water	1.000E+00	1.000E+00	---	FGWHH
R019	Livestock water fraction from ground water	not used	1.000E+00	---	FGWLW
R019	Irrigation fraction from ground water	not used	1.000E+00	---	FGWIR
R19B	Wet weight crop yield for Non-Leafy (kg/m**2)	7.000E-01	7.000E-01	---	YV(1)
R19B	Wet weight crop yield for Leafy (kg/m**2)	1.500E+00	1.500E+00	---	YV(2)
R19B	Wet weight crop yield for Fodder (kg/m**2)	not used	1.100E+00	---	YV(3)
R19B	Growing Season for Non-Leafy (years)	1.700E-01	1.700E-01	---	TE(1)
R19B	Growing Season for Leafy (years)	2.500E-01	2.500E-01	---	TE(2)
R19B	Growing Season for Fodder (years)	not used	8.000E-02	---	TE(3)
R19B	Translocation Factor for Non-Leafy	1.000E-01	1.000E-01	---	TIV(1)
R19B	Translocation Factor for Leafy	1.000E+00	1.000E+00	---	TIV(2)
R19B	Translocation Factor for Fodder	not used	1.000E+00	---	TIV(3)
R19B	Dry Foliar Interception Fraction for Non-Leafy	2.500E-01	2.500E-01	---	RDRY(1)
R19B	Dry Foliar Interception Fraction for Leafy	2.500E-01	2.500E-01	---	RDRY(2)
R19B	Dry Foliar Interception Fraction for Fodder	not used	2.500E-01	---	RDRY(3)
R19B	Wet Foliar Interception Fraction for Non-Leafy	2.500E-01	2.500E-01	---	RWET(1)
R19B	Wet Foliar Interception Fraction for Leafy	2.500E-01	2.500E-01	---	RWET(2)
R19B	Wet Foliar Interception Fraction for Fodder	not used	2.500E-01	---	RWET(3)
R19B	Weathering Removal Constant for Vegetation	2.000E+01	2.000E+01	---	WLAM
C14	C-12 concentration in water (g/cm**3)	not used	2.000E-05	---	C12WTR
C14	C-12 concentration in contaminated soil (g/g)	not used	3.000E-02	---	C12CZ
C14	Fraction of vegetation carbon from soil	not used	2.000E-02	---	CSOIL
C14	Fraction of vegetation carbon from air	not used	9.800E-01	---	CAIR
C14	C-14 evasion layer thickness in soil (m)	not used	3.000E-01	---	DMC

Site-Specific Parameter Summary (continued)

Menu	Parameter	User Input	Default	Used by RESRAD (If different from user input)	Parameter Name
C14	C-14 evasion flux rate from soil (1/sec)	not used	7.000E-07	---	EVSN
C14	C-12 evasion flux rate from soil (1/sec)	not used	1.000E-10	---	REVSN
C14	Fraction of grain in beef cattle feed	not used	8.000E-01	---	AVFG4
C14	Fraction of grain in milk cow feed	not used	2.000E-01	---	AVFG5
STOR	Storage times of contaminated foodstuffs (days):				
STOR	Fruits, non-leafy vegetables, and grain	1.400E+01	1.400E+01	---	STOR_T(1)
STOR	Leafy vegetables	1.000E+00	1.000E+00	---	STOR_T(2)
STOR	Milk	1.000E+00	1.000E+00	---	STOR_T(3)
STOR	Meat and poultry	2.000E+01	2.000E+01	---	STOR_T(4)
STOR	Fish	7.000E+00	7.000E+00	---	STOR_T(5)
STOR	Crustacea and mollusks	7.000E+00	7.000E+00	---	STOR_T(6)
STOR	Well water	1.000E+00	1.000E+00	---	STOR_T(7)
STOR	Surface water	1.000E+00	1.000E+00	---	STOR_T(8)
STOR	Livestock fodder	4.500E+01	4.500E+01	---	STOR_T(9)
R021	Thickness of building foundation (m)	not used	1.500E-01	---	FLOOR
R021	Bulk density of building foundation (g/cm**3)	not used	2.400E+00	---	DENSFL
R021	Total porosity of the cover material	not used	4.000E-01	---	TPCV
R021	Total porosity of the building foundation	not used	1.000E-01	---	TPFL
R021	Volumetric water content of the cover material	not used	5.000E-02	---	PH2OCV
R021	Volumetric water content of the foundation	not used	3.000E-02	---	PH2OFL
R021	Diffusion coefficient for radon gas (m/sec):				
R021	in cover material	not used	2.000E-06	---	DIFCV
R021	in foundation material	not used	3.000E-07	---	DIFFL
R021	in contaminated zone soil	not used	2.000E-06	---	DIFCZ
R021	Radon vertical dimension of mixing (m)	not used	2.000E+00	---	HMIX
R021	Average building air exchange rate (1/hr)	not used	5.000E-01	---	REXG
R021	Height of the building (room) (m)	not used	2.500E+00	---	HRM
R021	Building interior area factor	not used	0.000E+00	---	FAI
R021	Building depth below ground surface (m)	not used	-1.000E+00	---	DMFL
R021	Emanating power of Rn-222 gas	not used	2.500E-01	---	EMANA(1)
R021	Emanating power of Rn-220 gas	not used	1.500E-01	---	EMANA(2)

Summary of Pathway Selections

Pathway	User Selection
1 -- external gamma	active
2 -- inhalation (w/o radon)	active
3 -- plant ingestion	active
4 -- meat ingestion	suppressed
5 -- milk ingestion	suppressed
6 -- aquatic foods	suppressed
7 -- drinking water	active
8 -- soil ingestion	active
9 -- radon	suppressed
Find peak pathway doses	suppressed

Contaminated Zone Dimensions		Initial Soil Concentrations, pCi/g	
Area:	10000.00 square meters	Co-60	1.542E+02
Thickness:	2.00 meters		
Cover Depth:	0.00 meters		

Total Dose TDOSE(t), mrem/yr
Basic Radiation Dose Limit = 25 mrem/yr
Total Mixture Sum M(t) = Fraction of Basic Dose Limit Received at Time (t)

t (years):	0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	5.000E+01	1.000E+02	3.000E+02	1.000E+03
TDOSE(t):	1.298E+03	1.138E+03	8.746E+02	3.480E+02	2.500E+01	1.796E+00	2.483E-03	9.088E-15	0.000E+00
M(t):	5.193E+01	4.552E+01	3.498E+01	1.392E+01	9.998E-01	7.182E-02	9.934E-05	3.635E-16	0.000E+00

Maximum TDOSE(t): 1.298E+03 mrem/yr at t = 0.000E+00 years

Summary : Co60, resident adult, 10000m2 File: C060R10K.RAD

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 0.000E+00 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	1.266E+03	0.9752	7.935E-04	0.0000	0.000E+00	0.0000	3.212E+01	0.0247	0.000E+00	0.0000	0.000E+00	0.0000	5.336E-02	0.0000
Total	1.266E+03	0.9752	7.935E-04	0.0000	0.000E+00	0.0000	3.212E+01	0.0247	0.000E+00	0.0000	0.000E+00	0.0000	5.336E-02	0.0000

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)

As mrem/yr and Fraction of Total Dose At t = 0.000E+00 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.298E+03	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.298E+03	1.0000

*Sum of all water independent and dependent pathways.

Summary : Co60, resident adult, 10000m2 File: C060R10K.RAD

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+00 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	1.110E+03	0.9752	6.956E-04	0.0000	0.000E+00	0.0000	2.816E+01	0.0247	0.000E+00	0.0000	0.000E+00	0.0000	4.678E-02	0.0000
Total	1.110E+03	0.9752	6.956E-04	0.0000	0.000E+00	0.0000	2.816E+01	0.0247	0.000E+00	0.0000	0.000E+00	0.0000	4.678E-02	0.0000

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+00 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.138E+03	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.138E+03	1.0000

*Sum of all water independent and dependent pathways.

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 3.000E+00 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	8.529E+02	0.9752	5.345E-04	0.0000	0.000E+00	0.0000	2.164E+01	0.0247	0.000E+00	0.0000	0.000E+00	0.0000	3.595E-02	0.0000
Total	8.529E+02	0.9752	5.345E-04	0.0000	0.000E+00	0.0000	2.164E+01	0.0247	0.000E+00	0.0000	0.000E+00	0.0000	3.595E-02	0.0000

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 3.000E+00 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	8.746E+02	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	8.746E+02	1.0000

*Sum of all water independent and dependent pathways.

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+01 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	3.393E+02	0.9752	2.127E-04	0.0000	0.000E+00	0.0000	8.609E+00	0.0247	0.000E+00	0.0000	0.000E+00	0.0000	1.430E-02	0.0000
Total	3.393E+02	0.9752	2.127E-04	0.0000	0.000E+00	0.0000	8.609E+00	0.0247	0.000E+00	0.0000	0.000E+00	0.0000	1.430E-02	0.0000

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)

As mrem/yr and Fraction of Total Dose At t = 1.000E+01 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	3.480E+02	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	3.480E+02	1.0000

*Sum of all water independent and dependent pathways.

Summary : Co60, resident adult, 10000m2 File: CO60R10K.RAD

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 3.000E+01 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	2.438E+01	0.9752	1.528E-05	0.0000	0.000E+00	0.0000	6.184E-01	0.0247	0.000E+00	0.0000	0.000E+00	0.0000	1.027E-03	0.0000
Total	2.438E+01	0.9752	1.528E-05	0.0000	0.000E+00	0.0000	6.184E-01	0.0247	0.000E+00	0.0000	0.000E+00	0.0000	1.027E-03	0.0000

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)

As mrem/yr and Fraction of Total Dose At t = 3.000E+01 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.500E+01	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.500E+01	1.0000

*Sum of all water independent and dependent pathways.

Summary : Co60, resident adult, 10000m2 File: CO60R10K.RAD

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 5.000E+01 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	1.751E+00	0.9752	1.097E-06	0.0000	0.000E+00	0.0000	4.443E-02	0.0247	0.000E+00	0.0000	0.000E+00	0.0000	7.381E-05	0.0000
Total	1.751E+00	0.9752	1.097E-06	0.0000	0.000E+00	0.0000	4.443E-02	0.0247	0.000E+00	0.0000	0.000E+00	0.0000	7.381E-05	0.0000

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 5.000E+01 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.796E+00	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.796E+00	1.0000

*Sum of all water independent and dependent pathways.

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+02 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	2.422E-03	0.9752	1.518E-09	0.0000	0.000E+00	0.0000	6.144E-05	0.0247	0.000E+00	0.0000	0.000E+00	0.0000	1.021E-07	0.0000
Total	2.422E-03	0.9752	1.518E-09	0.0000	0.000E+00	0.0000	6.144E-05	0.0247	0.000E+00	0.0000	0.000E+00	0.0000	1.021E-07	0.0000

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+02 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.483E-03	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.483E-03	1.0000

*Sum of all water independent and dependent pathways.

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 3.000E+02 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	8.863E-15	0.9752	5.554E-21	0.0000	0.000E+00	0.0000	2.248E-16	0.0247	0.000E+00	0.0000	0.000E+00	0.0000	3.736E-19	0.0000
Total	8.863E-15	0.9752	5.554E-21	0.0000	0.000E+00	0.0000	2.248E-16	0.0247	0.000E+00	0.0000	0.000E+00	0.0000	3.736E-19	0.0000

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)

As mrem/yr and Fraction of Total Dose At t = 3.000E+02 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	9.088E-15	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	9.088E-15	1.0000

*Sum of all water independent and dependent pathways.

Summary : Co60, resident adult, 10000m2 File: C060R10K.RAD

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+03 years
Water Independent Pathways (Inhalation excludes radon)

Radio-	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
Nuclide	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)

As mrem/yr and Fraction of Total Dose At t = 1.000E+03 years
Water Dependent Pathways

Radio-	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
Nuclide	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000

*Sum of all water independent and dependent pathways.

Summary : Co60, resident adult, 10000m2 File: CO60R10K.RAD

Dose/Source Ratios Summed Over All Pathways
Parent and Progeny Principal Radionuclide Contributions Indicated

Parent (i)	Product (j)	Branch Fraction*	DSR(j,t) (mrem/yr)/(pCi/g) t= 0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	5.000E+01	1.000E+02	3.000E+02	1.000E+03
Co-60	Co-60	1.000E+00	8.418E+00	7.379E+00	5.671E+00	2.256E+00	1.621E-01	1.164E-02	1.610E-05	5.892E-17	0.000E+00

*Branch Fraction is the cumulative factor for the j't principal radionuclide daughter: CUMBRF(j) = BRF(1)*BRF(2)* ... BRF(j).
The DSR includes contributions from associated (half-life <= 0.5 yr) daughters.

Single Radionuclide Soil Guidelines G(i,t) in pCi/g

Basic Radiation Dose Limit = 25 mrem/yr

Nuclide (i)	t= 0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	5.000E+01	1.000E+02	3.000E+02	1.000E+03
Co-60	2.970E+00	3.388E+00	4.409E+00	1.108E+01	1.543E+02	2.147E+03	1.553E+06	*1.131E+15	*1.131E+15

*At specific activity limit

Summed Dose/Source Ratios DSR(i,t) in (mrem/yr)/(pCi/g)

and Single Radionuclide Soil Guidelines G(i,t) in pCi/g
at tmin = time of minimum single radionuclide soil guideline
and at tmax = time of maximum total dose = 0.000E+00 years

Nuclide (i)	Initial pCi/g	tmin (years)	DSR(i,tmin)	G(i,tmin) (pCi/g)	DSR(i,tmax)	G(i,tmax) (pCi/g)
Co-60	1.542E+02	0.000E+00	8.418E+00	2.970E+00	8.418E+00	2.970E+00

Individual Nuclide Dose Summed Over All Pathways
Parent Nuclide and Branch Fraction Indicated

Nuclide (j)	Parent (i)	BRF(i)	DOSE(j,t), mrem/yr t= 0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	5.000E+01	1.000E+02	3.000E+02	1.000E+03
Co-60	Co-60	1.000E+00	1.298E+03	1.138E+03	8.746E+02	3.480E+02	2.500E+01	1.796E+00	2.483E-03	9.088E-15	0.000E+00

BRF(i) is the branch fraction of the parent nuclide.

Individual Nuclide Soil Concentration
Parent Nuclide and Branch Fraction Indicated

Nuclide (j)	Parent (i)	BRF(i)	S(j,t), pCi/g t= 0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	5.000E+01	1.000E+02	3.000E+02	1.000E+03
Co-60	Co-60	1.000E+00	1.542E+02	1.352E+02	1.039E+02	4.134E+01	2.969E+00	2.133E-01	2.950E-04	1.080E-15	0.000E+00

BRF(i) is the branch fraction of the parent nuclide.

A.2.2 Cs-137: Resident Adult Summary Output File

Summary : Cs137, resident, 10000m2 File: C137R10K.RAD

Table of Contents

Part I: Mixture Sums and Single Radionuclide Guidelines

Dose Conversion Factor (and Related) Parameter Summary ... 2
Site-Specific Parameter Summary ... 3
Summary of Pathway Selections ... 6
Contaminated Zone and Total Dose Summary ... 7
Total Dose Components
Time = 0.000E+00 ... 8
Time = 1.000E+00 ... 9
Time = 3.000E+00 ... 10
Time = 1.000E+01 ... 11
Time = 3.000E+01 ... 12
Time = 5.000E+01 ... 13
Time = 1.000E+02 ... 14
Time = 3.000E+02 ... 15
Time = 1.000E+03 ... 16
Dose/Source Ratios Summed Over All Pathways ... 17
Single Radionuclide Soil Guidelines ... 17
Dose Per Nuclide Summed Over All Pathways ... 18
Soil Concentration Per Nuclide ... 18

Dose Conversion Factor (and Related) Parameter Summary
File: DOSFAC.BIN

Menu	Parameter	Current Value	Default	Parameter Name
B-1	Dose conversion factors for inhalation, mrem/pCi:			
B-1	Cs-137+D	3.190E-05	3.190E-05	DCF2(1)
D-1	Dose conversion factors for ingestion, mrem/pCi:			
D-1	Cs-137+D	5.000E-05	5.000E-05	DCF3(1)
D-34	Food transfer factors:			
D-34	Cs-137+D , plant/soil concentration ratio, dimensionless	4.000E-02	4.000E-02	RTF(1,1)
D-34	Cs-137+D , beef/livestock-intake ratio, (pCi/kg)/(pCi/d)	3.000E-02	3.000E-02	RTF(1,2)
D-34	Cs-137+D , milk/livestock-intake ratio, (pCi/L)/(pCi/d)	8.000E-03	8.000E-03	RTF(1,3)
D-5	Bioaccumulation factors, fresh water, L/kg:			
D-5	Cs-137+D , fish	2.000E+03	2.000E+03	BIOFAC(1,1)
D-5	Cs-137+D , crustacea and mollusks	1.000E+02	1.000E+02	BIOFAC(1,2)

Site-Specific Parameter Summary

Menu	Parameter	User Input	Default	Used by RESRAD (If different from user input)	Parameter Name
R011	Area of contaminated zone (m**2)	1.000E+04	1.000E+04	---	AREA
R011	Thickness of contaminated zone (m)	2.000E+00	2.000E+00	---	THICKD
R011	Length parallel to aquifer flow (m)	1.000E+02	1.000E+02	---	LCZPAQ
R011	Basic radiation dose limit (mrem/yr)	2.500E+01	3.000E+01	---	BRDL
R011	Time since placement of material (yr)	0.000E+00	0.000E+00	---	TI
R011	Times for calculations (yr)	1.000E+00	1.000E+00	---	T(2)
R011	Times for calculations (yr)	3.000E+00	3.000E+00	---	T(3)
R011	Times for calculations (yr)	1.000E+01	1.000E+01	---	T(4)
R011	Times for calculations (yr)	3.000E+01	3.000E+01	---	T(5)
R011	Times for calculations (yr)	5.000E+01	1.000E+02	---	T(6)
R011	Times for calculations (yr)	1.000E+02	3.000E+02	---	T(7)
R011	Times for calculations (yr)	3.000E+02	1.000E+03	---	T(8)
R011	Times for calculations (yr)	1.000E+03	0.000E+00	---	T(9)
R011	Times for calculations (yr)	not used	0.000E+00	---	T(10)
R012	Initial principal radionuclide (pCi/g): Cs-137	2.627E+01	0.000E+00	---	S1(1)
R012	Concentration in groundwater (pCi/L): Cs-137	not used	0.000E+00	---	W1(1)
R013	Cover depth (m)	0.000E+00	0.000E+00	---	COVER0
R013	Density of cover material (g/cm**3)	not used	1.500E+00	---	DENSCV
R013	Cover depth erosion rate (m/yr)	not used	1.000E-03	---	VCV
R013	Density of contaminated zone (g/cm**3)	1.500E+00	1.500E+00	---	DENSCZ
R013	Contaminated zone erosion rate (m/yr)	0.000E+00	1.000E-03	---	VCZ
R013	Contaminated zone total porosity	4.000E-01	4.000E-01	---	TPCZ
R013	Contaminated zone effective porosity	2.000E-01	2.000E-01	---	EPCZ
R013	Contaminated zone hydraulic conductivity (m/yr)	1.000E+01	1.000E+01	---	HCCZ
R013	Contaminated zone b parameter	5.300E+00	5.300E+00	---	BCZ
R013	Average annual wind speed (m/sec)	2.000E+00	2.000E+00	---	WIND
R013	Humidity in air (g/m**3)	not used	8.000E+00	---	HUMID
R013	Evapotranspiration coefficient	5.000E-01	5.000E-01	---	EVAPTR
R013	Precipitation (m/yr)	1.000E+00	1.000E+00	---	PRECIP
R013	Irrigation (m/yr)	2.000E-01	2.000E-01	---	RI
R013	Irrigation mode	overhead	overhead	---	IDITCH
R013	Runoff coefficient	2.000E-01	2.000E-01	---	RUNOFF
R013	Watershed area for nearby stream or pond (m**2)	1.000E+06	1.000E+06	---	WAREA
R013	Accuracy for water/soil computations	1.000E-03	1.000E-03	---	EPS
R014	Density of saturated zone (g/cm**3)	1.500E+00	1.500E+00	---	DENSAQ
R014	Saturated zone total porosity	4.000E-01	4.000E-01	---	TPSZ
R014	Saturated zone effective porosity	2.000E-01	2.000E-01	---	EPSZ
R014	Saturated zone hydraulic conductivity (m/yr)	1.000E+02	1.000E+02	---	HCSZ
R014	Saturated zone hydraulic gradient	2.000E-02	2.000E-02	---	HGWT
R014	Saturated zone b parameter	5.300E+00	5.300E+00	---	BSZ
R014	Water table drop rate (m/yr)	1.000E-03	1.000E-03	---	VWT
R014	Well pump intake depth (m below water table)	1.000E+01	1.000E+01	---	DWIBWT
R014	Model: Nondispersion (ND) or Mass-Balance (MB)	ND	ND	---	MODEL
R014	Well pumping rate (m**3/yr)	2.500E+02	2.500E+02	---	UW
R015	Number of unsaturated zone strata	1	1	---	NS

Site-Specific Parameter Summary (continued)

Menu	Parameter	User Input	Default	Used by RESRAD (If different from user input)	Parameter Name
R015	Unsat. zone 1, thickness (m)	4.000E+00	4.000E+00	---	H(1)
R015	Unsat. zone 1, soil density (g/cm**3)	1.500E+00	1.500E+00	---	DENSUZ(1)
R015	Unsat. zone 1, total porosity	4.000E-01	4.000E-01	---	TPUZ(1)
R015	Unsat. zone 1, effective porosity	2.000E-01	2.000E-01	---	EPUZ(1)
R015	Unsat. zone 1, soil-specific b parameter	5.300E+00	5.300E+00	---	BUZ(1)
R015	Unsat. zone 1, hydraulic conductivity (m/yr)	1.000E+01	1.000E+01	---	HCUZ(1)
R016	Distribution coefficients for Cs-137				
R016	Contaminated zone (cm**3/g)	1.000E+03	1.000E+03	---	DCNUCC(1)
R016	Unsaturated zone 1 (cm**3/g)	1.000E+03	1.000E+03	---	DCNUCU(1,1)
R016	Saturated zone (cm**3/g)	1.000E+03	1.000E+03	---	DCNUCS(1)
R016	Leach rate (/yr)	0.000E+00	0.000E+00	1.666E-04	ALEACH(1)
R016	Solubility constant	0.000E+00	0.000E+00	not used	SOLUBK(1)
R017	Inhalation rate (m**3/yr)	4.057E+03	8.400E+03	---	INHALR
R017	Mass loading for inhalation (g/m**3)	1.000E-04	1.000E-04	---	MLINH
R017	Exposure duration	3.000E+01	3.000E+01	---	ED
R017	Shielding factor, inhalation	4.000E-01	4.000E-01	---	SHF3
R017	Shielding factor, external gamma	7.000E-01	7.000E-01	---	SHF1
R017	Fraction of time spent indoors	6.550E-01	5.000E-01	---	FIND
R017	Fraction of time spent outdoors (on site)	8.000E-02	2.500E-01	---	FOTD
R017	Shape factor flag, external gamma	1.000E+00	1.000E+00	>0 shows circular AREA.	FS
R017	Radii of shape factor array (used if FS = -1):				
R017	Outer annular radius (m), ring 1:	not used	5.000E+01	---	RAD_SHAPE(1)
R017	Outer annular radius (m), ring 2:	not used	7.071E+01	---	RAD_SHAPE(2)
R017	Outer annular radius (m), ring 3:	not used	0.000E+00	---	RAD_SHAPE(3)
R017	Outer annular radius (m), ring 4:	not used	0.000E+00	---	RAD_SHAPE(4)
R017	Outer annular radius (m), ring 5:	not used	0.000E+00	---	RAD_SHAPE(5)
R017	Outer annular radius (m), ring 6:	not used	0.000E+00	---	RAD_SHAPE(6)
R017	Outer annular radius (m), ring 7:	not used	0.000E+00	---	RAD_SHAPE(7)
R017	Outer annular radius (m), ring 8:	not used	0.000E+00	---	RAD_SHAPE(8)
R017	Outer annular radius (m), ring 9:	not used	0.000E+00	---	RAD_SHAPE(9)
R017	Outer annular radius (m), ring 10:	not used	0.000E+00	---	RAD_SHAPE(10)
R017	Outer annular radius (m), ring 11:	not used	0.000E+00	---	RAD_SHAPE(11)
R017	Outer annular radius (m), ring 12:	not used	0.000E+00	---	RAD_SHAPE(12)
R017	Fractions of annular areas within AREA:				
R017	Ring 1	not used	1.000E+00	---	FRACA(1)
R017	Ring 2	not used	2.732E-01	---	FRACA(2)
R017	Ring 3	not used	0.000E+00	---	FRACA(3)
R017	Ring 4	not used	0.000E+00	---	FRACA(4)
R017	Ring 5	not used	0.000E+00	---	FRACA(5)
R017	Ring 6	not used	0.000E+00	---	FRACA(6)
R017	Ring 7	not used	0.000E+00	---	FRACA(7)
R017	Ring 8	not used	0.000E+00	---	FRACA(8)
R017	Ring 9	not used	0.000E+00	---	FRACA(9)
R017	Ring 10	not used	0.000E+00	---	FRACA(10)
R017	Ring 11	not used	0.000E+00	---	FRACA(11)
R017	Ring 12	not used	0.000E+00	---	FRACA(12)
R018	Fruits, vegetables and grain consumption (kg/yr)	1.936E+02	1.600E+02	---	DIET(1)

Site-Specific Parameter Summary (continued)

Menu	Parameter	User Input	Default	Used by RESRAD (If different from user input)	Parameter Name
R018	Leafy vegetable consumption (kg/yr)	0.000E+00	1.400E+01	---	DIET(2)
R018	Milk consumption (L/yr)	not used	9.200E+01	---	DIET(3)
R018	Meat and poultry consumption (kg/yr)	not used	6.300E+01	---	DIET(4)
R018	Fish consumption (kg/yr)	not used	5.400E+00	---	DIET(5)
R018	Other seafood consumption (kg/yr)	not used	9.000E-01	---	DIET(6)
R018	Soil ingestion rate (g/yr)	1.750E+01	3.650E+01	---	SOIL
R018	Drinking water intake (L/yr)	8.400E+02	5.100E+02	---	DWI
R018	Contamination fraction of drinking water	1.000E+00	1.000E+00	---	FDW
R018	Contamination fraction of household water	not used	1.000E+00	---	FHHW
R018	Contamination fraction of livestock water	not used	1.000E+00	---	FLW
R018	Contamination fraction of irrigation water	1.000E+00	1.000E+00	---	FIRW
R018	Contamination fraction of aquatic food	not used	5.000E-01	---	FR9
R018	Contamination fraction of plant food	-1	-1	0.500E+00	FPLANT
R018	Contamination fraction of meat	not used	-1	---	FMEAT
R018	Contamination fraction of milk	not used	-1	---	FMILK
R019	Livestock fodder intake for meat (kg/day)	not used	6.800E+01	---	LFI5
R019	Livestock fodder intake for milk (kg/day)	not used	5.500E+01	---	LFI6
R019	Livestock water intake for meat (L/day)	not used	5.000E+01	---	LWI5
R019	Livestock water intake for milk (L/day)	not used	1.600E+02	---	LWI6
R019	Livestock soil intake (kg/day)	not used	5.000E-01	---	LSI
R019	Mass loading for foliar deposition (g/m**3)	1.000E-04	1.000E-04	---	MLFD
R019	Depth of soil mixing layer (m)	1.500E-01	1.500E-01	---	DM
R019	Depth of roots (m)	9.000E-01	9.000E-01	---	DROOT
R019	Drinking water fraction from ground water	1.000E+00	1.000E+00	---	FGWDW
R019	Household water fraction from ground water	1.000E+00	1.000E+00	---	FGWHH
R019	Livestock water fraction from ground water	not used	1.000E+00	---	FGWLW
R019	Irrigation fraction from ground water	not used	1.000E+00	---	FGWIR
R19B	Wet weight crop yield for Non-Leafy (kg/m**2)	7.000E-01	7.000E-01	---	YV(1)
R19B	Wet weight crop yield for Leafy (kg/m**2)	1.500E+00	1.500E+00	---	YV(2)
R19B	Wet weight crop yield for Fodder (kg/m**2)	not used	1.100E+00	---	YV(3)
R19B	Growing Season for Non-Leafy (years)	1.700E-01	1.700E-01	---	TE(1)
R19B	Growing Season for Leafy (years)	2.500E-01	2.500E-01	---	TE(2)
R19B	Growing Season for Fodder (years)	not used	8.000E-02	---	TE(3)
R19B	Translocation Factor for Non-Leafy	1.000E-01	1.000E-01	---	TIV(1)
R19B	Translocation Factor for Leafy	1.000E+00	1.000E+00	---	TIV(2)
R19B	Translocation Factor for Fodder	not used	1.000E+00	---	TIV(3)
R19B	Dry Foliar Interception Fraction for Non-Leafy	2.500E-01	2.500E-01	---	RDRY(1)
R19B	Dry Foliar Interception Fraction for Leafy	2.500E-01	2.500E-01	---	RDRY(2)
R19B	Dry Foliar Interception Fraction for Fodder	not used	2.500E-01	---	RDRY(3)
R19B	Wet Foliar Interception Fraction for Non-Leafy	2.500E-01	2.500E-01	---	RWET(1)
R19B	Wet Foliar Interception Fraction for Leafy	2.500E-01	2.500E-01	---	RWET(2)
R19B	Wet Foliar Interception Fraction for Fodder	not used	2.500E-01	---	RWET(3)
R19B	Weathering Removal Constant for Vegetation	2.000E+01	2.000E+01	---	WLAM
C14	C-12 concentration in water (g/cm**3)	not used	2.000E-05	---	C12WTR
C14	C-12 concentration in contaminated soil (g/g)	not used	3.000E-02	---	C12CZ
C14	Fraction of vegetation carbon from soil	not used	2.000E-02	---	CSOIL
C14	Fraction of vegetation carbon from air	not used	9.800E-01	---	CAIR
C14	C-14 evasion layer thickness in soil (m)	not used	3.000E-01	---	DMC

Site-Specific Parameter Summary (continued)

Menu	Parameter	User Input	Default	Used by RESRAD (If different from user input)	Parameter Name
C14	C-14 evasion flux rate from soil (1/sec)	not used	7.000E-07	---	EVSN
C14	C-12 evasion flux rate from soil (1/sec)	not used	1.000E-10	---	REVSN
C14	Fraction of grain in beef cattle feed	not used	8.000E-01	---	AVFG4
C14	Fraction of grain in milk cow feed	not used	2.000E-01	---	AVFG5
STOR	Storage times of contaminated foodstuffs (days):				
STOR	Fruits, non-leafy vegetables, and grain	1.400E+01	1.400E+01	---	STOR_T(1)
STOR	Leafy vegetables	1.000E+00	1.000E+00	---	STOR_T(2)
STOR	Milk	1.000E+00	1.000E+00	---	STOR_T(3)
STOR	Meat and poultry	2.000E+01	2.000E+01	---	STOR_T(4)
STOR	Fish	7.000E+00	7.000E+00	---	STOR_T(5)
STOR	Crustacea and mollusks	7.000E+00	7.000E+00	---	STOR_T(6)
STOR	Well water	1.000E+00	1.000E+00	---	STOR_T(7)
STOR	Surface water	1.000E+00	1.000E+00	---	STOR_T(8)
STOR	Livestock fodder	4.500E+01	4.500E+01	---	STOR_T(9)
R021	Thickness of building foundation (m)	not used	1.500E-01	---	FLOOR
R021	Bulk density of building foundation (g/cm**3)	not used	2.400E+00	---	DENSFL
R021	Total porosity of the cover material	not used	4.000E-01	---	TPCV
R021	Total porosity of the building foundation	not used	1.000E-01	---	TPFL
R021	Volumetric water content of the cover material	not used	5.000E-02	---	PH2OCV
R021	Volumetric water content of the foundation	not used	3.000E-02	---	PH2OFL
R021	Diffusion coefficient for radon gas (m/sec):				
R021	in cover material	not used	2.000E-06	---	DIFCV
R021	in foundation material	not used	3.000E-07	---	DIFFL
R021	in contaminated zone soil	not used	2.000E-06	---	DIFCZ
R021	Radon vertical dimension of mixing (m)	not used	2.000E+00	---	HMIX
R021	Average building air exchange rate (1/hr)	not used	5.000E-01	---	REXG
R021	Height of the building (room) (m)	not used	2.500E+00	---	HRM
R021	Building interior area factor	not used	0.000E+00	---	FAI
R021	Building depth below ground surface (m)	not used	-1.000E+00	---	DMFL
R021	Emanating power of Rn-222 gas	not used	2.500E-01	---	EMANA(1)
R021	Emanating power of Rn-220 gas	not used	1.500E-01	---	EMANA(2)

Summary of Pathway Selections

Pathway	User Selection
1 -- external gamma	active
2 -- inhalation (w/o radon)	active
3 -- plant ingestion	active
4 -- meat ingestion	suppressed
5 -- milk ingestion	suppressed
6 -- aquatic foods	suppressed
7 -- drinking water	active
8 -- soil ingestion	active
9 -- radon	suppressed
Find peak pathway doses	suppressed

Contaminated Zone Dimensions		Initial Soil Concentrations, pCi/g	
Area:	10000.00 square meters	Cs-137	2.627E+01
Thickness:	2.00 meters		
Cover Depth:	0.00 meters		

Total Dose TDOSE(t), mrem/yr
Basic Radiation Dose Limit = 25 mrem/yr
Total Mixture Sum M(t) = Fraction of Basic Dose Limit Received at Time (t)

t (years):	0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	5.000E+01	1.000E+02	3.000E+02	1.000E+03
TDOSE(t):	5.025E+01	4.910E+01	4.687E+01	3.982E+01	2.500E+01	1.570E+01	4.903E+00	4.668E-02	3.931E-09
M(t):	2.010E+00	1.964E+00	1.875E+00	1.593E+00	1.000E+00	6.279E-01	1.961E-01	1.867E-03	1.572E-10

Maximum TDOSE(t): 5.025E+01 mrem/yr at t = 0.000E+00 years

Summary : Cs137, resident, 10000m2 File: C137R10K.RAD

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 0.000E+00 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	4.515E+01	0.8985	1.969E-05	0.0000	0.000E+00	0.0000	5.085E+00	0.1012	0.000E+00	0.0000	0.000E+00	0.0000	1.689E-02	0.0003
Total	4.515E+01	0.8985	1.969E-05	0.0000	0.000E+00	0.0000	5.085E+00	0.1012	0.000E+00	0.0000	0.000E+00	0.0000	1.689E-02	0.0003

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)

As mrem/yr and Fraction of Total Dose At t = 0.000E+00 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	5.025E+01	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	5.025E+01	1.0000

*Sum of all water independent and dependent pathways.

Summary : Cs137, resident, 10000m2 File: C137R10K.RAD

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+00 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	4.411E+01	0.8985	1.923E-05	0.0000	0.000E+00	0.0000	4.968E+00	0.1012	0.000E+00	0.0000	0.000E+00	0.0000	1.651E-02	0.0003
Total	4.411E+01	0.8985	1.923E-05	0.0000	0.000E+00	0.0000	4.968E+00	0.1012	0.000E+00	0.0000	0.000E+00	0.0000	1.651E-02	0.0003

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+00 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	4.910E+01	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	4.910E+01	1.0000

*Sum of all water independent and dependent pathways.

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 3.000E+00 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	4.211E+01	0.8985	1.836E-05	0.0000	0.000E+00	0.0000	4.742E+00	0.1012	0.000E+00	0.0000	0.000E+00	0.0000	1.576E-02	0.0003
Total	4.211E+01	0.8985	1.836E-05	0.0000	0.000E+00	0.0000	4.742E+00	0.1012	0.000E+00	0.0000	0.000E+00	0.0000	1.576E-02	0.0003

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 3.000E+00 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	4.687E+01	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	4.687E+01	1.0000

*Sum of all water independent and dependent pathways.

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+01 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	3.578E+01	0.8985	1.560E-05	0.0000	0.000E+00	0.0000	4.029E+00	0.1012	0.000E+00	0.0000	0.000E+00	0.0000	1.339E-02	0.0003
Total	3.578E+01	0.8985	1.560E-05	0.0000	0.000E+00	0.0000	4.029E+00	0.1012	0.000E+00	0.0000	0.000E+00	0.0000	1.339E-02	0.0003

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)

As mrem/yr and Fraction of Total Dose At t = 1.000E+01 years
Water Dependant Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	3.982E+01	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	3.982E+01	1.0000

*Sum of all water independent and dependent pathways.

Summary : Cs137, resident, 10000m2 File: C137R10K.RAD

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 3.000E+01 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	2.246E+01	0.8985	9.794E-06	0.0000	0.000E+00	0.0000	2.530E+00	0.1012	0.000E+00	0.0000	0.000E+00	0.0000	8.405E-03	0.0003
Total	2.246E+01	0.8985	9.794E-06	0.0000	0.000E+00	0.0000	2.530E+00	0.1012	0.000E+00	0.0000	0.000E+00	0.0000	8.405E-03	0.0003

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)

As mrem/yr and Fraction of Total Dose At t = 3.000E+01 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.500E+01	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.500E+01	1.0000

*Sum of all water independent and dependent pathways.

Summary : Cs137, resident, 10000m2 File: C137R10K.RAD

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 5.000E+01 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	1.410E+01	0.8985	6.149E-06	0.0000	0.000E+00	0.0000	1.588E+00	0.1012	0.000E+00	0.0000	0.000E+00	0.0000	5.277E-03	0.0003
Total	1.410E+01	0.8985	6.149E-06	0.0000	0.000E+00	0.0000	1.588E+00	0.1012	0.000E+00	0.0000	0.000E+00	0.0000	5.277E-03	0.0003

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 5.000E+01 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.570E+01	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.570E+01	1.0000

*Sum of all water independent and dependent pathways.

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+02 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	4.406E+00	0.8985	1.921E-06	0.0000	0.000E+00	0.0000	4.961E-01	0.1012	0.000E+00	0.0000	0.000E+00	0.0000	1.648E-03	0.0003
Total	4.406E+00	0.8985	1.921E-06	0.0000	0.000E+00	0.0000	4.961E-01	0.1012	0.000E+00	0.0000	0.000E+00	0.0000	1.648E-03	0.0003

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+02 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	4.903E+00	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	4.903E+00	1.0000

*Sum of all water independent and dependent pathways.

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 3.000E+02 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	4.194E-02	0.8985	1.829E-08	0.0000	0.000E+00	0.0000	4.723E-03	0.1012	0.000E+00	0.0000	0.000E+00	0.0000	1.569E-05	0.0003
Total	4.194E-02	0.8985	1.829E-08	0.0000	0.000E+00	0.0000	4.723E-03	0.1012	0.000E+00	0.0000	0.000E+00	0.0000	1.569E-05	0.0003

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)

As mrem/yr and Fraction of Total Dose At t = 3.000E+02 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	4.668E-02	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	4.668E-02	1.0000

*Sum of all water independent and dependent pathways.

Summary : Cs137, resident, 10000m2 File: C137R10K.RAD

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+03 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	3.532E-09	0.8985	1.540E-15	0.0000	0.000E+00	0.0000	3.977E-10	0.1012	0.000E+00	0.0000	0.000E+00	0.0000	1.321E-12	0.0003
Total	3.532E-09	0.8985	1.540E-15	0.0000	0.000E+00	0.0000	3.977E-10	0.1012	0.000E+00	0.0000	0.000E+00	0.0000	1.321E-12	0.0003

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)

As mrem/yr and Fraction of Total Dose At t = 1.000E+03 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	3.931E-09	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	3.931E-09	1.0000

*Sum of all water independent and dependent pathways.

Dose/Source Ratios Summed Over All Pathways
Parent and Progeny Principal Radionuclide Contributions Indicated

Parent (i)	Product (j)	Branch Fraction*	DSR(j,t) (mrem/yr)/(pCi/g) t= 0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	5.000E+01	1.000E+02	3.000E+02	1.000E+03
Cs-137	Cs-137	1.000E+00	1.913E+00	1.869E+00	1.784E+00	1.516E+00	9.517E-01	5.976E-01	1.867E-01	1.777E-03	1.496E-10

*Branch Fraction is the cumulative factor for the j't principal radionuclide daughter: CUMBRF(j) = BRF(1)*BRF(2)* ... BRF(j).
The DSR includes contributions from associated (half life <= 0.5 yr) daughters.

Single Radionuclide Soil Guidelines G(i,t) in pCi/g

Basic Radiation Dose Limit = 25 mrem/yr

Nuclide (i)	t= 0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	5.000E+01	1.000E+02	3.000E+02	1.000E+03
Cs-137	1.307E+01	1.338E+01	1.401E+01	1.649E+01	2.627E+01	4.184E+01	1.339E+02	1.407E+04	1.671E+11

Summed Dose/Source Ratios DSR(i,t) in (mrem/yr)/(pCi/g)

and Single Radionuclide Soil Guidelines G(i,t) in pCi/g

at tmin = time of minimum single radionuclide soil guideline
and at tmax = time of maximum total dose = 0.000E+00 years

Nuclide (i)	Initial pCi/g	tmin (years)	DSR(i,tmin) (pCi/g)	G(i,tmin) (pCi/g)	DSR(i,tmax)	G(i,tmax) (pCi/g)
Cs-137	2.627E+01	0.000E+00	1.913E+00	1.307E+01	1.913E+00	1.307E+01

Individual Nuclide Dose Summed Over All Pathways
Parent Nuclide and Branch Fraction Indicated

Nuclide (j)	Parent (i)	BRF(i)	DOSE(j,t), mrem/yr t= 0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	5.000E+01	1.000E+02	3.000E+02	1.000E+03
Cs-137	Cs-137	1.000E+00	5.025E+01	4.910E+01	4.687E+01	3.982E+01	2.500E+01	1.570E+01	4.903E+00	4.668E-02	3.931E-09

BRF(i) is the branch fraction of the parent nuclide.

Individual Nuclide Soil Concentration
Parent Nuclide and Branch Fraction Indicated

Nuclide (j)	Parent (i)	BRF(i)	S(j,t), pCi/g t= 0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	5.000E+01	1.000E+02	3.000E+02	1.000E+03
Cs-137	Cs-137	1.000E+00	2.627E+01	2.567E+01	2.450E+01	2.082E+01	1.307E+01	8.206E+00	2.563E+00	2.440E-02	2.055E-09

BRF(i) is the branch fraction of the parent nuclide.

A.3 Results of RESRAD Calculations for Resident Child Scenario

A.3.1 Co-60: Resident Child Summary Output File

Summary : Co6C, resident child, 10000m2
File : C0600000.RAD

Table of Contents

Part I: Mixture Sums and Single Radionuclide Guidelines

Dose Conversion Factor (and Related) Parameter Summary ... 2
Site-Specific Parameter Summary 3
Summary of Pathway Selections 7
Contaminated Zone and Total Dose Summary 8
Total Dose Components
Time = 0.000E+00 9
Time = 1.000E+00 10
Time = 3.000E+00 11
Time = 1.000E+01 12
Time = 3.000E+01 13
Time = 5.000E+01 14
Time = 1.000E+02 15
Time = 3.000E+02 16
Time = 1.000E+03 17
Dose/Source Ratios Summed Over All Pathways 18
Single Radionuclide Soil Guidelines 18
Dose Per Nuclide Summed Over All Pathways 19
Soil Concentration Per Nuclide 19

Dose Conversion Factor (and Related) Parameter Summary
File: DOSFAC.BIN

Menu	Parameter	Current Value	Default	Parameter Name
B-1	Dose conversion factors for inhalation, mrem/pCi:			
B-1	Co-60	2.190E-04	2.190E-04	DCF2(1)
D-1	Dose conversion factors for ingestion, mrem/pCi:			
D-1	Co-60	2.690E-05	2.690E-05	DCF3(1)
D-34	Food transfer factors:			
D-34	Co-60 , plant/soil concentration ratio, dimensionless	8.000E-02	8.000E-02	RTF(1,1)
D-34	Co-60 , beef/livestock-intake ratio, (pCi/kg)/(pCi/d)	2.000E-02	2.000E-02	RTF(1,2)
D-34	Co-60 , milk/livestock-intake ratio, (pCi/L)/(pCi/d)	2.000E-03	2.000E-03	RTF(1,3)
D-5	Bioaccumulation factors, fresh water, L/kg:			
D-5	Co-60 , fish	3.000E+02	3.000E+02	BIOFAC(1,1)
D-5	Co-60 , crustacea and mollusks	2.000E+02	2.000E+02	BIOFAC(1,2)

Site-Specific Parameter Summary

Menu	Parameter	User Input	Default	Used by RESRAD (If different from user input)	Parameter Name
R011	Area of contaminated zone (m**2)	1.000E+04	1.000E+04	---	AREA
R011	Thickness of contaminated zone (m)	2.000E+00	2.000E+00	---	THICKD
R011	Length parallel to aquifer flow (m)	1.000E+02	1.000E+02	---	LCZPAQ
R011	Basic radiation dose limit (mrem/yr)	2.500E+01	3.000E+01	---	BRDL
R011	Time since placement of material (yr)	0.000E+00	0.000E+00	---	TI
R011	Times for calculations (yr)	1.000E+00	1.000E+00	---	T(2)
R011	Times for calculations (yr)	3.000E+00	3.000E+00	---	T(3)
R011	Times for calculations (yr)	1.000E+01	1.000E+01	---	T(4)
R011	Times for calculations (yr)	3.000E+01	3.000E+01	---	T(5)
R011	Times for calculations (yr)	5.000E+01	1.000E+02	---	T(6)
R011	Times for calculations (yr)	1.000E+02	3.000E+02	---	T(7)
R011	Times for calculations (yr)	3.000E+02	1.000E+03	---	T(8)
R011	Times for calculations (yr)	1.000E+03	0.000E+00	---	T(9)
R011	Times for calculations (yr)	not used	0.000E+00	---	T(10)
R012	Initial principal radionuclide (pCi/g): Co-60	1.000E+00	0.000E+00	---	S1(1)
R012	Concentration in groundwater (pCi/L): Co-60	not used	0.000E+00	---	W1(1)
R013	Cover depth (m)	0.000E+00	0.000E+00	---	COVER0
R013	Density of cover material (g/cm**3)	not used	1.500E+00	---	DENSCV
R013	Cover depth erosion rate (m/yr)	not used	1.000E-03	---	VCV
R013	Density of contaminated zone (g/cm**3)	1.500E+00	1.500E+00	---	DENSCZ
R013	Contaminated zone erosion rate (m/yr)	0.000E+00	1.000E-03	---	VCZ
R013	Contaminated zone total porosity	4.000E-01	4.000E-01	---	TPCZ
R013	Contaminated zone effective porosity	2.000E-01	2.000E-01	---	EPCZ
R013	Contaminated zone hydraulic conductivity (m/yr)	1.000E+01	1.000E+01	---	HCCZ
R013	Contaminated zone b parameter	5.300E+00	5.300E+00	---	BCZ
R013	Average annual wind speed (m/sec)	2.000E+00	2.000E+00	---	WIND
R013	Humidity in air (g/m**3)	not used	8.000E+00	---	HUMID
R013	Evapotranspiration coefficient	5.000E-01	5.000E-01	---	EVAPTR
R013	Precipitation (m/yr)	1.000E+00	1.000E+00	---	PRECIP
R013	Irrigation (m/yr)	2.000E-01	2.000E-01	---	RI
R013	Irrigation mode	overhead	overhead	---	IDITCH
R013	Runoff coefficient	2.000E-01	2.000E-01	---	RUNOFF
R013	Watershed area for nearby stream or pond (m**2)	1.000E+06	1.000E+06	---	WAREA
R013	Accuracy for water/soil computations	1.000E-03	1.000E-03	---	EPS
R014	Density of saturated zone (g/cm**3)	1.500E+00	1.500E+00	---	DENSAQ
R014	Saturated zone total porosity	4.000E-01	4.000E-01	---	TPSZ
R014	Saturated zone effective porosity	2.000E-01	2.000E-01	---	EPSZ
R014	Saturated zone hydraulic conductivity (m/yr)	1.000E+02	1.000E+02	---	HCSZ
R014	Saturated zone hydraulic gradient	2.000E-02	2.000E-02	---	HGWT
R014	Saturated zone b parameter	5.300E+00	5.300E+00	---	BSZ
R014	Water table drop rate (m/yr)	1.000E-03	1.000E-03	---	VWT
R014	Well pump intake depth (m below water table)	1.000E+01	1.000E+01	---	DWIBWT
R014	Model: Nondispersion (ND) or Mass-Balance (MB)	ND	ND	---	MODEL
R014	Well pumping rate (m**3/yr)	2.500E+02	2.500E+02	---	UW
R015	Number of unsaturated zone strata	1	1	---	NS

Site-Specific Parameter Summary (continued)

Menu	Parameter	User Input	Default	Used by RESRAD (If different from user input)	Parameter Name
R015	Unsat. zone 1, thickness (m)	4.000E+00	4.000E+00	---	H(1)
R015	Unsat. zone 1, soil density (g/cm**3)	1.500E+00	1.500E+00	---	DENSUZ(1)
R015	Unsat. zone 1, total porosity	4.000E-01	4.000E-01	---	TPUZ(1)
R015	Unsat. zone 1, effective porosity	2.000E-01	2.000E-01	---	EPUZ(1)
R015	Unsat. zone 1, soil-specific b parameter	5.300E+00	5.300E+00	---	BUZ(1)
R015	Unsat. zone 1, hydraulic conductivity (m/yr)	1.000E+01	1.000E+01	---	HCUZ(1)
R016	Distribution coefficients for Co-60				
R016	Contaminated zone (cm**3/g)	1.000E+03	1.000E+03	---	DCUCC(1)
R016	Unsaturated zone 1 (cm**3/g)	1.000E+03	1.000E+03	---	DCUCU(1,1)
R016	Saturated zone (cm**3/g)	1.000E+03	1.000E+03	---	DCUCS(1)
R016	Leach rate (/yr)	0.000E+00	0.000E+00	1.666E-04	ALEACH(1)
R016	Solubility constant	0.000E+00	0.000E+00	not used	SOLUBK(1)
R017	Inhalation rate (m**3/yr)	2.720E+03	8.400E+03	---	INHALR
R017	Mass loading for inhalation (g/m**3)	1.000E-04	1.000E-04	---	MLINH
R017	Exposure duration	6.000E+00	3.000E+01	---	ED
R017	Shielding factor, inhalation	4.000E-01	4.000E-01	---	SHF3
R017	Shielding factor, external gamma	7.000E-01	7.000E-01	---	SHF1
R017	Fraction of time spent indoors	8.390E-01	5.000E-01	---	FIND
R017	Fraction of time spent outdoors (on site)	8.990E-02	2.500E-01	---	FOTD
R017	Shape factor flag, external gamma	1.000E+00	1.000E+00	>0 shows circular AREA.	FS
R017	Radii of shape factor array (used if FS = -1):				
R017	Outer annular radius (m), ring 1:	not used	5.000E+01	---	RAD_SHAPE(1)
R017	Outer annular radius (m), ring 2:	not used	7.071E+01	---	RAD_SHAPE(2)
R017	Outer annular radius (m), ring 3:	not used	0.000E+00	---	RAD_SHAPE(3)
R017	Outer annular radius (m), ring 4:	not used	0.000E+00	---	RAD_SHAPE(4)
R017	Outer annular radius (m), ring 5:	not used	0.000E+00	---	RAD_SHAPE(5)
R017	Outer annular radius (m), ring 6:	not used	0.000E+00	---	RAD_SHAPE(6)
R017	Outer annular radius (m), ring 7:	not used	0.000E+00	---	RAD_SHAPE(7)
R017	Outer annular radius (m), ring 8:	not used	0.000E+00	---	RAD_SHAPE(8)
R017	Outer annular radius (m), ring 9:	not used	0.000E+00	---	RAD_SHAPE(9)
R017	Outer annular radius (m), ring 10:	not used	0.000E+00	---	RAD_SHAPE(10)
R017	Outer annular radius (m), ring 11:	not used	0.000E+00	---	RAD_SHAPE(11)
R017	Outer annular radius (m), ring 12:	not used	0.000E+00	---	RAD_SHAPE(12)
R017	Fractions of annular areas within AREA:				
R017	Ring 1	not used	1.000E+00	---	FRACA(1)
R017	Ring 2	not used	2.732E-01	---	FRACA(2)
R017	Ring 3	not used	0.000E+00	---	FRACA(3)
R017	Ring 4	not used	0.000E+00	---	FRACA(4)
R017	Ring 5	not used	0.000E+00	---	FRACA(5)
R017	Ring 6	not used	0.000E+00	---	FRACA(6)
R017	Ring 7	not used	0.000E+00	---	FRACA(7)
R017	Ring 8	not used	0.000E+00	---	FRACA(8)
R017	Ring 9	not used	0.000E+00	---	FRACA(9)
R017	Ring 10	not used	0.000E+00	---	FRACA(10)
R017	Ring 11	not used	0.000E+00	---	FRACA(11)
R017	Ring 12	not used	0.000E+00	---	FRACA(12)

Site-Specific Parameter Summary (continued)

Menu	Parameter	User Input	Default	Used by RESRAD (If different from user input)	Parameter Name
R018	Fruits, vegetables and grain consumption (kg/yr)	1.930E+02	1.600E+02	---	DIET(1)
R018	Leafy vegetable consumption (kg/yr)	0.000E+00	1.400E+01	---	DIET(2)
R018	Milk consumption (L/yr)	not used	9.200E+01	---	DIET(3)
R018	Meat and poultry consumption (kg/yr)	not used	6.300E+01	---	DIET(4)
R018	Fish consumption (kg/yr)	not used	5.400E+00	---	DIET(5)
R018	Other seafood consumption (kg/yr)	not used	9.000E-01	---	DIET(6)
R018	Soil ingestion rate (g/yr)	7.300E+01	3.650E+01	---	SOIL
R018	Drinking water intake (L/yr)	4.745E+02	5.100E+02	---	DWI
R018	Contamination fraction of drinking water	1.000E+00	1.000E+00	---	FDW
R018	Contamination fraction of household water	not used	1.000E+00	---	FHHW
R018	Contamination fraction of livestock water	not used	1.000E+00	---	FLW
R018	Contamination fraction of irrigation water	1.000E+00	1.000E+00	---	FIRW
R018	Contamination fraction of aquatic food	not used	5.000E-01	---	FR9
R018	Contamination fraction of plant food	-1	-1	0.500E+00	FPLANT
R018	Contamination fraction of meat	not used	-1	---	FMEAT
R018	Contamination fraction of milk	not used	-1	---	FMILK
R019	Livestock fodder intake for meat (kg/day)	not used	6.800E+01	---	LFI5
R019	Livestock fodder intake for milk (kg/day)	not used	5.500E+01	---	LFI6
R019	Livestock water intake for meat (L/day)	not used	5.000E+01	---	LWI5
R019	Livestock water intake for milk (L/day)	not used	1.600E+02	---	LWI6
R019	Livestock soil intake (kg/day)	not used	5.000E-01	---	LSI
R019	Mass loading for foliar deposition (g/m**3)	1.000E-04	1.000E-04	---	MLFD
R019	Depth of soil mixing layer (m)	1.500E-01	1.500E-01	---	DM
R019	Depth of roots (m)	9.000E-01	9.000E-01	---	DROOT
R019	Drinking water fraction from ground water	1.000E+00	1.000E+00	---	FGWDW
R019	Household water fraction from ground water	1.000E+00	1.000E+00	---	FGWHH
R019	Livestock water fraction from ground water	not used	1.000E+00	---	FGWLW
R019	Irrigation fraction from ground water	not used	1.000E+00	---	FGWIR
R19B	Wet weight crop yield for Non-Leafy (kg/m**2)	7.000E-01	7.000E-01	---	YV(1)
R19B	Wet weight crop yield for Leafy (kg/m**2)	1.500E+00	1.500E+00	---	YV(2)
R19B	Wet weight crop yield for Fodder (kg/m**2)	not used	1.100E+00	---	YV(3)
R19B	Growing Season for Non-Leafy (years)	1.700E-01	1.700E-01	---	TE(1)
R19B	Growing Season for Leafy (years)	2.500E-01	2.500E-01	---	TE(2)
R19B	Growing Season for Fodder (years)	not used	8.000E-02	---	TE(3)
R19B	Translocation Factor for Non-Leafy	1.000E-01	1.000E-01	---	TIV(1)
R19B	Translocation Factor for Leafy	1.000E+00	1.000E+00	---	TIV(2)
R19B	Translocation Factor for Fodder	not used	1.000E+00	---	TIV(3)
R19B	Dry Foliar Interception Fraction for Non-Leafy	2.500E-01	2.500E-01	---	RDRY(1)
R19B	Dry Foliar Interception Fraction for Leafy	2.500E-01	2.500E-01	---	RDRY(2)
R19B	Dry Foliar Interception Fraction for Fodder	not used	2.500E-01	---	RDRY(3)
R19B	Wet Foliar Interception Fraction for Non-Leafy	2.500E-01	2.500E-01	---	RWET(1)
R19B	Wet Foliar Interception Fraction for Leafy	2.500E-01	2.500E-01	---	RWET(2)
R19B	Wet Foliar Interception Fraction for Fodder	not used	2.500E-01	---	RWET(3)
R19B	Weathering Removal Constant for Vegetation	2.000E+01	2.000E+01	---	WLAM
C14	C-12 concentration in water (g/cm**3)	not used	2.000E-05	---	C12WTR
C14	C-12 concentration in contaminated soil (g/g)	not used	3.000E-02	---	C12CZ
C14	Fraction of vegetation carbon from soil	not used	2.000E-02	---	CSOIL

Site-Specific Parameter Summary (continued)

Menu	Parameter	User Input	Default	Used by RESRAD (If different from user input)	Parameter Name
C14	Fraction of vegetation carbon from air	not used	9.800E-01	---	CAIR
C14	C-14 evasion layer thickness in soil (m)	not used	3.000E-01	---	DMC
C14	C-14 evasion flux rate from soil (1/sec)	not used	7.000E-07	---	EVSN
C14	C-12 evasion flux rate from soil (1/sec)	not used	1.000E-10	---	REVSN
C14	Fraction of grain in beef cattle feed	not used	8.000E-01	---	AVFG4
C14	Fraction of grain in milk cow feed	not used	2.000E-01	---	AVFG5
STOR	Storage times of contaminated foodstuffs (days):				
STOR	Fruits, non-leafy vegetables, and grain	1.400E+01	1.400E+01	---	STOR_T(1)
STOR	Leafy vegetables	1.000E+00	1.000E+00	---	STOR_T(2)
STOR	Milk	1.000E+00	1.000E+00	---	STOR_T(3)
STOR	Meat and poultry	2.000E+01	2.000E+01	---	STOR_T(4)
STOR	Fish	7.000E+00	7.000E+00	---	STOR_T(5)
STOR	Crustacea and mollusks	7.000E+00	7.000E+00	---	STOR_T(6)
STOR	Well water	1.000E+00	1.000E+00	---	STOR_T(7)
STOR	Surface water	1.000E+00	1.000E+00	---	STOR_T(8)
STOR	Livestock fodder	4.500E+01	4.500E+01	---	STOR_T(9)
R021	Thickness of building foundation (m)	not used	1.500E-01	---	FLOOR
R021	Bulk density of building foundation (g/cm**3)	not used	2.400E+00	---	DENSFL
R021	Total porosity of the cover material	not used	4.000E-01	---	TPCV
R021	Total porosity of the building foundation	not used	1.000E-01	---	TPFL
R021	Volumetric water content of the cover material	not used	5.000E-02	---	PH2OCV
R021	Volumetric water content of the foundation	not used	3.000E-02	---	PH2OFL
R021	Diffusion coefficient for radon gas (m/sec):				
R021	in cover material	not used	2.000E-06	---	DIFCV
R021	in foundation material	not used	3.000E-07	---	DIFFL
R021	in contaminated zone soil	not used	2.000E-06	---	DIFCZ
R021	Radon vertical dimension of mixing (m)	not used	2.000E+00	---	HMIX
R021	Average building air exchange rate (1/hr)	not used	5.000E-01	---	REXG
R021	Height of the building (room) (m)	not used	2.500E+00	---	HRM
R021	Building interior area factor	not used	0.000E+00	---	FAI
R021	Building depth below ground surface (m)	not used	-1.000E+00	---	DMFL
R021	Emanating power of Rn-222 gas	not used	2.500E-01	---	EMANA(1)
R021	Emanating power of Rn-220 gas	not used	1.500E-01	---	EMANA(2)

Summary of Pathway Selections

Pathway	User Selection
1 -- external gamma	active
2 -- inhalation (w/o radon)	active
3 -- plant ingestion	active
4 -- meat ingestion	suppressed
5 -- milk ingestion	suppressed
6 -- aquatic foods	suppressed
7 -- drinking water	active
8 -- soil ingestion	active
9 -- radon	suppressed
Find peak pathway doses	suppressed

Contaminated Zone Dimensions

Area:	10000.00 square meters
Thickness:	2.00 meters
Cover Depth:	0.00 meters

Initial Soil Concentrations, pCi/g

Co-60	1.000E+00

Total Dose TDOSE(t), mrem/yr

Basic Radiation Dose Limit = 25 mrem/yr
Total Mixture Sum M(t) = Fraction of Basic Dose Limit Received at Time (t)

t (years):	0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	5.000E+01	1.000E+02	3.000E+02	1.000E+03
TDOSE(t):	1.053E+01	9.233E+00	7.096E+00	2.823E+00	2.028E-01	1.457E-02	2.015E-05	7.373E-17	0.000E+00
M(t):	4.213E-01	3.693E-01	2.838E-01	1.129E-01	8.112E-03	5.827E-04	8.060E-07	2.949E-18	0.000E+00

Maximum TDOSE(t): 1.053E+01 mrem/yr at t = 0.000E+00 years

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 0.000E+00 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	1.052E+01	0.9801	4.291E-06	0.0000	0.000E+00	0.0000	2.077E-01	0.0197	0.000E+00	0.0000	0.000E+00	0.0000	1.824E-03	0.0002
Total	1.052E+01	0.9801	4.291E-06	0.0000	0.000E+00	0.0000	2.077E-01	0.0197	0.000E+00	0.0000	0.000E+00	0.0000	1.824E-03	0.0002

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 0.000E+00 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.053E+01	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.053E+01	1.0000

*Sum of all water independent and dependent pathways.

Summary : Co60, resident child, 10000m2
File : C0600000.RAD

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+00 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	9.050E+00	0.9801	3.762E-06	0.0000	0.000E+00	0.0000	1.821E-01	0.0197	0.000E+00	0.0000	0.000E+00	0.0000	1.599E-03	0.0002
Total	9.050E+00	0.9801	3.762E-06	0.0000	0.000E+00	0.0000	1.821E-01	0.0197	0.000E+00	0.0000	0.000E+00	0.0000	1.599E-03	0.0002

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)

As mrem/yr and Fraction of Total Dose At t = 1.000E+00 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	9.233E+00	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	9.233E+00	1.0000

*Sum of all water independent and dependent pathways.

Summary : Co60, resident child, 10000m2
File : C0600C00.RAD

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 3.000E+00 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	6.955E+00	0.9801	2.891E-06	0.0000	0.000E+00	0.0000	1.399E-01	0.0197	0.000E+00	0.0000	0.000E+00	0.0000	1.229E-03	0.0002
Total	6.955E+00	0.9801	2.891E-06	0.0000	0.000E+00	0.0000	1.399E-01	0.0197	0.000E+00	0.0000	0.000E+00	0.0000	1.229E-03	0.0002

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)

As mrem/yr and Fraction of Total Dose At t = 3.000E+00 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	7.096E+00	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	7.096E+00	1.0000

*Sum of all water independent and dependent pathways.

Summary : Co60, resident child, 10000m2
File : C0600000.RAD

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+01 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	2.767E+00	0.9801	1.150E-06	0.0000	0.000E+00	0.0000	5.566E-02	0.0197	0.000E+00	0.0000	0.000E+00	0.0000	4.889E-04	0.0002
Total	2.767E+00	0.9801	1.150E-06	0.0000	0.000E+00	0.0000	5.566E-02	0.0197	0.000E+00	0.0000	0.000E+00	0.0000	4.889E-04	0.0002

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+01 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.823E+00	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.823E+00	1.0000

*Sum of all water independent and dependent pathways.

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 3.000E+01 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	1.988E-01	0.9801	8.262E-08	0.0000	0.000E+00	0.0000	3.998E-03	0.0197	0.000E+00	0.0000	0.000E+00	0.0000	3.512E-05	0.0002
Total	1.988E-01	0.9801	8.262E-08	0.0000	0.000E+00	0.0000	3.998E-03	0.0197	0.000E+00	0.0000	0.000E+00	0.0000	3.512E-05	0.0002

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 3.000E+01 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.028E-01	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.028E-01	1.0000

*Sum of all water independent and dependent pathways.

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 5.000E+01 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	1.423E-02	0.9801	5.935E-09	0.0000	0.000E+00	0.0000	2.872E-04	0.0197	0.000E+00	0.0000	0.000E+00	0.0000	2.523E-06	0.0002
Total	1.428E-02	0.9801	5.935E-09	0.0000	0.000E+00	0.0000	2.872E-04	0.0197	0.000E+00	0.0000	0.000E+00	0.0000	2.523E-06	0.0002

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 5.000E+01 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.457E-02	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.457E-02	1.0000

*Sum of all water independent and dependent pathways.

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+02 years
Water Independent Pathways (Inhalation excludes radon)

Radio-	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
Nuclide	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	1.975E-05	0.9801	8.209E-12	0.0000	0.000E+00	0.0000	3.973E-07	0.0197	0.000E+00	0.0000	0.000E+00	0.0000	3.489E-09	0.0002
Total	1.975E-05	0.9801	8.209E-12	0.0000	0.000E+00	0.0000	3.973E-07	0.0197	0.000E+00	0.0000	0.000E+00	0.0000	3.489E-09	0.0002

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)

As mrem/yr and Fraction of Total Dose At t = 1.000E+02 years
Water Dependent Pathways

Radio-	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
Nuclide	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.015E-05	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.015E-05	1.0000

*Sum of all water independent and dependent pathways.

Summary : Co60, resident child, 10000m2
File : 00600000.RAD

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 3.000E+02 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	7.227E-17	0.9801	3.004E-23	0.0000	0.000E+00	0.0000	1.454E-18	0.0197	0.000E+00	0.0000	0.000E+00	0.0000	1.277E-20	0.0002
Total	7.227E-17	0.9801	3.004E-23	0.0000	0.000E+00	0.0000	1.454E-18	0.0197	0.000E+00	0.0000	0.000E+00	0.0000	1.277E-20	0.0002

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 3.000E+02 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	7.373E-17	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	7.373E-17	1.0000

*Sum of all water independent and dependent pathways.

Summary : Co60, resident child, 10000m2
File : C0600C00.RAD

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+03 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+03 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Co-60	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000

*Sum of all water independent and dependent pathways.

Dose/Source Ratios Summed Over All Pathways
Parent and Progeny Principal Radionuclide Contributions Indicated

Parent (i)	Product (j)	Branch Fraction*	DSR(j,t) (mrem/yr)/(pCi/g) t= 0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	5.000E+01	1.000E+02	3.000E+02	1.000E+03
Co-60	Co-60	1.000E+00	1.053E+01	9.233E+00	7.096E+00	2.823E+00	2.028E-01	1.457E-02	2.015E-05	7.373E-17	0.000E+00

*Branch Fraction is the cumulative factor for the j't principal radionuclide daughter: CUMBRF(j) = BRF(1)*BRF(2)* ... BRF(j).
The DSR includes contributions from associated (half-life <= 0.5 yr) daughters.

Single Radionuclide Soil Guidelines G(i,t) in pCi/g

Basic Radiation Dose Limit = 25 mrem/yr

Nuclide (i)	t= 0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	5.000E+01	1.000E+02	3.000E+02	1.000E+03
Co-60	2.374E+00	2.708E+00	3.523E+00	8.856E+00	1.233E+02	1.716E+03	1.241E+06	*1.131E+15	*1.131E+15

*At specific activity limit

Summed Dose/Source Ratios DSR(i,t) in (mrem/yr)/(pCi/g)

and Single Radionuclide Soil Guidelines G(i,t) in pCi/g
at tmin = time of minimum single radionuclide soil guideline
and at tmax = time of maximum total dose = 0.000E+00 years

Nuclide (i)	Initial pCi/g	tmin (years)	DSR(i,tmin) (pCi/g)	G(i,tmin) (pCi/g)	DSR(i,tmax)	G(i,tmax) (pCi/g)
Co-60	1.000E+00	0.000E+00	1.053E+01	2.374E+00	1.053E+01	2.374E+00

Individual Nuclide Dose Summed Over All Pathways
Parent Nuclide and Branch Fraction Indicated
DOSE(j,t), mrem/yr

Nuclide (j)	Parent (i)	BRF(i)	t= 0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	5.000E+01	1.000E+02	3.000E+02	1.000E+03
Co-60	Co-60	1.000E+00	1.053E+01	9.233E+00	7.096E+00	2.823E+00	2.028E-01	1.457E-02	2.015E-05	7.373E-17	0.000E+00

BRF(i) is the branch fraction of the parent nuclide.

Individual Nuclide Soil Concentration
Parent Nuclide and Branch Fraction Indicated
S(j,t), pCi/g

Nuclide (j)	Parent (i)	BRF(i)	t= 0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	5.000E+01	1.000E+02	3.000E+02	1.000E+03
Co-60	Co-60	1.000E+00	1.000E+00	8.766E-01	6.737E-01	2.680E-01	1.925E-02	1.383E-03	1.913E-06	7.000E-18	0.000E+00

BRF(i) is the branch fraction of the parent nuclide.

A.3.2 Cs-137: Resident Child Summary Output File

Summary : CS137, resident child, 10000m2
File : C1370000.RAD

Table of Contents

Part I: Mixture Sums and Single Radionuclide Guidelines

Dose Conversion Factor (and Related) Parameter Summary ... 2
Site-Specific Parameter Summary 3
Summary of Pathway Selections 7
Contaminated Zone and Total Dose Summary 8
Total Dose Components
Time = 0.000E+00 9
Time = 1.000E+00 10
Time = 3.000E+00 11
Time = 1.000E+01 12
Time = 3.000E+01 13
Time = 5.000E+01 14
Time = 1.000E+02 15
Time = 3.000E+02 16
Time = 1.000E+03 17
Dose/Source Ratios Summed Over All Pathways 18
Single Radionuclide Soil Guidelines 18
Dose Per Nuclide Summed Over All Pathways 19
Soil Concentration Per Nuclide 19

Dose Conversion Factor (and Related) Parameter Summary
File: DOSFAC.BIN

Menu	Parameter	Current Value	Default	Parameter Name
B-1	Dose conversion factors for inhalation, mrem/pCi:			
B-1	Cs-137+D	3.190E-05	3.190E-05	DCF2(1)
D-1	Dose conversion factors for ingestion, mrem/pCi:			
D-1	Cs-137+D	5.000E-05	5.000E-05	DCF3(1)
D-34	Food transfer factors:			
D-34	Cs-137+D , plant/soil concentration ratio, dimensionless	4.000E-02	4.000E-02	RTF(1,1)
D-34	Cs-137+D , beef/livestock-intake ratio, (pCi/kg)/(pCi/d)	3.000E-02	3.000E-02	RTF(1,2)
D-34	Cs-137+D , milk/livestock-intake ratio, (pCi/L)/(pCi/d)	8.000E-03	8.000E-03	RTF(1,3)
D-5	Bioaccumulation factors, fresh water, L/kg:			
D-5	Cs-137+D , fish	2.000E+03	2.000E+03	BIOFAC(1,1)
D-5	Cs-137+D , crustacea and mollusks	1.000E+02	1.000E+02	BIOFAC(1,2)

Site-Specific Parameter Summary

Menu	Parameter	User Input	Default	Used by RESRAD (If different from user input)	Parameter Name
R011	Area of contaminated zone (m**2)	1.000E+04	1.000E+04	---	AREA
R011	Thickness of contaminated zone (m)	2.000E+00	2.000E+00	---	THICK0
R011	Length parallel to aquifer flow (m)	1.000E+02	1.000E+02	---	LCZPAQ
R011	Basic radiation dose limit (mrem/yr)	2.500E+01	3.000E+01	---	BRDL
R011	Time since placement of material (yr)	0.000E+00	0.000E+00	---	TI
R011	Times for calculations (yr)	1.000E+00	1.000E+00	---	T(2)
R011	Times for calculations (yr)	3.000E+00	3.000E+00	---	T(3)
R011	Times for calculations (yr)	1.000E+01	1.000E+01	---	T(4)
R011	Times for calculations (yr)	3.000E+01	3.000E+01	---	T(5)
R011	Times for calculations (yr)	5.000E+01	1.000E+02	---	T(6)
R011	Times for calculations (yr)	1.000E+02	3.000E+02	---	T(7)
R011	Times for calculations (yr)	3.000E+02	1.000E+03	---	T(8)
R011	Times for calculations (yr)	1.000E+03	0.000E+00	---	T(9)
R011	Times for calculations (yr)	not used	0.000E+00	---	T(10)
R012	Initial principal radionuclide (pCi/g): Cs-137	1.000E+00	0.000E+00	---	S1(1)
R012	Concentration in groundwater (pCi/L): Cs-137	not used	0.000E+00	---	W1(1)
R013	Cover depth (m)	0.000E+00	0.000E+00	---	COVER0
R013	Density of cover material (g/cm**3)	not used	1.500E+00	---	DENSCV
R013	Cover depth erosion rate (m/yr)	not used	1.000E-03	---	VCV
R013	Density of contaminated zone (g/cm**3)	1.500E+00	1.500E+00	---	DENSCZ
R013	Contaminated zone erosion rate (m/yr)	0.000E+00	1.000E-03	---	VCZ
R013	Contaminated zone total porosity	4.000E-01	4.000E-01	---	TPCZ
R013	Contaminated zone effective porosity	2.000E-01	2.000E-01	---	EPCZ
R013	Contaminated zone hydraulic conductivity (m/yr)	1.000E+01	1.000E+01	---	HCCZ
R013	Contaminated zone b parameter	5.300E+00	5.300E+00	---	BCZ
R013	Average annual wind speed (m/sec)	2.000E+00	2.000E+00	---	WIND
R013	Humidity in air (g/m**3)	not used	8.000E+00	---	HUMID
R013	Evapotranspiration coefficient	5.000E-01	5.000E-01	---	EVAPTR
R013	Precipitation (m/yr)	1.000E+00	1.000E+00	---	PRECIP
R013	Irrigation (m/yr)	2.000E-01	2.000E-01	---	RI
R013	Irrigation mode	overhead	overhead	---	IDITCH
R013	Runoff coefficient	2.000E-01	2.000E-01	---	RUNOFF
R013	Watershed area for nearby stream or pond (m**2)	1.000E+06	1.000E+06	---	WAREA
R013	Accuracy for water/soil computations	1.000E-03	1.000E-03	---	EPS
R014	Density of saturated zone (g/cm**3)	1.500E+00	1.500E+00	---	DENSAQ
R014	Saturated zone total porosity	4.000E-01	4.000E-01	---	TPSZ
R014	Saturated zone effective porosity	2.000E-01	2.000E-01	---	EPSZ
R014	Saturated zone hydraulic conductivity (m/yr)	1.000E+02	1.000E+02	---	HCSZ
R014	Saturated zone hydraulic gradient	2.000E-02	2.000E-02	---	HGWT
R014	Saturated zone b parameter	5.300E+00	5.300E+00	---	BSZ
R014	Water table drop rate (m/yr)	1.000E-03	1.000E-03	---	VWT
R014	Well pump intake depth (m below water table)	1.000E+01	1.000E+01	---	DWIBWT
R014	Model: Nondispersion (ND) or Mass-Balance (MB)	ND	ND	---	MODEL
R014	Well pumping rate (m**3/yr)	2.500E+02	2.500E+02	---	UW
R015	Number of unsaturated zone strata	1	1	---	NS

Site-Specific Parameter Summary (continued)

Menu	Parameter	User Input	Default	Used by RESRAD (If different from user input)	Parameter Name
R015	Unsat. zone 1, thickness (m)	4.000E+00	4.000E+00	---	H(1)
R015	Unsat. zone 1, soil density (g/cm**3)	1.500E+00	1.500E+00	---	DENSUZ(1)
R015	Unsat. zone 1, total porosity	4.00UE-01	4.000E-01	---	TPUZ(1)
R015	Unsat. zone 1, effective porosity	2.000E-01	2.000E-01	---	EPUZ(1)
R015	Unsat. zone 1, soil-specific b parameter	5.300E+00	5.300E+00	---	BUZ(1)
R015	Unsat. zone 1, hydraulic conductivity (m/yr)	1.000E+01	1.000E+01	---	HCUZ(1)
R016	Distribution coefficients for Cs-137				
R016	Contaminated zone (cm**3/g)	1.000E+03	1.000E+03	---	DCNUCC(1)
R016	Unsaturated zone 1 (cm**3/g)	1.000E+03	1.000E+03	---	DCNUCU(1,1)
R016	Saturated zone (cm**3/g)	1.000E+03	1.000E+03	---	DCNUCS(1)
R016	Leach rate (/yr)	0.000E+00	0.000E+00	1.666E-04	ALEACH(1)
R016	Solubility constant	0.000E+00	0.000E+00	not used	SOLUBK(1)
R017	Inhalation rate (m**3/yr)	2.720E+03	8.400E+03	---	INHALR
R017	Mass loading for inhalation (g/m**3)	1.000E-04	1.000E-04	---	MLINH
R017	Exposure duration	6.000E+00	3.000E+01	---	ED
R017	Shielding factor, inhalation	4.000E-01	4.000E-01	---	SHF3
R017	Shielding factor, external gamma	7.000E-01	7.000E-01	---	SHF1
R017	Fraction of time spent indoors	8.390E-01	5.000E-01	---	FIND
R017	Fraction of time spent outdoors (on site)	8.990E-02	2.500E-01	---	FOTD
R017	Shape factor flag, external gamma	1.000E+00	1.000E+00	>0 shows circular AREA.	FS
R017	Radii of shape factor array (used if FS = -1):				
R017	Outer annular radius (m), ring 1:	not used	5.000E+01	---	RAD_SHAPE(1)
R017	Outer annular radius (m), ring 2:	not used	7.071E+01	---	RAD_SHAPE(2)
R017	Outer annular radius (m), ring 3:	not used	0.000E+00	---	RAD_SHAPE(3)
R017	Outer annular radius (m), ring 4:	not used	0.000E+00	---	RAD_SHAPE(4)
R017	Outer annular radius (m), ring 5:	not used	0.000E+00	---	RAD_SHAPE(5)
R017	Outer annular radius (m), ring 6:	not used	0.000E+00	---	RAD_SHAPE(6)
R017	Outer annular radius (m), ring 7:	not used	0.000E+00	---	RAD_SHAPE(7)
R017	Outer annular radius (m), ring 8:	not used	0.000E+00	---	RAD_SHAPE(8)
R017	Outer annular radius (m), ring 9:	not used	0.000E+00	---	RAD_SHAPE(9)
R017	Outer annular radius (m), ring 10:	not used	0.000E+00	---	RAD_SHAPE(10)
R017	Outer annular radius (m), ring 11:	not used	0.000E+00	---	RAD_SHAPE(11)
R017	Outer annular radius (m), ring 12:	not used	0.000E+00	---	RAD_SHAPE(12)
R017	Fractions of annular areas within AREA:				
R017	Ring 1	not used	1.000E+00	---	FRACA(1)
R017	Ring 2	not used	2.732E-01	---	FRACA(2)
R017	Ring 3	not used	0.000E+00	---	FRACA(3)
R017	Ring 4	not used	0.000E+00	---	FRACA(4)
R017	Ring 5	not used	0.000E+00	---	FRACA(5)
R017	Ring 6	not used	0.000E+00	---	FRACA(6)
R017	Ring 7	not used	0.000E+00	---	FRACA(7)
R017	Ring 8	not used	0.000E+00	---	FRACA(8)
R017	Ring 9	not used	0.000E+00	---	FRACA(9)
R017	Ring 10	not used	0.000E+00	---	FRACA(10)
R017	Ring 11	not used	0.000E+00	---	FRACA(11)
R017	Ring 12	not used	0.000E+00	---	FRACA(12)

Site-Specific Parameter Summary (continued)

Menu	Parameter	User Input	Default	Used by RESRAD (If different from user input)	Parameter Name
R018	Fruits, vegetables and grain consumption (kg/yr)	1.930E+02	1.600E+02	---	DIET(1)
R018	Leafy vegetable consumption (kg/yr)	0.000E+00	1.400E+01	---	DIET(2)
R018	Milk consumption (L/yr)	not used	9.200E+01	---	DIET(3)
R018	Meat and poultry consumption (kg/yr)	not used	6.300E+01	---	DIET(4)
R018	Fish consumption (kg/yr)	not used	5.400E+00	---	DIET(5)
R018	Other seafood consumption (kg/yr)	not used	9.000E-01	---	DIET(6)
R018	Soil ingestion rate (g/yr)	7.300E+01	3.650E+01	---	SOIL
R018	Drinking water intake (L/yr)	4.745E+02	5.100E+02	---	DWI
R018	Contamination fraction of drinking water	1.000E+00	1.000E+00	---	FDW
R018	Contamination fraction of household water	not used	1.000E+00	---	FHHW
R018	Contamination fraction of livestock water	not used	1.000E+00	---	FLW
R018	Contamination fraction of irrigation water	1.000E+00	1.000E+00	---	FIRW
R018	Contamination fraction of aquatic food	not used	5.000E-01	---	FR9
R018	Contamination fraction of plant food	-1	-1	0.500E+00	FPLANT
R018	Contamination fraction of meat	not used	-1	---	FMEAT
R018	Contamination fraction of milk	not used	-1	---	FMILK
R019	Livestock fodder intake for meat (kg/day)	not used	6.800E+01	---	LFI5
R019	Livestock fodder intake for milk (kg/day)	not used	5.500E+01	---	LFI6
R019	Livestock water intake for meat (L/day)	not used	5.000E+01	---	LWI5
R019	Livestock water intake for milk (L/day)	not used	1.600E+02	---	LWI6
R019	Livestock soil intake (kg/day)	not used	5.000E-01	---	LSI
R019	Mass loading for foliar deposition (g/m**3)	1.000E-04	1.000E-04	---	MLFD
R019	Depth of soil mixing layer (m)	1.500E-01	1.500E-01	---	DM
R019	Depth of roots (m)	9.000E-01	9.000E-01	---	DROOT
R019	Drinking water fraction from ground water	1.000E+00	1.000E+00	---	FGWDW
R019	Household water fraction from ground water	1.000E+00	1.000E+00	---	FGWHH
R019	Livestock water fraction from ground water	not used	1.000E+00	---	FGWLW
R019	Irrigation fraction from ground water	not used	1.000E+00	---	FGWIR
R19B	Wet weight crop yield for Non-Leafy (kg/m**2)	7.000E-01	7.000E-01	---	YV(1)
R19B	Wet weight crop yield for Leafy (kg/m**2)	1.500E+00	1.500E+00	---	YV(2)
R19B	Wet weight crop yield for Fodder (kg/m**2)	not used	1.100E+00	---	YV(3)
R19B	Growing Season for Non-Leafy (years)	1.700E-01	1.700E-01	---	TE(1)
R19B	Growing Season for Leafy (years)	2.500E-01	2.500E-01	---	TE(2)
R19B	Growing Season for Fodder (years)	not used	8.000E-02	---	TE(3)
R19B	Translocation Factor for Non-Leafy	1.000E-01	1.000E-01	---	TIV(1)
R19B	Translocation Factor for Leafy	1.000E+00	1.000E+00	---	TIV(2)
R19B	Translocation Factor for Fodder	not used	1.000E+00	---	TIV(3)
R19B	Dry Foliar Interception Fraction for Non-Leafy	2.500E-01	2.500E-01	---	RDRY(1)
R19B	Dry Foliar Interception Fraction for Leafy	2.500E-01	2.500E-01	---	RDRY(2)
R19B	Dry Foliar Interception Fraction for Fodder	not used	2.500E-01	---	RDRY(3)
R19B	Wet Foliar Interception Fraction for Non-Leafy	2.500E-01	2.500E-01	---	RWET(1)
R19B	Wet Foliar Interception Fraction for Leafy	2.500E-01	2.500E-01	---	RWET(2)
R19B	Wet Foliar Interception Fraction for Fodder	not used	2.500E-01	---	RWET(3)
R19B	Weathering Removal Constant for Vegetation	2.000E+01	2.000E+01	---	WLAM
C14	C-12 concentration in water (g/cm**3)	not used	2.000E-05	---	C12WTR
C14	C-12 concentration in contaminated soil (g/g)	not used	3.000E-02	---	C12CZ
C14	Fraction of vegetation carbon from soil	not used	2.000E-02	---	CSOIL

Site-Specific Parameter Summary (continued)

Menu	Parameter	User Input	Default	Used by RESRAD (If different from user input)	Parameter Name
C14	Fraction of vegetation carbon from air	not used	9.800E-01	---	CAIR
C14	C-14 evasion layer thickness in soil (m)	not used	3.000E-01	---	DMC
C14	C-14 evasion flux rate from soil (1/sec)	not used	7.300E-07	---	EVSN
C14	C-12 evasion flux rate from soil (1/sec)	not used	1.000E-10	---	REVSN
C14	Fraction of grain in beef cattle feed	not used	8.000E-01	---	AVFG4
C14	Fraction of grain in milk cow feed	not used	2.000E-01	---	AVFG5
STOR	Storage times of contaminated foodstuffs (days):				
STOR	Fruits, non-leafy vegetables, and grain	1.400E+01	1.400E+01	---	STOR_T(1)
STOR	Leafy vegetables	1.000E+00	1.000E+00	---	STOR_T(2)
STOR	Milk	1.000E+00	1.000E+00	---	STOR_T(3)
STOR	Meat and poultry	2.000E+01	2.000E+01	---	STOR_T(4)
STOR	Fish	7.000E+00	7.000E+00	---	STOR_T(5)
STOR	Crustacea and mollusks	7.000E+00	7.000E+00	---	STOR_T(6)
STOR	Well water	1.000E+00	1.000E+00	---	STOR_T(7)
STOR	Surface water	1.000E+00	1.000E+00	---	STOR_T(8)
STOR	Livestock fodder	4.500E+01	4.500E+01	---	STOR_T(9)
R021	Thickness of building foundation (m)	not used	1.500E-01	---	FLOOR
R021	Bulk density of building foundation (g/cm**3)	not used	2.400E+00	---	DENSFL
R021	Total porosity of the cover material	not used	4.000E-01	---	TPCV
R021	Total porosity of the building foundation	not used	1.000E-01	---	TPFL
R021	Volumetric water content of the cover material	not used	5.000E-02	---	PH2OCV
R021	Volumetric water content of the foundation	not used	3.000E-02	---	PH2OFL
R021	Diffusion coefficient for radon gas (m/sec):				
R021	in cover material	not used	2.000E-06	---	DIFCV
R021	in foundation material	not used	3.000E-07	---	DIFFL
R021	in contaminated zone soil	not used	2.000E-06	---	DIFCZ
R021	Radon vertical dimension of mixing (m)	not used	2.000E+00	---	HMIX
R021	Average building air exchange rate (1/hr)	not used	5.000E-01	---	REXG
R021	Height of the building (room) (m)	not used	2.500E+00	---	HRM
R021	Building interior area factor	not used	0.000E+00	---	FAI
R021	Building depth below ground surface (m)	not used	-1.000E+00	---	DMFL
R021	Emanating power of Rn-222 gas	not used	2.500E-01	---	EMANA(1)
R021	Emanating power of Rn-220 gas	not used	1.500E-01	---	EMANA(2)

Summary of Pathway Selections

Pathway	User Selection
1 -- external gamma	active
2 -- inhalation (w/o radon)	active
3 -- plant ingestion	active
4 -- meat ingestion	suppressed
5 -- milk ingestion	suppressed
6 -- aquatic foods	suppressed
7 -- drinking water	active
8 -- soil ingestion	active
9 -- radon	suppressed
Find peak pathway doses	suppressed

Contaminated Zone Dimensions		Initial Soil Concentrations, pCi/g	
Area:	10000.00 square meters	Cs-137	1.000E+00
Thickness:	2.00 meters		
Cover Depth:	0.00 meters		

Total Dose TDOSE(t), mrem/yr
Basic Radiation Dose Limit = 25 mrem/yr
Total Mixture Sum M(t) = Fraction of Basic Dose Limit Received at Time (t)

t (years):	0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	5.000E+01	1.000E+02	3.000E+02	1.000E+03
TDOSE(t):	2.358E+00	2.304E+00	2.199E+00	1.868E+00	1.173E+00	7.365E-01	2.301E-01	2.190E-03	1.844E-10
M(t):	9.431E-02	9.215E-02	8.796E-02	7.473E-02	4.692E-02	2.946E-02	9.203E-03	8.761E-05	7.377E-12

Maximum TDOSE(t): 2.358E+00 mrem/yr at t = 0.000E+00 years

Summary : CS137, resident child, 10000m2
File : C137C000.RAD

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 0.000E+00 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	2.161E+00	0.9167	6.250E-07	0.0000	0.000E+00	0.0000	1.930E-01	0.0819	0.000E+00	0.0000	0.000E+00	0.0000	3.390E-03	0.0014
Total	2.161E+00	0.9167	6.250E-07	0.0000	0.000E+00	0.0000	1.930E-01	0.0819	0.000E+00	0.0000	0.000E+00	0.0000	3.390E-03	0.0014

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 0.000E+00 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.358E+00	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.358E+00	1.0000

*Sum of all water independent and dependent pathways.

Summary : CS137, resident child, 10000m2
File : C137C000.RAD

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+00 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	2.112E+00	0.9167	6.107E-07	0.0000	0.000E+00	0.0000	1.886E-01	0.0819	0.000E+00	0.0000	0.000E+00	0.0000	3.312E-03	0.0014
Total	2.112E+00	0.9167	6.107E-07	0.0000	0.000E+00	0.0000	1.886E-01	0.0819	0.000E+00	0.0000	0.000E+00	0.0000	3.312E-03	0.0014

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+00 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.304E+00	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.304E+00	1.0000

*Sum of all water independent and dependent pathways.

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 3.000E+00 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	2.016E+00	0.9167	5.829E-07	0.0000	0.000E+00	0.0000	1.800E-01	0.0819	0.000E+00	0.0000	0.000E+00	0.0000	3.162E-03	0.0014
Total	2.016E+00	0.9167	5.829E-07	0.0000	0.000E+00	0.0000	1.800E-01	0.0819	0.000E+00	0.0000	0.000E+00	0.0000	3.162E-03	0.0014

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 3.000E+00 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.199E+00	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.199E+00	1.0000

*Sum of all water independent and dependent pathways.

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+01 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	1.713E+00	0.9167	4.953E-07	0.0000	0.000E+00	0.0000	1.529E-01	0.0819	0.000E+00	0.0000	0.000E+00	0.0000	2.687E-03	0.0014
Total	1.713E+00	0.9167	4.953E-07	0.0000	0.000E+00	0.0000	1.529E-01	0.0819	0.000E+00	0.0000	0.000E+00	0.0000	2.687E-03	0.0014

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+01 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.868E+00	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.868E+00	1.0000

*Sum of all water independent and dependent pathways.

Summary : CS137, resident child, 10000m2
File : C137C000.RAD

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 3.000E+01 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	1.075E+00	0.9167	3.110E-07	0.0000	0.000E+00	0.0000	9.602E-02	0.0819	0.000E+00	0.0000	0.000E+00	0.0000	1.687E-03	0.0014
Total	1.075E+00	0.9167	3.110E-07	0.0000	0.000E+00	0.0000	9.602E-02	0.0819	0.000E+00	0.0000	0.000E+00	0.0000	1.687E-03	0.0014

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)

As mrem/yr and Fraction of Total Dose At t = 3.000E+01 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.173E+00	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.173E+00	1.0000

*Sum of all water independent and dependent pathways.

Summary : CS137, resident child, 10000m2
File : C137C000.RAD

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 5.000E+01 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	6.752E-01	0.9167	1.952E-07	0.0000	0.000E+00	0.0000	6.029E-02	0.0819	0.000E+00	0.0000	0.000E+00	0.0000	1.059E-03	0.0014
Total	6.752E-01	0.9167	1.952E-07	0.0000	0.000E+00	0.0000	6.029E-02	0.0819	0.000E+00	0.0000	0.000E+00	0.0000	1.059E-03	0.0014

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)

As mrem/yr and Fraction of Total Dose At t = 5.000E+01 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	7.365E-01	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	7.365E-01	1.0000

*Sum of all water independent and dependent pathways.

Summary : CS137, resident child, 10000m2
File : C1370000.RAD

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+02 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	2.109E-01	0.9167	6.099E-08	0.0000	0.000E+00	0.0000	1.883E-02	0.0819	0.000E+00	0.0000	0.000E+00	0.0000	3.308E-04	0.0014
Total	2.109E-01	0.9167	6.099E-08	0.0000	0.000E+00	0.0000	1.883E-02	0.0819	0.000E+00	0.0000	0.000E+00	0.0000	3.308E-04	0.0014

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+02 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.301E-01	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.301E-01	1.0000

*Sum of all water independent and dependent pathways.

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 3.000E+02 years
Water Independent Pathways (Inhalation excludes radon)

Radio-	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
Nuclide	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	2.008E-03	0.9167	5.806E-10	0.0000	0.000E+00	0.0000	1.793E-04	0.0819	0.000E+00	0.0000	0.000E+00	0.0000	3.150E-06	0.0014
Total	2.008E-03	0.9167	5.806E-10	0.0000	0.000E+00	0.0000	1.793E-04	0.0819	0.000E+00	0.0000	0.000E+00	0.0000	3.150E-06	0.0014

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 3.000E+02 years
Water Dependent Pathways

Radio-	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
Nuclide	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.190E-03	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	2.190E-03	1.0000

*Sum of all water independent and dependent pathways.

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+03 years
Water Independent Pathways (Inhalation excludes radon)

Radio-Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	1.691E-10	0.9167	4.889E-17	0.0000	0.000E+00	0.0000	1.510E-11	0.0819	0.000E+00	0.0000	0.000E+00	0.0000	2.652E-13	0.0014
Total	1.691E-10	0.9167	4.889E-17	0.0000	0.000E+00	0.0000	1.510E-11	0.0819	0.000E+00	0.0000	0.000E+00	0.0000	2.652E-13	0.0014

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+03 years
Water Dependent Pathways

Radio-Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
Cs-137	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.844E-10	1.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.844E-10	1.0000

*Sum of all water independent and dependent pathways.

Summary : CS137, resident child, 10000m2
File : C137CCCC.RAD

Dose/Source Ratios Summed Over All Pathways
Parent and Progeny Principal Radionuclide Contributions Indicated

Parent (i)	Product (j)	Branch Fraction*	DSR(j,t) (mrem/yr)/(pCi/g) t= 0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	5.000E+01	1.000E+02	3.000E+02	1.000E+03
Cs-137	Cs-137	1.000E+00	2.358E+00	2.304E+00	2.199E+00	1.868E+00	1.173E+00	7.365E-01	2.301E-01	2.[illegible]0E-05	1.944E-10

*Branch Fraction is the cumulative factor for the j'th principal radionuclide daughter: CUMBRF(j) = BRF(1)*BRF(2)* ... BRF(j).
The DSR includes contributions from associated (half-life <= 0.5 yr) daughters.

Single Radionuclide Soil Guidelines G(i,t) in pCi/g
Basic Radiation Dose Limit = 25 mrem/yr

Nuclide (i)	t= 0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	5.000E+01	1.000E+02	3.000E+02	1.000E+03
Cs-137	1.060E+01	1.085E+01	1.137E+01	1.338E+01	2.131E+01	3.394E+01	1.087E+02	1.141E+04	1.356E+11

Summed Dose/Source Ratios DSR(i,t) in (mrem/yr)/(pCi/g)
and Single Radionuclide Soil Guidelines G(i,t) in pCi/g
at tmin = time of minimum single radionuclide soil guideline
and at tmax = time of maximum total dose = 0.000E+00 years

Nuclide (i)	Initial pCi/g	tmin (years)	DSR(i,tmin)	G(i,tmin) (pCi/g)	DSR(i,tmax)	G(i,tmax) (pCi/g)
Cs-137	1.000E+00	0.000E+00	2.358E+00	1.060E+01	2.358E+00	1.060E+01

Individual Nuclide Dose Summed Over All Pathways
Parent Nuclide and Branch Fraction Indicated

Nuclide (j)	Parent (i)	BRF(i)	DOSE(j,t), mrem/yr t= 0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	5.000E+01	1.000E+02	3.000E+02	1.000E+03
Cs-137	Cs-137	1.000E+00	2.358E+00	2.304E+00	2.199E+00	1.868E+00	1.173E+00	7.365E-01	2.301E-01	2.190E-03	1.844E-10

BRF(i) is the branch fraction of the parent nuclide.

Individual Nuclide Soil Concentration
Parent Nuclide and Branch Fraction Indicated

Nuclide (j)	Parent (i)	BRF(i)	S(j,t), pCi/g t= 0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	5.000E+01	1.000E+02	3.000E+02	1.000E+03
Cs-137	Cs-137	1.000E+00	1.000E+00	9.770E-01	9.326E-01	7.924E-01	4.975E-01	3.124E-01	9.757E-02	9.289E-04	7.822E-11

BRF(i) is the branch fraction of the parent nuclide.