

DRAWER 29
FILE 8

UNITED STATES DEPARTMENT OF THE INTERIOR
GEOLOGICAL SURVEY

INVESTIGATION OF AN AEROMAGNETIC ANOMALY
ON WEST SIDE OF YUCCA MOUNTAIN, NYE COUNTY, NEVADA

by

G. D. Bath and C. E. Jahren

Open-File Report 85-459

Prepared in cooperation with the
Nevada Operations Office
U.S. Department of Energy
(Interagency Agreement DE-AI08-73ET44802)

This report is preliminary and has not been reviewed for conformity with U.S. Geological Survey editorial standards and stratigraphic nomenclature. Company names are for descriptive purposes only and do not constitute endorsement by the U.S. Geological Survey.

Denver, Colorado
1985

Copies of this Open-File Report
may be purchased from

Open-File Services Section
Branch of Distribution
U.S. Geological Survey
Box 25425 Federal Center
Denver, Colorado 80225

PREPAYMENT IS REQUIRED

Price information will be published
in the monthly listing
"New Publications of the Geological Survey"

FOR ADDITIONAL ORDERING INFORMATION

CALL: Commercial: (303) 236-7476
FTS: 776-7476

Open-File Report 85-459

Open-File Report 85-459

UNITED STATES
DEPARTMENT OF THE INTERIOR
GEOLOGICAL SURVEY

INVESTIGATION OF AN AEROMAGNETIC ANOMALY
ON WEST SIDE OF YUCCA MOUNTAIN, NYE COUNTY, NEVADA

By

G. D. Bath and C. E. Jahren

CONTENTS

	Page
Abstract.....	1
Introduction.....	1
System of magnetic units.....	3
Acknowledgments.....	3
Observed, residual and smoothed anomalies.....	4
Magnetic properties.....	4
Estimate of magnetization.....	11
Anomaly interpretations.....	13
Aeromagnetic anomalies.....	15
Ground magnetic anomalies.....	15
Discussion.....	21
References Cited.....	22

ILLUSTRATIONS

	Page
Figure 1.--Residual aeromagnetic map of southern part of potential repository site.....	2
2.--Topographic map of area west of Yucca Mountain.....	5
3.--Aeromagnetic anomalies and altimeter records for traverses A, B, C, and D.....	6
4.--Residual and smoothed anomalies on traverse A82.....	7
5.--Residual and smoothed anomalies on traverse A83.....	8
6.--Residual and smoothed anomalies on traverse B83.....	9
7.--Residual and smoothed anomalies on traverse C83.....	10
8.--Geologic map of western part of Yucca Mountain.....	14
9.--Section along air traverse B-B' over models of dipole, source A, and source B.....	16
10.--Section along ground traverse C-C' over models of Solitario Canyon fault, source C and source D.....	19

TABLES

Table 1.--Magnetic properties and thicknesses of units penetrated in three drill holes.....	12
2.--Total magnetizations for models of figure 9.....	17
3.--Aeromagnetic anomalies computed for models of figure 9.....	17
4.--Total magnetizations of component parts of models of figure 10.....	20
5.--Ground magnetic and aeromagnetic anomalies computed for models of figure 10.....	20

UNITED STATES
DEPARTMENT OF THE INTERIOR
GEOLOGICAL SURVEY

INVESTIGATION OF AN AEROMAGNETIC ANOMALY ON WEST SIDE
OF YUCCA MOUNTAIN, NYE COUNTY, NEVADA

By

G. D. Bath and C. E. Jahren

ABSTRACT

Investigations of the source of a prominent aeromagnetic anomaly of 290 nT were undertaken at a potential repository site located in the Yucca Mountain area, Nevada Test Site. The anomaly was detected on a recent flight line of a survey flown north-south at 400 m (1,300 ft) spacing and 122 m (400 ft) above the surface. The anomaly was not detected on older lines flown east-west at the same spacing and altitude above the surface. The anomaly, which is on the high-standing side of a major fault, was interpreted previously as arising from either an increase of magnetization within a volcanic tuff or a small intrusive feature. Ground magnetic traverses were run to locate the ground maximum, and to delineate anomalies in a traverse that crosses the ground maximum and the nearby fault. Both air and ground anomalies were analyzed using geologic data from surface mapping and drill holes, and magnetic property data from drill holes. The anomaly is caused by contributions from at least three sources. The elevated topography gives a terrain effect since the altitude is decreased between the airplane and exposed Topopah Spring Member of the Paintbrush Tuff. Ground anomalies 300 m (1,000 ft) south of the air maximum indicate either an increase in magnetization or the presence of a small intrusive body. Finally, there is an increase in magnetic influence from the nearby Solitario Canyon fault.

INTRODUCTION

Studies of air and ground magnetic surveys by Bath and Jahren (1984) have provided structural information at and near the potential site for storage of radioactive waste at Yucca Mountain near the southwestern border of the Nevada Test Site for the Nevada Nuclear Waste Storage Investigations project of the U.S. Department of Energy. The air data consist of a high-altitude survey at a barometric elevation of 2,450 m (8,000 ft), and two low-altitude surveys about 120 m (400 ft) above the surface. One was flown east-west to investigate anomalies along major faults striking about north-south, and one was flown north-south to investigate anomalies trending across the site striking about east-west. The general distribution of contours in the two low-altitude surveys is similar, and differences in detail become plausible after considering difference in position and direction of flight paths. However, a prominent anomaly of 290 nT was detected on one north-south line that was not revealed on nearby east-west lines. It is located on figure 18 of Bath and Jahren (1984), and on figure 1 of this report.




Figure 1.--Residual aeromagnetic map of southern part of the potential repository (shaded) at Yucca Mountain showing prominent maximum of 290 nT that was detected on air traverse B. Also shown are air traverses A, C, and D; and maxima (triangles) and minima (inverted triangles) located on these and other flight traverses.

Yucca Mountain is underlain by a thick sequence of ash-flow tuffs and tuffaceous sediments, and subordinate amounts of lava flow and flow breccia. The volcanic rocks are of Tertiary age and attain a combined thickness of more than 1,829 m (6,000 ft). Pre-Tertiary rocks consist of sedimentary rocks with the possibility of igneous intrusions. A positive anomaly in the high-altitude aeromagnetic survey over exposures of strongly magnetized argillite of the Eleana Formation (Mississippian-Devonian Age) at Calico Hills extends westward 20 km into the site area where interpretations give an argillite thickness of 800 m (2,625 ft) at a depth of 2,250 m (7,400 ft). The high magnetite content of the argillite is not typical of the region, and was probably introduced by the heating effects of an underlying pluton. Pairs of positive and negative anomalies in the low-altitude aeromagnetic data are interpreted as major nearby faults that do not cross the site area.

This report was prepared to present 1982 and 1983 measurements of ground magnetic anomalies that were made to locate the position of the 290 nT anomaly on the ground, and to investigate the effects of magnetized geologic structures that could produce the anomaly. Bath and Jahren (1984) suggested the anomaly could be explained by either an increase of magnetization within normally magnetized ash flows or the presence of a small intrusive body. The suspected ash flows are the Topopah Spring Member of the Paintbrush Tuff, which is exposed at the surface, and the Bullfrog Member of the Crater Flat Tuff, which is at a depth of about 600 m (1,950 ft) below the surface. An investigation of flight records reveals another possibility. Radio altimeter measurements show an abrupt decrease in the flight altitude across the anomaly source, and this decrease in effective depth to the Topopah Spring Member will produce an increase in anomaly amplitude.

System Of Magnetic Units

All magnetic units are given in the International System of units (SI). Conversions to the older electromagnetic units (emu) are given in the following table:

Quantity	SI units	Equivalent unit (in emu)
Magnetic field	Nanotesla (nT)	1 nT = 1 gamma = 10^{-5} oersted
Magnetization	Ampere/meter (A/m)	1 A/m = 10^{-3} gauss

Acknowledgments

Several geologists and geophysicists have contributed to this report and their assistance is gratefully acknowledged. In particular, thanks go to R. B. Scott and R. W. Spengler for discussing relations of magnetic anomalies to the geologic features of Yucca Mountain, and Erick Esham for assistance in surveying locations of ground magnetic traverses. J. G. Rosenbaum provided the magnetic property data and wrote the three-dimensional forward program used in the model studies.

OBSERVED, RESIDUAL, AND SMOOTHED ANOMALIES

The observed data recorded by a magnetometer during an aeromagnetic or ground magnetic survey consist of the anomalies from the geologic features being studied plus the combined effects of the undisturbed geomagnetic field, magnetized sources deep within the Earth's crust, and man-made objects near the surface. Residual anomalies are those that remain after the Earth's field and effects of deep sources and man-made objects are removed from the observed data. The change in the Earth's field was eliminated from aeromagnetic data by removing the International Geomagnetic Reference Field (Barraclough and Fabiano, 1978), and from ground magnetic data by removing increases of 5.64 nT/km northward and 1.72 nT/km eastward. Effects of deep crustal sources were mostly eliminated by adjustment of observed data to an assumed zero field near Mercury in the southeastern corner of the Nevada Test Site (Bath and others, 1983). The zero field is the average value measured over a large area of nonmagnetic sedimentary rocks that are assumed to extend to great depths. Observed ground anomalies in areas near drill casing and other iron and steel objects are considered unreliable and therefore omitted from the data.

Residual anomalies were compiled for four air traverses and five ground traverses in the vicinity of the prominent aeromagnetic anomaly, and locations of traverses are given on the topographic map of figure 2. Figure 3 gives continuous measurements of anomalies and altimeter records for air traverses A, B, C, and D located on figures 1 and 2. The prominent anomaly is on traverse B. Ground anomalies were measured at 3-m (10-ft) intervals and are shown on figure 4 for traverse A82, figure 5 for traverse A83, figure 6 for traverse B83, and figure 7 for traverse C83. Ground traverse H82 is shown by Bath and Jahren (1984) on their figure 20. Ground anomalies measured close to magnetized rock have very irregular shapes, and a severe method of smoothing was used to convert them to a form resembling air anomalies. Each traverse was smoothed by continuation upward 122 m (400 ft) by the method of Henderson and Zietz (1949), and the resulting values were multiplied by a constant to restore the average value at ground level.

MAGNETIC PROPERTIES

The average total magnetization of a uniformly magnetized rock mass, denoted as the vector \vec{J}_t is defined as the vector sum of the induced magnetization, \vec{J}_i , and remanent magnetization, \vec{J}_r :

$$\vec{J}_t = \vec{J}_i + \vec{J}_r$$

Air and ground magnetic surveys will commonly detect an ash or lava flow when its average total magnetization is equal to or greater than 0.05 A/m (Bath, 1968). Therefore, units having intensities less than 0.05 A/m are herein designated nonmagnetic; and those having greater intensities are herein arbitrarily designated as either weakly, moderately, or strongly magnetized as defined by the following limits:

nonmagnetic \leq 0.05 A/m

0.05 A/m < weakly magnetized < 0.50 A/m

0.50 A/m < moderately magnetized < 1.50 A/m

1.50 A/m < strongly magnetized




Figure 2.--Topographic map of area west of Yucca Mountain showing air traverses A, B, C, and D; and ground traverses A82, H82, A83, B83, and C83. Traverse distances are in meters. Triangles give locations of anomaly maxima: 290 nT on air traverse B, and 584 nT on ground traverse C83. Contour interval is 20 feet.




Figure 3.--Residual anomalies and altimeter records for air traverses A, B, C, and D. The altimeter gives distance from airplane to ground surface in feet.




Figure 4.--Section along ground traverse A82 showing relation of residual and smoothed anomalies to volcanic strata, computed effects of Eleana Formation, and drill holes USW G-3, USW H-3, and USW H-5 along the crest of Yucca Mountain. Geologic symbols Tpc, Tpt, Tcb, and Tct are explained in the text.




Figure 5.--Profile of residual and smoothed anomalies along ground traverse A83 showing very irregular shapes with a thick zone of strongly magnetized rock extending from 625 to 740 m (2,051 to 2,428 ft). (See Fig. 4 for explanation of symbols.)




Figure 6.--Profile of residual and smoothed anomalies along ground traverse B83. (See fig. 4 for explanation of symbols.)




Figure 7.--Profile of residual and smoothed anomalies along ground traverse C83 showing a smoothed maximum of 584 nT at 707 m (2,320 ft). (See fig. 4 for explanation of symbols.)

Total magnetizations varying from nonmagnetic to strongly magnetic and of both normal and reversed polarities, were found in drill core samples from geologic exploration holes and surface samples in the Yucca Mountain area (Rosenbaum and Snyder, 1985). Large changes in magnetic intensity occur both laterally and vertically within the volcanic ash-flow sheets. Average magnetizations were determined for units mapped by Lipman and McKay (1965) and Scott and Bonk (1984), and penetrated in drill holes USW G-1 (Spengler and others, 1981), USW G-2 (Maldonado and Koether, 1983), and USW G-3 and USW GU-3 (Scott and Castellanos, 1984). The magnetic intensity values suggest the following eight units as possible anomaly producers in the vicinity of major faults in the Yucca Mountain area:

- Rainier Mesa Member of the Timber Mountain Tuff (Tmr)
- Tiva Canyon Member of the Paintbrush Tuff (Tpc)
- Pah Canyon Member of the Paintbrush Tuff (Tpp)
- Topopah Spring Member of the Paintbrush Tuff (Tpt)
- Bullfrog Member of the Crater Flat Tuff (Tcb)
- Tram Member of the Crater Flat Tuff (Tct)
- Lava flow and flow breccia (Tfb) Lava and flow breccia (Tll)

Considerations of thicknesses, and lateral extent of units has narrowed the eight to the Tiva Canyon, Topopah Spring, Bullfrog, and Tram Members. Their average magnetic properties and thicknesses are given in table 1. Modelling studies by Bath and Jahren (1984) showed a close resemblance between observed aeromagnetic anomalies in the Yucca Mountain area and theoretical anomalies computed for the faulted Topopah Spring Member. Their study designated the Topopah Spring Member as the most likely primary source of aeromagnetic anomalies.

Estimate of Magnetization

The method of estimating total magnetization by Smith (1961, equation 2.7) has been modified and applied to ground magnetic anomalies arising from near surface rocks in the NTS area by Bath and others (1983) and Bath and Jahren (1984). The estimates are based on the irregular and abrupt changes in anomaly amplitudes and shapes found in many ground traverses, and on the method of estimating depths to anomaly sources by Vacquier and others (1951). It is thus possible to use anomaly amplitudes to give minimum estimates of total magnetization within the following limits:

nonmagnetic \leq 15 nT

15 nT < weakly magnetized < 150 nT

150 nT < moderately magnetized < 450 nT

450 nT < strongly magnetized

Table 1.--Magnetic properties and thicknesses of four units that were penetrated in three holes drilled on Yucca Mountain. These are the most likely sources of magnetic anomalies

Unit	Drill hole	Magnetic polarity	J_t (A/m)	Thickness (m)	Comments
Tpc	USW G-3	Reversed	0.9	103	Entire unit
Tpt	USW G-1	Normal	1.3	335	Entire unit
Do.	do.	do.	0.7	169	upper part
Do.	do.	do.	2.0	166	lower part
Do.	USW G-2	do.	1.4	285	Entire unit
Do.	do.	do.	0.7	102	upper part
Do.	do.	do.	1.7	183	lower part
Do.	USW G-3	do.	1.2	272	Entire unit
			<u>1.3</u>	<u>297</u>	Average for entire unit
Tcb	USW G-1	Normal	1.0	130	Entire unit
Do.	USW G-2	do.	0.2	128	Entire unit (altered)
Do.	USW G-3	do.	3.0	182	Entire unit
			<u>1.4</u>	<u>147</u>	Average for entire unit
Tct	USW G-1	Reversed	1.2	268	Entire unit
Do.	do.	do.	2.0	142	upper part
Do.	do.	do.	0.1	126	lower part (altered)
Do.	USW G-2	do.	0.2	128	Entire unit (altered)
Do.	USW G-3	do.	1.8	369	Entire unit
			<u>1.1</u>	<u>255</u>	Average for entire unit

Estimates of magnetization based on ground anomalies range from weak to strong for near surface portions of rock units mapped along the crest and west of Yucca Mountain. Figure 2 shows the ground traverses, figure 8 shows the geologic units, and figures 4, 5, 6, and 7 show the anomalies.

Like the laboratory measurements of magnetic properties, estimates from ground magnetic anomalies indicate changes in magnetic intensity within the welded tuffs. For example, a lateral change from weak to moderate magnetization is revealed for the Tiva Canyon Member by abrupt changes in the residual anomalies shown in traverse A82 (fig. 4) which extend for a total distance of 6,860 m (22,500 ft) along the crest of Yucca Mountain. Amplitudes average about 50 nT for the first 4,900 m (16,075 ft) and about 200 nT for the remaining 1,960 m (6,425 ft). This northward increase is in the upper part of the member. As shown on figure 4, the anomaly datum is 115 m (375 ft) above the base of the member in drill hole USW G-3 (Scott and Castellanos, 1984), and 150 m (490 ft) above the base of the member of drill hole USW H-5 (Bentley and others, 1983).

Most of the irregular anomalies on traverses A83 (fig. 5) and B83 (fig. 6) and on the eastern part of traverse C83 (fig. 7), are above exposures of the Topopah Spring Member and indicate moderate magnetizations. A few isolated anomalies have amplitudes greater than 450 nT and indicate strongly magnetized rock. A prominent example is the thick zone that extends from 625 m (2,050 ft) to 740 m (2,428 ft) on traverse A83 (fig. 5). Estimates based on anomaly shapes (Vacquier and others, 1951) mark out a mass of strongly magnetized rock about 115 m (378 ft) wide with its top about 30 m (100 ft) below the surface.

ANOMALY INTERPRETATIONS

Aeromagnetic surveys in the Nevada Test Site region were flown close enough to the surface to detect anomalies produced by major faults (Bath, 1976, Bath and others, 1983; Bath and Jahren, 1984). While the terrain clearances shown on figure 3 average about 122 m (400 ft) for traverses A-A' and B-B' and about 183 m (600 ft) for traverses C-C' and D-D', there are some notable deviations. One obvious example is the interval decrease on traverse B-B' beneath the prominent aeromagnetic anomaly. Barometric altimeter readings indicate the elevation of the aircraft was about constant, and therefore, the decrease in terrain clearance reflects changes in topographic relief beneath the flight path. There is a small westward bend in the northward trend of Yucca Mountain, and figure 2 shows that the flight path crosses this feature.

A qualitative interpretation suggests the decrease in interval from airplane to surface is the source of the 290 nT-air anomaly. A positive anomaly would be expected in measurements closer to the surface of a normally magnetized ash flow; and there are approximate correlations of positions of anomaly maximum (fig. 1), westward bend of Yucca Mountain (fig. 2), and minimum interval (fig. 3). However, a depth estimate from anomaly B-B' of figure 3 places the source too deep to be explained by effects of terrain clearance alone. Some other source, or sources, must be present and have a contributing effect that distorts the shape of the anomaly that gives the misleading depth estimate. Quantitative interpretations were therefore undertaken to investigate other possible sources. The ground traverses proved




Figure 8.--Geologic map of western part of Yucca Mountain showing exposures of volcanic units (symbols are explained in text) and alluvium and colluvium (QTac); and the mapped trace of the Solitario Canyon fault that trends northward across the central part of the map. Also shown are locations of 290-nT air maximum and 584-nT ground maximum. Geology is from Scott and Bonk (1984).

to be an important part of the studies. They showed the effects of other possible sources, and not the effect of a decrease in interval between airplane and ground surface.

Aeromagnetic Anomalies

The pair of parallel aeromagnetic anomalies that trend north-south along the west side of Yucca Mountain (Bath and Jahren, 1984, figs. 12, 15, and 17) are explained as the edge effects of volcanic tuff displayed by the Solitario Canyon fault. The anomalies are negative over the low-standing side and positive over the high-standing side of the fault. The prominent anomaly on traverse B-B' of figure 3 is on the high-standing side, but its maximum of 290 nT is 135 nT greater than maximum on a east-west flight line that intersects B-B' to the south. Positive anomalies also are found on lines flown north-south near traverse B-B'. The broad positive anomalies on traverses A-A', C-C', and D-D' (fig. 3) were interpreted by Bath and Jahren as arising from a strongly magnetized source, possibly altered Eleana Formation like that at Calico Hills (Baldwin and Jahren, 1982), at a depth of about 2,250 m (7,400 ft) underlying the volcanic rocks penetrated in drill holes at Yucca Mountain.

As shown on figure 9, the 290 nT anomaly can be explained by three sources: a dipole or a finite prism (Source A) that represents an increase in magnetization, and an infinite vertical prism (Source B) that represents a small intrusive. This assumes no anomaly arising from the effect of terrain clearance. Table 2 gives the total magnetization of the sources, and table 3 gives the amplitudes of the computed anomalies.

The increase in magnetization indicated by the magnetic dipole and Source A is 625 m (2,050 ft) below the air datum, or at a depth of 549 m (1,800 ft). This places the source below the Topopah Spring Member of the Paintbrush Tuff and within the Bullfrog Member of the Crater Flat Tuff. The computed direction of magnetization in table 2 is about midway between normal and reverse, and suggests a body having a reversed thermoremanent magnetization with a substantial normal component of isothermal remanent magnetization. This is a significant deviation from the known normal polarity of the Bullfrog Member and from the intensity of 32.86 A/m, which is too large for the member. The reversed polarity would be more reasonable for a basaltic dike, and even the high intensity of 8.32 A/m is a possibility.

The models offer an opportunity to investigate effects of positioning the airplane closer to the ground surface. As shown in table 3, reducing the elevation by 61 m (200 ft) increases the amplitude by only about 70 nT, or about half of the 135 nT difference between maximum of 290 nT on north-south traverse B-B' of figure 1 and the maximum of 155 nT on the nearby east-west traverse (fig. 12 of Bath and Jahren, 1984). This indicates the sources are too deep for this change in elevation to explain the prominent aeromagnetic anomaly.

Ground Magnetic Anomalies

Another source for the prominent air anomaly is suggested by the positive values of smoothed ground anomalies that increase abruptly eastward with increasing distance from the Solitario Canyon fault. Figures 2 and 8 show the relationships between positions of traverses and faults, and also show that the small westward bend in the northward trend of Yucca Mountain is accompanied by a similar bend in the Solitario Canyon fault. The positive




Figure 9.--Section along part of air traverse B-B' showing close comparisons between the observed anomaly and anomalies computed for a dipole, source A, and source B. Source A represents an increase in magnetization within a vertical prism with length = width = 305 m (1,000 ft) and thickness = 152 m (500 ft). Source B represents a small intrusive with length = width = 305 m (1,000 ft) and thickness = infinity.

Table 2.--Total magnetizations for models of figure 9

Model	Intensity (A/m)	Azimuth ¹ (degrees)	Inclination ² (degrees)
Dipole ³	14,500	16	145
A	32.86	16	145
B	8.32	16	145

¹ Measured clockwise from north.

² Measured down from a northward horizontal axis.

³ Small cube with volume = $2.83 \times 10^4 \text{ m}^3$.

Table 3.--Aeromagnetic anomalies computed for models of figure 9

AMPLITUDE OF RESIDUAL AEROMAGNETIC ANOMALY						
Model	Depth to top		Maximum	Minimum	Total	Difference
	(m)	(ft)	(nT)	(nT)	(nT)	in total (nT)
Dipole	625	2,050 ¹	261	-36	297 ¹	71
Dipole	686	2,250 ¹	199	-27	226	
Source A	549	1,800	259	-38	297 ²	69
Source A	610	2,000	199	-29	228	
Source B	427	1,400	279	-18	297 ²	66
Source B	488	1,600	219	-12	321	

¹ Depth is to center of dipole.

² The anomaly shown in figure 9.

values average about 150 nT along traverse A83-A83' which is about parallel to and 100 m (300 ft) east of the fault. The average values increase to about 350 nT on traverse B83-B83' which is 120 m (400 ft) east of traverse A83-A83' and nearer the position of the air maximum. A similar increase is expected in the air anomalies. Air traverse B-B' was flown along a straight line and its distance from the fault increases in the area of the air maximum.

The ground data measured beneath air traverse B-B' supports an interpretation of change in terrain clearance, but not an interpretation of an increase in magnetization at a depth of 549 m (1,800 ft) or a basaltic dike at depth of 427 m (1,400 ft). The smoothed anomalies of traverse A83-A83' do not show a prominent anomaly having the amplitude to explain the air anomaly or the long wavelength to explain sources at depth. Almost all the unsmoothed anomalies have the short wavelengths that are produced by changes in rock magnetization in narrow zones near the surface. The thick zone of strongly-magnetized rock (fig. 5) is about 300 m (1,000 ft) south of the aeromagnetic maximum, and it is described in the section "Estimate of Magnetization". Traverse C83-C83' was measured in an east-west direction to delineate effects of the thick zone and the Solitario Canyon fault.

The smoothed anomaly of traverse C83-C83' was then analyzed to investigate the source of the thick zone and its effects at the elevations of the aeromagnetic anomalies. There is reliable information for the geology and magnetic properties of the flows displaced by the fault, and a three-dimensional forward program was used to model the high- and low-standing components of the fault and two possible sources for the thick zone.

As shown on figure 10, the anomaly over the fault has a total amplitude of 822 nT and can be modeled satisfactorily by the combined effects of faulted components of the Topopah Spring and Tiva Canyon Members, and the effects of either an increase in magnetization within the Topopah Spring Member (Source C) or a strongly magnetized intrusive body (Source D). Tuff magnetizations, estimated from anomaly amplitudes, are moderate near the fault and less than those from the drill-core samples of table 1, probably due to weathering; and the magnetizations of Sources C and D are strong. Aeromagnetic anomalies were computed from the models and compared with positive amplitudes measured 61 m (200 ft) and 122 m (400 ft) above the surface. Table 4 gives the magnetization directions and intensities for the models, and table 5 gives the amplitudes of computed anomalies.

The wide zone is interpreted as either a very large increase of magnetization within the Topopah Spring Member or a small intrusive body. The magnetization increase is designated more likely because abrupt changes are common within tuffs of the site area. Changes in drill-core samples are reported by Rosenbaum (1985), and other zones of strong magnetization were estimated from the anomaly amplitudes along ground traverse A83 of figure 5. Geological evidence in the area of the anomaly also argues against the possibility of a near surface intrusive. No change in appearance of surface rocks or effects of heating was noted. However, basalt float has been found at a lower elevation and about 1/2 mi south of the anomaly (R. B. Scott, USGS, oral commun., 1984).

As shown on figure 10 and in table 5, the wide zone as Source C will produce a significant anomaly of 86 nT at 122 m (400 ft) above the surface. This is the difference between the maximum effects of the fault, 186 nT; and the maximum effects of the fault plus Source C, 272 nT. A decrease in 61 m (200 ft) in flight elevation (table 5) increases computed maxima by 54 nT for the fault, 89 nT for Source C, and 101 nT for Source D.




Figure 10.--Section across the Solitario Canyon fault on ground traverse C83-C83' (projected) showing close comparisons between the observed anomaly and summations of anomalies computed for horizontal prisms that represent the Tiva Canyon Member (Tpc) and the Topopab Spring Member (Tpt); and for vertical prisms that represent an increase in magnetization (Source C) or magma pipe (Source D). The horizontal prisms extend along the fault strike for a distance of 3,048 m (10,000 ft). Source C has a width = 198 m (650 ft) along the traverse and its north-south length = depth extent = 152 m (500 ft). Source D has the same length and width as Source C, but its depth extent is infinite.

Table 4.--Total magnetizations of component parts of models of figure 10

Component	Intensity (A/m)	Azimuth ¹ (degrees)	Inclination ² (degrees)
Tiva Canyon tuff	0.75	167	-38
Topopah Spring tuff	0.75	326	62
Source C	3.73	326	62
Source D	3.26	0	62.5

¹ Measured clockwise from north.

² Measured down from a northward horizontal axis.

Table 5.--Ground magnetic and aeromagnetic anomalies computed for models of figure 10

AMPLITUDE OF RESIDUAL MAGNETIC ANOMALY					
Component	Type	Maximum (nT)	Difference in maximum (nT)	Minimum (nT)	Total (nT)
Faulted components plus either source C or D	Ground	584		-238	822 ¹
Faulted components only	Air ²	240		-66	306 ³
			54		
Faulted components only	Air ⁴	186		-11	197
Faulted components plus source C	Air ²	361		-84	445
			89		
Faulted components plus source C	Air ⁴	272		-20	292 ⁵
Faulted components plus source D	Air ²	413		-41	454
			101		
Faulted components plus source D	Air ⁴	312		13	299

¹ The ground anomaly shown on figure 10.

² 61 m (200 ft) above ground traverse.

³ The air anomaly shown for fault only on figure 10.

⁴ 122 m (400 ft) above ground traverse.

⁵ The air anomaly shown on figure 10.

DISCUSSION

The several single sources investigated did not give a satisfactory explanation of the prominent air anomaly. We believe the anomaly was produced by a combination of sources that include the magnetic effects of (1) reducing the interval from air datum to ground surface, (2) an increase of magnetization within the Topopah Spring Member of the Paintbrush Tuff, and (3) an increase in distance between the air datum and the Solitario Canyon fault. These interpretations do not require a significant change in the present understanding of the geology and magnetic properties of rocks within the area of the potential waste repository site. The presence of a small intrusive body would introduce a new concept, but the data now available do not favor this interpretation.

REFERENCES CITED

- Baldwin, M. J., and Jahren, C. E., 1982, Magnetic properties of drill core and surface samples from the Calico Hills area, Nye County, Nevada: U.S. Geological Survey Open-File Report 82-536, 27 p.
- Barracough, R. R., and Fabiano, E. B., 1978, Grid values and charts for the International Geomagnetic Reference Field 1975: U.S. Geological Survey Report PB-276 630, 139 p.: available only from U.S. Department of Commerce, National Technical Information Service, Springfield, VA, 22161.
- Bath, G. D., 1968, Aeromagnetic anomalies related to remanent magnetization in volcanic rock, Nevada Test Site, Nevada, in Nevada Test Site: Geological Society of America Memoir 110, p. 135-146.
- Bath, G. D., 1976, Interpretation of magnetic surveys in intermontane valleys at Nevada and southern New Mexico: U.S. Geological Survey Open-File Report 76-440, 36 p.
- Bath, G. D., Jahren, C. E., Rosenbaum, J. G., and Baldwin, M. J., 1983, Magnetic investigations, in Geologic and geophysical investigations of Climax stock intrusive, Nevada: U.S. Geological Survey Open-File Report 83-377, p. 40-77.
- Bath, G. D., and Jahren, C. E., 1984, Interpretations of magnetic anomalies at a potential repository site located in the Yucca Mountain area, Nevada Test Site: U.S. Geological Survey Open-File Report 84-120, 53 p.
- Bentley, C. B., Robison, J. H., and Spengler, R. W., 1983, Geohydrologic data for test well USW H-5, Yucca Mountain area, Nye County, Nevada: U.S. Geological Survey Open-File Report 83-853, 34 p.
- Henderson, R. G., and Zietz, Isidore, 1949, The upward continuation of anomalies in total magnetic intensity fields: Geophysics, v. 14, no. 4, p. 517-535.
- Lipman, P. W., and McKay, E. J., 1965, Geologic map of the Topopah Spring SW quadrangle, Nye County, Nevada: U.S. Geological Survey Geologic Quadrangle Map GQ-439, scale 1:24,000.
- Maldonado, Florian, and Koether, S. L., 1983, Stratigraphy, structure, and some petrographic features of Tertiary volcanic rocks at the USW G-2 drill hole, Yucca Mountain, Nye County, Nevada: U.S. Geological Survey Open-File Report 83-732, 83 p.
- Rosenbaum, J. G., and Snyder, D. B., 1985, Preliminary interpretation of paleomagnetic and magnetic property data from drill holes USW G-1, G-2, GU-3, and VH-1 and surface localities in the vicinity of Yucca Mountain, Nye County, Nevada: U.S. Geological Survey Open-File Report 85-49, 73 p.
- Scott, R. B., and Bonk, Jerry, 1984, Preliminary geologic map of Yucca Mountain, Nye County, Nevada with geologic sections: U.S. Geological Survey Open-File Report 84-494, scale 1:12,000.

Scott, R. B., and Castellanos, Mayra, 1984, Stratigraphic and structural relations of volcanic rocks in drill holes USW GU-3 and USW G-3 Yucca Mountain, Nye County, Nevada: U.S. Geological Survey Open-File Report 84-491, 121 p.

Smith, R. A., 1961, Some theorems concerning local magnetic anomalies: Geophysical Prospecting, v. 9, p. 399-410.

Spengler, R. W., Byers, F. M., Jr., and Warner, J. B., 1981, Stratigraphy and structure of volcanic rocks in drill hole USW-G1, Yucca Mountain, Nye County, Nevada: U.S. Geological Survey Open-File Report 81-1349, 50 p.

Vacquier, Victor, Steenland, N. C., Henderson, R. G., and Zietz, Isidore, 1951, Interpretation of aeromagnetic maps: Geological Society of America Memoir 47, 151 p.

```

C      itest = Error flag
C      1 = Next polygon was found
C      0 = No more polygons
C      -1 = A value in tree is less than 0.
C
C      Note: ngon and next are pointers to positions in the
C      topology arrays.
C
C *****
C      Subroutine walk(next,ngon,itest)
C      Common /topology/ info(100),iupper(100),idown(100),
C      & illeft(100),iright(100)
C      Common /commands/ nmax,epsiln,delin,delout
C
C      next=0
C      If (ngon.ge.1.and.ngon.le.nmax) then
C      npnt=ngon
C      next=idown(ngon)
C      If (next.GT.0) then
C      itest=1
C      Else If (next.EQ.0) Then
C      next=iright(ngon)
C      If (next.GT.0) Then
C      itest=1
C      Else If (next.EQ.0) Then
C      itest2=0
C      i=1
C      Do 20 while(itest2.EQ.0)
C      npnt=iupper(npnt)
C      If (npnt.GT.0) Then
C      next=iright(npnt)
C      If (next.GT.0) Then
C      itest=1
C      itest2=1
C      Else If (next.EQ.0) Then
C      itest2=0
C      Else
C      itest2=-1
C      itest=-1
C      print *, ' Error, iright was out of range (iright<=0)'
C      End If
C      Else If (npnt.EQ.0) Then
C      next=0
C      itest=0
C      itest2=1
C      Else
C      itest=-1
C      itest2=-1
C      print *, ' Error, lup of ',npnt,' less than zero'
C      End If
C      i=i+1
C      If (i.GT.nmax) Then
C      print *, ' Error, i (counter) was >',nmax
C      ites=-1
C      itest2=-1
C      End If
C      End Do
C      Else
C      print *, ' Error, iright of ',npnt,' was less than zero'
C      itest=-1
C      End If
C      Else
C      print *, ' Error, idown of ',ngon,' was less than zero'

```

```

      itest=-1
C      End If
C      Else
C      print *, ' Error, NGON passed to WALK out of range'
C      itest=-1
C      End If
C
C      Return
C      End
C+
C *****
C wcbp - World coordinate byte packing. Converts a world co-
C ordinate pair into the (hexadecimal) code required for
C the Envision terminal. See wcbp.inf and Envision manual.
C
C      Author: Robert Simpson, USGS, Menlo Park, CA, 10/83.
C-
C *****
C      Character*5 Function wcbp(i,j)
C      Character blank*1
C      Parameter (imax=16284,jmax=16284)
C      Parameter (imin=0,jmin=0)
C      Parameter (blank=' ',iblack=ichar(' '))
C
C - Force i,j into bounds...
C      iin=min(imax,max(imin,i))
C      jin=min(jmax,max(jmin,j))
C - Get hi and lo bytes and offset with blank...
C      ilo=mod(iin,64)+iblack
C      imed=iin/64+iblack
C      jlo=mod(jin,64)+iblack
C      jmed=jin/64+iblack
C - Put bytes together...
C      wcbp=blank//char(imed)//char(ilo)//char(jmed)//char(jlo)
C
C      Return
C      End
C+
C *****
C wrhead - Writes the header for a standard grid (new version).
C
C      Author: Robert W. Simpson, USGS, Menlo Park, CA.
C-
C *****
C      Subroutine wrhead(unit,id,pgm,ncol,nrow,nz,
C      &xo,dx,yo,dy,iproj,cm,b1,itest)
C      Character id*56,pgm*8
C      Integer unit
C
C      itest=1
C      Write (unit)id,pgm,ncol,nrow,nz,xo,dx,yo,dy,iproj,cm,b1
C
C      Return
C      End
C+
C *****
C wrtgrd - WRITe GRID - Writes out wrkgrd as a Denver standard
C grid file to 'name'.
C-
C *****
C      Subroutine wrtgrd(name,itest)
C      Common /work/wrkgrc:250000)
C      Common /gridspecs2/id2,pgm2,nc2,nr2,nz2,xo2,dx2,yo2,dy2,

```

```

&iproj2,cm2,b12
Character name*8#,id2*56,pgm2*8
C
  itest=1
  pgm2='polygon'
  dummy=#.#
  Open (11,file=name,status='new',form='unformatted')
  Call wrhead(11,id2,pgm2,nc2,nr2,nz2,xo2,dx2,yo2,dy2,iproj2,
&cm2,b12,itest)
  If (itest.GE.#) Then
    Do 2# j=1,nr2
      Write (11)dummy,(wrkgrd(1+(j-1)*nc2),f=1,nc2)
    2# Continue
  End If
  Close (11)
C
1## Continue
Return
End
C+
C *****
C wrtmod - Writes out a Polyogn model file.
C
C
C-
C *****
C
  Subroutine wrtmod(itest)
  Common /topology/info(1##),lupper(1##),ldown(1##),
&illeft(1##),ilright(1##)
  Common /screenloc/ntotal,numply(1##),xscr(1##,1##),yscr(1##,1##)
  Common /parameter /parm(1##,1#)
  Common /labels/label
  Common /commands/nmax,eps!in,delin,delout
  Common /junk/ngbtop,jnktop(1##),ngbloc,jnkloc(1##)
  Common /names/grdnam,modnam,modgrd
  Common /gr idspecs/id,pgm,nc,nr,nz,xo,dx,yo,dy,iproj,cm,b1
  Common /original/iwcorg,jwcorg,nxorgp,nyorgp
C
  Character label(1#)*15
  Character id*56,pgm*8,grdnam*8#,modnam*8#,modgrd*8#
C
  Open (1#,file=modnam,status='unknown',form='formatted',
&carriagecontrol='list')
C
C - Write out the current parms assigned to standard grid header
C information.
C
  write (1#,5) id,pgm,nc,nr,nz
C5 format(x,a56,x,a8,3(x,i3))
C write (1#,1#) xo,dx,yo,dy
C1# format(4(x,e15.8))
C write (1#,15) iproj,cm,b1
C15 format(x,i3,2(x,e15.8))
C
C - Write out the number of polygons in the model and the
C labels assigned to the parm arrays.
C
  Write (1#,2#)ntotal
  2# Format (x,i3)
  Write (1#,25)(label(k),k=1,5)
  Write (1#,25)(label(k),k=6,1#)
  25 Format (5(x,a15))
C
C - Write out the polygon number, the number of points in this

```

```

C polygon, the grid locations and the polygon parms.
C
  delxcn=dx/nxorgp
  delycn=dy/nyorgp
  Do 6# i=1,nmax
  npoly=info(i)
  If (npoly.GT.#) Then
    nbrpts=numply(npoly)
    Write (1#,33)npoly,nbrpts
  33 Format (2(x,i3))
    Write (1#,35)(parm(npoly,kk),kk=1,5)
    Write (1#,35)(parm(npoly,kk),kk=6,1#)
  35 Format (5(x,e15.8))
    Do 4# j=1,nbrpts
      xgrid=xo+((xscr(npoly,j)-iwcorg)*delxcn)
      ygrid=yo+((yscr(npoly,j)-jwcorg)*delycn)
    Write (1#,45)xgrid,ygrid
  45 Format (2(x,e16.8))
  4# Continue
    End If
  6# Continue
C
C - Write out the topology structure
C
  Do 7# ndpstn=1,nmax
  Write (1#,65)info(ndpstn),lupper(ndpstn),ldown(ndpstn),
&illeft(ndpstn),ilright(ndpstn)
  65 Format (5(x,i3))
  7# Continue
C
  Close (1#)
  itest=1
  Return
  End
C+
C *****
C wrtmsg - Writes out a text string to the terminal.
C-
C *****
C
  Subroutine wrtmsg(text)
  Character text*(*)
C
  ileng=itlen(text)
  Write (6,1#)text
  1# Format (x,a(ileng))
C
  Return
  End
C+
C *****
C zomcom - Zoom command mode. Controls the function of zooming
C and unzooming in the grid.
C
C
C Options:
C
C c = Clear zoom stack
C d = Draw the grid using a zoom value selected from the
C stack.
C r = Recall and draw the grid using the zoom values
C currently pointed to in stack by nzoom.
C s = Select a subgrid (using mouse, cursor, grid
C coords.)
C u = Unzoom grid, draw full grid on screen (does nothing
C to zoom values)

```

```

C      h = Help message
C      q = Quit and return to Polygon command level
C-----
C *****
C Subroutine zomcom(ifirst,itest)
C Character ans*2,quest*88,zcom*2,first*1
C Common /subgrid/icmin,icmax,irmin,irmax,ncmin,ncmax,nrmin,nrmax
C Common /scale/xsc,ysc,xstart,ystart,xinit,yinit
C Common /zoom/izoom,izval,nzoom,ncminz(5),ncmaxz(5),nrminz(5),
C &nrmaxz(5)
C Common /scalefacts/iwcB,jwcB,nxpix,nypix,pxdim
C Common /original/iwcorg,jwcorg,nxorgp,nyorgp
C Common /flags/mcflag,votflg
C Common /mode/mdflag
C Common /colors/plyclr,black,white
C Common /fill/open,solid,filtyp
C Character pyclr*1,black*1,white*1,open*1,solid*1,filtyp*1
C Character mcflag*2,votflg*2
C
C      Call enhmsg('*** Zoom mode ***')
C
C - Testing of input variables goes here
C
C      zcom='H'
C      iflag=#
C      Do 1# while(iflag.EQ.#)
C      quest=' Zoom command (c/d/r/s/u/h/q)'
C      ival=iaquest(quest,zcom,'(a2)',-2)
C
C      If (zcom.EQ.'C') Then
C      Call askans(ans)
C      If (ans.EQ.'Y') Then
C      Call intzom
C      Call setbnd
C      Else If (ans.EQ.'Q') Then
C      iflag=-1
C      End If
C      Else If (zcom.EQ.'D') Then
C      If (nzoom.GE.1) Then
C      Call zompck(itest)
C      If (itest.GE.#) Then
C      Call displa
C      Call scaleg2sc(#)
C      Call clrsgd(#,itest)
C      xstart=ncmin
C      ystart=nrmin
C      Call setbnd
C      End If
C      If (mdflag.EQ.1.AND.itest.GE.#) Then
C      Call setclr(pyclr)
C      Call setfill(open)
C      Call drawwalk
C      Call setfill(solid)
C      End If
C      Else
C      Write (6,5#)
C      Format (/, ' Sorry, zoom stack is empty...')
C      End If
C      Else If (zcom.EQ.'S') Then
C      If (ifirst.EQ.1) Then
C      Call intscr
C      Call clrsgd(1,itest)
C      Call clrply

```

```

xsc=float(nxpix)
ysc=float(nypix)
iwcorg=iwcB
jwcorg=jwcB
nxorgp=nxpix
nyorgp=nypix
Call setbnd
End If
If (mdflag.EQ.1.AND.ifirst.EQ.1) Then
Call setclr(pyclr)
Call setfill(open)
Call drawwalk
Call setfill(solid)
End If
ifirst=#
2# Continue
Call asktyp(itype,itest)
If (itest.EQ.-1)Go To 3#
If (nzoom.EQ.5)Call zomstk(itest)
If (itest.EQ.-1)Go To 3#
If (itype.EQ.1) Then
Call getsub(itest)
Else If (itype.EQ.2) Then
Call asksub(itest)
End If
If (itest.EQ.1)izoom=1
If (itest.EQ.-1)Go To 2#
3# Continue
If (itest.GE.#) Then
Call displa
Call scaleg2sc(#)
Call clrsgd(#,itest)
xstart=ncmin
ystart=nrmin
Call setbnd
End If
If (mdflag.EQ.1.AND.itest.GE.#) Then
Call setclr(pyclr)
Call setfill(open)
Call drawwalk
Call setfill(solid)
End If
Else If (zcom.EQ.'R') Then
If (nzoom.GE.1) Then
Call zomrc1(itest)
If (itest.GE.#) Then
Call displa
Call scaleg2sc(#)
Call clrsgd(#,itest)
Call clrply
xstart=ncmin
ystart=nrmin
Call setbnd
End If
If (mdflag.EQ.1.AND.itest.GE.#) Then
Call setclr(pyclr)
Call setfill(open)
Call drawwalk
Call setfill(solid)
End If
Else
Write (6,5#)
End If
Else If (zcom.EQ.'U') Then

```



```

      Call unzcom(itest)
    Else If (zcom.EQ.'H') Then
      Call hlpzom
    Else If (zcom.EQ.'Q'.OR.(ival.EQ.-1)) Then
      Iflag=-1
    Else
      Call errmsg
    End If
    zcom='Q'
  1# End Do
C
  If (Iflag.EQ.-1)itest=-1
  Return
  End
C+
C *****
C zompck - Allows the picking of a selected zoom value from the
C zoom stack (if any exist).
C-
C *****
C Subroutine zompck(itest)
C Common /zoom/izoom,izval,nzoom,ncminz(5),ncmaxz(5),nrminz(5),
C &nrmaxz(5)
C Common /subgrid/icmin,icmax,lrmin,lrmax,ncmin,ncmax,nrmin,nrmax
C Character quest*8#
C
C call hlpzpk
C
C itest=#
C Do 1# while(itest.EQ.#)
C   quest=' Stack position of zoom value (1-5, # or // to quit)'
C   ival=iquest(quest,izval,'(12)',#)
C
C   If (izval.GE.1.AND.izval.LE.5) Then
C     If (ncminz(izval).EQ.#.AND.ncmaxz(izval).EQ.#.AND.
C       & nrminz(izval).EQ.#.AND.nrmaxz(izval).EQ.#) Then
C       Write (6,15)
C 15 Format (/, ' Error, zoom values for this position not set',/)
C     Else
C       ncmin=ncminz(izval)
C       ncmax=ncmaxz(izval)
C       nrmin=nrminz(izval)
C       nrmax=nrmaxz(izval)
C       itest=1
C     End If
C   Else If (izval.EQ.#.OR.ival.EQ.-1) Then
C     itest=-1
C   Else
C     Call errmsg
C   End If
C 1# End Do
C
C Return
C End
C+
C *****
C zomrc1 - Recalls the subgrid values from position nzoom
C in the zoom stack.
C-
C *****
C Subroutine zomrc1(itest)
C Common /gridspecs/id,pgm,nc,nr,nz,xo,dx,yo,dy,iproj,cw,b1
C Common /subgrid/icmin,icmax,lrmin,lrmax,ncmin,ncmax,nrmin,nrmax
C Common /zoom/izoom,izval,nzoom,ncminz(5),ncmaxz(5),nrminz(5),

```

```

&nrmaxz(5)
Character id*56,pgm*8
C
  If (nzoom.GE.1.AND.nzoom.LE.5) Then
    ncmin=ncminz(nzoom)
    ncmax=ncmaxz(nzoom)
    nrmin=nrminz(nzoom)
    nrmax=nrmaxz(nzoom)
    itest=1
  Else If (nzoom.EQ.#) Then
    itest=#
  Else
    itest=-1
  End If
C
  Return
  End
C+
C *****
C zomstk - Controls the release of the zoom stack. Called by
C subroutine zomcom (ZOOM COMMAND) if nzoom equals five.
C-
C *****
C b = Bottom of stack (oldest zoom values)
C c = Clear stack (zeroes out zoom stack)
C t = Top of stack (youngest zoom values)
C h = Help message
C q = Quit z.id return to ZOOM COMMAND level.
C
C nzoom = Pointer to current position in zoom stack
C # = if zoom stack is empty.
C otherwise ranges between 1 and 5.
C-
C *****
C Subroutine zomstk(itest)
C Common /zoom/izoom,izval,nzoom,ncminz(5),ncmaxz(5),nrminz(5),
C &nrmaxz(5)
C Character quest*8#,ans*2
C
C itest=#
C Call msgstk
C Call hlpstk
C ans='H'
C Do 1# while(itest.EQ.#)
C   quest=' What method of stack release (b/c/t/h/q)'
C   ival=iquest(quest,ans,'(a2)',#)
C
C   If (ans.EQ.'B') Then
C     itest=1
C     Do 2# i=1,4
C       ncminz(i)=ncminz(i+1)
C       ncmaxz(i)=ncmaxz(i+1)
C       nrminz(i)=nrminz(i+1)
C       nrmaxz(i)=nrmaxz(i+1)
C 2# Continue
C     ncminz(5)=#
C     ncmaxz(5)=#
C     nrminz(5)=#
C     nrmaxz(5)=#
C     nzoom=4
C   Else If (ans.EQ.'T') Then
C     itest=1
C     ncminz(5)=#
C     ncmaxz(5)=#
C     nrminz(5)=#

```

```

        nrmaxz(5)=#
        nzoom=4
    Else If (ans.EQ.'C') Then
        Call intzom
        itest=1
    Else If (ans.EQ.'Q'.OR.ival.EQ.-1) Then
        itest=-1
    Else If (ans.EQ.'H') Then
        Call hipstk
    Else
        Call errmsg
    End If
1# End Do
C
    Return
    End
C+
C*****
C zomstr - Stores the values of ncmn,ncmax,nrmin,nrmax, used for
C zooming, in the zoom stack at position nzoom.
C
C      See intzom.for for a description of the variables in
C      the zoom common block.
C*****
C Subroutine zomstr(itest)
C Common /zoom/izoom,izval,nzoom,ncminz(5),ncmaxz(5),nrminz(5),
C &nrmaxz(5)
C Common /subgrid/icmin,icmax,irmin,irmax,ncmin,ncmax,nrmin,nrmax
C
C If (nzoom.GE.#.AND.nzoom.LF.5) then
C   itest=1
C   nzoom=nzoom+1
C   ncmnz(nzoom)=ncmn
C   ncmaxz(nzoom)=ncmax
C   nrminz(nzoom)=nrmin
C   nrmaxz(nzoom)=nrmax
C Else If (nzoom.GE.5) Then
C   Call wrtmsg(' Zoom stack is full')
C   itest=-1
C End If
C
C Return
C End

```