

FFTF FSAR REFERENCE DOCUMENT

SECTION NO.	REFERENCE NO.	DOCUMENT NO.
2.6	023	JABE-WADCO 04

00000000112

**SUPPLEMENTARY GEOLOGIC
 INVESTIGATIONS FOR SEISMIC EVALUATION
 OF THE FFTF SITE NEAR RICHLAND
 WASHINGTON**

APPLIED TECHNOLOGY

Any further distribution by any holder of this document or of the data therein to third parties representing foreign interests, foreign governments, foreign companies and foreign subsidiaries of foreign divisions of U.S. companies should be coordinated with the Director, Division of Reactor Development and Demonstration, U.S. Energy Research and Development Administration.

Hanford Engineering Development Laboratory

Operated by the
**Westinghouse
 Hanford Company**

A Subsidiary of
 Westinghouse Electric
 Corporation

for the United States
 Energy Research and
 Development Administration
 Contract No. E(45-1)2170

**8603270141 851216
 PDR WASTE
 WM-10 PDR**

REC'D W/CTR DTG 8603270132 851216

FOR
DRAWINGS
SEE
APERTURE
CARD

July-Wadeo - 04

Sec. No.

2.6

Pl. No.

023

1. Geologic Map and Cross Sections
Along a portion of the Horn
Rapids Lineament Near
Richland Washington Plate III
2. Photographs of the Ringold Formation
on Bluffs East of the Columbia River Part 2, Plate 2
3. Geologic Map and Cross Sections Along
A portion of the Horn Rapids Lineament
Near Richland, Washington Plate IV
4. Geologic Map and Cross Sections Along
A portion of the Horn Rapids Lineament Near
Richland Washington Plate V
5. RECONNAISSANCE Geologic Map of Fast Flux Test
Facility Site and Vicinity. Part 2 Plate 1

76.99

Supplementary
Geologic Investigations for Seismic
Evaluation of the FFTF Site
Near Richland, Washington

Prepared for WADCO
of
Richland, Washington

February 1971

0000000115

SUPPLEMENTARY GEOLOGIC INVESTIGATIONS FOR
SEISMIC EVALUATION OF THE FFTF SITE
NEAR RICHLAND, WASHINGTON

0000000116
000000001

Prepared for WADCO
of
Richland, Washington

February 1971

by

John A. Blume & Associates, Engineers
612 Howard Street
San Francisco, California 94105

SUPPLEMENTARY GEOLOGIC INVESTIGATIONS FOR
SEISMIC EVALUATION OF THE FFTF SITE
NEAR RICHLAND, WASHINGTON

CONTENTS

Preface

Part 1, Section 1.

Geology along a portion of the Wallula Gap fault in the Horse Heaven Hills near Benton City, Washington. With a section on the Horn Rapids Lineament.

000000117

	<u>Page</u>
INTRODUCTION -----	1
GENERAL GEOLOGY AND SUMMARY -----	3
CHEMICAL ANALYSIS	
<u>General Statement</u> -----	5
<u>Discussion and Results</u> -----	5
STRATIGRAPHY	
<u>General Statement</u> -----	9
<u>Priest Rapids Flow No. 3</u> -----	10
<u>The Habton Interbed</u> -----	12
<u>Priest Rapids Flow No. 4 (Umatilla)</u> -----	13
<u>The Selah Interbed</u> -----	14
<u>The Pomona Flow</u> -----	15
<u>The Rattlesnake Ridge Interbed</u> -----	17
<u>The Elephant Mountain Flow</u> -----	17
<u>Surficial Deposits</u>	
Basalt Gravels -----	18
Older Alluvium -----	19
Loess -----	19
Residual Soil and Colluvium -----	19
Younger Alluvium -----	19
Glaciofluvial Gravels -----	20

Page

GEOLOGIC STRUCTURE

<u>Folding</u> -----	21
<u>Faulting</u> -----	23

0.00000000118

Illustrations, Part 1, Section 1.

	<u>Following Page</u>
11. Photograph of a Portion of the Wallula Gap Fault Zone in Badger Canyon -----	23
12. Brecciated Basalt adjacent to the Wallula Gap Fault In Badger Canyon. (photograph) -----	25

0.88888888120

TABLE 1. Results of Chemical Analysis of Samples from the Badger Coulee Area showing Sample Location and Flow Designation. -----	8
--	---

Part 2

Geologic Reconnaissance in the Vicinity of the Fast Flux Test Facility
Site near Richland, Washington.

	<u>Page</u>
INTRODUCTION -----	45
SUMMARY AND CONCLUSIONS -----	47
AREAL GEOLOGY	
<u>General Statement</u> -----	48
<u>The Central Area</u> -----	48
<u>The Eastern Area</u> -----	49
<u>The Western Basalt Area</u> -----	50
<u>Horn Rapids Lineament</u> -----	51
BIBLIOGRAPHY -----	52

0 0 0 0 0 0 0 0 0 0 1 2 3

Illustrations, Part 2

- PLATE
1. Reconnaissance Geologic Map of Fast Flux Test Facility Site and Vicinity. (in pocket)
 2. Photographs showing the Ringold formation along cliffs east of the Columbia River. (in pocket)

08080808124

PREFACE

This document contains results of two recent geologic investigations and updates a report on an earlier investigation, all of which were located in or near the Hanford Atomic Energy Commission reservation in Washington. The investigations were required for seismic evaluation of the Fast Flux Test Facility (FFTF) site. The primary purpose was to identify and delineate faults which are near enough to the FFTF site to be considered in its seismic evaluation.

Two types of possible seismic hazard must be considered in seismic evaluation. One is vibratory motion due to a distant earthquake and the other is due to surface rupture along a fault which passes beneath, or very close to the facility. Therefore, geologic studies were carried out near the site to locate any possible evidence of surface faulting and geologic mapping was done at some distance from the site to evaluate faults or possible faults along which a major earthquake might be localized.

The first geologic study was carried out in May 1970 and consisted of geologic reconnaissance within 5 to 10 miles of the site. This survey was intended to reveal any possible evidence of faulting which might pass directly under the site or very close to it. No such evidence was found. A report on this investigation was submitted to Battelle Memorial Institute Pacific Northwest Laboratory (BNW) of Richland, Washington on June 17, 1970. It was titled, "Geologic Reconnaissance in the Vicinity of the Fast Flux Test Facility Site at Hanford, Washington." It has been updated and comprises Part 2 of this document.

Part 1 of this report is titled, "Geology Along A Portion of the Wallula Gap Fault Near Benton City, Washington With A Section on

00000008125

In August 1970, a report titled "Seismic and Geologic Siting Evaluation Fast Flux Test Facility Near Richland, Washington" was submitted to WADCO.

In August and September 1970, thirty-six man days were spent in geologic mapping along a portion of the Wallula Gap fault near Badger Coulee and Benton City, Washington. Results of this work are described in the first section of Part I of this document.

Field work was done by Blume geologists Larry Tabor and Andrew Cunningham under the direction of Andrew Cunningham who also prepared the report.

00000000127

Part 1, Section 1

GEOLOGY ALONG A PORTION OF THE WALLULA GAP FAULT
IN THE
HORSE HEAVEN HILLS NEAR BENTON CITY, WASHINGTON

with a section on
The Horn Rapids Lineament

Prepared for
WADCO
of
Richland, Washington

February 1971

By
John A. Blume & Associates, Engineers
612 Howard Street
San Francisco, California 94105

0000000120

INTRODUCTION

This report describes a geologic investigation in the vicinity of Badger Coulee, about 7 miles southwest of Richland, Washington. The investigation was carried out by John A. Blume & Associates, Engineers (Blume), of San Francisco for WADCO of Richland, Washington.

The purpose of the study was to identify and delineate faults along the northeast slope of the Horse Heaven Hills. Knowledge of faulting in this region is necessary in determining seismic design criteria for the Fast Flux Test Facility (FFTF). Seismic and geologic evaluation of the FFTF site is presented in a prior report by Blume (August, 1970).

Because of the special aspect of this investigation which is concerned particularly with faulting, some topics normally included in a geologic study are omitted or abbreviated. Also, the mapped area was not examined in its entirety, but those areas which appeared to contain a maximum of structural information were mapped more completely than others.

The mapped area extends along the north slope of the Horse Heaven Hills from Badger Canyon northwest to the Yakima River and continues 2 miles beyond the river. It is about 14 miles long and varies in width from 1 to 2 miles. The hills are paralleled by Badger Coulee in the eastern and central part of the area. Thirty-six man days were spent during July and August, 1970, in geologic mapping which was done on aerial photographs at a scale of 1 in. equals 500 ft. Fifty-one samples of basalt were taken and submitted for chemical analysis of iron, calcium, potassium and barium content.

The cooperation of Randall E. Brown of Battelle Northwest Institute and Donald J. Brown of Atlantic Richfield Hanford Corporation

00000000129

In providing unpublished data and other assistance is gratefully acknowledged. However, John A. Blume & Associates, Engineers is responsible for the contents and conclusions of this report.

0000000130

GENERAL GEOLOGY AND SUMMARY

Geology of the Columbia Plateau has been the subject of numerous studies, most of which were concerned with broad regional aspects. In recent years the work of Laval (1956), Schmincke (1967), and others has provided a background of structural and stratigraphic information which is applicable to the Horse Heaven Hills area. Laval did limited geologic mapping in the area as part of a study which included a much larger region. Schmincke's work was primarily concerned with correlating stratigraphic units. His sample locations and cross sections traversed the area. The impetus for the present investigation was provided partly by recent work of Bingham, Londquist and Baltz (1970) who noted that the projected extension of the Wallula Gap fault mapped by them at Yellepit would lie along the north margin of the Horse Heaven Hills where it would be buried by Touchet beds.

Figure 1 shows the location of the mapped area in relation to the FTF Site and of geologic maps Plate I and Plate II.

The rocks exposed in the study area consist of Mio-Pliocene age basalt flows which are the uppermost part of the Yakima basalt formation. The flows are separated by intercalated beds of unconsolidated volcanoclastic sediments belonging to the Ellensburg formation. Stratigraphic terminology adopted by Schmincke (1967) and Bingham, Londquist and Baltz (1970) has been followed in this study with minor changes in each. From oldest to youngest the mapped units are the Priest Rapids Flow No. 3, the Mabton Interbed, Priest Rapids Flow No. 4 (Umatilla), the Selah Interbed, the Pomona Flow, the Rattlesnake Ridge Interbed, and the Elephant Mountain Flow.

Figure 2 is a simplified stratigraphic section and shows the relationship of flows and interbeds.

0.8888000133






Figure 2. Simplified stratigraphic section showing upper basalt flows of the Yakima basalt formation and interstratified sediments of the Ellensburg formation. Modified after Schmincke (1967). Field mapping included all of the units shown below the upper Ellensburg.

0688888134

The geologic structures in the area are of three distinct types. A small breached anticline is located in the northwest sector with an axial plane fault having about 50 ft displacement downward to the northeast (See section A-A', Figure 3). A monocline forms part of the crest of the Horse Heaven Hills in the central sector. Limited exposures do not provide sufficient data to determine whether or not faulting which might have contributed to elevation of the hills occurs along the steep northeast flank of the monocline (See section B-B', Figure 3). The southeasterly sector is occupied by flat-lying basalt flows and interbeds which are transected by the Wallula Gap fault. Displacement of about 300 ft downward to the northeast has taken place on the fault. The geologic structure is simplified and illustrated in Figure 3, section C-C'.

This investigation therefore confirms that the Wallula Gap fault extends northwest from Yellepit to the Horse Heaven Hills as suggested by the U.S. Geological Survey (Bingham, Londquist and Baltz, 1970). It is concluded that the Wallula Gap fault is a lesser fault or at most, it might be considered equal in size. As the Wallula Gap fault is a greater distance from the FFTF site it is concluded that the Rattlesnake-Wallula fault is the fault capable of producing maximum vibratory ground motion at the site.

00000000135



APPROX. SCALE 1 inch = 2 miles

John A. Stone and Associates,
Engineers. JABE-WADCO-04




Figure 3

CHEMICAL ANALYSIS

General Statement

0.000000136 ;


Mapping of the layered basalt flows is difficult because the flows are similar in appearance, and a prominent feature seen in one flow may occur in others. It is often necessary or desirable to identify a flow where only parts of it are exposed in isolated outcrops. Schmincke's work was addressed specifically to the problem of flow identification. Schmincke, (1957) studied most of the geologic properties of the flows including their physical appearance in the field, their megascopic and microscopic petrography, primary structures, weathering characteristics, optical properties of certain minerals within the flows, refractive index of fused beads from the flows, and chemical composition. Of the many properties studied, chemical composition is fundamental because other properties are influenced by or determined by the chemical composition. However, it is important to consider all properties to determine which of them if any are diagnostic of certain flows, as Schmincke has done.

Schmincke has noted that field criteria for identification of the flows is not diagnostic and, "a flow so identified should be correlated at new outcrops only if the over and underlying flows can also be identified. Although helpful, this type of identification and correlation is highly controversial."

In the present structural geology investigation, no microscopic or detailed petrographic studies were made due to time and budgetary limitations. Reliance was placed upon features of the rocks which could be observed in the field and upon chemical analysis.

Discussion of Results


Chemical analyses of the basalt flows of South Central Washington by Schmincke (1967) and others have shown that some of the flows



Chemical Analysis of Basalt Flows from Central Washington after Schmincke (1967)


Fig. 4

0000000139



Chemical analysis of iron and calcium content of samples from Badger Coulee area.
Showing sample numbers


0000000717



Chemical analysis of iron and potassium content of samples from Badger Coulee area showing sample numbers.

Fig. 6

0000000141



Chemical analysis of iron and barium content of samples from Badger Coulee area.
Showing sample numbers

0000000

0000000142




Fig. 8

0 0 0 0 0 0 0 1 4 3

Barium content of samples is plotted in Figure 7. Nine samples contain 2000 or more parts per million (ppm) of barium while the remaining samples contain 700 ppm or less. The nine samples are from the two flows of the upper part of the Priest Rapids member of the Yakima Basalt formation.

The chemical data show that two flows sampled in sec. 15, T.9 N., R.29 E., which are separated by a sedimentary interbed are both high in barium. The uppermost of these flows lies below the Pomona flow and corresponds to the Umatilla flow of Schmincke and Laval. They considered the sedimentary interbed to be the Mabton bed. Bingham, Londquist and Baltz designate the flow below the Pomona as "Priest Rapids flow No. 4" and accordingly the 2nd flow would correspond to their Priest Rapids flow No. 3. These workers placed the Mabton bed below flow No. 3 rather than above it, as Schmincke and Laval had done. Laval (1965, p. 43) refers to the red alteration zone at the top of the Mabton interbed in this locality which clearly corresponds to the interbed between Priest Rapids flow No. 3 and 4 in sec. 15. In this report the terminology of Bingham, Londquist and Baltz will be followed. The uppermost flow is therefore the Priest Rapids flow No. 4 (Umatilla) and the second barium-rich flow is the Priest Rapids flow No. 3. The interbed which separates these two flows will be called the Mabton interbed in accordance with the work of Schmincke and Laval.

Three samples were run in duplicate to determine approximately what amount of variation is present within a single sample. Duplicate samples were prepared by selecting two identical 1/2 lb. samples from an original 10 lb. sample. The samples which were run in duplicate were numbers 73, 74, and 75. Variations in iron and calcium content were on the order of several tenths of a percent and variations in potassium content amounted to several hundredths of a percent. The spectrographic method used in determining barium is not highly accurate but shows that the values for duplicated determinations of barium are comparable. The results are in agreement with Schmincke (1967) who found that analyses of basalt flows were reproduceable within similar limits.

Four analyses gave anomalous results. That is, the results did not correspond to results of other samples or to flow identification based upon geologic evidence. The samples are numbered 31, 35, 41 and 42. The results are being evaluated based upon more comprehensive chemical data.

Results of analyses and flow identification derived from the data are compiled in Table 1. With exceptions noted above these determinations are in agreement with determinations which were made independently based upon geologic field evidence.

0000000000
00000000144

TABLE 1

RESULTS OF CHEMICAL ANALYSIS OF SAMPLES FROM THE BADGER COULEE AREA,
SHOWING SAMPLE LOCATION AND FLOW DESIGNATION

Sample Number	Location				Chemical Analysis				Flow Designation
	1/4 Sec.	Sec.	Township (North)	Range (East)	FeO (%)	CaO (%)	K ₂ O (%)	Barium (ppm)	
29	NE	22	9	26	17.0	8.4	2.4	500	
30	NE	22	9	26	16.9	7.7	2.7	500	
31	NE	22	9	26	13.4	9.1	1.3	300	
32	SE	15	9	26	14.7	5.6	5.1	2,000	Typr3 or Typr4
33	SE	15	9	26	16.0	5.2	5.5	3,000	- same -
34	SE	15	9	26	14.4	5.2	5.3	3,000	- same -
35	SE	33	9	27	14.2	5.7	1.1	300	
36	NW	4	8	27	15.8	5.2	5.1	3,000	Typr3 or Typr4
37	NE	8	8	27	15.8	8.4	4.8	3,000	- same -
38	SE	33	9	27	17.6	7.0	2.4	700	
39	NW	29	8	28	11.8	9.8	0.7	300	Pomona
40	NW	29	8	28	13.8	5.8	2.7	3,000	Typr3 or Typr4
41	SE	30	8	28	14.2	9.1	1.1	500	
42	NE	19	8	28	15.3	8.7	1.1	200	
43	SE	18	8	28	11.5	10.5	0.7	200	Pomona
44	SE	18	8	28	11.1	11.2	0.7	500	Pomona
46	SE	12	8	27	11.6	9.4	0.7	300	Pomona
48	NE	11	8	27	11.6	10.5	0.8	300	Pomona
49	NW	15	9	26	13.8	5.7	2.6	3,000	Typr3 or Typr4
50	NW	15	9	26	12.7	5.1	2.8	3,000	- same -
51	NW	15	9	26	13.5	6.4	2.0	700	
53	SE	15	9	26	11.8	10.1	0.6	300	Pomona
54	SE	15	9	26	11.0	10.5	0.7	300	Pomona
55	SE	15	9	26	14.4	6.4	1.8	700	
56	NW	19	9	27	11.5	9.4	0.8	300	Pomona

00000001
0000000145

00000000146

Sample Number	Location				Chemical Analysis				Flow Designation
	1/4 Sec.	Sec.	Township (North)	Range (East)	FeO (%)	CaO (%)	K ₂ O (%)	Barium (ppm)	
57	NE	24	9	26	15.6	8.0	1.0	500	
58	NE	25	9	26	11.3	10.5	0.7	200	Pomona
59	SW	33	9	27	11.3	9.8	0.7	300	Pomona
60	SE	32	9	27	15.7	8.0	1.1	700	
61	NW	32	9	27	16.2	7.1	1.3	500	
63	NW	25	9	26	11.3	9.4	0.7	300	Pomona
64	NE	33	9	27	10.6	11.9	0.7	300	Pomona
65	NW	34	9	27	10.4	10.8	0.6	300	Pomona
66	NE	8	8	27	11.3	8.4	0.5	300	Pomona
67	NW	4	8	27	11.2	9.4	0.5	300	Pomona
68	NW	33	9	27	10.9	11.9	0.4	300	Pomona
69	SW	30	9	27	14.6	8.0	1.0	700	
70	NE	23	9	26	11.2	9.8	0.6	300	Pomona
73	NE	22	9	26	11.6	10.5	0.78	150	Pomona
73B	NE	22	9	26	11.5	10.1	0.75	100	
74	NW	23	9	26	11.3	10.5	0.60	500	Pomona
74B	NW	23	9	26	10.9	9.4	0.61	500	
75	NW	23	9	26	11.9	9.1	0.49	200	Pomona
75B	NW	23	9	26	11.3	9.1	0.41	200	
76	SE	33	9	27	11.6	9.1	0.6	200	Pomona
77	SE	8	8	27	15.9	7.7	1.1	500	
78	NE	8	8	27	11.8	9.8	0.5	300	Pomona
79	NE	8	8	27	11.9	9.4	0.5	300	Pomona
80	NE	8	8	27	13.6	5.1	2.6	2,000	Typr3 or Typr4
81	SE	29	9	27	15.5	7.3	1.3	500	
82	SW	28	9	27	11.8	9.1	0.5	300	Pomona

STRATIGRAPHY

General Statement

The nomenclature adopted by the U.S. Geological Survey (Bingham, Londquist and Baltz, et al, 1970) is followed in this work with one minor modification which places the Mabton interbed between Priest Rapids flows No. 3 and 4 instead of below Priest Rapids flow No. 3. This nomenclature also corresponds closely to that followed by Schmincke (1967). (See Figure 9).

The stratigraphic units which were observed and mapped in the field from oldest to youngest are the Priest Rapids flow No. 3, the Mabton interbed, Priest Rapids flow No. 4, the Selah interbed, the Ponona flow, the Rattlesnake Ridge interbed, and the Elephant Mountain flow. Surficial materials which are not listed in order of age consist of residual soil and colluvium, loess, younger alluvium, older alluvium, basalt gravels, landslide debris, and glaciofluvial deposits.

Sedimentary interbeds are not generally useful as recognizable stratigraphic horizons because of their great lateral variability in petrography and thickness and the scarcity of outcrops.

The Ward Gap flow, a thin unit lying upon the Elephant Mountain flow was not differentiated. The petrography of the basalt flows and sedimentary interbeds has been described most recently in detail by Schmincke and previously by others. The flows are tholeiitic basalt. Tholeiitic basalt is common to most continental flood basalt provinces throughout the world and its occurrence is usually continental. As described by Schmincke, the chief mineral constituents in the flows are "plagioclase, clinopyroxene, iron titanium ores and olivine. Apatite occurs in all the flows but cristobalite and zeolite are rare. A mineraloid (green, yellow or brown chlorophaeite) is ubiquitous in vesicles, cracks and along grain boundaries."

0.0000000147

FIGURE SHOWING PRINCIPAL MAP UNITS, THEIR THICKNESS AND AGE RELATIONSHIPS

SYSTEM	SERIES	GROUP	FORMATION	MEMBER OR UNIT	ESTIMATED THICKNESS (feet)	
Quaternary	Holocene		Not in order of age	Loess & wind blown sand	0-30	
				Younger alluvium	0-15	
				Older alluvium	0-30	
				Residual soil & colluvium	0-30	
				Landslide debris	0-40	
Pleistocene	?		Ringold	Undifferentiated glacio-fluvial deposits	0-50	
				Sand, silt & clay	Absent in mapped area	
Tertiary	Columbia River Group		Ellensberg	Basalt gravels	0-180	
				Yakima Basalt	Hard Gap Flow	
					Elephant Mountain Flow	30-60
					Rattlesnake Ridge Bed	30-40
					Pomona Flow	100-300
					Selah Bed	30-40
					Flow No. 4 (Umatilla)	75-150
					Mabton bed	30-40
					Flow No. 3	220?

Modified after Bingham, Lordquist and Baltz (1970)

FIGURE 9

Differences in crystallinity and other textural variations as well as compositional differences occur within flows. Schmincke (1957) states, "In addition to intraflow variations significant interflow differences in textures and modal, mineral, and chemical compositions also exist between most flows. They are most obvious in comparisons of rocks of similar crystallinity."

Useful criteria in field identification of flows are the usually coarser grain size and lighter color of the Elephant Mountain flow and the finer grain size and darker, nearly blue-black color of the Pomona flow. Sinuous columnar jointing forming columns 6" to 9" in diameter is typical of the Pomona while the Elephant Mountain flow is often more massive. The Pomona tends to weather into equidimensional, angular blocks and the Elephant Mountain flow often weathers to sub-rounded, bouldery shapes. The Priest Rapids flow No. 4 resembles the Pomona flow in most of its megascopic features. Priest Rapids flow No. 3 is also a dense, dark bluish-black, fine grained basalt. A distinctive feature of the central part of this flow in the mapped area consists of closely spaced fracturing which forms elongate, prismatic, or tabular pieces. The fragments are typically 6" to 9" in width and often up to 2 or 3 ft in length. These fragments usually produce a distinctive ringing sound when struck.

Priest Rapids Flow No. 3

The Priest Rapids flow No. 3 is the oldest known rock unit exposed in the mapped area. Its base is not known to be exposed. It crops out in Webber Canyon, in sec. 15 and sec. 22, T.9N., R.29 E., and possibly in Badger Canyon.

The upper portion of the flow is widely exposed in sec. 15 and 22 about 3 miles west of Kiona where it consists of highly scoriaceous and vesicular basalt or clinker. It forms rough, jagged, spinose outcrops consisting of aphanitic, dark blue grey basalt intermixed with red to purple scoria pieces and pale brown silica gel. Most outcrops are isolated and appear to project vertically,

00000000149

perpendicular to the surrounding flow surface. They have a distinctly elongate form which parallels the structural trend of the Horse Heaven Hills. Similar scoriaceous material or clinker was not observed at other localities where the Priest Rapids flow No. 3 crops out.

The interior of the flow exhibits distinctive, closely spaced fracturing which produces smooth sided tabular or prismatic fragments up to 2 or 3 ft in length.

In the north half of sec. 15, three miles west of Benton City, the uppermost part of the flow crops out at the base of a well exposed, steeply dipping sequence of rocks. Successively exposed above the Priest Rapids flow No. 3 are the Mabton bed, Priest Rapids flow No. 4 (Umatilla), and the Selah bed. The Priest Rapids flows at this locality were identified by their high barium content.

A few hundred feet south of this sequence the scoriaceous crust of flow No. 3 crops out again dipping gently to the north. The distinctive prismatic interior of the flow is exposed in the deep canyon near the center of section 15.

The Priest Rapids flow No. 3 crops out along Webber Canyon Road in sec. 4 and sec. 5, T.8 N., R.27 E. The distinctively fractured prismatic phase of the flow is well exposed in several road cuts. The basalt flows in Webber Canyon comprise a flat lying, layered sequence and successively younger flows are exposed in the canyon as it rises to the south.

Northward in sec. 33 nearer the mouth of the canyon the rocks exposed at the base of the canyon walls belong to the Pomona flow. This abrupt transition from older to younger rocks is explained by movement on the Wallula Gap fault which has displaced the northeasterly block downward. The fault is not exposed in Webber Canyon.

0 0 0 0 0 0 0 0 0 0 1 5 0

In Badger Canyon a high barium flow crops out on the south side of the Wallula Gap Fault in the NW/4 of sec. 29, T.8 N., R.28 E. The basalt is dark greyish to blueish black, dense and fine grained. Columnar jointing is well developed in outcrops along the banks of the stream channel. Thirty to fifty feet above the stream, scoriaceous and vesicular basalt crops out irregularly. The Mabton bed is not exposed and other evidence which might serve to distinguish between Priest Rapids flow No. 3 and 4 is lacking. Therefore, it is concluded that the high barium rocks represent the upper part of either Priest Rapids flow No. 3 or Priest Rapids flow No. 4.

Thickness of the Priest Rapids flow No. 3 cannot be estimated because the base of the flow is not exposed in the map area. However, core drilling in the Hanford reservation shows that the combined thickness of the high barium flows is about 300 ft. If 80 ft is deducted for the thickness of flow No. 4, which is its thickness where exposed 3 miles west of Benton City, it appears that the Priest Rapids flow No. 3 is about 220 ft thick.

Age of the various basalt flows is being determined by the potassium-argon method and results will be published in the future. (Donald Brown, 1970). These determinations are part of continuing investigations being carried out by Atlantic Richfield Hanford Co. Bingham, Londquist and Baltz (1970) assign a Mio-Pliocene age (about 12 million years) to the upper Priest Rapids flows.

The Mabton Interbed

The Mabton interbed is well-exposed in the north half of sec. 15 about 3 miles west of Benton City. It consists of well-bedded white tuff with minor amounts of sand and silt. The uppermost 2 to 3 feet of the tuff are baked and stained pale grey. Below this greyish baked zone is a brightly reddish stained zone about 3 or 4 feet in thickness. The total thickness of the interbed at this locality is about 35 feet. The Mabton is overlain by Priest

00000000151

Rapids flow No. 4 and is underlain by Priest Rapids flow No. 3. Soft sedimentary interbeds such as the Mabton outcrop very poorly. No other exposures of the Mabton are known in the map area.

Priest Rapids Flow No. 4 (Umatilla)

The Priest Rapids flow No. 4 is the uppermost of two of barium-rich flows which have been identified by Donald Brown (1970). It is separated from the Pomona flow above by the Selah interbed. Flow No. 4 is separated from flow No. 3 below it by the Mabton interbed. It is called the Priest Rapids flow No. 4 by the U.S. Geologic Survey and it corresponds to the Umatilla of Schmincke (1967).

The flow consists of dark greyish black to dark blueish black fine-grained basalt exhibiting well developed columnar jointing which is generally more regular and of larger diameter than that of the overlying Pomona flow. In local areas where the rock has been stressed and folded as in parts of sec. 15, the Priest Rapids flow No. 4 becomes highly brecciated. Small fractures in the breccia pieces vary from about one inch to four inches in size. The rock is weathered and stained yellowish, reddish brown, and greenish.

Priest Rapids flow No. 4 comprises the central flow in a sequence of well-exposed rocks in the north half of sec. 15, T.9 N., R.26 E. The flow has a steep northerly dip in sec. 15. Its outcrop continues to the southeast where it forms a prominent ridge for a distance of 1-1/2 miles.

Flow No. 4 also outcrops in Webber Canyon in sec. 5, T.9 N., R.27 E. At this locality the flat lying flow is in contact with the overlying Selah interbed, which in turn is in contact with the Pomona flow above. The Pomona flow and Priest Rapids flow No. 4 were sampled and identified by their diagnostic chemical characteristics. The Mabton interbed underlying flow No. 4 does not outcrop well in Webber Canyon but flow No. 3 beneath the Mabton interbed is well-exposed at lower elevations in the canyon.

00000000152

Priest Rapids flow No. 4 may also outcrop in Badger Canyon along the southwest side of the Wallula Gap fault. Local outcrops of basalt at this locality which is in sec. 29, T.8 N., R.28 E. were sampled and found to be high in barium content indicating that the basalt belongs to one of the uppermost Priest Rapids flows. There is insufficient evidence to indicate whether the rocks belong to Priest Rapids flow No. 4 or flow No. 3.

In the north half of section 15 about three miles west of Benton City, the base and top of flow No. 4 are both well exposed. The thickness of the flow here is about 80 feet. In Webber Canyon the base of the Priest Rapids flow No. 4 is not well exposed. However, estimates of its approximate elevation indicate a thickness for the flow here on the order of 100 feet.

Most recent estimates of age of these rocks are contained in a U.S. Geological Survey report by Bingham, Londquist and Baltz (1970). These writers assigned a Mio-Pliocene age to the Priest Rapids flow No. 4.

The Selah Interbed

The Selah interbed lies between Priest Rapids flow No. 4 and the Pomona flow. It consists of poorly consolidated tuff with varying amounts of sand and silts. Local occurrences of gravel in the Selah were described by Schmincke (1967). One such occurrence was observed within the mapped area of this survey in the SE/4 of sec. 12, T.8 N., R.27 E. A zone of dark grey vitreous fused tuff about 2 feet thick usually marks the top of the Selah interbed. Fused tuff at the base of the Pomona and at the top of the Selah interbed crops out in sec. 15, T.8 N., R.26 E., sec. 4, 8, and 12, T.8 N., R.27 E., and in sec. 18 and 19, T.8 N., R. 28 E.

Where surface soils are underlain by the Selah interbed the soil frequently exhibits well-developed shrinkage cracks and clay bloom indicating a high content of montmorillonite clay.

00
25
10888888880

The thickness of the Selah Interbed is difficult to estimate due to poor exposures. It appears to be about 30 to 40 feet in thickness.

The age of the Selah Interbed as given by Bingham, Lundquist and Baltz (1970) is Mio-Pliocene.

The Pomona Flow

The Pomona flow was named by Schmincke (1967), who showed that the flow could be identified over large areas and that it has distinctive petrographic and chemical characteristics. The Pomona flow is underlain by the Selah Interbed and overlain by the Rattlesnake Ridge Interbed. The petrography and chemistry of this flow have been described elsewhere in this report. The reader is referred to the work of Schmincke (1967) for detailed descriptions.

A well sampled outcrop of the Pomona flow is located in a road cut in Highway 81 about 3/4 of a mile northwest of Kiona. This locality was sampled by Schmincke (1967) for analysis and also by Blume geologists. The Pomona flow outcrops extensively on the northwest side of the Yakima River in sections 7, 8 and 9, T.9 N., R.26 E., where it comprises the lowermost exposed unit. It is overlain by the Rattlesnake Ridge Interbed which in turn is overlain by the Elephant Mountain flow. The Pomona flow also outcrops in sec. 10 on the southeast side of the Yakima River and from there intermittently along the base of the Horse Heaven Hills. A quarry situated at the base of the hill in sec. 23, T.9 N., R.26 E., has exposed the Pomona flow. The flow dips from 50 to 65° to the northeast and strikes N 60°W parallel to the mountain ridge. Parts of the flow at this locality have a scoriaceous, vesicular and brecciated crust similar to the crust of flow No. 3. This clinker, as it is called by Laval (1956), or scoriaceous crust outcrops intermittently along the base of the hills for about two miles southeast of the quarry. The Pomona flow was also identified along the crest of the Horse Heaven Hills in sec. 22 and 23, T.9 N., R.26 E., and sec. 31, T.9 N., R.27 E.

0 0 0 0 0 0 0 0 1 5 6

a broad flat plateau situated between Webber Canyon and Badger Canyon and outcrops extensively along the cliffs at the north margin of this plateau. The Pomona flow outcrops near the mouth of Badger Canyon and upstream as far as sec. 9, T.8 N., R.28 E., where it is terminated on the Wallula Gap fault. The lower contact of the Pomona flow is exposed intermittently along the base of the cliffs which face Badger Coulee between Badger Canyon and Webber Canyon. The base of the flow is also well exposed in sec. 8, T.8 N., R.27 E., in the upper part of the canyon. At locations where the elevation of the base of the Pomona is known the elevation of the top of the flow has been estimated giving thicknesses of from 250 to 300 feet.

Bingham, Londquist and Baltz (1970) give a lower Pliocene age for the Pomona flow.

The Rattlesnake Ridge Interbed

The Rattlesnake Ridge Interbed crops out in five locations northwest of the Yakima River but it is not known to crop out in the map area southeast of the river. Good exposures of the bed are located immediately north of the road in sec. 9, T.9 N., R.26 E. Farther to the northwest along the trend of this small breached anticline, the Rattlesnake Ridge bed appears in small isolated outcrops of well-cemented grey tuffaceous sand. The thickness of the bed in this locality is from 30 to 40 feet. Its age is given by Bingham, Londquist and Baltz (1970) as lower Pliocene.

The Elephant Mountain Flow

The Elephant Mountain flow lies upon the Rattlesnake Ridge Interbed and is the uppermost rock unit mapped during this survey.

The petrography of the Elephant Mountain flow is summarized elsewhere in this report and has been treated in detail by Schmincke (1967). Chemical analyses made during the course of this study indicate that the chemical characteristics of the Elephant Mountain

flow using the analyses employed for this study are not sufficiently distinctive to serve as a means of identifying the flow.

The flow crops out extensively along the trend of the small breached anticline on the northwest side of the Yakima River. The flow crops out locally southeast of the Yakima River in sec. 9, 15, and 16, T.9 N., R.26 E.

A localized outcrop of the Elephant Mountain flow forms the top of the rounded hill in sec. 22, T.9 N., R.26 E. The flow also crops out locally at other places along the ridge forming the crest of the Horse Heaven Hills southeast toward Webber Canyon. The Elephant Mountain flow is also present, although poorly exposed, as the uppermost rock unit on parts of the plateau at 1100 ft elevation northwest and southeast of Webber Canyon.

The Elephant Mountain flow is best exposed in the vicinity of a small breached anticline northwest of the Yakima River where it is approximately 80 feet thick. Elsewhere in the map area where the Elephant Mountain flow is exposed its thickness appears to be comparable.

The age of the Elephant Mountain flow according to Bingham, Lundquist and Baltz (1970) is lower Pliocene.

Surficial Deposits

Basalt Gravels - Basalt gravel crops out at two localities along the crest of the Horse Heaven Hills within the mapped area. One of these is in sec. 23, T.9 N., R.26 E. where the gravels appear to be about 20 feet thick and consist predominantly of basalt clasts averaging 3 to 4 inches in diameter. The second locality is in sec. 32 and 33, T.9 N., R.27 E. and sec. 4 and 5, T.8 N., R.27 E. on the crest of a high ridge overlooking Webber Canyon. Here the gravels are similar in size and composition. They outcrop over an area of about 1/4 square mile and appear to be about 100 ft thick.

00000000157

05188888880

The basalt gravels are unusual in that their composition is predominantly basalt indicating a local origin within the Columbia plateau itself for most of the gravel constituents. Secondly, the gravels are of interest because no basalt gravels have been found at low elevations. The known outcrops are restricted to higher elevations along the ridge top. It is clear that the gravels indicate the former presence of a stream or river at their place of deposition. A possible explanation for the presence of these gravels is that they were deposited by a pre-glacial southwesterly flowing river which existed before elevation of the Horse Heaven Hills.

Older Alluvium - Older alluvium was mapped at a single locality in sec. 16, T.9 N., R.26 E. It forms an old river terrace at an elevation of 830 feet or about 330 feet above the present elevation of the Yakima River.

Loess - Loess has been deposited widely throughout the area. It is present on most level ground and many steep slopes are also blanketed.

Residual Soil and Colluvium - In many places residual soil forms a thin blanket 1 or 2 feet thick above the underlying rock units. Such soil is typically developed on gently sloping hillsides. Areas covered by residual soil have been identified and are indicated on the geologic map.

Younger Alluvium - Younger alluvium has been deposited along the Yakima River and in Badger Canyon and Webber Canyon in places where the stream gradient becomes low. Some alluvium has been deposited at higher elevations where drainage has been trapped by landslides, especially in sec. 15, 16, 21 and 22, T.9 N., R.26 E. Two landslides can be definitely identified on the north slopes of Horse Heaven Hills in sec. 15 and 16, T.9 N., R.26 E. In many other places closed depressions suggest the presence of landsliding but the actual slide and its margin cannot be located.

The structure along portions of the north slope of the Horse Heaven Hills where it consists of moderately to steeply dipping basalt flows underlain by poorly consolidated sediments is highly conducive to landsliding. In the past when river erosion cut away the basalt flows at the base of the hills the flows themselves became unsupported and landsliding must have resulted.

Glaciofluvial Gravels - These gravels are intermixed with sand and silt, and individual clasts are composed of many different rock types. Large boulders called glacial erratics which may have in part been transported by ice are common in this formation. Glaciofluvial gravels crop out in an area about 2 miles southwest of Goose Hill.

00000000059

not well exposed. Laval (1967) described the major structure of the Horse Heaven Hills as having been superimposed upon minor east-westerly trending folds. The minor structural features observed in this area could be explained in this way and are considered to lend support to Laval's interpretation.

Overtured bedding occurs at the base of the monocline in sec. 15, about 3 miles west of Benton City. Here the lower contact of the Pomona flow is exposed adjacent to fused tuff of the Selah Interbed, both of which dip about 70° to the southwest and strike northwesterly. The Pomona basalt is cut by two fault zones about six inches in width which parallel the attitude of the flow. These features could be interpreted as overturned folding and related thrust faulting, but they could also be explained as the result of landsliding and slumping at the base of the hill.

The monoclinial section of the Horse Heaven Hills structure lying within the mapped area extends from Webber Canyon northwesterly to a point about 3 miles west of Benton City.

The difference in elevation between the crest of Horse Heaven Hills and the floor of the canyon to the north is greatest in this monoclinial section and less in the section to the southeast where the bedding is flat.

The basalt flows south of the Horse Heaven Hills crest slope gently upward to the northeast on the broad plateau until reaching the crest of the hill where they bend abruptly and plunge down the north face of the hill until reaching the base where they again flatten out. The generally heavy cover of loess along the north slope of the hill and overall paucity of outcrops have rendered most of this portion of the structure obscure. Therefore, in this area it could not be determined within the scope of this survey whether or not the flows have been displaced by faulting between the crest of the hill and the valley floor. The structural conditions which are known

1918888888.0

cut by numerous vertical fault structures about 1 inch in thickness which are filled with fine sand sized and pebble sized breccia. Three such fault structures are 1 to 2 feet in width. They strike from north 60 to north 70 west and are normally spaced at 1 or 2 foot intervals. Locally these individual fault structures are filled with from 1 to 5 inches of chalcedony.

Some apparent flow banding occurs within the fault zone which is parallel to the general strike and dip of the fault. The vertical flow structures could represent irregularities in an otherwise normal horizontal basalt flow. It might also indicate flowage of a dike-like intrusion which invaded the fault zone. Or it is possible that fault movements taking place before final solidification of the flow could produce vertically oriented banded structures parallel to the fault. Figure 11 is a photograph of a portion of the fault zone which shows the vertical fault structures, fault breccia and intervening fine-grained basalt.




Figure 11: A portion of the Wallula Gap fault zone in Badger Canyon.

rock and that those movements which did take place were downward toward the east throughout the mass. Figure 12 shows a close-up view of the breccia.




Figure 12. Brecciated basalt adjacent to the Wallula Gap fault in Badger Canyon.

The dark fine-grained basalt with columnar jointing which is exposed near the fault zone and at the breccia zone continues upstream in outcrop about 200 yards beyond the breccia zone. The columnar basalt is exposed vertically up the northwest canyon wall for a distance of 50 feet where it is capped by a scoriaceous and vesicular crust. This basalt was sampled about 50 feet southwest of the breccia outcrop in the wall of the canyon and analysis showed a high barium content indicative of the two uppermost flows of the Priest Rapids member of the Yakima basalt.

The structural conditions indicate displacement on the fault zone with downward movement on the northeast side of the fault. The amount of offset on the fault is estimated to be from 200 to 300 ft. The structure is illustrated on Cross Section D shown on Geologic Map Plate II.

0-000000160

suggests the presence of a northeasterly trending cross fault. However, no fault could be located on the ground and in fact the structural relations are such as to preclude any significant fault movement along the lineation. It is most probable that the lineation represents a fracture zone along which there has been no displacement. It is indicated on the geologic map as an alignment of fractures. Other similar lineations were observed on aerial photos at the apex of the Horse Heaven Hills about 3 miles west of Benton City. These also may be fracture zones without significant displacement. They are indicated on the geologic map as dotted lines with question marks.

Age of faulting on the Wallula Gap fault where exposed in the southeastern area was not determined. It is known to be younger than the basalt gravels which were deposited during initial uplift of the Horse Heaven Hills. It is probable that trenching across the trend of the Wallula Gap fault in the surficial materials to exposed Holocene ash beds or other stratigraphic horizons would produce more conclusive evidence regarding the age of last movement. Such additional investigation is not recommended because the data which might be obtained is not critical to the FFTF project.

The vertical component of movement on the Wallula Gap fault in the southeastern area has been determined, but if any horizontal movement took place it would not be observable in the flat lying sequence of flows and interbeds. It might, however, be observable on the ground surface where any possible recent horizontal movement would be expected to displace stream channels. There is no clearcut evidence of displacement of stream channels, and one stream flowing to the northeast through sec. 11, T.8 N., R.27 E., does not show any curvature at all but on the contrary is quite straight. It is concluded therefore, that the predominant movement on the Wallula Gap fault has probably been vertical without significant horizontal component. The Wallula Gap fault is therefore probably a normal fault.

Part 1, Section 2

GEOLOGY ALONG THE
HORN RAPIDS LINEAMENT

Prepared for WADCO
of
Richland, Washington

February 1971

by

John A. Blume & Associates, Engineers
612 Howard Street
San Francisco, California 94105


0-0000000170

By far the greatest part of the map area is blanketed by surficial deposits consisting of wind blown sand, loess and alluvium. No significant mappable features were observed in the surficial deposits and mapping was confined to areas of basalt outcrop. For convenience the hills are numbered 1 through 6 beginning with the northerly most hill which is called Hill No. 1 and ending with the southerly most Hill, 2 miles south of Richland, which was called Hill No. 6.

Geology and cross sections through each hill are shown on geologic maps, Plate III, IV and V. Figure 13 shows locations of the cross sections, the hills, and the geologic map plates in relation to the FTF site.

0888888872

Figure 13. Location Map Showing Geologic Plates and Geologic Cross Sections, FFTF Site, and Hills 1 thru 6. (map taken from U.S. Geological Survey Topographic Maps titled Badger Mtn. & Richland Quadrangles)



It was concluded that with two exceptions all of the basalt exposures in the area are of the Elephant Mountain flow. The Ward Gap flow, which is a thin basalt flow overlying the Elephant Mountain flow, might be present in the area but was not distinguished. The two exceptions are small outcrops of basalt identified as belonging to the Pomona flow by chemical analysis. One of these outcrops is situated a short distance west of Hill No. 1 and the other is located about 1 mile northwest of Hill No. 3. In both instances the exposure of the older Pomona flow was due to a combination of geologic structure which brought the Pomona flow into a relatively high position and erosion which stripped away the overlying Elephant Mountain flow.

0.0000000175

STRATIGRAPHY

The principal rock type exposed along the Horn Rapids Lineament is the Elephant Mountain basalt. The Ward Gap flow, which is a thin flow overlying the Elephant Mountain, may also be exposed in the area but it was not distinguished. Several different phases of the Elephant Mountain flow crop out in the map area. Most common is a massive, medium grained highly crystalline phase. This rock as well as the other types of Elephant Mountain basalt typically exhibits red-brown weathering color and indistinct banding. Subrounded boulders of this type of basalt reaching 10 to 15 ft in diameter crop out on Hill No. 6. A darker, dense, fine-grained to aphanitic phase of the Elephant Mountain basalt also crops widely in the area. Locally, this type of basalt is brecciated as described on Hill No. 1, Hill No. 3 and north of Hill No. 3. The breccia is probably due to stress created by folding of the basalt flows.

Jagged, spinose outcrops of very fine to aphanitic basalt crop out north and west of Hill No. 2. This rock is highly vesicular, dark bluish-black in color and often exhibits ropy structure. It is probably the upper surface of the Elephant Mountain flow or possibly the Ward Gap flow.

A less common phase of the Elephant Mountain flow is basalt which weathers into spherical shapes varying from about 6 to 12 inches in diameter. This rock was exposed in a quarry at the south end of Hill No. 6.

The base of the Elephant Mountain flow was exposed at one locality on the south bank of the Yakima River a short distance north of Hill No. 2. Here the flow rests upon slightly baked tuff of the Rattlesnake-Ridge interbed.

0888800176

Two outcrops of dark dense fine-grained basalt of the Pomona flow were identified. One of these is west of Hill No. 1 and the other is about 1 mile northwest of Hill No. 3.

Flow identification was based on hand specimens examination and chemical analysis. The method of analysis and the basis of flow identification is discussed in detail in Part I of this report. Briefly, the chemical analysis methods of identification used in this study are based upon the work of Schmincke (1967) and more recent unpublished work by D. J. Brown (1970). The basalt analyses used for flow identification were provided by D. J. Brown of Atlantic Richfield Hanford Company (ARHCO). Results of analyses and sample locations are presented in Table 2. Figure 14 is a graphical plot of the iron, potassium and calcium content of samples which can be compared with a similar plot of Schmincke's data (Part I, Figure 4). Sample locations are also plotted on the geologic map (Plates III, IV and V).

Although the method of analyses was not considered to provide a diagnostic identification of the Elephant Mountain flow, the flow can be identified with a reasonable degree of assurance by a process of elimination.


The Pomona flow lying below the Elephant Mountain can be distinguished chemically. Likewise, the uppermost two flows of the Priest Rapids member lying below the Pomona can be distinguished chemically. Therefore, if a flow does not belong to either one of these two, then it must either be from the lower part of the Priest Rapids member or the Elephant Mountain or Ward Gap flow. It is highly unlikely that any of the basalts exposed in the area could belong to the lower Priest Rapids member because this would require removal by erosion of approximately 800 feet of overlying basalt flows and interbeds. Therefore, it can be concluded that the rocks belong to the Elephant Mountain flow. This conclusion is supported by the fact that in two instances the Pomona flow was identified in a stratigraphically lower position. The reader

0-100000177

TABLE 2

CHEMICAL ANALYSIS OF BASALT SAMPLES FROM HORN RAPIDS LINEMENT

Sample No.	1-18	3-18	4-34	5-34	6-34	9-10	11-10	12-15	13-10	14-15	15-23	16-23
SiO ₂ (%)	51	47	50	49	48	50	49	52	50	50	49	52
Al ₂ O ₃ (%)	12.9	13.0	12.4	12.6	12.7	13.0	13.0	13.7	13.6	13.7	14.5	12.5
FeO (%)	14.7	15.1	15.0	15.1	14.6	13.8	13.3	14.2	15.0	13.1	11.0	14.4
MgO (%)	4.5	7.1	6.0	6.0	4.7	6.3	4.4	5.8	7.0	5.6	8.0	4.3
CaO (%)	10.9	10.1	9.4	8.4	8.6	9.5	8.6	10.1	10.3	10.4	9.7	8.4
Na ₂ O (%)	2.3	2.3	2.7	2.4	2.5	2.4	2.5	2.3	2.4	2.5	2.4	2.9
K ₂ O (%)	1.2	0.7	0.8	1.3	1.2	0.5	1.0	0.5	0.6	0.6	0.6	1.3
TiO ₂ (%)	3.1	2.9	3.1	3.2	3.2	2.9	3.0	3.2	3.1	2.9	1.6	3.2
Ca (ppm)	551	525	525	582	525	674	525	753	661	615	342	588
<u>Location</u>												
1/4 Section	SE	NW	NW	NW	SW	NE	NE	NE	SE	NE	SE	SE
Section	18	17	34	34	34	10	10	15	10	15	23	23
Township (N)	10	10	11	11	11	10	10	10	10	10	10	10
Range (E)	27	27	27	27	27	27	27	27	27	27	27	27
Hill No.	-	-	1	1	1	-	2	2	2	2	-	-
<u>Flow Designation</u>												
Elephant Mtn.	X	X	X	X	X	X	X	X	X	X		X
Porona											X	



IRON, CALCIUM AND POTASSIUM CONTENT OF BASALT SAMPLES FROM THE HORN RAPIDS LINEAMENT.

FIGURE 14
JOHN A. BLUME & ASSOCIATES, ENGINEERS

GEOLOGIC STRUCTURE

Introduction

Two different aspects of geologic structure along the Horn Rapids Lineament are considered. One of these is a broader regional structure which is formed by the alignment of the hills. The other aspect of structure consists of the individual structure of each hill.

There are two linear features of the regional structure which are of particular interest. One of these linear features extends along the northeast sides of Hills 3, 4 and 5, all of which appear to be truncated abruptly along the northeast side. This truncation could have been accomplished by erosion but the well-defined linear aspect sides of the hills suggest also that it might have been caused by faulting or by erosion along a fault or fracture zone. The second linear feature of interest is a north-south trending bench which extends along the east side of Hill No. 2 and about 3 miles south of Hill No. 2. Both of these features can be best observed on the topographic map of Figure 1. Although there are no outcrops which completely transect either one of these features, some general observations can be made regarding them. If the linear feature passing near Hill No. 2 is due to a concealed fault, it is probable that the fault has little or no vertical displacement because basalt outcrops east of Hill No. 2 near the Yakima River show no apparent dislocation or offset. If the linear feature along Hills No. 3, 4, 5 and 6 is due to a concealed fault, it could not be continuous beyond Hill No. 3 because it would be exposed on the flanks of Hill No. 2 or in the basalt outcrops north of Hill No. 2 along the Yakima River. Furthermore, it may be noted that neither of these linear features trends toward the FFTF site.

The internal structure of the hills is poorly exposed except in the case of Hill No. 2, where basalt outcrops are fairly continuous

around the north end of the hill and the structure could be determined. In general, the evidence which was observed suggests that the basalt flows on the tops of the hills are more or less flat lying and the basalt flows around the sides of the hills dip outward away from the hilltops. Chemical analysis of samples collected from different parts of the hills shows that the flow which is exposed is the Elephant Mountain flow. One exception to this is an outcrop of the Pomona flow at Hill No. 1. This evidence is nowhere conclusive and is sometimes highly inconclusive. It is nevertheless compatible with the hypothesis that all of the hills are similar in their structure to Hill No. 2, which is a flat topped dome or anticline.

The hypothesized structure of the hills is not indicated on the geologic maps with the exception of Hill No. 2 where structural evidence is more conclusive. Hill No. 2 is shown on the map as an anticlinal structure. Cross sections shown on the geologic map, Plates III, IV and V illustrate the probable structure of the hills.

Seismic refraction surveys which would determine the profile of the buried basalt surface in various places have been recommended. This information would be helpful in defining the internal structure of the hills and in evaluating the larger linear features of the lineament.

Hill No. 1

Hill No. 1 is surrounded and partly covered by alluvial sand and gravel and by wind blown sand. The basalt flows strike about N 10W parallel to the long axes of the outcrops. Inclination, or dip, of the flows varies from about 5°E to 0. These flows probably represent the eastern flank of a very subdued domal uplift or anticline whose axes is situated somewhere to the west of Hill No. 1. The missing part of this structure was removed by erosion.

The basalt is broken in places by steep easterly trending fractures

0000000104

and small fracture zones. There is no evidence of movement along these fractures. Local outcrops of brecciated basalt occur along the west margin of the hill. This rock has the appearance of tectonic breccia as contrasted to a flow breccia. There is nothing to suggest that these isolated outcrops of breccia were produced by faulting or are related to faulting. It appears likely that the breccia was produced by folding and is similar to brecciated basalt at the north base of Hill No. 3.

All of the rocks exposed along the higher portion of Hill No. 1 belong to the Elephant Mountain flow or possibly the overlying Ward Gap flow. A small outcrop of basalt belonging to the Pomona flow is situated about 1 quarter of a mile west of Hill No. 1 nearer to the probable axis of the structure. The elevation of these outcrops is about 80 feet lower than that of basalt on top of Hill No. 1 and its location nearer the crest of the anticline is responsible for the flows being in relatively higher position.

Hill No. 2

Hill No. 2 is relatively well exposed at its northern end. Basalt flows dip gently away from the hill top to the northwest, north, and northeast. A small flexure strikes northeasterly from the hill top and is associated with northeasterly trending steep fractures. The top of the hill is about 1/2 mile long in its longest direction and is flat. Outcrops at the south end of Hill No. 2 are very sparse and the structure there cannot be determined. However, the topography of the south part of the hill is quite similar to that of the north part of the hill and the structure is probably similar. Hill No. 2 is a flat topped dome. Total vertical movement associated with uplift is on the order of 100 feet.

The basalt flows exposed on Hill No. 2 belong to the Elephant Mountain flow. The same flow is exposed on the Yakima River about 1/2 mile north of the hill and can be traced almost continuously exposed upstream on the south bank of the river for 5 or 6 miles. The base of the flow is exposed locally on the bank of the Yakima

slope of the hill appear to dip down the slope but this cannot be confirmed by measurement of reliable attitudes. The attitude of the breccia occurrence at the base of the north slope is not conformable to a postulated domal structure for Hill No. 3. There appears to be an easterly or northeasterly trending fold at the north end of Hill No. 3 which is responsible for the reversed dip at the base of the hill. Alternatively, the south dipping basalt could be a remnant of a hill, now eroded away, which once was situated north of Hill No. 3.

An Eroded Hill

Some subdued outcrops of basalt located to the north of Hill No. 3 are of particular interest. These outcrops are in the W/2 of the SE/4 of S. 23, T. 10N, R. 27E, about 1 mile north of Hill No. 3.

Two elongate, northwesterly trending outcrop areas are separated by a narrow strip of alluvium. The westernmost outcrop consists of Elephant Mountain basalt which strikes N 55W and dips 20° southwest. The easternmost area consists of dense, dark grey basalt which also strikes northwest and has a low south west dip. This rock is brecciated and is similar to the brecciated basalt on the north face of Hill No. 3. It has been identified as Pomona basalt.

The breccia pieces are angular, uncentred, and average about 4 inches in size. Individual pieces do not have baked surfaces or show evidences of rapid cooling as might be seen in a flow breccia. Many fractures are aligned in a common direction varying from N20 to N50 west and all fractures are steeply dipping. Individual fractures can rarely be traced more than a few feet and they often terminate abruptly on cross fractures. Many fracture surfaces are planar, while some are gently curved and others have steps of 1/8 or 1/4 inch thickness. No slickensiding or other evidence of movement could be found on fracture surfaces. The breccia zone is at least 150 ft wide perpendicular to the strike and is probably wider.

It is concluded that this breccia is similar to that at the base of Hill No. 3 and has been produced by folding. The two west dipping flows are one limb of an anticline or dome which has been eroded away. South dipping basalt near Hill No. 3 may be part of the same structure. The hill which once occupied this area probably extended to the east beyond the present site of the Yakima River. The probable structure existing today is shown on cross section C-C', geologic map Plate IV.

Hill No. 4

Hill No. 4 rises to an elevation of 760 ft above sea level or about 350 ft above the adjacent small community of West Richland. The scattered small outcrops on the north and east slopes of Hill No. 4 exhibit low to moderate dips away from the hill top. Analyses indicate that the basalt outcroppings are part of the Elephant Mountain flow. This evidence is not conclusive but suggests that the hill is an asymmetric domal structure. The hill is asymmetrical in that the south and west slopes are very gradual and flatten out on a relatively high plateau. The north and east sides of the hills slope down to much lower elevations and the basalt flows there dip below surficial deposits.

Hill No. 5

Hill No. 5, located 2 miles west of Richland rises to an elevation of about 630 ft, or 250 ft above the adjacent river valley. The northeast side of Hill No. 5 dips steeply down toward the Yakima River. The west side of Hill No. 5 slopes very gradually down to a broad plateau about 60 ft below the elevation of the hill top. The northwest and southeast slopes of the hill are gradual and the top is flat or gently undulating topography over a length of about 1/3 of a mile. Basalt outcrops are scattered over the northwest and southeast sides and on top of Hill No. 5. These rocks belong to the Elephant Mountain flow. Where dips can be observed on the flows, they dip gradually away from the hill top. Hill No. 5 appears to be an asymmetrical domal structure similar to Hill No. 4.

Hill No. 6

Hill No. 6 is a long low hill rising to an elevation of 745 ft at its south-easterly end. It is about 1-1/2 miles long in a north-westerly direction and about 1/2 mile wide. Basalt is widely exposed over most of the hill top. The rocks belong to the Elephant Mountain flow. An unusual phase of this flow is exposed over much of Hill No. 6 which consist of basalt blocks exposed and isolated by erosion. These blocks weather to irregular shapes which are 12 to 15 ft in diameter. The rock has a reddish-brown color and with poorly defined banded flow structures. Most of the structures observed in the basalt indicate very low dips but no reliable attitudes were obtained except in the quarry on the hill top at the south end where flat lying basalt flows are well exposed. Quarrying operations have exposed a 50-foot thick vertical section of the Elephant Mountain basalt which consists of an upper section 10 ft in thickness of columnar basalt, underlain by a 15-foot thickness of spherical weathering basalt, which in turn is underlain by a 25-foot section of columnar basalt. Samples taken from the uppermost and lowermost sections in the quarry indicate that the rock is probably the Elephant Mountain flow.

Basalt outcropping along the southwest margin of the hill is broken by many northwesterly trending steep fractures. These fractures parallel the margin of basalt outcropping along the edge of the hill in many places and suggest that the margin of outcrop itself, which is linear, may be controlled by a fracture zone. This possibility cannot be evaluated further because there are no outcrops within the possible fracture zone itself.

Although the evidence is inconclusive regarding the structure of Hill No. 6, it is postulated that the hill is an elongate symmetrical anticline plunging downward to the northwest and to the southeast.

INTRODUCTION

This report describes a reconnaissance field geologic investigation carried out in the area surrounding the Fast Flux Test Facility (FFTF) site near Richland, Washington. The work was required for earthquake engineering analysis of the site which was done by John A. Blume & Associates, Engineers (Blume) of San Francisco, California. This analysis was done for Battelle Memorial Institute, Pacific Northwest Laboratory (BNW) of Richland, Washington..

A preliminary report titled, "Preliminary Report: Seismic and Geologic Siting Evaluation and Seismic Design Criteria for the Fast Flux Test Facility Site near Richland, Washington," was submitted to BNW on December 1, 1969. This preliminary study was based upon a partial review of the literature, a site exploration program by others which was then in progress, and field work carried out by the United States Geologic Survey (USGS) which covered selected areas in and adjacent to the Hanford Reservation. The USGS field investigations were carried out in an area on Saddle Mountain, on Gable Mountain and Gable Butte, and near Wallula Gap. The final site evaluation report was submitted by Blume in August 1970.

Prior work has delineated the major structural trends in the region but has not included studies to evaluate the possible presence of minor faults in the vicinity of the site. Faults in the vicinity of the site might pass directly under the plant or very close to it and thus would present a hazard of damage to the structure due to possible surface movement on the fault. Therefore, this area was examined in order to determine insofar as possible whether faults are present which might pass under or very close to the facilities.

This work was carried out by reconnaissance geologic mapping of

SUMMARY AND CONCLUSIONS

A reconnaissance geologic survey of the area within 5 to 10 miles of the FFTF site has not revealed evidence of faulting which might pass directly under the site or close to it. With the exception of one minor fault, no known surface rupture or displacement by faulting was observed.

One minor fault was observed in the Yakima River gap striking N 20°E which would pass within 3 miles of the site at its nearest point if it continued northward beyond the slopes of Rattlesnake Mountain. No evidence was found to indicate that it does continue northward. It therefore does not constitute a hazard to the plant due to surface rupture.

A geologic survey of the Horn Rapids lineament was carried out which revealed an occurrence of brecciated basalt and an occurrence of fractured basalt. It was recommended that both of those tectonic structures be studied in more detail and that the lineament be mapped using 1:6,000 scale aerial photographs which were being obtained. It was noted that it might be advisable to augment future detailed studies of tectonic structures on the lineament with subsurface exploration. The detailed geologic mapping has since been completed and is described in Part I of this document. Conclusions based upon this more detailed mapping confirmed earlier work. No evidence of faulting which might pass beneath or very close to the FFTF facilities was found.

00000000194

topography. The dunes near the site, in contrast to dunes five to eight miles to the north which are largely active, are stabilized by a pervasive covering of sage brush, grass and a variety of small plants. All of the dunes, both active and stabilized, exhibit a marked linear orientation in directions varying from north 75° east in the northern part of the area to north 55° east in the southern part of the area.

A few man-made excavations have exposed stratified glaciofluvial sediments which lie beneath the wind blown sand. If faulting were present in these exposures it could be observed in the stratified materials because the strata would be disturbed and offset. No such evidence of faulting was observed during this survey.

If geologically young faulting of perhaps a few hundred years in age had caused surface rupture in the vicinity of the site some linear trace of the rupture might still be visible. No such linear traces could be observed.

The Eastern Area

The bluffs along the east bank of the Columbia River consist of well stratified, flat lying deposits of the Ringold Formation. A gravel and cobble conglomerate unit of the Ringold Formation outcrops prominently, extending from the water surface at elevation of about 345 ft up to elevation 455 and locally as high as 470 ft. These beds were examined where they outcrop almost continuously between Ringold Flats and the pumping plant about 8 miles south of Ringold Flats, and no evidence of disruption by faulting was observed.

Other prominent but less persistent beds crop out on the high cliffs above the conglomerate to elevations of 700 ft and higher. These strata were viewed from the river and no breaks or disruptions such as might be caused by faulting could be seen. The continuous exposures along the bluffs east of the site are illustrated in Figure 1. Locally, exposures on the cliffs are masked by alluvium, colluvium, or slide material where erosion of the cliffs by stream action or landsliding has occurred.

0-8888888888-98

BIBLIOGRAPHY

BIBLIOGRAPHY

Beck, G. F., Late Tertiary Stratigraphy and Paleontology of South-Central Washington and Adjacent Oregon (abst.), Geol. Soc. American Bull., Vol. 51, No. 12, Pt. 2, 1940.

Bingham, J. W., Londquist, C. J., Baltz, E. H., Geologic Investigation of Faulting in the Hanford Region, Washington, U.S. Geol. Survey open-file report, 1970.

Blume, John A., Earthquake Ground Motion and Engineering Procedures for Important Installations Near Active Faults, Proc. Third World Conf. on Earthquake Engineering, New Zealand, 1965.

Blume, John A., and Associates, Engineers, Geologic Reconnaissance in the Vicinity of the Fast Flux Test Facility Site at Hanford Washington, Report draft, June 1970.

Bonilla, M. G., Historic Surface Faulting in the Continental United States and Adjacent Parts of Mexico, U.S. Geol. Survey Interagency Rept., 1967.

Bonilla, M. G., Surface Faulting and Related Effects, in, Weigel, R. L., Earthquake Engineering, Englewood Cliffs, N. J., Prentiss Hall Inc., 518 P., 1970.

Bretz, J. H., H. T. V. Smith and G. E. Neff, Channeled Scablands of Washington: New Data and Interpretations, Geological Society of America Bull., Vol. 67, No. 8, 1956.

Brown, B. H., State Line Earthquake at Milton and Walla Walla, Seism. Soc. America Bull., Vol. 231, No. 3, 1937.

Brown, D. J., Personal Communication, 1970.

Brown, D. J., and Brown, R. E., Subsurface Eolian Deposit at the Hanford Project, Washington (abst.), Geol. Soc. America Bull., Vol. 69, No. 12, Pt. 2, 1958.

Brown, R. E., and Brown, D. J., The Surface of the Basalt Bedrock Beneath the Hanford Works Area, USAEC Rept. HW-57935, 1958.

Brown, R. E., An Introduction to the Surface of the Rinold Formation Beneath the Hanford Works Area, USAEC Rept. HW-66289, 1960.

Brown, R. E., and Raymond, J. R., Geophysical Evaluation at Hanford, USAEC Research and Development Rept. BNWL-47, 1964.

0 Brown, R. E., A Study of Reported Faulting in the Pasco Basin, USAEC Research and Development Rept. BNWL-662, 1968.

0 Brown, R. E., Personal communication, 1970.

0 Cantwell, T., Deep Resistivity Measurements in the Pacific Northwest, Jour. Geophys. Research, Vol. 70, No. 8, p. 1931-1937, 1965.

0 Clark, T. H., and Stearn, C. W., The Geologic Evolution of North America, Ronald Press Company, New York, 1960.

0 Crandell, D. R., Mullineaus, D. R., and Waldron, H. H., Age and Origin of the Puget Sound Trough in Western Washington, U.S. Geol. Survey Prof. Paper 525-B, 1965.

De Sitter, L. U., Structural Geology, McGraw-Hill Book Company, Inc., New York, 1956.

Eardly, A. J., Structural Geology of North America, Harper & Bros., New York, 1951.

Eaton, J. P., Personal communication, November 1969.

Eppley, R. A., Earthquake History of the United States, Part I, No. 41-1, U.S. Gov't. Printing Office, 1965.

Fifer, H. F., Earthquake Studies of the Hanford Area, Douglas United Nuclear, DUN-3625, February 1968.

Flint, R. F., Summary of Late Cenozoic Geology of Southeastern Washington, Amer. Jour. Sci., 5th series, Vol. 35, No. 207, 1938.

Fuller, R. E., Structural Features in the Columbia River Lavas of Central Washington, A Criticism, Jour. Geol. Vol. 42, 1934.

Gilluly, J., Waters, A. C., and Woodford, A. D., Principles of Geology, 3rd Edit., W. H. Freeman and Company, San Francisco, 1968.

Griggs, A. B., The Columbia Basin, in Mineral and Water Resources of Washington, U.S. Congress Committee Report prepared by the U.S.G.S., 1966.

Hamilton, Warren, and Myers, Cenozoic Tectonics of the Western United States, Review of Geophysics, Am. Geophys. Union, Vol. 4, No. 4, 1966.

Hammer, A. A., Rattlesnake Hills Gas Field, Benton County, Washington, Amer. Assoc. Petroleum Geologists Bull., Vol. 18, 1934.

Hodge, E. T., Columbia River Fault, Geol. Soc. America Bull., Vol. 42, 1931.

Hodge, E. T., Geology of the Lower Columbia River, Geol. Soc. America Bull., Vol. 49, 1938.

Hoffman, M. G., The Geology of Bald Butte Ridge, Jour. Geol. Vol. 40, No. 7, 1932.

Housner, G. W., Recommended Seismic Design Criteria for Hanford Nuclear Facility, Douglas United Nuclear, Inc. Rept. DUN-1403, 1966.

Jahns, R. H., Geologic Factors Relating to Engineering Seismology in the Hanford Area, Washington, Douglas United, Inc. Rept. DUN-3100, 1967.

Jones, Fred O. and Deacon, Robert J., Geology and Tectonic History of the Hanford Area, Douglas United Nuclear, DUN-1410, June 1966.

Kummel, B., History of the Earth, W. H. Freeman and Company, San Francisco, 1961.

Laval, W. N., Stratigraphy and Structural Geology of Portions of South-Central Washington, Thesis, Univ. of Washington, 1956.

Laval, W. N., Anticlines of the Southwestern Columbia Basin (abst.), Geol. Soc. America Bull., Vol. 69, No. 12, 1958.

Lowry, W. D., and Baldwin, E. M., Late Cenozoic Geology of the Lower Columbia River Valley, Geol. Soc. America, Vol. 63, 1952.

Lupher, R. L., Elastic Dikes of Columbia Basin Region, Geol. Soc. America Bull., Vol. 55, No. 12, 1944.

Merriam, J. C., Nature and Extent of Tertiary Formations Immediately Following the Columbia Lava Flows of the Northwest (abst.), Science (New Series), Vol. 80, 1934.

Neumann, F., Seismological Investigations at the Hanford Area, USAEC Rept. HW-63832, 1959.

Newcomb, R. C., Rinold Formation of Pleistocene Age in Type Locality, the White Bluffs, Washington, Amer. Jour., Sci., Vol. 256, 1958.

Newcomb, R. C., Geology and Groundwater Resources of the Walla Walla River Basin, Washington-Oregon, Wash. Div. of Water Res. Water Supply Bull., No. 1, 1965.

Newcomb, R. C., The Dalles-Umatilla Syncline, Oregon and Washington, U.S. Geological Survey Professional Paper 575-B, 1967.

Pakister, L. C., Structure of the Crust and Upper Mantle in the Western States, Jour. Geophys. Research, Vol. 68, No. 20, p. 5747-5756, 1963.

Plett, A. H., Preliminary Epicenter Map of Earthquakes in the Region of the AEC Hanford Works, U.S. Geological Survey, 1969.

Raisz, E., The Olympic-Wallowa Lineament, Amer. Jour. Sci., Vol. 243-A, 1945.

Rasmussen, M. H., Seismology Report on Washington, Idaho, Northern Oregon, and Western Montana and the Hanford Area, Washington, Douglas United Nuclear, Inc., Rept. DUN-1409, 1966.

Rasmussen, M. H., Washington State Earthquakes, 1840 through 1965, PISA Vol. 57, No. 3, June 1967.

Ross, R. J., Jr., Distribution of Ordovician Graptolites in Eugeosynclinal Facies in Western North America, and Its Paleogeographic Implications, Amer. Assoc. Petroleum Geologists Bull., Vol. 45, No. 3, 1961.

Russell, I. C., A Geological Reconnaissance in Central Washington, U.S. Geological Survey Bull., 108, 1893.

Ryall, A., Slemmons, D. B., and Gedney, L. D., Seismicity, Tectonism, and Surface Faulting in the Western United States During Historic Times, Bull. Seism. Soc. America, Vol. 56, No. 5, 1966.

Schmincke, H. V., Stratigraphy and Petrography of Four Upper Yakima Basalt Flows in South-Central Washington, Geol. Soc. America Bull., Vol. 28, P. 1385-1422, 1967.

Skehan, J. W., A Continental-Oceanic Crustal Boundary in the Pacific Northwest, U.S. Air Force Cambridge Research Lab. Rept. AFCRL-65-904, 1965.

Smith, G. O., Anticlinal Mountain Ridges in Central Washington, Jour. Geol. Vol. 11, 1903.

Snavely, P. D., Jr., and Wagner, H. C., Tertiary Geologic History of Western Oregon and Washington (abst.), Amer. Assoc. Petroleum Geologists Bull., Vol. 46, No. 2, 1962.

Taubeneck, W. H., An Evaluation of Tectonic Rotation In the Pacific Northwest, Jour. Geophys. Research, Vol. 70, No. 8, 1966.

Thayer, T. P., Some Relations of later Tertiary Volcanology and Structure In Eastern Oregon, Science, Vol. 124, 1956.

Uniform Building Code, Volume 1, 1967 Edition, International Conference of Building Officials, Pasadena, California.

U.S. Atomic Energy Commission, Nuclear Reactors and Earthquakes, TID-7024, 1963.

U.S. Atomic Energy Commission, Seismic and Geological Siting Criteria for Nuclear Power Plants, March 19, 1969, Tentative Regulatory Criteria.

Waldron, H. H., and Bonilla, M. G., Field Review of Possible Young Faulting In the Hanford Area, U.S. Geol. Survey Interagency Rept., 1968.

Waters, A. C., Volcanic Rocks and the Tectonic Cycle, Geol. Soc. America Spec. Paper 62, 1955.

Waters, A. C., Basalt Magma Types and Their Tectonic Associations, Pacific Northwest of the United States, Amer. Geophysical Union, Geophysical Monograph No. 6, p. 195, 1963.

Watkins, N. D., Paleomagnetism of the Columbia Plateaus, Jour. Geophys. Research, Vol. 70, 1965.

Weaver, C. E., Geology of Oregon and Washington and Its Relation to Occurrence of Oil and Gas, Amer. Assoc. Petroleum Geologists Bull., Vol. 29, 1955.

Weissenborn, A. E., Yates, R. G., Griggs, A. B., Wagner, H. C., and Snavely, P. D., Jr., Mineral and Water Resources of Washington, U.S. Geol. Survey report prepared in collaboration with the Washington State Dept. of Conservation and other agencies, p. 436, 1966.

Wiggins, J. H., Effect of Site Conditions on Earthquake Intensity, Jour. Structural Division, Proc. ASCE, Vol. 90, No. ST2, Part 1, 1964.

Willis, C. L., The Chiwaukum Graben, A Major Structure of Central Washington, Amer. Jour. Sci., Vol. 251, No. 11, 1953.

Wilson, B. W., Earthquake Occurrence and Effects in Ocean Areas, U.S.N. Civil Engineering Laboratory, Tech. Rept. CR-69.027, AD-859931, Feb. 1969.

Wise, D. U., An Outrageous Hypothesis for the Tectonic Pattern of the North American Cordillera, Geol. Soc. America Bull., Vol. 74, 1963.

0.000000205