



January 17, 2005

JAN 24 2005

United States Nuclear Regulatory Commission
Region IV
611 Ryan Plaza Drive
Suite 400
Arlington, Texas 78011

Dear Rachel S. Browder,

This letter is to provide an update on our progress in the decommissioning process of Carroll College materials license (25-07093-01) according to 10CFR30.36(C)(D).

We currently house the sealed sources that are indicated on the updated inventory (Table 1.), one source remains unidentified. Please note that initially 5 unknown sealed sources were indicated, however upon further review of the materials records with the former RSO, Father Harrington Ph.D. Four of the sources that I identified as sealed sources were actually homemade samples in which a drop of either CsCl or CoCl₂ was placed between two sheets of plastic and mounted on a small piece of cardboard. We have also drain disposed of all materials that are not check sources in the inventory. I have attached a copy of the drain disposal records, Table 2.

I have contacted one commercial company (qualtech) regarding disposal and plan to contact others. They cannot complete a cost estimate for disposal until all sources are identified. Additionally Bobby Walton from the state of California has expressed interest in a possible transfer of many of our sealed sources. The transfer would be from the Carroll college materials license to the license of the State of California. The major problem we are running into in the disposal is the unidentified source. I have not been able to locate anyone locally with a sodium iodide multichannel analyzer such that we could identify the unidentified sealed source.

Finally in regards to materials that were buried on the Carroll College Campus, June 30, 1961. I am including the RESRAD modeling output for those materials. Given that the materials buried at the site contained Iodine 131, with a half-life of 8.04 days, Gold 198 with a half-life of 2.69 days, and Carbon 14, with a half-life of 5,568 years. The calculations were only performed for Carbon 14.

I appreciate your time and consideration in these matters.



Sincerely,

A handwritten signature in black ink, appearing to read "Sam Alvey", with a long horizontal flourish extending to the right.

Sam Alvey

RSO

Carroll College

1601 N. Benton Ave.

Helena, MT 59625

406-447-4313

salvey@carroll.edu

Table 2. Radioactive Materials Disposal Log (Drain Disposal)

| Isotope | Chemical Form | Quantity and Date Received | Amount Disposed and Date | Volume of Water For Dilutions 24-Hours | Concentration of Material | Allowable Concentration |
|-------------------|-----------------|------------------------------|---------------------------|--|---|--|
| ^3H | Thymidine | 0.25 mCi 3-14-83 | 0.25 mCi 9-3-04 | 24,483 liters | 1.02×10^{-5} $\mu\text{Ci/mL}$ | 1.0×10^{-2} $\mu\text{Ci/mL}$ |
| ^{14}C | d-glucose | 50 μCi 1-22-86 | 50 μCi 10-4-04 | 24,483 liters | 2.04×10^{-6} $\mu\text{Ci/mL}$ | 3.0×10^{-4} $\mu\text{Ci/mL}$ |
| ^{14}C | Sodium acetate | 30 μCi 2-19-85 | 30 μCi 10-5-04 | 24,483 liters | 1.23×10^{-6} $\mu\text{Ci/mL}$ | 3.0×10^{-4} $\mu\text{Ci/mL}$ |
| ^{14}C | Palmitic acid | 15 μCi 2-03-86 | 15 μCi 10-5-04 | 24,483 liters | 6.12×10^{-7} $\mu\text{Ci/mL}$ | 3.0×10^{-4} $\mu\text{Ci/mL}$ |
| ^{14}C | Sodium heptate | 10 μCi 2-03-84 | 10 μCi 10-5-04 | 24,483 liters | 4.08×10^{-7} $\mu\text{Ci/mL}$ | 3.0×10^{-4} $\mu\text{Ci/mL}$ |
| ^{60}Co | CoCl_2 | 1 mCi 5-06-61 | | | | |
| | | | 0.5 mCi 9-7-04 | 24,483 liters | 2.04×10^{-5} $\mu\text{Ci/mL}$ | 3.0×10^{-5} $\mu\text{Ci/mL}$ |
| | | | 0.5 mCi 10-11-04 | 24,483 liters | 2.04×10^{-5} $\mu\text{Ci/mL}$ | 3.0×10^{-5} $\mu\text{Ci/mL}$ |
| ^{137}Cs | CsCl | 1 mCi 2-25-61 | | | | |
| | | | 0.2 mCi 8-3-04 | 24,483 liters | 8.17×10^{-6} $\mu\text{Ci/mL}$ | 1.0×10^{-5} $\mu\text{Ci/mL}$ |
| | | | 0.2 mCi 9-7-04 | 24,483 liters | 8.17×10^{-6} $\mu\text{Ci/mL}$ | 1.0×10^{-5} $\mu\text{Ci/mL}$ |

| | | | | | | | |
|--|--|--|---------|---------|---------------|-----------------------------------|----------------------------------|
| | | | 0.2 mCi | 10-5-04 | 24,483 liters | 8.17×10^{-6} μ Ci/mL | 1.0×10^{-5} μ Ci/mL |
| | | | 0.2 mCi | 11-2-04 | 24,483 liters | 8.17×10^{-6} μ Ci/mL | 1.0×10^{-5} μ Ci/mL |
| | | | 0.2 mCi | 12-7-04 | 24,483 liters | 8.17×10^{-6} μ Ci/mL | 1.0×10^{-5} μ Ci/mL |

* Isotopes were disposed down the drain of a stainless steel sink in Simperman Hall room 218 with the tap running (measured at approximately 7.5 liters per minute). The tap remained on for 1-hour each time material was disposed of. The waste line from the sink joins into the building sewer line that connects to the sewer lines from two dormitories on the Carroll College campus. The mixture of effluent from these three buildings is discharged into the city sewer system. Based upon water meter readings for these buildings during the lowest water usage months of the year approximately 24,483 liters of water was estimated to leave the campus through this sewer pipe in a 24 hour period. This value (24,483 liters) was used to estimate the dilution of radioactive material such that the material could be drain disposed under NRC guidelines.

Section 5. Radioactive Material

Inventory of stored isotopes stored in the lead brick housing within in the locked cabinet below the fume hood in room 218 Simperman Hall. If the date of receipt is not included in the table below it is not known at this time. All quantities currently held by Carroll College are the maximum amounts that will be stored on site until they can be disposed of through the decommissioning process and termination of license 25-07093-01. No new radioactive materials will be acquired by Carroll College.

Table 1. Radioactive Materials Inventory 2005

| Isotope | Chemical Form | Quantity and Date Recieved | Maximum Amount Housed | Disposal |
|-------------------|-------------------|----------------------------|-----------------------|----------|
| ³ H | Thymidine | 0.25 mCi 3-14-83 | CAO* | Disposed |
| ¹⁴ C | d-glucose | 50 μCi 1-22-86 | CAO | Disposed |
| ¹⁴ C | Sodium acetate | 30 μCi 2-19-85 | CAO | Disposed |
| ¹⁴ C | Palmitic acid | 15 μCi 2-03-86 | CAO | Disposed |
| ¹⁴ C | Sodium hectate | 10 μCi 2-03-84 | CAO | Disposed |
| ⁶⁰ Co | CoCl ₂ | 1 mCi 5-06-61 | CAO | Disposed |
| ¹³⁷ Cs | CsCl | 1 mCi 2-25-61 | CAO | Disposed |
| ⁶⁰ Co | Sealed Source | Pre 1960 (donation) | CAO | |
| Uranium | Uranium Nitrate | Approx. 1.3 lbs | CAO | |
| Uranium | Uranium Acetate | Approx 1.0 lbs | CAO | |
| ¹³⁷ Cs | Sealed Source | Pre 1960 (donation) | CAO | |
| ¹³⁷ Cs | Sealed Source | 0.05 μCi | CAO | |
| ¹³⁷ Cs | Sealed Source | 0.5 μCi | CAO | |
| ⁶⁰ Co | Sealed Source | 0.5 μCi | CAO | |
| ¹³³ Ba | Sealed Source | 0.05 μCi | CAO | |
| ¹³⁷ Cs | Sealed Source | 0.05 μCi | CAO | |
| ¹³⁷ Cs | Sealed Source | 0.5 μCi | CAO | |
| ⁶⁰ Co | Sealed Source | 0.05 μCi | CAO | |
| ⁶⁰ Co | Sealed Source | 0.5 μCi | CAO | |
| ¹³³ Ba | Sealed Source | 0.05 μCi | CAO | |
| ¹³³ Ba | Sealed Source | 0.5 μCi | CAO | |
| ¹³³ Ba | Sealed Source | 0.5 μCi | CAO | |
| ¹⁴ C | Sealed Source | 17,000 cpm | CAO | |
| ¹⁴ C | Sealed Source | Unkown | CAO | |
| ⁹⁰ Sr | Sealed Source | Unknown | CAO | |
| ²¹⁰ Pb | Sealed Source | Unknown | CAO | |
| ²³⁸ U | Sealed Source | Unknown | CAO | |
| Radium | Sealed Source | 1.0 μCi | CAO | |
| Radium | Sealed Source | Unknown | CAO | |
| | Sealed Source | Unknown | CAO | |
| Unk 1 | Dry sample | Unknown | CAO | Disposed |
| Unk 2 | Dry sample | Unknown | CAO | Disposed |

| | | | | |
|-------------------|---------------|---------|-----|----------|
| Unk 3 | Dry sample | Unknown | CAO | Disposed |
| Unk 4 | Dry sample | Unknown | CAO | Disposed |
| Unk 5 | Sealed Source | Unknown | CAO | |
| ⁶⁰ Co | Sealed Source | 1.0 μCi | CAO | |
| ²⁰⁴ Ti | Sealed Source | 0.8 μCi | CAO | |
| ²⁰⁴ Ti | Sealed Source | 0.8 μCi | CAO | |
| ²⁰⁴ Ti | Sealed Source | 0.8 μCi | CAO | |
| ²¹⁰ Po | Sealed Source | 0.1 μCi | CAO | |
| ¹³⁷ Cs | Sealed Source | Unknown | CAO | |
| ⁹⁰ Sr | Sealed Source | 0.1 μCi | CAO | |

Unknowns 1-4 were small spots of CoCl₂ or CsCl placed between two sheets of plastic and used in student experiments in the early 1960's. Since they were a very small amount of the total quantity received the dried materials (spot) were reconstituted in water and disposed of down the drain as part of the CoCl₂ or CsCl disposals.

Unk 5 has still not been identified but is a sealed source.

CAO = Current Amount Only, no additional materials will be obtained or housed at Carroll College.

Table of Contents

Part I: Mixture Sums and Single Radionuclide Guidelines

| | |
|--|----|
| Dose Conversion Factor (and Related) Parameter Summary ... | 2 |
| Site-Specific Parameter Summary | 3 |
| Summary of Pathway Selections | 7 |
| Contaminated Zone and Total Dose Summary | 8 |
| Total Dose Components | |
| Time = 0.000E+00 | 9 |
| Time = 1.000E+00 | 10 |
| Time = 1.000E+01 | 11 |
| Time = 2.000E+01 | 12 |
| Time = 4.000E+01 | 13 |
| Time = 1.000E+02 | 14 |
| Dose/Source Ratios Summed Over All Pathways | 15 |
| Single Radionuclide Soil Guidelines | 15 |
| Dose Per Nuclide Summed Over All Pathways | 16 |
| Soil Concentration Per Nuclide | 16 |

Dose Conversion Factor (and Related) Parameter Summary
 File: FGR 13 Morbidity

| Menu | Parameter | Current Value | Default | Pa |
|------|--|---------------|-----------|-------|
| B-1 | Dose conversion factors for inhalation, mrem/μCi: | | | |
| B-1 | C-14 | 2.090E+00 | 2.090E+00 | ✓ DCF |
| D-1 | Dose conversion factors for ingestion, mrem/μCi: | | | |
| D-1 | C-14 | 2.090E+00 | 2.090E+00 | ✓ DCF |
| D-34 | Food transfer factors: | | | |
| D-34 | C-14 , plant/soil concentration ratio, dimensionless | 5.500E+00 | 5.500E+00 | ✓ RTF |
| D-34 | C-14 , beef/livestock-intake ratio, (μCi/kg)/(μCi/d) | 3.100E-02 | 3.100E-02 | ✓ RTF |
| D-34 | C-14 , milk/livestock-intake ratio, (μCi/L)/(μCi/d) | 1.200E-02 | 1.200E-02 | ✓ RTF |
| D-5 | Bioaccumulation factors, fresh water, L/kg: | | | |
| D-5 | C-14 , fish | 5.000E+04 | 5.000E+04 | ✓ BIO |
| D-5 | C-14 , crustacea and mollusks | 9.100E+03 | 9.100E+03 | ✓ BIO |

Site-Specific Parameter Summary

| Menu | Parameter | User Input | Default | Used b (If different) |
|------|---|------------|-----------|-----------------------|
| R011 | Area of contaminated zone (m**2) | 1.000E-01 | 1.000E+04 | - |
| R011 | Thickness of contaminated zone (m) | 2.000E+00 | 2.000E+00 | ✓ |
| R011 | Length parallel to aquifer flow (m) | 1.000E+02 | 1.000E+02 | ✓ |
| R011 | Basic radiation dose limit (mrem/yr) | 2.500E+01 | 2.500E+01 | ✓ |
| R011 | Time since placement of material (yr) | 0.000E+00 | 0.000E+00 | - |
| R011 | Times for calculations (yr) | 1.000E+00 | 1.000E+00 | - |
| R011 | Times for calculations (yr) | 1.000E+01 | 3.000E+00 | - |
| R011 | Times for calculations (yr) | 2.000E+01 | 1.000E+01 | - |
| R011 | Times for calculations (yr) | 4.000E+01 | 3.000E+01 | - |
| R011 | Times for calculations (yr) | 1.000E+02 | 1.000E+02 | - |
| R011 | Times for calculations (yr) | not used | 3.000E+02 | - |
| R011 | Times for calculations (yr) | not used | 1.000E+03 | - |
| R011 | Times for calculations (yr) | not used | 0.000E+00 | - |
| R011 | Times for calculations (yr) | not used | 0.000E+00 | - |
| R012 | Initial principal radionuclide (µCi/g): C-14 | 2.500E-01 | 0.000E+00 | - |
| R012 | Concentration in groundwater (µCi/L): C-14 | not used | 0.000E+00 | - |
| R013 | Cover depth (m) | 1.200E+00 | 0.000E+00 | - |
| R013 | Density of cover material (g/cm**3) | 1.500E+00 | 1.500E+00 | - |
| R013 | Cover depth erosion rate (m/yr) | 1.000E-03 | 1.000E-03 | - |
| R013 | Density of contaminated zone (g/cm**3) | 1.500E+00 | 1.500E+00 | - |
| R013 | Contaminated zone erosion rate (m/yr) | 1.000E-03 | 1.000E-03 | - |
| R013 | Contaminated zone total porosity | 4.000E-01 | 4.000E-01 | - |
| R013 | Contaminated zone field capacity | 2.000E-01 | 2.000E-01 | - |
| R013 | Contaminated zone hydraulic conductivity (m/yr) | 1.000E+01 | 1.000E+01 | - |
| R013 | Contaminated zone b parameter | 5.300E+00 | 5.300E+00 | - |
| R013 | Average annual wind speed (m/sec) | 2.000E+00 | 2.000E+00 | - |
| R013 | Humidity in air (g/m**3) | not used | 8.000E+00 | - |
| R013 | Evapotranspiration coefficient | 5.000E-01 | 5.000E-01 | - |
| R013 | Precipitation (m/yr) | 1.000E+00 | 1.000E+00 | - |
| R013 | Irrigation (m/yr) | 2.000E-01 | 2.000E-01 | - |
| R013 | Irrigation mode | overhead | overhead | - |
| R013 | Runoff coefficient | 2.000E-01 | 2.000E-01 | - |
| R013 | Watershed area for nearby stream or pond (m**2) | 1.000E+06 | 1.000E+06 | - |
| R013 | Accuracy for water/soil computations | 1.000E-03 | 1.000E-03 | - |
| R014 | Density of saturated zone (g/cm**3) | 1.500E+00 | 1.500E+00 | ✓ |
| R014 | Saturated zone total porosity | 4.000E-01 | 4.000E-01 | - |
| R014 | Saturated zone effective porosity | 2.000E-01 | 2.000E-01 | - |
| R014 | Saturated zone field capacity | 2.000E-01 | 2.000E-01 | - |
| R014 | Saturated zone hydraulic conductivity (m/yr) | 1.000E+02 | 1.000E+02 | - |
| R014 | Saturated zone hydraulic gradient | 2.000E-02 | 2.000E-02 | - |
| R014 | Saturated zone b parameter | 5.300E+00 | 5.300E+00 | - |
| R014 | Water table drop rate (m/yr) | 1.000E-03 | 1.000E-03 | - |
| R014 | Well pump intake depth (m below water table) | 1.000E+01 | 1.000E+01 | - |
| R014 | Model: Nondispersion (ND) or Mass-Balance (MB) | ND | ND | - |
| R014 | Well pumping rate (m**3/yr) | 2.500E+02 | 2.500E+02 | - |
| R015 | Number of unsaturated zone strata | 1 | 1 | ✓ |

Assuming coffee can size is 200g - this is reasonable and conservative. RSBrowder

Site-Specific Parameter Summary (continued)

| Menu | Parameter | User Input | Default | Used b (If different) |
|------|--|------------|-----------|--------------------------|
| R015 | Unsat. zone 1, thickness (m) | 4.000E+00 | 4.000E+00 | ✓ - |
| R015 | Unsat. zone 1, soil density (g/cm**3) | 1.500E+00 | 1.500E+00 | - |
| R015 | Unsat. zone 1, total porosity | 4.000E-01 | 4.000E-01 | - |
| R015 | Unsat. zone 1, effective porosity | 2.000E-01 | 2.000E-01 | - |
| R015 | Unsat. zone 1, field capacity | 2.000E-01 | 2.000E-01 | - |
| R015 | Unsat. zone 1, soil-specific b parameter | 5.300E+00 | 5.300E+00 | - |
| R015 | Unsat. zone 1, hydraulic conductivity (m/yr) | 1.000E+01 | 1.000E+01 | ✓ - |
| R016 | Distribution coefficients for C-14 | | | |
| R016 | Contaminated zone (cm**3/g) | 0.000E+00 | 0.000E+00 | - |
| R016 | Unsat. zone 1 (cm**3/g) | 0.000E+00 | 0.000E+00 | - |
| R016 | Saturated zone (cm**3/g) | 0.000E+00 | 0.000E+00 | - |
| R016 | Leach rate (/yr) | 0.000E+00 | 0.000E+00 | 7.79 |
| R016 | Solubility constant | 0.000E+00 | 0.000E+00 | not |
| R017 | Inhalation rate (m**3/yr) | 8.400E+03 | 8.400E+03 | ✓ - |
| R017 | Mass loading for inhalation (g/m**3) | 1.000E-04 | 1.000E-04 | ✓ - |
| R017 | Exposure duration | 4.500E+01 | 3.000E+01 | - |
| R017 | Shielding factor, inhalation | 4.000E-01 | 4.000E-01 | - |
| R017 | Shielding factor, external gamma | 7.000E-01 | 7.000E-01 | - |
| R017 | Fraction of time spent indoors | 5.000E-01 | 5.000E-01 | - |
| R017 | Fraction of time spent outdoors (on site) | 2.500E-01 | 2.500E-01 | - |
| R017 | Shape factor flag, external gamma | 1.000E+00 | 1.000E+00 | >0 shows ci |
| R017 | Radii of shape factor array (used if FS = -1): | | | |
| R017 | Outer annular radius (m), ring 1: | not used | 5.000E+01 | - |
| R017 | Outer annular radius (m), ring 2: | not used | 7.071E+01 | - |
| R017 | Outer annular radius (m), ring 3: | not used | 0.000E+00 | - |
| R017 | Outer annular radius (m), ring 4: | not used | 0.000E+00 | - |
| R017 | Outer annular radius (m), ring 5: | not used | 0.000E+00 | - |
| R017 | Outer annular radius (m), ring 6: | not used | 0.000E+00 | - |
| R017 | Outer annular radius (m), ring 7: | not used | 0.000E+00 | - |
| R017 | Outer annular radius (m), ring 8: | not used | 0.000E+00 | - |
| R017 | Outer annular radius (m), ring 9: | not used | 0.000E+00 | - |
| R017 | Outer annular radius (m), ring 10: | not used | 0.000E+00 | - |
| R017 | Outer annular radius (m), ring 11: | not used | 0.000E+00 | - |
| R017 | Outer annular radius (m), ring 12: | not used | 0.000E+00 | - |
| R017 | Fractions of annular areas within AREA: | | | |
| R017 | Ring 1 | not used | 1.000E+00 | - |
| R017 | Ring 2 | not used | 2.732E-01 | - |
| R017 | Ring 3 | not used | 0.000E+00 | - |
| R017 | Ring 4 | not used | 0.000E+00 | - |
| R017 | Ring 5 | not used | 0.000E+00 | - |
| R017 | Ring 6 | not used | 0.000E+00 | - |
| R017 | Ring 7 | not used | 0.000E+00 | - |
| R017 | Ring 8 | not used | 0.000E+00 | - |
| R017 | Ring 9 | not used | 0.000E+00 | - |
| R017 | Ring 10 | not used | 0.000E+00 | - |
| R017 | Ring 11 | not used | 0.000E+00 | - |
| R017 | Ring 12 | not used | 0.000E+00 | - |

Site-Specific Parameter Summary (continued)

| Menu | Parameter | User Input | Default | Used b (If different) |
|------|--|------------|-----------|--------------------------|
| R018 | Fruits, vegetables and grain consumption (kg/yr) | 1.600E+02 | 1.600E+02 | ✓ - |
| R018 | Leafy vegetable consumption (kg/yr) | 1.400E+01 | 1.400E+01 | - |
| R018 | Milk consumption (L/yr) | 9.200E+01 | 9.200E+01 | - |
| R018 | Meat and poultry consumption (kg/yr) | 6.300E+01 | 6.300E+01 | - |
| R018 | Fish consumption (kg/yr) | 5.400E+00 | 5.400E+00 | - |
| R018 | Other seafood consumption (kg/yr) | 9.000E-01 | 9.000E-01 | - |
| R018 | Soil ingestion rate (g/yr) | 3.650E+01 | 3.650E+01 | - |
| R018 | Drinking water intake (L/yr) | 5.100E+02 | 5.100E+02 | - |
| R018 | Contamination fraction of drinking water | 1.000E+00 | 1.000E+00 | - |
| R018 | Contamination fraction of household water | not used | 1.000E+00 | - |
| R018 | Contamination fraction of livestock water | 1.000E+00 | 1.000E+00 | - |
| R018 | Contamination fraction of irrigation water | 1.000E+00 | 1.000E+00 | - |
| R018 | Contamination fraction of aquatic food | 5.000E-01 | 5.000E-01 | ✓ - |
| R018 | Contamination fraction of plant food | -1 | -1 | 0.50 |
| R018 | Contamination fraction of meat | -1 | -1 | 0.50 |
| R018 | Contamination fraction of milk | -1 | -1 | 0.50 |
| R019 | Livestock fodder intake for meat (kg/day) | 6.800E+01 | 6.800E+01 | ✓ - |
| R019 | Livestock fodder intake for milk (kg/day) | 5.500E+01 | 5.500E+01 | - |
| R019 | Livestock water intake for meat (L/day) | 5.000E+01 | 5.000E+01 | - |
| R019 | Livestock water intake for milk (L/day) | 1.600E+02 | 1.600E+02 | - |
| R019 | Livestock soil intake (kg/day) | 5.000E-01 | 5.000E-01 | - |
| R019 | Mass loading for foliar deposition (g/m**3) | 1.000E-04 | 1.000E-04 | - |
| R019 | Depth of soil mixing layer (m) | 1.500E-01 | 1.500E-01 | - |
| R019 | Depth of roots (m) | 9.000E-01 | 9.000E-01 | - |
| R019 | Drinking water fraction from ground water | 1.000E+00 | 1.000E+00 | - |
| R019 | Household water fraction from ground water | not used | 1.000E+00 | - |
| R019 | Livestock water fraction from ground water | 1.000E+00 | 1.000E+00 | - |
| R019 | Irrigation fraction from ground water | 1.000E+00 | 1.000E+00 | - |
| R19B | Wet weight crop yield for Non-Leafy (kg/m**2) | 7.000E-01 | 7.000E-01 | ✓ - |
| R19B | Wet weight crop yield for Leafy (kg/m**2) | 1.500E+00 | 1.500E+00 | - |
| R19B | Wet weight crop yield for Fodder (kg/m**2) | 1.100E+00 | 1.100E+00 | - |
| R19B | Growing Season for Non-Leafy (years) | 1.700E-01 | 1.700E-01 | - |
| R19B | Growing Season for Leafy (years) | 2.500E-01 | 2.500E-01 | - |
| R19B | Growing Season for Fodder (years) | 8.000E-02 | 8.000E-02 | - |
| R19B | Translocation Factor for Non-Leafy | 1.000E-01 | 1.000E-01 | - |
| R19B | Translocation Factor for Leafy | 1.000E+00 | 1.000E+00 | - |
| R19B | Translocation Factor for Fodder | 1.000E+00 | 1.000E+00 | - |
| R19B | Dry Foliar Interception Fraction for Non-Leafy | 2.500E-01 | 2.500E-01 | - |
| R19B | Dry Foliar Interception Fraction for Leafy | 2.500E-01 | 2.500E-01 | - |
| R19B | Dry Foliar Interception Fraction for Fodder | 2.500E-01 | 2.500E-01 | - |
| R19B | Wet Foliar Interception Fraction for Non-Leafy | 2.500E-01 | 2.500E-01 | - |
| R19B | Wet Foliar Interception Fraction for Leafy | 2.500E-01 | 2.500E-01 | - |
| R19B | Wet Foliar Interception Fraction for Fodder | 2.500E-01 | 2.500E-01 | - |
| R19B | Weathering Removal Constant for Vegetation | 2.000E+01 | 2.000E+01 | - |
| C14 | C-12 concentration in water (g/cm**3) | 2.000E-05 | 2.000E-05 | ✓ - |
| C14 | C-12 concentration in contaminated soil (g/g) | 3.000E-02 | 3.000E-02 | - |
| C14 | Fraction of vegetation carbon from soil | 2.000E-02 | 2.000E-02 | ✓ - |
| C14 | Fraction of vegetation carbon from air | 9.800E-01 | 9.800E-01 | - |

Site-Specific Parameter Summary (continued)

| Menu | Parameter | User Input | Default | Used b (If different) |
|------|--|------------|------------|--------------------------|
| C14 | C-14 evasion layer thickness in soil (m) | 3.000E-01 | 3.000E-01 | - |
| C14 | C-14 evasion flux rate from soil (1/sec) | 7.000E-07 | 7.000E-07 | - |
| C14 | C-12 evasion flux rate from soil (1/sec) | 1.000E-10 | 1.000E-10 | - |
| C14 | Fraction of grain in beef cattle feed | 8.000E-01 | 8.000E-01 | - |
| C14 | Fraction of grain in milk cow feed | 2.000E-01 | 2.000E-01 | - |
| C14 | DCF correction factor for gaseous forms of C14 | 8.894E+01 | 8.894E+01 | - |
| STOR | Storage times of contaminated foodstuffs (days): | | | |
| STOR | Fruits, non-leafy vegetables, and grain | 1.400E+01 | 1.400E+01 | - |
| STOR | Leafy vegetables | 1.000E+00 | 1.000E+00 | - |
| STOR | Milk | 1.000E+00 | 1.000E+00 | - |
| STOR | Meat and poultry | 2.000E+01 | 2.000E+01 | - |
| STOR | Fish | 7.000E+00 | 7.000E+00 | - |
| STOR | Crustacea and mollusks | 7.000E+00 | 7.000E+00 | - |
| STOR | Well water | 1.000E+00 | 1.000E+00 | - |
| STOR | Surface water | 1.000E+00 | 1.000E+00 | - |
| STOR | Livestock fodder | 4.500E+01 | 4.500E+01 | - |
| R021 | Thickness of building foundation (m) | not used | 1.500E-01 | - |
| R021 | Bulk density of building foundation (g/cm**3) | not used | 2.400E+00 | - |
| R021 | Total porosity of the cover material | not used | 4.000E-01 | - |
| R021 | Total porosity of the building foundation | not used | 1.000E-01 | - |
| R021 | Volumetric water content of the cover material | not used | 5.000E-02 | - |
| R021 | Volumetric water content of the foundation | not used | 3.000E-02 | - |
| R021 | Diffusion coefficient for radon gas (m/sec): | | | |
| R021 | in cover material | not used | 2.000E-06 | - |
| R021 | in foundation material | not used | 3.000E-07 | - |
| R021 | in contaminated zone soil | not used | 2.000E-06 | - |
| R021 | Radon vertical dimension of mixing (m) | not used | 2.000E+00 | - |
| R021 | Average building air exchange rate (1/hr) | not used | 5.000E-01 | - |
| R021 | Height of the building (room) (m) | not used | 2.500E+00 | - |
| R021 | Building interior area factor | not used | 0.000E+00 | - |
| R021 | Building depth below ground surface (m) | not used | -1.000E+00 | - |
| R021 | Emanating power of Rn-222 gas | not used | 2.500E-01 | - |
| R021 | Emanating power of Rn-220 gas | not used | 1.500E-01 | - |
| TITL | Number of graphical time points | 32 | --- | - |
| TITL | Maximum number of integration points for dose | 17 | --- | - |
| TITL | Maximum number of integration points for risk | 257 | --- | - |

Summary of Pathway Selections

| Pathway | User Selection |
|-----------------------------|------------------------|
| 1 -- external gamma | active |
| 2 -- inhalation (w/o radon) | active |
| 3 -- plant ingestion | active |
| 4 -- meat ingestion | active |
| 5 -- milk ingestion | active |
| 6 -- aquatic foods | active |
| 7 -- drinking water | active |
| 8 -- soil ingestion | active |
| 9 -- radon | suppressed ✓ <i>AL</i> |
| Find peak pathway doses | active |

| Contaminated Zone Dimensions | | Initial Soil Concentrations, µCi/g | |
|------------------------------|--------------------|------------------------------------|-----------|
| Area: | 0.10 square meters | C-14 | 2.500E-01 |
| Thickness: | 2.00 meters | | |
| Cover Depth: | 1.20 meters | | |

Total Dose TDOSE(t), mrem/yr

Basic Radiation Dose Limit = 2.500E+01 mrem/yr

Total Mixture Sum M(t) = Fraction of Basic Dose Limit Received at Time (t)

| | | | | | | |
|------------|-----------|-----------|-----------|-----------|-----------|-----------|
| t (years): | 0.000E+00 | 1.000E+00 | 1.000E+01 | 2.000E+01 | 4.000E+01 | 1.000E+02 |
| TDOSE(t): | 9.506E-24 | 7.046E+00 | 7.269E-01 | 3.011E-04 | 5.164E-11 | 0.000E+00 |
| M(t): | 3.802E-25 | 2.818E-01 | 2.908E-02 | 1.204E-05 | 2.066E-12 | 0.000E+00 |

Maximum TDOSE(t): 3.996E+01 mrem/yr at t = 4.286 ± 0.009 years

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and (p) as mrem/yr and Fraction of Total Dose At t = 4.286E+00 years

Water Independent Pathways (Inhalation excludes radon)

| Radio- Nuclide Nuclide | Ground | | Inhalation | | Radon | | Plant | | Meat | |
|------------------------------|-----------|--------|------------|--------|-----------|--------|-----------|--------|-----------|--------|
| | mrem/yr | fract. | mrem/yr | fract. | mrem/yr | fract. | mrem/yr | fract. | mrem/yr | fract. |
| C-14 | 4.071E-25 | 0.0000 | 0.000E+00 | 0.0000 | 0.000E+00 | 0.0000 | 0.000E+00 | 0.0000 | 0.000E+00 | 0.0000 |
| Total | 4.071E-25 | 0.0000 | 0.000E+00 | 0.0000 | 0.000E+00 | 0.0000 | 0.000E+00 | 0.0000 | 0.000E+00 | 0.0000 |

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and (p) as mrem/yr and Fraction of Total Dose At t = 4.286E+00 years

Water Dependent Pathways

| Radio- Nuclide Nuclide | Water | | Fish | | Radon | | Plant | | Meat | |
|------------------------------|-----------|--------|-----------|--------|-----------|--------|-----------|--------|-----------|--------|
| | mrem/yr | fract. |
| C-14 | 2.981E+01 | 0.7459 | 1.015E+01 | 0.2541 | 0.000E+00 | 0.0000 | 2.179E-05 | 0.0000 | 9.819E-06 | 0.0000 |
| Total | 2.981E+01 | 0.7459 | 1.015E+01 | 0.2541 | 0.000E+00 | 0.0000 | 2.179E-05 | 0.0000 | 9.819E-06 | 0.0000 |

*Sum of all water independent and dependent pathways.

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and
 As mrem/yr and Fraction of Total Dose At t = 0.000E+00 years

Water Independent Pathways (Inhalation excludes radon)

| Radio- Nuclide | Ground | | Inhalation | | Radon | | Plant | | Meat | |
|-------------------|------------------|---------------|------------------|---------------|------------------|---------------|------------------|---------------|------------------|---------------|
| | mrem/yr | fract. |
| C-14 | 9.506E-24 | 1.0000 | 0.000E+00 | 0.0000 | 0.000E+00 | 0.0000 | 0.000E+00 | 0.0000 | 0.000E+00 | 0.0000 |
| Total | 9.506E-24 | 1.0000 | 0.000E+00 | 0.0000 | 0.000E+00 | 0.0000 | 0.000E+00 | 0.0000 | 0.000E+00 | 0.0000 |

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and
 As mrem/yr and Fraction of Total Dose At t = 0.000E+00 years

Water Dependent Pathways

| Radio- Nuclide | Water | | Fish | | Radon | | Plant | | Meat | |
|-------------------|------------------|---------------|------------------|---------------|------------------|---------------|------------------|---------------|------------------|---------------|
| | mrem/yr | fract. |
| C-14 | 0.000E+00 | 0.0000 |
| Total | 0.000E+00 | 0.0000 |

*Sum of all water independent and dependent pathways.

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and As mrem/yr and Fraction of Total Dose At t = 1.000E+00 years

Water Independent Pathways (Inhalation excludes radon)

| Radio-Nuclide | Ground | | Inhalation | | Radon | | Plant | | Meat | |
|---------------|-----------|--------|------------|--------|-----------|--------|-----------|--------|-----------|--------|
| | mrem/yr | fract. | mrem/yr | fract. | mrem/yr | fract. | mrem/yr | fract. | mrem/yr | fract. |
| C-14 | 4.558E-24 | 0.0000 | 0.000E+00 | 0.0000 | 0.000E+00 | 0.0000 | 0.000E+00 | 0.0000 | 0.000E+00 | 0.0000 |
| Total | 4.558E-24 | 0.0000 | 0.000E+00 | 0.0000 | 0.000E+00 | 0.0000 | 0.000E+00 | 0.0000 | 0.000E+00 | 0.0000 |

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and As mrem/yr and Fraction of Total Dose At t = 1.000E+00 years

Water Dependent Pathways

| Radio-Nuclide | Water | | Fish | | Radon | | Plant | | Meat | |
|---------------|-----------|--------|-----------|--------|-----------|--------|-----------|--------|-----------|--------|
| | mrem/yr | fract. |
| C-14 | 5.311E+00 | 0.7537 | 1.735E+00 | 0.2463 | 0.000E+00 | 0.0000 | 3.525E-06 | 0.0000 | 1.503E-06 | 0.0000 |
| Total | 5.311E+00 | 0.7537 | 1.735E+00 | 0.2463 | 0.000E+00 | 0.0000 | 3.525E-06 | 0.0000 | 1.503E-06 | 0.0000 |

*Sum of all water independent and dependent pathways.

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and
 As mrem/yr and Fraction of Total Dose At t = 1.000E+01 years

Water Independent Pathways (Inhalation excludes radon)

| Radio- Nuclide | Ground | | Inhalation | | Radon | | Plant | | Meat | |
|-------------------|------------------|---------------|------------------|---------------|------------------|---------------|------------------|---------------|------------------|---------------|
| | mrem/yr | fract. |
| C-14 | 0.000E+00 | 0.0000 |
| Total | 0.000E+00 | 0.0000 |

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and
 As mrem/yr and Fraction of Total Dose At t = 1.000E+01 years

Water Dependent Pathways

| Radio- Nuclide | Water | | Fish | | Radon | | Plant | | Meat | |
|-------------------|------------------|---------------|------------------|---------------|------------------|---------------|------------------|---------------|------------------|---------------|
| | mrem/yr | fract. |
| C-14 | 5.403E-01 | 0.7433 | 1.866E-01 | 0.2567 | 0.000E+00 | 0.0000 | 4.074E-07 | 0.0000 | 1.868E-07 | 0.0000 |
| Total | 5.403E-01 | 0.7433 | 1.866E-01 | 0.2567 | 0.000E+00 | 0.0000 | 4.074E-07 | 0.0000 | 1.868E-07 | 0.0000 |

*Sum of all water independent and dependent pathways.

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and
 As mrem/yr and Fraction of Total Dose At t = 2.000E+01 years

Water Independent Pathways (Inhalation excludes radon)

| Radio- Nuclide | Ground | | Inhalation | | Radon | | Plant | | Meat | |
|-------------------|------------------|---------------|------------------|---------------|------------------|---------------|------------------|---------------|------------------|---------------|
| | mrem/yr | fract. |
| C-14 | 0.000E+00 | 0.0000 |
| Total | 0.000E+00 | 0.0000 |

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and
 As mrem/yr and Fraction of Total Dose At t = 2.000E+01 years

Water Dependent Pathways

| Radio- Nuclide | Water | | Fish | | Radon | | Plant | | Meat | |
|-------------------|------------------|---------------|------------------|---------------|------------------|---------------|------------------|---------------|------------------|---------------|
| | mrem/yr | fract. |
| C-14 | 2.238E-04 | 0.7433 | 7.728E-05 | 0.2567 | 0.000E+00 | 0.0000 | 1.688E-10 | 0.0000 | 7.736E-11 | 0.0000 |
| Total | 2.238E-04 | 0.7433 | 7.728E-05 | 0.2567 | 0.000E+00 | 0.0000 | 1.688E-10 | 0.0000 | 7.736E-11 | 0.0000 |

*Sum of all water independent and dependent pathways.

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and
 As mrem/yr and Fraction of Total Dose At t = 4.000E+01 years

Water Independent Pathways (Inhalation excludes radon)

| Radio- Nuclide | Ground | | Inhalation | | Radon | | Plant | | Meat | |
|-------------------|------------------|---------------|------------------|---------------|------------------|---------------|------------------|---------------|------------------|---------------|
| | mrem/yr | fract. |
| C-14 | 0.000E+00 | 0.0000 |
| Total | 0.000E+00 | 0.0000 |

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and
 As mrem/yr and Fraction of Total Dose At t = 4.000E+01 years

Water Dependent Pathways

| Radio- Nuclide | Water | | Fish | | Radon | | Plant | | Meat | |
|-------------------|------------------|---------------|------------------|---------------|------------------|---------------|------------------|---------------|------------------|---------------|
| | mrem/yr | fract. |
| C-14 | 3.839E-11 | 0.7433 | 1.326E-11 | 0.2567 | 0.000E+00 | 0.0000 | 2.895E-17 | 0.0000 | 1.327E-17 | 0.0000 |
| Total | 3.839E-11 | 0.7433 | 1.326E-11 | 0.2567 | 0.000E+00 | 0.0000 | 2.895E-17 | 0.0000 | 1.327E-17 | 0.0000 |

*Sum of all water independent and dependent pathways.

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and
 As mrem/yr and Fraction of Total Dose At t = 1.000E+02 years

Water Independent Pathways (Inhalation excludes radon)

| Radio- Nuclide | Ground | | Inhalation | | Radon | | Plant | | Meat | |
|-------------------|-----------|--------|------------|--------|-----------|--------|-----------|--------|-----------|--------|
| | mrem/yr | fract. | mrem/yr | fract. | mrem/yr | fract. | mrem/yr | fract. | mrem/yr | fract. |
| C-14 | 0.000E+00 | 0.0000 | 0.000E+00 | 0.0000 | 0.000E+00 | 0.0000 | 0.000E+00 | 0.0000 | 0.000E+00 | 0.0000 |
| Total | 0.000E+00 | 0.0000 | 0.000E+00 | 0.0000 | 0.000E+00 | 0.0000 | 0.000E+00 | 0.0000 | 0.000E+00 | 0.0000 |

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and
 As mrem/yr and Fraction of Total Dose At t = 1.000E+02 years

Water Dependent Pathways

| Radio- Nuclide | Water | | Fish | | Radon | | Plant | | Meat | |
|-------------------|-----------|--------|-----------|--------|-----------|--------|-----------|--------|-----------|--------|
| | mrem/yr | fract. |
| C-14 | 0.000E+00 | 0.0000 |
| Total | 0.000E+00 | 0.0000 |

*Sum of all water independent and dependent pathways.

Dose/Source Ratios Summed Over All Pathways
 Parent and Progeny Principal Radionuclide Contributions Indicated

| Parent (i) | Product (j) | Branch Fraction* | DSR(j,t) (mrem/yr)/(µCi/g) | | | | | |
|---------------|----------------|---------------------|----------------------------|-----------|-----------|-----------|-----------|-----------|
| | | | t= 0.000E+00 | 1.000E+00 | 1.000E+01 | 2.000E+01 | 4.000E+01 | 1.000E+02 |
| C-14 | C-14 | 1.000E+00 | 3.802E-23 | 2.818E+01 | 2.908E+00 | 1.204E-03 | 2.066E-10 | 1.043E-30 |

*Branch Fraction is the cumulative factor for the j't principal radionuclide daughter: CUMBRF(j)
 The DSR includes contributions from associated (half-life ≤ 0.5 yr) daughters.

Single Radionuclide Soil Guidelines G(i,t) in µCi/g
 Basic Radiation Dose Limit = 2.500E+01 mrem/yr

| Nuclide (i) | t= | 0.000E+00 | 1.000E+00 | 1.000E+01 | 2.000E+01 | 4.000E+01 | 1.000E+02 |
|----------------|------------|-----------|-----------|-----------|------------|------------|-----------|
| C-14 | *4.454E+06 | 8.870E-01 | 8.598E+00 | 2.076E+04 | *4.454E+06 | *4.454E+06 | |

*At specific activity limit

Summed Dose/Source Ratios DSR(i,t) in (mrem/yr)/(µCi/g)
 and Single Radionuclide Soil Guidelines G(i,t) in µCi/g
 at tmin = time of minimum single radionuclide soil guideline
 and at tmax = time of maximum total dose = 4.286 ± 0.009 years

| Nuclide (i) | Initial (µCi/g) | tmin (years) | DSR(i,tmin) | G(i,tmin) (µCi/g) | DSR(i,tmax) | G(i,tmax) (µCi/g) |
|----------------|--------------------|-----------------|-------------|----------------------|-------------|----------------------|
| C-14 | 2.500E-01 | 4.286 ± 0.009 | 1.599E+02 | 1.564E-01 | 1.599E+02 | 1.564E-01 |

Maximum Dose was at 4.286 years
 The material was disposed of
 on June 30, 1961 which is
 approximately 44.5 years ago,
 today Jan 12, 2006. RG broader
 1-12-06

Individual Nuclide Dose Summed Over All Pathways
 Parent Nuclide and Branch Fraction Indicated

| Nuclide (j) | Parent (i) | BRF(i) | DOSE(j,t), mrem/yr | | | | | |
|----------------|---------------|-----------|--------------------|-----------|-----------|-----------|-----------|-----------|
| | | | t= 0.000E+00 | 1.000E+00 | 1.000E+01 | 2.000E+01 | 4.000E+01 | 1.000E+02 |
| C-14 | C-14 | 1.000E+00 | 9.506E-24 | 7.046E+00 | 7.269E-01 | 3.011E-04 | 5.164E-11 | 0.000E+00 |

BRF(i) is the branch fraction of the parent nuclide.

Individual Nuclide Soil Concentration
 Parent Nuclide and Branch Fraction Indicated

| Nuclide (j) | Parent (i) | BRF(i) | S(j,t), µCi/g | | | | | |
|----------------|---------------|-----------|---------------|-----------|-----------|-----------|-----------|-----------|
| | | | t= 0.000E+00 | 1.000E+00 | 1.000E+01 | 2.000E+01 | 4.000E+01 | 1.000E+02 |
| C-14 | C-14 | 1.000E+00 | 2.500E-01 | 1.147E-01 | 1.033E-04 | 4.271E-08 | 7.295E-15 | 3.636E-35 |

BRF(i) is the branch fraction of the parent nuclide.

RESCALC.EXE execution time = 3.32 seconds