

Entergy Nuclear South Entergy Operations, Inc. 17265 River Road Killona, LA 70057-3093 Tel 504-739-6715 Fax 504-739-6698 rmurill@entergy.com

Robert J. Murillo Licensing Manager, Acting Waterford 3

W3F1-2005-0026

April 20, 2005

U.S. Nuclear Regulatory Commission Attn: Document Control Desk Washington, DC 20555-0001

Subject: Annual Radiological Environmental Operating Report - 2004 Waterford Steam Electric Station, Unit 3 (Waterford 3) Docket No. 50-382 License No. NPF-38

Dear Sir or Madam:

Attached is the Annual Radiological Environmental Operating Report for the period of January 1 through December 31, 2004. This report is submitted pursuant to the requirements of Waterford 3 Technical Specification Section 6.9.1.7.

If there are any questions please contact S.T. Fontenot at (504) 739-6656.

There are no new commitments contained in this submittal.

Sincerely,

Mun

R.J. Murillo Acting Licensing Manager

RJM/STF/stf

Attachment(s)

TE2S

W3F1-2005-0026 Page 2

cc: Mr. Bruce S. Mallett Regional Administrator U. S. Nuclear Regulatory Commission Region IV 611 Ryan Plaza Drive, Suite 400 Arlington, TX 76011-8064

> NRC Senior Resident Inspector Waterford Steam Electric Station Unit 3 P.O. Box 822 Killona, LA 70066-0751

U. S. Nuclear Regulatory Commission Attn: Mr. N. Kalyanam Mail Stop O-07D1 Washington, DC 20555-0001

Wise, Carter, Child & Caraway ATTN: J. Smith P.O. Box 651 Jackson, MS 39205

Winston & Strawn ATTN: N.S. Reynolds 1700 K Street, NW Washington, DC 20006-3817 Attachment 1 To

-

W3F1-2005-0026

Annual Radiological Environmental Operating Report - 2004



Annual Radiological Environmental Operating Report

January 1, 2004 - December 31, 2004



Waterford 3 Steam Electric Station Entergy Operations, Inc.

Docket Number 50-382 License Number NPF-38

Originator:

Una V. Dulpis

Ann V. Dubois, Chemistry Technician

03-24-05 Date

Reviewed By:

John M. Amato, Chemistry Supervisor (Acting)

David M. Rieder, Quality Assurance

<u>3/3//05</u> Date

3

Date

Approved By:

Hornsby, Chemistry Superintendent John L

4/12/05 Date

۰.

1

TABLE OF CONTENTS

SU	MM	ARY	1
1.0	IN	ITRODUCTION	5
1	1.1	Radiological Environmental Monitoring Program	5
1	.2	Pathways Monitored	5
1	.3	Land Use Census	5
2.0	IN	ITERPRETATION AND TRENDS OF RESULTS	20
2	2.1	Air Particulate and Radioiodine Sample Results	20
2	2.2	Thermoluminescent DosimetrySample Results	20
2	2.3	Water Sample Results	21
2	2.4	Sediment Sample Results	21
2	2.5	Milk Sample Results	22
2	2.6	Fish Sample Results	22
2	2.7	Broad Leaf Vegetation Sample Results	22
2	2.8	Land Use Census Results	22
2	2.9	Interlaboratory Comparison Results	22
3.0	R	ADIOLOGICAL ENVIRONMENTAL MONITORING PROGRAM SUMMARY	25
3	3.1	2004 Program Results Summary	25

LIST OF TABLES

TABLE 1.1	RADIOLOGICAL ENVIRONMENTAL SAMPLING PROGRAM	6
TABLE 2.1	BIANNUAL LAND USE CENSUS RESULTS	23
TABLE 3.1	RADIOLOGICAL ENVIRONMENTAL MONITORING PROGRAM SUMMARY	26

. ·

•

LIST OF FIGURES

FIGURE 1-1	REMP SAMPLES WITHIN 2 MILES OF WATERFORD 3	17
FIGURE 1-2	REMP SAMPLES WITHIN 10 MILES OF WATERFORD 3	18
FIGURE 1-3	REMP SAMPLES WITHIN 50 MILES OF WATERFORD 3	19
FIGURE 2-1	TLD RADIATION DOSE COMPARISON (BY YEAR)	24

.

• • • •

LIST OF ATTACHMENTS

ATTACHMENT 1	2004 RADIOLOGICAL MONITORING REPORT SUMMARY OF MONITORING RESULTS	30
ATTACHMENT 2	STATISTICAL COMPARISONS	47
ATTACHMENT 3	REVISED ODCM TABLES	55

Summary

The Annual Radiological Environmental Operating Report presents data obtained through analyses of environmental samples collected for Waterford 3's (W3) Radiological Environmental Monitoring Program (REMP) for the period January 1 through December 31, 2004. This report fulfills the requirements of W3 Technical Specification 6.9.1.7.

During 2004, gross beta radioactivity was detected in air and drinking/surface water locations. Results obtained at the indicator locations were similar to those obtained at the control location. Therefore, levels continue to remain at background.

Cesium-137, a man-made nuclide, was detected during 2004 at an indicator and control sediment location (SHWK-1 and SHWQ-6). This is consistent with results obtained from the preoperational program and previous years of operation. Studies in Louisiana indicate that Cesium-137 is commonly found in soils and sediments as a result of atmospheric weapons testing. Because the Cesium-137 levels are consistent with preoperational values, the Cesium-137 level detected in 2004 is more than likely attributable to weapons testing fallout.

Radiological Environmental Monitoring Program

W3 established the REMP prior to the station becoming operational (1985) to provide data on background radiation and radioactivity normally present in the area. W3 has continued to monitor the environment by sampling air, water, sediment, milk, fish and broad leaf vegetation, as well as measuring radiation directly.

The REMP includes sampling indicator and control locations within a 38-mile radius of the plant. The REMP utilizes indicator locations near the site to show any increases or buildup of radioactivity that might occur due to station operation, and control locations farther away from the site to indicate the presence of only naturally occurring radioactivity. W3 compares indicator results with control, preoperational, and previous years operational results to assess any impact W3 might have on the surrounding environment.

In 2004, W3 collected environmental samples for radiological analysis. Based on the comparison results of indicator locations with control locations and previous studies, it was concluded that overall W3 operations had no significant impact on plant environs. The review of 2004 data, in many cases, showed undetectable radiation levels in the environment and near background level in significant pathways associated with W3.

Harmful Effects or Irreversible Damage

The REMP did not detect any harmful effects or evidence of irreversible damage in 2004. Therefore, no analysis or planned course of action to alleviate problems was necessary.

Reporting Levels

W3's review indicates that no samples equaled or exceeded reporting levels for radioactivity concentration in environmental samples, as outlined in Technical Requirements Manual (TRM) Table 3.12-2 when averaged over any calendar quarter, due to W3 effluents. Therefore, 2004 results did not trigger any Radiological Monitoring Program Special Reports.

Radioactivity Not Attributable to W3

The W3 REMP detected radioactivity attributable to other sources twice. These include the 25th Chinese nuclear test explosion in 1980, and the radioactivity plume release due to reactor core degradation at the Chernobyl Nuclear Power Plant in 1986.

Comparison to State Program

W3 compared REMP data to the monitoring program of the Environmental Radiological Laboratory – Department of Environmental Quality Laboratory Services Division (ERL-DEQLSD). The ERL-DEQLSD and the W3 REMP entail similar radiological environmental monitoring program requirements. Both programs have obtained similar results over previous years.

Sample Deviations

• Milk Samples

Milk samples for the 1st, 3rd and 4th quarter of 2004 were unavailable from indicator location MKE-3 due to cows not producing enough milk. However, a sample was available during the 2nd quarter. One sample from control location MKR-40 was unavailable during the 4th quarter due to owner selling the cows. With the absence of milk samples at these locations, broad leaf vegetation sampling was performed as required by TRM Table 3.12-1.

• Air Samples

The air sample locations listed below failed to meet the requirement for sample continuity. As described in footnote (1) of TRM Table 3.12-1, deviations are permitted from the required sampling schedule due to malfunction of sampling equipment and other legitimate reasons.

Location	Sample period	Explanation of Deviation
APC-1	06/01/04 — 06/14/04	Sample pump tripped

Missed Samples

TLDs located at stations D-2, R-1, F-9 and E-15 were missing at the time of the third quarter TLD exchange.

• Required Lower Limit of Detection (LLD) Values

All LLDs during this reporting period were within the acceptable limits required by the W3 TRM.

Unavailable Results

W3 received analytical results in adequate time for inclusion in this report. In addition, W3's review identified no missing results.

• Program Modifications

The following changes were made to the ODCM during 2004:

- The description for TLD location R-1 was revised.
- Milk location MKQ-5 was deleted from the program.
- Milk location MKE-3 was added to the program.

<u>Attachments</u>

Attachment 1 contains results of air, TLD, water, sediment, milk, fish and broad leaf vegetation collected in 2004. TLDs were analyzed by Waterford-3 Dosimetry. All remaining samples were analyzed by the River Bend (RBS) Environmental Laboratory. Attachment 1 also contains River Bend's participation in the interlaboratory comparison program during 2004.

Attachment 2 contains statistical comparisons of:

- TLD measurements from stations grouped by distance
- TLD radiation dose to historical data by location
- Gross beta activity measurements on air particulate filters
- Gross beta activity measurements in surface/drinking water samples

Attachment 3 contains revised ODCM tables.

1.0 Introduction

1.1 Radiological Environmental Monitoring Program

W3 established the REMP to ensure that plant operating controls properly function to minimize any associated radiation endangerment to human health or the environment. The REMP is designed for:

- Analyzing important pathways for anticipated types and quantities of radionuclides released into the environment.
- Considering the possibility of a buildup of long-lived radionuclides in the environment and identifying physical and biological accumulations that may contribute to human exposures.
- Considering the potential radiation exposure to plant and animal life in the environment surrounding W3.
- Correlating levels of radiation and radioactivity in the environment with radioactive releases from station operation.

1.2 Pathways Monitored

The airborne, direct radiation, waterborne and ingestion pathways are monitored as required by W3 TRM Table 3.12-1. A description of the W3 REMP utilized to monitor the exposure pathways is described in Table 1.1 and shown in Figures 1-1, 1-2 and 1-3.

Section 2.0 of this report provides a discussion of 2004 sampling results with Section 3.0 providing a summary of results for the monitored exposure pathways.

1.3 Land Use Census

W3 conducts a land use census biennially, as required by Section 3.12.2 of the TRM. The purpose of this census is to identify changes in uses of land within five miles of W3 that would require modifications to the REMP and the Offsite Dose Calculation Manual (ODCM). The most important criteria during this census are to determine the location in each sector of the nearest:

- 1) Residence
- 2) Animal milked for human consumption
- 3) Garden of greater than 50 m² (500 ft²) producing broad leaf vegetation.

W3 conducts the land use census by:

- Field surveys in each meteorological sector out to five miles in order to confirm:
 - > Nearest permanent residence
 - > Nearest garden and approximate size
 - > Nearest beef cow
 - Nearest food product
 - Nearest milking animal
- Identifying locations on maps, measuring distances to W3 and recording results on data sheets.
- Comparing current census results to previous results.

- -

Table 1.1

Exposure Pathway	Requirement	Sample Point Description, Distance and Direction	Sampling and Collection Frequency	Type and Frequency Of Analyses
Airborne	Radiolodine and Particulates Three samples from close to the three SITE BOUNDARY locations, in different sectors, in or near sectors having the highest calculated annual average ground level D/Q.	 APQ-1 (NW, 0.81 Miles) – (West bank) Located in soybean/sugarcane field off LA 18 approximately 0.6 miles east of LA 18/3141 intersection. APF-1 (ESE, 0.35 Miles) – (West bank) Located on north side of Secondary Meteorological Tower. APC-1 (NE, 0.67 Miles) – (East bank) Located inside the Little Gypsy Cooling Water Intake Structure fence enclosure. 	Continuous sampler operation with sample collection bi- weekly, or more frequently if required by dust loading.	Radioiodine Canister – I-131 analysis bi-weekly. Particulate Sampler – Gross beta radioactivity analysis following filter change. Gamma isotopic analysis of composite (by location) quarterly.
	Radioiodine and Particulates One sample from the vicinity of a community having the highest calculated annual average ground level D/Q.	APP-1 (WNW, 0.84 Miles) – (West bank) Located in soybean/sugarcane field at northwest corner of Short St. in Killona.		
	Radioiodine and Particulates One sample from a control location, as for example 15 -30 km distant and in the least prevalent wind direction.	APE-30 (E, 25.2 Miles) – (West bank) Located on the roof of the Entergy Office building on Delaronde St. in Algiers. (Control)		

Table 1.1

n Frequency Of Analyses
arterly Gamma dose quarterly.

Table 1.1

Exposure	Requirement	Sample Point Description,	Sampling and	Type and Frequency
Pathway		Distance and Direction	Collection Frequency	Of Analyses
Direct Radiation	TLDs An inner ring of stations, one in each meteorological sector in the general area of the SITE BOUNDARY.	 E-1 (E, 0.41 Miles) – (West bank) Located on utility pole along LA 18 approximately 0.3 miles east of Waterford 3 plant entrance. F-2 (ESE, 1.15 Miles) – (West bank) Located on southeast corner of fence enclosure surrounding the Entergy sub station 0.2 miles south of LA 18 on LA 3142. G-2 (SE, 1.26 Miles) – (West bank) Located on fence east of LA 3142 approximately 0.3 miles north of railroad overpass. H-2 (SSE, 1.54 Miles) – (West bank) Located off LA 3142 on southwest edge of fence along shell road 0.4 miles north of LA 3127/3142 intersection. 	Quarterly	Gamma dose quarterly.

Table 1.1

Exposure Pathway	Requirement	Sample Point Description, Distance and Direction	Sampling and Collection Frequency	Type and Frequency Of Analyses
Direct Radiation	TLDs An inner ring of stations, one in each meteorological sector in the general area of the SITE BOUNDARY.	J-2 (S, 1.38 Miles) – (West bank) Located on fence enclosure for valve station south of LA 3127 approximately 0.6 miles west of LA 3127/3142 intersection.	Quarterly	Gamma dose quarterly.
		K-1 (SSW, 1.06 Miles) – (West bank) Located on stop sign at entrance to Entergy Education Center on LA 3127.		
		L-1 (SW, 1.06 Miles) – (West bank) Located on gated entrance off of LA 3127 approximately 1.6 miles west of LA 3127/3142 intersection.		
		M-1 (WSW, 0.76 Miles) – (West bank) Located on south gate of Waterford 1 and 2 fuel oil storage tank enclosure.		
		N-1 (W, 0.98 Miles) – (West bank) Located on pole at corner of Railroad Avenue and School House Road.		

Table 1.1

Radiological Environmental Sampling Program

Exposure Pathway	Requirement	Sample Point Description, Distance and Direction	Sampling and Collection Frequency	Type and Frequency Of Analyses
Direct Radiation	TLDs An inner ring of stations, one in each meteorological sector in the general area of the SITE BOUNDARY.	P-1 (WNW, 0.84 Miles) – (West bank) Located on fence enclosing air sample station APP-1.	Quarterly	Gamma dose quarterly.
		Q-1 (NW, 0.81 Miles) – ((West bank) Located on fence enclosing air sample station APQ-1.		
		R-1 (NNW, 0.51 Miles) – (West bank) Located at Waterford 1 and 2 Cooling Water Intake Structure on the catwalk.		
	TLDs An outer ring of stations, 1 in 10 of the meteorological sectors in the 6- to 8- km range from the site.	A-5 (N, 4.59 Miles) – (East bank) Located on utility pole at intersection of Oswald Avenue and US 61.		

.

.

Table 1.1

Exposure	Requirement	Sample Point Description,	Sampling and	Type and Frequency
Pathway		Distance and Direction	Collection Frequency	Of Analyses
Direct Radiation	TLDs An outer ring of stations, 1 in 10 of the meteorological sectors in the 6- to 8- km range from the site.	 B-4 (NNE, 3.75 Miles) – (East bank) Located on utility pole guidewire next to transmission tower south of weigh station on US 61 at St. John/St. Charles Parish line. D-5 (ENE, 4.09 Miles) – (East bank) Located on gate on shell road approximately 0.1 miles north of US61/LA48 intersection. F-4 (ESE, 3.53 Miles) – (West bank) Located on utility pole behind house at 646 Aquarius St. in Hahnville. 	Quarterly	Gamma dose quarterly.

Table 1.1

Exposure Pathway	Requirement	Sample Point Description, Distance and Direction	Sampling and Collection Frequency	Type and Frequency Of Analyses
Direct Radiation	TLDs An outer ring of stations, 1 in 10 of the meteorological sectors in the 6- to 8- km range from the site.	 E-5 (E, 4.08 Miles) – (East bank) Located on the Norco Substation fence enclosure at the end of Wesco Street off of LA 48. G-4 (SE, 3.30 Miles) – (West bank) Located on railroad sign on LA 3160 approximately 0.1 miles north of railroad track. H-8 (SSE, 8.13 Miles) – (West bank) Located on a road sign on south side of HWY 90 directly in front of Hahnville High School 	Quarterly	Gamma dose quarterly.
	-	 approximately 0.1 miles east of Tiger Drive. P-6 (WNW, 5.58 Miles) – (West bank) Located on a fence surrounding the communications tower at the LA 640/railroad track intersection. Q-5 (NW, 5.01 Miles) – (West bank) Located on utility pole along LA 18 across from Mississippi River marker 137. 		

Table 1.1

Exposure Pathway	Requirement	Sample Point Description, Distance and Direction	Sampling and Collection Frequency	Type and Frequency Of Analyses
Direct Radiation	TLDs An outer ring of stations, 1 in 10 of the meteorological sectors in the 6- to 8- km range from the site.	R-6 (NNW, 5.52 Miles) (East bank) Located on fence enclosure approximately 0.2 miles west of US 61 on LA 3223 near railroad crossing.	Quarterly	Gamma dose quarterly.
	TLDs The balance of the stations to be in special interest areas such as population centers, nearby residences, schools, and in 1 or 2 areas to serve as control stations.	F-9 (ESE, 8.18 Miles) – (East bank) Located on entrance gate to Destrehan Substation just north of railroad tracks on Jonathan Street approximately 1.5 miles east of Luling-Destrehan Bridge, off of LA 48.		
		G-8 (SE, 7.74 Miles) – (West bank) Located on southern most corner of the back fence of Entergy Office in Luling.		
		E-15 (E, 11.7 Miles) – (East bank) Located on Kenner Substation fence enclosure on Alliance Ave approximately 0.1 miles from LA 48.		•

Table 1.1

Exposure Pathway	Requirement	Sample Point Description, Distance and Direction	Sampling and Collection Frequency	Type and Frequency Of Analyses
Direct Radiation	<u>TLDs</u> The balance of the stations to be in special interest areas such as population centers, nearby residences, schools, and in 1 or 2 areas to serve as control stations.	J-15 (S, 11.7 Miles) - (West bank) Located on utility pole near the LA 631/Hwy 90 intersection in Des Allemands. E-30 (E, 25.2 Miles) - (West bank) Located at entrance to the Entergy Office on Delaronde St. in Algiers. (Control)	Quarterly	Gamma dose quarterly.
Waterborne .	Surface Water One sample upstream	SWP-7 (WNW, 7.37 Miles) - (West bank) Located at St. John Parish Waterworks off LA 18 in Edgard. (Control)	Composite sample over one quarter period.	Gamma isotopic analysis quarterly. Composite for tritium analysis quarterly.
	One sample downstream	 SWF-2 (ESE, 1.51 Miles) - (West bank) Located at Dow Chemical Plant drinking water canal. SWE-5 (E, 4.59 Miles) - (East bank) Located at St. Charles Parish Waterworks off LA 48 in New Sarpy. SWK-1 (SSW, 0.49 Miles) - (West bank) Located at 40 Arpent Canal south of the plant. The canal is northwest of the shell access road/railroad track intersection. 		

Exposure	Requirement	Sample Point Description,	Sampling and	Type and Frequency		
Pathway		Distance and Direction	Collection Frequency	Of Analyses		
Waterborne	Drinking Water One sample upstream One sample downstream	 DWP-7 (WNW, 7.37 Miles) - (West bank) Located at St. John Parish Waterworks off LA 18 in Edgard. (Control) DWF-2 (ESE, 1.51 Miles) - (West bank) Located at Dow Chemical Plant drinking water canal. DWE-5 (E, 4.59 Miles) - (East bank) Located at St. Charles Parish Waterworks off of LA 48 in New Sarpy. 	Composite sample over one month period when I-131 analysis is performed, quarterly composite otherwise.	I-131 analysis on each composite when the dose calculated for the consumption of the water is greater than one mrem per year. Composite for gross beta and gamma isotopic analyses quarterly. Composite for tritium analysis quarterly.		
	Sediment from Shoreline One sample upstream	SHWQ-6 (NW, 5.99 Miles) – (East bank) Located of LA 628 approximately 0.1 miles east of Reserve ferry landing. (Control)	Annualiy	Gamma isotopic analysis annually.		
	One sample downstream	 SHWE-3 (E, 2.99 Miles) – (West bank) Located at Foot Ferry landing off LA 18 in Taft. SHWK-1 (SSW, 0.49 Miles) – (West bank) Located at 40 Arpent Canal south of plant. The canal is northwest of the shell access road/railroad track intersection. 				
Ingestion	Milk Samples from milking animals in the three locations within 5 km distance having the highest dose potential. If there are none, then, one sample from milking animals in each of the three areas between 5 to 8 km distant where doses are calculated to be greater than 1 mrem per year.	MKE-3 (E, 2.4 Miles) - (West bank) Located at the Zeringue's house on LA 18 in Taft.	Quarterly	Gamma isotopic and I-131 analysis quarterly.		

	Table 1.1	
Radiological	Environmental Sam	pling Program

Table 1.1

Exposure Pathway	Requirement	Sample Point Description, Distance and Direction	Sampling and Collection Frequency	Type and Frequency Of Analyses
Ingestion	Milk One sample from milking animals at a control location 15 – 30 km distant and in the least prevalent wind direction.	MKR-40 (NNW, 40.70 Miles) – (East bank) Located at 24254 LA Highway 442, Holden. La. (Control)	Quarterly	Gamma isotopic and I-131 analysis quarterly.
	Fish and Invertebrates One sample of each commercially and recreational important species in vicinity of plant discharge area.	- FH-2 (Distance/Direction Not Applicable) – Downstream of the plant intake structure. FH-3 (Distance/Direction Not Applicable)– (Westbank) Waterways downstream of plant discharge directed to 40 Arpent Canal.	Sample in season, or annually if they are not seasonal	Gamma isotopic analysis on edible portion.
	One sample of same species in area not influenced by plant discharge.	FH-1 (Distance/Direction Not Applicable) – Upstream of the plant discharge structure. (Control)		
	Broadleaf Samples of one to three different kinds of broadleaf vegetation grown nearest each of two different off-site locations of highest predicted annual average groundlevel D/Q if milk sampling is not performed. One sample of each of the similar broadleaf vegetation grown 15 – 30 km distant in the least prevalent wind direction if milk sampling is not performed.	 BLQ-1 (NW, 0.83 Miles) – (West bank) Located near air sample station APQ-1. BLB-1 (NNE, 0.81 Miles) – (East bank) Located near transmission tower west of Little Gypsy on LA 628. BLE-20 (E, 19.7 Miles) – (West bank) Located on property of Nine Mile Point in Westwego, LA. (Control) 	Quarterly	Gamma isotopic and I-131 analysis.



FIGURE 1-2





2.0 Interpretation and Trends of Results

2.1 Air Particulate and Radioiodine Sample Results

Samples of airborne particulate and radioiodine were collected at four indicator locations and one control location and analyzed for gross beta radionuclides, lodine-131 and gamma radionuclides (quarterly air particulate filter composites only). W3 did not detect any gamma radionuclides in the quarterly air particulate composites or lodine-131 in the radioiodine cartridges during the reporting period, as has been the case in previous years. Indicator gross beta air particulate results for 2004 were similar to those background levels obtained in previous years of the operational REMP and well below preoperational levels as seen below. Results are reported as annual average pCi/m³.

Monitoring Period	<u>Result</u>
Preoperational	0.080
1983 – 2003	0.019
2004	0.019

Table 3.1, which includes gross beta concentrations for 2004, provides a comparison of the indicator and control means further emphasizes that the airborne pathway continues to remain at background levels. In addition, as shown in Attachment 2, the standard "t" test was used to compare average gross beta activity from each indicator station to the average gross beta activity at the control station. The results from this test show the average activity detected at all indicator stations is statistically the same as the average activity detected at the control station. Therefore, W3 concluded that plant operations had no significant impact on "this pathway during 2004.

2.2 Thermoluminescent DosimetrySample Results

The average exposure rates during 2004 are consistent with those from the preoperational program and the previous five years of operation as seen in Figure 2-1. In particular, the preoperational survey indicates that exposure rates ranged between 11 and 33 mrem/standard quarter with an average of 20 mrem/standard quarter. The range during the previous five years of operation was 9 to 15 mrem/standard quarter with an average exposure rate of 12 mrem/standard quarter.

A comparison of the indicator results to the control results, as seen in Table 3.1, shows that the average indicator is slightly higher than that of the control. As shown in Attachment 1, Table 2.1, several indicator locations are higher than the control by a few mrem with a maximum difference of three mrem at one location (F-4).

As shown in Attachment 2, Table 2.1, the standard "t" test was used to compare average exposure rates for TLD stations located in groups 0-2 miles and 2-5 miles from the plant to those > 5 miles. The results indicate that the average exposure rates 0-2 miles and 2-5 miles from the plant are statistically the same as those > 5 miles.

The differences between indicator locations and the control, and TLD stations grouped by distance from the plant are expected due to a variety of factors not related to W3 plant operations that can affect background radiation in the vicinity of each TLD station. Direct radiation measurements at each TLD station have remained statistically the same in 2004 as previous years of operation as evidenced on Attachment 2, Table 2.2. In addition, Radiological Gaseous Effluents for 2004 were only a small fraction of the limits as is typical in previous years of operation and are not expected to have any impact on environmental TLD measurements.

2.3 Water Sample Results

Analytical results for 2004 drinking/surface water samples were similar to those reported in previous years.

Drinking/Surface Water

Drinking water samples also serve as surface water samples for W3. Therefore, monthly and quarterly gamma spectroscopy and tritium analyses of drinking water also satisfy the surface water sampling requirement.

Composite drinking/surface water samples were collected from two indicator and one control location and analyzed for lodine-131, gamma radionuclides and tritium. Results indicate that all measurements were below the calculated LLDs.

Although gross beta was detected in the drinking/surface water samples, results for the indicator locations were below previous operational and preoperational years as seen below. Results are reported as annual average pCi/l.

Monitoring Period	<u>Result</u>
Preoperational	7.0
1983 – 2003	5.1
2004	2.8

Table 3.1, which includes gross beta concentrations for 2004, provides a comparison of the indicator and control means shows that the waterborne pathway continues to remain at background levels. In addition, as shown in Attachment 2, the standard "t" test was used to compare average gross beta activity from each indicator station to the average gross beta activity from the control station. The results from this test show average activity detected at all indicator stations is statistically the same as the average activity detected at the control station. Therefore, W3 concluded that plant operations had no significant impact on this pathway during 2004.

Surface Water

Surface water samples were collected from one indicator location and analyzed for gamma radionuclides and tritium. Results indicate that all measurements were below the calculated LLDs.

4

2.4 Sediment Sample Results

Sediment samples were collected from two indicator locations and one control location and analyzed for gamma radionuclides. Cesium-137, a man-made nuclide, was detected in sample SHWK-1 with a concentration of 32 pCi/kg and sample SHWQ-6 with a concentration of 83 pCi/kg. No other man-made radionuclides were detected in any of the samples.

The Cesium-137 results obtained during 2004 are consistent with those from the preoperational program and previous years of operation. In particular, the preoperational survey indicates that Cesium-137 was detected in 9 of 14 soil samples at concentrations ranging between 30 and 890 pCi/kg with an average concentration of 164 pCi/kg. Similarly, the range indicated during the previous years of operation was 18 to 142 pCi/kg with an average activity of 57 pCi/kg.

W3 has detected Cs-137 in wastewater tanks discharged to the Mississippi River at concentrations typically below 1E-6 uCi/ml in the past few years. After dilution by Circulating Water, this concentration is reduced to well below 1E-10 uCi/ml prior to mixing with the Mississippi River where it is diluted even further. At the minimal concentrations being discharged from W3 as compared to the typical Cs-137 concentrations commonly found in

soils and sediments in Louisiana as a result of atmospheric fallout from nuclear weapons testing as noted in the preoperational study, plant operations is not expected to result in any appreciable quantities of radioactivity in sediment collected from the bank of the Mississippi River. In addition, the radioactivity detected in sediment this year is consistent with preoperational data even after applying a correction for natural decay of Cs-137.

Therefore, W3 concluded that plant operations had no significant impact on this pathway during 2004.

2.5 Milk Sample Results

Milk samples were collected from one indicator and one control location and analyzed for lodine-131 and gamma radionuclides. Results indicate that all measurements were below the calculated LLDs. Therefore, W3 concluded that plant operations had no significant impact on this pathway during 2004.

2.6 Fish Sample Results

Fish samples were collected from two indicators and one control location and analyzed for gamma radionuclides. Results indicate that all measurements were below the calculated LLDs. Therefore, W3 concluded that plant operations had no significant impact on this pathway during 2004.

2.7 Broadleaf Vegetation Sample Results

Broadleaf vegetation samples were collected from two indicators and one control location and analyzed for lodine-131 and gamma radionuclides. Results indicate that all measurements were below the calculated LLDs. Therefore, W3 concluded that plant operations had no significant impact on this pathway during 2004.

2.8 Land Use Census Results

In compliance with the Waterford 3 ODCM and TRM, the land use census was conducted on September 27, 28 and 29, 2004. The nearest residence, garden, beef cow, food product and milk animal in each sector within a five mile radius of the plant was located by visual inspection and verbal inquiry.

While residence, milk cow and food product locations remained unchanged for 2004, one location of goats (sector E) and two locations of beef cows (sector G) were removed. Two new garden locations (sector C and D), and three new goat locations (sector E, F and Q) were identified in 2004. Based upon the locations identified in this survey, the locations identified in previous surveys and the locations currently being used to calculate dose commitments from liquid and gaseous effluents released from W3, no REMP sampling location changes are necessary. Results of the 2004 biennial census are shown in Table 2.1.

2.9 Interlaboratory Comparison Results

The River Bend Station Environmental Laboratory analyzed interlaboratory comparison samples for W3 to fulfill the requirements of Section 5.7.2 of the ODCM. Attachment 1 contains these results.

Sector	Direction	Distance from Plant in Miles					
		Residence	Garden	Milk Cows	Beef Cows	Goats	Food Products
A	N	1.3	1.7	* 4.6	4.6	^	4.1
В	NNE	1.1	1.3	۸	۸	۸	1.3
С	NE	0.9	1.0	٨	۸	۸	^
D	ENE	0.9	0.9	۸	۸	۸	^
E	E	2.2	2.2	**2.3	2.3	* 3.2	0.3
F	ESE	3.1	2.2	۸	2.3	* 3.5	0.3
G	SE	4.0	4.1	^	2.4	^	0.3
н	SSE	۸	۸	۸	۸	۸	0.3
J	S	۸	۸	۸	۸	٨	0.5
к	SSW	۸	۸	٨	۸	۸	0.5
L	sw	۸	^	۸	۸	^	0.5
М	wsw	۸	1.4	۸	1.2	۸	0.5
N	W	1.0	1.1	۸	1.0	٨	0.6
Р	WNW	0.9	0.9	۸	0.9	۸	0.6
Q	NW	0.9	1.0	٨	0.9	* 4.9	0.6
R	NNW	3.0	3.0	۸	4.9	۸	2.6

TABLE 2.1Biennial Land Use Census Results

^ Indicates that nothing was found in the Sector within a five mile radius of Waterford 3

* Animals were located at this distance from Waterford 3, but the milk is not currently used for human consumption

** Samples are being obtained from animals at this location (MKE-3) for REMP

FIGURE 2-1 TLD RADIATION DOSE COMPARISON (BY YEAR)

3.0 Radiological Environmental Monitoring Program Summary

3.1 2004 Program Results Summary

Table 3.1 summarizes the 2004 REMP results. W3 did not use values reported as less than the lower limit of detection (< LLD) when determining ranges and means for indicator and control locations.

Radiological Environmental Monitoring Program Summary

Name of Facility: <u>Waterford 3 SES</u> Docket No: <u>50-382</u> Location of Facility: <u>St. Charles, Louisiana</u> Reporting Period: <u>January - December 2004</u>

Sample Type (Units)	Type & Number Of Analyses ^a	LLD b	Indicator Locations Mean(F) ^C [Range]	Location with High	nest Annual Mean	Control Locations Mean(F) ^C [Range]	Number of Nonroutine Results ^e
				Location d	Mean(F) ^C [Range]		
Airborne Particulates (pCi/m ³)	GB 130	0.01	0.019(104/104) [0.009 - 0.034]	APP-1 (WNW, 0.84 mi.)	0.019 (26 / 26) [0.009 - 0.032]	0.019 (26 / 26) [0.006 - 0.036]	0
	GS 20 Cs-134 Cs-137	0.05 0.06	<lld <lld< td=""><td>N/A N/A</td><td>N/A N/A</td><td><lld <lld< td=""><td>0 0</td></lld<></lld </td></lld<></lld 	N/A N/A	N/A N/A	<lld <lld< td=""><td>0 0</td></lld<></lld 	0 0
Airborne lodine (pCl/m ³)	I-131 130	0.07	<lld< td=""><td>N/A</td><td>N/A</td><td><lld< td=""><td>0</td></lld<></td></lld<>	N/A	N/A	<lld< td=""><td>0</td></lld<>	0
Indicator TLDs (mrem/Std. Qtr)	Gamma 120	(f)	12(116/120) [8.0-14.0]	F-4 (ESE, 3.53 mi.)	14 (4 / 4) [13.7 – 14.9]	N/A	0
Control TLDs (mrem/Std. Qtr)	Gamma 4	(f)	N/A	N/A	N/A	11 (4 / 4) [10.0 – 12.2]	0

Radiological Environmental Monitoring Program Summary

Name of Facility: <u>Waterford 3 SES</u> Docket No: <u>50-382</u> Location of Facility: <u>St. Charles, Louisiana</u> Reporting Period: <u>January - December 2004</u>

Sample Type (Units)	Type & Number LLD b of Analyses ^a		Type & Number of Analyses ^a [Range]		Indicator Location Mean(F) ^C [Range]	Location with Highest Annual Mean		Control Locations Mean (F) ^C [Range]	Number of Nonroutine Results ^e
					Location ^d	Mean(F) ^C [Range]			
Surface Water & Drinking Water (pCi/l)	Gross Beta	12	4	3.1 (7/8) [1.2 – 5.2]	DWF-2 (ESE, 1.51 mi.)	3.9 (4/4) [1.2 – 5.2]	2.1 (3 / 4) [1.7 – 2.4]	0	
	I-131	42	1	<lld< td=""><td>N/A</td><td>N/A</td><td><lld< td=""><td>0</td></lld<></td></lld<>	N/A	N/A	<lld< td=""><td>0</td></lld<>	0	
	Н-3	12	2000	<lld< td=""><td>N/A</td><td>N/A</td><td><lld< td=""><td>0</td></lld<></td></lld<>	N/A	N/A	<lld< td=""><td>0</td></lld<>	0	
	GS Mn-5 Fe-59 Co-56 Zo-66 Zn-69 Zr-95 Nb-9 Cs-13 Cs-13 Ba-14 La-14	12 54 9 8 0 5 5 5 5 34 37 40	15 30 15 15 15 15 15 15 15	<lld <lld <lld <lld <lld <lld <lld <lld< td=""><td>N/A N/A N/A N/A N/A N/A N/A N/A N/A N/A</td><td>N/A N/A N/A N/A N/A N/A N/A N/A N/A N/A</td><td><lld <lld <lld <lld <lld <lld <lld <lld< td=""><td>0 0 0 0 0 0 0 0 0 0 0 0 0</td></lld<></lld </lld </lld </lld </lld </lld </lld </td></lld<></lld </lld </lld </lld </lld </lld </lld 	N/A N/A N/A N/A N/A N/A N/A N/A N/A N/A	N/A N/A N/A N/A N/A N/A N/A N/A N/A N/A	<lld <lld <lld <lld <lld <lld <lld <lld< td=""><td>0 0 0 0 0 0 0 0 0 0 0 0 0</td></lld<></lld </lld </lld </lld </lld </lld </lld 	0 0 0 0 0 0 0 0 0 0 0 0 0	

Radiological Environmental Monitoring Program Summary

Name of Facility: <u>Waterford 3 SES</u> Docket No: <u>50-382</u> Location of Facility: <u>St. Charles, Louisiana</u> Reporting Period: <u>January - December 2004</u>

Sample Type (Units)	Type Numb of Analys	& er ses ^a	LLD b	Indicator Locations Mean(F) ^C [Range]	Location with Highest Annual Mean		Control Locations Mean(F) ^C [Range]	Number of Nonroutine Results ^e
					Location d	Mean(F) ^C [Range]		
Surface Water (pCi/l)	H-3	4	3000	<lld< td=""><td>N/A</td><td>NA</td><td>N/A</td><td>0</td></lld<>	N/A	NA	N/A	0
	GS	15						
	Mn-5 Fe-5 Co-5 Co-6 Zn-6 Zr-9 Nb-9 Cs-13 Cs-13 Ba-14 La-14	4 9 8 0 5 5 5 5 34 37 40	15 30 15 15 30 15 15 15 15 15 15	<lld <lld <lld <lld <lld <lld <lld <lld< td=""><td>N/A N/A N/A N/A N/A N/A N/A N/A N/A</td><td>N/A N/A N/A N/A N/A N/A N/A N/A N/A</td><td>N/A N/A N/A N/A N/A N/A N/A N/A N/A N/A</td><td>0 0 0 0 0 0 0 0 0 0 0 0</td></lld<></lld </lld </lld </lld </lld </lld </lld 	N/A N/A N/A N/A N/A N/A N/A N/A N/A	N/A N/A N/A N/A N/A N/A N/A N/A N/A	N/A N/A N/A N/A N/A N/A N/A N/A N/A N/A	0 0 0 0 0 0 0 0 0 0 0 0
Shoreline Sediment (pCi/kg dry)	GS Cs-13 Cs-13	3 34 37	150 180	<lld 32.2 (1 / 2) N/A</lld 	N/A SHWK-1 (SSW, 0.49 mi.)	N/A 32.2 (1 / 2) N/A	<lld 82.5 (1 / 1) N/A</lld 	0 0

•

Radiological Environmental Monitoring Program Summary

Name of Facility: Waterford	<u>3 SES</u> Docket No: <u>50-382</u>
Location of Facility: <u>St. Charles, Louisiana</u>	Reporting Period: January - December 2004

Sample Type (Units)	Type & Number of Analyses ^a	LLD b	Indicator Location Mean (F) ^C [Range]	Location with High	hest Annual Mean	Control Locations Mean(F) ^C [Range]	Number of Nonroutine Results ^e
				Location d	Mean(F) ^C [Range]		
Milk (pCi/l)	I-131 4	1	<lld< td=""><td>N/A</td><td>N/A</td><td><lld< td=""><td>0</td></lld<></td></lld<>	N/A	N/A	<lld< td=""><td>0</td></lld<>	0
	Cs-134 Cs-137 Ba-140 La-140	15 18 15 15	<lld <lld <lld <lld< td=""><td>N/A N/A N/A N/A</td><td>N/A N/A N/A N/A</td><td><lld <lld <lld <lld< td=""><td>0 0 0 0</td></lld<></lld </lld </lld </td></lld<></lld </lld </lld 	N/A N/A N/A N/A	N/A N/A N/A N/A	<lld <lld <lld <lld< td=""><td>0 0 0 0</td></lld<></lld </lld </lld 	0 0 0 0
Fish (pCi/kg wet)	GS 12 Mn-54 Fe-59 Co-58 Co-60 Zn-65 Cs-134 Cs-137	130 260 130 130 260 130 150	<lld <lld <lld <lld <lld <lld <lld< td=""><td>N/A N/A N/A N/A N/A N/A N/A</td><td>N/A N/A N/A N/A N/A N/A N/A</td><td><lld <lld <lld <lld <lld <lld <lld< td=""><td>0 0 0 0 0 0</td></lld<></lld </lld </lld </lld </lld </lld </td></lld<></lld </lld </lld </lld </lld </lld 	N/A N/A N/A N/A N/A N/A N/A	N/A N/A N/A N/A N/A N/A N/A	<lld <lld <lld <lld <lld <lld <lld< td=""><td>0 0 0 0 0 0</td></lld<></lld </lld </lld </lld </lld </lld 	0 0 0 0 0 0
Broadleaf Vegetation (pCi/kg wet)	I-131 12 GS 12 Cs-134	60 60	<lld< td=""><td>N/A N/A</td><td>N/A</td><td><lld< td=""><td>0</td></lld<></td></lld<>	N/A N/A	N/A	<lld< td=""><td>0</td></lld<>	0
	Cs-137	80	<lld< td=""><td>N/A</td><td>N/A</td><td><lld< td=""><td><u> </u></td></lld<></td></lld<>	N/A	N/A	<lld< td=""><td><u> </u></td></lld<>	<u> </u>

^a GB = Gross beta; I-131 = Iodine-131; H-3 = Tritium; GS = Gamma scan.

b LLD = required lower limit of detection based on Waterford 3 TRM.

^c Mean and range based upon detectable measurements only. Fraction of detectable measurements at specified locations is indicated in parenthesis (F).

d Locations are specified (1) by name and (2) degrees relative to reactor site.

• Non-routine results are those which exceed ten times the control station value. If no control station value is available, the result is considered non-routine if it exceeds ten times the preoperational value for the location.

f LLD is not defined in Waterford 3 TRM.

TABLE OF CONTENTS

TABLE 1.1	AIR PARTICULATE FILTER	32
TABLE 1.2	RADIOIODINE CARTRIDGE	33
TABLE 1.3	AIR PARTICULATE FILTER	34
TABLE 2.1	THERMOLUMINESCENT DOSIMETERS	35
TABLE 3.1	DRINKING/SURFACE WATER	36
TABLE 3.2	DRINKING/SURFACE WATER	37
TABLE 3.3	DRINKING/SURFACE WATER	38
TABLE 3.4	DRINKING/SURFACE WATER	39
TABLE 3.5	SURFACE WATER	40
TABLE 4.1	SEDIMENT	41
TABLE 5.1	MILK	42
TABLE 6.1	FISH	43
TABLE 7.1	BROADLEAF VEGETATION	44
TABLE 8.1	INTERLABORATORY COMPARISON	45

.

Annual Radiological Environmental Operating Report

• • •

Table 1.1 Sample Type: <u>Air Particulate Filter</u> Analysis: Gross Beta Units: pCi/m³

End Date	APF-1	APQ-1	APP-1	APC-1	APE-30
	(Indicator)	(Indicator)	(Indicator)	(Indicator)	(Control)
Required LLD	<u>0.01</u>	<u>0.01</u>	<u>0.01</u>	<u>0.01</u>	<u>0.01</u>
01-12-04 01-26-04 02-09-04 02-25-04 03-08-04 03-22-04 04-05-04 04-20-04 05-03-04 05-17-04 06-01-04 06-14-04 06-29-04 07-12-04 07-26-04 08-09-04 08-23-04 09-07-04 09-20-04 10-04-04 10-18-04	0.0207 0.0339 0.0261 0.0232 0.0104 0.0180 0.0154 0.0183 0.0162 0.0198 0.0141 0.0093 0.0221 0.0100 0.0174 0.0195 0.0175 0.0175 0.0145 0.0131 0.0333 0.0134	0.0219 0.0314 0.0244 0.0228 0.0105 0.0166 0.0142 0.0179 0.0159 0.0200 0.0169 0.0104 0.0301 0.0100 0.0180 0.0176 0.0191 0.0137 0.0125 0.0307 0.0210	0.0223 0.0315 0.0283 0.0232 0.0119 0.0184 0.0147 0.0203 0.0192 0.0210 0.0163 0.0102 0.0320 0.0094 0.0179 0.0192 0.0183 0.0121 0.0096 0.0290 0.0217	0.0217 0.0316 0.0254 0.0232 0.0119 0.0193 0.0140 0.0193 0.0155 0.0216 0.0170 0.0113 0.0262 0.0098 0.0186 0.0176 0.0189 0.0130 0.0127 0.0298 0.0232	0.0220 0.0363 0.0271 0.0233 0.0131 0.0191 0.0156 0.0196 0.0159 0.0190 0.0179 0.0107 0.0208 0.0060 0.0164 0.0208 0.0164 0.0208 0.0194 0.0142 0.0149 0.0311 0.0228
11-01-04	0.0182	0.0150	0.0172	0.0172	0.0167
11-15-04	0.0194	0.0201	0.0198	0.0207	0.0210
11-29-04	0.0158	0.0095	0.0149	0.0140	0.0143
12-13-04	0.0230	0.0215	0.0220	0.0230	0.0220
12-29-04	0.0230	0.0210	0.0220	0.0210	0.0230

Table 1.2 Sample Type: <u>Radiolodine Cartridge</u> Analysis: lodine-131 Units: pCi/m³

End Date	APF-1 (Indicator)	APQ-1 (Indicator)	APP-1 (Indicator)	APC-1 (Indicator)	APE-30 (Control)	
Required LLD	► <u>0.07</u>	0.07	<u>0.07</u>	<u>0.07</u>	0.07	
01-12-04	< 0.013	<0.014	<0.013	<0.015	<0.014	
01-26-04	< 0.016	<0.010	< 0.015	< 0.012	<0.011	
02-09-04	< 0.017	<0.015	<0.016	<0.018	<0.015	
02-25-04	< 0.014	< 0.014	<0.012	< 0.009	< 0.014	
03-08-04	< 0.018	<0.016	<0.019	<0.016	<0.015	
03-22-04	< 0.015	< 0.014	<0.016	< 0.013	< 0.014	
04-05-04	< 0.016	< 0.013	<0.017	<0.011	<0.014	
04-20-04	< 0.014	<0.014	<0.014	<0.012	<0.013	
05-03-04	< 0.018	<0.013	<0.017	<0.015	<0.015	
05-17-04	< 0.014	<0.016	<0.017	<0.013	<0.013	
06-01-04	< 0.021	<0.020	<0.019	<0.014	<0.023	
06-14-04	< 0.013	<0.014	<0.016	<0.016	<0.014	
06-29-04	< 0.014	<0.015	<0.014	<0.012	<0.013	
07-12-04	< 0.018	<0.012	<0.015	<0.014	<0.016	
07-26-04	< 0.017	<0.014	<0.017	<0.010	<0.014	
08-09-04	< 0.014	<0.015	<0.015	<0.012	<0.016	
08-23-04	< 0.017	<0.009	<0.017	<0.009	<0.012	
09-07-04	< 0.024	<0.023	<0.022	<0.015	<0.019	
09-20-04	< 0.013	<0.013	<0.014	<0.013	<0.014	
10-04-04	< 0.017	<0.017	<0.014	<0.011	<0.013	
10-18-04	< 0.017	<0.015	<0.018	<0.014	<0.014	
11-01-04	< 0.016	<0.013	<0.016	<0.013	<0.013	
11-15-04	< 0.015	<0.016	<0.016	<0.011	<0.013	
11-2 9-04	< 0.015	<0.015	<0.018	<0.014	<0.013	
12-13-04	< 0.028	<0.024	<0.026	<0.015	<0.022	
12-29-04	< 0.012	<0.012	<0.013	<0.011	<0.015	

ı,

Table 1.3 Sample Type: <u>Air Particulate Filter</u> Analysis: Gamma Isotopic Units: pCi/m³

Location	Quarteriy Composite	Cs-134	Cs-137		
	Required LLD ->	<u>0.05</u>	<u>0.06</u>		
APF-1 (Indicator)	1st	< 0.003	< 0.002		
APQ-1 (Indicator)	1st	< 0.002	< 0.002		
APP-1 (Indicator)	1st	< 0.001	< 0.001		
APC-1 (Indicator)	1st	< 0.002	< 0.001		
APE-30 (Control)	1st	< 0.002	< 0.002		
APF-1 (Indicator)	2nd	< 0.002	< 0.002		
APQ-1 (Indicator)	2nd	< 0.001	< 0.002		
APP-1 (Indicator)	2nd	< 0.001	< 0.001		
APC-1 (Indicator)	2nd	< 0.001	< 0.002		
APE-30 (Control)	2nd	< 0.001	< 0.001		
APF-1 (Indicator)	3rd	< 0.003	< 0.002		
APQ-1 (Indicator)	3rd	< 0.002	< 0.002		
APP-1 (Indicator)	3rd	< 0.002	< 0.002		
APC-1 (Indicator)	3rd	< 0.002	< 0.001		
APE-30 (Control)	3rd	< 0.002	< 0 .002		
•					
APF-1 (Indicator)	4th	< 0.002	< 0.001		
APQ-1 (Indicator)	4th	< 0.002	< 0.002		
APP-1 (Indicator)	4th	< 0.002	< 0.002		
APC-1 (Indicator)	4th	< 0.001	< 0.002		
APE-30 (Control)	4th	< 0.002	< 0.001		

Annual Radiological Environmental Operating Report

Table 2.1	
Sample Type: T	nermoluminescent Dosimeters
Analysis: Gamn	na Dose

Units: mrem/Std. Qtr.

		indicator Locatio	115		
Station	1st Qtr '04	2nd Qtr '04	3rd Qtr '04	4th Qtr '04	Annual Mean '0
A-2	12	13	14	13	13
A-5	13	13	13	13	13
B-1	12	13	14	13	13
B-4	14	12	15	13	13
C-1	10	10	12	10	10
D-2	14	13	**	13	13
D-5	12	12	13	11	12
E-1	11	10	13	11	11
E-5	11	11	13	11	12
E-15	10	10	**	10	10
F-2	12	12	14	12	12
F-4*	14	14	15	15	14
F-9	12	13	**	12	13
G-2	15	15	14	13	14
G-4	10	11	11	10	11
G-8	11	9	10	10	10
H-2	12	12	13	12	12
H-8	11	12	12	11	12
J-2	12	12	13	12	12
J-15	14	14	14	13	14
K-1	12	12	13	14	13
L-1	14	14	14	13	14
M-1	12	11	12	12	12
N-1	13	12	13	13	13
P-1	10	9	10	10	10
P-6	14	13	14	13	13
Q-1	11	11	13	12	12
Q-5	11	10	11	11	11
R-1	9	8	**	7	8
R-6	10	10	11	10	10
Station	1st Qtr '04	Control Location 2nd Qtr '04	3rd Qtr '04	4th Qtr '04	Annual Mean '04
E-30	10	10	12	10	11

* Location with highest annual mean.

** TLDs at location D-2, E-15, F-9 and R-1 were missing for 3Q04.

Table 3.1 Sample Type: <u>Drinking/Surface Water</u> Analysis: Gross Beta Units: pCi/I

Quarterly Composite		Quarterly Composite		DWF/SWF-2 (Indicator)	DWE/SWE-5 (Indicator)	DWP/SWP-7 (Control)		
Required LLD	→	<u>4</u>	<u>4</u>	<u>4</u>				
1st		4.73	<3.41	2.28				
2nd		1.16	2.82	<1.65				
3rd		5.15	1.44	2.38				
Ath		4.69	1.36	1.73				

- --

Table 3.2 Sample Type: <u>Drinking/Surface Water</u> Analysis: Iodine-131 Units: pCi/I

Collection Date	SWK-1 (Indicator)	DWF/SWF-2 (Indicator)	DWE/SWE-5 (Indicator)	DWP/SWP-7 (Control)		
LLD	1	1	1	1		
12-29-03	< 6.28	< 0.88	<0.74	<0.86		
01-26-04	< 5.55	< 0.78	<0.90	<0.90		
02-25-04	< 3.94	< 0.89	<0.89	<0.79		
03-22-04	< 10.6	< 0.90	<0.89	<0.88		
04-20-04	< 4.98	< 0.74	<0.85	<0.90		
05-17-04	< 4.85	< 0.88	<0.86	<0.82		
06-14-04	< 4.14	< 0.90	<0.84	<0.88		
07-12-04	< 4.22	< 0.90	<0.90	<0.89		
08-09-04	< 4.14	< 0.86	<0.90	<0.87		
09-07-04	< 5.17	< 0.85	<0.87	<0.83		
10-04-04	< 5.32	< 0.90	<0.85	<0.87		
11-01-04	< 4.34	< 0.89	<0.83	<0.84		
11-29-04	< 4.22	< 0.88	<0.74	· <0.90		
12-29-04	< 5.38	< 0.90	<0.90	<0.90		

•

Table 3.3 Sample Type: <u>Drinking/Surface Water</u> Analysis: Gamma Isotopic Units: pCi/I

Loc	ation	Collection Date	Mn-54	Co-58	Fe-59	Co-60	Zn-65	Nb-95	Zr-95	Cs-134	Cs-137	Ba-140	La-140
<u> </u>	Required LLC	<u>0</u> →	<u>15</u>	<u>15</u>	<u>30</u>	<u>15</u>	<u>30</u>	<u>15</u>	<u>15</u>	<u>15</u>	<u>18</u>	<u>15</u>	<u>15</u>
DWF/SWF-2	(Indicator)	03-22-04	<1.20	<1.37	<3.32	<1.37	<2.41	<1.80	<2.34	<1.26	<1.22	<14.34	<5.12
DWE/SWE-5	(Indicator)	03-22-04	<1.21	<1.36	<3.23	<1.27	<2.53	<1.98	<2.57	<1.31	<1.28	<14.78	<5.33
DWP/SWP-7	(Control)	03-22-04	<1.18	<1.36	<3.22	<1.25	<2.33	<1.75	<2.40	<1.20	<1.16	<14.33	<4.59
DWF/SWF-2	(Indicator)	06-14-04	<2.73	<2.60	<3.99	<2.90	<6.86	<3.59	<5.27	<3.09	<3.40	<14.23	<3.68
DWE/SWE-5	(Indicator)	06-14-04	<4.41	<3.31	<7.60	<3.23	<6.74	<3.57	<5.02	<4.27	<4.18	<14.14	<3.10
DWP/SWP-7	(Control)	06-14-04	<3.54	<3.31	<7.47	<2.52	<7.93	<4.80	<7.26	<3.76	<3.33	<14.77	<4.11
DWF/SWF-2	(Indicator)	09-07-04	<3.39	<3.50	<7.26	<3.62	<8.55	<4.39	<6.25	<3.67	<3.62	<14.48	<6.28
DWE/SWE-5	(Indicator)	09-07-04	<3.03	<3.49	<6.89	<2.02	<7.67	<3.68	<5.68	<2.85	<2.84	<14.93	<6.97
DWP/SWP-7	(Control)	09-07-04	<3.18	<2.87	<6.53	<3.08	<6.38	<3.24	<5.62	<3.19	<3.02	<14.97	<4.80
DWF/SWF-2	(Indicator)	12-29-04	<4.13	<3.55	<5.69	<3.24	<8.55	<4.13	<6.57	<4.20	<3.81	<14.98	<4.70
DWE/SWE-5	(Indicator)	12-29-04	<3.60	<2.99	<8.46	<2.80	<7.88	<3.90	<6.66	<4.83	<3.29	<14.68	<3.72
DWP/SWP-7	(Control)	12-29-04	<4.28	<3.59	<7.77	<3.14	<9.99	<4.99	<5.81	<4.19	<3.30	<14.79	<2.19

Table 3.4 Sample Type: <u>Drinking/Surface Water</u> Analysis: Tritium Units: pCi/I

Quarter	DWF/SWF-2 (Indicator)	DWE/SWE-5 (Indicator)	SWK-1 (Indicator)	DWP/SWP-7 (Control)	
Required LLD	2000	2000	<u>3000</u>	2000	
1 st	< 595.01	< 579.05	<584.93	< 588.58	
2 nd	< 560.80	< 551.46	< 551.38	< 553.56	
3 rd	< 570.26	< 570.87	< 553.46	< 553.56	
4 th	< 573.09	< 558.27	< 564.00	< 560.20	

Table 3.5 Sample Type: <u>Surface Water</u> Analysis: Gamma Isotopic Units: pCi/I

Lc	ocation	Collection Date	Mn-54	Co-58	Fe-59	Co-60	Zn-65	Nb-95	Zr-95	Cs-134	Cs-137	Ba-140	La-140
	Required LLD	->	<u>15</u>	<u>15</u>	<u>30</u>	<u>15</u>	<u>30</u>	<u>15</u>	<u>15</u>	<u>15</u>	<u>18</u>	<u>15</u>	<u>15</u>
S (In	WK-1 dicator)	12-29-03 01-26-04 02-25-04 03-22-04 04-20-04 05-17-04 06-14-04 07-12-04 08-09-04 09-07-04 10-04-04 11-01-04 11-29-04 12-29-04	< 5.40 < 4.32 < 4.13 < 2.93 < 1.38 < 4.60 < 3.18 < 3.49 < 3.27 < 2.94 < 2.80 < 2.94 < 4.61 < 3.40 < 4.83	< 3.39 < 5.39 < 5.59 < 3.86 < 1.62 < 4.34 < 2.25 < 3.66 < 2.82 < 3.22 < 3.22 < 3.20 < 3.44 < 3.51 < 5.37	< 8.25 < 11.6 < 11.4 < 7.16 < 3.98 < 7.41 < 6.85 < 8.38 < 6.54 < 5.65 < 7.16 < 6.38 < 7.75 < 8.39 < 8.28	< 4.39 < 7.15 < 5.82 < 3.36 < 1.34 < 4.88 < 2.36 < 3.81 < 2.41 < 2.98 < 3.19 < 2.42 < 3.55 < 4.12 < 5.80	< 7.21 < 11.0 < 7.66 < 9.10 < 3.11 < 9.85 < 9.46 < 3.75 < 6.77 < 5.28 < 6.95 < 6.91 < 7.52 < 9.13	< 5.35 < 7.62 < 5.26 < 4.94 < 2.32 < 4.34 < 3.67 < 4.31 < 3.49 < 4.01 < 3.22 < 3.54 < 4.18 < 3.40 < 4.62	< 10.31 < 9.80 < 10.44 < 6.93 < 2.60 < 6.59 < 5.67 < 6.08 < 5.97 < 8.33 < 5.21 < 5.65 < 6.42 < 5.35 < 8.79	< 5.27 < 6.11 < 5.35 < 3.94 < 1.42 < 4.25 < 3.60 < 4.36 < 3.45 < 3.24 < 3.40 < 4.29 < 3.68 < 5.09	< 4.01 < 6.57 < 5.99 < 4.48 < 1.36 < 4.74 < 2.52 < 3.46 < 2.59 < 5.01 < 3.40 < 3.56 < 4.30 < 4.13 < 2.32	< 19.24 < 11.41 < 13.88 < 11.12 < 14.78 < 14.36 < 14.88 < 14.89 < 14.85 < 14.85 < 14.85 < 14.85 < 14.85 < 14.18 < 14.11 < 14.01	< 6.84 < 7.49 < 6.37 < 4.26 < 5.92 < 3.20 < 4.02 < 4.50 < 5.38 < 4.38 < 5.15 < 5.69 < 4.91 < 5.52 < 4.31

* Duplicate sample

.

Table 4.1Sample Type:SedimentAnalysis:Gamma IsotopicUnits:pCi/kg (dry)

 Loc	ation	Collection Date	Mn-54	Co-58	Co-60	Cs-134	Cs-137
	Required LLD	->	<u>n/a</u>	<u>n/a</u>	<u>n/a</u>	<u>150</u>	<u>180</u>
SHWK-1	(Indicator)	04-29-04	<43.9	<38.8	<39.1	<48.5	32.2
SHWE-3	(Indicator)	05-03-04	<36.8	<33.8	<39.5	<36.2	<39.4
SHWQ-6	(Control)	05-03-04	<36.1	<32.0	<29.8	<28.9	82.5

1

Annual Radiological Environmental Operating Report

Table 5.1 Sample Type: Milk Analysis: Iodine-131 and Gamma Isotopic Units: pCi/l

Location	Collection Date	I-131	Cs-134	Cs-137	Ba-140	La-140
Required LLC	$2 \rightarrow$	<u>1</u>	<u>15</u>	<u>18</u>	<u>15</u>	<u>15</u>
*MKE-3 (Indicator)	03-16-04 04-21-04 06-15-04 09-21-04 12-21-04	n/a <0.80 n/a n/a	n/a <4.18 n/a n/a n/a	n/a <4,64 n/a n/a n/a	n/a <3.54 n/a n/a n/a	n/a <14.94 n/a n/a
**MKR-40 (Control)	03-15-04 06-14-04 09-20-04 12-20-04	<0.87 <0.79 <0.89 n/a	<0.85 <4.01 <4.19 n/a	<4.80 <4.64 <5.27 n/a	<14.86 <13.56 <14.76 n/a	<5.66 <4.17 <3.86 n/a

* Sample not available. Cow not producing enough milk. ** Sample not available. Owner sold dairy.

Table 6.1 Sample Type: <u>Fish</u> Analysis: Gamma Isotopic Units: pCi/kg (wet)

Location	Collection Date	Species	Mn-54	Fe-59	Co-58	Co-60	Zn-65	Cs-134	Cs-137
Required LLD	→		<u>130</u>	<u>260</u>	<u>130</u>	<u>130</u>	<u>260</u>	<u>130</u>	<u>150</u>
FH-1 (Control)	10-19-04	Buffalo	<16.20	<13.86	<40.31	<12.74	<43.57	<12.20	<12.34
FH-1 (Control)	10-19-04	Carp	<11.25	<19.63	<44.71	<11.53	<39.31	<15.31	<12.10
FH-1 (Control)	10-19-04	Catfish	<15.26	<17.70	<44.44	<12.55	<37.63	<13.57	<12.27
FH-1 (Control)	10-1 9 -04	Mullet	<17.35	<17.78	<52.51	<26.72	<51.03	<15.90	<12.87
FH-2 (Indicator)	10-21-04	Buffalo	<13.49	<158.43	<45.98	<16.38	<38.06	<13.11	<8.80
FH-2 (Indicator)	10-21-04	Carp	<12.22	<16.60	<26.61	<21.39	<39.73	<10.92	<14.17
FH-2 (Indicator)	10-21-04	Catfish	<11.57	<16.66	<44.63	<18.99	<44.70	<15.41	<11.35
FH-2 (Indicator)	10-21-04	Mullet	<17.63	<21.75	<54.54	<18.02	<37.67	<16.27	<13.37
FH-3 (Indicator)	10-20-04	Buffalo	<11.75	<19.84	<38.65	<19.29	<47.42	<17.24	<12.78
FH-3 (Indicator)	10-20-04	Carp	<15.28	<14.64	<44.66	<12.77	<38.95	<13.00	<10.78
FH-3 (Indicator)	10-20-04	Catfish	<11.15	<15.17	<33.97	<14.16	<29.07	<11.02	<11.14
FH-3 (Indicator)	10-20-04	Mullet	<17.02	<18.33	<44.46	<23.79	<36.98	<13.74	<16.40

.

,

Table 7.1Sample Type:Broad Leaf VegetationAnalysis:Iodine-131 and Gamma IsotopicUnits:pCi/kg (wet)

Location	Collection Date	I-131	Cs-134	Cs-137
<u>Required LL</u>	<u>.D</u> >	<u>60</u>	<u>60</u>	<u>80</u>
BLQ-1 (Indicator)	03-02-04	<59.7	<38.8	<36.7
BLQ-1 (Indicator)	06-02-04	<52.8	<46.0	<43.4
BLQ-1 (Indicator)	09-01-04	<59.3	<30.6	<20.8
BLQ-1 (Indicator)	12-01-04	<49.3	<56.4	<43.6
BLB-1 (Indicator)	03-02-04	<56.5	<35.6	<43.1
BLB-1 (Indicator)	06-02-04	<53.8	<33.2	<53.8
BLB-1 (Indicator)	09-01-04	<59.7	<44.0	<43.9
BLB-1 (Indicator)	12-01-04	<56.8	<31.2	<41.5
BLE-20 (Control)	03-02-04	<32.0	<46.3	<25.5
BLE-20 (Control)	06-02-04	<58.5	<39.0	<41.6
BLE-20 (Control)	09-01-04	<58.7	<35.4	<48.0
BLE-20 (Control)	12-01-04	<57.4	<27.3	<29.9

Table 8.1Sample Type: Interlaboratory ComparisonAnalysis: Gross Beta, Iodine-131, Tritium and Gamma Isotopic

Sample Type (units)	Analytics #	Date	Analysis	Known value	RBS Value	RBS N-DEV	RBS N-RANGE
Charcoal Cartridge (pCi/filter)	E4171-125	6/17/2004	I-131	81.9	84	0.44	0.361
Water	E4170-125	6/17/2004	BETA	286	251	-1.43	0.028
(pCI/liter)	E4169-125	6/17/2004	CR-51	250	234	-1.09	0.142
			MN-54	77.4	84.3	1.55	0.916
			CO-58	50.8	46.3	-1.52	0.465
			FE-59	48.9	55.3	2.28	0.725
			CO-60	189	198	0.86	0.188
			ZN-65	109	107	-0.26	0.108
			I-131	84.3	87.0	0.55	0.841
			CS-134	111	109	-0.36	0.319
			CS-137	171	176	0.51	0.069
			CE-141	172.00	181	0.94	0.618
	E4242-125	9/16/2004	H-3	12000	11685	-0.45	0.224
Air Filter	E4240-125	9/16/2004	BETA	67.7	66.7	-0.26	0.174
(pCl/filter)	E4243-125	9/16/2004	CR-51	129	115	-1.83	0.458
			MN-54	105	104.7	-0.05	0.338
			CO-58	54.8	55.5	0.23	0.614
			FE-59	53.0	56.4	1.12	0.256
			CO-60	72.5	72.3	-0.05	0.807
			ZN-65	103	97.1	-0.99	0.241
			CS-134	55.8	50.6	-1.60	0.191
			CS-137	124	117	-1.02	0.143
			CE-141	145.0	143	-0.20	0.244
Sediment	E4241-125	9/16/2004	CR-51	0.348	0.293	-2.72	0.424
(pCi/gram)			MN-54	0.252	0.264	0.82	0.141
			CO-58	0.132	0.128	-0.52	0.492
			FE-59	0.127	0.140	1.77	0.558
			CO-60	0.174	0.174	0.03	0.339
			ZN-65	0.248	0.264	1.09	0.333
			CS-134	0.134	0.130	-0.52	0.485
			CS-137	0.403	0.426	1.00	0.132
			CE-141	0.348	0.352	0.20	0.424

Annual Radiological Environmental Operating Report

Table 8.1 Sample Type: Interlaboratory Comparison

Analysis: Gross Beta, Iodine-131, Tritium and Gamma Isotopic

Sample Type (units)	Analytics #	Date	Analysis	Known Value	RBS Value	RBS N-DEV	RBS N-RANGE
Milk	E4172-125	6/17/2004	CR-51	228	251	1.72	0.181
(pCi/liter)			MN-54	70.5	74.0	0.86	0.335
			CO-58	46.2	46.0	-0.07	0.384
			FE-59	44.5	44.4	-0.04	1.818
			CO-60	172	173	0.13	0.206
ï			ZN-65	99.3	91.7	-1.33	0.535
			I-131	58.2	53.7	-1.35	0.710
			CS-134	101	93.3	-1.31	0.234
			CS-137	156	147	-1.04	0.265
			CE-141	157	145	-1.29	0.263

NOTES:

(a) The known value as determined by Analytics.

(b) The normalized deviation from the known value is computed from the deviation and the standard error of the mean; ±2.00 is the warning limit, and ±3.00 is the control limit. This is a measure of accuracy of the analytical methods.

(c) The normalized range is computed from the mean range, the control limit, and the standard error of the range; +2.000 is the warning limit and +3.000 is the control limit. This is a measure of precision of the analytical methods.

-- - -

ATTACHMENT 2

Statistical Comparisons

TABLE OF CONTENTS

- - -

TABLE 2.1	STATISTICAL COMPARISON OF 2004 TLD MEASUREMENTS FROM STATIONS GROUPED BY DISTANCE	51
TABLE 2.2	STATISTICAL COMPARISON OF 2004 TLD RADIATION DOSE TO HISTORICAL DATA BY LOCATION	52
TABLE 2.3	STATISTICAL COMPARISON OF 2004 GROSS BETA ACTIVITY MEASUREMENTS ON AIR PARTICULATE FILTERS	53
TABLE 2.4	STATISTICAL COMPARISON OF 2004 GROSS BETA ACTIVITY MEASUREMENTS IN DRINKING/SURFACE WATER SAMPLES	54

Statistical Analyses

• Calculation of the Mean and Standard Deviation

The mean and standard deviation for different groups of analyses are calculated using the following equations:

$$\overline{X} = \sum_{i=1}^{n} \frac{X_i}{n}$$

and

$$S = \left(\frac{\sum_{i=l}^{n} (X_i - \overline{X})^2}{(n-l)}\right)^{0.5}$$

where:

 \overline{X} = mean of sample population,

S = standard deviation of sample population,

n = number of samples in sample population, and

 X_i = value of the i'th sample.

• Comparing Two Sample Population Means

The means of two sample populations are compared for statistical difference using the standard "t" test. The use of the test requires the assumption that the data within the populations are normally distributed and that the true standard deviations of the mean are equal for both populations. The standard "t" test tests the hypothesis that the true means of both populations are equal. The "t" value can be calculated from the equation below (obtained from the <u>CRC Standard Mathematical Tables</u>, 26th Edition (1981)):

$$t = \frac{\overline{X} - \overline{Y}}{\left(\frac{(n_x - 1)S_x^2 + (n_y - 1)S_y^2}{n_x + n_y - 2}\right)^{0.5} \left(\frac{1}{n_x} + \frac{1}{n_y}\right)^{0.5}}$$

where:

t = calculated "t" value,

 \overline{X} = mean of first data set.

- \overline{Y} = mean of second data set,
- η_x = number of variables in first data set,
- S_x = standard deviation of first data set,
- η_y = number of variables in second data set, and
- S_y = standard deviation of second data set.

The calculated "t" value is used to test the hypothesis that the true mean of the first population (m $_x$) is equal to the true mean of the second population (m $_y$) assuming that the true standard deviation of both populations are equal

 $(m_x = m_y)$. The calculated "t" value is compared to a tabular "t" value such that:

- a if t > $t_{u,n}$ then reject the hypothesis when $m_x > m_y$,
- b. if $t < -t_{\mu,n}$ then reject the hypothesis when $m_x < m_y$,
- c. if $t > t_{u/2,n}$ then reject the hypothesis when $m_x = m_y$,

where $t_{\mu/2,n}$ and $t_{\mu,n}$ are the tabular "t" values, with a preselected error (5%), confidence level $(1 - \mu)$ or $(1 - \mu/2)$, and degrees of freedom $n = n_x + n_y - 2$. Tabular values of the "t" were obtained from the <u>CRC</u> <u>Standard Mathematical Tables</u>, 26th Edition (1981).

ST	ATISTICAL COMPARISON OF STATIONS GROU	2004 TLD MEASUREMENTS FROM JPED BY DISTANCE				
	Stations Located 0-2 Miles from the Plant	Stations Located 2-5 Miles from the Plant	Stations Located more than 5 Miles from the Plant			
Mean (mRem/std.qtr.)	12.03	12.27	11.60			
Standard Deviation (mRem/std. qtr.)	1.60	1.42	1.52			
Number in Sample	62	28	26			
Calculated "t" Value (comparison of stations 0-2 and 2-5 miles from the plant to stations >5 miles from the plant)	1.15	1.67	NA*			
Tabular "t" Value at 95% Confidence(t _{0.025,n})	1.991(a)	2.008(a)	NA*			

TABLE 2.1

(a) Results indicate the mean for stations located 0-2 miles and 2-5 miles from the plant are statistically identical to the mean for stations located more than 5 miles from the plant.

* Not Applicable

C

				TABLE 2.2				
	STATISTICAL COM	PARISON OF 2004 TLD RA	DIATIO	N DOSE TO HIST	ORICAL DATA BY	LOCATION		
	an a	and an ann an Anna an A Anna an Anna an Anna an Anna an			مىچىنىدۇ ئولغۇم بىلەر مەرىكى بىلەر بىلەر بىلەر بىلەر بىلەر بىلەر بىلەر بىلەر بىل بىلەر بىلەر بىل		Units: mrem	/Std. Qtr.
Station	1990 - 2003 Avg**	1990 - 2003 Std Dev**	1990	- 2003 Range**	2004 Avg**	2004 Std Dev**	2004	Range**
A-2	14	1.4	11	18	13	0.8	12	14
A-5	13	1.5	10	17	13	0.0	13	13
B-1	13	1.6	11	19	13	0.8	12	14
B-4	13	1.1	12	17	14	1.3	12	15
C-1	9	1.3	7	13	11	1.0	10	12
D-2	12	2.1	8	19	13	0.6	13	14
D-5	12	1.7	9	18	12	0.8	11	13
E-1	11	1.3	10	16	11	1.3	10	13
E-5	12	1.7	9	17	12	1.0	11	13
E-15	11	1.9	8	16	10	0.0	10	10
E-30*	11	1.7	8	17	11	1.0	10	12
F-2	12	1.2	10	17	13	1.0	12	14
F-4	14	1.6	11	19	15	0.6	14	15
F-9	13	1.6	7	17	12	0.6	12	13
G-2	15	1.3	12	19	14	1.0	13	15
G-4	11	1.4	9	16	11	0.6	10	11
G-9	12	2.1	9	19	10	0.8	9	11
H-2	13	1.3	11	18	12	0.5	12	13
H-6	12	1.2	10	17	12	0.6	11	12
J-2	13	1.5	11	17	12	0.5	12	13
J-15	13	1.4	11	17	14	0.5	13	14
K-1	11	1.4	9	16	13	1.0	12	14
L-1	13	1.3	10	16	14	0.5	13	14
M-1	12	1.6	10	18	12	0.5	11	12
N-1	13	1.6	8	18	13	0.5	12	13
P-1	10	1.4	8	15	10	0.5	9	10
P-6	14	1.5	11	19	14	0.6	13	14
Q-1	12	1.3	10	16	12	1.0	11	13
Q-5	14	2.1	9	18	11	0.5	10	11
R-1	11	1.7	6	15	8	1.0	7	9
R-6	13	2.7	9	18	10	0.5	10	11

* Control Location

.

** Significant outliers were removed from data sets.

PERS data indicates an average of 20 mrem for all indicator locations with a range of 11 to 33 and an average control of 18 mrem.

S	STATISTICAL COMPARISON OF 2004 GROSS BETA ACTIVITY MEASUREMENTS ON AIR PARTICULATE FILTERS							
SAMPLE STATION	APF-1	APQ-1	APP-1	APC-1	APE-30			
Mean (10 ⁻³ pCi/m ³)	18.7	18.6	19.3	19.1	19.3			
Standard Deviation (10 ⁻³ pCi/m ³)	6.00	5.98	6.12	5.48	6.06			
Number in Sample	26	26	26	26	26			
Calculated "t" Value (comparison of the indicator stations to the control station)	0.40	0.47	0.01	0.13	NA*			
Tabular "t" Value at 95% Confidence(t _{0.025,n})	2.011(a)	2.011(a)	2.011(a)	2.011(a)	NA*			

TABLE 2.3

(a) Results indicate the mean for the indicator stations is statistically identical to the mean for the control station.

. ·

* Not Applicable

.

STATI MEAS	STATISTICAL COMPARISON OF 2004 GROSS BETA ACTIVITY MEASUREMENTS IN DRINKING/SURFACE WATER SAMPLES							
	DWF-2	DWE-5	DWP-7					
Mean (pCi/liter)	3.9	1.9	2.1					
Standard Deviation (pCi/liter)	1.61	0.67	0.29					
Number in Sample	4	3	3					
Calculated "t" Value (comparison of the indicator stations to the control station)	2.26	0.61	NA*					
Tabular "t" Value at 95% Confidence(t _{o.o25,n})	2.571(a)	2.776(a)	NA*					

TABLE 2.4

(a) Results indicate the mean for the indicator stations is statistically identical to the mean for the control station.

* Not Applicable

ATTACHMENT 3

.

2004 Revised ODCM Tables

SAMPLE	LOCATION	ANALYSIS	FREQUENCY	VOLUME	
TLD	A-2, B-1, C-1, D-2, E-1, F-2, G-2, H-2, J-2, K-1, L-1, M-1, N-1, P-1, Q-1, R-1, A-5, B-4, D-5, E-5 F-4, G-4, H-8, P-6, Q-5 R-6, F-9, G-8, E-15, J-15 E-30	πo [™]	Quarterly	N/A	
Radioiodine and	APP-1, APQ-1, APF-1,	Gross beta ^m , I-131	Bi-Weekly	285m³/wk	
Particulates	APC-1, APE-30	γ isotopic ²³	Quarterly composite	3700m³/qtr	
Ground Water	NONE	NONE	NONE	NONE	
·	DWF-2(4)/SWF-2(4)	H-3	Quarterly composite		
Drinking Water(Surface Water	DWP-7/SWP-7 DWE-5 ⁽⁴⁾ /SWE-5 ⁽⁴⁾	Gross beta, y isotopic	Quarterly composite ⁽⁵⁾	Homogeneous 8 liters	
	SWK-1	I-131 ⁰⁷	Monthly composite		
Shoreline Sediment	SHWE-3, SHWK-1, SHWQ-6	γ isotopic	Annually	2 Kilograms	
Milk	MKE-3 	γ isotopic, I-131	Quarterly	8 liters	
Fish	FH-1, FH-2, FH-3	y isotopic	In season or Annually	500 grams	
Broad Leaf	BLQ-1, BLB-1, BLE-20	γ isotopic, I-131	Quarterly	500 grams	
Sanitary System ⁽¹¹⁾	SWR-1	y isotopic	Monthly Composite ⁽¹⁹⁾	Homogeneous 1 Liter	

RADIOLOGICAL ENVIRONMENTAL MONITORING PROGRAM

"Sample collection at specific locations may be increased at any time in order to increase the effectivenes of the REMP program.

UNT-005-014 Revision 7

Attachment 7.13 (1 of 3)

56

RADIOLOGICAL ENVIRONMENTAL MONITORING PROGRAM

LOCATION NUMBER	LOCATION DESCRIPTION	BEARING/ MILES TO PLANT	Position Latitude Longitude	יי 11. 11
	DIRECT RADIATION (TLD) (continued)			
P-1	(Westbank) Located on fence enclosing air sample station APP-1.	119° 0.84	N 30.00158 W 90.48323	
Q-1	(Westbank) Located on fence enclosing air sample station APQ-1.	132° 0.81	N 30.00355 W 90.48091	
R-1	(Westbank) Located at Waterford 1 and 2 Cooling Water Intake Structure on east-hand- rail approximately a quarter of the way down the catwalk.	147° 0.51	N 30.00181 W 90.47564	2.
A-5	(Eastbank) Located on utility pole at intersection of Oswald Avenue and US 61.	177° 4.59	N 30.06212 W 90.47334	
B-4	(Eastbank) Located on utility pole guidewire next to transmission tower south of weigh station on US 61 at St. John/ St. Charles Parish line.	197° 3.75	N 30.04717 W 90.45130	

SAMPLE LOCATION TABLE (Continued)

UNT-005-014 Revision 7

Attachment 7.14 (4 of 12)

)

RADIOLOGICAL ENVIRONMENTAL MONITORING PROGRAM

LOCATION NUMBER	LOCATION	BEARING/ MILES TO PLANT	Position Latitude Longitude		
	INGESTION				
	MILK				4.00
MKE-3 	Zeringue's (Westbank) Located at the Wobro's house on LA 18 across from Mississppi River marke r 137. in Taft.	279° 429° 4.99 2.35	N 29.98926 N 30.04224 W 90.53467 W 90.13243	4 C C C	4 FEB :
MKR-40*	(Eastbank) Located at 24254 LA Hwy 442, Holden, LA.	166° 40.7	N 30.57102 W 90.62381		FEB 24
	FISH				
FH-1*	Upstream of the plant intake structure.	N/A	N/A]	
FH-2	Downstream of the plant discharge structure.	N/A	N/A		
FH-3	(Westbank) Waterways downstream of plant discharge directed to 40 Arpent Canal.	N/A	N/A		

SAMPLE LOCATION TABLE (Continued)

* DENOTES CONTROL LOCATIONS

N/A - Not Applicable for this sampling location.

UNT-005-014 Revision 7

Attachment 7.14 (10 of 12)

213