

Spatial Analysis and Decision Assistance (SADA) Version 4

Presented by
George E. Powers
Office of Nuclear Regulatory Research
Nuclear Regulatory Commission

Environmental Assessment Methods in SADA
Denver, CO
May 25, 2005

SADA General Information

Windows--based freeware designed to integrate scientific models with decision and cost analysis frameworks in a seamless, easy to use environment.

- Visualization/GIS
- Statistical Analysis
- Geospatial Interpolation
- Geospatial Uncertainty Analysis
- Human Health Risk Assessment
- Ecological Risk Assessment
- Custom Analysis
- MARSSIM Module
- Area of Concern Frameworks
- Cost Benefit Analysis
- Sampling Designs
- Export to Arcview/Earthvision

SADA has been supported by both the DOE, EPA, and the NRC. SADA Version 3.0 had about 11000 downloads. Version 4.0 has had 800+ since December, 2004.

SADATM

Spatial Analysis and Decision Assistance

SADA General Information (cont.)

Free stand-alone package for Windows 98, 98SE, NT SP4 or higher, 2000, ME, and XP.

Contact information, updates, documentation, and downloads are available on-line at <http://www.tiem.utk.edu/~sada/>

A SADA user's group, email, annual conferences, and on-site training.

A substantial help file is included.

Conduct "black and white box" testing internally as well as an external beta release period.

Publish verification document on the website.

SADATM

Spatial Analysis and Decision Assistance

Data Formats

- SADA can accept data in two formats: comma delimited files (csv) and Microsoft Access.
- SADA requires the presence of certain fields in the data set.
 - Easting
 - Northing
 - Depth
 - Value
 - Name
- SADA can use other forms of information as well
 - Media
 - Detection
 - Date
 - CAS Number
- Any other form of meta data can be imported as well. User can plot and retrieve this meta data during an analysis.
- SADA recognizes soil, sediment, surfacewater, groundwater, air, biota, and background, and the “basic” media type. Basic is assigned to data that have no media type.

SADATM

Spatial Analysis and Decision Assistance

Importing Data Into SADA

The next step is to match the columns of information in the ascii data file to information categories that are required or may be useful in SADA.

Required information categories are followed by an (*) and must be assigned to a column in the ascii data file. A category is not assigned if the (none) option is selected in the drop down box. The Depth category is required only when data exist at varying depths. If the Detect Qualifier is not assigned, the data are assumed to be all detects.

Matching Headers with Categories

SADA has found the following columns of data in your file. These columns need to be matched to information types: required (*) or useful during analysis.

Information Category	Column Headers
Easting*	Easting
Northing*	Northing
Depth*	Depth
CAS Number	Casnumber
Contaminant Name*	Name
Values*	Value
Detect Qualifier	Detected
Media Id	Media
Date	(None)

<< Back Cancel Next >> Help

If Media ID, which denotes the type of media the contaminants are sampled in (e.g. soil or groundwater) is not assigned, SADA adds an artificial media column titled 'Basic' and the human health risk and/or ecological risk modules cannot be setup later.

After the columns have been set, press **Next>>**. SADA begins the conversion process and presents the data as it will be imported into the Data Editor.


SADATM

Spatial Analysis and Decision Assistance

Data Editor

The Data Editor provides a chance to identify errors in the data set and correct them during the import process. It may also be accessed from the **Tools Menu** at any time later for data corrections or additions.

SADA highlights cells with red if they contain an unacceptable value. To determine the exact error, place the mouse over the red cell and the yellow text box near the top explains the problem. Once there are no red cells, the process continues.



The screenshot shows the SADA Data Editor window. At the top, there is a dropdown menu and a checkbox for "Automatic Error Checking" which is checked. Below this, a yellow text box displays the error message: "Easting coordinates has a non numerical or empty value." The main area contains a table with the following columns: ID, Name, Casnumber, Easting, Northing, Depth, Value, Detected, and Media. Row 18 is highlighted, and its "Easting" cell contains "NA" and is highlighted in red. The bottom of the window has buttons for "Submit", "Check Errors", "Cancel", "Delete Record", and "Help". A "Finished" status bar is at the very bottom.

ID	Name	Casnumber	Easting	Northing	Depth	Value	Detected	Media
1	Ac-225	14265851	27596.25	21900	0	1.99657	1	SD
2	Ac-225	14265851	28310.25	21900	0	1.63026	1	SW
3	Ac-225	14265851	28935	21900	0	0.86914	1	SW
4	Ac-225	14265851	27685.5	22200	0	2.053298	1	SW
5	Ac-225	14265851	28131.75	22200	0	4.185278	1	SD
6	Ac-225	14265851	29202.75	22500	0	1.49788	1	SD
7	Ac-225	14265851	27150	23160	0	1.70351	1	SD
8	Ac-225	14265851	27685.5	22920	0	2.306226	1	SD
9	Ac-225	14265851	28042.5	23100	0	4.965262	1	GW
10	Ac-225	14265851	28221	23100	0	4.232573	1	GW
11	Ac-225	14265851	28667.25	23220	0	2.951485	1	GW
12	Ac-225	14265851	29113.5	22980	0	0.99677	1	SD
13	Ac-225	14265851	27417.75	23580	0	1.92566	1	SD
14	Ac-225	14265851	27774.75	23640	0	2.939644	1	SD
15	Ac-225	14265851	28310.25	23400	0	3.121663	1	SD
16	Ac-225	14265851	28935	23460	0	0.83135	1	SD
17	Ac-225	14265851	28200	22560	0	4.8	1	SD
18	Ac-225	14265851	NA	22500	0	3.3	1	SD
19	Ac-225	14265851	27200	22380	0	2.03	1	SD
20	Barium	7440393	27596.25	21900	0	42.77435607	1	SD
21	Barium	7440393	28310.25	21900	0	35.12087239	1	GW
22	Barium	7440393	28935	21900	0	18.48691343	1	SD
23	Barium	7440393	27685.5	22200	0	43.55465395	1	SD

It is recommended that the **Automatic Error Checking** box remain checked so SADA looks for mistakes as you type. When the user is entering or pasting large amounts of data and does not wish the process to be slowed, however, it may be preferable to uncheck the **Automatic Error Checking** box and check errors later with the **Check Errors** button.

SADATM

Spatial Analysis and Decision Assistance

The New SADA Look: Scalable Interfacing

Analysis Box


Data Type Box

Data Name Box

Labels Box

Layers Box

Interviews



Steps Window

Parameters Window

Results Window

The New SADA Look

The image shows a screenshot of the SADA software interface with several components labeled by arrows:

- Analysis Box**: Points to the top toolbar area.
- Data Type Box**: Points to the 'Soil' dropdown menu.
- Data Name Box**: Points to the 'Anthracene' dropdown menu.
- Labels Box**: Points to the '(None)' dropdown menu.
- Layers Box**: Points to the 'Z = 0' dropdown menu.
- Interviews**: Points to the 'Plot my data' dropdown menu.
- Steps Window**: Points to the left sidebar containing a list of steps (1-9) and navigation buttons.
- Parameters Window**: Points to the 'Data Query' section, which includes options for 'Duplicate' and 'Non Detected' data.
- Results Window**: Points to the scatter plot showing 'Anthracene Sample Locations (Z = 0)' with a color scale on the right.

The main window displays a list of data types: Ecological, General, Human Health, Ecological, and NRC Screening Values. The 'General' option is currently selected. The scatter plot shows data points with Easting coordinates on the x-axis (ranging from 27,026.00 to 29,202.7) and Northing coordinates on the y-axis (ranging from 21,900.00 to 22,944.00). A color scale on the right indicates values from 0.80 to 5.50.

The New SADA Look

The image shows the SADA software interface with several components labeled and arrows pointing to them:

- Analysis Box**: Points to the top toolbar.
- Data Type Box**: Points to the 'Soil' dropdown menu.
- Data Name Box**: Points to the 'Anthracene' dropdown menu.
- Labels Box**: Points to the '(None)' dropdown menu.
- Layers Box**: Points to the 'Z = 0' dropdown menu.
- Interviews**: Points to the 'Plot my data' button.
- Steps Window**: Points to the 'Steps' panel on the left, which contains a list of steps: 1. Choose your data, 2. Set vertical layers, 3. Set GIS layers, 4. Set polygons, 5. Show the results, 6. Import sampled data, 7. Format picture, 8. Auto-document, 9. Add to results gallery. It also has '<< Back', 'Next >>', and 'Show The Results' buttons.
- Parameters Window**: Points to the 'Data Query' panel, which includes options for 'Data Query' (All, Interval), 'Duplicate Data' (Use all values, Use only detected values, Use most recent value, Use most recent detected value), and 'Non Detects' (Use zero, Use half the detection limit, Use the full detection limit).
- Results Window**: Points to the 'Anthracene Sample Locations (Z = 0)' plot, which is a scatter plot showing sample locations on a grid. The y-axis is labeled 'Northings' and the x-axis is labeled 'Easting'. A color scale on the right indicates values from 0.80 to 5.50.

The New SADA Look

The image shows the SADA software interface with several callout boxes pointing to specific features:

- Analysis Box**: Points to the 'General' dropdown menu.
- Data Type Box**: Points to the 'Soil' dropdown menu.
- Data Name Box**: Points to the 'Anthracene' dropdown menu.
- Labels Box**: Points to the '(None)' dropdown menu.
- Layers Box**: Points to the 'Z = 0' dropdown menu.
- Interviews**: Points to the 'Plot my data' button.
- Steps Window**: Points to the 'Steps' panel on the left, which contains a list of 9 steps: 1. Choose your data, 2. Set vertical layers, 3. Set GIS layers, 4. Set polygons, 5. Show the results, 6. Import sampled data, 7. Format picture, 8. Auto-document, 9. Add to results gallery. It also includes 'Back', 'Next', and 'Show The Results' buttons.
- Parameters Window**: Points to the 'Data Query' panel, which includes a 'Data Query' section with 'All' selected, an 'Interval' dropdown set to '9/26/1970 to 8/7/19...', and 'Add' and 'Delete' buttons. Below this is a 'Duplicate Data' section with options: 'Use all values.', 'Use only detected values.' (selected), 'Use most recent value.', 'Use most recent detected value.', 'Use maximum', 'Use average', and 'Use minimum'. At the bottom is a 'Non Detects' section with options: 'Use zero', 'Use half the detection limit.' (selected), and 'Use the full detection limit.'
- Results Window**: Points to the 'Anthracene Sample Locations (Z = 0)' window, which displays a scatter plot of sample locations. A callout box highlights the legend, listing: Ac-225, Anthracene, Arsenic, Barium, and Pooled Data. The plot has 'Easting' on the x-axis (ranging from 27,026.00 to 29,202.7) and 'Northings' on the y-axis (ranging from 21,900.00 to 22,596.00). A color scale on the right ranges from 0.80 to 2.68.

The New SADA Look

The image shows the SADA software interface with several key components labeled:

- Analysis Box**: Points to the top toolbar area.
- Data Type Box**: Points to the 'Soil' dropdown menu.
- Data Name Box**: Points to the 'Anthracene' dropdown menu.
- Labels Box**: Points to the '(None)' dropdown menu.
- Layers Box**: Points to the 'Z = 0' dropdown menu.
- Interviews**: Points to the 'Plot my data' button.
- Steps Window**: A vertical panel on the left containing a numbered list of steps: 1. Choose your data, 2. Set vertical layers, 3. Set GIS layers, 4. Set polygons, 5. Show the results, 6. Import sampled data, 7. Format picture, 8. Auto-document, 9. Add to results gallery. It includes 'Back' and 'Next' buttons and a 'Show The Results' button.
- Parameters Window**: A central window titled 'Data Query' showing a date interval '9/26/1970 to 8/7/199' and a list of data fields: '(None)', 'Value', 'Detected', 'Media', 'Date', 'SampleID', 'Region', 'Plotted Value', and '(None)'. A green arrow points to this window.
- Results Window**: A scatter plot titled 'Anthracene Sample Locations (Z = 0)'. The x-axis is 'Easting' (ranging from 27,026.00 to 29,202.7) and the y-axis is 'Northing' (ranging from 40.00 to 98.00). A color scale on the right indicates values from 0.80 to 5.50.

The New SADA Look

Analysis Box


Data Type Box

Data Name Box

Labels Box

Layers Box

Interviews



The New SADA Look

The image shows a screenshot of the SADA software interface with several components labeled by arrows:

- Analysis Box**: Points to the top-left toolbar area.
- Data Type Box**: Points to the 'Soil' dropdown menu.
- Data Name Box**: Points to the 'Anthracene' dropdown menu.
- Labels Box**: Points to the '(None)' dropdown menu.
- Layers Box**: Points to the 'Z = 0' dropdown menu.
- Interviews**: Points to the 'Plot my data' dropdown menu.
- Steps Window**: Points to the 'Steps' panel on the left, which contains a list of steps: 1. Choose your data, 2. Set vertical layers, 3. Set GIS layers, 4. Set polygons, 5. Show the results, 6. Import sampled data, 7. Format picture, 8. Auto-document, 9. Add to results gallery. It also has '<< Back' and 'Next >>' buttons and a 'Show The Results' button.
- Parameters Window**: Points to the 'Data Query' panel, which includes a 'Data Query' section with 'All' selected, an 'Interval' dropdown set to '9/26/1970 to 8/7/199', and 'Add' and 'Delete' buttons.
- Results Window**: Points to the main plot area, which displays a map titled 'Anthracene Sample Locations (Z = 0)' with a color scale on the right ranging from 0.80 to 5.50.

The New SADA Look

The image shows the SADA software interface with several components labeled and highlighted:

- Analysis Box**: Points to the top-left menu area.
- Data Type Box**: Points to the 'Soil' dropdown menu.
- Data Name Box**: Points to the 'Anthracene' dropdown menu.
- Labels Box**: Points to the '(None)' dropdown menu.
- Layers Box**: Points to the 'Z = 0' dropdown menu.
- Interviews**: Points to the 'Plot my data' button.
- Steps Window**: A blue window on the left containing a list of 9 steps: 1. Choose your data, 2. Set vertical layers, 3. Set GIS layers, 4. Set polygons, 5. Show the results, 6. Import sampled data, 7. Format picture, 8. Auto-document, 9. Add to results gallery. It includes '<< Back' and 'Next >>' buttons and a 'Show The Results' button.
- Parameters Window**: A central window titled 'Steps' with a red 'X' close button. It contains the same 9-step list as the Steps Window, with step 6 highlighted in red. It also includes '<< Back', 'Next >>', and 'Show The Results' buttons.
- Results Window**: A window on the right showing a scatter plot of 'Sample Locations' with a color scale from 0.80 to 5.50. The x-axis is labeled 'Easting' with values 28,332.05, 28,767.40, and 29,202.7.

The New SADA Look

The image shows the SADA software interface with several callout boxes pointing to specific features:

- Analysis Box**: Points to the top toolbar.
- Data Type Box**: Points to the 'Soil' dropdown menu.
- Data Name Box**: Points to the 'Anthracene' dropdown menu.
- Labels Box**: Points to the '(None)' dropdown menu.
- Layers Box**: Points to the 'Z=0' dropdown menu.
- Interviews**: Points to the 'Plot my data' button.
- Steps Window**: Points to the 'Steps' panel on the left, which contains a numbered list of tasks: 1. Choose your data, 2. Set vertical layers, 3. Set GIS layers, 4. Set polygons, 5. Show the results, 6. Import sampled data, 7. Format picture, 8. Auto-document, 9. Add to results gallery. It also includes '<< Back', 'Next >>', and 'Show The Results' buttons.
- Parameters Window**: Points to the 'Data Query' panel, which contains:
 - Data Query**: Radio buttons for 'All' (selected) and 'Interval' (with a date range dropdown '9/26/1970 to 8/7/199'). 'Add' and 'Delete' buttons.
 - Duplicate Data**: Radio buttons for 'Use all values.', 'Use only detected values.' (selected), 'Use most recent value.', and 'Use most recent detected value.'.
 - Non Detects**: Radio buttons for 'Use zero', 'Use half the detection limit.' (selected), and 'Use the full detection limit.'

A large black-bordered box on the right side of the image provides a detailed view of the 'Data Query' and 'Duplicate Data' sections of the Parameters Window, showing the radio button options and the 'Add' and 'Delete' buttons.

The New SADA Look



Analysis Box

Data Type Box

Data Name Box


Labels Box

Layers Box



Steps Window

Parameters Window



Results Window

What exactly can you do in SADA?

Create initial sample designs

Import data

Plot data

Import GIS layers

Aggregate sections of the site

Calculate statistics (univariate, bivariate)

Model spatial correlation

Create contour maps

Create a kriging variance map

Perform traditional HH and Eco risk assessments
(tabular risk, screens, prgs, benchmarks)

Create a HH or Eco contoured risk map

Create a HH or Eco point risk map

Create a data screen map for HH, Eco, Custom

Create an eco point dose map

Create an contoured eco dose map

Create probability maps

Define areas of concern

Calculate cost vs cleanup

Draw a LISA Map

Develop secondary sample designs

Perform a MARSSIM data analysis

Detect and Define MARSSIM elevated area

Visualize results in 3d

Autodocument results

Create a geobayesian site conceptual model

Draw area of concern maps based on conceptual model

Calculate cost vs cleanup based on conceptual model



Update the site conceptual model

Export to ESRI or Earthvision or common window applications


Data Exploration

Spatial Data Screens

Data Plot/GIS Overlays



Polygon Selection/Cutaways



Statistics

A "Statistics" dialog box with a table of data. A yellow arrow points to the table.

Name	CAS Number	Mean	Variance	Number of Data
Ac-225	14265851	3.4	1.3	10
Beryllium and compo	7440417	75.6	640.9	10
Arsenic, Inorganic	7440382	8.3	6.	10

Statistics

- Additional univariate statistics
- Non-parametric hypothesis testing

One-Sided Sign Test

Pooled Data -
One-Sided Test

alpha = 0.05

0 + 30 - N=30 (0 Ties)
alpha = 0.05 upper=20

Ho = The median of the population of all possible differences is 0.
Ha = The median difference does not equal 0.

At alpha = 0.05 , Ho is accepted and Ha is rejected. The median of the population of all possible differences is 0. Pooled Data is as likely to exceed than is as likely to exceed Pooled Data.

OK

Univariate Statistics

Univariate Statistics

Detects Standard Deviation Range Geometric Mean

Mean Mean Absolute Deviation UCL95 Geometric UCL95

Median Coefficient of Variance Skewness Minimum Detect

Variance Interquartile Range Kurtosis Maximum Detect

Analyte	Detects	Mean	Median	Variance	Coefficient of Variance	Interquartile Range	UCL95	Geometric Mean
Ac-225	28/28	2.6368	2.6	1.2114	0.4174	1.5	2.9911	2.3925
Barium	28/28	65.3536	74.6	792.5255	0.4308	51.5	74.4157	58.036
Arsenic	30/30	31.8867	31.2	89.5295	0.2967	11.2	34.822	30.5044
Anthracene	28/28	2.6536	2.65	1.3944	0.445	1.7	3.0337	2.3812

Wilcoxon Rank Sum Test

Ac-225 - Ac-225

alpha = 0.1

Sum of Ranks for Ac-225 = 798

Sum of Ranks for Ac-225 = 798

m = 28 n = 28

WRS Critical Value = 0

Ho = The median concentration in the survey area exceeds that in the reference area by more than the decision criterion.
Ha = The median concentration in the survey unit exceeds that in the reference area by less than the LBGR.


At alpha = 0.1 , Ho is accepted and Ha is rejected. Ac-225 exceeds that of Ac-225 by more than the decision criterion.

Value	DataSet	Rank	Tied?
0.8	Ac-225	1.5	True
0.8	Ac-225	1.5	False
0.9	Ac-225	3.5	True
0.9	Ac-225	3.5	False
1	Ac-225	5.5	True
1	Ac-225	5.5	False
1.5	Ac-225	7.5	True
1.5	Ac-225	7.5	False
1.6	Ac-225	9.5	True
1.6	Ac-225	9.5	False
1.7	Ac-225	12.5	True
1.7	Ac-225	12.5	True
1.7	Ac-225	12.5	False
1.7	Ac-225	12.5	True
1.9	Ac-225	15.5	True
1.9	Ac-225	15.5	False
2	Ac-225	18.5	True
2	Ac-225	18.5	True
2	Ac-225	18.5	True

OK

MARSSIM Functionality

- Calculate sample size based on Sign Test and WRS Test
- Develop initial sample design incorporating DCGLS, Area Factors, Instrument sensitivity
- Post sampling analysis (A site passes or fails)
- Detecting and Defining Elevated Areas



Perform MARSSIM Analysis

MARSSIM Analysis for Sign Test- CI-36


Step 1- Compare all measurements to DCGLw
Fail One or more values exceed the DCGLw.

Step 2- Compare survey area mean to DCGLw
Pass Area mean is less than the DCGLw.

Step 3- Conduct statistical test versus DCGLw
Fail Sign Test comparison between the survey data and the DCGLw failed.

**Elevated areas were detected in the survey area data.
Elevated areas can be defined using the area of concern tool.**

OK



Instrumentation Check

MARSSIM Parameters for Default

Grid Area (Survey area/ Sample N) 185.3054 (10006.49/54)

Area Factor (AF) Corresponding to Grid Area 1.2 Retrieve AF from RESRAD-MARSSIM

DCGLw 5

DCGLemc 6

Minimum Detectable Concentration for Instrument 12

Current check of instrumentation has failed.



When the instrumentation check has failed, one can also find the sample size that is necessary to meet the MDC for the instrument. Enter or retrieve the grid area that corresponds to the needed Area Factor below to find the alternative sample size that meets the MDC. In the next step, you will be asked to choose between the original sample size and this new sample size.

Grid Area Corresponding to needed AF 146 Retrieve Grid Area from RESRAD-MARSSIM


Sample N Corresponding to Grid Area equal to MDC 69

Instrument Check Cancel Next



Spatial Analysis





Spatial Uncertainty



Spatial Estimation



Version 4.0 Correlation Modeling Tools



Correlation Autofit

Choose the number of correlation structures to use.

Single Structure Both Structures


Select the models to be considered during Autofit

Spherical

Exponential

Gaussian

OK Cancel



Geostatistics

SADA provides two kriging (geostatistical) models: Ordinary and Indicator kriging. Ordinary kriging assumes a normal or lognormal distribution for the data. Indicator kriging is a non parametric approach that does not assume any distribution.


Like the methods discussed in Basic Spatial Analysis Tools, both methods are based on a weighted combination of nearby samples. However, the development and expression of these weights is quite complex and beyond the scope of this training guide.

It may be helpful to think of kriging as an advanced form of the inverse distance method. Recall that the inverse distance method weights sampled values by their distance from the unsampled location.

Kriging approaches the problem in much the same way. However, rather than distance (d), the weights are based on the amount of *spatial correlation* or *spatial covariance* that samples exhibit at varying distances $C(d)$.

Spatial Correlation

If data are spatially correlated, then on average, sample points that are close to each other are more alike than sample points further away. (More complex spatial correlations exist but this type is the most common).




Spatial Correlation

The degree to which data are more or less “alike” for any given distance can be calculated. SADA uses the *semi-variogram* method, which returns a measure of variance for any given distance of separation. This measure is defined as half of the average squared difference between values separated by distance **h**. The term **h** is referred to as the *lag* or *lag distance*.


$$\gamma(\mathbf{h}) = \frac{1}{2N(\mathbf{h})} \sum_{i=1}^{N(\mathbf{h})} (x_i - y_i)^2$$

where $N(\mathbf{h})$ is the number of pairs separated by distance **h**, x_i is the starting sample point (tail), and y_i is the ending sample point (head).



Spatial Correlation


Rarely in practice, will you ever have any sample points separated by exactly a lag distance h . Therefore, a lag tolerance centered about the lag distance will permit a capture of more data points in the calculation of $\gamma(h)$. In the figure below, all data points within the blue shaded area will be used.



So if we are interested in the variance of all data points separated by 10 feet and we permit a lag tolerance of 2 feet. We will actually be calculating the variance of all pairs of data between 9 and 11 feet apart.

Spatial Correlation


Although assigning a lag tolerance helps, most cases will never have enough samples separated by a lag - tol/2 to lag + tol/2 along a straight line to calculate the semivariogram value. Therefore, an angle tolerance, θ , is also introduced to expand the region and to include more points in the calculation of the semivariogram value for the specified lag distance. In the figure below, all data points within the blue shaded area will be used.



θ = Angle Tolerance


Spatial Correlation

If we repeat this operation for a number of lag distances, we would generate a cone shaped object expanding outward from the point of interest. This cone would be partitioned by lag groups centered about our lag distances.



Spatial Correlation

As the cone stretches farther out, it opens up increasingly wide, capturing more and more data points as it moves away. In practice, geostatisticians will often apply a constraint called the *bandwidth*. This bandwidth limits the expansion of the cone to a certain width. If you do not wish to constrain the cone's expansion, specify a very large bandwidth.




SADATM

Spatial Analysis and Decision Assistance

Spatial Correlation

Our final parameter is the angle. The angle specifies in what direction you will be calculating the semi-variogram values. This is sometimes referred to as the angle of anisotropy. Now we are constraining our semi-variogram values to a certain direction.




Lag	Distance	Value
1	32	2.3
2	67	3.4
3	98	8.9
4	123	8.9

SADATM

Spatial Analysis and Decision Assistance


Spatial Correlation



The semi-variogram calculation is performed for every sampled point.


Spatial Correlation

The semi-variogram calculation is performed for every sampled point.




Spatial Correlation

The semi-variogram calculation is performed for every sampled point.




Spatial Correlation

The semi-variogram calculation is performed for every sampled point.




Spatial Correlation

The semi-variogram values are then plotted.



Lag	Distance	Value
1	32	2.3
2	67	3.4
3	98	8.9
4	123	8.9

Semi-variogram Plot




SADATM

Spatial Analysis and Decision Assistance

Spatial Correlation

Note though, by specifying an angle α , we are excluding all those data points located outside of the cone from $\alpha - \theta$ degrees to $\alpha + \theta$ degrees. In other words, we are exploring how data are correlated in a particular direction. If we find that data are more correlated in one direction than another, the data are said to be *anisotropic*. This means that data in the direction α are more alike than in other directions.




SADATM

Spatial Analysis and Decision Assistance

Spatial Correlation

In fact, if anisotropic conditions exist, the direction of highest correlation is considered the *major direction* of anisotropy. The perpendicular direction is referred to as the *minor direction* of anisotropy. The *major direction* of correlation will exhibit semi-variogram values that increase at a slower rate than any other direction.




Spatial Correlation

Theoretically, the semi-variogram values will continue to rise until they reach the *sill* value. The sill is the point at which the data are now far enough apart to be independent. The sill value should be roughly equivalent to the variance of the data set. A *semi-variogram plot* is useful in detecting the sill value and location.

Semi-variogram plot


Lag	Major (30)	Minor (120)
1	0.56	0.6
2	0.95	3.4
3	2.3	4.5
4	3.2	4.8
5	4.3	5
6	4.7	5
7	4.9	5
8	5	4.95
9	5.01	5.05
10	4.99	5



In the above example, we see a major direction at 30 degrees and the corresponding minor direction at 120 degrees. A sill value of approximately 5 is detected around 6 feet of separation.

Isotropic Variograms

In order to calculate an isotropic or *omni-directional* variogram, simply set the angle tolerance to 90 degrees and make the bandwidth significantly larger than the site. This will force the cone to consider the entire spectrum of data points.




Spatial Correlation

What about in the opposite direction?

It is assumed that correlation is symmetrical. If data are varying a certain amount at 30 degrees, then they are varying the same amount at 120 degrees.

Why don't we include those sample points in the 120 degrees direction to improve our semi-variogram calculation?

We do. Our current point of interest will be captured by the cone of those points behind it.




Three-Dimensional Variography

Three-dimensional semi-variogram calculation is the same approach as in the two-dimensional case. In addition to the previously defined parameters, a z angle (dip), z tolerance, and z bandwidth must be specified.

Z Angle (Dip) – The angle below the horizontal plane that the cone should dip.

Z Tolerance – The tolerance on this dip angle.

Z Bandwidth – The maximum distance the vertical component of the cone is permitted to go.




SADATM

Spatial Analysis and Decision Assistance

Setting Variography Model

To calculate semivariogram values, select **Correlation Modeling** from the **Steps Window** and enter the appropriate information on the **Parameters Window**. The results of two separate cones are viewed at once to provide visual comparison and check for anisotropic correlation. Press **Show Me**.



Correlation Modeling

Correlation Modeling

Variogram Type

OK Edit

Variography

Variogram Both Rose

Name	Major	Minor
Caption	Major	Minor
Lag Number	6	6
Lag Distance	150	150
Lag Tol	100	85
Angle	90	45
Tol	45	70
Band	1000	1500
Dip	0	0
ZTol	90	90
ZBand	1000	1000


Modeling

Model Not Used Not Used

Major Range		
Minor Range		
Angle		
Contribution		
Z Angle		
Z Range		
Rotation		

Autofit Nugget

Show Me





SADATM

Spatial Analysis and Decision Assistance

Rose Diagrams

Rather than viewing only on angle at a time, users can view semivariogram values in all directions at once. They can then choose an angle of interest by clicking on the rose diagram map. SADA will show the semivariogram values for that direction.




SADATM

Spatial Analysis and Decision Assistance

Secondary Information

- The term secondary information describes a collection of information that may be either quantitative or qualitative in nature. This collection of information is not the direct subject of interest. It is however related and may assist in characterization of the primary subject, particularly within a spatial context.
- Direct measurements of the subject may be costly or perhaps dangerous to obtain. This results in only a few explicit samples.
- If secondary information is available in great quantities, it may improve heterogeneity in the final results.

Impact of Secondary Information



*Taken from Pierre Goovaerts' Presentation "Performance comparison of geostatistical algorithms for incorporating elevation into the mapping of precipitation"

SADATM

Spatial Analysis and Decision Assistance

Geobayesian History

- The U.S. Nuclear Regulatory Commission is interested in explicitly using all relevant information about a contaminated site to create a better design strategy for subsurface (3d) sampling.
- The interest originates from final status decommissioning surveys conducted by NRC.
- Issues in two-dimensional applications have been worked out in the MARSSIM guidance developed by NRC, EPA, and DOE.
- MARSSIM guidance is best suited for 2d applications because of the role that walk over radiological scans play in the process.
- The goal is to identify an analogous approach to MARSSIM for 3d, particularly when faced with sparse data sets.

Geobayesian History

- In many cases, potentially useful information is known about the site that can drive sampling and characterization strategies. These may include site history, geology, and previous sampling.
- An approach is needed to explicitly use these varying sources of information in a formal geospatial framework to drive the location of final survey samples, to characterize the radiological risk, and to support closure decisions.
- A number of approaches are being evaluated. The first is a method originally formalized as the Adaptive Sampling and Analysis Program (ASAP) at Argonne National Laboratory. This method integrates a standard bayesian approach with indicator kriging.
- This approach was the basis for the SADA geobayesian module, created by NRC and the University of Tennessee in the freeware program SADA.
- Other models under consideration include Co-kriging, and Markov Bayes.

SADATM

Spatial Analysis and Decision Assistance

The Geobayesian Model

Historical Documents

EMPLOYEE/VENDOR/PAYEE IDENTIFIED BELOW HAS OVERPAID THE NUCLEAR REGULATORY COMMISSION FOR GOODS AND/OR SERVICES PROVIDED AND IS DUE A REFUND

EMPLOYEE/VENDOR/PAYEE CODE: _____

NAME: *Northside Community Hospital*

ADDRESS: *4742 Allen Maryland, CO2*

ADDRESS: *7846 National Lodge Blvd*

CITY: *H. Arma* STATE: *MO* ZIP: *63121*

TRANS CODE: *21*

TRANS TYPE: _____ FUND: _____ JOB CODE: _____ AMOUNT: *\$400.00*

TRANS TYPE: *IR* FUND: *81523* JOB CODE: *INTR* AMOUNT: _____

TRANS TYPE: *IR* FUND: *81023* JOB CODE: *ADSH* AMOUNT: _____


TRANS TYPE: *IR* FUND: *81023* JOB CODE: *FINE* AMOUNT: _____

TOTAL REFUND AMOUNT: *\$400.00*

COMMENTS: *App dtd 6/20/97/OK 2761/75 expmt*


PREPARED BY: *Shirley Burtfield* DATE: *July 11, 1997*

Geotechnical data



Previous data
(secondary or direct)

X	Y	Z	CAS	Names	Values	Detect
27273.89	23261.16	2.6	1234123	Chlordane	0.00015	
27273.89	23261.16	5.2	1234123	Chlordane	0.00027	
27273.89	23261.16	7.8	1234123	Chlordane	0.00046	
27273.89	23261.16	10.4	1234123	Chlordane	0.00045	
27273.89	23261.16	13	1234123	Chlordane	0.00028	
27273.89	23261.16	15.6	1234123	Chlordane	0.00012	
27273.89	23261.16	18.2	1234123	Chlordane	0.00004	
27273.89	23261.16	20.8	1234123	Chlordane	0	
27273.89	23261.16	23.4	1234123	Chlordane	0	
27273.89	23261.16	26	1234123	Chlordane	0	
27512.91	23260.99	2.6	1234123	Chlordane	0.46147	
27512.91	23260.99	5.2	1234123	Chlordane	0.69144	
27512.91	23260.99	7.8	1234123	Chlordane	0.73024	
27512.91	23260.99	10.4	1234123	Chlordane	0.38323	



Probability of Exceedance




SADA™

Spatial Analysis and Decision Assistance

Case Study: Site Description

The KISKI Data Set

- Used as an example data set to test Geobayesian modeling.
- 1261 samples in shallow sediment.
- ~90 boreholes.
- Values range from near zero to 900 pCi/g.
- Contaminant name was changed.
- Large number of data, but typical spatial distribution.
- Good starting point for evaluating the new Geobayesian approach.




SADA™

Spatial Analysis and Decision Assistance





Case Study: Iterative Sampling

Depth



Decision Analysis

- Spatial Screens
- Sampling Strategies






- Spatial Risk
- Area of Concern
- Cost Benefit

SADATM

Spatial Analysis and Decision Assistance

3D Visualization

True 3d Views: Points, Blocks, and Isosurfaces



SADATM

Spatial Analysis and Decision Assistance

Sample Designs

SADA has a number of sample design strategies in Version 4.0. These strategies include initial and secondary designs. Some are based on data alone while others are based on modeling results. With the exception of a couple of exclusively 2d designs all are available in 3d dimensions.


Initial Sample Designs

- Judgmental
- Simple Random
- Simple Grid
- Simple Unaligned Grid
- Standard Grid
- Standard Unaligned Grid
- ***MARSSIM Design***
- 2d and 3d Hot Spot search designs

Secondary Sample Designs

- Threshold Radial
- Adaptive Fill
- High Value
 - (soft, simulated & unsimulated)
- High Variance
 - (soft, simulated & unsimulated)
- Extreme Value
 - (soft, simulated & unsimulated)
- Area of Concern Boundary Design
 - (soft, simulated & unsimulated)
- Minimize/Maximize Area of Concern
- LISA Designs
 - (Ripley's K, Moran's I, Geary's C)

I want to create a MARSSIM sample design



- (1) Identify the survey area
- (2) Set Class I, II, or III based on extent of contamination suspected/known
- (3) Set WRS or Sign (background or not)
- (4) View/edit DCGL and associated values (DCGL_w, LBGR, alpha, beta, sigma)
- (5) Show power curve, return N, alpha, beta
- (6) Get grid area (survey area/N)
- (7) Get grid area-area factor curve
- (8) Update AF for new grid area, calculate DCGL_{emc}, get MDC
- (9) Instrument sensitivity check
 - (1) If pass
 - (1) Show 2D Elipgrid results for circular hot spot of size grid area
 - (2) If fail
 - (1) Query for area factor based on updated grid area of (needed scan factor/DCGL)
 - (2) Recalculate N based on updated grid area and survey area
 - (3) Show elipgrid probabilities for both Ns and update grid area
 - (4) Accept original N and higher risk of missing circular hotspot or new N and lower risk of missing same hotspot size
- (10) Show MARSSIM grid or simple random sample design based on Class type

SADATM

Spatial Analysis and Decision Assistance

Calculate Grid Area and Enter Area Factor

Grid area is calculated based on the number of samples and the area of the site

Area Factor can be entered or retrieved from an excel file generated in RESRAD-MARSSIM

Click on Retrieve AF from RESRAD-MARSSIM

MARSSIM Parameters for Default


Step 1- Determine DCGLw and Sample Size

Sign Test		Sigma	0.2
DCGLw	1	alpha	0.05
LBGR	0.8	beta	0.05
		Sample Size	29

Step 2- Enter Area Factor

Grid Area (Survey area / Sample Size) 0.0344828 (1/29)

Area Factor (AF) for Grid Area >=1




SADATM

Spatial Analysis and Decision Assistance

Determining Number of Samples – Wilcoxon Rank Sum

- User inputs DCGL, LBGR, and acceptable Type I and II error rates
- Appropriate for grid designs and simple random sampling
- Used when no background is available



SADA4

S W R B C D






You need a total of 36 samples. You need 18 in your area of interest and 18 in your reference area. SADA will now plot the 18 samples in your area of interest.

OK

SADA™


Spatial Analysis and Decision Assistance

Some Example Initial Designs




Some Example Secondary Designs


Threshold Radial




High Value Design



AOC Boundary Design



Min/Max AOC




SADA Overview: Autodocumentation

- Provides transparency in the modeling process and facilitates reproducibility of results.
- SADA automatically analyzes any current result and determines what the “ingredients” of that result are. These ingredients are presented to the user, who can choose the level of documentation to create.
- Self-documentation of all parameters, models, and other relevant information.
 - Exposure concentrations
 - Risk models
 - Exposure variables
 - Geospatial parameters
 - Toxicity data
 - Images as bitmaps
- HTML format, can be exported to popular word processors

SADA Overview: Autodocumentation

- Area of concern map
- Based on HH Risk
- Utilized inverse distance as geospatial model
- Block based area of concern framework.



SADA Overview: Autodocumentation

The screenshot displays the SADA software interface. On the left, the 'Steps' window shows a list of 13 steps, with '12. Auto-document' highlighted in red. The 'Report Wizard' window is open, showing the 'Auto Documentation' section. The 'Active Report' name is 'test'. Below this, the 'Current Information to Add to the Report' section lists various options, many of which are checked. On the right, the 'test' window displays the resulting documentation output, including 'Layer Extents', 'Grid Information', and 'Spatial Parameters'.

Steps

1. Choose your data
2. Set vertical layers
3. Set GIS layers
4. Set polygons
5. Set grid specs
6. Interpolation methods
7. Search neighborhood
8. Specify decision criteria
9. Show the results
10. Cross validation
11. Format picture
12. Auto-document
13. Add to results gallery

Report Wizard

Auto Documentation

Active Report

Name: test

Change Active Report

Current Information to Add to the Report

- Picture
- GIS Files
- SHP Files
- Layering Design
- Grid Dimensions
- Variography Model Parameters
- Correlation Model Parameters
- Geospatial Parameters
- Decision Framework
- Human Health Risk Model
- Ecological Risk Model
- Decision Criteria
- ASCII Results
- Media Data

test

Layer Extents

Min Z: 0

Max Z: 10

Grid Information

Dir	Start	Size	Num
East	26900	46.055	50
North	21900	34.8	50

Spatial Parameters

Name	Value
Major Search Radius	1151.375
Minor Search Radius	1151.375
Vertical Search Radius	1
Horizontal Angle	0
Vertical Angle	0
Rotation Angle	0
Min Number of Data Values	2
Max Number of Data Values	20





Step

Analysis of Model
Elements

Documentation Output

Results Gallery

- Users can now save “static” results to the results gallery
- Users can view them, format them, and change various viewing properties
- Prevents users from having to regenerate a picture each time they want to see it
- Version 5.0 will allow dynamic results to be saved for further modeling



SADA™

Spatial Analysis and Decision Assistance

Human Health and Ecological Risk

- SADA implements EPA methods for conducting ecological and human health risk assessments
- Calculation of site-specific preliminary remediation goals
- Benchmark database for contaminant effects on ecological receptors
- Exposure modeling for humans and over 20 other terrestrial species
- Contains IRIS/HEAST toxicity databases for calculating risk from exposure
- Contains EPA default exposure parameters for the risk models
- Tabular screening and risk results
- Point screens
- Risk and dose mapping

Human Health Risk Calculations

- For each media
 - Soil, Sediment, Surface Water, Groundwater
- Exposure Scenarios
 - Residential, Industrial, Recreational, Agricultural, Excavation
- Exposure Pathways
 - Ingestion, Inhalation, Dermal Contact, Food Chain (Beef, Milk, and Vegetable Ingestion)
- IRIS and HEAST Toxicity Databases for Carcinogenic and Noncarcinogenic Effects
- Physical Parameters for Modeling
 - Bioaccumulation Factors
 - Volatilization, Particulate Emission Factors
 - Permeability Constants, Absorption Factors
 - Saturation Coefficients, Radionuclide Half-Lives

Scenario Parameters

Food Ingestion

Description	Symbol	Unit	Value
Body Weigh	BWn	kg	70
Body Weigh	BWn	kg	15
Exposure Di	ED	year	30
Exposure Di	EDn	year	24
Exposure Di	EDn	year	6
Exposure Fr	EF	day/year	350
Fraction Ing	Fim	unitless	1
Fraction Ing	Fiv	unitless	0.4
Ingestion R	IRb	kg/day	0.075
Ingestion R	IRm	kg/day	0.305
Ingestion R	IRm	kg/day	0.509
Ingestion R	IRv	kg/day	0.2
Body Weigh	BW	kg	70
Fraction Ing	Fib	unitless	1
LifeTime	LT	year	70

Chemical Parameters

Names	CAS	Analyte	Vol_org	Type	Oral RfD Wt	Oral SF Wat	Oral RfD So	Oral SF Soil	Oral RfD Die	Oral SF Diet	Inhalation R	Inhalation L
Ac-225	14265851	Radionuclid	NO	Carcinogen		3000000189		3000000518		3000000271		300000028
Barium	7440393	Inorganics	NO	Noncarcinog	0.07		0.07		0.07		0.0005	
Arsenic, Inoi	7440382	Inorganics	NO	Both	0.0003	1.5	0.0003	1.5	0.0003	1.5		4.3
Anthracene	120127	Organics	YES	Noncarcinog	0.3		0.3		0.3			

SADA™

Spatial Analysis and Decision Assistance

Human Health Risk

- PRG Calculation
- PRG Screens
- Human Health Risk

Risk Based Screening Goals: Target risk = 0.0001...

Pathways

Ingestion Dermal Fish Beef All

Inhalation External Vegetables Dairy

Rads and Nonrads/Soil/Residential/Carcinogenic

Name	CAS	Ingestion	Inhalation	All
Ac-225	14265851	5.6E+2	1.5E+5	5.6E+2
Arsenic, Inol	7440382	4.3E+1	7.4E+4	4.3E+1

Human Health Risk Results

Pathways

Ingestion Dermal Fish Beef All

Inhalation External Vegetables Dairy

Rads and Nonrads/Soil/Residential/Carcinogenic

Name	CAS	Ingestion	Inhalation	External	All
Ac-225	14265851	5.5E-7	2.E-9	5.5E-7	1.1E-6
Arsenic, Inol	7440382	1.8E-5	1.E-8		1.8E-5
Total		1.8E-5	1.2E-8	5.5E-7	1.9E-5

Screening Results: Target risk = 0.0001/Target ...

Pathways

Ingestion Dermal Fish Beef All

Inhalation External Vegetables Dairy

Rads and Nonrads/Soil/Residential/Carcinogenic

Name	CAS	Ingestion	Vegetables	All
Ac-225	14265851		Yes	Yes
Arsenic, Inol	7440382		Yes	Yes

SADATM



Spatial Analysis and Decision Assistance

Spatial Implementation of Risk Assessment

- Conventional Risk Assessment Limitations
 - Typically regulatory exposure assessment guidance recommends a summary statistic for the exposure concentration
 - Spatial information is lost when a summary statistic is used in the RA-exposure is assumed to be continuous in space and time
 - Often this lost info not recovered in the rest of the remediation process
- Reasons for incorporating spatial statistics into risk assessment
 - Maximize the use of limited resources
 - Efficiently collect data
 - Retain collected spatial info in the risk assessment
 - Use all types of available data, including expert judgment
 - To more adequately characterize the exposure distribution
 - Extrapolate from known data to cover data gaps
 - Account for spatial processes related to exposure
 - Better understand uncertainties in the exposure assessment

Human Health Spatial Risk Maps

- SADA calculates risk for each sampling point based on contaminant and exposure scenario
- Legend scale changes to risk



Human Health Risk Scenario for Pool...

Analyte		
<input type="radio"/> Rad	<input checked="" type="radio"/> Nonrad	
<input type="radio"/> Both		
Nonrad Type		
<input checked="" type="radio"/> Carcinogen	<input type="radio"/> Noncarcinogen	
Age		
<input type="radio"/> Child	<input checked="" type="radio"/> Adult	
Landuse		
<input type="radio"/> Agricultural	<input type="radio"/> Industrial	<input checked="" type="radio"/> Residential
<input type="radio"/> Excavation	<input type="radio"/> Recreational	
Pathway		
<input checked="" type="radio"/> Ingestion	<input type="radio"/> External	<input type="radio"/> Beef
<input type="radio"/> Inhalation	<input type="radio"/> Fish	<input type="radio"/> Milk
<input type="radio"/> Dermal	<input type="radio"/> Vegetables	<input type="radio"/> Total
Total Pathway Components		
<input type="checkbox"/> Ingestion	<input type="checkbox"/> External	<input type="checkbox"/> Beef
<input type="checkbox"/> Inhalation	<input type="checkbox"/> Fish	<input type="checkbox"/> Milk
<input type="checkbox"/> Dermal	<input type="checkbox"/> Vegetables	

OK



SADA™

Spatial Analysis and Decision Assistance

Ecological Risk


Ecological Risk Benchmarks

- Suitable for screening ERAs
- Compilation of ecological benchmarks for surface water, soil, and sediment
- Benchmarks a function of environmental variables where appropriate



Ecological Dose Exposures

- SADA calculates dose (mg/kg BW d) from food ingestion, soil ingestion, dermal contact, and inhalation for terrestrial exposures
- SSL, Female, Male, or Juvenile
- Over 20 different species



Set Species-Specific Terrestrial Exposure Parameters

Select a species to view (and change if necessary) the default exposure parameters used in determining the daily contaminant dose received from exposure to soil.

Long-tailed weasel SSL

Mustela frenata



Food Ingestion Parameters		Mammalian Prey Diet	
Food ingestion rate	0.1 kg dw / kg bw day	Fraction foliage	0.485 0-1
Fraction foliage	0 0-1	Fraction seed	0 0-1
Fraction seed	0 0-1	Fraction invert	0.485 0-1
Fraction invert	0 0-1	Fraction soil	0.029 0-1
Fraction mammal	1 0-1		
Soil Ingestion Parameters		Soil Inhalation Parameters	
Soil ingestion	0.039 fraction of food IR	Inhalation Rate	0.456 m ³ /day
Dermal Contact Parameters		Physical Parameters	
Adherence Factor	0.000001 kg/cm ²	Body weight	0.202 kg
Surface area	388 cm ²	Area usage factor	1 fraction

Range:
Extending from just north of the United States-Canadian border through Central America to northern South America.

Save Changes Exit

Set Terrestrial Modeling Contaminant Parameters

PCBs (total)

Chemical Constants		Dermal Contact	
Log Octanol/Water Partitioning Coefficient (Log Kow)	7.31 (mg/L)/(mg/L)	Absorption Fraction	0.06 mg/mg
Inhalation		Soil -> Invertebrate Concentration	
<input checked="" type="radio"/> Volatile <input type="radio"/> Non-Volatile		<input type="radio"/> Custom BAF	(mg/kg)/(mg/kg)
Volatilization Factor (VF)	m ³ /kg	<input checked="" type="radio"/> Kow-based BAF	33.4187 (mg/kg)/(mg/kg)
Particulate Emission Factor (PEF)	1316239339 kg/m ³	<input type="radio"/> Tissue Regression	Log-linear slope
			Log-linear intercept
Soil -> Plant Concentration: Foliage		Soil -> Small Mammal Concentration	
<input type="radio"/> Custom BAF	(mg/kg)/(mg/kg)	<input type="radio"/> Custom BAF	(mg/kg)/(mg/kg)
<input checked="" type="radio"/> Kow-based BAF	0.0313 (mg/kg)/(mg/kg)	<input type="radio"/> Tissue Regression	Log-linear slope
<input type="radio"/> Tissue Regression	Log-linear slope		Log-linear intercept
	Log-linear intercept	Diet -> Small Mammal Concentration	
Soil -> Plant Concentration: Seed		<input checked="" type="radio"/> Custom BAF	2.63 (mg/kg)/(mg/kg)
<input type="radio"/> Custom BAF	(mg/kg)/(mg/kg)	<input type="radio"/> Tissue Regression	Log-linear slope
<input checked="" type="radio"/> Kow-based BAF	0.0313 (mg/kg)/(mg/kg)		Log-linear intercept
<input type="radio"/> Tissue Regression	Log-linear slope		
	Log-linear intercept		

Save Changes Exit