

316 DEMONSTRATION


VERMONT YANKEE NUCLEAR POWER STATION

CONNECTICUT RIVER
VERNON, VERMONT



BIOLOGICAL, HYDROLOGICAL &
ENGINEERING INFORMATION

aquatec INC.



316 DEMONSTRATION

VERMONT YANKEE NUCLEAR POWER STATION
CONNECTICUT RIVER
VERNON, VERMONT

BIOLOGICAL, HYDROLOGICAL & ENGINEERING INFORMATION

AND

ENVIRONMENTAL IMPACT ASSESSMENT

(For the period 16 May to 14 October)

VERMONT YANKEE
NUCLEAR POWER CORPORATION

Philip C. Downey
Roger C. Binkerd
aquatec INC.

and

Daniel J. Marx
Vermont Yankee
Nuclear Power Corporation

June 1990

aquatec INC.

75 Green Mountain Drive, South Burlington, Vermont 05403

SUMMARY

This report presents biological, engineering, and hydrological information in support of Vermont Yankee Nuclear Power Corporation's request, pursuant to Sections 316a and 316b of the Federal Pollution Control Act Amendments of 1972, for alternate temperature standards for its electric generating station located along the Connecticut River in Vernon, Vermont, during 16 May to 14 October. The alternate temperature standards requested in this 316 Demonstration are for increases in mixed river temperatures of 2 to 5°F based upon upstream ambient temperatures:

<u>Ambient Water Temperature (°F)</u>	<u>Temperature Increase Standard (°F)</u>
Above 63	2
>59, ≤63	3
>55, ≤59	4
Below 55	5

This 316 Demonstration is the culmination of a nearly ten year effort referred to as Project SAVE (Save Available Vermont Energy). Biological data obtained as part of Project SAVE and studies conducted during the past 20 years to fulfill monitoring requirements form the basis of the bioassessment of the aquatic community. The bioassessment focused on potential effects of Vermont Yankee's open or hybrid cycle operation during 16 May to 14 October.

Changes in the phytoplankton, zooplankton, and macroinvertebrate communities as a result of Project SAVE operation were not detected during the ten year study. Fish impingement of resident fish was low during Vermont Yankee operation with negligible effect on the fish community. Resident fish, smallmouth bass, white perch, yellow perch, walleye and spottail shiner, were not excluded from any habitats in the river due to plant operation nor was growth of these species adversely affected by open or hybrid operation during Project SAVE. There was no blockage of upstream migration of adult anadromous fish, Atlantic salmon and American shad. Downstream migrating Atlantic salmon smolts moved into and through Vermont Yankee's thermal plume without any observed delay.

The results of these studies have documented that Vermont Yankee Project SAVE operation, 16 May through 14 October, has not altered the distribution, abundance or diversity of aquatic biota of the Connecticut River near Vernon. The conclusion of this study is that continued Vermont Yankee operation under the sliding scale temperature standards, presented above, will occur in such a manner that the protection and propagation of a balanced, indigenous community of shellfish, fish, and wildlife in and on the Connecticut River in the Vernon area will be assured.

TABLE OF CONTENTS

	<u>Page</u>
SUMMARY.....	ii
LIST OF TABLES.....	iv
LIST OF FIGURES.....	vi
1.0 INTRODUCTION.....	1
1.1 Historical Overview	1
1.2 Operational History of Project SAVE.....	3
2.0 316 DEMONSTRATION.....	6
2.1 Alternate Temperature Standards.....	6
2.2 Demonstration Approach.....	6
2.3 Bioassessment Summary.....	7
3.0 BIOLOGICAL INFORMATION.....	10
3.1 Phytoplankton.....	10
3.1.1 316 Demonstration Criteria.....	10
3.1.2 Phytoplankton Bioassessment.....	10
3.2 Zooplankton.....	10
3.2.1 316 Demonstration Criteria.....	10
3.2.2 Zooplankton Bioassessment.....	15
3.3 Macroinvertebrates.....	16
3.3.1 316 Demonstration Criteria.....	16
3.3.2 Macroinvertebrate Bioassessment.....	16
3.4 Fish.....	27
3.4.1 316 Demonstration Criteria.....	27
3.4.2 Fish Bioassessment.....	27
3.4.2.1 Community Composition.....	27
3.4.2.2 Fish Impingement.....	34
3.4.2.3 Fish Entrainment.....	36
3.4.2.4 Representative Important Species.....	40
ATLANTIC SALMON.....	40
AMERICAN SHAD.....	46
SMALLMOUTH BASS.....	49
WHITE PERCH.....	50
WALLEYE.....	52
YELLOW PERCH.....	54
SPOTTAIL SHINER.....	56
4.0 ENGINEERING AND HYDROLOGICAL INFORMATION.....	58
4.1 Introduction.....	58
4.2 Vermont Yankee Facilities and Operation.....	58
4.3 Vernon Dam Facilities and Operation.....	58
4.4 Effect of Discharged Heat on Connecticut River Temperatures.....	59
4.5 Thermal Impact Assessment.....	59
5.0 LITERATURE CITED.....	67

LIST OF TABLES

		<u>Page</u>
Table 1.1	Operational history of Vermont Yankee during Project SAVE and the corresponding additional energy production.....	4
Table 2.1	Alternate temperature standards applied for in this 316 Demonstration.....	6
Table 3.1	Checklist of phytoplankton collected in the Connecticut River near Vernon, Vermont.....	11
Table 3.2	Checklist of zooplankton collected in the Connecticut River near Vernon, Vermont.....	17
Table 3.3	Checklist of macroinvertebrates collected in the Connecticut River near Vernon, Vermont.....	21
Table 3.4	Historical fishing effort for juvenile and adult fish, 1968-1989.....	28
Table 3.5	Checklist of fish collected in the Connecticut River near Vernon, 1968-1989.....	30
Table 3.6	Fish species which were not observed or were first observed during 1981-1989.....	29
Table 3.7	Number and weight of fish collected upstream of Vernon Dam, 1968-1989.....	32
Table 3.8	Number and weight of fish collected downstream of Vernon Dam, 1968-1989.....	33
Table 3.9	Numbers, weights, and corresponding percentages of fish from circulating water traveling screen impingement collections, 1974-1989.....	35
Table 3.10	Collection dates for Atlantic salmon impinged on the Vermont Yankee circulating water traveling screens, 1981-1989.....	34
Table 3.11	Ichthyoplankton sampling effort, number of 0.5 meter net samples, 1982-1989.....	37
Table 3.12	Relative abundance of ichthyoplankton collected near Vernon, 1982-1989.....	38
Table 3.13	Earliest and latest dates of capture of ichthyoplankton for four game fish species, 1982-1989.....	38

LIST OF TABLES
(continued)

	<u>Page</u>
Table 3.14 Annual upstream passage of smallmouth bass, Atlantic salmon, American shad, and walleye at the Vernon Dam fishway.....	41
Table 3.15 Adult Atlantic salmon passage at Vernon Dam fishway and Vermont Yankee operation, 1985-1989.....	45
Table 3.16 Adult American shad passage at three Connecticut River fishways, 1980-1990. Percent efficiency of the previous facility passage is also expressed.....	46
Table 3.17 Sex composition of adult American shad populations from three drainages.....	48
Table 4.1 Example of data on 18 May, 0800 hours, statistics and calculations for thermal impact assessment.....	61

LIST OF FIGURES

	<u>Page</u>
Figure 1.1 Location of Vermont Yankee and biological monitoring stations.....	2
Figure 3.1 River discharge, fishway temperature, and Vermont Yankee operation during passage of adult Atlantic salmon in the Vernon Dam fishway, 1987.....	42
Figure 3.2 River discharge, fishway temperature, and Vermont Yankee operation during passage of adult Atlantic salmon in the Vernon Dam fishway, 1988.....	43
Figure 3.3 River discharge, fishway temperature, and Vermont Yankee operation during passage of adult Atlantic salmon in the Vernon Dam fishway, 1990.....	44
Figure 3.4 Mean back calculated total length in millimeters for Smallmouth Bass, 1969-1989.....	51
Figure 3.5 Mean back calculated total length in millimeters for White perch, 1969-1989.....	53
Figure 3.6 Mean back calculated total length in millimeters for walleye, 1969-1989.....	55
Figure 3.7 Mean back calculated total length in millimeters for Yellow perch, 1969-1989.....	57
Figure 4.1 Range of Station 7 temperatures, 1970-1989.....	62
Figure 4.2 Range of Connecticut River Discharge at Vernon, Vermont, 1973-1989.....	63
Figure 4.3 Distribution of simulated temperature increase using 2-5°F standard.....	64
Figure 4.4 Comparison of temperature differences with simulated increases.....	65
Figure 4.5 Mean and maximum Station 7 temperatures compared to 1989 temperatures.....	66

1.0 INTRODUCTION

1.1 Historical Overview

Vermont Yankee Nuclear Power Station is located on the west shore of the Connecticut River in Vernon, Vermont. Vernon Dam and Vernon Station, a hydroelectric generating facility, are located approximately one-half mile downstream of Vermont Yankee (Figure 1.1).

The rated reactor core thermal power level of Vermont Yankee is 1593 Megawatts (MW) providing a gross electrical output of 537 MW. The remaining 1056 MW are removed as waste heat by the circulating cooling water as it passes through the condenser.

Vermont Yankee was originally designed in 1967 with a once through condenser cooling water system. This system would have utilized Connecticut River water pumped once through the condenser and the heat would have been transferred directly to the river. During public hearings ending in 1972, which preceded the licensing and operation of Vermont Yankee, concerns were expressed of possible environmental impact of a once through mode of condenser cooling. In response to these concerns two mechanical draft cooling towers were installed. These towers were designed to dissipate directly to the atmosphere all, or part of, the heat added to the circulating water from cooling the condenser. Any use of the cooling towers, however, results in a reduction in plant generation due to losses in the plant's thermodynamic efficiency and to the energy needed to operate the cooling tower fans and pumps.

Under the conditions of the U.S. Atomic Energy Commission's full power operating license, issued early in 1973, Vermont Yankee was required to use closed cycle condenser cooling until determinations could be made concerning the possible environmental impact from the discharge of heat to the river. For the first two years of plant operation (1972-1973), only the closed cycle mode of condenser cooling was utilized. In 1974, to obtain empirical data to make a determination of environmental impact, Vermont Yankee was permitted to discharge heat in compliance with existing State of Vermont temperature standards, concurrent with an intensive biological and hydrological testing program. This program was developed with the aid of the Technical Advisory Committee (informally organized around 1968 or 1969) that consisted of fisheries and water quality experts from the States of Vermont, New Hampshire, and Massachusetts.

These studies, referred to as the "Phase Studies," were conducted during 1974-1978. As a result of these Phase Studies and the pre-operational studies, a 316 Demonstration was made for alternate temperature standards (Binkerd et al. 1978). Subsequently in 1978, Vermont Yankee was granted an amended NPDES Permit to discharge condenser cooling water to the river from 15 October to 15 May according to the following criteria:




Figure 1.1 Location of Vermont Yankee and biological monitoring stations.

1. The temperature at Station 3 during open cycle operation shall not exceed 65°F;
2. The rate of temperature change at Station 3 shall not exceed 5°F per hour; and
3. The increase in temperature above ambient at Station 3 shall not exceed 13.4°F.

During the remainder of the year the plant operated in the closed cycle mode.

In 1981, Vermont Yankee proposed a program and a study plan to the State of Vermont's Department of Water Resources for evaluating alternate operational condenser cooling modes from 16 May to 14 October. The study plan was for evaluation of the thermal discharge and impingement/entrainment effects on the Connecticut River and its aquatic biota. The goal of this program, referred to as Project SAVE (Save Available Vermont Energy), was to maximize Vermont Yankee's energy production without objectionable environmental impact.

1.2 Operational History of Project SAVE

The groundwork for Project SAVE was formulated in 1981 at meetings between Vermont Yankee representatives and the Technical Advisory Committee. To begin Project SAVE, State of Vermont temperature standards were assumed to define a level of thermal impact which was not objectionable. Test programs were discussed at these meetings and experimental studies were applied for under Section A1.6.(1) of Vermont Yankee's NPDES Permit No. VT0000264. These studies focused on intensive biological investigations, particularly fisheries, and the optimization of plant operation within temperature standards. The temperature standards in effect for this area of the Connecticut River in 1982, were the warm water fish habitat standards which allowed 1 to 5°F in mixed river temperature increase, depending upon upstream ambient water temperatures.

Although "background" biological studies and intensive planning occurred in 1981, Project SAVE investigations began in earnest in 1982 with a major riverine ichthyoplankton sampling program. Approximately 830 ichthyoplankton samples were obtained during May through August at stations upstream, adjacent to, and downstream of Vermont Yankee's discharge. Monthly fishing efforts for resident fish were increased and included monthly summer trapnetting at 16 locations and the employment of an additional method for collecting fish, electrofishing, at these same locations. Riverine sampling of fish using these two methods was supplemented by gill nets and seines.

As part of these intensified studies, tagging of adult fish began in 1982. Three species of fish, smallmouth bass (Micropterus dolomieu), walleye (Stizostedion vitreum vitreum), and white perch (Morone

americana) were part of this initial effort. Due to a high mortality rate of tagged white perch and low tag returns of walleye, this program was modified to concentrate on smallmouth bass during subsequent years of Project SAVE.

During 1982-1985, impingement of fish at the intake structure was monitored in accordance with Vermont Yankee's NPDES Permit. This Permit required sampling of the plants circulating cooling water traveling screens and service water traveling screens during summer closed cycle operation.

In 1984, the success of the American shad (Alosa sapidissima) reproducing in upper Turners Falls pool, an area within one mile downstream of Vermont Yankee, was documented. Electrofishing efforts targeted the young-of-the-year shad to investigate growth and out-migration of shad in a region receiving heated water from Vermont Yankee.

During the early years of Project SAVE, the plant was operated in either open or hybrid mode for various time periods from 16 May to 14 October (Table 1.1). Due to the increased frequency of movement of the recirculation gate demanded by Project SAVE, operational difficulties with the gate were encountered and the reliability of the gate was reduced which placed constraints on planned thermal discharges during Project SAVE; however, an increased energy production of 5,000 to 18,000 MW hours (MWh) per year was realized during 1982-1985.

Table 1.1 Operational history of Vermont Yankee during Project SAVE and the corresponding additional energy production.

<u>Year</u>	<u>Operational Dates Project SAVE</u>	<u>Five Month Additional Energy Production (MWh)</u>
1982	16 May to 31 May; and 15 September to 13 October	11,000
1983	24 June to 6 July	5,000
1984	16 May to 15 June	18,000
1985	16 May to 29 May	6,000
1986	Not Operated	0
1987	16 May to 7 August	13,877
1988	16 May to 19 July	10,120
1989	16 May to 14 October	<u>68,315</u>
	Total	132,312

In 1985, the goals of Project SAVE were reassessed since the current NPDES Permit would expire in January 1986 and would be reissued for another five year period. Also, during 1985, revised Vermont Water Quality Standards were being implemented by the Vermont Water Resources Board. As part of these new standards, the habitat designation of the Connecticut River near Vernon was changed from warmwater to coldwater fish habitat. This change meant that the coldwater fish habitat temperature standards, 1°F above ambient for thermal discharges, were now to be applied to the Connecticut River near Vernon. Presumably, this change in habitat designation was not in response to the resident fishery, but to the restoration goals for anadromous fish, Atlantic salmon (Salmo salar) and American shad, to the Connecticut River.

In the 1986 NPDES Permit, temperature standards for 15 October to 15 May remained the same as in the previous NPDES Permit; however, a 1°F temperature standard for the period of 16 May to 14 October was now permitted. Biological monitoring program requirements of the Permit were drafted in two sections. Part I, Task Oriented Monitoring, defined specific plankton, fish, and invertebrate sampling requirements. Part II, Goal Oriented Monitoring, broadly defined five goals which were to be addressed during the five year duration of the 1986 NPDES Permit. Vermont Yankee was required to design and present study proposals annually which addressed one or more of these goals. The study plans were developed in close conjunction with the Vermont Yankee Environmental Advisory Committee (VYEAC), now, a formal advisory body whose function was defined in the 1986 NPDES Permit. Also, a five year experimental program was permitted instead of the year-to-year authorization required prior to 1986.

Many of the goal oriented studies have focused on the anadromous fish being established in the Connecticut River. Juvenile American shad have been monitored annually for relative year class strength, growth, and distribution in the upper Turners Falls pool. Age and sex composition of the returning adult American shad population have been studied at the Vernon Dam fishway during 1989-1990. Beginning in the spring of each year, 1988-1990, the downstream migratory behavior of Atlantic salmon smolts was investigated by radiotelemetry. Other studies have included habitat documentation of juvenile American shad, plankton, and macroinvertebrates. Monitoring of fish and other biota, as required by Part I of the 1986 NPDES Permit, was similar in scope and methodology to previous monitoring.

In 1986, Vermont Yankee underwent an extended maintenance and refueling outage to replace pipes in the containment building. In early 1987, the recirculation gate and one intake gate were upgraded with hydraulic operators to withstand repeated cycling. Open or hybrid operation for Project SAVE continued in 1987. Energy production resulting from Project SAVE peaked in 1989 with 68,315 MWh produced.

2.0 316 DEMONSTRATION

2.1 Alternate Temperature Standards

This 316 Demonstration is the culmination of the nearly ten year effort of Project SAVE and the previous effort extended for the 316 Demonstration submitted in 1978. The data collected as part of these two studies, and the analysis of additional data obtained as part of the NPDES Permit requirements, provided the basis for this Demonstration.

This petition for alternative temperature standards is consistent with Section 316 in the Federal Water Pollution Control Act Amendments of 1972, Public Law 92-500, and with Vermont Water Quality Standards, 27 April 1990, Section 3-01,d. Both Federal and State statutes allow for alternative temperature standards provided that the present standards can be shown to be more stringent than necessary to assure the protection and propagation of a balanced indigenous population of shellfish, fish, and wildlife in and on the Connecticut River near Vernon, Vermont.

The alternate temperature standards applied for in this 316 Demonstration are for increases in mixed river temperatures of 2 to 5°F based upon upstream ambient temperatures (Table 2.1).

Table 2.1 Alternate temperature standards applied for in this 316 Demonstration.

<u>Ambient Water Temperature (°F)</u>	<u>Temperature Increase Standard (°F)</u>
Above 63	2
>59, ≤63	3
>55, ≤59	4
Below 55	5

These alternate temperature standards would apply during 16 May to 14 October and at river temperatures noted above. Temperature standards for the period of 15 October to 15 May have been addressed in the 1978-316 Demonstration.

2.2 Demonstration Approach

The effects of both Vermont Yankee's thermal discharge and cooling water intake were considered in this report with regard to requirements in Sections 316(a) and 316(b), respectively. The ~~draft~~ 316 Technical Guidance Manuals developed by the U.S. Environmental Protection Agency (EPA) (EPA 1977a, 1977b) have been utilized to format this Demonstration. In these draft Technical Guidance Manuals, the EPA has outlined three predictive approaches for 316 a and b Demonstrations:

This is most current!

1. Type 1: Lack of Prior Appreciable Harm;
2. Type 2: Protection of Representative Important Species (RIS); and
3. Type 3: Low Potential Impact.

By definition, and in concurrence with members of VYEAC in 1986, the potential impacts for the phytoplankton, zooplankton, and macroinvertebrate communities were deemed low (Type 3) in accordance with the Technical Guidance Manual decision criteria. However, additional studies have been conducted on these three communities. Interpretation of these data was presented following the Type 1 format, Lack of Prior Appreciable Harm.

The Type 2 format, Representative Important Species, was used for the fish community Demonstration. With the approval of the VYEAC, seven species of fish were selected as the RIS: Atlantic salmon, American shad, smallmouth bass, walleye, yellow perch, white perch, and spottail shiner.

2.3 Bioassessment Summary

The bioassessment of the aquatic community focused on the potential effects of Vermont Yankee's open or hybrid cycle operation from 16 May to 14 October. The proposed alternate temperature standards, a sliding scale of 2 to 5°F increase in mixed river temperatures, are discussed in Section 4, Engineering and Hydrological Information. Based upon measured temperatures and river discharges during the past 20 years, the likelihood of possible temperature increases for each day (and hour) that could be expected from Vermont Yankee operation are presented. The bioassessment of these proposed temperature standards, 2 to 5°F, was viewed in light of the fluctuating ambient temperatures. The increase in mixed river temperature due to Vermont Yankee operation was relatively low compared with these natural variabilities in ambient temperatures.

A Type 1 Demonstration narrative was conducted for the phytoplankton, zooplankton, and macroinvertebrates. With the concurrence of the VYEAC and the 316 Technical Guidance Manual's definition of low potential impact for these three communities, the impact of Vermont Yankee's operation was deemed as low. However, routine sampling and analyses of these communities have been conducted throughout Project SAVE. This sampling did not detect any changes in these communities, either in shifts towards undesirable species or loss of species as a result of plant operation.

A Type 2 Demonstration, Representative Important Species, was conducted for the fish community near Vernon. During the 18 year Vermont Yankee monitoring history of the Connecticut River while Vermont Yankee was operational, fish mortality due to cold and/or heat shock has never been observed. The fish community composition near Vernon has changed since 1981. New species of fish, American shad, Gizzard

shad, sea lamprey, Atlantic salmon, and blueback herring have been captured in the region. The presence of these fish was a result of the operation of the Turners Falls Dam and Vernon Dam fishways which began in 1980 and 1981, respectively. Resident fish community composition was similar to historical conditions, although natural long-term population cycles were noted.

Fish impingement of resident fish was low during Project SAVE, which was consistent with historical analyses of previous Vermont Yankee impingement rates. The average impingement rate of fish was 26 fish per day, averaging 320 grams. Fifty-nine juvenile Atlantic salmon were impinged during the 1981-1989 impingement studies. Four of these salmon were impinged during the period of this 316 Demonstration, 16 May to 14 October. Only one American shad was impinged during all of Project SAVE. Impingement limits for these two species of anadromous fish are currently regulated in Vermont Yankee's 1986 NPDES Permit. These limits were not approached, nor exceeded, during Project SAVE.

Ichthyoplankton entrainment has also been investigated since 1982. Ichthyoplankton samples at all locations were routinely dominated by minnows and white perch larvae. Densities in front of and in the intake structure were low, typically less than one fish per cubic meter of water, which was much less than densities observed in upstream littoral areas sampled by ichthyoplankton seine. Larval walleye were a sparse component of the ichthyoplankton near the intake. Walleye larvae were present for about two weeks in May. During the eight years of Project SAVE operation and ichthyoplankton entrainment, there were no detectable changes in the fish community near Vernon as a result of ichthyoplankton entrainment and plant operation.

Migration of anadromous fish was evaluated during Project SAVE. The analysis of American shad upstream passage at Vernon Dam did not detect any correlation between plant operation and the size of the shad run at Vernon. Major upstream migrations of American shad through the Vernon Dam fishway were documented while Vermont Yankee was operating in open or hybrid cycle modes. Juvenile American shad have inhabited upper Turners Falls pool annually since 1984. American shad were not excluded from any large areas as a result of Vermont Yankee operation.

Adult Atlantic salmon have also migrated upstream and through Vernon Dam fishway while Vermont Yankee was operating in open or hybrid cycle modes as part of Project SAVE with no blockage of migration observed. The Vernon Dam fishway passed 75 percent of the Atlantic salmon that were passed through Turners Falls fishway. Radiotelemetry studies of the Atlantic salmon smolts demonstrated rapid downstream movement of these fish into and through Vermont Yankee's thermal plume without any observed delays.

Growth of resident fish, smallmouth bass, white perch, yellow perch, and walleye, was comparable among locations and over time. Decreases in growth rate due to plant operation were not observed.

Growth rates for these species were rated as average to above average when compared with growth rates reported for the same species in other North American drainages. The resident RIS fish species were found in areas within and outside of the thermal plume with no documented exclusion of habitat due to plant operation. There was no blockage of movement of resident fish due to the thermal plume.

3.0 BIOLOGICAL INFORMATION

3.1 Phytoplankton

3.1.1 316 Demonstration Criteria

In accordance with the EPA draft Technical Guidance Manual, the decision criteria for the 316 Demonstration will be successful if it can be shown that the site is a low potential impact area for phytoplankton. The EPA defined open systems in which phytoplankton are not the major food chain base as low potential impact areas. Accordingly, most rivers, including the Connecticut River, are in this category. By definition, the area of the Connecticut River near Vermont Yankee is an area of low potential impact. This Demonstration was made with a narrative which summarized the phytoplankton analyses conducted as part of the Vermont Yankee monitoring program.

3.1.2 Phytoplankton Bioassessment

Twenty years of studies have been conducted on the phytoplankton community. Binkerd et al. (1978) summarized the 1968-1977 analyses conducted on entrainment and thermal effects on the phytoplankton community due to Vermont Yankee operation. The conclusion of this study was that the "operation of Vermont Yankee in hybrid or open cycle condenser cooling modes had not effected a demonstrable change in the phytoplankton community of the river." In addition, that study found that entrainment does not usually result in a complete loss of viability of phytoplankters.

Sampling of phytoplankton during 1981-1985 was conducted monthly at Stations 7 and 3 and at the Vermont Yankee intake and discharge structures (Aquatec et. segg. 1982). Phytoplankton sampling was required monthly, April through November, at the intake and discharge structures as part of the 1986 NPDES Permit. Two other studies conducted on phytoplankton in the Turners Falls and Vernon pools documented riverine phytoplankton composition (Shambaugh 1987, Shambaugh and Downey 1990).

Observations on phytoplankton composition in the Connecticut River near Vernon during 1981-1989 revealed a continued diversity in the number of species present (Table 3.1). Seasonal succession of the algal community and physical effects, such as river discharge, were noticeable as in previous studies. During Project SAVE, there was no shift observed in the community composition towards nuisance algae as a result of Vermont Yankee operation.

3.2 Zooplankton

3.2.1 316 Demonstration Criteria

The criteria for a Demonstration of low potential impact for zooplankton listed by the 316 draft Technical Guidance Manual are:

Table 3.1 Checklist of phytoplankton collected in the Connecticut River near Vernon, Vermont.

CYANOCHLORONTA

Cyanophyceae

Chroococcales

Chroococcaceae

Aphanocapsa sp.

Aphanocapsa elachista

Aphanothece sp.

Chroococcus sp.

Gomphosphaeria sp.

Gomphosphaeria aponina

Gomphosphaeria lacustris

Gomphosphaeria naegeliana

Merismopedia sp.

Microcystis sp.

Microcystis aeruginosa

Nostocales

Nostocaceae

Anabaena sp.

Anabaena planctonica

Anabaena spiroides

Anabaena wisconsinense

Aphanizomenon sp.

Aphanizomenon flos-aquae

Raphidiopsis sp.

Oscillatoriaceae

Lyngbya sp.

Oscillatoria sp.

Oscillatoria limnetica

Oscillatoria limosa

Oscillatoria rubescens

Oscillatoria splendida

Phormidium sp.

Rivulariaceae

Calothrix sp.

CHLOROPHYCOPHYTA

Chlorophyceae

Volvocales

Chlamydomonaceae

Carteria sp.

Chlamydomonas sp.

Volvocaceae

Eudorina sp.

Gonium sp.

Pandorina sp.

Pandorina morum

Volvox sp.

Volvox aureus

Chlorococcales

Chlorococcaceae

Characium sp.

Schroederia sp.

Tetraedron

Tetraedron caudatum

Tetraedron caudatum

var. longispinum

Tetraedron minimum

Tetraedron trigonum

Palmellaceae

Gloeocystis sp.

Gloeocystis gigas

Sphaerocystis sp.

Sphaerocystis Schroeteri

Oocystaceae

Ankistrodesmus sp.

Ankistrodesmus convolutus

Ankistrodesmus falcatus

Ankistrodesmus falcatus

var. acicularis

Ankistrodesmus spiralis

Closteriopsis sp.

Kirchneriella sp.

Kirchneriella obesa

Kirchneriella obesa var. major

Kirchneriella subsolitaria

Lagerheimia sp.

Lagerheimia ciliata

Lagerheimia quadriseta

Nephrocytium sp.

Oocystis sp.

Quadrigula sp.

Quadrigula closteroides

Quadrigula lacustris

Selenastrum sp.

Selenastrum minutum

Selenastrum Westii

Treubaria sp.

Treubaria triappendicularia

Westella sp.

Westella linearis

Micractinaceae

Golenkinia sp.

Micractinium pusillum

Dictyosphaeriaceae

Dictyosphaerium sp.

Dictyosphaerium pulchellum

Table 3.1 (continued).

Scenedesmaceae	
<u>Actinastrum</u> sp.	<u>Spirogyra</u> sp.
<u>Actinastrum hantzschii</u>	<u>Zygnema</u> sp.
<u>Actinastrum hantzschii</u> var. <u>fluviatile</u>	<u>Zygnemopsis</u> sp.
<u>Coelastrum</u> sp.	Desmidiaceae
<u>Crucigenia</u> sp.	<u>Anthrodesmus</u> sp.
<u>Crucigenia crucifera</u>	<u>Closterium</u> sp.
<u>Crucigenia fenestrata</u>	<u>Cosmarium</u> sp.
<u>Crucigenia quadrata</u>	<u>Desmidium</u> sp.
<u>Crucigenia rectangularis</u>	<u>Micrasterias</u> sp.
<u>Crucigenia tetrapedia</u>	<u>Spondylosium</u> sp.
<u>Scenedesmus</u> sp.	<u>Staurastrum</u> sp.
<u>Scenedesmus acuminatus</u>	Oedogoniales
<u>Scenedesmus arcuatus</u>	Oedogoniaceae
<u>Scenedesmus arcuatus</u> var. <u>platydisca</u>	<u>Bulbochaeta</u> sp.
<u>Scenedesmus bijuga</u>	<u>Oedogonium</u> sp.
<u>Scenedesmus dimorphus</u>	Ulotrichales
<u>Scenedesmus falcatus</u>	Ulotricaceae
<u>Scenedesmus obliquus</u>	<u>Ulothrix</u> sp.
<u>Scenedesmus quadricauda</u>	Chaetophorales
<u>Scenedesmus quadricauda</u> var. <u>westii</u>	Chaetophoraceae
<u>Tetradesmus</u> sp.	<u>Stigeoclonium</u> sp.
<u>Tetrastrum</u> sp.	Cladophorales
<u>Tetrastrum heteracanthum</u>	Cladophoraceae
Hydrodictyaceae	<u>Cladophora</u> sp.
<u>Pediastrum</u> sp.	<u>Rhizoclonium</u> sp.
<u>Pediastrum biradiatum</u>	Tetrasporales
<u>Pediastrum boryanum</u>	Tetrasporaceae
<u>Pediastrum duplex</u>	<u>Schizochlamys compacta</u>
<u>Pediastrum duplex</u> var. <u>coharens</u>	<u>Tetraspora</u> sp.
<u>Pediastrum duplex</u> var. <u>gracilimum</u>	<u>Tetraspora lacustris</u>
<u>Pediastrum duplex</u> var. <u>rotundatum</u>	<u>Tetraspora lamellosa</u>
<u>Pediastrum obtusum</u>	Microsporales
<u>Pediastrum simplex</u>	Microsporaceae
<u>Pediastrum simplex</u> var. <u>duodenarium</u>	<u>Microspora</u> sp.
<u>Pediastrum tetras</u>	EUGLENOPHYCOPHYTA
<u>Sorastrum</u> sp.	Euglenophyceae
Coccomyxaceae	Euglenales
<u>Elakatothrix</u> sp.	Euglenaceae
<u>Elakatothrix gelatinosa</u>	<u>Euglena</u> sp.
<u>Elakatothrix viridis</u>	<u>Phacus</u> sp.
Zygnematales	<u>Trachelomonas</u> sp.
Zygnemataceae	CHRYSOPHYCOPHYTA
<u>Mougeotia</u> sp.	Chrysophyceae
	Ochromonadales
	Ochromonadaceae
	<u>Uroglena</u> sp.
	Dinobryaceae
	<u>Dinobryon</u> sp.
	<u>Dinobryon bavaricum</u>

Table 3.1 (continued).

<u>Dinobryon cylindricum</u>	<u>Hannaea arcus</u>
<u>Dinobryon divergens</u>	<u>Meridion</u> sp.
<u>Dinobryon sertularia</u>	<u>Meridion circulare</u>
<u>Dinobryon Vanhoeffenii</u>	<u>Synedra</u> sp.
Synuraceae	<u>Synedra acus</u>
<u>Mallomonas</u> sp.	<u>Synedra ulna</u>
<u>Mallomonas akrokomos</u>	<u>Tabellaria</u> sp.
<u>Mallomonas pseudocoronata</u>	<u>Tabellaria fenestrata</u>
<u>Mallomonas urniformis</u>	<u>Tabellaria flocculosa</u>
<u>Synura</u> sp.	Achnanthales
<u>Synura uvella</u>	Achnanthaceae
Chromulinales	<u>Achnanthes</u> sp.
Hydruraceae	<u>Cocconeis</u> sp.
<u>Hydrurus foetidus</u>	Naviculales
<u>Rhizochrysidales</u>	Naviculaceae
<u>Rhizochrysidaceae</u>	<u>Frustulia</u> sp.
<u>Chrysamoebae</u> sp.	<u>Gyrosigma</u> sp.
<u>Diceras</u> sp.	<u>Navicula</u> sp.
Chrysocapsales	Gomphonemaceae
Chrysocapsaceae	<u>Gomphonema</u> sp.
<u>Chrysocapsa planktonica</u>	Cymbellaceae
Prymnesiophyceae	<u>Cymbella</u> sp.
Isochrysidales	Surirellales
Derepyxidaceae	Surirellaceae
<u>Cladomonas fruticulosa</u>	<u>Campylodiscus hibernicus</u>
Xanthophyceae	<u>Surirella</u> sp.
Heterochlondales	<u>Surirella ovalis</u>
Stipitococceaceae	Nitzschiales
<u>Stipitococcus</u> sp.	Nitzschiaceae
Heterococcales	<u>Nitzschia</u> sp.
Chlorotheciaceae	Coscinodiscales
<u>Centritractus</u> sp.	Coscinodiscaceae
<u>Ophiocytium</u> sp.	<u>Melosira</u> sp.
BACILLARIOPHYCOPHYTA	<u>Melosira granulata</u>
Bacillariophyceae	<u>Melosira italica</u>
Fragilariales	<u>Melosira varians</u>
Fragilariaceae	Rhizosoleniales
<u>Asterionella</u> sp.	Rhizosoleniaceae
<u>Asterionella formosa</u>	<u>Rhizosolenia</u> sp.
<u>Centronella reichelti</u>	<u>Rhizosolenia eriensis</u>
<u>Diatoma</u> sp.	Biddulphiales
<u>Diatoma vulgare</u>	Chaetoceraceae
<u>Fragilaria</u> sp.	<u>Attheya</u> sp.
<u>Fragilaria capucina</u>	PYRRROPHYCOPHYTA
<u>Fragilaria crotonensis</u>	Dinophyceae
<u>Hannaea</u> sp.	Peridinales

Table 3.1 (continued).

Peridinaceae
Peridinium sp.
Peridinium inconspicuum
Ceratiaceae
Ceratium sp.
Ceratium hirundinella
Glenodinaceae
Glenodinium sp.
CRYPTOPHYCOPHYTA
Cryptophyceae
Cryptomonadales
Cryptomonadaceae
Cryptomonads
Cryptomonads sm.
Cryptomonads lg.
Chroomonas sp.
Cryptomonas sp.
Cryptomonas ovata
INDETERMINATE
Indeterminate cells
Indeterminate colonies
Indeterminate filaments
Indeterminate flagellates
RHODOPHYCOPHYTA
Rhodophyceae
Nemalionales
Batrachospermaceae
Batrachospermum sp.

Table 3.2 (continued).

Monogononta

Flosculariacea

Conochilidae

Conochiloides sp.
Conochilus sp.
Conochilus hippocrepis
Conochilus unicornis

Hexarthridae

Hexarthra sp.

Testudinellidae

Filinia sp.
Filinia longiseta
Filinia terminalis
Pompholyx sulcata
Testudinella sp.
Testudinella patina

Collothecacea

Collothecidae

Collotheca sp.
Collotheca mutabilis
Collotheca pelagica

Ploima

Notommatidae

Cephalodella gibba
Notommata sp.

Synchaetidae

Polyarthra sp.
Polyarthra dolichoptera
Polyarthra euryptera
Polyarthra maior
Polyarthra remata
Polyarthra vulgaris
Synchaeta sp.

Ploesomatidae

Ploesoma sp.
Ploesoma hudsoni
Ploesoma lenticulare
Ploesoma truncatum

Gastropodidae

Ascomorpha sp.
Gastropus sp.
Gastropus stylifer

Trichocercidae

Trichocerca sp.
Trichocerca cylindrica
Trichocerca longiseta

Trichocerca multicornis

Trichocerca rattus

Trichocerca rousseleti

Asplanchnidae

Asplanchna sp.

Asplanchna priodonta

Brachionidae

Brachionus sp.

Brachionus angularis

Brachionus budapestinensis

Brachionus calyciflorus

Brachionus pterodinoides

Brachionus quadridentata

Brachionus rubens

Brachionus variabilis

Euchlanis sp.

Euchlanis alata

Euchlanis meneta

Euchlanis parva

Euchlanis pellucida

Kellicottia sp.

Kellicottia bostoniensis

Kellicottia longispina

Keratella sp.

Keratella cochlearis

Keratella hiemalis

Keratella quadrata

Keratella serrulata

Keratella taurocephala

Lecane sp.

Lecane flexilis

Lecane leontina

Lecane luna

Lecane mira

Lecane ohioensis

Lepadella sp.

Lepadella ovalis

Lepadella patella

Macrochaetus sp.

Monostyla sp.

Monostyla bulla

Monostyla copeis

Monostyla crenata

Monostyla lunaris

Monostyla quadridentata

Monostyla stenroosi

Table 3.2 (continued).

	<u>Mytilina</u> sp.	<u>Alona</u> <u>guttata</u>
	<u>Mytilina</u> <u>ventralis</u>	<u>Alona</u> <u>quadrangularis</u>
	<u>Mytilina</u> <u>ventralis</u> var. <u>brevispina</u>	<u>Alonella</u> sp.
	<u>Notholca</u> sp.	<u>Alonella</u> <u>excisa</u>
	<u>Notholca</u> <u>acuminata</u>	<u>Alonella</u> <u>exigua</u>
	<u>Platytias</u> sp.	<u>Alonella</u> <u>nana</u>
	<u>Platytias</u> <u>patulus</u>	<u>Camptocercus</u> sp.
	<u>Trichotria</u> sp.	<u>Camptocercus</u> <u>rectirostris</u>
	<u>Trichotria</u> <u>pocillum</u>	<u>Chydorus</u> sp.
	<u>Trichotria</u> <u>tetractis</u>	<u>Chydorus</u> <u>gibbus</u>
TARDIGRADA		<u>Chydorus</u> <u>sphaericus</u>
ANNELIDA		<u>Graptoleberis</u> <u>testudinaria</u>
Oligochaeta		<u>Leydiga</u> <u>acanthocercoides</u>
ARTHROPODA		<u>Leydiga</u> <u>quadrangularis</u>
Crustacea		<u>Monospilus</u> <u>dispar</u>
Cladocera		<u>Pleuroxus</u> sp.
Leptodoridae		<u>Pleuroxus</u> <u>denticulatus</u>
<u>Leptodora</u> sp.		<u>Pleuroxus</u> <u>hamulatus</u>
<u>Leptodora</u> <u>kindtii</u>		<u>Pleuroxus</u> <u>hastatus</u>
Sididae		<u>Pleuroxus</u> <u>procurvus</u>
<u>Diaphanosoma</u> <u>brachyurum</u>		<u>Pleuroxus</u> <u>striatus</u>
<u>Sida</u> sp.		Polyphemidae
<u>Sida</u> <u>crystallina</u>		<u>Polyphemus</u> sp.
Daphnidae		<u>Polyphemus</u> <u>pediculus</u>
<u>Ceriodaphnia</u> sp.		Ostracoda
<u>Ceriodaphnia</u> <u>lacustris</u>		Eucopepoda
<u>Ceriodaphnia</u> <u>reticulata</u>		Copepoda
<u>Daphnia</u> sp.		Copepoda nauplii
<u>Daphnia</u> <u>galeata</u> mendotae		Copepoda adult
<u>Daphnia</u> <u>retrocurva</u>		Calanoida
<u>Scapholeberis</u> sp.		Cyclopoida
<u>Scapholeberis</u> <u>kingi</u>		Insecta
Bosminidae		Ephemeroptera
<u>Bosmina</u> sp.		GASTROTRICHA
<u>Bosmina</u> <u>longirostris</u>		Chaetonotoidea
<u>Eubosmina</u> <u>coregoni</u>		Chaetonotida
Macrothricidae		Chaetonotidae
<u>Ilyocryptus</u> sp.		<u>Chaetonotus</u> sp.
<u>Ilyocryptus</u> <u>spinifer</u>		INDETERMINATE
<u>Macrothrix</u> <u>laticornis</u>		
<u>Streblocercus</u> <u>serricaudatus</u>		
Chydoridae		
<u>Acroperus</u> <u>harpae</u>		
<u>Alona</u> sp.		
<u>Alona</u> <u>affinis</u>		
<u>Alona</u> <u>costata</u>		

November. In the 1986 NPDES Permit, macroinvertebrate sampling effort at the same four stations was required three times per year, June, August, and October. A study of spatial and temporal distribution of macroinvertebrates downstream of Vernon Dam was conducted in 1986 (Wood 1988).

Over two hundred taxa, representing nine phyla, were obtained during Project SAVE's macroinvertebrate sampling (Table 3.3). No commercial, endangered, or threatened species were identified.

The discharge of heat during the 16 May to 14 October period resulted in buoyant plumes which remain near the surface in lower Vernon pool (Binkerd et al. 1978, Binkerd et al. 1990b). The potential impact of this discharge on benthos in Vernon pool is low during this time period, being limited to the area at the discharge and some near-shore areas in shallower water downstream from the discharge along the Vermont shore of Station 4. The macroinvertebrate community at Station 4 was diverse with dipterans and oligochaetes being predominant (Aquatec, et. segg. 1978). Burrowing mayflies, caddisflies, and other mayflies were consistently collected at these stations throughout the NPDES monitoring program.

Downstream of Vernon Dam, the macroinvertebrate community was also represented by a diversity of organisms. Densities of macroinvertebrates at five locations have varied seasonally ranging from 10 to 3,000 organisms per square meter (Wood 1988, Aquatec et. segg. 1982). Dipterans (Chironomids) and oligochaetes were the predominate organisms at the downstream stations. Caddisflies, particularly Hydropsychidae, were a significant component at Station 3. Other caddisflies and mayflies were represented routinely in macroinvertebrate collections at these stations. Freshwater clams were consistently a part of the macroinvertebrate community downstream during the late summer (Wood 1988).

Macroinvertebrate diurnal drift was investigated during 1986 (Wood 1988). During the daytime, chironomids were the predominate drift organisms captured. At night, the hydropsychid densities increased and these organisms were the most abundant organisms captured.

A diverse community of macroinvertebrates was present at all stations during Project SAVE. The operation of Vermont Yankee did not result in changes in the benthic community downstream which were outside the natural seasonal fluctuations. The absence of commercial and endangered species from this region and the lack of measurable harm to the community indicated that appreciable harm to the macroinvertebrate community by Project SAVE operation did not occur.

Table 3.3 Checklist of macroinvertebrates collected in the Connecticut River near Vernon, Vermont.

PROTOZOA	
PORIFERA	
Demospongea	<u>Dero</u> sp.
Haplosclerina	<u>Nais</u> sp.
Spongillidae	<u>Ophidonais serpentina</u>
COELENTERATA	<u>Paranais</u> sp.
Hydrozoa	<u>Pristina</u> sp.
Hydroida	<u>Slavina appendiculata</u>
Hydridae	<u>Stylaria</u> sp.
<u>Hydra</u> sp.	<u>Stylaria lacustris</u>
PLATYHELMINTHES	Tubificidae
Turbellaria	<u>Branchiura sowerbyi</u>
Tricladida	Lumbriculida
Planariidae	Lumbriculidae
<u>Dugesia</u> sp.	<u>Lumbriculus</u> sp.
<u>Dugesia tigrina</u>	Hirudinea
NEMATODA	Rhynchobdellida
BRYOZOA	Glossiphonidae
Gymnolaemata	<u>Actinobdella</u> sp.
Ctenostomata	<u>Helobdella</u> sp.
Paludicellidae	<u>Helobdella elongata</u>
<u>Paludicella</u> sp.	<u>Helobdella stagnalis</u>
<u>Paludicella articulata</u>	<u>Placobdella</u> sp.
Phylactolaemata	Piscicolidae
Cristatellidae	<u>Piscicola</u> sp.
<u>Cristatella mucedo</u>	Polychaeta
Fredericellidae	Sabellidae
<u>Fredericella sultana</u>	<u>Manayunkia speciosa</u>
Plumatellidae	ARTHROPODA
<u>Hyalinella</u> sp.	Arachnoidea
<u>Plumatella</u> sp.	Acariformes
<u>Plumatella repens</u>	Hydrachnidae
Lophopodidae	Arrenuridae
<u>Lophopodella carteri</u>	<u>Arrenurus</u> sp.
Pectinatellidae	Araneae
<u>Pectinatella magnifica</u>	Hydracarina
ANNELIDA	Crustacea
Oligochaeta	Cladocera
Haplotaxida	Chydoridae
Aeolosomatidae	Daphnidae
<u>Aeolosoma</u> sp.	Leptodoridae
Enchytraeidae	<u>Leptodora kindti</u>
Naididae	Sididae
<u>Allonais</u> sp.	<u>Sida crystallina</u>
<u>Chaetogaster</u> sp.	Polyphemidae
	Isopoda
	Asellidae
	<u>Asellus</u> sp.

Table 3.3 (continued).

	<u>Asellus racovitzai</u>		<u>Baetis intercalaris</u>
	<u>Caecidotea</u> sp.		<u>Baetis levitans</u>
Trichoniscidae			<u>Centroptilum</u> sp.
	<u>Trichoniscus demivirge</u>		<u>Cloeon</u> sp.
Amphipoda			<u>Pseudocloeon</u> sp.
Gammaridae		Caenidae	
	<u>Crangonyx</u> sp.		<u>Brachycercus</u> sp.
	<u>Gammarus</u> sp.		<u>Caenis</u> sp.
	<u>Synurella</u> sp.	Ephemerellidae	
Talitridae			<u>Attenella</u> sp.
	<u>Hyalella</u> sp.		<u>Ephemerella</u> sp.
	<u>Hyalella azteca</u>		<u>Ephemerella attenuata</u>
Decapoda			<u>Ephemerella needhami</u>
Cambaridae			<u>Eurylophella</u> sp.
Cambarinae			<u>Eurylophella bicolor</u>
	<u>Orconectes</u> sp.	Ephemeridae	
	<u>Orconectes limosus</u>		<u>Ephemera</u> sp.
	<u>Orconectes obscurus</u>		<u>Hexagenia</u> sp.
Ostracoda			<u>Hexagenia bilineata</u>
Eucopepoda			<u>Hexagenia limbata</u>
Insecta			<u>Hexagenia munda</u>
Collembola			<u>Hexagenia recurvata</u>
Entomobryidae			<u>Hexagenia rigida</u>
	<u>Tomocerus</u> sp.		<u>Litobrancha recurvata</u>
Isotomidae		Heptageniidae	
	<u>Isotomurus tricolor</u>		<u>Epeorus</u> sp.
Plecoptera			<u>Heptagenia</u> sp.
Taeniopterygidae			<u>Heptagenia flavescens</u>
	<u>Taenionema</u> sp.		<u>Leucrocuta</u> sp.
	<u>Taeniopteryx</u> sp.		<u>Nixe</u> sp.
Leuctridae			<u>Serratella</u> sp.
Perlidae			<u>Stenacron</u> sp.
	<u>Acroneuria</u> sp.		<u>Stenonema</u> sp.
	<u>Neoperla</u> sp.		<u>Stenonema ares</u>
	<u>Perlesta</u> sp.		<u>Stenonema bipunctatum</u>
	<u>Phasganophora</u> sp.		<u>Stenonema femoratum</u>
Perlodidae			<u>Stenonema fuscum</u>
Chloroperlidae			<u>Stenonema ithaca</u>
	<u>Hastaperla</u> sp.		<u>Stenonema luteum</u>
Ephemeroptera			<u>Stenonema tripunctatum</u>
Baetiscidae		Leptophlebiidae	
	<u>Baetisca</u> sp.		Polymitarcidae
Baetidae			<u>Ephoron</u> sp.
	<u>Acentrella</u> sp.	Siphonuridae	
	<u>Acerpenna</u> sp.		<u>Ameletus</u> sp.
	<u>Baetis</u> sp.		<u>Isonychia</u> sp.

Table 3.3 (continued).

Tricorythidae	<u>Tricorythodes</u> sp.	<u>Macronema</u> sp.
Odonata		Hydroptilidae
Aeschnidae		Hydroptilinae
<u>Basiaeschna</u> sp.		<u>Agraylea</u> sp.
<u>Boyeria</u> sp.		<u>Hydroptila</u> sp.
Coenagrionidae		<u>Ochrotrichia</u> sp.
<u>Anomalagrion</u> sp.		<u>Oxyethira</u> sp.
<u>Anomalagrion hastatum</u>		Orthotrichiinae
<u>Argia</u> sp.		<u>Orthotrichia</u> sp.
<u>Chromagrion</u> sp.		Lepidostomatidae
<u>Enallagma</u> sp.		<u>Lepidostoma</u> sp.
<u>Ischnura</u> sp.		Leptoceridae
Corduliidae		<u>Ceraclea</u> sp.
<u>Epicordulia</u> sp.		<u>Mystacides</u> sp.
<u>Neurocordulia</u> sp.		<u>Nectopsyche</u> sp.
<u>Somatochlora</u> sp.		<u>Oecetis</u> sp.
<u>Tetragoneuria</u> sp.		<u>Triaenodes</u> sp.
Gomphidae		Limnephilidae
<u>Dromogomphus</u> sp.		<u>Hydatophylax</u> sp.
<u>Gomphus</u> sp.		<u>Neophylax</u> sp.
<u>Gomphus confraternus</u>		<u>Pycnopsyche</u> sp.
<u>Ophiogomphus</u> sp.		Molannidae
<u>Stylurus</u> sp.		<u>Molanna</u> sp.
Libellulidae		Philopotamidae
<u>Pantala</u> sp.		<u>Chimarra</u> sp.
Macromiidae		Phryganeidae
<u>Didymops</u> sp.		<u>Phryganea</u> sp.
<u>Macromia</u> sp.		Polycentropodidae
Megaloptera		<u>Cernotina</u> sp.
Corydalidae		<u>Cyrnellus</u> sp.
Sialidae		<u>Neureclipsis</u> sp.
<u>Sialis</u> sp.		<u>Phylocentropus</u> sp.
Neuroptera		<u>Polycentropus</u> sp.
Sisyridae		Psychomyiidae
<u>Climacia</u> sp.		<u>Psychomyia</u> sp.
<u>Sisyra</u> sp.		Lepidoptera
Trichoptera		Pyralidae
Brachycentridae		<u>Acentria</u> sp.
<u>Brachycentrus</u> sp.		Hymenoptera
Glossosomatidae		Coleoptera
<u>Glossosoma</u> sp.		Carabidae
Hydropsychidae		<u>Brachinus</u> sp.
<u>Cheumatopsyche</u> sp.		Cerambycidae
<u>Hydropsyche</u> sp.		Chrysomelidae
<u>Hydropsyche phalerata</u>		<u>Galerucella</u> sp.
		Dytiscidae

Table 3.3 (continued).

<u>Deronectes</u> sp.	<u>Potthastia</u> sp.
Elmidae	<u>Potthastia longimanus</u>
<u>Ancyronyx</u> sp.	<u>Pseudodiamesa pertinax</u>
<u>Dubiraphia</u> sp.	Prodiamesinae
<u>Gonielmis</u> sp.	<u>Monodiamesa</u> sp.
<u>Macronychus</u> sp.	Orthoclaadiinae
<u>Neoelmis</u> sp.	<u>Acricotopus</u> sp.
<u>Optioservus</u> sp.	<u>Corynoneura</u> sp.
<u>Stenelmis</u> sp.	<u>Corynoneura taris</u>
Gyrinidae	<u>Cricotopus</u> sp.
<u>Dineutes</u> sp.	<u>Cricotopus bicinctus</u>
<u>Gyrinus</u> sp.	<u>Cricotopus intersectus</u>
Haliplidae	<u>Cricotopus sylvestris</u>
<u>Haliplus</u> sp.	<u>Cricotopus tremulus</u>
Hydrophilidae	<u>Diplocladius</u> sp.
<u>Berosus</u> sp.	<u>Eukiefferiella</u> sp.
<u>Helophorus</u> sp.	<u>Eukiefferiella potthastia</u>
Melyridae	<u>Heterotrissocladus marcidus</u>
Psephenidae	<u>Hydrobaenus</u> sp.
<u>Psephenus</u> sp.	<u>Nanocladius</u> sp.
Staphylinidae	<u>Nanocladius distinctus</u>
Corrodentia	<u>Orthocladus</u> sp.
Liposcelidae	<u>Parametricnemus lundbecki</u>
Diptera	<u>Psectrocladius</u> sp.
Anthomyiidae	<u>Psectrocladius psilopterus</u>
<u>Limnophora</u> sp.	<u>Rheocricotopus</u> sp.
Ceratopogonidae	<u>Symposiocladus</u> sp.
Chaoboridae	<u>Synorthocladus</u> sp.
<u>Chaoborus</u> sp.	<u>Thienemanniella</u> sp.
Chironomidae	<u>Tvetenia</u> sp.
Tanypodinae	<u>Tvetenia bavarica</u>
<u>Ablabesmyia</u> sp.	<u>Tvetenia discoloripes</u>
<u>Ablabesmyia annulata</u>	Chironominae
<u>Ablabesmyia mallochi</u>	<u>Axarus</u> sp.
<u>Ablabesmyia parajanta</u>	<u>Chironomus</u> sp.
<u>Alotanypus</u> sp.	<u>Chironomus decorus</u>
<u>Clinotanypus</u> sp.	<u>Cladopelma</u> sp.
<u>Coelotanypus</u> sp.	<u>Cladotanytarsus</u> sp.
<u>Djalmabatista</u> sp.	<u>Cryptochironomus</u> sp.
<u>Labrundinia</u> sp.	<u>Cryptochironomus fulvus</u>
<u>Paramerina</u> sp.	<u>Cryptotendipes</u> sp.
<u>Procladius sublettei</u>	<u>Demicryptochironomus</u> sp.
<u>Procladius</u> sp.	<u>Dicrotendipes</u> sp.
<u>Thienemannimyia</u> sp.	<u>Dicrotendipes neomodestus</u>
Diamesinae	<u>Dicrotendipes nervosus</u>
<u>Paqastia</u> sp.	<u>Einfeldia</u> sp.

Table 3.3 (continued).

<u>Endochironomus</u> sp.	<u>Tribelos iucundus</u>
<u>Endochironomus nigricans</u>	<u>Xenochironomus</u> sp.
<u>Endochironomus subtendens</u>	<u>Zavrelia</u> sp.
<u>Glyptotendipes</u> sp.	Dolichopodidae
<u>Glyptotendipes lobiferus</u>	Empididae
<u>Harnischia</u> sp.	<u>Chelifera</u> sp.
<u>Harnischia curtilamellata</u>	<u>Chelifera precatoria</u>
<u>Lauterborniella</u> sp.	<u>Hemerodromia</u> sp.
<u>Lenziella</u> sp.	Ephydridae
<u>Micropsectra</u> sp.	Heleidae
<u>Microtendipes</u> sp.	Heleinae
<u>Microtendipes caelum</u>	<u>Bezzia glabra</u>
<u>Nilothauma</u> sp.	<u>Culicoides</u> sp.
<u>Nilothauma babiye</u>	<u>Palpomyia</u> sp.
<u>Pagastiella</u> sp.	<u>Palpomyia tibialis</u>
<u>Parachironomus</u> sp.	<u>Probezzia</u> sp.
<u>Parachironomus abortivus</u>	<u>Stilobezzia</u> sp.
<u>Parachironomus carinatus</u>	Psychodidae
<u>Parachironomus frequens</u>	<u>Psychoda</u> sp.
<u>Paracladopelma</u> sp.	Simuliidae
<u>Paracladopelma longanae</u>	<u>Simulium</u> sp.
<u>Paralauterborniella</u> sp.	Tabanidae
<u>Paratanytarsus</u> sp.	<u>Chrysops</u> sp.
<u>Paratendipes</u> sp.	Tipulidae
<u>Phaenopsectra</u> sp.	<u>Antocha</u> sp.
<u>Phaenopsectra dyari</u>	Hemiptera
<u>Phaenopsectra flavipes</u>	Corixidae
<u>Polypedilum</u> sp.	Gerridae
<u>Polypedilum convictum</u>	<u>Metrobates</u> sp.
<u>Polypedilum fallax</u>	Hydrometridae
<u>Polypedilum illinoense</u>	<u>Hydrometra</u> sp.
<u>Polypedilum</u> nr. <u>scalaenum</u>	Nepidae
<u>Pseudochironomus</u> sp.	<u>Ranatra</u> sp.
<u>Rheotanytarsus</u> sp.	Homoptera
<u>Rheotanytarsus distinctissimus</u>	Aphididae
<u>Rheotanytarsus exiguus</u>	Chermidae
<u>Robackia</u> sp.	Membracidae
<u>Robackia demeigerei</u>	MOLLUSCA
<u>Stempellina</u> sp.	Gastropoda
<u>Stempellinella</u> sp.	Basommatophora
<u>Stenochironomus</u> sp.	Ancylidae
<u>Stictochironomus</u> sp.	<u>Ferrissia</u> sp.
<u>Tanytarsus</u> sp.	Lymnaeidae
<u>Tanytarsus glabrescens</u>	<u>Lymnaea</u> sp.
<u>Tanytarsus querlus</u>	Physidae
<u>Tribelos</u> sp.	<u>Physa</u> sp.

Table 3.3 (continued).

Planorbidae	<u>Gyraulus</u> sp.
	<u>Gyraulus parvus</u>
	<u>Helisoma</u> sp.
	<u>Helisoma anceps</u>
Mesogastropoda	
Hydrobiidae	<u>Amnicola</u> sp.
	<u>Amnicola limosa</u>
	<u>Birgella subglobosa</u>
Valvatidae	<u>Valvata piscinalis</u>
	<u>Valvata tricarinata</u>
Stylommatophora	
Pelecypoda	
Prionodesmacea	
Sphaeriidae	<u>Musculium</u> sp.
	<u>Pisidium</u> sp.
	<u>Pisidium henslowanum</u>
	<u>Sphaerium</u> sp.
Unionidae	<u>Alasmidonta undulata</u>
	<u>Elliptio</u> sp.
	<u>Elliptio complanata</u>
	<u>Lampsilis</u> sp.
	<u>Ligumia</u> sp.
ROTATORIA	
Monogononta	
Flosculariacea	
Conochilidae	<u>Conochilus</u> sp.

3.4 Fish

3.4.1 316 Demonstration Criteria

According to the EPA's 316 draft Technical Guidance Manual, the fish section of the 316(a) Demonstration is judged successful if the applicant can show that fish communities will not suffer appreciable harm from:

1. Direct or indirect mortality from cold shocks;
2. Direct or indirect mortality from excess heat;
3. Reduced reproductive success or growth as a result of plant discharges;
4. Exclusion from unacceptably large areas; and
5. Blockage of migration.

The first two criteria, mortality resulting from thermal shock, are addressed in Section 3.4.2.1, Community Composition, for all species of fish. The remaining three criteria, growth, habitat exclusion, and blockage of migration, are addressed where appropriate in Section 3.4.2.4, Representative Important Species. All Demonstration criteria are not always appropriate for each RIS. For example, the growth of Atlantic salmon near Vernon cannot be applied to the two migratory lifestages present, either smolt or adult, since these fish were transient. Blockage of both upstream and downstream migration is an important topic for this species and was fully addressed.

Entrainment of ichthyoplankton and impingement of fish are discussed in Sections 3.4.2.2 and 3.4.2.3. The Demonstration focused on documenting low impingement rates at the intake structure, and on changes in the fish community near Vernon in relation to Vermont Yankee Project SAVE operation.

3.4.2 Fish Bioassessment

3.4.2.1 Community Composition

The fish community of the Connecticut River near Vermont Yankee has been studied extensively as part of this 316 Demonstration and the previous 1978-316 Demonstration. Approximately 3,500 riverine fish collections, exclusive of ichthyoplankton samples, have been made since 1968 (Table 3.4). About 60 percent of this effort involved the use of trapnets, which have been used routinely throughout the 22 year period. Seines and gillnets have been used during 1968 through the early 1980's, but in recent monitoring, these methods have not been used systematically. Electrofishing was first used extensively in 1982 and subsequently became a routine methodology. Over 83,000 fish have been captured during the 22 years of monitoring.

The riverine fishing effort was part of the larger monitoring program conducted by Vermont Yankee in which over 13,000 biological samples have been collected. Biologists conducting studies as part of

Table 3.4 Historical fishing effort for juvenile and adult fish, 1968-1989. (CPE = number of fish caught per hour).

Year	Trap Net			Gill Net*			Seine**		Electro-fishing		Other		
	Coll's	Hours	Fish	CPE	Coll's	Hours	Fish	CPE	Coll's	Fish	Coll's	Fish	
1968	-	-	-	-	-	-	-	-	-	-	-	-	
1969	24	719	199	0.28	27	641	177	0.28	1	288	2	43	
1970	162	6,101	2,242	0.37	109	2,602	638	0.25	12	3,330	159	2,740	
1971	29	612	142	0.23	3	57	14	0.25	6	1,241	13	12	
1972	46	1,061	400	0.38	24	532	42	0.08					
1973	61	1,405	694	0.49	29	667	142	0.21	6	839	1	82	
1974	86	2,220	1,454	0.65	63	1,775	36	0.02	17	511	91	8	
1975	151	3,919	2,109	0.54	58	1,360	319	0.23	63	2,858	8	8	
1976	85	2,279	1,620	0.71	35	922	80	0.09	10	1,366	9	11	
1977	55	1,414	958	0.68	27	654	94	0.14	5	1,163			
1978	57	1,445	1,035	0.72	32	781	199	0.25	16	536	1	30	
1979	38	884	473	0.54	19	449	203	0.45	9	1,785			
1980	68	1,553	1,170	0.75	27	607	284	0.47	3	148			
1981	100	2,440	2,144	0.88	42	1,008	326	0.32	6	889			
1982	307	7,084	6,162	0.87	13	305	27	0.08	6	344			
1983	195	4,164	3,381	0.81	4	88	0	0.00					
1984	189	3,780	2,709	0.72					23	3,336	24	1,987	
1985	115	2,408	2,274	0.94					13	1,877	60	3,994	
1986	112	2,287	1,546	0.68					18	4,055	62	1,825	
1987	112	2,342	1,814	0.78							98	2,132	
1988	112	2,378	1,913	0.81							44	2,300	
1989	112	2,427	1,803	0.74							44	1,630	
											44	2,081	
											44	2,076	
Total	2,216				512				214		421		291

* Method not used after 1983.
 ** Method not used after 1985.

the NPDES monitoring requirements were on the Connecticut River sampling approximately 60 to 120 days annually, from May through October. During this 18 year sampling history, there has been no mortality of fish observed, either due to heat and/or cold shock, resulting from Vermont Yankee's operation.

Fish capture rates have varied from year-to-year, partly as a function of the different levels of effort and methodologies employed. The most consistent methodology employed has been trapnetting. Capture rates for trapnets, expressed as catch per unit effort (CPE), did not display any reduction in catch rates during the years. There has been a slight but discernible increase in fish capture rates during 1981-1989 as compared to previous years. *not necessarily better*

Plume attraction studies conducted as part of the earlier "Phase Studies" were summarized in the 1978-316 Demonstration. There was no apparent attraction of resident fish to the warmer waters along the Vermont shore downstream of Vermont Yankee's discharge. These studies and other more recent data indicate that the potential for attracting fish to the thermal plume areas and subsequent mortality resulting from thermal shock from increasing or decreasing temperature were negligible.

An annotated checklist of selected Connecticut River fish was presented in Binkerd et al. (1978). Biological data collected on all fish have been computerized (Park 1990). The 83,000 fish captured during 1968-1989 were represented by 39 species of fish (Table 3.5). During the Project SAVE experimental studies, 1981-1989, thirty-six species of fish were collected. Three species of fish not observed during Project SAVE were represented by very few individuals in previous years and their presence in the Connecticut River near Vernon was most likely due to incidental displacement from upstream tributaries. There were seven new species of fish recorded during 1981-1989 (Table 3.6).

Table 3.6 Fish species which were not observed or were first observed during 1981-1989.

<u>Species Not Observed</u>	<u>Common Name</u>
<u>Catostomus catostomus</u>	longnose sucker
<u>Semotilus atromaculatus</u>	creek chub
<u>Couesius plumbeus</u>	lake chub
<u>Species First Observed</u>	
<u>Notropis volucellus</u>	mimic shiner
<u>Petromyzon marinus</u>	sea lamprey
<u>Alosa sapidissima</u>	American shad
<u>Alosa aestivalis</u>	blueback herring
<u>Dorosoma cepedianum</u>	gizzard shad
<u>Rhinichthys atratulus</u>	blacknose dace
<u>Notropis atherinoides</u>	emerald shiner

Table 3.5 Checklist of fish collected in the Connecticut River near Vernon, 1968-1989. Taxonomy is in accordance with Robins et al. 1980.

<u>Scientific Name</u>	<u>Common Name</u>
Petromyzontidae	Lampreys
<u>Petromyzon marinus</u> Linnaeus	Sea lamprey
Anguillidae	Freshwater eels
<u>Anguilla rostrata</u> (Lesueur)	American eel
Clupeidae	Herrings
<u>Alosa aestivalis</u> (Mitchill)	Blueback herring
<u>Alosa sapidissima</u> (Wilson)	American shad
<u>Dorosoma cepedianum</u> (Lesueur)	Gizzard shad
Salmonidae	Trouts
<u>Salmo gairdneri</u> Richardson	Rainbow trout
<u>Salmo salar</u> Linnaeus	Atlantic salmon
<u>Salmo trutta</u> Linnaeus	Brown trout
<u>Salvelinus fontinalis</u> (Mitchill)	Brook trout
Osmeridae	Smelts
<u>Osmerus mordax</u> (Mitchill)	Rainbow smelt
Esocidae	Pikes
<u>Esox lucius</u> Linnaeus	Northern pike
<u>Esox niger</u> (Lesueur)	Chain pickerel
Cyprinidae	Carps and minnows
<u>Couesius plumbeus</u> (Agassiz)	Lake chub
<u>Cyprinus carpio</u> Linnaeus	Common carp
<u>Hybognathus regius</u> Girard	Eastern silvery minnow
<u>Notemigonus crysoleucas</u> (Mitchill)	Golden shiner
<u>Notropis atherinoides</u> Rafinesque	Emerald shiner
<u>Notropis cornutus</u> (Mitchill)	Common shiner
<u>Notropis hudsonius</u> (Clinton)	Spottail shiner
<u>Notropis volucellus</u> (Cope)	Mimic shiner
<u>Rhinichthys atratulus</u> (Hermann)	Blacknose dace
<u>Semotilus corporalis</u> (Mitchill)	Fallfish
<u>Semotilus atromaculatus</u> (Mitchill)	Creek chub
Catostomidae	Suckers
<u>Catostomus commersoni</u> (Lacepede)	White sucker
<u>Catostomus catostomus</u> (Forster)	Longnose sucker
Ictaluridae	Bullhead catfishes
<u>Ictalurus natalis</u> (Lesueur)	Yellow bullhead
<u>Ictalurus nebulosus</u> (Lesueur)	Brown bullhead
Cyprinodontidae	Killifishes
<u>Fundulus diaphanus</u> (Lesueur)	Banded killifish
Percichthyidae	Temperate basses
<u>Morone americana</u> (Gmelin)	White perch
Centrarchidae	Sunfishes
<u>Ambloplites rupestris</u> (Rafinesque)	Rock bass
<u>Lepomis gibbosus</u> (Linnaeus)	Pumpkinseed
<u>Lepomis macrochirus</u> Rafinesque	Bluegill
<u>Micropterus dolomieu</u> Lacepede	Smallmouth bass
<u>Micropterus salmoides</u> (Lacepede)	Largemouth bass
<u>Pomoxis nigromaculatus</u> (Lesueur)	Black crappie
Percidae	Perches
<u>Etheostoma olmstedi</u> Storer	Tessellated darter
<u>Perca flavescens</u> (Mitchill)	Yellow perch
<u>Stizostedion vitreum vitreum</u> (Mitchill)	Walleye
Cottidae	Sculpins
<u>Cottus cognatus</u> Heckel	Slimy sculpin

Four of these species, sea lamprey, American shad, blueback herring, and the gizzard shad, have appeared as a direct result of the installation and operation of fishways required for the Connecticut River Anadromous Fish Restoration Program. Fishways at Turners Falls and Vernon Dams had direct bearing on the passage of resident and anadromous fish into regions near Vernon. Fishway operation began in 1980 and 1981 at Turners Falls and Vernon, respectively, which coincided with the initiation of Project SAVE in 1981.

The American shad and, to a lesser extent, the blueback herring have become an established population in the upper Turners Falls pool with young-of-the-year fish being present from July through early-October. Ammocoetes (larval sea lamprey) have been collected in bottom sediments from the Connecticut River near Vernon. Gizzard shad were first recorded in the Connecticut River near Vernon in 1989. Gizzard shad spawning success in Vernon pool was documented in 1989 with the capture of about 120 young gizzard shad (Briggs and Downey 1990).

The mimic shiner was first identified ^{taxonomy may have accounted for} upstream of Vernon Dam in the West River near Brattleboro, Vermont in 1983. This species was observed in 1984 near Vernon and has since become a common minnow in fish collections.

Only one specimen of the emerald shiner was obtained in an impingement sample during Project SAVE. Two blacknose dace were collected at Station 5, upstream of Vermont Yankee, in seine samples obtained on two successive days in 1985. The presence of these two species probably was a result of incidental displacement from upstream drainages.

Although sampling methodologies have varied during the 22 years of studies, the overall relative composition can be inferred from data collected with the various gears used for juvenile/adult fish collection. The fish community near Vernon was characterized as a coolwater fish community (Tables 3.7 and 3.8) (Downey 1990e). Sunfishes, including the basses, have historically represented about 30 percent of the total fish sampled. Yellow perch, white suckers, white perch, and minnows were all numerically important components of the fish community. Overall, the population has generally stayed consistent with some minor annual variations observed. American shad in the river near Vernon, a direct result of fisheries management goals, have become an important community downstream of Vernon Dam. Mimic shiners, a recently identified species from the area, were an important component of the minnow community.

Long-term cycles in resident populations were also noticeable. A strong recruitment of white perch was quite apparent in 1982 and 1983. This species, which had represented five to ten percent of the fish community in the 1970's, accounted for nearly 20 percent of the community during 1982 and 1983. Relative numbers of yellow perch in the community have increased in recent years, also reflecting improved

Figure 3.7 Number and weight of fish collected upstream of Vernon Dam, 1968-1989. Impingement and ichthyoplankton samples are not included.

	1968-1980				1981-1989			
	Total Number (No.)	(%)	Total Weight (g)	(%)	Total Number (No.)	(%)	Total Weight (g)	(%)
Sea Lamprey					1	<0.1		
American eel	20	0.1	15,310	0.5	123	0.3	58,176	1.0
<i>Alosa</i> sp.					253	0.7	1,225	<0.1
Blueback herring					270	0.7	1,065	<0.1
American shad					46	0.1	6,133	0.1
Atlantic salmon	1	<0.1	77	<0.1	1	<0.1	195	<0.1
Rainbow trout	1	<0.1	108	<0.1				
Brown trout					11	<0.1	3,616	0.1
Brook trout	1	<0.1	135	<0.1				
Northern pike					10	<0.1	9,121	0.1
Chain pickerel	11	0.1	3,619	0.1	21	0.1	8,399	0.1
Cyprinidae	158	1.0	87	<0.1	1,605	4.1	7,505	0.1
<i>Cyprinus</i> sp.					81	0.2	382	<0.1
Common carp	309	1.9	1,229,677	44.1	311	0.8	1,171,820	19.4
Eastern silvery minnow	168	1.0	1,070	<0.1	159	0.4	1,289	<0.1
Golden shiner	1,072	6.5	35,305	1.3	992	2.6	52,159	0.9
<i>Notropis</i> sp.					3	<0.1	2	<0.1
Common shiner					69	0.2	813	<0.1
Spottail shiner	1,216	7.4	9,840	0.4	4,441	11.4	18,511	0.3
Mimic shiner					1,086	2.8	286	<0.1
Blacknose dace					2	<0.1	1	<0.1
Fallfish	3	<0.1	91	<0.1	2	<0.1	28	<0.1
Catostomidae					1,037	2.7		
Longnose sucker	2	<0.1	352	<0.1				
White sucker	1,386	8.4	478,597	17.2	3,638	9.4	1,580,832	26.2
Yellow bullhead	5	<0.1	170	<0.1	9	<0.1	1,222	<0.1
Brown bullhead	52	0.3	3,588	0.1	93	0.2	24,454	0.4
Banded killifish	752	4.6	1,501	0.1	115	0.3	297	<0.1
White perch	2,828	17.1	386,484	13.8	5,077	13.1	981,298	16.2
Centrarchidae					12	<0.1		
Rock bass	466	2.8	43,989	1.6	1,570	4.0	146,860	2.4
<i>Lepomis</i> sp.	2,067	12.5	1,524	0.1	494	1.3	1,356	<0.1
Pumpkinseed	1,765	10.7	21,632	0.8	3,372	8.7	233,561	3.9
Bluegill	375	2.3	11,242	0.4	1,298	3.3	84,849	1.4
<i>Micropterus</i> sp.					6	<0.1	304	<0.1
Smallmouth bass	583	3.5	119,821	4.3	2,229	5.7	440,864	7.3
Largemouth bass	271	1.6	11,308	0.4	846	2.2	273,048	4.5
Percidae					29	<0.1		
Tessellated darter	57	0.3	64	<0.1	134	0.3	99	<0.1
Yellow perch	2,665	16.1	277,509	9.9	9,027	23.2	659,399	10.9
Walleye	280	1.7	137,485	4.9	385	1.0	270,246	4.5
Indeterminate					3	<0.1	654	<0.1
Total	16,514		2,790,585		38,861		6,040,069	

Table 3.8 Number and weight of fish collected downstream of Vernon Dam, 1968-1989. Ichthyoplankton samples are not included.

	1968-1980				1981-1989			
	Total Number (No.)	(%)	Total Weight (g)	(%)	Total Number (No.)	(%)	Total Weight (g)	(%)
Sea lamprey					13	0.1	4,441	0.2
American eel.	34	0.3	15,084	1.1	146	1.0	42,014	1.6
Clupeidae					5	<0.1	50	<0.1
<u>Alosa</u> spp.					2	<0.1	1,194	<0.1
Blueback herring					82	0.5	4,486	0.2
American shad					1,428	9.5	81,140	3.0
Atlantic salmon					1	<0.1	162	<0.1
Brook trout					1	<0.1	129	<0.1
Rainbow trout	11	0.1	1,092	0.3				
Northern pike	1	<0.1	400	<0.1	14	0.1	22,881	0.9
Chain pickerel	29	0.2	12,389	0.9	15	0.1	5,729	0.2
Cyprinidae	3,400	25.8	2,790	0.2	431	2.9	446	<0.1
Common carp	46	0.3	228,490	16.7	47	0.3	232,973	8.7
Eastern silvery minnow	1	<0.1	16	<0.1	30	0.2	41	<0.1
Golden shiner	320	2.4	5,956	0.4	98	0.7	4,822	0.2
Lake Chub	2	<0.1	12	<0.1				
Spottail shiner	2,112	16.1	1,586	0.1	843	5.6	4,954	0.2
Golden shiner					769	5.1	1,163	<0.1
Fallfish	9	0.1	4,046	0.3	181	1.2	12,873	0.5
Catostomidae					201	1.3		
Longnose sucker	9	0.1	3,473	0.3				
White sucker	1,769	13.4	359,722	26.3	3,241	21.5	1,204,344	44.8
Yellow bullhead	3	<0.1	77	<0.1	1	<0.1	492	<0.1
Brown bullhead	133	1.0	25,316	1.9	91	0.1	34,560	1.3
Banded killifish	299	2.3	724	0.1	93	0.6	138	<0.1
White perch	832	6.3	162,917	11.9	252	1.7	65,945	2.5
Centrarchidae					10	<0.1		
Rock bass	922	7.0	125,626	9.2	2,139	14.2	292,832	10.9
<u>Lepomis</u> sp.	254	1.9	164	<0.1	404	2.7	731	<0.1
Pumpkinseed	375	2.9	25,273	1.8	492	3.3	29,847	1.1
Bluegill	467	3.6	63,349	4.6	686	4.6	96,353	3.6
<u>Micropterus</u> sp.					20	0.1	1,949	0.1
Smallmouth bass	782	5.9	132,757	9.7	1,783	11.8	347,387	12.9
Largemouth bass	68	0.5	6,840	0.5	142	0.9	18,149	0.7
Black crappie	48	0.4	6,732	0.5	1	<0.1	15	<0.1
Tessellated darter	31	0.2	29	<0.1	6	<0.1	26	<0.1
Yellow Perch	1,082	8.2	144,007	10.5	1,260	8.4	108,144	4.0
Walleye	114	0.9	37,274	2.7	116	0.8	64,629	2.4
Indeterminate					4	<0.1	38	<0.1
Total	13,153		1,366,141		15,048		2,685,079	

recruitment success of previous years. These changes were most likely natural cycles. There has not been any significant changes in the fish community species composition during Project SAVE, which were attributable to Vermont Yankee operation.

3.4.2.2 Fish Impingement

Approximately 40,000 fish have been collected in 1,560 impingement samples (Table 3.9) (Downey and Haro 1985, Briggs and Downey 1990). Seventy-five service water intake traveling screen samples obtained during 1981-1983, yielded a total of 20 fish; less than one fish per three days of sampling (Downey and Haro 1985). Due to this very low impingement rate, service water traveling screen monitoring was not required in the 1986 NPDES Permit.

Fish impingement on the circulating water traveling screens averaged about 26 fish and 320 grams per day (Briggs and Downey 1990). More than 80 percent of these fish were small sunfishes, rock bass, minnows, and yellow perch which were typically less than 100 millimeters in total length.

Fifty-nine Atlantic salmon juveniles have been collected from the circulating water traveling screen samples since 1981 (Table 3.10). Nearly all (about 95 percent) of these smolts were caught between 20 April and 15 May; a time period outside of this 316 Demonstration.

Table 3.10 Collection dates for Atlantic salmon smolts impinged on Vermont Yankee's circulating water traveling screens, 1981-1989.

Collection Date	Number of Atlantic Salmon Smolts Impinged								
	1981	1982	1983	1984	1985	1986	1987	1988	1989
20 April							1		
25 April								1	
27 April							1		
1 May								18	
2 May				1					
4 May				1					
5 May									4
7 May									13
8 May									1
9 May								2	5
10 May								4	
11 May								2	
15 May									1
20 May							2		
22 May		1							
31 May				1					
Annual Total	0	1	0	3	0	0	4	27	24

Table 3. Numbers, weights, and corresponding percentages of fish from circulating water traveling screen impingement collections, 1974-1989. Weights have been rounded to the nearest gram.

	1974-1984			1985-1989			1974-1989		
	Total Number (No.)	(%)	Total Weight (g)	Total Number (No.)	(%)	Total Weight (g)	Total Number (No.)	(%)	Total Weight (g)
Sea lamprey	5	<0.1	599	2	<0.1	8	2	<0.1	8
American eel	11	<0.1	48	3	<0.1	630	8	<0.1	1,229
Blueback herring	1	<0.1	16	1	<0.1	2	11	<0.1	48
American shad	2	<0.1	13	55	1.6	2,724	1	<0.1	16
Gizzard shad	4	<0.1	198	2	<0.1	169	2	<0.1	13
Rainbow trout	4	<0.1	289	11	0.3	55	59	0.1	2,922
Atlantic salmon	4	<0.1	289	2	<0.1	144	4	<0.1	289
Brown trout	2	<0.1	144	2	<0.1	179	4	<0.1	313
Brook trout	67	0.2	790	2	<0.1	277	78	0.2	845
Rainbow smelt	40	0.1	9,619	2	<0.1	243	42	0.1	9,896
Common carp	519	1.4	3,227	37	1.1	243	556	1.4	3,470
Eastern silvery minnow	2	<0.1	733	1	<0.1	4	1	<0.1	4
Emerald shiner	543	1.5	6,604	19	0.6	258	562	1.4	6,862
Golden shiner	9	<0.1	49	4	0.1	26	13	<0.1	75
Common shiner	12,936	35.5	56,150	240	6.9	1,000	13,176	33.0	57,150
Spottail shiner	17	<0.1	25	14	0.4	16	31	<0.1	41
Mimic shiner	2	<0.1	733	6	0.2	2	2	<0.1	733
Fallfish	3	<0.1	7	22	0.6	2,469	3	<0.1	7
Greek chub	226	0.6	1,659	6	0.2	2	232	0.6	1,661
Unidentified cyprinids	291	0.8	13,289	2	<0.1	61	313	0.8	15,758
White sucker	2	<0.1	61	12	0.3	65	2	<0.1	61
Longnose sucker	26	0.1	294	467	13.5	3,040	38	0.1	359
Yellow bullhead	673	1.8	8,767	4	0.1	12	1,140	2.9	11,807
Brown bullhead	381	1.0	1,134	73	2.1	3,336	385	1.0	1,146
Banded killifish	2,182	6.0	51,658	776	22.4	4,281	2,255	5.6	54,994
White perch	2,615	7.2	21,368	179	5.2	2,442	3,391	8.5	25,649
Rock bass	3,559	9.8	16,040	443	12.8	7,022	3,738	9.4	18,482
Pumpkinseed	1,152	3.2	8,347	589	17.0	1,292	1,595	4.0	15,369
Bluegill	6,681	18.3	15,719	61	1.8	2,169	7,270	18.2	17,011
Unidentified <i>Lepomis</i> spp.	781	2.1	25,898	33	1.0	258	842	2.1	28,067
Smallmouth bass	181	0.5	3,740	25	0.7	61	214	0.5	3,998
Largemouth bass	151	0.4	414	377	10.9	7,100	176	0.4	475
Tessellated darter	3,361	9.2	66,634	1	<0.1	5	3,738	9.4	73,734
Yellow perch	21	0.1	3,784	1	<0.1	5	21	<0.1	3,784
Walleye	3	<0.1	11	1	<0.1	5	4	<0.1	16
Slimy sculpin	33	0.1	1,622	3	<0.1	16	33	<0.1	1,622
Unidentified									
Total	36,484		318,950	3,459		38,966	39,943		357,916

Vermont Yankee has anadromous fish impingement limits placed upon their operational mode by the 1986 and previous NPDES Permits. If Atlantic salmon impingement rates exceeded 0.1 percent of the estimated smolt equivalents migrating past Vermont Yankee, the plant would have reverted to a completely closed cycle mode until 15 June. American shad impingement rate was limited to one impinged shad for each adult shad passed at the Vernon Dam fishway and/or transported by State/Federal fisheries personnel upstream of Vernon Dam. Impingement of these two anadromous species never approached nor exceeded any of these impingement limits during the nine years of Project SAVE.

Analyses of 16 years of impingement data indicated that impingement of resident and anadromous fish by Vermont Yankee had little effect on the resident fish community and the anadromous fish, Atlantic salmon and American shad.

3.4.2.3 Fish Entrainment

Ichthyoplankton enumeration began in 1977 as part of the evaluation of the Vermont Yankee's phase studies. Sampling in 1977 focused on ichthyoplankton sampling at various depths in the intake structure forebay. Larval fish collected were enumerated and ichthyoplankton densities reported by Binkerd et al. (1978) were generally less than one fish per cubic meter.

In 1982, an intensive investigation of ichthyoplankton was undertaken at the intake structure and at various locations in the river, both upstream and downstream of Vernon Dam (Binkerd et al. 1983). Ichthyoplankton were collected at various depths during 1982-1985 at the intake structure and at selected locations as part of Project SAVE. Ichthyoplankton sampling in front of the intake structure was required weekly, 1 May to 15 July, as part of the 1986 NPDES Permit. During the eight years, 1982-1989, over 1,800 samples have been collected with a 0.5 meter net, while 63 seine samples using a 500 micron mesh have been obtained (Table 3.11).

Nearly 37,000 ichthyoplankton were collected representing 14 different fish taxa (Table 3.12). Differences in the total number of ichthyoplankton captured in Vernon and Turners Falls pools were primarily the result of the significantly higher sampling effort in Vernon pool. Minnows and white perch were the most abundant ichthyoplankton, representing about 95 percent of the larvae collected. Sunfishes (*Lepomis* spp.), suckers, yellow perch, and walleye were collected consistently, but usually in low densities. Other fish, particularly smallmouth bass, were captured infrequently. The relatively low abundance of bass captured using a towed 0.5 meter net in the main portion of the river was not surprising and was most likely the result of the shallow water nesting behavior of this species.

Table 3.11 Ichthyoplankton sampling effort, number of 0.5 meter net samples, 1982-1989.

Depth (m)	Year								
	1982	1983	1984	1985	1986	1987	1988	1989	
Station 2									
<5		8	27	5					
5-10									
>10									
Stebbin Island									
<5				10	6	2	3		
5-10									
>10									
Station 3									
<5	45	28	54	10	7	3	3		
5-10	18						2		
>10	9						2		
Station 4									
<5	98	29	18	10	21	11	2		
5-10	40	12	18	10	16	6	2		
>10						2			
VY Boat Launch									
<5						6			
5-10						6			
>10						6			
VY Intake									
<5	2			1	10	44	12	11	
5-10	37		17	7	10	21	11	11	
>10	127		32	12	3		10	11	
Station 5									
<5	176	64	84	45	33	12	3		
5-10	123	36	46	43	33	12	3		
>10	96								
Other Locations									
-	58					15			
Annual Total	829	177	296	153	139	146	53	33	

Table 3.12 Relative abundance of ichthyoplankton collected near Vernon, 1982-1989. Vernon fish were obtained from 1,584 samples while Turners Falls fish represented 242 samples.

	<u>Vernon Pool</u>		<u>Turners Falls Pool</u>	
	<u>(No.)</u>	<u>(%)</u>	<u>(No.)</u>	<u>(%)</u>
Clupeidae	2	<0.1	1	<0.1
American shad	15	<0.1	2	0.1
Blueback herring	7	<0.1		
Cyprinidae	4,018	12.1	288	8.5
Common carp	41	0.1	25	0.7
Golden shiner	20	<0.1	3	0.1
<u>Notropis</u> spp.	15,020	45.2	1,828	53.8
Catostomidae	87	0.3	7	0.2
White sucker	541	1.6	23	0.7
White perch	12,348	37.2	1,057	31.1
Centrarchidae	7	<0.1	1	<0.1
Rock bass	1	<0.1		
<u>Lepomis</u> spp.	434	1.3	120	3.5
Smallmouth bass	2	<0.1		
Percidae	11	<0.1		
Tessellated darter	13	<0.1	1	<0.1
Yellow perch	339	1.0	28	0.8
Walleye	225	0.7	9	0.3
Indeterminate	95	0.3	5	0.1
Total	33,226		3,398	

A distinct pattern of seasonal succession of ichthyoplankton was apparent during these studies. The yellow perch were usually the first larvae to be encountered in early to mid-May (Table 3.13). Walleye larvae were consistently captured during mid-May, 15 May to 28 May, while white perch were collected during mid-May to early-July. Minnows, usually collected beginning in early-June, were the principal component of ichthyoplankton during late-June and July.

Table 3.13 Earliest and latest dates of capture of ichthyoplankton for four game fish species, 1982-1989.

<u>Year</u>	<u>Date of Capture</u>							
	<u>Smallmouth Bass</u>		<u>White Perch</u>		<u>Walleye</u>		<u>Yellow Perch</u>	
	<u>Earliest</u>	<u>Latest</u>	<u>Earliest</u>	<u>Latest</u>	<u>Earliest</u>	<u>Latest</u>	<u>Earliest</u>	<u>Latest</u>
1982	26 Jun	30 Jun	17 May	21 Jul	17 May	09 Jun	11 May	10 Jun
1983	22 Jun	23 Jun	20 May	06 Jul	16 May	23 May	12 May	02 Jun
1984			18 May	11 Jul	14 May	08 Jun	07 May	08 Jun
1985			16 May	11 Jul	14 May	21 May	03 May	21 May
1986			14 May	08 Jul	14 May	20 May	14 May	29 May
1987			14 May	09 Jun			04 May	26 May
1988			16 May	13 Jul			10 May	23 Jun
1989			23 May	14 Jul			12 May	19 May

Yellow perch and walleye larvae were only collected during a three week interval, indicating that spawning occurred during a relatively narrow time frame. White perch ichthyoplankton, on the other hand, appeared in samples from mid-May through July, with peak densities observed in late-May. These observations indicated that the white perch did not spawn all at once, but continued to spawn throughout the spring and early-summer.

Ichthyoplankton densities were typically less than one fish per cubic meter, although densities of minnows during late-June and July of three to five fish per cubic meter were occasionally observed. Near the shore in shallower regions of Vernon pool, densities of ichthyoplankton, specifically minnows collected by seining, have been estimated as high as 3,000 fish per cubic meter (Downey 1990d).

Overall, density of ichthyoplankton were found not to be dependent upon sampling depth, but a marked depth preference among the different species was observed. Minnows were generally more abundant in surface (less than 5 feet) samples, while white perch were most numerous in deeper (greater than 5 feet) samples (Downey 1990d).

Densities of the two most abundant fish taxa, minnows and white perch, near the Vermont Yankee intake structure were comparable to other sampling locations. Walleye were also collected at low densities in and near the intake. Walleye densities were very low and more than 90 percent of the larval walleye were captured annually during 15 May to 28 May (Downey 1990d). Densities of walleye appear to be highest in the mid-depth strata while surface and near-bottom samples contained a lower abundance of larval walleye.

The operation of Vermont Yankee did result in fish being entrained in the circulating water cooling system. Densities of the abundant taxa were relatively low in intake water and were comparable to densities observed in river samples. Shallow shoreline areas were more productive and contained much higher densities of the abundant taxa. Because of the low densities in the intake samples observed during the eight year sampling program, the effect of entrainment on walleye and other game species, such as smallmouth bass, was minimal. If ichthyoplankton entrainment resulted in an adverse effect on the fish community, these effects would have been noticeable as either major changes in the fish community and/or decreases in the relative abundance of the Representative Important Species. As is demonstrated in Section 3.4.2.1, Community Composition, and Section 3.4.2.4, Representative Important Species, no changes in the fisheries in the Connecticut River near Vernon were documented as a result of Project SAVE operation. The evaluation of ichthyoplankton near Vernon indicates that the entrainment of ichthyoplankton through Vermont Yankee's circulating water cooling system did not result in appreciable harm to the fish community during Project SAVE.

3.4.2.4 Representative Important Species

Based on a consensus of the VYEAC, seven species of fish were selected to be Representative Important Species: Atlantic salmon, American shad, smallmouth bass, white perch, walleye, yellow perch, and spottail shiner. The objective of this section is to discuss, when applicable, the distribution of the species in the river near Vernon, migration of the species relative to the thermal plume areas, and growth.

ATLANTIC SALMON

Atlantic salmon, an anadromous fish, are presently undergoing extensive restoration efforts in the Connecticut River. The Atlantic salmon were historically residents of the Connecticut River and its tributaries, but with the construction of Turners Falls Dam in 1798 and dams at Vernon, Vermont and Holyoke, Massachusetts, the salmon were extirpated. The recent construction of fishways at dams on the Connecticut River and the hatchery stocking program have resulted in Atlantic salmon returning to the Connecticut River drainage.

Atlantic salmon spawn in the gravelly reaches of tributaries and smaller streams in the fall. There the eggs overwinter and hatch into young salmon, termed parr. These young salmon remain in the fresh-water for one to two years before they undergo smoltification; a physiological change preparing the fish for life in salt water. Once smoltification begins in April and May, the salmon migrate into the main stem of the Connecticut River and then move downstream into the Atlantic Ocean. The salmon will usually spend two winters at sea before returning as adults to the same stream in the spring, May through July, to spawn. Unlike the Pacific salmon, Atlantic salmon do not necessarily die after spawning, and can spawn again in following years.

Since this species is presently being restored, the production of parr and smolts is totally dependent upon hatchery operations. Most of the fry and smolts are stocked in the West and White Rivers in Vermont and in the Ammoonusic River in New Hampshire. Generally, several hundred thousand parr and smolts are released upstream of Vermont Yankee in these three rivers. When a wild population of salmon becomes established in the Connecticut River, spawning will most likely occur in these tributaries, and others, which contain suitable habitat for spawning and parr survival. Spawning in the main stem of the Connecticut River is unlikely due to lack of available habitat and their spawning behavior. The interaction between Vermont Yankee operation and Atlantic salmon will occur during the times of migration through the Vernon region, i.e., upstream migration of adults and downstream migration of smolts.

During Project SAVE, the returns of adult Atlantic salmon to the Connecticut River have been low, with a maximum annual return of 529 salmon in 1981 (Rideout and McLaughlin 1985). Many of these returning

State and

adult Atlantic salmon are trapped at Holyoke Dam and retained for brood stock in the federal hatchery system. Since 1985, about ten percent of the adult Atlantic salmon have passed through Holyoke Dam fishway. A total of 59 Atlantic salmon adults have been passed through the Turners Falls Dam fishway gatehouse between 1981-1990. Forty-four of these salmon were passed through Vernon Dam fishway (Table 3.14), resulting in a 75 percent passage efficiency.

Table 3.14 Annual upstream passage of smallmouth bass, Atlantic salmon, American shad and walleye at the Vernon Dam fishway. These data were obtained from Cox (1990) and unpublished data (K. Cox, Vermont Fish and Wildlife; personal communication).

Year	Number of Fish			
	Smallmouth Bass	American Shad	Atlantic Salmon	Walleye
1981	2,529	97	8	498
1982	1,542	9	0	363
1983	467	2,597	0	211
1984	262	335	0	56
1985	181	833	4	83
1986	703	982	4	71
1987	660	3,459	13	96
1988	281	1,370	5	72
1989	466	2,953	0	20
1990	440	10,868	10	89

Behavioral delays of these salmon were not observed during Vermont Yankee Project SAVE operation (Table 3.15). During many of the documented Atlantic salmon passages at Vernon Dam fishway, Vermont Yankee was operating in a hybrid or open cycle mode as part of Project SAVE (Figures 3.1, 3.2, and 3.3).

The behavior of outmigrating smolts in the vicinity of Vermont Yankee's thermal plume was investigated in 1980 and 1981 (McAvoy 1980, 1981) and again in 1988 and 1989 (Downey et al. 1990b). These studies used radiotelemetry to monitor juvenile Atlantic salmon movement through and past regions affected by Vermont Yankee's thermal plume. With temperature modulated transmitters used in the latter study, the movement of smolts into the thermal plume and downstream to Vernon Dam without delay was documented. The conclusion by Downey et al. (1990b) was that fully smoltified Atlantic salmon move through and downstream of Vermont Yankee's thermal plume without any appreciable avoidance behavior. These observations of adult and Atlantic salmon smolt migration patterns relative to Vermont Yankee operation indicate that blockage of Atlantic salmon migration by the thermal plume did not occur.




Figure 3.1 River discharge, fishway temperature, and Vermont Yankee operation during passage of adult Atlantic Salmon in the Vernon Dam fishway, 1987.




Figure 3.2 River discharge, fishway temperature, and Vermont Yankee operation during passage of adult Atlantic Salmon in the Vernon Dam fishway, 1988.




Figure 3.3 River discharge, fishway temperature, and Vermont Yankee operation during passage of adult Atlantic Salmon in the Vernon Dam fishway, 1990.

Table 3.15 Adult Atlantic salmon observed at the Vernon Dam fishway and Vermont Yankee operation, 1985-1989. Atlantic salmon passage times and dates were obtained from Cox (1990) and unpublished data (K. Cox; Department of Fish and Wildlife, personal communication).

<u>Time of First Observation</u>	<u>VY Operation (% of open cycle)</u>	<u>Time of First Observation</u>	<u>VY Operation (% of open cycle)</u>
<u>1985</u>		<u>1988</u>	
1230h 27 May	75	1045h 29 May	40
1115h 29 May	<5	0820h 01 Jun	46
		0825h 01 Jun	47
		0750h 14 Jun	5
<u>1986</u>		1750h 14 Jun	29
1810h 20 May	Not Operating		
1308h 22 May	Not Operating		
1345h 23 May	Not Operating	<u>1990</u>	
1025h 01 Jul	Not Operating	0900h 08 Jun	42
		0900h 09 Jun	78
<u>1987</u>		1000h 09 Jun	86
1705h 19 May	72	0800h 10 Jun	79
1208h 20 May	32	1400h 12 Jun	100
0740h 27 May	20	1600h 12 Jun	100
1306h 27 May	11	0900h 14 Jun	100
0800h 31 May	<5	0700h 18 Jun	<5
1700h 01 Jun	37	0700h 18 Jun	<5
0845h 02 Jun	38	1900h 18 Jun	50
0725h 03 Jun	14		
0950h 05 Jun	43		
1030h 05 Jun	43		
1945h 05 Jun	48		
0800h 14 Jun	38		

AMERICAN SHAD

American shad, an anadromous fish, are also part of the Anadromous Fish Restoration Program. Management efforts to restore these fish to their native Connecticut River range began in the late 1950's and the shad passage at the Holyoke Dam fishway in the 1980's have ranged from 275,000 to 530,000 per year.

The American shad begin ascending the Connecticut River in April and continue through early-July. Males usually enter the river first. Spawning takes place near the water's surface with eggs and sperm released in open water. The fertilized eggs, which are slightly heavier than water, then settle to the bottom.

Larval shad emerge from the egg at about nine to ten millimeters (mm) and spend the summer in freshwater. During the late-summer and fall, the juveniles migrate downstream. American shad will spend

three to four years in the ocean before returning to their native river to spawn. From 1955-1986, between 10 and 65 percent of the Connecticut River adult American shad returned as repeat spawners (Crecco and Savoy 1987).

Connecticut River shad migration, the number of fish passed at fishways, declines steadily as the run progressed upstream. Passage at the Turners Falls fishway was only about one to seven percent of Holyoke Dam fishway passage (Table 3.16). Passage at Vernon Dam fishway ranged from about 8 to 40 percent of Turners Falls fishway passage between 1983-1990. The percent passage efficiencies of Vernon Dam in 1981 and 1982 were even higher, but low numbers of American shad were passed at Turners Falls Dam. The reduction in fish progressing upstream was, in part, the result of American shad maturing and spawning in the river between fishways.

Table 3.16 Adult American shad passage at three Connecticut River fishways, 1980-1990. Percent efficiency (% Eff) relative to the immediate downstream facility passage is also expressed. Data were obtained from Kynard and O'Leary (1989a, 1989b), Cox (1990), Meyers (1990) unpublished data (Cox, Vermont Department of Fish and Wildlife personal communication), and S. Amaral, Massachusetts Cooperative Fishery Research Unit).

<u>Year</u>	<u>Holyoke</u>	<u>Turners Falls</u>		<u>Vernon Dam</u>	
	<u>Fishway</u>	<u>Fishway</u>		<u>Fishway</u>	
	<u>(No.)</u>	<u>(No.)</u>	<u>(% Eff)</u>	<u>(No.)</u>	<u>(% Eff)</u>
1980	380,000	298	<0.1	Not Constructed	
1981	380,000	200	<0.1	97	48.5
1982	290,000	11	<0.1	9	81.8
1983	530,000	12,705	2.4	2,597	20.4
1984	500,000	4,333	0.9	335	7.7
1985	480,000	3,855	0.8	833	21.6
1986	350,000	17,858	5.1	982	5.5
1987	280,000	18,959	6.8	3,459	18.2
1988	290,000	15,787	5.4	1,370	8.7
1989	354,000	9,511	2.7	2,953	31.0
1990	363,788	27,908	7.7	10,868	38.9

The passage at the upstream fishways, Turners Falls and Vernon, began in mid- to late-May when the fishways are first operated. The runs at these two fishways typically peaked between 25 May and 10 June, but low numbers of shad continue to pass up to the closing of the fishways in early-July (Downey 1990f).

The passage efficiency of the Vernon Dam fishway was compared with the increased energy production by Vermont Yankee during Project SAVE, 1981-1990. The American shad passage efficiency at Vernon Dam

fishway was high during years of high thermal discharge (1989 and 1990) and during 1986 when Vermont Yankee was not operating. No correlation between Vernon Dam fishway shad passage efficiency and Vermont Yankee Project SAVE operation was observed during this study.

Sex ratios of the adult American shad population was heavily skewed in favor of males at the Vernon Dam (Staats 1990). The male component of the Vernon Dam fishway population was estimated at 82 percent which was comparable to the sex ratio of the American shad population sampled in the river below Vernon Dam. Males also represented about 71 percent of Turners Falls fishway population (Amaral 1990). Predominance of males at upstream locations was typical of many American shad populations in other rivers of North America (Table 3.17). The hypotheses for these sex ratio differences in migration include the higher energetic cost to the female in upstream migration, behavioral differences, and/or size (sex) selectivity of the fishways.

The young-of-the-year American shad abundance and growth in the upper Turners Falls pool have been monitored since 1984 (Downey 1985a, 1985e, 1987, 1988a, 1990a, and 1990b). The growth rates observed during these studies range from 0.4mm per day to 1mm per day which were comparable with the growth rates of young-of-the-year American shad in the lower Connecticut River (Crecco et al. 1981). Changes in growth rates of young American shad were not observed as a result of Project SAVE.

Reproductive success of the American shad in the Vernon pool was documented in 1982 and 1983 by the capture of young-of-the-year American shad (Downey 1990f). These two years of juvenile shad recruitment coincided with the trucking of adult American shad by State and Federal personnel from the Holyoke Dam fishway to a release point in the Vernon pool, several miles upstream of Vermont Yankee. The reasons for the lack of documented spawning success by adult American shad in Vernon pool in subsequent years are not known. The low proportion of female to male American shad and their sexual maturity when passed into the Vernon pool may be factors. The lower Vernon pool region, between Vermont Yankee's discharge and the dam, does not contain appreciable spawning habitat for American shad. Since significant recruitment of juvenile shad occurs each year immediately downstream of Vernon Dam, it is unlikely that Vermont Yankee operation was a significant factor for the lack of spawning success in Vernon pool.

SMALLMOUTH BASS

Smallmouth bass are an important game fish in the Connecticut River. Although the smallmouth bass are not native to the Connecticut River (Scarola 1973), these fish are now common near Vernon. The smallmouth bass represented about four percent of the fish captured upstream of Vernon Dam and about 11 percent downstream.

This species typically inhabits the same area during its life. In recapture investigations of smallmouth bass in other drainages, individuals were usually recaptured within one-half to three miles of

Table 3.17 Sex composition of adult American shad populations from three drainages. Data were obtained from Chittenden (1969), RMC (1988, 1989, 1990), Moffitt (1979), Sherer (1975), Henry and Kaska (1984), Henry and Krska (1984), Amaral (1990), Cox (1990) and Staats (1990).

	<u>Year</u>	<u>Sample Size</u>	<u>Composition (%)</u>	
			<u>Male</u>	<u>Female</u>
Delaware River	1961	198	86	14
	1962	220	99	1
	1963	302	62	38
	1964	199	38	62
	1965	23	52	48
Susquahanna River, Conowingo Dam	1987	5,304	76	24
	1988	3,375	74	26
	1989	4,077	77	23
Farmington River, Rainbow Dam	1976	916	73	27
	1977	297	78	22
Connecticut River, Holyoke Dam	1970	1,611	75	25
	1971	672	66	34
	1972	3,808	40	60
	1973	3,652	20	80
	1982	837	63	37
	1986	-	50	50
	1988	-	69	31
	1989	680	46	54
Connecticut River, Turners Falls Dam	1983	-	91	9
	1986	75	60	40
	1988	51	78	22
	1989	46	71	29
Connecticut River, Vernon Dam	1988	43	98	2
	1989	176	82	18

their original location (Scott and Crossman 1973). Tag and recapture studies conducted on this species near Vernon confirmed this behavior (Binkerd et al. 1990a). Movement upstream and downstream through the Vernon Dam fishway has been documented (Table 3.14).

The distribution of smallmouth bass appeared to be unaffected by the presence of the thermal plume. These fish were routinely collected at Station 4 along the Vermont shore, an area where the thermal plume was located. Although this species resides at Station 4, studies by Binkerd et al. (1978) have demonstrated that the thermal plume did not attract these fish.

Downstream of Vernon Dam, smallmouth bass were an important component of the fish community. This upper Turners Falls pool region, which was less than one mile downstream of Vermont Yankee, receives thermal effluent when Vermont Yankee was operating. The presence of a viable smallmouth bass population in the thermal plume region, and downstream, indicated that smallmouth bass are not excluded from any area as a result of Vermont Yankee's thermal discharge.

The general health, parasitic burden, and overall condition of the smallmouth bass near Vernon were comparable to smallmouth bass collected in Bellows Falls pool (Downey 1984a).

Growth of juvenile and adult smallmouth bass have been monitored for 22 years (Downey 1985b, 1990c). Smallmouth bass near Vernon grew to a size of about 90 to 100mm the first year (Downey 1990c). Back-calculated lengths at each scale annulus were computed and grouped according to location of capture, and three time periods (Figure 3.4). Annual growth, defined as the mean length at annulus formation, was similar for all time periods and locations, with one exception. The calculated size of smallmouth bass at age one, for bass captured in the two locations affected by Vermont Yankee's thermal discharge, and at the one location upstream of Vermont Yankee, was statistically smaller for 1981-1989 when compared with corresponding estimates for the 1968-1973 and 1974-1980 periods. The mean difference was six to eight millimeters. Comparison of mean length at age one of bass captured in the two locations receiving thermal discharge and the upstream location not affected by thermal discharge were essentially the same for the 1981-1989 age one calculations. This observation suggests that the apparent smaller size at age one was not directly related to thermal discharges. Lengths at age one for Connecticut River smallmouth bass were above the average of other back-calculated lengths at age one for smallmouth bass populations from other drainages in North America (Downey 1990c).

WHITE PERCH

White perch are found along the Atlantic coast in North America from the St. Lawrence River to South Carolina. These fish are usually found in brackish water but can survive as freshwater landlocked populations. White perch thrive in a variety of habitats but seem to be more successful in warmer waters that reach 75°F or more in the summer (Scott and Crossman 1973).





Figure 3.4 Mean back calculated total length in millimeters for Smallmouth Bass, 1969-1989.

The white perch population in the Connecticut River near Vernon is not native. Fish surveys conducted in 1939 did not collect any of these fish, but in 1968 this species was a component of the fish community (Aquatec et seqq. 1972).

Sexually mature adults have been captured in late-May through June near Vernon. White perch larvae have been collected in ichthyoplankton samples from 14 May to 21 July, although the peak densities of white perch collected in the river occur during the end of May through early-June (Binkerd et al. 1983). This pattern of extended larval recruitment, presumably indicative of recent spawning, indicated that the Connecticut River white perch population spawned over a six to eight week period. This protracted spawning period has been observed in other white perch populations. Mansueti (1961) noted that only a portion of the total number of eggs are released at one time and eggs ripen progressively over time being released in two or three separate spawning acts.

In ichthyoplankton samples collected near Vernon, white perch accounted for half of the larval fish captured (Downey 1990d). White perch larvae were rarely caught in surface water samples but represent a major component of ichthyoplankton captured at a depth of five feet or more. White perch have been collected at all stations sampled in the Turners Falls and Vernon pools, including at the Vermont Yankee intake. White perch densities typically average less than one fish per cubic meter.

White perch are a large component of the fish community in the Connecticut River near Vernon, particularly in the lower Vernon pool. In this pool, white perch represented about 5 to 25 percent of the fish captured annually (Downey 1990e). The white perch represented about five to eight percent of the population during 1968-1980 in the Vernon pool. Several strong year classes resulted in the white perch becoming a dominant component in the early 1980's, peaking at over 25 percent of the fish captured in 1982-1983. The white perch component of the fish community declined over subsequent years resulting in the relative composition of the white perch returning to previous levels. Long-term population fluctuations in white perch are not uncommon and have been reported in other populations.

In the upper Turners Falls pool, white perch were numerically less dominant. The relative composition of white perch in this pool ranged from five percent during 1968-1980 to about two percent in 1981-1989.

The growth of white perch has been consistent throughout the three regions studied (Figure 3.5). Differences of growth during Project SAVE, 1981-1989, were not evident in any region when compared to time periods of 1965-1973, and 1974-1980. The mean size at annulus of white perch from the three Connecticut River regions was comparable to other white perch populations in North America (Downey 1985c, 1990c).





Figure 3.5 Mean back calculated total length in millimeters for White Perch, 1969-1989.

These evaluations of white perch distribution and growth demonstrate that exclusion of habitat, or changes in growth, did not occur during Project SAVE operation.

WALLEYE

The walleye are an important game fish in the Connecticut River near Vernon. This species is not indigenous to the Connecticut River, but was presumably introduced from Lake Champlain (Bailey 1938). The walleye represented about one-half to two percent of the fish captured in the Turners Falls and Vernon pools. Walleye have been captured consistently at Station 4 near the Vermont shore. Individuals of this species were collected throughout the summer and early fall at Station 4 during closed cycle operation and at times when Vermont Yankee was operating in an open or hybrid cycle during Project SAVE (Binkerd et al. 1983). Walleye have been documented moving through the Vernon Dam fishway in the spring while heat was being discharged to Vernon pool (Table 3.14).

Walleye spawning in the Connecticut River support what has been reported for other populations (Scott and Crossman 1973). In the ichthyoplankton samples collected between 1982-1989, walleye larvae were only obtained in a relatively short time period; primarily from 14 May to 23 May. If a 10 to 15 day egg maturation and larval hatching process is assumed, this would indicate that most of the walleye spawning near Vernon takes place in late-April to the first week of May.

Although walleye larvae have been collected throughout the Vernon and Turners Falls pools, walleye spawning areas have not been documented. Station 4, being composed of predominately silty substrates, contains little, if any, suitable habitat for walleye spawning.

Growth of walleye in the various regions of the Connecticut River near Vernon have shown a statistically non-significant but noticeable trend of increased growth in regions affected by thermal discharge during the 22 years of these studies (Figure 3.6) (Downey 1984b, Downey 1990c). The comparison of mean size at annulus among the three regions of capture during the same time intervals did not reveal any trends. Overall, mean size at annulus was above the average reported for other walleye populations in the northeast (Downey 1990c). The analyses of walleye distribution and growth during 1981-1989 indicate that Project SAVE operation did not result in an adverse effect on this species.

YELLOW PERCH

Yellow perch have a circumpolar distribution in fresh water in the northern hemisphere. This species is very adaptable and can utilize a wide variety of warm to cooler habitats. They are most abundant in open water lakes with moderate vegetation and softer sand/mud bottoms.





Figure 3.6 Mean back calculated total length in millimeters for Walleye, 1969-1989.

In the Vernon pool, yellow perch were a major component of the fish community. Yellow perch represented about 16 percent of the community during 1968-1980 and about 23 percent of the community during 1981-1989. These fish were found throughout the lower Vernon pool including those locations affected by thermal effluent. In the upper Turners Falls pool, the relative abundance of yellow perch has remained relatively constant, about eight percent, for the two time periods.

Spawning of yellow perch typically occurs mid-April through early-May. In ichthyoplankton samples, yellow perch larvae were most often obtained between 3 May and 26 May, with larvae/small juveniles captured as late as 10 June. The yellow perch ichthyoplankton represented less than one percent of the ichthyoplankton collected annually.

Parasitic burdens of yellow perch collected near Vernon included digenetic trematodes, monogenetic trematodes, fish leeches, and nematodes encysted in the visceral mesenteries which were comparable to infestations observed in yellow perch located in Bellows Falls pool (Downey 1984a). One striking difference observed was the heavy infestation of yellow grub (Clinostomum marginatum) in the yellow perch captured in the Bellows Falls pool, a 100 percent incidence, whereas the yellow grub was not isolated in yellow perch captured near Vernon. Since that study (Downey 1984a), yellow perch captured near Vernon were routinely examined and only a few fish have been recorded with light yellow grub infestation.

Yellow perch growth has been generally comparable among fish captured in three locations during 1969-1989 (Figure 3.7). Statistically significant lower growth rates of 1974-1980 cohorts were observed for yellow perch captured in upper Vernon pool; an area outside the influence of the thermal plume (Downey 1985d, 1990c). Significant differences in the growth of fish captured among the three locations were not observed during Project SAVE, 1981-1989. The analysis of the growth and distribution of yellow perch near Vernon did not reveal any significant effect of Project SAVE operation on this species.

SPOTTAIL SHINER

The spottail shiner occurs in parts of Canada south into the United States to Georgia in the east and Iowa and Missouri in the west (Scott and Crossman 1973). This species is generally regarded as a fish of relatively large lakes and rivers.

Spottail shiners were reported to be the most abundant fish in the Connecticut River near Vernon (Binkerd et al. 1978). This species continued to be abundant in the Connecticut River near Vernon. In Vernon pool, the spottail shiner accounted for eight percent of the fish captured during 1968-1980 and 11 percent of the fish captured (exclusive of 500 micrometer seine samples) during 1981-1989. In upper Turners Falls pool, about 16 percent of the fish captured during 1968-1980 were spottails while they represented about five percent of the fish captured during 1981-1989.





Figure 3.7 Mean back calculated total length in millimeters for Yellow Perch, 1969-1989.

Minnow larvae first appeared in late-May and were the dominant ichthyoplankton during June through August. The minnow larvae (Notropis spp., Hybognathus regius, and Notemigonus crysoleucas) were not known well enough for positive differentiation. These ichthyoplankton were found in the largest densities in surface samples, less than five feet deep.

At least three age groups of spottail shiners were collected in the Vernon and Turners Falls pools during 1968-1989. Observed sizes were consistent with reported size at annulus formation for spottail shiners (Scott and Crossman 1978).

The presence and abundance of spottail shiners in Vernon and Turners Falls pools demonstrate that no significant reduction in growth and distribution of spottail shiner have resulted from Project SAVE operation.

4.0 ENGINEERING AND HYDROLOGICAL INFORMATION

4.1 Introduction

This section describes the thermal effects on the Connecticut River due to the discharge of heat from Vermont Yankee during the period 16 May to 14 October. The thermal effects on the Connecticut River are described in two regions: in Vernon Pool where thermal plumes were observed, and downstream of Vernon Dam where the heated water was well mixed with ambient river water.

4.2 Vermont Yankee Facilities and Operation

Vermont Yankee facilities are essentially unchanged since 1978. The location of gates are the same as previously described by Binkerd et al. (1978); however, gear-driven gate operators on four gates, one intake gate, two cooling tower bypass gates, and the recirculation gate, have been replaced with hydraulic gate operators to withstand frequent cycling. A temperature data acquisition system was installed in 1984 as a replacement to a system first installed in 1976 (King 1985). The system was modified and now includes pH recording and monitoring of Vernon Dam fishway temperatures (King 1990).

Heat has been discharged from Vermont Yankee to the Connecticut River since 1982 between 16 May and 14 October without exceeding temperature standards permitted for Project SAVE. Plant operation within the test temperature standards has been achieved in part due to a computer simulation (Binkerd et al. 1983). In 1981 and early in 1982, the computer simulation of plant operation was developed based on plant design information independent of actual operational data.

The simulator was verified through extensive physical experiments in the spring and summer of 1982. The following year, the computerized version of the simulator was used to develop a manual for plant operators (Johnston and Binkerd 1984). Several revisions to this manual have been written and provided to the Vermont Yankee operators. Revision 4, dated May 1990, is currently being used for the control of heat discharged to the river.

4.3 Vernon Dam Facilities and Operation

The Vernon Dam and Vernon Station were described in Binkerd et al. (1978). Many changes have been made by the New England Power Company to the dam and hydroelectric station since 1978. A fishway was constructed and first operated in 1981. Also, the wooden flashboards along the crest of the dam were increased from six feet high to eight feet high in 1984. Most of these wooden flashboards were replaced in 1987 with hydraulically operated gates. Presently, there are nine gates along the crest of the dam and only two sections of wooden flashboards remain.

Since the installation of these gates, the typical pool elevation has increased by about two feet; pool elevations of over 219.5 feet

above mean sea level (msl) are now common. The hydraulically operated gates have also reduced the frequency with which the pool water elevation is reduced to dam crest elevation (about 212 feet msl). Hydraulically operated gates allow more control on the location of spillage over the dam's crest, which affects the shape of the thermal plume in Vernon Pool.

4.4 Effect of Discharged Heat on Connecticut River Temperatures

The heat discharged by Vermont Yankee results in thermal plumes in Vernon pool and increased mixed river temperatures downstream. A summary of temperature increases at Station 3 was produced for the years 1968-1983 (Johnston 1984). An update of Johnston (1984) by Luxenberg (1990a) summarized temperature increases at Station 3 for 1984-1989.

Temperature gradients were quite apparent in Vernon pool, typically between the discharge structure and Vernon Dam. At low river discharges, thermal plumes extended upstream of the discharge structure. Thermal plumes measured from May to October, 1989, were displayed in Binkerd et al. (1990b). Some differences were observed in these plumes compared with the plumes measured by Binkerd et al. (1978) during the late fall, winter, and early spring. In general, thermal plumes observed in the summer had smaller areas of higher excess temperatures, and larger areas of lower excess temperatures; but the shape and characteristics of the 1989 summer plumes were similar to previously measured thermal plumes in Vernon pool. These differences were most likely due to increased pool elevation, selective hydraulic gate operation, and higher ambient river temperatures.

Temperature patterns near the Vernon Dam fishway during high river discharge indicated that exit fishway temperatures were often equal to or less than Station 3 temperatures during high river discharge (Binkerd 1985a). During lower river discharge, however, exit fishway temperatures were often 1 to 2°F higher than Station 3 temperatures and differences as much as 3 to 4°F have occurred.

The heated water became well mixed with ambient water by Station 3 as a result of Vernon Dam and Station operation. As this heated water was transported downstream to Long Island Sound, temperature increases were reduced due to longitudinal mixing and transfer of excess heat to the atmosphere. The temperature increase near Holyoke Dam (58 miles downstream) could have been 0.1 to 1.5°F for initial increases of up to 5°F at Station 3 during the period 16 May to 14 October (Luxenberg 1985).

4.5 Thermal Impact Assessment

Temperature increases resulting from Vermont Yankee operation under the proposed alternate temperature standards are limited by the river discharge (discharged based) or the temperature standard (criteria based). Criteria based temperature increases were

determined by hourly upstream Station 7 temperatures (Figure 4.1) and proposed alternate temperature standards. Discharged based temperature increases were determined by available river discharge records from 1973-1989 (Figure 4.2) and the assumption that Vermont Yankee discharged all of its excess heat. The minimum of these two values, either criteria based or discharged based temperature increases, formed a data set of temperature increases that would have occurred had Vermont Yankee operated using the proposed alternate temperature standards. Hourly probabilities of creating a river temperature increase were calculated using this data set. An example of these calculations is illustrated in Table 4.1. Interpolated probability values for the occurrence of 0.5, 1.5, 2, 3, 4, and 5°F temperature increases are plotted in Figure 4.3.

According to this analysis, in May when river discharge is high, the maximum proposed temperature standard (5°F) will not be reached. Also, there is less than a five percent chance of achieving a 4°F increase during the mid-May period. During the summer (July-August) there is a 60 to 80 percent chance that a 2°F increase will occur. In this mid-summer period, the proposed temperature based standard of 2°F increase will dominate. In October, 5°F increases will occur about ten percent of the time.

The mean hourly temperature at Station 7 was subtracted from the 20 year maximum measured temperature and plotted in Figure 4.4 along with the standard deviation of temperature calculated from Station 7 data. The simulated maximum temperature increase and the simulated increase not exceeded 68.3 percent of the time are also plotted in Figure 4.4 for comparison.

The historical mean and maximum temperatures are plotted along with temperatures measured at Stations 3 and 7 in 1989 (Figure 4.5). During 1989, the plant was discharging heat to the river the entire time under Project SAVE. Station 3 temperatures were usually warmer than Station 7 temperatures during 1989. Both Stations 3 and 7 temperatures fluctuated frequently about the mean sometimes by 5°F or greater. The historical daily minimum temperature occurred in mid-June, 1989, followed by historical daily maximum temperature in late-June, indicating the wide fluctuations in ambient temperatures that can occur in a very short time period.

This analysis illustrates that expected temperature increases will be typically less than the historical standard deviation from mid-May to late-September. During late-September to mid-October, expected temperature increases will exceed the historical standard deviation of the mean ambient temperature. It is anticipated that river temperature increases will not always be at the level of proposed temperature standards, but, in many instances, will be much lower due to expected high river discharges.

Table 4.1 Example of data on 18 May, 0800 hours, statistics, and calculations for thermal impact assessment.

Year	18 May at 0800 hours		Temperature Increases, °F				Rank of Column 3	Percent Probability of Temperature Increase in Column 3
	Station 7	Connecticut River	1#	2@	3	4\$		
	Temperature °F	Discharge cfs						
1970	54.0		5.0	0.9	0.9	12	67	
1971	51.2	17,920	4.0	0.7	0.7	16	89	
1972	51.5	23,260	3.0	1.4	1.4	9	50	
1973	52.5	11,690	4.0	1.0	1.0	10	56	
1974	55.2	16,240	3.0	2.0	2.0	4	22	
1975	59.0	7,860	5.0	0.7	0.7	17	94	
1976	57.2	24,310	3.0	1.7	1.7	6	33	
1977	59.0	9,430	4.0	11.6	4.0	1	6	
1978	50.2	1,380	4.0	0.9	0.9	11	61	
1979	62.1	17,030	4.0	1.6	1.6	7	39	
1980	55.4	10,030	5.0	0.8	0.8	15	83	
1981	55.6	20,420	5.0	0.8	0.8	14	78	
1982	57.6	20,340	3.0	10.8	3.0	2	11	
1983	50.0	1,490	3.0	11.7	3.0	3	17	
1984	47.9	1,370	3.0	2.0	2.0	5	28	
1985	59.3	8,150	3.0	1.4	1.4	8	44	
1986	60.9	11,180	4.0	0.8	0.8	13	72	
1987	60.0	19,720						
1988	60.2							
1989	55.9							
Maximum	62.1 ^{A,D,E}	24,310 ^B			4.0 ^D			
Mean	55.7 ^{A,D,E}	13,048 ^B						
Minimum	47.9 ^A	1,370 ^B						
Standard Deviation	4.1 ^{A,D}							

1.8^{D,!}}

Use 2-5 °F criteria
 @ Use 16040/discharge
 \$ Highest increase has rank = 1,
 * Lowest increase has rank =17
 - 100 x rank/18
 ! Temperature increase with 68.3 percent probability of being equal to or less than actual increase

A Statistics used in Figure 4.1
 B Statistics used in Figure 4.2
 C Statistics used in Figure 4.3
 (Interpolate to get probabilities at specific temperature increases)
 D Statistics used in Figure 4.4
 (Interpolate to get temperature increases at specific probability)
 E Statistics used in Figure 4.5




Figure 4.1 Range of Station 7 temperatures, 1970-1989.




Figure 4.2 Range of Connecticut River discharge at Vernon, Vermont, 1973-1989.

Percent probability of equivalence or exceedance




Figure 4.3 Distribution of simulated temperature increase using 2 to 5 °F temperature standard.




Figure 4.4 Comparison of temperature differences with simulated increases.

5.0 LITERATURE CITED

- Aquatec, Inc. 1972 et seqq. Ecological studies of the Connecticut River, Vernon, Vermont. Reports I through XIX, January 1972 through December 1989. Aquatec, Inc., South Burlington, Vermont.
- Bailey, R.M. 1938. The fishes of the Merrimack watershed. Pages 149-185 in Biological survey of the Merrimack watershed. New Hampshire Fish and Game Department, Concord.
- Binkerd, Roger C. 1985a. Temperature patterns near Vernon Dam fish passage during high river discharge. Aquatec, Inc., South Burlington, Vermont.
- Binkerd, Roger C., William D. Countryman, R. Mason McNeer and Daniel J. Marx. 1978. 316 demonstration engineering, hydrological and biological information and environmental impact assessment. Aquatec, Inc., South Burlington, Vermont.
- Binkerd, Roger C., H. Gregory Johnston, Philip C. Downey, Stephen C. McAvoy, and Hudson R. DeYoe. 1983. Operational and biological studies of the Vermont Yankee/Connecticut River system. Aquatec, Inc., South Burlington, Vermont.
- Binkerd, Roger C., Michael T. Hewlett, and Philip C. Downey. 1990a. Tag and recapture studies of smallmouth bass (Micropterus dolomieu Lacepede), 1981-1989. Aquatec Inc., South Burlington, Vermont.
- Binkerd, Roger C., Roland R. Luxenberg, and Stephen P. Farrington. 1990b. Thermal plumes in the lower Vernon pool, 1989. Aquatec Inc., South Burlington, Vermont.
- Briggs, Errol C., and Philip C. Downey. 1990. Fish impingement on intake screens at Vermont Yankee, 1985-1989. Aquatec Inc., South Burlington, Vermont.
- Chittenden, Mark E., Jr. 1969. Life history and ecology of the American shad, *Alosa sapidissima*, in the Delaware River. Doctoral dissertation. Rutgers University, New Brunswick, New Jersey.
- Cox, Kenneth. 1990. Job Performance Report job I-4 Vernon Fishway Project F-12-R-22. Department of Fish and Game, Roxbury, Vermont.
- Crecco, Victor A., and Thomas Savoy, 1987. Fisheries management plan for American shad in the Connecticut River. Connecticut Bureau of Fisheries, Waterford.
- Crecco, Victor A., Linda Gunn, and Thomas Savoy. 1981. Connecticut River shad study, 1980. Connecticut Bureau of Fisheries, Waterford.

- Downey, Philip, C. 1984a. Notes on the health of fishes of the Connecticut River near Vernon, Vermont. Aquatec, Inc., South Burlington, Vermont.
- Downey, Philip C. 1984b. Age and growth of walleye (Stizostedion vitreum vitreum (Mitchill)) of the Connecticut River near Vernon, Vermont. Aquatec Inc., South Burlington, Vermont.
- Downey, Philip C. 1985a. Growth of 1984 juvenile American shad (Alosa sapidissima (Wilson)) of the Connecticut River near Vernon, Vermont. Aquatec Inc., South Burlington, Vermont.
- Downey, Philip C. 1985b. Age and growth of smallmouth bass (Micropterus dolomieu Lacepede) of the Connecticut River near Vernon, Vermont. Aquatec Inc., South Burlington, Vermont.
- Downey, Philip C. 1985c. Age and growth of white perch (Morone americana (Gmelin)) of the Connecticut River near Vernon, Vermont. Aquatec Inc., South Burlington, Vermont.
- Downey, Philip C. 1985d. Age and growth of yellow perch (Perca flavescens (Mitchill)) of the Connecticut River near Vernon, Vermont. Aquatec Inc., South Burlington, Vermont.
- Downey, Philip C. 1985e. Growth of 1985 juvenile American shad (Alosa sapidissima (Wilson)) of the Connecticut River near Vernon, Vermont. Aquatec Inc., South Burlington, Vermont.
- Downey, Philip C. 1987. Spatial distribution of 1986 juvenile shad (Alosa sapidissima (Wilson)) of the Connecticut River near Vernon, Vermont. Aquatec Inc., South Burlington, Vermont.
- Downey, Philip C. 1988a. Age and growth of 1986 juvenile American shad (Alosa sapidissima (Wilson)) of the Connecticut River near Vernon, Vermont. Aquatec Inc., South Burlington, Vermont.
- Downey, Philip C. 1990a. Microhabitats of juvenile American shad (Alosa sapidissima (Wilson)) in the Connecticut River near Vernon, Vermont. Aquatec Inc., South Burlington, Vermont.
- Downey, Philip C. 1990b. Age and growth of juvenile American shad (Alosa sapidissima (Wilson)) of the Connecticut River near Vernon, Vermont. Aquatec Inc., South Burlington, Vermont.
- Downey, Philip C. 1990c. Age and growth of selected resident fish of the Connecticut River near Vernon, Vermont. Aquatec Inc., South Burlington, Vermont.
- Downey, Philip C. 1990d. Abundance, density and composition of ichthyoplankton of the Connecticut River near Vernon, Vermont. Aquatec Inc., South Burlington, Vermont.

- Downey, Philip C. 1990e. Composition of the fish community of the Connecticut River Near Vernon, Vermont. Aquatec Inc., South Burlington, Vermont.
- Downey, Philip C. 1990f. Adult American shad (Alosa sapidissima (Wilson)) migration in the Connecticut River near Vernon, Vermont. Aquatec Inc., South Burlington, Vermont.
- Downey, Philip C., and Alexander J. Haro. 1985. Fish impingement on intake screens at Vermont Yankee, 1974-1984. Aquatec Inc., South Burlington, Vermont.
- Downey, Philip C., Nickolas R. Staats, and Roger C. Binkerd. 1990b. Downstream movement of Atlantic salmon (Salmo salar Linnaeus) smolts in Vernon pool. Aquatec Inc., South Burlington, Vermont.
- Henry, S. and R. J. Krska. 1984. Project Performance Report Job 2 - Turners Falls fish passage facility of Project F-45-R-2. Massachusetts Cooperative Fishery Research Unit, Amherst.
- Johnston, H. Gregory. 1984. Thermal experience of the Connecticut River near Vernon, Vermont. Aquatec, Inc., South Burlington, Vermont.
- Johnston, H. Gregory, and Roger C. Binkerd. 1984. Determination of optimal settings of condenser cooling system facilities. Aquatec Inc., South Burlington, Vermont.
- King, David E. 1985. Vermont Yankee environmental temperature system. Aquatec Inc., South Burlington, Vermont.
- King, David E. 1990. The Vermont Yankee environment data acquisition systems; an update. Aquatec Inc., South Burlington, Vermont.
- Kynard, Boyd, and John O'Leary. 1989a. Job Performance Report Job 1-Holyoke fish passage facility of Project F-45-R-8. Massachusetts Cooperative Fishery Research Unit, Amherst.
- Kynard, Boyd, and John O'Leary. 1989b. Job Performance Report Job 2-Turners Falls fish passage facility of Project F-45-R-8. Massachusetts Cooperative Fishery Research Unit, Amherst.
- Luxenberg, Roland R. 1985. Connecticut River temperature increase. Aquatec Inc., South Burlington, Vermont.
- Luxenberg, Roland R. 1987. Temporal and spatial distribution of water quality parameters in upper Turners Falls pool. Aquatec Inc., South Burlington, Vermont.
- Luxenberg, Roland R. 1990a. Thermal history of the Connecticut River, 1984-1989. Aquatec Inc., South Burlington, Vermont.

- Luxenberg, Roland R. 1990b. Analysis of the thermal history of the Connecticut River, 1970-1989. Aquatec, Inc., South Burlington, Vermont.
- Mansueti, R. J. 1961. Movements, reproduction and mortality of the white perch, Roccus americanus, in the Patuxent estuary, Maryland. Chesapeake Science 2(3-4): 142-205.
- McAvoy, Stephen C. 1980. A study of Atlantic salmon smolt migration in the Connecticut River near Vermont Yankee Nuclear Power Station, spring 1980. Aquatec, Inc., South Burlington, Vermont.
- McAvoy, Stephen C. 1981. A study of Atlantic salmon smolt migration in the Connecticut River near Vermont Yankee Nuclear Power Station, spring 1981. Aquatec, Inc., South Burlington, Vermont.
- Meyers, Theodore. 1990. 1990 salmon and shad returns [letter to the Connecticut River Anadromous Fish Commission dated July 9, 1990].
- Moffitt, Christine M. 1979. Recolonization of American shad, Alosa sapidissima (Wilson), above Rainbow Dam fish ladder, Farmington River, Connecticut. Doctoral dissertation. University of Massachusetts, Amherst.
- Park, Janice P. 1990. Acquisition of biological data and their translation to a computer database. Aquatec Inc., South Burlington, Vermont.
- Rideout, Stephen G., and Everett A. McLaughlin. 1985. Progress Report Connecticut River Anadromous Fish Restoration Program January 1, 1977 - December 31, 1982. U.S. Fish and Wildlife Service, Hadley, Massachusetts.
- RMC Environmental Services, Inc. 1988. Job I. Summary of Conowingo Dam fish passage facility operation in spring 1989 in Susquehanna River Anadromous Fish Restoration Committee, Richard St. Pierre, Coordinator. Restoration of American shad to the Susquehanna River annual report 1989. Harrisburg, Pennsylvania.
- RMC Environmental Services, Inc. 1989. Job VI. Summary of Conowingo Dam fish passage facility operation in spring 1989 in Susquehanna River Anadromous Fish Restoration Committee, Richard St. Pierre, Coordinator. Restoration of American shad to the Susquehanna River annual report 1989. Harrisburg, Pennsylvania.
- RMC Environmental Services, Inc. 1990. Job I. Summary of Conowingo Dam Fish passage facility operation in spring 1989 in Susquehanna River Anadromous Fish Restoration Committee, Richard St. Pierre, Coordinator. Restoration of American shad to the Susquehanna River annual report 1989. Harrisburg, Pennsylvania.

- Robins, C.R., R.M. Baily, Carl E. Bond, James R. Booker, Ernest A. Lachner, Robert N. Lea, and W.B. Scott. 1980. A list of common and scientific names of fishes from the United States and Canada (Fourth Edition). American Fisheries Society Special Publication No. 12, Bethesda, Maryland.
- Scarola, J.F. 1973. Freshwater fishes of New Hampshire. New Hampshire Fish and Game Department, Concord.
- Scott, W.B., and E.J. Crossman. 1973. Freshwater fishes of Canada. Fisheries Research Board of Canada, Bulletin 184, Ottawa.
- Shambaugh, Angela d. 1987. Temporal and spatial distribution of phytoplankton in upper Turners Falls pool. Aquatec Inc., South Burlington, Vermont.
- Shambaugh, Angela d., and Philip C. Downey. 1990. The occurrence of Leptodora kindti in the Connecticut River near Vernon, Vermont during early summer 1988. Aquatec Inc., South Burlington, Vermont.
- Staats, Nickolas R. 1990. Age and sex composition of adult American shad (Alosa sapidissima (Wilson)) at Vernon Dam fishway, 1989. Aquatec Inc., South Burlington, Vermont.
- Timmons, Maria J. 1988. Temporal and spatial distribution of zooplankton in upper Turners Falls pool. Aquatec Inc., South Burlington, Vermont.
- Wood, Susan M. 1988. Temporal and spatial distribution of macroinvertebrates in upper Turners Falls pool. Aquatec Inc., South Burlington, Vermont.
- U.S. Environmental Protection Agency. 1977a. Interagency 316(a) technical guidance manual and guide for thermal effects sections of nuclear facilities. Environmental impact statements (draft). U.S. Environmental Protection Agency, Office of Water Enforcement, Permits Division, Industrial Permits Brank, Washington, D.C.
- U.S. Environmental Protection Agency. 1977b. Guidance for evaluating the adverse impact of cooling water intake structures on the aquatic environment: Section 316(b) P.L. 92-500 (draft). U.S. Environmental Protection Agency, Office of Water Enforcement, Permits Division, Industrial Permits Brank, Washington, D.C.

90043D31MAY90