


**FIRSTENERGY NUCLEAR OPERATING COMPANY  
BEAVER VALLEY POWER STATION**



**2007 ANNUAL ENVIRONMENTAL OPERATING REPORT  
NON-RADIOLOGICAL  
UNITS NO. 1 AND 2  
LICENSES DPR-66 AND NPF-73**

**BEAVER VALLEY POWER STATION  
ENVIRONMENTAL & CHEMISTRY SECTION**


**Technical Report Approval:**


**2007 ANNUAL NON-RADIOLOGICAL ENVIRONMENTAL OPERATING REPORT**

**UNITS NO. 1 AND 2**

**LICENSES DPR-66 AND NPF-73**

Prepared by: Cameron L. Lange (Via E-mail) Date: 04/16/08

Prepared by: Michael D. Banko III  Date: 04/17/08

Reviewed by: Julie A. Firestone  Date: 04/17/08

Approved by: Donald J. Salera  Date: 4-17-08

# TABLE OF CONTENTS

	<u>Page</u>
<b>1.0 EXECUTIVE SUMMARY .....</b>	<b>1</b>
1.1 INTRODUCTION.....	1
1.2 SUMMARY & CONCLUSIONS .....	1
1.3 ANALYSIS OF SIGNIFICANT ENVIRONMENTAL CHANGE.....	2
1.4 AQUATIC MONITORING PROGRAM EXECUTIVE SUMMARY .....	2
<b>2.0 ENVIRONMENTAL PROTECTION PLAN NON-COMPLIANCES .....</b>	<b>4</b>
<b>3.0 CHANGES INVOLVING UNREVIEWED ENVIRONMENTAL QUESTIONS ...</b>	<b>4</b>
<b>4.0 NONROUTINE ENVIRONMENTAL REPORTS.....</b>	<b>4</b>
<b>5.0 AQUATIC MONITORING PROGRAM.....</b>	<b>5</b>
5.1 SITE DESCRIPTION .....	5
5.2 STUDY AREA .....	6
5.3 METHODS .....	6
5.3.1 <u>Benthic Macroinvertebrate Monitoring</u> .....	6
5.3.2 <u>Fish Monitoring</u> .....	7
5.3.3 <u>Corbicula/Zebra Mussel Density Determinations</u> .....	8
5.3.4 <u>Corbicula Juvenile Monitoring</u> .....	9
5.3.5 <u>Zebra Mussel Monitoring</u> .....	10
5.3.6 <u>Reports</u> .....	11
5.4 RESULTS OF THE AQUATIC MONITORING PROGRAM .....	11
5.4.1 <u>Benthic Macroinvertebrate Monitoring Program</u> .....	11
5.4.2 <u>Fish Sampling Program</u> .....	14
5.4.3 <u>Corbicula Monitoring Program</u> .....	16
5.4.4 <u>Corbicula Juvenile Monitoring</u> .....	17
5.2.4 <u>Zebra Mussel Monitoring Program</u> .....	18
<b>6.0 ZEBRA MUSSEL AND CORBICULA CONTROL ACTIVITIES.....</b>	<b>19</b>
<b>7.0 REFERENCES.....</b>	<b>20</b>
<b>8.0 TABLES</b>	
<b>9.0 FIGURES</b>	
<b>10.0 PERMITS</b>	

## APPENDIX A SCIENTIFIC AND COMMON NAME OF FISH COLLECTED IN THE NEW CUMBERLAND POOL OF THE OHIO RIVER, 1970 THROUGH 2007 BVPS

## LIST OF TABLES

- 5.1 Beaver Valley Power Station (BVPS)  
Sampling Dates For 2007
- 5.2 Systematic List of Macroinvertebrates Collected From 1973 through 2007 in the Ohio River near BVPS (6 sheets)
- 5.3 Benthic Macroinvertebrate Counts for Triplicate Samples Taken at Each Sample Station by Sample for May and September 2007
- 5.4 Mean Number of Macroinvertebrates (Number/m<sup>2</sup>) and Percent Composition of Oligochaeta, Chironomidae, Mollusca and Other Organisms, 2007 - BVPS
- 5.5 Mean Number of Macroinvertebrates (Number/m<sup>2</sup>) and Percent Composition of Oligochaeta, Chironomidae, Mollusca and Other Organisms for the Control Station (1) and the Average for Non-control Stations (2B1, 2B2, and 2B3), 2006 BVPS
- 5.6 Shannon-Weiner Diversity, Evenness and Richness Indices for Benthic Macroinvertebrates Collected in the Ohio River, 2007
- 5.7 Benthic Macroinvertebrate Densities (Number/m<sup>2</sup>) for Station 1 (Control) and Station 2B (Non-Control) During Preoperational and Operational Years through 2007 BVPS
- 5.8 Total Fish Catch, Electrofishing and Seine Net Combined During the BVPS 2007 Fisheries Survey
- 5.9 Comparison of Control vs. Non-Control Electrofishing Catches, During the BVPS 2007 Fisheries Survey
- 5.10 Comparison of Control vs. Non-Control Seine Catches, During the BVPS 2007 Fisheries Survey
- 5.11 Fish Species Collected During the May 2007 Sampling of the Ohio River in the Vicinity of BVPS
- 5.12 Fish Species Collected During the July 2007 Sampling of the Ohio River in the Vicinity of BVPS
- 5.13 Fish Species Collected During the September 2007 Sampling of the Ohio River in the Vicinity of BVPS
- 5.14 Fish Species Collected During the November 2007 Sampling of the Ohio River in the Vicinity of BVPS
- 5.15 Estimated Number of Fish Observed During Electrofishing Operations

## LIST OF TABLES

- 5.16 Catch Per Unit of Effort (CPUE as Fish/Electrofishing Minute) by Season During the BVPS 2004 Fisheries Survey
- 5.17 Catch Per Unit of Effort (CPUE as Fish/Electrofishing Minute) by Season During the BVPS 2005 Fisheries Survey
- 5.18 Catch Per Unit of Effort (CPUE as Fish/Electrofishing Minute) by Season During the BVPS 2006 Fisheries Survey
- 5.19 Catch Per Unit of Effort (CPUE as Fish/Electrofishing Minute) by Season During the BVPS 2007 Fisheries Survey
- 5.20 Unit 1 Cooling Reservoir Monthly Sampling *Corbicula* Density Data for 2007 from BVPS
- 5.21 Unit 2 Cooling Reservoir Monthly Sampling *Corbicula* Density Data for 2007 from BVPS

## LIST OF FIGURES

- 5.1 Location Map for the 2007 Beaver Valley Power Station Aquatic Monitoring Program Sampling Control and Non-Control Sampling Stations
- 5.2 Location Map for Beaver Valley Power Station Benthic Organism Survey Sampling Sites for the 2007 Study
- 5.3 Location Map for Beaver Valley Power Station Fish Population Survey Fish Sampling Sites for the 2007 Study
- 5.4 Location of Study Area, Beaver Valley Power Station Shippingport, Pennsylvania BVPS
- 5.5 Comparison of Live *Corbicula* Clam Density Estimates Among BVPS Unit 1 Cooling Tower Reservoir Sample Events for Various Clam Shell Size Groups, 2007.
- 5.6 Comparison of Live *Corbicula* Clam Density Estimates Among Unit 2 Cooling Tower Reservoir Sample Events for Various Clam Shell Size Groups, 2007.
- 5.7 Comparison of Live *Corbicula* Clam Density Estimates Among Intake Structure Sample Events for Various Clam Shell Size Groups, 2007.
- 5.8 Water Temperature and River Elevation Recorded on the Ohio River at the BVPS Intake Structure, During Monthly Sampling Dates, 2007.
- 5.9 Density of Zebra Mussel Veligers ( $\#/m^3$ ) Collected at Beaver Valley Power Station, Intake Structure, Unit 1 Cooling Tower Reservoir and Unit 2 Cooling Tower Reservoir, 2007.
- 5.10 Density of Zebra Mussel Veligers ( $\#/m^3$ ) Collected at Beaver Valley Power Station, Barge Slip, Splash Pool and Emergency Outfall Basin, 2007.
- 5.11 Density ( $\#/m^2$ ) of Settled Zebra Mussels at Beaver Valley Power Station Intake Structure, Unit 1 Cooling Tower Reservoir and Unit 2 Cooling Tower Reservoir, 2007.
- 5.12 Density ( $\#/m^2$ ) of Settled Zebra Mussels at Beaver Valley Power Station, Barge Slip, Splash Pool and Emergency Outfall Basin, 2007.

## **1.0 EXECUTIVE SUMMARY**

### **1.1 INTRODUCTION**

This report is submitted in accordance with Section 5.4.1 of Appendix B To Facility Operating License No. NPF-73, Beaver Valley Power Station Unit 2, Environmental Protection Plan (Non-Radiological). Beaver Valley Power Station (BVPS) is operated by FirstEnergy Nuclear Operating Company (FENOC). The Objectives of the Environmental Protection Plan (EPP) are:

- Verify that the facility is operated in an environmentally acceptable manner, as established by the Final Environmental Statement-Operating License Stage (FES-OL) and other NRC environmental impact assessments.
- Coordinate NRC requirements and maintain consistency with other Federal, State, and local requirements for environmental protection.
- Keep NRC informed of the environmental effects of facility construction and operation and of actions taken to control those effects.

To achieve the objectives of the EPP, FirstEnergy Corporation, FENOC, and BVPS, have written programs and procedures to comply with the EPP, protect the environment, and comply with governmental requirements primarily including the US Environmental Protection Agency (EPA), and the Pennsylvania Department of Environmental Protection (PA DEP) requirements. Water quality matters identified in the Final Environmental Statements-Operating License Stage (FES-OL) are regulated under the National Pollutants Discharge Elimination System (NPDES) Permit No. PA0025615. Waste is regulated under EPA Identification No. PAR000040485. Attachment 10.1 contains a listing of permits and certificates for environmental compliance.

The BVPS programs and procedures include pre-work and pre-project environmental evaluations, operating procedures, pollution prevention and response programs procedures and plans, process improvement and corrective action programs, and human performance programs. Technical and managerial monitoring of tasks, operations, and other activities are performed. Any identified challenges, concerns, or questions are captured in the FENOC Problem Identification and Resolution Program with a Condition Report. Condition Reports include investigations, cause determinations, and corrective actions.

During 2007 BVPS continued an Aquatic Monitoring Program to evaluate its potential impact on the New Cumberland Pool of the Ohio River, and to provide information on potential impacts to BVPS operation from macrofoulers such as Asian clams and Zebra mussels.

### **1.2 SUMMARY AND CONCLUSIONS**

There were no significant environmental events during 2007.

During 2007, no significant changes to operations that could affect the environment were made at

Beaver Valley Power Station. *As in previous years, results of the BVPS environmental programs did not indicate any adverse environmental impacts from station operation.*

### 1.3 ANALYSIS OF SIGNIFICANT ENVIRONMENTAL CHANGE

During 2007, no significant changes were made at BVPS to cause significant negative affect on the environment.

### 1.4 AQUATIC MONITORING PROGRAM

The 2007 Beaver Valley Power Station (BVPS) Units 1 and 2 Non-Radiological Monitoring Program consisted of an Aquatic Program that included surveillance and field sampling of the Ohio River's aquatic life in the vicinity of the station. The Aquatic Program is an annual program conducted to provide baseline aquatic resources data, to assess the impact of the operation of BVPS on the aquatic ecosystem of the Ohio River, and to monitor for potential impacts of biofouling organisms (*Corbicula* and zebra mussels) on BVPS operations. This is the 32<sup>nd</sup> year of operational environmental monitoring for Unit 1 and the 21<sup>st</sup> year for Unit 2. As in previous years, the results of the program did not indicate any adverse environmental impact to the aquatic life in the Ohio River associated with the operation of BVPS.

The results of the 2007 benthic macroinvertebrate survey conducted in May and September indicated a normal community structure exists in the Ohio River both upstream and downstream of the BVPS. These benthic surveys are also a continuation of a Fate and Effects Study conducted from 1990 through 1992 for the Pennsylvania Department of Environmental Protection (PADEP) to assess the ecosystem impacts of the molluscicides Betz Clamtrol CT-1, CT-2, and Powerline 3627 that are used to control biofouling organisms at BVPS. To date the results of the benthic studies have not indicated any impacts of operation at the BVPS including the use these biocides on the benthic community below the BVPS discharge.

Substrate was probably the most important factor influencing the distribution and abundance of the benthic macroinvertebrates in the Ohio River near BVPS. The generally soft muck-type substrate along the shoreline found in 2007 and previous years was conducive to segmented worm (oligochaete) and midge (chironomid) proliferation. Forty-two (42) macroinvertebrate taxa were identified during the 2007 monitoring program. One new taxon was added to the cumulative taxa list of macroinvertebrates collected near BVPS. This was the chironomid (midge fly), *Stempellina* sp. No state or Federal threatened or endangered macroinvertebrate species were collected during 2007. In May and September, oligochaetes were the most frequently collected group of macroinvertebrate. *There were no major differences in the community structure between control and non-control stations that could be attributed to operation of BVPS. The overall community structure has changed little since pre-operational years, and program results did not indicate that BVPS operations were affecting the benthic community of the Ohio River.*



The fish community of the Ohio River near the BVPS was sampled in May, July, September and November of 2007 with nighttime electrofishing and daytime seining. Since monitoring began in the early 1970's, the number of identified fish taxa has increased from 43 to 78 for the New Cumberland Pool.

Benthivores (bottom feeders including suckers and buffalo) were generally collected in the highest numbers in 2007, although smallmouth bass was the most frequently collected species. As in 2006, forage species, although common, were collected in lower numbers than in previous years. Variations in annual catch were probably attributable to normal fluctuations in the population size of the forage species and the predator populations that rely on them. Forage species, such as gizzard shad and emerald shiner with high reproductive potentials, frequently respond to changes in natural environmental factors (competition, food availability, cover, and water quality) with large fluctuations in population size, which may be the reason for the lower numbers of these species observed in 2006 and 2007. This, in turn, influences their appearance in the sample populations during annual surveys. Spawning/rearing success due to abiotic factors is usually the determining factor of the size and composition of a fish community.

In 2007, the annual catch rate for electrofishing (number of fish per minute of effort, or CPUE) was 1.17 fish per minute. The greatest catch rate in 2007 occurred in November (winter) (1.63 fish/electrofishing minute). Sauger, smallmouth bass, and shorthead redhorse suckers contributed the most to this total. The lowest catch rate occurred in September (fall) with a rate of 0.85 fish/electrofishing minute. The annual catch rate in 2007 was consistent with the previous two years ranging from a high of 1.60 fish per minute in 2006 to 1.16 in 2005.

Little difference in the species composition of the catch was observed between the control (Station 1) and non-control (Stations 2A, 2B and 3) stations. Habitat preference and availability were probably the most important factors affecting where and when fish were collected. ***Results from the 2007 fish surveys indicated that a normal community structure for the Ohio River exists near BVPS based on species composition and relative abundance. In 2007, there was no indication of negative impact to the fish community in the Ohio River from the operation of BVPS.***

The monthly reservoir ponar samples collected in Units 1 and 2 cooling towers and the four samples collected at the intake during 2007 indicated that *Corbicula* were entering and colonizing the station. ***Overall, the numbers of Corbicula collected in the samples were comparatively low, which continued the trend over the past few years of fewer *Corbicula* and reflected a water-body-wide trend observed in the Ohio River.***

In 1995, live zebra mussels were collected for the first time by divers in the BVPS main intake and auxiliary intake structures during scheduled cleanings. Overall, both the number of observations and densities of settled mussels were similar during 2004-2007. The density of veligers in 2007 was much lower than in 2006 but was comparable to 2005 and 2004. ***Although densities of settled mussels in the vicinity of BVPS are low compared to other populations such as in the Lower Great Lakes, densities comparable to those in the Ohio River are sufficient to cause problems in the operation of untreated cooling water intake systems.***

## **2.0 ENVIRONMENTAL PROTECTION PLAN NON-COMPLIANCES**

There were no Environmental Protection Plan non-compliances identified in 2007.

## **3.0 CHANGES INVOLVING UNREVIEWED ENVIRONMENTAL QUESTIONS**

No Unreviewed Environmental Questions were identified in 2007. Therefore, there were no changes involving an Unreviewed Environmental Question.

## **4.0 NON-ROUTINE ENVIRONMENTAL REPORT**

A letter report was submitted to the Pennsylvania Department of Environmental Protection (PA DEP) On June 11, 2007. It was a follow-up written report to describe a condition initially reported to the PA DEP via telephone on June 11, 2007.

A corrugated pipe ruptured that channels water discharged from our Unit 2 Emergency Outfall Structure (EOF), NPDES Permit No. PA0025615 Outfall 010, into an Impact Basin (basin is to reduce velocity and prevent erosion). That rupture caused the discharging water to create a sinkhole, washing out soil and fill material that slid down the hill towards the Ohio River. Most of the washed-out material appears to have been captured on a flat area between the outfall and the river, and in the Impact Basin itself. Some material may have made it past the Impact Basin to the river. Note that the materials washed out were sandy and rocky, typical of the fill used during construction of BVPS. There was no further communication with the PA DEP regarding this matter.

There were no other non-routine environmental reports in 2007.

## 5.0 AQUATIC MONITORING PROGRAM

This section of the report summarizes the Non-Radiological Environmental Program conducted for the BVPS 1 Units 1 and 2; Operating License Numbers DPR-66 and NPF-73. This is a non-mandatory program, because on February 26, 1980, the Nuclear Regulatory Commission (NRC) granted BVPS's request to delete all of the Aquatic Monitoring Program, with the exception of the fish impingement program (Amendment No. 25), from the Environmental Technical Specifications (ETS). In 1983, BVPS was permitted to also delete the fish impingement studies from the ETS program of required sampling along with non-radiological water quality requirements. However, in the interest of providing an uninterrupted database, BVPS has continued the Aquatic Monitoring Program.

The objectives of the 2007 environmental program were:

- To monitor for any possible environmental impact of BVPS operation on the benthic macroinvertebrate and fish communities in the Ohio River;
- To provide a low level sampling program to continue an uninterrupted environmental database for the Ohio River near BVPS, pre-operational to present; and
- To evaluate the presence, growth, and reproduction of macrofouling *Corbicula* (Asiatic clam) and zebra mussels (*Dreissena* spp.) at BVPS.

## 5.1 SITE DESCRIPTION

BVPS is located on an approximately 453-acre tract of land on the south bank of the Ohio River in the Borough of Shippingport, Beaver County, Pennsylvania. The Shippingport Atomic Power Station once shared the site with BVPS before being decommissioned. Figure 5.1 is a plan view of BVPS. The site is approximately 1 mile (1.6 km) from Midland, Pennsylvania; 5 miles (8 km) from East Liverpool, Ohio; and 25 miles (40 km) from Pittsburgh, Pennsylvania. The population within a 5-mile (8 km) radius of the plant is approximately 18,000. The Borough of Midland, Pennsylvania has a population of approximately 3,500.

The station is situated at Ohio River Mile 34.8 (Latitude: 40° 36' 18"; Longitude: 80° 26' 02") at a location on the New Cumberland Pool that is 3.1 river miles (5.3 km) downstream from Montgomery Lock and Dam and 19.6 miles (31.2 km) upstream from New Cumberland Lock and Dam. The Pennsylvania-Ohio-West Virginia border is 5.2 river miles (8.4 km) downstream from the site. The river flow is regulated by a series of dams and reservoirs on the Beaver, Allegheny, Monongahela, and Ohio Rivers and their tributaries.

The study site lies along the Ohio River in a valley, which has a gradual slope that extends from the river (Elevation 665 ft (203 m) above mean sea level) to an elevation of 1,160 ft (354 m) along a ridge south of BVPS. The plant entrance elevation at the station is approximately 735 ft (224 m) above mean sea level.

BVPS Units 1 and 2 have a thermal rating of 2,660 megawatts (MW). Units 1 & 2 have a design electrical rating of 835 MW and 836 MW, respectively. The circulating water systems for each unit are considered a closed cycle system with continuous overflow, using a cooling tower to minimize heat released to the Ohio River. Commercial operation of BVPS Unit 1 began in 1976 and Unit 2 began operation in 1987.

## 5.2 STUDY AREA

The environmental study area, established to assess potential impacts, consists of four sampling stations each having a north and south shore (Figure 5.1). Station 1 is located at River Mile (RM) 34.5, approximately 0.3 miles (0.5 km) upstream of BVPS and is the control station. Station 2A is located approximately 0.5 miles (0.8 km) downstream of the BVPS discharge structure in the main channel. Station 2B is located in the back channel of Phillis Island, also 0.5 miles downstream of the BVPS discharge structure. Station 2B is the principal non-control station because the majority of discharges from BVPS Units 1 and 2 are released to this back channel. Station 3 is located approximately two miles (3.2 km) downstream of BVPS

## 5.3 METHODS

Shaw Environmental, Inc. (Shaw) was contracted to perform the 2007 Aquatic Monitoring Program as specified in BVBP-ENV-001-Aquatic Monitoring (procedural guide). This procedural guide references and describes in detail the field and laboratory procedures used in the various monitoring programs, as well as the data analysis and reporting requirements. These procedures are summarized according to task in the following subsections. Sampling was conducted according to the schedule presented in Table 5.1.

### 5.3.1 Benthic Macroinvertebrate Monitoring

The benthic macroinvertebrate monitoring program consisted of river bottom sampling using a Ponar grab sampler at four stations on the Ohio River. Prior to 1996, duplicate sampling occurred at Stations 1, 2A, and 3, while triplicate sampling occurred at Station 2B (i.e., one sample at each shoreline and mid-channel) (Figures 5.1 and 5.2). In 1996, a review of the sampling design indicated that sampling should be performed in triplicate at each station to conform to standardized U.S. Environmental Protection Agency (USEPA) procedures. Therefore, starting in 1996, triplicate samples were taken at Stations 1, 2A, and 3, as in 1995, with triplicate samples also collected at each shore and mid-channel location at Station 2B. A petite Ponar dredge was used to collect these samples, replacing the standard Ponar dredge used in prior studies. This sampling was conducted in May and September 2007. For each 2007 field effort, 18 benthic samples were collected and processed in the laboratory. All field procedures and data analyses were conducted in accordance with the procedural guide.

The contents of each Ponar grab sample were gently washed through a U.S. Standard No. 30 sieve and the retained contents were placed in a labeled bottle and preserved in ethanol. In the laboratory, rose bengal stain was added to aid in sorting and identifying the benthic organisms. Macroinvertebrates were sorted from each sample, identified to the lowest taxon practical and counted. Mean density (number/m<sup>2</sup>) for each taxon was calculated for each replicate. Four indices used to describe the benthic community were calculated: Shannon-Weiner diversity index, evenness (Pielou, 1969), species richness, and the number of taxa. These estimates provide an indication of the relative quality of the macroinvertebrate community.

### 5.3.2 Fish Monitoring

Fish sampling was conducted in 2007 to provide a continuous baseline of data and to detect possible changes that may have occurred in the fish populations in the Ohio River near BVPS. Fish population surveys have been conducted in the Ohio River near BVPS annually from 1970 through 2007. These surveys have resulted in the collection of 73 fish species and five different hybrids.

Adult fish surveys were scheduled and performed in May, July, September, and November 2007. During each survey, fish were sampled by standardized electrofishing techniques at four stations (Stations 1, 2A, 2B and 3) (Figure 5.3). Seining was performed at Station 1 (north shore) and Station 2B (south shore of Phillis Island) to sample species that are generally under-represented in electrofishing catches (e.g., young-of-the-year fish and small cyprinids).

Night electrofishing was conducted using a boat-mounted electroshocker with floodlights attached to the bow. A Smith-Root Type VI A variable voltage, pulsed-DC electrofishing unit powered by a 5-kW generator was used. The voltage selected depended on water conductivity and was adjusted to provide a constant amperage (4-6 amps) of the current through the water. The north and south shoreline areas at each station were shocked for at least 10 minutes of unit "on" time (approximately five minutes along each shore) during each survey.

When large schools of fish of a single non-game species such as gizzard shad and shiners were encountered during electrofishing efforts, all of the stunned fish were not netted and retrieved onboard the boat. A few fish were netted for verification of identity, and the number of observed stunned fish remaining in the water was estimated. The size range of the individual fish in the school was also estimated and recorded. This was done in an effort to expedite sample processing and cover a larger area during the timed electrofishing run. Regardless of the number of individuals, all game fish were boated when observed.

Fish seining was performed at Station 1 (control) and Station 2B (non-control) during each scheduled 2007 BVPS fishery survey. A 30-ft long bag seine made of 1/4-inch nylon mesh netting was used to collect fish located close to shore in 1 to 4 ft of water. Three seine hauls were performed at both Station 1 (north shore) and Station 2B (south shore of Phillis Island) during each survey.

Fish collected during electrofishing and seining efforts were processed according to standardized procedures. All captured game fishes were identified, counted, measured for total length (nearest 1 mm), and weighed (nearest 1 g for fish less than or equal to 1000 g and the nearest 5 g for all other fish). Non-game fishes were counted, and a random subsample of lengths was taken. Live fish were returned to the river immediately after processing was completed. All fish that were unidentifiable or of questionable identification and were obviously not on the endangered or threatened species list were placed in plastic sample bottles, preserved, labeled and returned to the laboratory for identification. Any species of fish that had not previously been collected at BVPS was retained for the voucher collection. Any threatened or endangered species (if collected) would be photographed and released.

### 5.3.3 Corbicula Density Determinations for Cooling Tower Reservoirs

The *Corbicula* Monitoring Program at BVPS includes sampling the circulating river water and the service water systems of the BVPS (intake structure and cooling towers). This report describes this Monitoring Program and the results of the field and plant surveys conducted in 2007.

The objectives of the ongoing Monitoring Program are to evaluate the presence of *Corbicula* at BVPS, and to evaluate the potential for and timing of infestation of the BVPS. This program is conducted in conjunction with a program to monitor for the presence of macrofouling zebra mussels (see Section 5.3.5).

*Corbicula* enter the BVPS from the Ohio River by passing through the water intakes, and eventually settle in low flow areas including the lower reservoirs of the Units 1 and 2 cooling towers. The density and growth of these *Corbicula* were monitored by collecting monthly samples from the lower reservoir sidewalls and sediments. The sampler used on the sidewalls consisted of a D-frame net attached behind a 24-inch long metal scraping edge. This device was connected to a pole long enough to allow the sampler to extend down into the reservoir area from the outside wall of the cooling tower. Sediments were sampled with a petite Ponar.

The Cooling Tower Reservoir Sampling was historically conducted once per month, annually. Beginning in December 1997, it was decided to forego sampling in cold water months since buildup of *Corbicula* does not occur then. Monthly sampling has been maintained throughout the warmer water months of the year. In 2007 sampling began in March and ended in early November.

In 2007, once each month (March through November), a single petite Ponar grab sample was taken in the reservoir of each cooling tower to obtain density and growth information on *Corbicula* present in the bottom sediment. The samples collected from each cooling tower were returned to the laboratory and processed. Samples were individually washed, and any *Corbicula* removed and rinsed through a series of stacked U.S. Standard sieves that ranged in mesh size from 1.00 mm to 9.49 mm. Live and dead clams retained in each sieve were counted and the

numbers were recorded. The size distribution data obtained using the sieves reflected clam width, rather than length. Samples containing a small number of *Corbicula* were not sieved; individuals were measured and placed in their respective size categories. A scraping sample of about 12 square feet was also collected at each cooling tower during each monthly sampling effort. This sample was processed in a manner consistent with the petit Ponar samples.

Population surveys of both BVPS cooling tower reservoirs have been conducted during scheduled outages (1986 through 2005) to estimate the number of *Corbicula* present in these structures. During the scheduled shutdown period for each unit, each cooling tower reservoir bottom is sampled by petite Ponar at standardized locations within the reservoir. Counts of live and dead clams and determination of density were made. There were no scheduled outages during 2007 when samples were collected.

#### 5.3.4 *Corbicula* Juvenile Monitoring

The *Corbicula* juvenile study was designed to collect data on *Corbicula* spawning activities and growth of individuals entering the intake from the Ohio River. From 1988 through 1998, clam cages were deployed in the intake forebay to monitor for *Corbicula* that entered the BVPS.

Observational-based concerns that the clam cages could quickly clog with sediment during high sediment periods and, as a result, would not be sampled effectively, led to an evaluation of an alternate sampling technique. From April through June 1997, a study was conducted to compare the results of the clam cage samplers to a petite Ponar dredge technique to determine *Corbicula* presence and density in the BVPS intake bays. It was hypothesized that using a Ponar sampler to collect bottom sediments and analysis of those sediments would provide a more representative sample of *Corbicula* settlement and growth rates, and had the added benefit of not requiring confined space entry to conduct the sampling. Results of the study confirmed this hypothesis.

During the 1998 sampling season, at the request of BVPS personnel, all clam cages were removed after the May 18<sup>th</sup> collection. Monthly petite Ponar grabs from the forebay in the intake building continued thereafter. Samples were processed in the same manner as Cooling Tower Samples (Section 5.3.3).

From 2002 to present, because of site access restrictions, sampling with the petite Ponar has been moved to the Ohio River directly in front of the Intake Structure Building. Collections are presently made in conjunction with the fisheries sampling (May, July, September, and November). During each sampling month two Ponar grabs are taken approximately 20 feet off shore of the intake building. These grab samples are processed in the same manner as when they were collected from within the Intake Structure Building.

### 5.3.5 Zebra Mussel Monitoring

The Zebra Mussel Monitoring Program includes sampling the Ohio River and the circulating river water system of the BVPS.

The objectives of the Monitoring Program were:

- (1) To identify if zebra mussels were in the Ohio River adjacent to BVPS and provide early warning to operations personnel as to their possible infestation;
- (2) To provide data as to when the larvae were mobile in the Ohio River and insights as to their vulnerability to potential treatments; and
- (3) To provide data on their overall density and growth rates under different water temperatures and provide estimates on the time it requires these mussels to reach the size and density that could impact the plant.

The zebra mussel sampling for settled adults was historically conducted once per month, yearlong. Beginning in December 1997, it was decided to forego sampling in the colder water months of each year, since buildup of zebra mussels, does not occur then. Monthly sampling has been maintained throughout the balance of the year. In 2007 sampling occurred from March through November.

A pump sample for zebra mussel veligers was collected at the barge slip location monthly from April through October in 1996 and 1997. The scope of the sampling was expanded in 1998 to also include the intake structure. In June 1998, the emergency outfall and emergency outfall impact basin locations were also added. Additional pump samples were collected from the cooling tower of Unit 1 and Unit 2 in October 1998. In 2007 veliger sampling began in April and was conducted monthly through October.

At the Intake Structure and Barge Slip the following surveillance techniques were used:

- Wall scraper sample collections on a monthly basis (March through November) from the barge slip and the riprap near the intake structure to detect attached adults; and
- Pump sample collections from the barge slip and outside the intake structure, to detect the planktonic early life forms (April through October).

At each of the Cooling Towers the following techniques were used:

- Monthly reservoir scraper sample collections in each cooling tower (March through November); and
- Pump samples in April through October to detect planktonic life forms.



At the Emergency Outfall and the Splash Pool the following techniques were used:

- Monthly scraper sample collections in each (March through November); and
- Pump samples in each from April through October to detect planktonic life forms.

#### 5.3.6 Reports

Each month, activity reports that summarized the activities that took place the previous month were prepared and submitted. These reports included the results of the monthly *Corbicula*/zebra mussel monitoring including any trends observed and any preliminary results available from the benthic and fisheries programs. The reports addressed progress made on each task, and reported any observed biological activity of interest.

### 5.4 RESULTS OF THE AQUATIC MONITORING PROGRAM

The following sections summarize the findings for each of the program elements. Sampling dates for each of the program elements are presented in Table 5.1.

#### 5.4.1 Benthic Macroinvertebrate Monitoring Program

Benthic surveys were scheduled and performed in May and September 2007. Benthic samples were collected at Stations 1, 2A, 2B, and 3 (Figure 5.2), using a petite Ponar grab sampler. Triplicate samples were taken off the south shore at Stations 1, 2A, and 3. Sampling at Station 2B, in the back channel of Phillis Island, consisted of triplicate petite Ponar grabs at the south side, middle, and north side of the channel (i.e., Sample Stations 2B1, 2B2, and 2B3, respectively).

Substrate type is an important factor in determining the composition of the benthic community. The habitats in the vicinity of BVPS are the result of damming, channelization, and river traffic. Shoreline habitats at the majority of sampling locations were generally soft muck substrates composed of sand, silt, and detritus. An exception was along the north shoreline of Phillis Island at Station 2A where hard pan clay dominated. The other distinct habitat, hard substrate (gravel and cobble), was located in mid-channel of the back channel of Phillis Island. The hard substrate was probably the result of channelization and scouring by river currents.

Forty-two (42) macroinvertebrate taxa were identified during the 2007 monitoring program (Tables 5.2 and 5.3). A mean density of 727 macroinvertebrates/m<sup>2</sup> was collected in May and 2,183/m<sup>2</sup> in September (Table 5.4). As in previous years, the macroinvertebrate assemblage during 2007 was dominated by burrowing organisms typical of soft unconsolidated substrates. Oligochaetes (segmented worms), mollusks (clams and snails) and chironomid (midge fly) larvae were abundant (Table 5.4).

Twenty-five (25) taxa were present in the May samples, and thirty-seven (37) taxa in the September samples (Table 5.3.1 and 5.3.2). Twenty (20) of the 42 taxa were present in both May and September.

The Asiatic clam (*Corbicula*) has been observed in the Ohio River near BVPS from 1974 to present. Zebra mussels were first collected in the BVPS benthic samples in 1998. Adult zebra mussels, however, were detected in 1995 and 1996 by divers in the BVPS main and auxiliary intake structures during scheduled cleaning operations. Zebra mussel veligers, adults and juveniles were collected during the 1997-2007 sampling programs (see Sections 5.4.5 Zebra Mussel Monitoring Program). Both live *Corbicula* and zebra mussels were collected in benthic macroinvertebrate samples in 2007.

In 2007, one taxon was added to the cumulative taxa list of macroinvertebrates collected near BVPS (Table 5.2). The new taxon was *Stempellina* sp., which is a chironomid (midge fly) larvae. No state or Federal threatened or endangered macroinvertebrate species were collected during 2007.

In the May 2007 samples, oligochaetes accounted for the highest mean density of macroinvertebrates and chironomids had the second highest (552/m<sup>2</sup> or 76 percent of the total density and 98/m<sup>2</sup> or 13 percent, respectively) (Table 5.4). Mollusks had a mean density of only 29/m<sup>2</sup>. Organisms other than oligochaetes, chironomids and mollusks were present at a density of 48/m<sup>2</sup> in May.

In September 2007 samples, oligochaetes accounted for the highest mean density of macroinvertebrates and mollusks had the second highest (1488/m<sup>2</sup> or 68 percent of the total density and 351/m<sup>2</sup> or 16 percent, respectively) (Table 5.4). Chironomids had the third highest mean density in September 2007 (291/m<sup>2</sup> or 13 percent) while the "others" category had the fourth highest mean density (53/m<sup>2</sup> or 2 percent).

In May, the highest density of macroinvertebrates (1,547 organisms/m<sup>2</sup>) occurred at Station 2B1. In September, the highest density of macroinvertebrates occurred at Station 2B3 (8,586/m<sup>2</sup>). In May the lowest mean density of organisms occurred at Station 3 (415/m<sup>2</sup>). In September, the lowest mean density of organisms occurred at Station 2A (530/m<sup>2</sup>).

For a comparison of the control to non-control stations, Station 1 was designated the control station, because it is always out of the influence of the BVPS discharge and Station 2B (mean density of Station 2B1, 2B2, and 2B3) was designated as the non-control station, since it is the

station most regularly subjected to BVPS's discharge. Stations 3 and 2A may be under the influence of the plume under certain conditions, but it is unlikely that they are regularly influenced by BVPS.

The mean density of macroinvertebrates in the non-control station was 1.63 times higher ( $912/m^2$ ) than that of the control station ( $559/m^2$ ) in May (Table 5.5). The density of each of the major groups (chironomids, oligochaetes, mollusks, and others) was higher at the non-control station than at the control station. Overall the differences probably reflect the natural differences in substrate and natural heterogeneous distributions of these organisms between the stations rather than project-related impacts.

In September, the density of macroinvertebrates present at the non-control ( $3,793/m^2$ ) was about 6.8 times greater than at the control station ( $560/m^2$ ). The density of oligochaetes was about 11 times higher at the non-control than the control stations. The density of chironomids was 1.8 times higher at the non-control than the control stations. The density of mollusks was 6.7 times higher in the non-control station ( $487/m^2$ ) than in the control ( $72/m^2$ ). Other taxa were present only in the non control stations. As in May, the differences observed between Station 1 (control) and Station 2B (non-control) were probably related to observed differences in habitat at each station. Differences were within the expected range of variation for natural populations of macroinvertebrates.

Indices that describe the relative diversity, evenness, and richness of the macroinvertebrate population structure among stations and between control and non-control sites were calculated. A higher Shannon-Weiner diversity index indicates a relatively better structured assemblage of organisms, while a lower index generally indicates a low quality or stressed community. Evenness is an index that estimates the relative contribution of each taxon to the community assemblage, the closer to 1.00, the healthier the community. The community richness is another estimate of the quality of the macroinvertebrate community with a higher richness number indicating a healthier community.

The Shannon-Weiner diversity indices in May 2007 collections ranged from 0.42 at Station 2A to 0.88 at Station 2B3 (Table 5.6). In May evenness ranged from 0.49 at Station 2A to 0.90 at Station 3. Richness was greatest at Station 2B3 (3.52) and lowest at Station 2A (1.05). The overall low indices at Station 2A is probably attributed to the dominant substrate, hard pan clay, which generally is a low quality benthic habitat. The range in diversity of the macroinvertebrate community in September was generally comparable to that in May. Diversity ranged from 0.45 at Station 2B3 to 0.95 at Station 2B1. Evenness was also comparable in September to May and ranged from 0.34 at Station 2B3 to 0.90 at Station 3. Richness was greatest at Station 3 (2.53) and lowest at Station 1 (0.55).

In May, the number of taxa, diversity, evenness and richness indices were somewhat lower in the control station (Station 1) and in the non-control stations (2B1, 2B2, 2B3) (Table 5.6). In September the same pattern held true. These differences were also apparent in 2006 and were likely due to natural annual variations in the local populations at these locations. No impacts of the BVPS on the benthic community, as measured by differences between control and non-

control zones, were evident in either May or September.

Substrate was probably the most important factor controlling the distribution and abundance of the benthic macroinvertebrates in the Ohio River near BVPS. Soft, mucky substrates that generally existed along the shoreline are conducive to oligochaete, chironomid, and mollusk habitation and limit species of macroinvertebrates that require a more stable bottom.

The density of macroinvertebrates in May and September 2007 fell within the range of densities of macroinvertebrates collected at BVPS in previous years (Table 5.7). *The community structure has changed little since pre-operational years, and the available evidence does not indicate that BVPS operations have affected the benthic community of the Ohio River.*

#### 5.4.2 Fish Sampling Program

In 2007, 216 fishes representing 24 taxa were collected (i.e., handled) during BVPS surveys by electrofishing and seining (Table 5.8). All taxa collected in 2007 were previously encountered at BVPS. The most common species in the 2007 BVPS surveys, collected by electrofishing and seining combined, were smallmouth bass (14.4% of the total catch), followed by shorthead redhorse sucker (12.5 %), smallmouth buffalo (12.0%), gizzard shad (11.1%), sauger (10.2%), golden redhorse sucker (9.7%) and spotted bass (8.3%). The remaining 17 species combined accounted for 8.3% of the total handled catch. The most frequently observed but not handled fish in 2007 were gizzard shad (Table 5.15). Game fishes collected during 2007 included channel catfish, flathead catfish, white bass, black crappie, bluegill, smallmouth bass, sauger, walleye, rock bass and spotted bass. Game fishes represented 40.7 % of the total handled catch, 24.5% of which were smallmouth bass and sauger.

A total of 189 fish, representing 23 taxa, was collected by electrofishing in 2007 (Table 5.9). Shorthead redhorse suckers and smallmouth bass accounted for the largest portion of the 2007 electrofishing catch (13.8% and 13.2%, respectively) followed by smallmouth buffalo (12.2%), sauger (11.6%), golden redhorse sucker (11.1%), gizzard shad (9.5%) and spotted bass (5.8%). No other species collected contributed to greater than five percent of the total catch. Fish observed and not collected in the 2007 electrofishing study are presented in Table 5.15.

A total of 27 fish representing 8 taxa was collected by seining in 2007 (Table 5.10). The most abundant taxa collected was spotted bass (25.9% of the total catch) followed by gizzard shad (22.2%) and smallmouth bass (18.5%). The only other game species collected by seining was a white bass.

A total of 50 fish representing 14 species was captured during the May 2007 sampling event (Table 5.11). All of the fish were collected during electrofishing; none during seine netting. Golden redhorse sucker (32.0% of the total catch), shorthead redhorse sucker (14.0%) and sauger (12.0%) were the most common species boated during the electrofishing effort. Sauger, bluegill, rock bass, white bass, smallmouth bass, spotted bass, and black crappie were the game species collected in May.

A total of 56 fish representing 14 species was captured during the July 2007 sampling event (Table 5.12). A total of 37 fish representing 12 species was collected during electrofishing efforts. Gizzard shad (32.4% of the total catch), smallmouth buffalo (13.5%), carp (13.5%) and smallmouth bass (10.8%) were the most common species boated during the electrofishing effort. Hundreds of small gizzard shad were observed throughout the study reach during electrofishing, but not boated due to their small size (Table 5.15). Smallmouth bass, flathead catfish, sauger, walleye, and spotted bass were the only game species collected during the July electrofishing study (Table 5.12). A total of 19 fish representing seven species was collected by the seines. Gizzard shad were also the most abundant species in seine netting and represented 32.4% of the total catch. Juvenile spotted bass (26.3% of the total catch) and smallmouth buffalo (15.8% of the catch) were the next most abundant species in seine netting. The spotted bass and a juvenile white bass were the only game fish collected.

During the September sampling event, 43 fish representing 114 taxa were collected during fish sampling efforts (Table 5.13). A total of 35 fish representing 14 species was collected during electrofishing in September. Smallmouth bass and smallmouth buffalo were the most abundant species, each contributing to 20% of the total catch. Longnose gar (14.3% of the electrofishing catch) and shorthead redhorse sucker (8.6%) were also relatively abundant. White bass, channel catfish, sauger and bluegill were game species collected during electrofishing efforts in September. Eight fish; four smallmouth bass, two spotted bass and two shorthead redhorse suckers, were the only fish collected in seines. All of the fish collected in the seines were juveniles.

During the November sampling event, 67 fish representing 15 taxa were captured (Table 5.14). All of the fish were collected during electrofishing; none during seine netting. Sauger and smallmouth bass were the most abundant species collected by electrofishing, each contributing to 19.4% of the total. Other relatively abundant species were shorthead redhorse sucker (17.9% of the total), spotted bass (10.5%), and smallmouth buffalo (9.0%). Game species collected by electrofishing included walleye, white bass, smallmouth bass, bluegill, sauger and spotted bass. Fish observed and not collected in the November electrofishing study are presented in Table 5.15.

Electrofishing catch rates are presented in Tables 5.16, 5.17, and 5.18 for fish that were boated and handled during the 2005 through 2007 surveys by season (FENOC 2006 and 2007). In 2007, the annual catch rate was 1.17 fish per minute. The greatest catch rate in 2007 occurred in November (winter) (1.63 fish/ electrofishing minute). Sauger, smallmouth bass, and shorthead redhorse suckers contributed the most to this total. The lowest catch rate occurred in September (fall) with a rate of 0.85 fish/ electrofishing minute. The annual catch rates were consistent over the three years ranging from a high of 1.60 fish per minute in 2006 to 1.16 in 2005. Over the three years, the highest seasonal catch rate was 2.85 fish per minute, which occurred in May 2006. The lowest seasonal catch rate was 0.70 fish per minute, which occurred in July 2006. In 2006 the highest catch rate was in the spring whereas in 2005 and 2007 the highest catch rate was during the winter survey.

The results of the electrofishing sampling effort (Table 5.9) did not indicate any major differences in species composition between the control station (1) and the non-control Stations 2A, 2B, and 3. A greater number of fish representing more species was captured at non-control stations than control stations. This was most likely due to the extra effort expended at non-control stations versus control stations (i.e., there are three non-control stations and only one control station). In 2007, few individuals and species were collected by seines at both control and non-control stations (Table 5.10). The lower numbers at both locations than in previous years could be attributed to river conditions at the time of sampling rather than to any impacts attributable to BVPS operation.

In 2007, species composition remained comparable among stations. Common taxa collected in the 2007 surveys by all methods included gizzard shad, redhorse sucker species, smallmouth buffalo, sauger, smallmouth bass, and walleye. Little difference in the species composition of the catch was observed between the control (1) and non-control stations (2A, 2B and 3). Habitat preference and availability were probably the most important factors affecting where and when different species of fish are collected.

The results of the 2007 fish surveys indicated that there is a normal community structure in the Ohio River in the vicinity of BVPS based on species composition and relative abundance of fish observed during the surveys. Benthivores (bottom feeders including suckers and buffalo) were collected in the highest numbers. Forage species, although common were collected in lower numbers than in other study years. Variations in annual catch were probably attributable to normal fluctuations in the population size of the forage species and the predator populations that rely on them. Forage species, such as gizzard shad and emerald shiner with high reproductive potentials, frequently respond to changes in natural environmental factors (competition, food availability, cover, and water quality) with large fluctuations in population size, which could be the reason for the lower numbers of these species observed in 2007. This, in turn, influences their appearance in the sample populations during annual surveys. Spawning/rearing success due to abiotic factors is usually the determining factor of the size and composition of a fish community. In addition, differences in electrofishing catch rate can be attributed to environmental conditions that prevail during sampling efforts. High water, increased turbidity, and swift currents that occur during electrofishing efforts in some years can decrease the collection efficiency of this year.

#### 5.4.3 Corbicula Monitoring Program

In 2007, 20 *Corbicula* (35.0 percent alive) were collected from the Unit 1 cooling tower basin during monthly reservoir sampling. No *Corbicula* were collected in the scraping samples. The largest *Corbicula* collected was retained in a sieve with a 6.30-9.49 mm length size range (Table 5.19 and Figure 5.5). The greatest numbers of *Corbicula* were collected in March (six individuals). The mean density of total *Corbicula* (live and dead) in Unit 1 in 2007 was 108/m<sup>2</sup>. *Corbicula* were collected in all months sampled except November. No sampling was conducted in September because the Unit 1 Cooling Tower could not be accessed due to unit shutdown.

In 2007, 24 *Corbicula* (91.7 percent alive) were collected from the Unit 2 cooling tower reservoir

during monthly sampling. The largest *Corbicula* collected was within the 3.35-4.74 mm length size range (Table 5.20 and Figure 5.6). Individuals were collected in May through September. The mean density of *Corbicula* in Unit 2 in 2007 was 128/m<sup>2</sup>. The greatest numbers of *Corbicula* (nine individuals) were collected in September.

In 2007, BVPS continued its *Corbicula* control program (Year 15), which included the use of a molluscicide (CT-1) to prevent the proliferation of *Corbicula* within BVPS. BVPS was granted permission by the Pennsylvania Department of Environmental Protection to use CT-1 to target the Unit 1 river water system and the Unit 2 service water system.

In 1990 through 1993, the molluscicide applications (CT-1) focused on reducing the *Corbicula* population throughout the entire river water system of each BVPS plant (Units 1 and 2). In 1994 and 1995, the CT-1 applications targeted the internal water systems; therefore, the CT-1 concentrations in the cooling towers were reduced during CT-1 applications. Consequently, adult and juvenile *Corbicula* in the cooling towers often survived the CT-1 applications. Reservoir sediment samples taken after CT-1 applications represent mortality of *Corbicula* in the cooling tower only and do not reflect mortality in BVPS internal water systems.

The monthly reservoir sediment samples collected in Units 1 and 2 Cooling Towers during 2007 demonstrated that *Corbicula* were entering and colonizing the reservoirs. Overall, densities in Unit 2 were somewhat less than those in 2006 and in Unit 1 densities were comparable to in 2006. The recent decrease of *Corbicula* at the BVPS returns densities to levels more consistent with densities in the Ohio River in the mid-1990's, but well below those present during the 1980's.

#### 5.4.4 *Corbicula* Juvenile Monitoring Program

Figure 5.7 presents the abundance and size distribution data for samples collected in the Ohio River near the intake structure by petite ponar dredge in 2007. Three live individuals were collected during May, nine in July and one each in September and November. They ranged in size from the 1.00-1.99mm size range that were spawned in 2007 to greater than 9.50mm that were spawned in prior years. The number of individuals collected in 2007 (14) was comparable with the number collected in 2005 (17 individuals) and higher than in 2006 (three individuals).

A spring/early-summer spawning period typically occurs in the Ohio River near BVPS each year when preferred spawning temperatures (60-65° F) are reached (Figure 5.8). The offspring from this spawning event generally begin appearing in the sample collections in May. The settled clams generally increase in size throughout the year. ***The overall low numbers of live Corbicula collected in the sample collected outside the intake and cooling towers in 2007, compared to levels in the 1980's, likely reflects a natural decrease in the density of Corbicula in the Ohio River near BVPS.***

#### 5.4.5 Zebra Mussel Monitoring Program

Zebra mussels (*Dreissena polymorpha*) are exotic freshwater mollusks that have ventrally flattened shells generally marked with alternating dark and lighter bands. They are believed to have been introduced into North America through the ballast water of ocean-going cargo vessels probably from Eastern Europe. They were first identified in Lake St. Clair in 1988 and rapidly spread to other Great Lakes and the Mississippi River drainage system, becoming increasingly abundant in the lower, middle, and upper Ohio River. They use strong adhesive byssal threads, collectively referred to as the byssus, to attach themselves to any hard surfaces (e.g., intake pipes, cooling water intake systems, and other mussels). Responding to NRC Notice No. 89-76 (Biofouling Agent-Zebra Mussel, November 21, 1989), BVPS instituted a Zebra Mussel Monitoring Program in January 1990. Studies have been conducted each year since then.

Zebra mussels were detected in both pump samples (Figures 5.9 and 5.10) and substrate samples (Figure 5.11 and 5.12) in 2007. Zebra mussel veliger pump samples were collected from April through October (Figures 5.9 and 5.10). Veligers were collected at all of the six sites that were sampled in 2007. Densities of veligers generally increased through the year, and peaked at all sample locations in September or October. The greatest density of veligers was present in the sample collected from the Emergency Outfall Facility in September (3,293/m<sup>3</sup>). Veligers were present in all months sampled. In March, they were present only in the cooling towers, which had greater than ambient water temperature that were within the required temperature range for zebra mussel spawning. Overall, veliger densities were much lower than in 2006 but comparable in magnitude to 2005 and 2004. This is likely due to annual variability in numbers of veligers in the Ohio River.

In 2007, settled zebra mussels were collected in scrape samples at the barge slip, and the intake structure (Figures 5.11 and 5.12). The highest density of mussels was present in the sample collected at the barge slip (22 mussels/m<sup>2</sup>) in November. The mussels collected at each of the sites included individuals that were capable of reproducing. The density of collected adult zebra mussels was consistent with past years.

Overall, both the number of observations and densities of settled mussels were similar in 2003-2007. The density of veligers in 2007 was much lower than in 2006 or 2003, but was comparable to 2005 and 2004. Although densities of settled mussels are low compared to other populations such as the Lower Great Lakes, densities comparable to those in the Ohio River are sufficient to cause problems in the operation of untreated cooling water intake systems. ***Whether the population of zebra mussels in this reach of the Ohio River is resurging or only yearly fluctuations are present cannot be determined. In any case, the densities of mussels that presently exist are more than sufficient to impact the BVPS, if continued prudent monitoring and control activities are not conducted.***



## 6.0 ZEBRA MUSSEL AND *CORBICULA* CONTROL ACTIVITIES

In 2007, BVPS continued its *Corbicula* and zebra mussel control program (14<sup>th</sup> year), which included the use of a molluscicide (CT-1) to prevent the proliferation of *Corbicula* within BVPS.

BVPS was granted permission by the Pennsylvania Department of Environmental Protection to use CT-1 to target the Unit 1 river water system and the Unit 2 service water system.

In 1990 through 1993, the molluscicide applications (CT-1) focused on reducing the *Corbicula* population throughout the entire river water system of each BVPS plant (Units 1 and 2). In 1994 through 2007, the CT-1 or 2 applications targeted zebra mussels and *Corbicula* in the internal water systems; therefore the molluscicide concentrations in the cooling towers were reduced during CT-1 or 2 applications. Consequently, adult and juvenile *Corbicula* in the cooling towers often survived the applications. Reservoir sediment samples taken after CT-1 or 2 applications represented mortality of *Corbicula* in the cooling tower only and do not reflect mortality in BVPS internal water systems.

In addition to clamicide treatments, preventive measures were taken that included quarterly cleaning of the Intake Bays. The bay cleanings are intended to minimize the accumulation and growth of mussels within the bays. This practice prevents creating an uncontrolled internal colonization habitat.

## 7.0 REFERENCES

- Commonwealth of Pennsylvania, 1994. Pennsylvania's Endangered Fishes, Reptiles and Amphibians. Published by the Pennsylvania Fish Commission.
- Counts, C. C. III, 1985. Distribution of *Corbicula fluminea* at Nuclear Facilities. Division of Engineering, U.S. Nuclear Regulatory Commission. NUREGLCR. 4233. 79 pp.
- Dahlberg, M. D. and E. P. Odum, 1970. Annual cycles of species occurrence, abundance and diversity in Georgia estuarine fish populations. Am. Midl. Nat. 83:382-392.
- FENOC, 2003. Annual Environmental Operating Report, Non-radiological. First Energy Nuclear Operating Company, Beaver Valley Power Station, Unit No. 1 & 2. 113 pp.
- FENOC, 2004. Annual Environmental Operating Report, Non-radiological. First Energy Operating Company, Beaver Valley Power Station, Unit No. 1&2. 82 pp.
- FENOC, 2005. Annual Environmental Operating Report, Non-radiological. First Energy Operating Company, Beaver Valley Power Station, Unit No. 1&2. 82 pp.
- FENOC, 2006. Annual Environmental Operating Report, Non-radiological. First Energy Operating Company, Beaver Valley Power Station, Unit No. 1&2. 82 pp.
- Hutchinson, G. E., 1967. A treatise on limnology. Vol. 2, Introduction to lake biology and the limnoplankton. John Wiley and Sons, Inc., New York. 1115 pp.
- Hynes, H. B. N., 1970. The ecology of running waters. Univ. Toronto Press, Toronto.
- NRC, IE Bulletin 81-03: Flow Blockage of Cooling Tower to Safety System Components by *Corbicula sp.* (Asiatic Clam) and *Mytilus sp.* (Mussel).
- Pielou, E. C., 1969. An introduction to mathematical ecology. Wiley Interscience, Wiley & Sons, New York, NY.
- Robins, C. R., R. M. Bailey, C. E. Bond, J. R. Brooker, E. A. Lachner, R. N. Lea, and W. B. Scott, 1991. Common and Scientific Names of Fishes from the United States and Canada (fifth edition). American Fisheries Society Special Publication No. 20:1-183.
- Shiffer, C., 1990. Identification Guide to Pennsylvania Fishes. Pennsylvania Fish Commission, Bureau of Education and Information. 51 pp.
- Winner, J. M., 1975. Zooplankton. In: B. A. Whitton, ed. River ecology. Univ. Calif. Press, Berkeley and Los Angeles. 155-169 pp.

**8.0**

**TABLES**

TABLE 5.1

BEAVER VALLEY POWER STATION (BVPS)  
SAMPLING DATES FOR 2007

Study	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Benthic Macroinvertebrate					22				24		8	
Fish					22,29		31		24		8	
<i>Corbicula</i> and Zebra Mussel			22	23	22	13	31	27	24	17	8	
<i>Corbicula</i> CT Density												
Zebra Mussel Veliger				23	22	13	31	27	24	17		

Table 5.2

## Systematic List of Macroinvertebrates Collected From 1973 Through 2007 in The Ohio River Near BVPS

Phylum	Class	Family	Sub-Family	Genus and Species	Previous Collections	Collected in 2007	New in 2007
Porifera				<i>Spongilla fragilis</i>	X		
Cnidaria							
	Hydrozoa						
		Clavidae		<i>Cordylophora lacustris</i>	X		
		Hydridae		<i>Craspedacusta sowerbii</i>	X		
				<i>Hydra sp.</i>	X		
Platyhelminthes							
	Tricladida				X		
	Rhabdocoela				X		
Nemertea					X		
Nematoda					X		
Entoprocta				<i>Urnatella gracilis</i>	X		
Ectoprocta							
				<i>Fredericella sp.</i>	X		
				<i>Paludicella articulata</i>	X		
				<i>Pectinatella sp.</i>	X		
				<i>Plumatella sp.</i>	X		
Annelida							
	Oligochaeta				X		
		Aeolosomatidae			X		
		Enchytraeidae			X	X	
		Naididae			X	X	
				<i>Allonais pectinata</i>	X		
				<i>Amphichaeta leydigi</i>	X		
				<i>Amphichaeta sp.</i>	X		
				<i>Arcteonais lomondi</i>	X		
				<i>Aulophorus sp.</i>	X		
				<i>Chaetogaster diaphanus</i>	X		
				<i>C. diastrophus</i>	X		
				<i>Dero digitata</i>	X		
				<i>Dero flabelliger</i>	X		
				<i>D. nivea</i>	X		
				<i>Dero sp.</i>	X		
				<i>Nais barbata</i>	X		
				<i>N. behningi</i>	X		
				<i>N. bretscheri</i>	X		
				<i>N. communis</i>	X		
				<i>N. elinguis</i>	X		
				<i>N. pardalis</i>	X		
				<i>N. pseudobtusa</i>	X		
				<i>N. simplex</i>	X		
				<i>N. variabilis</i>	X	X	
				<i>Nais sp.</i>	X		
				<i>Ophidonais serpentina</i>	X		
				<i>Paranais frici</i>	X	X	
				<i>Paranais litoralis</i>	X		
				<i>Paranais sp.</i>	X		
				<i>Piquetiella michiganensis</i>	X		
				<i>Pristina idrensis</i>	X		
				<i>Pristina longisoma</i>	X		
				<i>Pristina longiseta</i>	X		
				<i>P. osborni</i>	X	X	
				<i>P. sima</i>	X		
				<i>Pristina sp.</i>	X		
				<i>Pristinella sp.</i>	X		

Table 5.2 (continued)

Systematic List of Macroinvertebrates Collected From 1973 Through 2007 in The Ohio River Near BVPS

Phylum	Class	Family	Sub-Family	Genus and Species	Previous Collections	Collected in 2007	New in 2007		
Annelida	Oligochaeta	Naididae		<i>Pristinella jenkiniae</i>	X				
				<i>Pristinella idrensis</i>	X				
				<i>Pristinella osborni</i>	X				
				<i>Ripistes parasita</i>	X				
				<i>Slavina appendiculata</i>	X				
				<i>Specaria josinae</i>	X	X			
				<i>Stephensoniana trivandranana</i>	X				
				<i>Stylaria fossularis</i>	X				
				<i>S. lacustris</i>	X				
				<i>Uncinais uncinata</i>	X				
				<i>Vejdovskiyella comata</i>	X				
				<i>Vejdovskiyella intermedia</i>	X				
				<i>Vejdovskiyella sp.</i>	X				
			Tubificida	Tubificidae			X		
						<i>Aulodrilus limnobius</i>	X	X	
						<i>A. pigueti</i>	X		
						<i>A. pluriseta</i>	X		
						<i>Aulodrilus sp.</i>	X		
						<i>Bothrioneurum vejdovskyanum</i>	X		
						<i>Branchiura sowerbyi</i>	X	X	
						<i>Ilyodrilus templetoni</i>	X		
						<i>Limnodrilus cervix</i>	X		
						<i>L. cervix (variant)</i>	X	X	
						<i>L. claparedianus</i>	X		
						<i>L. hoffmeisteri</i>	X	X	
						<i>L. maumeensis</i>	X	X	
						<i>L. profundicla</i>	X	X	
						<i>L. spiralis</i>	X		
						<i>L. udekemianus</i>	X		
						<i>Limnodrilus sp.</i>	X		
						<i>Peloscolex multisetosus longidentus</i>	X		
						<i>P. m. multisetosus</i>	X		
						<i>Potamothrix moldaviensis</i>	X		
						<i>Potamothrix sp.</i>	X		
						<i>P. vejdovskyi</i>	X	X	
						<i>Psammoryctides curvisetosus</i>	X		
						<i>Tubifex tubifex</i>	X		
						Unidentified immature forms:			
						with hair chaetae		X	
						without hair chaetae		X	X
						Lumbriculidae		X	
		Hirudinae		X					
		Glossiphoniidae		X					
			<i>Helobdella elongata</i>	X					
			<i>H. stagnalis</i>	X					
			<i>Helobdella sp.</i>	X					
		Erpobdellidae							
			<i>Erpobdella sp.</i>	X					
			<i>Mooreobdella microstoma</i>	X					
		Haplotaxidae							
			<i>Stylodrilus heringianus</i>	X					
	Lumbricina			X					
		Lumbricidae		X					

Table 5.2 (continued)  
 Systematic List of Macroinvertebrates Collected From 1973 Through 2007 in The Ohio River Near BVPS

Phylum	Class	Family	Sub-Family	Genus and Species	Previous Collections	Collected in 2007	New in 2007
<b>Arthropoda</b>							
	Acarina				X		
	Ostracoda				X		
	Isopoda						
				<i>Asellus sp.</i>	X		
<b>Arthropoda</b>							
	Amphipoda						
		Talitridae					
				<i>Hyaella azteca</i>	X		
		Gammaridae					
				<i>Crangonyx pseudogracilis</i>	X		
				<i>Crangonyx sp.</i>	X		
				<i>Gammarus fasciatus</i>	X		
				<i>Gammarus sp.</i>	X	X	
		Pontoporeiidae					
				<i>Monoporeia affinis</i>	X		
<b>Decapoda</b>							
					X		
<b>Collembola</b>							
					X		
<b>Ephemeroptera</b>							
		Heptageniidae			X		
				<i>Stenacron sp.</i>	X		
				<i>Stenonema sp.</i>	X		
		Ephemeridae					
				<i>Ephemera sp.</i>	X		
				<i>Hexagenia sp.</i>	X	X	
				<i>Ephron sp.</i>	X		
		Baetidae					
				<i>Baetis sp.</i>	X		
		Caenidae					
				<i>Caenis sp.</i>	X	X	
				<i>Serratella sp.</i>	X		
		Tricorythidae					
				<i>Tricorythodes sp.</i>	X		
<b>Megaloptera</b>							
				<i>Stalis sp.</i>	X		
<b>Odonata</b>							
		Gomphidae					
				<i>Argia sp.</i>	X		
				<i>Dromogomphus spoliatus</i>	X		
				<i>Dromogomphus sp.</i>	X		
				<i>Gomphus sp.</i>	X		
		Libellulidae					
				<i>Libellula sp.</i>	X		
<b>Plecoptera</b>							
					X		
<b>Trichoptera</b>							
		Hydropsychidae					
				<i>Cheumatopsyche sp.</i>	X		
				<i>Hydropsyche sp.</i>	X		
				<i>Parapsyche sp.</i>	X		
		Hydroptilidae					
				<i>Hydroptila sp.</i>			
				<i>Orthotrichia sp.</i>			
				<i>Oxyethira sp.</i>			
		Leptoceridae					
				<i>Ceraclea sp.</i>	X		
				<i>Oecetis sp.</i>	X		
		Polycentropodidae					
				<i>Cyrnellus sp.</i>	X		
<b>Trichoptera</b>							
		Polycentropodidae		<i>Polycentropus sp.</i>	X		

Table 5.2 (continued)  
 Systematic List of Macroinvertebrates Collected From 1973 Through 2007 in The Ohio River Near BVPS

Phylum	Class	Family	Sub-Family	Genus and Species	Previous Collections	Collected in 2007	New in 2007
Coleoptera		Hydrophilidae			X		
Coleoptera		Elmidae		<i>Ancyronyx variegatus</i>	X		
				<i>Dubiraphia sp.</i>	X		
				<i>Helichus sp.</i>	X		
				<i>Optioserus sp.</i>	X		
				<i>Stenelmis sp.</i>	X		
		Psephenidae			X		
Diptera				Unidentified Diptera	X		
		Psychodidae			X		
				<i>Pericoma sp.</i>	X		
				<i>Psychoda sp.</i>	X		
				<i>Telmatoscopus sp.</i>	X		
				Unidentified Psychodidae pupae	X		
		Chaoboridae					
				<i>Chaoborus sp.</i>	X		
		Simuliidae					
				<i>Simulium sp.</i>	X		
		Chironomidae			X		
		Chironominae			X		
				Tanytarsini pupa	X		
				Chironominae pupa	X	X	
				<i>Axarus sp.</i>	X		
				<i>Chironomus sp.</i>	X	X	
				<i>Cladopelma sp.</i>	X	X	
				<i>Cladotanytarsus sp.</i>	X	X	
				<i>Cryptochironomus sp.</i>	X	X	
				<i>Dicrotendipes nervosus</i>	X		
				<i>Dicrotendipes sp.</i>	X	X	
				<i>Glyptotendipes sp.</i>	X		
				<i>Harnischia sp.</i>	X		
				<i>Microchironomus sp.</i>	X		
				<i>Micropsectra sp.</i>	X		
				<i>Microtendipes sp.</i>	X		
				<i>Parachironomus sp.</i>	X		
				<i>Paracladopelma sp.</i>	X		
				<i>Paratanytarsus sp.</i>	X		
				<i>Paratendipes sp.</i>	X	X	
				<i>Phaenopsectra sp.</i>	X		
				<i>Polypedilum (s.s.) convictum</i> type	X		
				<i>P. (s.s.) simulans</i> type	X		
				<i>Polypedilum sp.</i>	X	X	
				<i>Rheotanytarsus sp.</i>	X		
				<i>Stempelina sp.</i>		X	X
				<i>Stenochironomus sp.</i>	X		
				<i>Stictochironomus sp.</i>	X		
				<i>Tanytarsus coffinani</i>	X		
				<i>Tanytarsus sp.</i>	X	X	
				<i>Tribelos sp.</i>	X		
				<i>Xenochironomus sp.</i>	X	X	
		Tanypodinae			X		
				Tanypodinae pupae	X		
				<i>Ablabesmyia sp.</i>	X		
				<i>Clinotanypus sp.</i>	X	X	
				<i>Coelotanypus scapularis</i>	X		
				<i>Coelotanypus sp.</i>	X	X	
				<i>Djalmabatista pulcher</i>	X		
				<i>Djalmabatista sp.</i>	X		
				<i>Procladius sp.</i>	X	X	
				<i>Tanypus sp.</i>	X		



Table 5.2 (continued)

## Systematic List of Macroinvertebrates Collected From 1973 Through 2007 in The Ohio River Near BVPS

Phylum	Class	Family	Sub-Family	Genus and Species	Previous Collections	Collected in 2007	New in 2007	
Diptera		Tanypodinae		<i>Thienemannimyia</i> group	X			
				<i>Zavrelimyia</i> sp.	X			
		Orthoclaadiinae		<i>Orthoclaadiinae</i> pupae	X			
				<i>Cricotopus bicinctus</i>	X			
				<i>C. (s.s.) trifascia</i>	X			
				<i>Cricotopus (Isocladus)-sylvestris</i> Group	X			
				<i>C. (Isocladus)</i> sp.	X			
				<i>Cricotopus (s.s.)</i> sp.	X	X		
				<i>Eukiefferiella</i> sp.	X			
				<i>Hydrobaenus</i> sp.	X			
				<i>Limnophyes</i> sp.	X			
				<i>Nanocladius (s.s.) distinctus</i>	X			
				<i>Nanocladius</i> sp.	X			
				<i>Orthocladus</i> sp.	X			
				<i>Parametricnemus</i> sp.	X			
				<i>Paraphaenocladus</i> sp.	X			
				<i>Psectrocladius</i> sp.	X			
				<i>Pseudorthocladus</i> sp.	X			
				<i>Pseudosmittia</i> sp.	X			
				<i>Smittia</i> sp.	X			
				<i>Thienemannimyia</i> sp.	X			
			Diamesinae					
					<i>Diamesa</i> sp.	X		
					<i>Pothastia</i> sp.	X		
			Ceratopogonidae			X		
					<i>Probezzia</i> sp.	X	X	
					<i>Bezzia</i> sp.	X	X	
					<i>Culicoides</i> sp.	X	X	
			Dolichopodidae			X		
			Empididae			X	X	
					<i>Clinocera</i> sp.	X		
					<i>Wiedemannia</i> sp.	X		
			Ephydriidae			X		
	Muscidae			X				
	Rhagionidae			X				
	Tipulidae			X				
	Stratiomyidae			X				
	Syrphidae			X				
Lepidoptera				X				
Hydrachnidia				X				
Mollusca								
	Gastropoda			X				
		Hydrobiidae		X				
		Amnicolinae						
			<i>Amnicola</i> sp.	X	X			
			<i>Aminicola binneyana</i>	X	X			
			<i>Amnicola limosa</i>	X	X			
			<i>Stagnicola elodes</i>	X				
	Physacea			X				
		Pleuroceridae						
			<i>Pleurocera acuta</i>	X				
			<i>Goniobasis</i> sp.	X	X			
		Physidae		X				
			<i>Physa</i> sp.	X				
			<i>Physa ancillaria</i>	X				
			<i>Physa integm</i>	X				

Table 5.2 (continued)

## Systematic List of Macroinvertebrates Collected From 1973 Through 2007 in The Ohio River Near BVPS

Phylum	Class	Family	Sub-Family	Genus and Species	Previous Collections	Collected in 2007	New in 2007			
Mollusca	Physacea	Ancyliidae		Ferrissia sp.	X					
			Planorbidae		Gillia atilis	X				
				Valvatidae		Valvata perdepressa	X			
					Valvata piscinalis	X				
					Valvata sincera sincera	X				
					Valvata sp.	X				
			Pelecypoda		Sphaeriacea	Corbiculidae		Corbicula fluminea	X	X
							Corbicula sp.	X		
				Sphaeriidae				Pisidium ventricosum	X	
	Pisidium sp.	X					X			
	Sphaerium sp.	X								
	Unidentified immature Sphaeriidae	X								
Dreissenidae		Dreissena polymorpha		X			X			
	Unionidae			Anodonta grandis			X			
		Anodonta (immature)		X						
		Elliptio sp.		X						
		Quadrula pustulosa		X						
		Unidentified immature Unionidae		X						

TABLE 5.3

**BENTHIC MACROINVERTEBRATE COUNTS FOR TRIPPLICATE SAMPLES  
TAKEN AT EACH SAMPLE STATION FOR MAY AND SEPTEMBER 2007**

Scientific name	May							Sept							2007 Total
	Location						May Total	Location						Sept Total	
	1	2A	2B1	2B2	2B3	3		1	2A	2B1	2B2	2B3	3		
<i>Amnicola binneyana</i>	0	0	0	0	0	0	0	0	0	0	1	1	0	2	2
<i>Amnicola limosa</i>	0	0	0	0	0	0	0	0	0	2	2	0	7	11	11
<i>Aulodrilus limnobius</i>	0	0	2	0	0	0	2	0	0	3	0	0	0	3	5
<i>Branchiura sowerbyi</i>	0	0	0	0	0	0	0	0	0	2	0	2	3	7	7
<i>Caenis</i> sp.	0	0	0	0	0	0	0	0	0	1	0	0	0	1	1
Chironomid pupae	0	0	0	0	1	0	1	1	0	1	1	1	0	4	5
<i>Chironomus</i> sp.	0	2	3	17	1	0	23	0	1	0	0	0	0	1	24
<i>Cladopelma</i> sp.	0	0	0	0	0	0	0	0	0	1	0	0	0	1	1
<i>Cladotanytarsus</i> sp.	0	0	0	0	0	0	0	0	0	16	0	0	0	16	16
<i>Coelotanytus</i> sp.	0	0	0	0	0	0	0	0	0	0	2	0	5	7	7
<i>Corbicula fluminea</i>	0	0	0	0	0	0	0	2	14	8	2	1	5	32	32
<i>Cricotopus</i> (s.s.) sp.	0	1	0	0	0	0	1	0	0	1	0	0	0	1	2
<i>Cryptochironomus</i> sp.	0	0	0	2	0	0	2	0	1	15	1	19	0	36	38
<i>Culicoides</i> sp.	0	0	0	0	0	0	0	0	0	0	0	7	0	7	7
<i>Dicrotentipides</i> sp.	0	1	0	0	0	0	1	0	0	3	0	0	0	3	4
<i>Dreissena polymorpha</i>	0	0	0	0	0	0	0	0	0	0	0	0	6	6	6
Empididae	1	0	0	0	0	0	1	0	0	0	0	0	0	0	1
Enchytraeidae	0	0	0	0	1	0	1	0	0	0	0	0	0	0	1
Ephemeroptera	0	0	0	0	0	0	0	0	0	0	0	1	0	1	1
<i>Gammarus</i> sp.	0	0	1	0	0	0	1	0	6	0	0	0	0	6	7
<i>Goniobasis</i> sp.	0	0	0	0	0	0	0	1	3	1	1	0	2	8	8
<i>Hexagenia</i> sp.	0	0	0	4	7	0	11	0	0	0	1	0	0	1	12
Immature tubificid without	27	4	63	7	13	10	124	16	0	15	9	452	14	506	630
<i>Limnodrilus cervix</i>	1	0	0	0	0	0	1	0	0	0	0	0	0	0	1
<i>Limnodrilus hoffmeisteri</i>	2	0	9	4	5	7	27	1	0	1	2	66	5	75	102
<i>Limnodrilus maumeensis</i>	0	0	0	0	1	4	5	0	0	0	0	19	0	19	24
<i>Limnodrilus profundicola</i>	0	0	2	0	0	6	8	0	0	0	0	2	0	2	10
Naididae	0	0	1	0	0	0	1	0	0	0	0	0	0	0	1
<i>Nais variabilis</i>	0	34	16	0	2	0	52	0	0	3	0	1	0	4	56
<i>Paranais frici</i>	0	0	0	0	0	0	0	3	0	2	0	0	0	5	5
<i>Paratendipes</i> sp.	0	0	0	1	0	0	1	0	0	0	0	0	0	0	1
<i>Pisidium</i> sp.	0	2	9	1	0	0	12	2	1	60	15	8	2	88	100
<i>Polypedilum</i> sp.	3	0	1	2	2	1	9	13	0	15	0	1	2	31	40
<i>Potamothrix vejdvovskiyi</i>	0	0	1	2	3	0	6	0	0	0	0	1	1	2	8
<i>Pristina osborni</i>	0	0	0	0	0	0	0	0	0	0	0	3	0	3	3
<i>Proboezzia</i> sp.	1	0	0	0	4	0	5	0	0	0	0	6	0	6	11
<i>Procladius</i> sp.	0	0	0	1	0	0	1	0	1	1	0	5	0	7	8
<i>Specaria josinae</i>	0	0	0	0	0	0	0	0	0	0	0	1	0	1	1
<i>Stempellina</i> sp.	0	1	0	1	0	0	2	0	0	1	0	0	0	1	3
<i>Tanytarsus</i> sp.	0	0	0	0	0	0	0	0	0	6	0	1	0	7	7
Tubificidae	4	0	0	0	0	0	4	0	0	0	0	1	0	1	5
<i>Xenochironomus</i> sp.	0	0	0	0	0	0	0	0	2	0	0	0	0	2	2
Monthly Total	39	45	108	42	40	28	302	39	29	158	37	599	52	914	1216

TABLE 5.4

**MEAN NUMBER OF MACROINVERTEBRATES (NUMBER/M<sup>2</sup>) AND PERCENT COMPOSITION  
OF OLIGOCHAETA, CHIRONOMIDAE, MOLLUSCA, AND OTHER ORGANISMS, 2007 BVPS**

May	1 (Control)		2A		2B1 (Non-control)		2B2 (Non-control)		2B3 (Non-control)		3		Total Mean	
	#/m <sup>2</sup>	%	#/m <sup>2</sup>	%	#/m <sup>2</sup>	%	#/m <sup>2</sup>	%	#/m <sup>2</sup>	%	#/m <sup>2</sup>	%	#/m <sup>2</sup>	%
Oligochaeta	487	87	545	84	1347	87	186	30	358	62	387	93	552	76
Chironomidae	43	8	58	9	57	4	344	56	57	10	14	3	98	13
Mollusca	0	0	43	7	129	8	14	2	0	0	0	0	29	4
Others	29	5	0	0	14	1	72	12	158	28	14	3	48	7
<b>Total</b>	<b>559</b>	<b>100</b>	<b>646</b>	<b>100</b>	<b>1547</b>	<b>100</b>	<b>616</b>	<b>100</b>	<b>573</b>	<b>100</b>	<b>415</b>	<b>100</b>	<b>727</b>	<b>100</b>

September	1 (Control)		2A		2B1 (Non-control)		2B2 (Non-control)		2B3 (Non-control)		3		Total Mean	
	#/m <sup>2</sup>	%	#/m <sup>2</sup>	%	#/m <sup>2</sup>	%	#/m <sup>2</sup>	%	#/m <sup>2</sup>	%	#/m <sup>2</sup>	%	#/m <sup>2</sup>	%
Oligochaeta	244	44	0	0	344	15	158	30	7855	91	330	44	1488	68
Chironomidae	244	44	72	17	889	39	57	11	387	5	100	13	291	13
Mollusca	72	13	258	62	1018	45	301	57	143	2	315	42	351	16
Others	0	0	86	21	14	1	14	3	201	2	0	0	53	2
<b>Total</b>	<b>560</b>	<b>100</b>	<b>416</b>	<b>100</b>	<b>2265</b>	<b>100</b>	<b>530</b>	<b>100</b>	<b>8586</b>	<b>100</b>	<b>745</b>	<b>100</b>	<b>2183</b>	<b>100</b>

TABLE 5.5

MEAN NUMBER OF MACROINVERTEBRATES (NUMBER/M<sup>2</sup>) AND PERCENT COMPOSITION OF OLIGOCHAETA, CHIRONOMIDAE, MOLLUSCA, AND OTHER ORGANISMS FOR THE CONTROL STATION (1) AND THE AVERAGE FOR NON-CONTROL STATIONS ( 2B1, 2B2, AND 2B3), 2007 BVPS

May	Control Station (Mean)		Non-Control Station (Mean)	
	#/m <sup>2</sup>	%	#/m <sup>2</sup>	%
Oligochaeta	487	87	630	69
Chironomidae	43	8	153	17
Mollusca	0	0	48	5
Others	29	5	81	9
TOTAL	559	100	912	100

September	Control Station (Mean)		Non-Control Station (Mean)	
	#/m <sup>2</sup>	%	#/m <sup>2</sup>	%
Oligochaeta	244	44	2786	73
Chironomidae	244	44	444	12
Mollusca	72	13	487	13
Others	0	0	76	2
TOTAL	560	100	3793	100

TABLE 5.6

SHANNON-WEINER DIVERSITY, EVENNESS AND RICHNESS INDICES  
FOR BENTHIC MACROINVERTEBRATES COLLECTED IN THE OHIO RIVER, 2007

May	Station					
	1	2A	2B1	2B2	2B3	3
No. of Taxa	7	7	11	11	11	5
Shannon-Weiner Index	0.49	0.42	0.62	0.83	0.88	0.63
Evenness	0.58	0.49	0.60	0.79	0.84	0.90
Richness	1.34	1.05	2.56	3.21	3.52	1.80

September	Station					
	1	2A	2B1	2B2	2B3	3
No. of Taxa	8	8	21	11	21	11
Shannon-Weiner Index	0.66	0.68	0.95	0.79	0.45	0.94
Evenness	0.73	0.75	0.72	0.76	0.34	0.90
Richness	0.55	2.38	1.98	3.32	2.35	2.53

**Table 5.7. Benthic Macroinvertebrate Densities for Stations 1 (Control) and 2B (Noncontrol), BVPS, 1973-2007.**

	Preoperational					
	1973		1974		1975	
	1	2B	1	2B	1	2B
May	248	508	1116	2197		
August	99	244	143	541	1017	1124
<b>Mean</b>	<b>173</b>	<b>376</b>	<b>630</b>	<b>1369</b>	<b>1017</b>	<b>1124</b>

	Operational					
	1976		1977		1978	
	1	2B	1	2B	1	2B
May	927	3660	674	848	351	126
August	851	785	591	3474	601	1896
<b>Mean</b>	<b>889</b>	<b>2223</b>	<b>633</b>	<b>2161</b>	<b>476</b>	<b>1011</b>

	Operational					
	1979		1980		1981	
	1	2B	1	2B	1	2B
May	1004	840	1041	747	209	456
Aug/Sept	1185	588	1523	448	2185	912
<b>Mean</b>	<b>1095</b>	<b>714</b>	<b>1282</b>	<b>598</b>	<b>1197</b>	<b>684</b>

	Operational					
	1982		1983		1984	
	1	2B	1	2B	1	2B
May	3490	3026	3590	1314	2741	621
September	2958	3364	4172	4213	1341	828
<b>Mean</b>	<b>3223</b>	<b>3195</b>	<b>3881</b>	<b>2764</b>	<b>2041</b>	<b>725</b>

	Operational					
	1985		1986		1987	
	1	2B	1	2B	1	2B
May	2256	867	601	969	1971	2649
September	1024	913	849	943	2910	2780
<b>Mean</b>	<b>1640</b>	<b>890</b>	<b>725</b>	<b>956</b>	<b>2440</b>	<b>2714</b>

**Table 5.7. Benthic Macroinvertebrate Densities for Stations 1 (Control) and 2B (Noncontrol), BVPS, 1973-2007 (Continued).**

	Operational					
	1988		1989		1990	
	1	2B	1	2B	1	2B
May	1804	1775	3459	2335	15135	5796
September	1420	1514	1560	4707	5550	1118
<b>Mean</b>	<b>1612</b>	<b>1645</b>	<b>2510</b>	<b>3274</b>	<b>10343</b>	<b>3457</b>

	Operational					
	1991		1992		1993	
	1	2B	1	2B	1	2B
May	7760	6355	7314	10560	8435	2152
September	3588	2605	2723	4707	4693	2143
<b>Mean</b>	<b>5808</b>	<b>4480</b>	<b>5019</b>	<b>7634</b>	<b>6564</b>	<b>2148</b>

	Operational					
	1994		1995		1996	
	1	2B	1	2B	1	2B
May	6980	2349	8083	9283	1987	1333
September	1371	2930	1669	3873	1649	2413
<b>Mean</b>	<b>4176</b>	<b>2640</b>	<b>4876</b>	<b>6578</b>	<b>1814</b>	<b>1873</b>

	Operational					
	1997		1998		1999	
	1	2B	1	2B	1	2B
May	1411	2520	6980	2349	879	1002
September	1944	2774	1371	2930	302	402
<b>Mean</b>	<b>1678</b>	<b>2647</b>	<b>4176</b>	<b>2640</b>	<b>591</b>	<b>702</b>

	Operational					
	2000		2001		2002	
	1	2B	1	2B	1	2B
May	2987	2881	3139	5232	1548	2795
September	3092	2742			8632	14663
<b>Mean</b>	<b>3040</b>	<b>2812</b>	<b>3139</b>	<b>5232</b>	<b>5090</b>	<b>8729</b>



**Table 5.7. Benthic Macroinvertebrate Densities for Stations 1 (Control) and 2B (Noncontrol), BVPS, 1973-2007 (Continued).**

	Operational					
	2003		2004		2005	
	1	2B	1	2B	1	2B
May	7095	10750	2752	4558	516	1146
September	2193	6464	10062	7604	4773	6435
<b>Mean</b>	<b>4644</b>	<b>8607</b>	<b>6407</b>	<b>6121</b>	<b>2645</b>	<b>3791</b>

	Operational					
	2006		2007			
	1	2B	1	2B	1	2B
May	143	1242	559	912	---	---
September	229	2199	560	3794	---	---
<b>Mean</b>	<b>186</b>	<b>1721</b>	<b>560</b>	<b>2353</b>	---	---

TABLE 5.8

TOTAL FISH CATCH; ELECTROFISHING AND SEINE NET  
COMBINED DURING THE BVPS 2007 FISHERIES SURVEY

Common Name	Scientific Name	Number	Percent
Smallmouth buffalo	<i>Ictiobus bubalus</i>	26	12.04
Black crappie	<i>Pomoxis nigromaculatus</i>	1	0.46
Bluegill	<i>Lepomis macrochirus</i>	3	1.39
Carp	<i>Cyprinus carpio</i>	6	2.78
Channel catfish	<i>Ictalurus punctatus</i>	2	0.93
Emerald shiner	<i>Notropis atherinoides</i>	2	0.93
Flathead catfish	<i>Pylodictis olivaris</i>	2	0.93
Freshwater drum	<i>Aplodinotus grunniens</i>	2	0.93
Gizzard shad	<i>Dorosoma cepedianum</i>	24	11.11
Golden redhorse sucker	<i>Moxostoma erythrurum</i>	21	9.72
Log perch	<i>Percina caprodes</i>	2	0.93
Longnose gar	<i>Lepisosteus osseus</i>	7	3.24
Mooneye	<i>Hiodon tergisus</i>	2	0.93
Northern hog sucker	<i>Hypentelium nigricans</i>	1	0.46
Quillback	<i>Carpiodes cyprinus</i>	2	0.93
River carpsucker	<i>Carpiodes carpio</i>	1	0.46
Rock bass	<i>Ambloplites rupestris</i>	1	0.46
Sauger	<i>Sander canadense</i>	22	10.19
Shorthead redhorse sucker	<i>Moxostoma macrolepidotum</i>	27	12.50
Silver redhorse	<i>Moxostoma anisurum</i>	5	2.31
Smallmouth bass	<i>Micropterus dolomieu</i>	31	14.35
Spotted bass	<i>Micropterus punctulatus</i>	18	8.33
Walleye	<i>Sander vitreum</i>	3	1.39
White bass	<i>Morone chrysops</i>	5	2.31
<b>Total Fish Collected in 2007</b>		<b>216</b>	<b>100.00</b>

**TABLE 5.9**

**COMPARISON OF CONTROL VS. NON-CONTROL ELECTROFISHING CATCHES  
DURING THE BVPS 2007 FISHERIES SURVEY**

<b>Common Name</b>	<b>Control</b>	<b>%</b>	<b>Non-control</b>	<b>%</b>	<b>Total fish</b>	<b>%</b>
Smallmouth buffalo	5	8.62	18	13.7	23	12.17
Black crappie	1	1.72			1	0.53
Bluegill	2	3.45	1	0.8	3	1.59
Carp	1	1.72	5	3.8	6	3.17
Channel catfish	2	3.45			2	1.06
Flathead catfish			2	1.5	2	1.06
Freshwater drum			2	1.5	2	1.06
Gizzard shad	5	8.62	13	9.9	18	9.52
Golden redhorse sucker	7	12.07	14	10.7	21	11.11
Log perch			1	0.8	1	0.53
Longnose gar	1	1.72	6	4.6	7	3.70
Mooneye	1	1.72	1	0.8	2	1.06
Northern hog sucker			1	0.8	1	0.53
Quillback			2	1.5	2	1.06
River carpsucker	1	1.72			1	0.53
Rock bass			1	0.8	1	0.53
Sauger	9	15.52	13	9.9	22	11.64
Shorthead redhorse sucker	6	10.34	19	14.5	25	13.23
Silver redhorse	2	3.45	3	2.3	5	2.65
Smallmouth bass	8	13.79	18	13.7	26	13.76
Spotted bass	7	12.07	4	3.1	11	5.82
Walleye			3	2.3	3	1.59
White bass			4	3.1	4	2.12
<b>Electrofishing Totals</b>	<b>58</b>	<b>100.00</b>	<b>131</b>	<b>100</b>	<b>189</b>	<b>100.00</b>

TABLE 5.10

COMPARISON OF CONTROL VS. NON-CONTROL SEINE CATCHES  
DURING THE BVPS 2007 FISHERIES SURVEY

Common Name	Control	%	Non-control	%	Total fish	%
Smallmouth buffalo		0.00	3	14.29	3	11.11
Emerald shiner	1	16.67	1	4.76	2	7.41
Gizzard shad		0.00	6	28.57	6	22.22
Log perch		0.00	1	4.76	1	3.70
Shorthead redhorse sucker	2	33.33		0.00	2	7.41
Smallmouth bass	1	16.67	4	19.05	5	18.52
Spotted bass	2	33.33	5	23.81	7	25.93
White bass		0.00	1	4.76	1	3.70
<b>Seine Totals</b>	<b>6</b>	<b>100.00</b>	<b>21</b>	<b>100.00</b>	<b>27</b>	<b>100.00</b>

TABLE 5.11

FISH SPECIES COLLECTED DURING THE MAY 2007 SAMPLING  
OF THE OHIO RIVER IN THE VICINITY OF BVPS

Common Name	Sample locations *						Seine		Electrofishing	
	S-1	S-2	E-1	E-2A	E-2B	E-3	Total	%	Total	%
Smallmouth buffalo	-	-	2	2	-	1	0	-	5	10.00
Black crappie	-	-	1	-	-	-	0	-	1	2.00
Bluegill	-	-	1	-	-	-	0	-	1	2.00
Carp	-	-	-	-	-	-	0	-	0	0.00
Channel catfish	-	-	-	-	-	-	0	-	0	0.00
Emerald shiner	-	-	-	-	-	-	0	-	0	0.00
Flathead catfish	-	-	-	-	-	-	0	-	0	0.00
Freshwater drum	-	-	-	-	-	-	0	-	0	0.00
Gizzard shad	-	-	-	2	-	1	0	-	3	6.00
Golden redhorse sucker	-	-	6	5	4	1	0	-	16	32.00
Log perch	-	-	-	-	-	-	0	-	0	0.00
Longnose gar	-	-	-	-	1	-	0	-	1	2.00
Mooneye	-	-	-	-	-	-	0	-	0	0.00
Northern hog sucker	-	-	-	1	-	-	0	-	1	2.00
Quillback	-	-	-	-	-	-	0	-	0	0.00
River carpsucker	-	-	-	-	-	-	0	-	0	0.00
Rock bass	-	-	-	-	1	-	0	-	1	2.00
Sauger	-	-	-	1	2	3	0	-	6	12.00
Shorthead redhorse sucker	-	-	-	3	-	4	0	-	7	14.00
Silver redhorse	-	-	1	2	-	-	0	-	3	6.00
Smallmouth bass	-	-	1	-	1	-	0	-	2	4.00
Spotted bass	-	-	2	-	-	-	0	-	2	4.00
Walleye	-	-	-	-	-	-	0	-	0	0.00
White bass	-	-	-	1	-	-	0	-	1	2.00
<b>Total</b>	<b>0</b>	<b>0</b>	<b>14</b>	<b>17</b>	<b>9</b>	<b>10</b>	<b>0</b>	<b>-</b>	<b>50</b>	<b>100.00</b>

\* Gear = (E) Fish captured by electrofishing; (S) captured by seining

TABLE 5.12

**FISH SPECIES COLLECTED DURING THE JULY 2007 SAMPLING  
OF THE OHIO RIVER IN THE VICINITY OF BVPS**

Common Name	Sample locations *						Seine		Electrofishing	
	S-1	S-2	E-1	E-2A	E-2B	E-3	Total	%	Total	%
Smallmouth buffalo	-	3	1	1	1	2	3	15.79	5	13.51
Black crappie	-	-	-	-	-	-	0	0.00	0	0.00
Bluegill	-	-	-	-	-	-	0	0.00	0	0.00
Carp	-	-	1	-	4	-	0	0.00	5	13.51
Channel catfish	-	-	-	-	-	-	0	0.00	0	0.00
Emerald shiner	1	1	-	-	-	-	2	10.53	0	0.00
Flathead catfish	-	-	-	-	1	-	0	0.00	1	2.70
Freshwater drum	-	-	-	-	-	-	0	0.00	0	0.00
Gizzard shad	-	6	5	5	2	-	6	31.58	12	32.43
Golden redhorse sucker	-	-	-	-	-	-	0	0.00	0	0.00
Log perch	-	1	-	-	-	-	1	5.26	0	0.00
Longnose gar	-	-	-	-	-	-	0	0.00	0	0.00
Mooneye	-	-	1	-	-	-	0	0.00	1	2.70
Northern hog sucker	-	-	-	-	-	-	0	0.00	0	0.00
Quillback	-	-	-	1	-	-	0	0.00	1	2.70
River carpsucker	-	-	-	-	-	-	0	0.00	0	0.00
Rock bass	-	-	-	-	-	-	0	0.00	0	0.00
Sauger	-	-	1	-	-	-	0	0.00	1	2.70
Shorthead redhorse sucker	-	-	2	1	-	-	0	0.00	3	8.11
Silver redhorse	-	-	1	-	-	-	0	0.00	1	2.70
Smallmouth bass	-	1	-	2	2	-	1	5.26	4	10.81
Spotted bass	1	4	-	2	-	-	5	26.32	2	5.41
Walleye	-	-	-	1	-	-	0	0.00	1	2.70
White bass	-	1	-	-	-	-	1	5.26	0	0.00
<b>Total</b>	<b>2</b>	<b>17</b>	<b>12</b>	<b>13</b>	<b>10</b>	<b>2</b>	<b>19</b>	<b>100.00</b>	<b>37</b>	<b>100.00</b>

\* Gear = (E) Fish captured by electrofishing; (S) captured by seining

TABLE 5.13

FISH SPECIES COLLECTED DURING THE SEPTEMBER 2007 SAMPLING  
OF THE OHIO RIVER IN THE VICINITY OF BVPS

Common Name	Sample locations*						Seine		Electrofishing	
	S-1	S-2	E-1	E-2A	E-2B	E-3	Total	%	Total	%
Smallmouth buffalo	-	-	-	-	3	4	0	0.00	7	20.00
Black crappie	-	-	-	-	-	-	0	0.00	0	0.00
Bluegill	-	-	1	-	-	-	0	0.00	1	2.86
Carp	-	-	-	-	-	-	0	0.00	0	0.00
Channel catfish	-	-	2	-	-	-	0	0.00	2	5.71
Emerald shiner	-	-	-	-	-	-	0	0.00	0	0.00
Flathead catfish	-	-	-	-	1	-	0	0.00	1	2.86
Freshwater drum	-	-	-	1	1	-	0	0.00	2	5.71
Gizzard shad	-	-	-	-	1	-	0	0.00	1	2.86
Golden redhorse sucker	-	-	-	1	-	-	0	0.00	1	2.86
Log perch	-	-	-	1	-	-	0	0.00	1	2.86
Longnose gar	-	-	1	-	4	-	0	0.00	5	14.29
Mooneye	-	-	-	-	-	-	0	0.00	0	0.00
Northern hog sucker	-	-	-	-	-	-	0	0.00	0	0.00
Quillback	-	-	-	-	-	-	0	0.00	0	0.00
River carpsucker	-	-	-	-	-	-	0	0.00	0	0.00
Rock bass	-	-	-	-	-	-	0	0.00	0	0.00
Sauger	-	-	1	-	1	-	0	0.00	2	5.71
Shorthead redhorse sucker	2	-	-	3	-	-	2	25.00	3	8.57
Silver redhorse	-	-	-	1	-	-	0	0.00	1	2.86
Smallmouth bass	1	3	4	1	1	1	4	50.00	7	20.00
Spotted bass	1	1	-	-	-	-	2	25.00	0	0.00
Walleye	-	-	-	-	-	-	0	0.00	0	0.00
White bass	-	-	-	-	-	1	0	0.00	1	2.86
<b>Total</b>	<b>4</b>	<b>4</b>	<b>9</b>	<b>8</b>	<b>12</b>	<b>6</b>	<b>8</b>	<b>100.00</b>	<b>35</b>	<b>100.00</b>

\* Gear = (E) Fish captured by electrofishing; (S) captured by seining

TABLE 5.14

**FISH SPECIES COLLECTED DURING THE NOVEMBER 2007 SAMPLING  
OF THE OHIO RIVER IN THE VICINITY OF BVPS**

Common Name	Sample locations *						Seine		Electrofishing	
	S-1	S-2	E-1	E-2A	E-2B	E-3	Total	%	Total	%
Smallmouth buffalo	-	-	2	2	-	2	0	-	6	8.96
Black crappie	-	-	-	-	-	-	0	-	0	0.00
Bluegill	-	-	-	1	-	-	0	-	1	1.49
Carp	-	-	-	-	1	-	0	-	1	1.49
Channel catfish	-	-	-	-	-	-	0	-	0	0.00
Emerald shiner	-	-	-	-	-	-	0	-	0	0.00
Flathead catfish	-	-	-	-	-	-	0	-	0	0.00
Freshwater drum	-	-	-	-	-	-	0	-	0	0.00
Gizzard shad	-	-	-	-	1	1	0	-	2	2.99
Golden redhorse sucker	-	-	1	-	2	1	0	-	4	5.97
Log perch	-	-	-	-	-	-	0	-	0	0.00
Longnose gar	-	-	-	-	1	-	0	-	1	1.49
Mooneye	-	-	-	1	-	-	0	-	1	1.49
Northern hog sucker	-	-	-	-	-	-	0	-	0	0.00
Quillback	-	-	-	-	1	-	0	-	1	1.49
River carpsucker	-	-	1	-	-	-	0	-	1	1.49
Rock bass	-	-	-	-	-	-	0	-	0	0.00
Sauger	-	-	7	-	4	2	0	-	13	19.40
Shorthead redhorse sucker	-	-	4	3	2	3	0	-	12	17.91
Silver redhorse	-	-	-	-	-	-	0	-	0	0.00
Smallmouth bass	-	-	3	3	4	3	0	-	13	19.40
Spotted bass	-	-	5	1	1	-	0	-	7	10.45
Walleye	-	-	-	1	-	1	0	-	2	2.99
White bass	-	-	-	-	1	1	0	-	2	2.99
<b>Total</b>	<b>0</b>	<b>0</b>	<b>23</b>	<b>12</b>	<b>18</b>	<b>14</b>	<b>0</b>	<b>-</b>	<b>67</b>	<b>100.00</b>

\* Gear = (E) Fish captured by electrofishing; (S) captured by seining



TABLE 5.15

ESTIMATED NUMBER OF FISH OBSERVED \* DURING  
ELECTROFISHING OPERATIONS, 2007

Common Name	May	July	Sept	Nov	Total
Unidentified suckers			2	6	8
Carp		6			6
Smallmouth buffalo		3	2	2	7
Gizzard shad		100's			100s
Walleye/sauger		1		1	2
Smallmouth/spotted bass				3	3
Longnose gar				1	1
Total	0	10	2	13	25+100's

\* = Not boated or handled

**Table 5.16**

**CATCH PER UNIT EFFORT (CPUE AS FISH/ELECTROFISHING  
MINUTE)  
BY SEASON DURING THE BVPS 2004 FISHERIES SURVEY**

Season	Effort (min)	Common Name	Count of species	CPUE (fish/min)
Spring	40.7	Black buffalo	7	0.172
		Bluegill	0	0.000
		Channel catfish	6	0.147
		Common carp	1	0.025
		Emerald shiner	0	0.000
		Flathead catfish	2	0.049
		Freshwater drum	3	0.074
		Gizzard shad	4	0.098
		Golden redhorse sucker	16	0.393
		Longnose gar	1	0.025
		Mooneye	1	0.025
		Northern hogsucker	1	0.025
		Quillback	5	0.123
		Sauger	5	0.123
		Shorthead redhorse sucker	10	0.246
		Silver redhorse	3	0.074
		Smallmouth bass	1	0.025
		Striped bass hybrid	0	0.000
		Walleye	2	0.049
White bass	2	0.049		
Season Total			70	1.720
Season	Effort (min)	Common Name	Count of species	CPUE (fish/min)
Summer	40.4	Black buffalo	7	0.1733
		Bluegill	0	0.0000
		Channel catfish	3	0.0743
		Common carp	1	0.0248
		Emerald shiner	1	0.0248
		Flathead catfish	1	0.0248
		Freshwater drum	1	0.0248
		Gizzard shad	1	0.0248
		Golden redhorse sucker	7	0.1733
		Longnose gar	0	0.0000
		Mooneye	1	0.0248
		Northern hogsucker	0	0.0000
		Quillback	2	0.0495
		Sauger	0	0.0000
		Shorthead redhorse sucker	4	0.0990
		Silver redhorse	6	0.1485
		Smallmouth bass	1	0.0248
		Striped bass hybrid	0	0.0000
		Walleye	1	0.0248
White bass	1	0.0248		
Season Total			38	0.9406

Table 5.16 Continued

Season	Effort (min)	Common Name	Number Collected	CPUE (fish/min)
Fall	40.70	Black Buffalo	1	0.0248
		Bluegill	0	0.0000
		Channel catfish	0	0.0000
		Common carp	1	0.0248
		Emerald shiner	0	0.0000
		Flathead catfish	0	0.0000
		Freshwater drum	4	0.0990
		Gizzard shad	2	0.0495
		Golden redhorse sucker	9	0.2228
		Longnose gar	0	0.0000
		Mooneye	0	0.0000
		Northern hogsucker	0	0.0000
		Quillback	1	0.0248
		Sauger	5	0.1238
		Shorthead redhorse sucker	1	0.0248
		Silver redhorse	1	0.0248
		Smallmouth bass	1	0.0248
		Striped bass hybrid	0	0.0000
		Walleye	4	0.0990
		White catfish	0	0.0000
White bass	0	0.0000		
Season Total			30	0.7426
Season	Effort (min)	Common Name	Number Collected	CPUE (fish/min)
Winter	40.40	Black Buffalo	17	0.4208
		Bluegill	0	0.0000
		Channel catfish	1	0.0248
		Common carp	0	0.0000
		Emerald shiner	0	0.0000
		Flathead catfish	0	0.0000
		Freshwater drum	3	0.0743
		Gizzard shad	6	0.1485
		Golden redhorse sucker	8	0.1980
		Longnose gar	1	0.0248
		Mooneye	4	0.0990
		Northern hogsucker	0	0.0000
		Quillback	3	0.0743
		Sauger	3	0.0743
		Shorthead redhorse sucker	8	0.1980
		Silver redhorse	2	0.0495
		Smallmouth bass	0	0.0000
		Striped bass hybrid	2	0.0495
		Walleye	10	0.2475
		White catfish	0	0.0000
White bass	1	0.0248		
Season Total			69	1.7079
162.20		207		1.2762

**Table 5.17**

**CATCH PER UNIT EFFORT (CPUE AS FISH/ELECTROFISHING MINUTE)  
BY SEASON DURING THE BVPS 2005 FISHERIES SURVEY**

Season	Effort (min)	Common Name	Number Collected	CPUE (fish/min)
Spring	41	Black buffalo	5	0.1220
		Black crappie	1	0.0244
		Channel catfish	3	0.0732
		Freshwater drum	1	0.0244
		Gizzard shad	8	0.1951
		Golden redhorse sucker	19	0.4634
		Highfin carpsucker	1	0.0244
		Longnose gar	2	0.0488
		Quillback	1	0.0244
		Sauger	3	0.0732
		Shorthead redhorse sucker	4	0.0976
		Silver redhorse	3	0.0732
		Smallmouth bass	1	0.0244
		Spotfin shiner	1	0.0244
		Walleye	1	0.0244
		Season Total	54	1.3171
Season	Effort (min)	Common Name	Number Collected	CPUE (fish/min)
Summer	40	Black buffalo	1	0.0250
		Channel catfish	1	0.0250
		Flathead catfish	1	0.0250
		Gizzard shad	14	0.3500
		Golden redhorse sucker	5	0.1250
		Longnose gar	1	0.0250
		Mooneye	1	0.0250
		Shorthead redhorse sucker	4	0.1000
		Smallmouth bass	1	0.0250
		White bass	2	0.0500
		Season Total	31	0.7750

**Table 5.17 continued**

Season	Effort (min)	Common Name	Number Collected	CPUE (fish/min)
Fall	40	Black Buffalo	4	0.1000
		Channel catfish	1	0.0250
		Flathead catfish	2	0.0500
		Gizzard shad	24	0.6000
		Golden redhorse	3	0.0750
		Mooneye	1	0.0250
		Quillback	4	0.1000
		Sauger	2	0.0500
		Shorthead redhorse sucker	6	0.1500
		Silver redhorse	2	0.0500
		Smallmouth bass	1	0.0250
		Walleye	1	0.0250
		White bass	25	0.6250
				Season Total
Season	Effort (min)	Common Name	Number Collected	CPUE (fish/min)
Winter	41	Black buffalo	12	0.2927
		Bluegill	2	0.0488
		Common carp	2	0.0488
		Creek chub	2	0.0488
		Freshwater drum	1	0.0244
		Gizzard shad	18	0.4390
		Golden redhorse sucker	8	0.1951
		Mooneye	4	0.0976
		Quillback	8	0.1951
		River carsucker	2	0.0488
		Sauger	14	0.3415
		Shorthead redhorse sucker	20	0.4878
		Silver redhorse	1	0.0244
		Smallmouth bass	5	0.1220
		Spotted bass	2	0.0488
Walleye	2	0.0488		
White bass	17	0.4146		
		Season Total	120	2.9268
	162		281	1.7346

**Table 5.18**

**CATCH PER UNIT EFFORT (CPUE AS FISH/ELECTROFISHING MINUTE)  
BY SEASON DURING THE BVPS 2006 FISHERIES SURVEY**

Season	Effort (min)	Common Name	Number Collected	CPUE (fish/min)
Spring	41	Black buffalo	10	0.2439
		Black crappie	1	0.0244
		Gizzard shad	9	0.2195
		Golden redhorse sucker	19	0.4634
		Quillback	2	0.0488
		River carpsucker	0	0.0000
		Sauger	17	0.4146
		Shorthead redhorse sucker	16	0.3902
		Skipjack herring	1	0.0244
		Smallmouth bass	11	0.2683
		Spotted bass	5	0.1220
		Striped bass x White bass	1	0.0244
		Walleye	10	0.2439
		White bass	15	0.3659
		Season Total	117	2.8537
Season	Effort (min)	Common Name	Number Collected	CPUE (fish/min)
Summer	40	Black buffalo	2	0.0500
		Black crappie	1	0.0250
		Gizzard shad	3	0.0750
		Golden redhorse sucker	4	0.1000
		Highfin carpsucker	0	0.0000
		Longnose gar	2	0.0500
		Shorthead redhorse sucker	12	0.3000
		Silver redhorse	1	0.0250
		Smallmouth bass	3	0.0750
		Season Total	28	0.7000

**Table 5.18 Continued**

Season	Effort (min)	Common Name	Number Collected	CPUE (fish/min)
Fall	40	Black buffalo	3	0.0732
		Bluegill	1	0.0244
		Carp	3	0.0732
		Gizzard shad	5	0.1220
		Golden redhorse sucker	2	0.0488
		Longnose gar	4	0.0976
		Mooneye	1	0.0244
		Quillback	2	0.0488
		Sauger	5	0.1220
		Shorthead redhorse sucker	14	0.3415
		Silver redhorse	1	0.0244
		Smallmouth bass	2	0.0488
		Spotted bass	1	0.0244
		White bass	1	0.0244
		Season Total	45	1.1250
Season	Effort (min)	Common Name	Number Collected	CPUE (fish/min)
Winter	40	Black buffalo	18	0.4500
		Carp	1	0.0250
		Channel catfish	1	0.0250
		Flathead catfish	1	0.0250
		Freshwater drum	4	0.1000
		Gizzard shad	18	0.4500
		Golden redhorse sucker	2	0.0500
		Mooneye	3	0.0750
		Northern hog sucker	0	0.0000
		Quillback	6	0.1500
		River carsucker	1	0.0250
		Shorthead redhorse sucker	11	0.2750
		Walleye	1	0.0250
		White bass	1	0.0250
		Season Total	68	1.7000
	161		258	1.6025

**Table 5.19**

**CATCH PER UNIT EFFORT (CPUE AS FISH/ELECTROFISHING MINUTE)  
BY SEASON DURING THE BVPS 2007 FISHERIES SURVEY**

Season	Effort (min)	Common Name	Count of species	CPUE (fish/min)
Spring	41	Smallmouth buffalo	5	0.122
		Black crappie	1	0.024
		Bluegill	1	0.024
		Gizzard shad	3	0.073
		Golden redhorse sucker	16	0.390
		Longnose gar	1	0.024
		Northern hog sucker	1	0.024
		Rock bass	1	0.024
		Sauger	6	0.146
		Shorthead redhorse sucker	7	0.171
		Silver redhorse	3	0.073
		Smallmouth bass	2	0.049
		Spotted bass	2	0.049
White bass	1	0.024		
Season Total			50	1.220
Season	Effort (min)	Common Name	Count of species	CPUE (fish/min)
Summer	41	Smallmouth buffalo	5	0.1220
		Carp	5	0.1220
		Flathead catfish	1	0.0244
		Gizzard shad	12	0.2927
		Mooneye	1	0.0244
		Quillback	1	0.0244
		Sauger	1	0.0244
		Shorthead redhorse sucker	3	0.0732
		Silver redhorse	1	0.0244
		Smallmouth bass	4	0.0976
		Spotted bass	2	0.0488
Walleye	1	0.0244		
Season Total			37	0.9024



**Table 5.19 (Cont'd)**  
**CATCH PER UNIT EFFORT (CPUE AS FISH/ELECTROFISHING MINUTE)**  
**BY SEASON DURING THE BVPS 2007 FISHERIES SURVEY**

Season	Effort (min)	Common Name	Number Collected	CPUE (fish/min)
Fall	40.00	Smallmouth buffalo	7	0.1707
		Bluegill	1	0.0244
		Channel catfish	2	0.0488
		Flathead catfish	1	0.0244
		Freshwater drum	2	0.0488
		Gizzard shad	1	0.0244
		Golden redhorse sucker	1	0.0244
		Log perch	1	0.0244
		Longnose gar	5	0.1220
		Sauger	2	0.0488
		Shorthead redhorse sucker	3	0.0732
		Silver redhorse	1	0.0244
		Smallmouth bass	7	0.1707
		White bass	1	0.0244
Season Total			35	0.8537
Season	Effort (min)	Common Name	Number Collected	CPUE (fish/min)
Winter	40.00	Smallmouth buffalo	6	0.1463
		Bluegill	1	0.0244
		Carp	1	0.0244
		Gizzard shad	2	0.0488
		Golden redhorse sucker	4	0.0976
		Longnose gar	1	0.0244
		Mooneye	1	0.0244
		Quillback	1	0.0244
		River carpsucker	1	0.0244
		Sauger	13	0.3171
		Shorthead redhorse sucker	12	0.2927
		Smallmouth bass	13	0.3171
		Spotted bass	7	0.1707
		Walleye	2	0.0488
		White bass	2	0.0488
Season Total			67	1.6341
	162.00		189	1.1667

TABLE 5.20

UNIT 1 COOLING RESERVOIR MONTHLY SAMPLING  
CORBICULA DENSITY DATA FOR  
 2007 FROM BVPS

Collection Date	Area sampled (sq ft)	Live or Dead	Count	Maximum Length Range (mm)	Minimum length Range(mm)	Estimated number (per sq m)
3/22/2007	0.25	Dead	4	1.00-1.99	0.001-0.99	172
		Live	2	2.00-3.34	1.00-1.99	86
4/23/2007	0.25	Dead	3	6.30-9.49	1.00-1.99	129
		Live	0	-	-	0
5/22/2007	0.25	Dead	0	-	-	0
		Live	2	2.00-3.34	2.00-3.34	86
6/13/2007	0.25	Dead	4	4.75-6.29	1.00-1.99	172
		Live	1	1.00-1.99	1.00-1.99	43
7/31/2007	0.25	Dead	0	-	-	0
		Live	1	4.75-6.29	4.75-6.29	43
8/27/2007	0.25	Dead	1	1.00-1.99	1.00-1.99	43
		Live	1	4.75-6.29	4.75-6.29	43
9/24/2007*	0.25	Dead	--	-	-	--
		Live	--	-	-	--
10/17/2007	0.25	Dead	1	2.00-3.34	2.00-3.34	43
		Live	0			0
11/8/2007	0.25	Dead	0	-	-	0
		Live	0	-	-	0
Unit summary		Dead	13	6.30-9.49	0.001-0.99	70
		Live	7	4.75-6.29	1.00-1.99	38

\*Cooling Tower dewatered; not sampled.


**TABLE 5.21**

**UNIT 2 COOLING RESERVOIR MONTHLY SAMPLING  
CORBICULA DENSITY DATA FOR  
2007 FROM BVPS**


<b>Collection Date</b>	<b>Area sampled (sq ft)</b>	<b>Live or Dead</b>	<b>Count</b>	<b>Maximum Length Range (mm)</b>	<b>Minimum length Range(mm)</b>	<b>Estimated number (per sq m)</b>
3/22/2007	0.25	Dead	0	---	---	0
		Live	0	---	---	0
4/23/2007	0.25	Dead	0	---	---	0
		Live	0	---	---	0
5/22/2007	0.25	Dead	1	1.00-1.99	1.00-1.99	43
		Live	3	2.00-3.34	1.00-1.99	129
6/13/2007	0.25	Dead	0	---	---	0
		Live	4	2.00-3.34	1.00-1.99	172
7/31/2007	0.25	Dead	1	1.00-1.99	1.00-1.99	43
		Live	3	1.00-1.99	1.00-1.99	129
8/27/2007	0.25	Dead	0	---	---	0
		Live	3	2.00-3.34	0.001-0.99	129
9/24/2007	0.25	Dead	0	---	---	0
		Live	9	3.35-4.74	1.00-1.99	387
10/17/2007	0.25	Dead	0	---	---	0
		Live	0	---	---	0
11/8/2007	0.25	Dead	0	---	---	0
		Live	0	---	---	0
Unit summary		Dead	2	1.00-1.99	1.00-1.99	10
		Live	22	3.35-4.74	0.001-0.99	118

**9.0**

**FIGURES**




**Figure 5.1** 2007 Beaver Valley Power Station Aquatic Monitoring Program Sampling Control and Non-Control Sampling Stations



ANNUAL ENVIRONMENTAL REPORT

Figure 5.2 Location Map for Beaver Valley Power Station Benthic Organism Survey Sampling Sites for the 2007 Study



ANNUAL ENVIRONMENTAL REPORT

Figure 5.3 Location Map for Beaver Valley Power Station Fish Population Survey Fish Sampling Sites for the 2007 Study

ANNUAL ENVIRONMENTAL REPORT





Figure 5.4 Location of Study Area, Beaver Valley Power Station Shippingport, Pennsylvania BVPS



Comparison of live Corbicula clam density estimates among 2007  
BVPS Unit 1 cooling tower reservoir events, for various clam shell groups.




	3/22	4/23	5/22	6/13	7/31	8/27	9/24	10/17	11/8
■ 0.01-0.99 mm	0	0	0	0	0	0	0	0	0
■ 1.00-1.99 mm	43	0	0	43	0	0	0	0	0
■ 2.00-3.34 mm	43	0	86	0	0	0	0	0	0
■ 3.35-4.74 mm	0	0	0	0	0	0	0	0	0
■ 4.75-6.29 mm	0	0	0	0	43	43	0	0	0
■ 6.30-9.49 mm	0	0	0	0	0	0	0	0	0
■ >9.50 mm	0	0	0	0	0	0	0	0	0
TOTAL #/m2	86	0	86	43	43	43	0	0	0

Note: Not Sampled in Sept due to Outage

Figure 5.5



Comparison of live Corbicula clam density estimates among 2007  
 BVPS Unit 1 cooling tower reservoir events, for various clam shell groups.




	3/22	4/23	5/22	6/13	7/31	8/27	9/24	10/17	11/8
■ 0.01-0.99 mm	0	0	0	0	0	0	0	0	0
■ 1.00-1.99 mm	43	0	0	43	0	0	0	0	0
■ 2.00-3.34 mm	43	0	86	0	0	0	0	0	0
■ 3.35-4.74 mm	0	0	0	0	0	0	0	0	0
■ 4.75-6.29 mm	0	0	0	0	43	43	0	0	0
■ 6.30-9.49 mm	0	0	0	0	0	0	0	0	0
■ >9.50 mm	0	0	0	0	0	0	0	0	0
TOTAL #/m2	86	0	86	43	43	43	0	0	0

Note: Not Sampled in Sept due to Outage

Figure 5.5



Comparison of live Corbicula clam density estimates among 2007 BVPS Intake Structure sample events, for various clam shell groups.



Intake structure bottom samples are collected from the Ohio River at the Intake Building.

Figure 5.7

**Water Temperature and River Elevation Recorded at the Ohio River at BVPS Intake Structure During 2007 on Monthly Sample Dates.**





Figure 5.8



### Density of zebra mussels veligers collected at Beaver Valley Power Station, 2007

Note: No Cooling Tower 1 sample in September due to outage




Sample location

Figure 5.9

### Density of zebra mussels veligers collected at Beaver Valley Power Station, 2007

Note: Splash pool and EOF not sampled in June due to lack of access



Sample location

Figure 5.10

### Density of settled zebra mussels at Beaver Valley Power Station, 2007

Note: No Intake sample in March due to high water.

Note: No Cooling Tower 1 sample in September due to outage.




Figure 5.11



### Density of settled zebra mussels at Beaver Valley Power Station, 2007

Note: No Barge Slip sample in March due to high water

Note: No Splash pool or EOF samples in June due to lack of access.




Figure 5.12



**10.0**

**PERMITS**

## Attachment 10.1:

## PERMITS &amp; CERTIFICATES FOR ENVIRONMENTAL COMPLIANCE

Registration Number	Regulator/Description	Expiration
PAR000040485	BVPS EPA generator identification Resource Conservation & Recovery Act (RCRA) Identification number for regulated waste activity. Also used by PA DEP to monitor regulated waste activity under the Pennsylvania Solid Waste Management Act (SWMA).	Indefinite
04-02474	BVPS EPA Facility Identification Number for CERCLA/EPCRA/SARA. Used for SARA Tier II reporting and emergency planning.	Indefinite
04-02475	FE Long Term Distribution Center/Warehouse (22) EPA Facility Identification Number for CERCLA/EPCRA/SARA. Used for SARA Tier II reporting and emergency planning.	Indefinite
PA0025615	BVPS NPDES Permit number under US EPA and PA DEP.	12/27/2006 <i>Continued pending approval of renewal application.</i>
04-13281	BVPS Unit 1 PA DEP Facility Identification & certificate number for regulated storage tanks.	Indefinite
04-13361	BVPS Unit 2 PA DEP Facility Identification & certificate number for regulated storage tanks.	Indefinite
OP-04-00086	PA DEP State Only Synthetic Minor Permit for emergency auxiliary boilers, emergency diesel generators, paint shop and other miscellaneous sources..	10/12/2012
N/A	PA DEP Open Burning Permit for operation of the BVPS Fire School- annual application and renewal	12/31/2008
060503 4450 004L	US Department of Transportation Hazardous Materials Registration	06/30/2009
200100242	US Army Permit for maintenance dredging (With Encroachment/Submerged Lands Agreement #0477705, this allows maintenance dredging.).	12/31/2011
0477705	Encroachment Permit/Submerged Lands Agreement for construction and maintenance of current barge slip. (With US Army Permit #200100242, this allows maintenance dredging.)	Indefinite
06786A	Encroachment Permit/Submerged Lands Agreement for transmission line over Ohio River @ Mile 34.5	Indefinite
18737	Encroachment Permit/Submerged Lands Agreement for Unit 1 intake and discharge (main combined intake and outfall structures)	Indefinite
0475711	Encroachment Permit/Submerged Lands Agreement for construction and maintenance of Unit 2 auxiliary intake	Indefinite
- End Table -		

**APPENDIX A**

**SCIENTIFIC AND COMMON NAME<sup>1</sup>  
OF FISH COLLECTED IN THE NEW CUMBERLAND  
POOL OF THE OHIO RIVER, 1970 THROUGH 2007  
BVPS**

<sup>1</sup>Nomenclature follows Robins, et al. (1991)

<u>Family and Scientific Name</u>	<u>Common Name</u>
Lepisosteidae (gars) <i>Lepisosteus osseus</i>	Longnose gar
Hiodontidae (mooneyes) <i>Hiodon alosoides</i> <i>H. tergisus</i>	Goldeye Mooneye
Clupeidae (herrings) <i>Alosa chrysochloris</i> <i>A. pseudoharengus</i> <i>Dorosoma cepedianum</i>	Skipjack herring Alewife Gizzard shad
Cyprinidae (carps and minnows) <i>Campostoma anomalum</i> <i>Carassius auratus</i> <i>Ctenopharyngodon idella</i> <i>Notropis spilopterus</i> <i>Cyprinus carpio</i> <i>C. carpio</i> x <i>C. auratus</i> <i>Luxilus chrysocephalus</i> <i>Macrhybopsis storeriana</i> <i>Nocomis micropogon</i> <i>Notemigonus crysoleucas</i> <i>Notropis atherinoides</i> <i>N. buccatus</i> <i>N. hudsonius</i> <i>N. rubellus</i> <i>N. stramineus</i> <i>N. volucellus</i> <i>Pimephales notatus</i> <i>P. promelas</i> <i>Rhinichthys atratulus</i> <i>Semotilus atromaculatus</i>	Central stoneroller Goldfish Grass carp Spotfin shiner Common carp Carp-goldfish hybrid Striped shiner Silver chub River chub Golden shiner Emerald shiner Silverjaw minnow Spottail shiner Rosyface shiner Sand shiner Mimic shiner Bluntnose minnow Fathead minnow Blacknose dace Creek chub
Catostomidae (suckers) <i>Carpiodes carpio</i> <i>C. cyprinus</i> <i>C. velifer</i> <i>Catostomus commersonii</i> <i>Hypentelium nigricans</i> <i>Ictiobus bubalus</i> <i>I. niger</i> <i>Minytrema melanops</i>	River carpsucker Quillback Highfin carpsucker White sucker Northern hogsucker Smallmouth buffalo Black buffalo Spotted sucker

<u>Family and Scientific Name</u>	<u>Common Name</u>
<i>Moxostoma anisurum</i>	Silver redhorse
<i>M. carinatum</i>	River redhorse
<i>M. duquesnei</i>	Black redhorse
<i>M. erythrurum</i>	Golden redhorse
<i>M. macrolepidotum</i>	Shorthead redhorse
Ictaluridae (bullhead catfishes)	
<i>Ameiurus catus</i>	White catfish
<i>A. furcatus</i>	Blue catfish
<i>A. melas</i>	Black bullhead
<i>A. natalis</i>	Yellow bullhead
<i>A. nebulosus</i>	Brown bullhead
<i>Ictalurus punctatus</i>	Channel catfish
<i>Noturus flavus</i>	Stonecat
<i>Pylodictis olivaris</i>	Flathead catfish
Esocidae (pikes)	
<i>Esox lucius</i>	Northern pike
<i>E. masquinongy</i>	Muskellunge
<i>E. lucius x E. masquinongy</i>	Tiger muskellunge
Salmonidae (trouts)	
<i>Oncorhynchus mykiss</i>	Rainbow trout
Percopsidae (trout-perches)	
<i>Percopsis omiscomaycus</i>	Trout-perch
Cyprinodontidae (killifishes)	
<i>Fundulus diaphanus</i>	Banded killifish
Atherinidae (silversides)	
<i>Labidesthes sicculus</i>	Brook silverside
Percichthyidae (temperate basses)	
<i>Morone chrysops</i>	White bass
<i>M. saxatilis</i>	Striped bass
<i>M. saxatilis x M. chrysops</i>	Striped bass hybrid
Centrarchidae (sunfishes)	
<i>Ambloplites rupestris</i>	Rock bass
<i>Lepomis cyanellus</i>	Green sunfish
<i>L. gibbosus</i>	Pumpkinseed
<i>L. macrochirus</i>	Bluegill
<i>L. microlophus</i>	Redear sunfish
<i>L. gibbosus x L. microlophus</i>	Pumpkinseed-redear sunfish hybrid
<i>Micropterus dolomieu</i>	Smallmouth bass
<i>M. punctulatus</i>	Spotted bass
<i>M. salmoides</i>	Largemouth bass
<i>Pomoxis annularis</i>	White crappie
<i>P. nigromaculatus</i>	Black crappie

**Family and Scientific Name**

**Common Name**

Percidae (perches)

*Etheostoma blennioides*

*E. nigrum*

*E. zonale*

*Perca flavescens*

*Percina caprodes*

*P. copelandi*

*Sander canadense*

*S. vitreum*

*S. canadense* x *S. vitreum*

Greenside darter

Johnny darter

Banded darter

Yellow perch

Logperch

Channel darter

Sauger

Walleye

Saugeye

Sciaenidae (drums)

*Aplodinotus grunniens*

Freshwater drum