

Generic Environmental Impact Statement for In-Situ Leach Uranium Milling Facilities

Chapters 1 through 4

Draft Report for Comment

Office of Federal and State Materials and
Environmental Management Programs

Wyoming Department of Environmental Quality
Land Quality Division

AVAILABILITY OF REFERENCE MATERIALS IN NRC PUBLICATIONS

NRC Reference Material

As of November 1999, you may electronically access NUREG-series publications and other NRC records at NRC's Public Electronic Reading Room at <http://www.nrc.gov/reading-rm.html>.

Publicly released records include, to name a few, NUREG-series publications; *Federal Register* notices; applicant, licensee, and vendor documents and correspondence; NRC correspondence and internal memoranda; bulletins and information notices; inspection and investigative reports; licensee event reports; and Commission papers and their attachments.

NRC publications in the NUREG series, NRC regulations, and *Title 10, Energy*, in the Code of *Federal Regulations* may also be purchased from one of these two sources.

1. The Superintendent of Documents
U.S. Government Printing Office
Mail Stop SSOP
Washington, DC 20402-0001
Internet: bookstore.gpo.gov
Telephone: 202-512-1800
Fax: 202-512-2250
2. The National Technical Information Service
Springfield, VA 22161-0002
www.ntis.gov
1-800-553-6847 or, locally, 703-605-6000

A single copy of each NRC draft report for comment is available free, to the extent of supply, upon written request as follows:

Address: U.S. Nuclear Regulatory Commission
Office of Administration
Mail, Distribution and Messenger Team
Washington, DC 20555-0001
E-mail: DISTRIBUTION@nrc.gov
Facsimile: 301-415-2289

Some publications in the NUREG series that are posted at NRC's Web site address <http://www.nrc.gov/reading-rm/doc-collections/nuregs> are updated periodically and may differ from the last printed version. Although references to material found on a Web site bear the date the material was accessed, the material available on the date cited may subsequently be removed from the site.

Non-NRC Reference Material

Documents available from public and special technical libraries include all open literature items, such as books, journal articles, and transactions, *Federal Register* notices, Federal and State legislation, and congressional reports. Such documents as theses, dissertations, foreign reports and translations, and non-NRC conference proceedings may be purchased from their sponsoring organization.

Copies of industry codes and standards used in a substantive manner in the NRC regulatory process are maintained at—

The NRC Technical Library
Two White Flint North
11545 Rockville Pike
Rockville, MD 20852-2738

These standards are available in the library for reference use by the public. Codes and standards are usually copyrighted and may be purchased from the originating organization or, if they are American National Standards, from—

American National Standards Institute
11 West 42nd Street
New York, NY 10036-8002
www.ansi.org
212-642-4900

Legally binding regulatory requirements are stated only in laws; NRC regulations; licenses, including technical specifications; or orders, not in NUREG-series publications. The views expressed in contractor-prepared publications in this series are not necessarily those of the NRC.

The NUREG series comprises (1) technical and administrative reports and books prepared by the staff (NUREG-XXXX) or agency contractors (NUREG/CR-XXXX), (2) proceedings of conferences (NUREG/CP-XXXX), (3) reports resulting from international agreements (NUREG/IA-XXXX), (4) brochures (NUREG/BR-XXXX), and (5) compilations of legal decisions and orders of the Commission and Atomic and Safety Licensing Boards and of Directors' decisions under Section 2.206 of NRC's regulations (NUREG-0750).

Generic Environmental Impact Statement for In-Situ Leach Uranium Milling Facilities

Chapters 1 through 4

Draft Report for Comment

Manuscript Completed: June 2008

Date Published: July 2008

Prepared by:

**U.S. Nuclear Regulatory Commission
Office of Federal and State Materials and
Environmental Management Programs**

**Wyoming Department of Environmental Quality
Land Quality Division**

COMMENTS ON DRAFT REPORT

Any interested party may submit comments on this report for consideration by the NRC staff. Comments may be accompanied by additional relevant information or supporting data. Please specify the report number NUREG-1910, draft, in your comments, and send them postmarked by September 26, 2008, to the following address:

Chief, Rulemaking, Directives and Editing Branch
U.S. Nuclear Regulatory Commission
Mail Stop T6-D59
Washington, DC 20555-0001

Comments postmarked after September 26, 2008, will be considered to the extent practical.

Electronic comments may be submitted to the NRC by the Internet at
NRCREP.Resource@nrc.gov.

For any questions about the material in this report, please contact:

J. Park
TWFN T8-F5
U.S. Nuclear Regulatory Commission
Washington, DC 20555-0001
Phone: 301-415-6935
E-mail: James.Park@nrc.gov

**NUREG-1910, Vol. 1, has been reproduced
from the best available copy.**

ABSTRACT

The U.S. Nuclear Regulatory Commission (NRC) has prepared a Draft Generic Environmental Impact Statement (Draft GEIS) to identify and evaluate potential environmental impacts associated with the construction, operation, aquifer restoration, and decommissioning of *in-situ* leach (ISL) uranium recovery facilities for identified regions in the western United States. Based on discussions between uranium mining companies and the NRC staff, ISL facilities could be located in portions of Wyoming, Nebraska, South Dakota, and New Mexico. NRC is the licensing authority for ISL facilities in these states.

NRC developed this Draft GEIS using (1) knowledge gained during the past 30 years licensing and regulating ISL facilities, (2) the active participation of the State of Wyoming Department of Environmental Quality as a cooperating agency, and (3) public comments received during the scoping period for the GEIS. NRC's research indicates that the technology used for ISL uranium recovery is relatively standardized throughout the industry and therefore appropriate for a programmatic evaluation in a GEIS.

As a framework for the analyses presented in this GEIS, NRC has identified four geographic regions based on

- Past and existing uranium milling sites are located within States where NRC has regulatory authority over uranium recovery;
- Potential new sites are identified based on NRC's understanding of where the uranium recovery industry has plans to develop uranium deposits using ISL technology; and
- Locations of historical uranium deposits within portions of Wyoming, Nebraska, South Dakota, and New Mexico.

The purpose behind developing the GEIS is to improve the efficiency of NRC's environmental reviews for ISL license applications required under the National Environmental Policy Act of 1969, as amended (NEPA). NRC regulations that implement NEPA and discuss environmental reviews are found in Title 10, "Energy," of the Code of Federal Regulations (10 CFR) Part 51. The NRC staff plans to use the GEIS as a starting point for its NEPA analyses for site-specific license applications for new ISL facilities. Additionally, the NRC staff plans to use the GEIS, along with applicable previous site-specific environmental review documents, in its NEPA analysis for the restart or expansions of existing facilities.

Paperwork Reduction Act Statement

This NUREG contains information collection requirements that are subject to the Paperwork Reduction Act of 1995 (44 U.S.C. 3501 et seq.) These information collections were approved by the Office of Management and Budget, approval numbers 3150-0020; 3150-0014.

Public Protection Notification

The NRC may not conduct or sponsor, and a person is not required to respond to, a request for information or an information collection requirement unless the requesting documents displays a currently valid OMB control number.

CONTENTS

Section	Page
ABSTRACT	iii
FIGURES	xxiii
TABLES	xxviii
EXECUTIVE SUMMARY	xxxiii
ABBREVIATIONS/ACRONYMS	li
SI* (MODERN METRIC) CONVERSION FACTORS	liii
 1 INTRODUCTION	 1-1
1.1 Purpose of the GEIS	1-1
1.2 The Proposed Federal Action	1-3
1.3 Purpose and Need for the Action	1-4
1.4 Structure of the GEIS	1-4
1.4.1 Describing the ISL Process	1-5
1.4.2 Describing the Affected Environment	1-5
1.4.3 Identifying Environmental Issues and Characterizing Significance ..	1-5
1.5 Scope of the GEIS	1-6
1.5.1 The GEIS Scoping Process	1-6
1.5.2 Issues To Be Studied in Detail	1-8
1.5.3 Issues Eliminated From Detailed Study	1-12
1.5.4 Issues Outside of the Scope of the GEIS	1-13
1.6 Agencies Involved in Uranium ISL Facility Licensing	1-13
1.6.1 Federal Agencies	1-13
1.6.1.1 NRC	1-13
1.6.1.2 EPA	1-14
1.6.1.3 Occupational Safety and Health Administration	1-14
1.6.1.4 U.S. Department of Transportation	1-14
1.6.1.5 Other Federal Agencies	1-15
1.6.2 Tribal Agencies	1-15
1.6.3 State Agencies	1-15
1.6.3.1 Wyoming Department of Environmental Quality	1-15
1.6.3.2 Nebraska Department of Environmental Quality	1-16
1.6.3.3 South Dakota Department of Environmental and	
Natural Resources	1-16
1.6.3.4 New Mexico Environmental Department	1-16
1.7 Licensing and Permitting Process for a Uranium ISL Facility	1-16
1.7.1 The NRC Licensing Process	1-17
1.7.2 EPA Permitting	1-17
1.7.2.1 Water Resources	1-17
1.7.2.2 Air Quality	1-20
1.7.3 Other Federal Agencies	1-21
1.7.4 Tribal Agencies	1-22
1.7.5 State Agencies	1-22
1.7.5.1 Wyoming	1-22
1.7.5.2 Nebraska	1-23
1.7.5.3 South Dakota	1-23
1.7.5.4 New Mexico	1-23
1.8 Use of the GEIS in the NRC Licensing Process	1-24
1.8.1 Applicant's or Licensee's Environmental Report	1-25

CONTENTS (continued)

Section	Page
1.8.2	Acceptance Review of the License Application and Environmental Report 1-25
1.8.3	NRC's Site-Specific Environmental Review..... 1-26
1.8.4	Public Participation Activities..... 1-27
1.8.5	NRC's Final Environmental Review Document and Findings 1-27
1.9	References..... 1-27
2	<i>IN-SITU</i> LEACH URANIUM RECOVERY AND ALTERNATIVES..... 2-1
2.1	Overview of ISL Uranium Recovery..... 2-1
2.1.1	Geochemistry of Uranium..... 2-1
2.1.2	Physical Characteristics of Uranium Deposits 2-2
2.1.3	General Description of ISL Facilities..... 2-4
2.2	Pre-Construction..... 2-6
2.3	Construction..... 2-7
2.3.1	Underground Infrastructure 2-7
2.3.1.1	Well Fields..... 2-7
2.3.1.2	Pipelines..... 2-13
2.3.2	Surface Facilities..... 2-14
2.4	Operations..... 2-15
2.4.1	Uranium Mobilization..... 2-15
2.4.1.1	Lixiviant Chemistry 2-15
2.4.1.2	Lixiviant Injection and Production 2-16
2.4.1.3	Excursions..... 2-18
2.4.1.4	Excursion Monitoring..... 2-18
2.4.2	Uranium Processing..... 2-20
2.4.2.1	Ion Exchange 2-20
2.4.2.2	Elution..... 2-20
2.4.2.3	Precipitation, Drying, and Packaging..... 2-22
2.4.3	Management of Production Bleed and Other Liquid Effluents..... 2-25
2.5	Aquifer Restoration..... 2-26
2.5.1	Groundwater Transfer..... 2-26
2.5.2	Groundwater Sweep 2-27
2.5.3	Reverse Osmosis, Permeate Injection, and Recirculation 2-28
2.5.4	Stabilization..... 2-29
2.6	Decontamination, Decommissioning, and Reclamation 2-30
2.7	Effluents and Waste Management..... 2-32
2.7.1	Gaseous or Airborne Particulate Emissions 2-32
2.7.2	Liquid Wastes 2-35
2.7.3	Solid Wastes..... 2-37
2.8	Transportation 2-38
2.9	Radiological Health and Safety..... 2-40
2.10	Financial Surety..... 2-41
2.11	Information From Historical Operation of ISL Uranium Milling Facilities 2-42
2.11.1	Area of ISL Uranium Milling Facilities 2-42
2.11.2	Spills and Leaks..... 2-43
2.11.3	Groundwater Use..... 2-46
2.11.4	Excursions 2-46

CONTENTS (continued)

Section	Page
2.11.5	Aquifer Restoration 2-48
2.11.6	Socioeconomic Information 2-51
2.12	Alternatives Considered and Included in the Impact Analysis 2-51
2.12.1	The No-Action Alternative 2-51
2.13	Alternatives Considered and Excluded From the Impact Analysis 2-52
2.14	References 2-52
3	DESCRIPTION OF THE AFFECTED ENVIRONMENT 3.1-1
3.1	Introduction 3.1-1
3.1.1	Geographic Scope—Defining Uranium Milling Regions 3.1-1
3.1.2	General Information for All Uranium Milling Regions 3.1-7
3.1.3	References 3.1-8
3.2	Wyoming West Uranium Milling Region 3.2-1
3.2.1	Land Use 3.2-1
3.2.2	Transportation 3.2-4
3.2.3	Geology and Soils 3.2-7
3.2.4	Water Resources 3.2-12
3.2.4.1	Surface Waters 3.2-12
3.2.4.2	Wetlands and Waters of the United States 3.2-15
3.2.4.3	Groundwater 3.2-17
3.2.4.3.1	Regional Aquifer Systems 3.2-17
3.2.4.3.2	Aquifer Systems in the Vicinity of Uranium Milling Sites 3.2-18
3.2.4.3.3	Uranium-Bearing Aquifers 3.2-19
3.2.4.3.4	Other Important Surrounding Aquifers for Water Supply 3.2-21
3.2.5	Ecology 3.2-22
3.2.5.1	Terrestrial 3.2-22
3.2.5.2	Aquatic 3.2-27
3.2.5.3	Threatened and Endangered Species 3.2-37
3.2.6	Meteorology, Climatology, and Air Quality 3.2-44
3.2.6.1	Meteorology and Climatology 3.2-44
3.2.6.2	Air Quality 3.2-45
3.2.7	Noise 3.2-49
3.2.8	Historical and Cultural Resources 3.2-53
3.2.8.1	Cultural Resources Overview 3.2-53
3.2.8.2	Historic Properties Listed in the National And State Registers 3.2-57
3.2.8.3	Tribal Consultation 3.2-59
3.2.8.4	Places of Cultural Significance 3.2-60
3.2.9	Visual/Scenic Resources 3.2-63
3.2.10	Socioeconomic 3.2-66
3.2.10.1	Demographics 3.2-67
3.2.10.2	Income 3.2-71
3.2.10.3	Housing 3.2-71
3.2.10.4	Employment Structure 3.2-75
3.2.10.4.1	State Data 3.2-76
3.2.10.4.1.1	Montana 3.2-76

CONTENTS (continued)

Section	Page
3.2.10.4.1.2	Wyoming 3.2-76
3.2.10.4.2	County Data 3.2-77
3.2.10.5	Local Finance 3.2-78
3.2.10.6	Education 3.2-79
3.2.10.7	Health and Social Services 3.2-80
3.2.11	Public and Occupational Health 3.2-80
3.2.11.1	Background Radiological Conditions 3.2-80
3.2.11.2	Public Health and Safety 3.2-81
3.2.11.3	Occupational Health and Safety 3.2-82
3.2.12	References 3.2-82
3.3	Wyoming East Uranium Milling Region 3.3-1
3.3.1	Land Use 3.3-1
3.3.2	Transportation 3.3-4
3.3.3	Geology and Soils 3.3-6
3.3.4	Water Resources 3.3-12
3.3.4.1	Surface Waters 3.3-12
3.3.4.2	Wetlands and Waters of the United States 3.3-15
3.3.4.3	Groundwater 3.3-16
3.3.4.3.1	Regional Aquifer Systems 3.3-16
3.3.4.3.2	Aquifer Systems in the Vicinity of Uranium Milling Sites 3.3-17
3.3.4.3.3	Uranium-Bearing Aquifers 3.3-18
3.3.4.3.4	Other Important Surrounding Aquifers for Water Supply 3.3-19
3.3.5	Ecology 3.3-20
3.3.5.1	Wyoming East Uranium Milling Flora 3.3-20
3.3.5.2	Aquatic 3.3-24
3.3.5.3	Threatened and Endangered Species 3.3-32
3.3.6	Meteorology, Climatology, and Air Quality 3.3-35
3.3.6.1	Meteorology and Climatology 3.3-35
3.3.6.2	Air Quality 3.3-37
3.3.7	Noise 3.3-39
3.3.8	Historical and Cultural Resources 3.3-41
3.3.8.1	Cultural Resources Overview 3.3-41
3.3.8.2	Historic Properties Listed in the National and State Registers 3.3-41
3.3.8.3	Tribal Consultation 3.3-42
3.3.8.4	Places of Cultural Significance 3.3-43
3.3.9	Visual/Scenic Resources 3.3-43
3.3.10	Socioeconomic 3.3-45
3.3.10.1	Demographics 3.3-45
3.3.10.2	Income 3.3-48
3.3.10.3	Housing 3.3-51
3.3.10.4	Employment Structure 3.3-51
3.3.10.4.1	State Data 3.3-52
3.3.10.4.1.1	Colorado 3.3-52
3.3.10.4.1.2	South Dakota 3.3-52
3.3.10.4.1.3	Wyoming 3.3-53

CONTENTS (continued)

Section	Page
3.3.10.4.2	County Data 3.3-53
3.3.10.5	Local Finance 3.3-55
3.3.10.6	Education..... 3.3-57
3.3.10.7	Health and Social Services 3.3-59
3.3.11	Public and Occupational Health 3.3-59
3.3.11.1	Background Radiological Conditions..... 3.3-59
3.3.11.2	Public Health and Safety..... 3.3-60
3.3.11.3	Occupational Health and Safety..... 3.3-60
3.3.12	References..... 3.3-60
3.4	Nebraska-South Dakota-Wyoming Uranium Milling 3.4-1
3.4.1	Land Use..... 3.4-1
3.4.2	Transportation..... 3.4-4
3.4.3	Geology and Soils 3.4-6
3.4.3.1	The Black Hills (Western South Dakota-Northeastern Wyoming) 3.4-6
3.4.3.2	The Crawford Basin(Northwestern Nebraska) 3.4-12
3.4.4	Water Resources 3.4-17
3.4.4.1	Surface Waters 3.4-17
3.4.4.2	Wetlands and Waters of the United States..... 3.4-22
3.4.4.3	Groundwater 3.4-22
3.4.4.3.1	Regional Aquifer Systems 3.4-22
3.4.4.3.2	Aquifer systems in the Vicinity of Uranium Milling Sites 3.4-25
3.4.4.3.3	Uranium-Bearing Aquifers 3.4-26
3.4.4.3.4	Other Important Surrounding Aquifers for Water Supply 3.4-27
3.4.5	Ecology 3.4-28
3.4.5.1	Nebraska-South Dakota-Wyoming Uranium Milling Region Flora 3.4-28
3.4.5.2	Aquatic..... 3.4-31
3.4.5.3	Threatened and Endangered Species..... 3.4-43
3.4.6	Meteorology, Climatology, and Air Quality..... 3.4-47
3.4.6.1	Meteorology and Climatology..... 3.4-47
3.4.6.2	Air Quality 3.4-48
3.4.7	Noise 3.4-49
3.4.8	Historical and Cultural Resources 3.4-53
3.4.8.1	Cultural Resources Overview..... 3.4-53
3.4.8.1.1	Cultural Resources of Western and Southwestern South Dakota..... 3.4-53
3.4.8.1.2	Cultural Resources of Western Nebraska 3.4-56
3.4.8.2	Historic Properties Listed in the National and State Registers 3.4-58
3.4.8.2.1	Historic Properties in Western South Dakota 3.4-58
3.4.8.2.2	Historic Properties in Western Nebraska 3.4-63
3.4.8.3	Tribal Consultations 3.4-64

CONTENTS (continued)

Section	Page
3.4.8.3.1	South Dakota Tribal Consultation..... 3.4-64
3.4.8.3.2	Nebraska Tribal Consultation 3.4-64
3.4.8.4	Places of Cultural Significance 3.4-65
3.4.8.4.1	Places of Cultural Significance in South Dakota 3.4-66
3.4.8.4.2	Places of Cultural Significance in Nebraska..... 3.4-66
3.4.9	Visual/Scenic Resources..... 3.4-66
3.4.10	Socioeconomic..... 3.4-67
3.4.10.1	Demographics..... 3.4-68
3.4.10.2	Income..... 3.4-68
3.4.10.3	Housing 3.4-78
3.4.10.4	Employment Structure 3.4-81
3.4.10.4.1	State Data..... 3.4-81
3.4.10.4.1.1	Nebraska..... 3.4-81
3.4.10.4.1.2	South Dakota 3.4-82
3.4.10.4.1.3	Wyoming 3.4-83
3.4.10.4.2	County Data..... 3.4-83
3.4.10.4.2.1	Nebraska..... 3.4-83
3.4.10.4.2.2	South Dakota 3.4-83
3.4.10.4.2.3	Wyoming 3.4-84
3.4.10.4.3	Native American Communities 3.4-85
3.4.10.5	Local Finance 3.4-85
3.4.10.5.1	Nebraska 3.4-85
3.4.10.5.2	South Dakota 3.4-85
3.4.10.5.3	Wyoming..... 3.4-86
3.4.10.5.4	Native American Communities 3.4-86
3.4.10.6	Education..... 3.4-87
3.4.10.6.1	Nebraska 3.4-87
3.4.10.6.2	South Dakota 3.4-87
3.4.10.6.3	Wyoming..... 3.4-88
3.4.10.7	Health and Social Services..... 3.4-89
3.4.11	Public and Occupational Health 3.4-89
3.4.11.1	Background Radiological Conditions..... 3.4-89
3.4.11.2	Public Health and Safety..... 3.4-90
3.4.11.3	Occupational Health and Safety..... 3.4-90
3.4.12	References 3.4-90
3.5	Northwestern New Mexico Uranium Milling Region 3.5-1
3.5.1	Land Use..... 3.5-1
3.5.2	Transportation..... 3.5-4
3.5.3	Geology and Soils 3.5-6
3.5.4	Water Resources 3.5-14
3.5.4.1	Surface Waters 3.5-14
3.5.4.2	Wetlands and Waters of the United States..... 3.5-14
3.5.4.3	Groundwater..... 3.5-17
3.5.4.3.1	Regional Aquifer Systems..... 3.5-18
3.5.4.3.2	Aquifer Systems in the Vicinity of Uranium Milling Sites 3.5-19

CONTENTS (continued)

Section		Page
	3.5.4.3.3 Uranium-Bearing Aquifers	3.5-20
	3.5.4.3.4 Other Important Surrounding Aquifers for Water Supply	3.5-21
3.5.5	Ecology	3.5-21
	3.5.5.1 Northwestern New Mexico Flora	3.5-21
	3.5.5.2 Aquatic.....	3.5-26
	3.5.5.3 Threatened and Endangered Species.....	3.5-28
3.5.6	Meteorology, Climatology, and Air Quality.....	3.5-32
	3.5.6.1 Meteorology and Climatology.....	3.5-32
	3.5.6.2 Air Quality	3.5-34
3.5.7	Noise	3.5-36
3.5.8	Historical and Cultural Resources	3.5-37
	3.5.8.1 New Mexico Historic and Cultural Resources.....	3.5-39
	3.5.8.2 Historic Properties Listed in the National and State Registers	3.5-52
	3.5.8.3 New Mexico Tribal Consultation.....	3.5-52
	3.5.8.4 Traditional Cultural Landscapes.....	3.5-55
3.5.9	Visual/Scenic Resources.....	3.5-59
3.5.10	Socioeconomic.....	3.5-61
	3.5.10.1 Demographics.....	3.5-62
	3.5.10.2 Income.....	3.5-65
	3.5.10.3 Housing	3.5-65
	3.5.10.4 Employment Structure	3.5-70
	3.5.10.4.1 State Data.....	3.5-70
	3.5.10.4.1.1 Arizona	3.5-70
	3.5.10.4.1.2 New Mexico.....	3.5-70
	3.5.10.4.2 County Data.....	3.5-71
	3.5.10.5 Local Finance	3.5-72
	3.5.10.6 Education.....	3.5-75
	3.5.10.7 Health and Social Services	3.5-76
3.5.11	Public and Occupational Health	3.5-77
	3.5.11.1 Background Radiological Conditions.....	3.5-77
	3.5.11.2 Public Health and Safety.....	3.5-78
	3.5.11.3 Occupational Health and Safety.....	3.5-78
3.5.12	References	3.5-78
4	ENVIRONMENTAL IMPACTS OF CONSTRUCTION, OPERATION, AQUIFER RESTORATION, AND DECOMMISSIONING ACTIVITIES	4.1-1
4.1	Introduction.....	4.1-1
4.2	Wyoming West Uranium Milling Region.....	4.2-1
	4.2.1 Land Use Impacts	4.2-1
	4.2.1.1 Construction Impacts to Land Use	4.2-1
	4.2.1.2 Operation Impacts to Land Use.....	4.2-4
	4.2.1.3 Aquifer Restoration Impacts to Land Use.....	4.2-4
	4.2.1.4 Decommissioning Impacts to Land Use	4.4-4
4.2.2	Transportation Impacts.....	4.4-5
	4.2.2.1 Construction Impacts to Transportation.....	4.2-5
	4.2.2.2 Operation Impacts to Transportation.....	4.2-5

CONTENTS (continued)

Section	Page
4.2.2.3	Aquifer Restoration Impacts to Transportation 4.2-8
4.2.2.4	Decommissioning Impacts to Transportation..... 4.2-8
4.2.3	Geology and Soils Impacts..... 4.2-9
4.2.3.1	Construction Impacts to Geology and Soils..... 4.2-9
4.2.3.2	Operation Impacts to Geology and Soils 4.2-10
4.2.3.3	Aquifer Restoration Impacts to Geology and Soils 4.2-12
4.2.3.4	Decommissioning Impacts to Geology and Soils..... 4.2-13
4.2.4	Water Resources Impacts 4.2-14
4.2.4.1	Surface Water Impacts..... 4.2-14
4.2.4.1.1	Construction Impacts to Surface Water 4.2-14
4.2.4.1.2	Operation Impacts to Surface Water 4.2-15
4.2.4.1.3	Aquifer Restoration Impacts to Surface Water 4.2-16
4.2.4.1.4	Decommissioning Impacts to Surface Water 4.2-17
4.2.4.2	Groundwater Impacts..... 4.2-17
4.2.4.2.1	Construction Impacts to Groundwater 4.2-18
4.2.4.2.2	Operation Impacts to Groundwater 4.2-18
4.2.4.2.2.1	Operation Impacts to Shallow (Near-Surface) Aquifers 4.2-18
4.2.4.2.2.2	Operation Impacts to Production and Surrounding Aquifers 4.2-19
4.2.4.2.2.3	Operation Impacts to Deep Aquifers Below the Production Aquifers 4.2-24
4.2.4.2.3	Aquifer Restoration Impacts to Groundwater 4.2-24
4.2.4.2.4	Decommissioning Impacts to Groundwater 4.2-26
4.2.5	Ecological Resources Impacts 4.2-27
4.2.5.1	Construction Impacts to Ecological Resources 4.2-27
4.2.5.2	Operation Impacts to Ecological Resources..... 4.2-32
4.2.5.3	Aquifer Restoration Impacts to Ecological Resources 4.2-33
4.2.5.4	Decommissioning Impacts to Ecological Resources 4.2-33
4.2.6	Air Quality Impacts 4.2-33
4.2.6.1	Construction Impacts to Air Quality 4.2-34
4.2.6.2	Operation Impacts to Air Quality 4.2-35
4.2.6.3	Aquifer Restoration Impacts to Air Quality..... 4.2-35
4.2.6.4	Decommissioning Impacts to Air Quality 4.2-36

CONTENTS (continued)

Section	Page
4.2.7	Noise Impacts 4.2-36
4.2.7.1	Construction Impacts to Noise 4.2-36
4.2.7.2	Operation Impacts to Noise..... 4.2-40
4.2.7.3	Aquifer Restoration Impacts to Noise 4.2-41
4.2.7.4	Decommissioning Impacts to Noise 4.2-41
4.2.8	Historical and Cultural Resources Impacts 4.2-42
4.2.8.1	Construction Impacts to Historical and Cultural Resources..... 4.2-42
4.2.8.2	Operation Impacts to Historical and Cultural Resources 4.2-43
4.2.8.3	Aquifer Restoration Impacts to Historical and Cultural Resources 4.2-43
4.2.8.4	Decommissioning Impacts to Historical and Cultural Resources..... 4.2-44
4.2.9	Visual/Scenic Resources Impacts 4.2-44
4.2.9.1	Construction Impacts to Visual/ Scenic Resources 4.2-44
4.2.9.2	Operation Impacts to Visual/Scenic Resources..... 4.2-46
4.2.9.3	Aquifer Restoration Impacts to Visual/Scenic Resources..... 4.2-46
4.2.9.4	Decommissioning Impacts to Visual/Scenic Resources 4.2-47
4.2.10	Socioeconomic Impacts 4.2-47
4.2.10.1	Construction Impacts to Socioeconomics..... 4.2-48
4.2.10.2	Operation Impacts to Socioeconomics 4.2-49
4.2.10.3	Aquifer Restoration Impacts to Socioeconomics 4.2-50
4.2.10.4	Decommissioning Impacts to Socioeconomics..... 4.2-51
4.2.11	Public and Occupational Health and Safety Impacts 4.2-51
4.2.11.1	Construction Impacts to Public and Occupational Health and Safety 4.2-51
4.2.11.2	Operation Impacts to Public and Occupational Health and Safety 4.2-52
4.2.11.2.1	Radiological Impacts to Public and Occupational Health and Safety for Normal Operations..... 4.2-52
4.2.11.2.2	Radiological Impacts to Public and Occupational Health and Safety From Accidents..... 4.2-54
4.2.11.2.3	Non-radiological Impacts on Public and Occupational Health and Safety From Normal Operations..... 4.2-55
4.2.11.2.4	Non-radiological Impacts to Public and Occupational Health and Safety From Accidents..... 4.2-55
4.2.11.3	Aquifer Restoration Impacts to Public and Occupational Health and Safety..... 4.2-57
4.2.11.4	Decommissioning Impacts to Public and Occupational Health and Safety..... 4.2-59

CONTENTS (continued)

Section	Page
4.2.12	Waste Management Impacts 4.2-59
4.2.12.1	Construction Impacts to Waste Management 4.2-60
4.2.12.2	Operations Impacts to Waste Management 4.2-60
4.2.12.3	Aquifer Restoration Impacts to Waste Management... 4.2-61
4.2.12.4	Decommissioning Impacts to Waste Management 4.2-61
4.2.13	References 4.2-63
4.3	Wyoming East Uranium Milling Region 4.3-1
4.3.1	Land Use Impacts 4.3-1
4.3.1.1	Construction Impacts to Land Use 4.3-1
4.3.1.2	Operation Impacts to Land Use 4.3-1
4.3.1.3	Aquifer Restoration Impacts to Land Use 4.3-2
4.3.1.4	Decommissioning Impacts to Land Use 4.3-2
4.3.2	Transportation Impacts 4.3-2
4.3.2.1	Construction Impacts to Transportation 4.3-2
4.3.2.2	Operation Impacts to Transportation 4.3-3
4.3.2.3	Aquifer Restoration Impacts to Transportation 4.3-3
4.3.2.4	Decommissioning Impacts to Transportation 4.3-3
4.3.3	Geology and Soils Impacts 4.3-4
4.3.3.1	Construction Impacts to Geology and Soils 4.3-4
4.3.3.2	Operation Impacts to Geology and Soils 4.3-5
4.3.3.3	Aquifer Restoration Impacts to Geology and Soils 4.3-7
4.3.3.4	Decommissioning Impacts to Geology and Soils 4.3-8
4.3.4	Water Resources Impacts 4.3-9
4.3.4.1	Surface Water Impacts 4.3-9
4.3.4.1.1	Construction Impacts to Surface Waters 4.3-9
4.3.4.1.2	Operation Impacts to Surface Waters 4.3-9
4.3.4.1.3	Aquifer Restoration Impacts to Surface Waters 4.3-9
4.3.4.1.4	Decommissioning Impacts to Surface Water 4.3-9
4.3.4.2	Groundwater Impacts 4.3-10
4.3.4.2.1	Construction Impacts to Groundwater 4.3-10
4.3.4.2.2	Operation Impacts to Groundwater 4.3-10
4.3.4.2.2.1	Operation Impacts to Shallow (Near-Surface) Aquifers 4.3-11
4.3.4.2.2.2	Operation Impacts to Production and Surrounding Aquifers 4.3-12
4.3.4.2.2.3	Operation Impacts to Deep Aquifers Below the Production Aquifers 4.3-15
4.3.4.2.3	Aquifer Restoration Impacts to Groundwater 4.3-16
4.3.4.2.4	Decommissioning Impacts to Groundwater 4.3-18

CONTENTS (continued)

Section	Page
4.3.5	Ecological Resources Impacts 4.3-19
4.3.5.1	Construction Impacts to Ecological Resources 4.3-19
4.3.5.2	Operation Impacts to Ecological Resources 4.3-20
4.3.5.3	Aquifer Restoration Impacts to Ecological Resources 4.3-21
4.3.5.4	Decommissioning Impacts to Ecological Resources 4.3-21
4.3.6	Air Quality Impacts 4.3-21
4.3.6.1	Construction Impacts to Air Quality 4.3-21
4.3.6.2	Operation Impacts to Air Quality 4.3-22
4.3.6.3	Aquifer Restoration Impacts to Air Quality 4.3-22
4.3.6.4	Decommissioning Impacts to Air Quality 4.3-22
4.3.7	Noise Impacts 4.3-23
4.3.7.1	Construction Impacts to Noise 4.3-23
4.3.7.2	Operation Impacts to Noise 4.3-23
4.3.7.3	Aquifer Restoration Impacts to Noise 4.3-23
4.3.7.4	Decommissioning Impacts to Noise 4.3-24
4.3.8	Historical and Cultural Resources Impacts 4.3-24
4.3.8.1	Construction Impacts to Historical and Cultural Resources 4.3-25
4.3.8.2	Operation Impacts to Historical and Cultural Resources 4.3-26
4.3.8.3	Aquifer Restoration Impacts to Historical and Cultural Resources 4.3-26
4.3.8.4	Decommissioning Impacts to Historical and Cultural Resources 4.3-26
4.3.9	Visual/Scenic Resources Impacts 4.3-27
4.3.9.1	Construction Impacts to Visual/Scenic Resources 4.3-27
4.3.9.2	Operation Impacts to Visual/Scenic Resources 4.3-27
4.3.9.3	Aquifer Restoration Impacts to Visual/Scenic Resources 4.3-27
4.3.9.4	Decommissioning Impacts to Visual/Scenic Resources 4.3-28
4.3.10	Socioeconomic Impacts 4.3-28
4.3.10.1	Construction Impacts to Socioeconomics 4.3-28
4.3.10.2	Operation Impacts to Socioeconomics 4.3-30
4.3.10.3	Aquifer Restoration Impacts to Socioeconomics 4.3-31
4.3.10.4	Decommissioning Impacts to Socioeconomics 4.3-31
4.3.11	Public and Occupational Health and Safety Impacts 4.3-32
4.3.11.1	Construction Impacts to Public and Occupational Health and Safety 4.3-32
4.3.11.2	Operation Impacts to Public and Occupational Health and Safety 4.3-32
4.3.11.2.1	Radiological Impacts to Public and Occupational Health and Safety 4.3-32
4.3.11.2.2	Radiological Impacts to Public and Occupational Health and Safety From Accidents 4.3-33

CONTENTS (continued)

Section	Page
4.3.11.2.3	Non-radiological Impacts to Public and Occupational Health and Safety for Normal Operations..... 4.3-33
4.3.11.2.4	Non-radiological Impacts to Public and Occupational Health and Safety for Accidents 4.3-33
4.3.11.3	Aquifer Restoration Impacts to Public and Occupational Health and Safety 4.3-33
4.3.11.4	Decommissioning Impacts to Public and Occupational Health and Safety 4.3-33
4.3.12	Waste Management Impacts..... 4.3-34
4.3.12.1	Construction Impacts to Waste Management..... 4.3-34
4.3.12.2	Operation Impacts to Waste Management 4.3-34
4.3.12.3	Aquifer Restoration Impacts to Waste Management .. 4.3-34
4.3.12.4	Decommissioning Impacts to Waste Management..... 4.3-35
4.3.13	References..... 4.3-35
4.4	Nebraska-South Dakota-Wyoming Uranium Milling Region 4.4-1
4.4.1	Land Use Impacts 4.4-1
4.4.1.1	Construction Impacts to Land Use 4.4-1
4.4.1.2	Operation Impacts to Land Use..... 4.4-1
4.4.1.3	Aquifer Restoration Impacts to Land Use..... 4.4-2
4.4.1.4	Decommissioning Impacts to Land Use 4.4-2
4.4.2	Transportation Impacts..... 4.4-2
4.4.2.1	Construction Impacts to Transportation..... 4.4-2
4.4.2.2	Operation Impacts to Transportation 4.4-3
4.4.2.3	Aquifer Restoration Impacts to Transportation 4.4-3
4.4.2.4	Decommissioning Impacts to Transportation..... 4.4-3
4.4.3	Geology and Soils Impacts..... 4.4-4
4.4.3.1	Construction Impacts to Geology and Soils..... 4.4-4
4.4.3.2	Operation Impacts to Geology and Soils 4.4-5
4.4.3.3	Aquifer Restoration Impacts to Geology and Soils 4.4-7
4.4.3.4	Decommissioning Impacts to Geology and Soils..... 4.4-8
4.4.4	Water Resources Impacts 4.4-9
4.4.4.1	Surface Water Impacts..... 4.4-9
4.4.4.1.1	Construction Impacts to Surface Water 4.4-9
4.4.4.1.2	Operation Impacts to Surface Water 4.4-9
4.4.4.1.3	Aquifer Restoration Impacts to Surface Water..... 4.4-9
4.4.4.1.4	Decommissioning Impacts to Surface Water..... 4.4-10

CONTENTS (continued)

Section	Page
4.4.4.2	Groundwater Impacts..... 4.4-10
4.4.4.2.1	Construction Impacts to Groundwater..... 4.4-10
4.4.4.2.2	Operation Impacts to Groundwater 4.4-11
4.4.4.2.2.1	Operation Impacts to Shallow (Near-Surface) Aquifers 4.4-11
4.4.4.2.2.2	Operation Impacts to Production and Surrounding Aquifers..... 4.4-12
4.4.4.2.2.3	Operation Impacts to Deep Aquifers Below the Production Aquifers..... 4.4-16
4.4.4.2.3	Aquifer Restoration Impacts to Groundwater..... 4.4-17
4.4.4.2.4	Decommissioning Impacts to Groundwater..... 4.4-19
4.4.5	Ecological Resources Impacts 4.4-20
4.4.5.1	Construction Impacts to Ecological Resources 4.4-20
4.4.5.2	Operation Impacts to Ecological Resources..... 4.4-21
4.4.5.3	Aquifer Restoration Impacts to Ecological Resources 4.4-21
4.4.5.4	Decommissioning Impacts to Ecological Resources 4.4-21
4.4.6	Air Quality Impacts 4.4-21
4.4.6.1	Construction Impacts to Air Quality 4.4-22
4.4.6.2	Operation Impacts to Air Quality 4.4-23
4.4.6.3	Aquifer Restoration Impacts to Air Quality..... 4.4-23
4.4.6.4	Decommissioning Impacts to Air Quality 4.4-23
4.4.7	Noise Impacts 4.4-24
4.4.7.1	Construction Impacts to Noise 4.4-24
4.4.7.2	Operation Impacts to Noise..... 4.4-24
4.4.7.3	Aquifer Restoration Impacts to Noise..... 4.4-24
4.4.7.4	Decommissioning Impacts to Noise 4.4-25
4.4.8	Historical and Cultural Resources Impacts..... 4.4-25
4.4.8.1	Construction Impacts to Historical and Cultural Resources 4.4-26
4.4.8.2	Operation Impacts to Historical and Cultural Resources 4.4-27
4.4.8.3	Aquifer Restoration Impacts to Historical and Cultural Resources 4.4-27
4.4.8.4	Decommissioning Impacts to Historical and Cultural Resources 4.4-27
4.4.9	Visual/Scenic Resources Impacts 4.4-28
4.4.9.1	Construction Impacts to Visual/Scenic Resources 4.4-28
4.4.9.2	Operation Impacts to Visual/Scenic Resources..... 4.4-28
4.4.9.3	Aquifer Restoration Impacts to Visual/Scenic Resources 4.4-28
4.4.9.4	Decommissioning Impacts to Visual/ Scenic Resources..... 4.4-29

CONTENTS (continued)

Section	Page
4.4.10	Socioeconomic Impacts 4.4-29
4.4.10.1	Construction Impacts to Socioeconomics..... 4.4-29
4.4.10.2	Operation Impacts to Socioeconomics 4.4-31
4.4.10.3	Aquifer Restoration Impacts to Socioeconomics 4.4-32
4.4.10.4	Decommissioning Impacts to Socioeconomics..... 4.4-33
4.4.11	Public and Occupational Health and Safety Impacts 4.4-33
4.4.11.1	Construction Impacts to Public and Occupational Health and Safety 4.4-33
4.4.11.2	Operation Impacts to Public and Occupational Health and Safety 4.4-33
4.4.11.2.1	Radiological Impacts to Public and Occupational Health and Safety for Normal Operation 4.4-33
4.4.11.2.2	Radiological Impacts to Public and Occupational Health and Safety From Accidents 4.4-34
4.4.11.2.3	Non-radiological Impacts to Public and Occupational Health and Safety for Normal Operations 4.4-34
4.4.11.2.4	Non-radiological Impacts to Public and Occupational Health and Safety for Accidents 4.4-34
4.4.11.3	Aquifer Restoration Impacts to Public and Occupational Health and Safety 4.4-34
4.4.11.4	Decommissioning Impacts to Public and Occupational Health and Safety 4.4-35
4.4.12	Waste Management Impacts..... 4.4-35
4.4.12.1	Construction Impacts to Waste Management..... 4.4-35
4.4.12.2	Operation Impacts to Waste Management 4.4-35
4.4.12.3	Aquifer Restoration Impacts to Waste Management .. 4.4-36
4.4.12.4	Decommissioning Impacts to Waste Management..... 4.4-36
4.4.13	References..... 4.4-36
4.5	Northwestern New Mexico Uranium Milling Region 4.5-1
4.5.1	Land Use Impacts 4.5-1
4.5.1.1	Construction Impacts to Land Use 4.5-1
4.5.1.2	Operation Impacts to Land Use..... 4.5-1
4.5.1.3	Aquifer Restoration Impacts to Land Use..... 4.5-2
4.5.1.4	Decommissioning Impacts to Land Use 4.5-2
4.5.2	Transportation Impacts..... 4.5-2
4.5.2.1	Construction Impacts to Transportation..... 4.5-2
4.5.2.2	Operation Impacts to Transportation 4.5-3
4.5.2.3	Aquifer Restoration Impacts to Transportation 4.5-3
4.5.2.4	Decommissioning Impacts to Transportation..... 4.5-3
4.5.3	Geology and Soils Impacts..... 4.5-4
4.5.3.1	Construction Impacts to Geology and Soils 4.5-4
4.5.3.2	Operation Impacts to Geology and Soils 4.5-5
4.5.3.3	Aquifer Restoration Impacts to Geology and Soils 4.5-7
4.5.3.4	Decommissioning Impacts to Geology and Soils..... 4.5-8

CONTENTS (continued)

Section	Page
4.5.4	Water Resources Impacts 4.5-9
4.5.4.1	Surface Water Impacts..... 4.5-9
4.5.4.1.1	Construction Impacts to Surface Water 4.5-9
4.5.4.1.2	Operational Impacts to Surface Water 4.5-9
4.5.4.1.3	Aquifer Restoration Impacts to Surface Waters..... 4.5-9
4.5.4.1.4	Decommissioning Impacts to Surface Water 4.5-10
4.5.4.2	Groundwater Impacts..... 4.5-10
4.5.4.2.1	Construction Impacts to Groundwater 4.5-10
4.5.4.2.2	Operation Impacts to Groundwater 4.5-11
4.5.4.2.2.1	Operation Impacts to Shallow (Near-Surface) Aquifers..... 4.5-11
4.5.4.2.2.2	Operation Impacts to Production and Surrounding Aquifers..... 4.5-12
4.5.4.2.2.3	Operation Impacts to Deep Aquifers Below the Production Aquifers..... 4.5-15
4.5.4.2.3	Aquifer Restoration Impacts to Groundwater..... 4.5-16
4.5.4.2.4	Decommissioning Impacts to Groundwater..... 4.5-18
4.5.5	Ecological Resources Impacts 4.5-18
4.5.5.1	Construction Impacts to Ecological Resources 4.5-18
4.5.5.2	Operation Impacts to Ecological Resources..... 4.5-22
4.5.5.3	Aquifer Restoration Impacts to Ecological Resources 4.5-23
4.5.5.4	Decommissioning Impacts to Ecological Resources 4.5-23
4.5.6	Air Quality Impacts 4.5-23
4.5.6.1	Construction Impacts to Air Quality 4.5-24
4.5.6.2	Operation Impacts to Air Quality 4.5-24
4.5.6.3	Aquifer Restoration Impacts to Air Quality..... 4.5-24
4.5.6.4	Decommissioning Impacts to Air Quality 4.5-25
4.5.7	Noise Impacts 4.5-25
4.5.7.1	Construction Impacts to Noise 4.5-25
4.5.7.2	Operation Impacts to Noise..... 4.5-25
4.5.7.3	Aquifer Restoration Impacts to Noise 4.5-26
4.5.7.4	Decommissioning Impacts to Noise 4.5-26
4.5.8	Historical and Cultural Resources Impacts 4.5-26
4.5.8.1	Construction Impacts to Historical and Cultural Resources 4.5-27
4.5.8.2	Operation Impacts to Historical and Cultural Resources 4.5-28
4.5.8.3	Aquifer Restoration Impacts to Historical and Cultural Resources 4.5-28

CONTENTS (continued)

Section	Page
4.5.8.4	Decommissioning Impacts to Historical and Cultural Resources 4.5-28
4.5.9	Visual/Scenic Resources Impacts 4.5-29
4.5.9.1	Construction Impacts to Visual/Scenic Resources 4.5-29
4.5.9.2	Operation Impacts to Visual/Scenic Resources 4.5-29
4.5.9.3	Aquifer Restoration Impacts to Visual/Scenic Resources 4.5-29
4.5.9.4	Decommissioning Impacts to Visual/Scenic Resources 4.5-30
4.5.10	Socioeconomic Impacts 4.5-30
4.5.10.1	Construction Impacts to Socioeconomics 4.5-30
4.5.10.2	Operation Impacts to Socioeconomics 4.5-32
4.5.10.3	Aquifer Restoration Impacts to Socioeconomics 4.5-33
4.5.10.4	Decommissioning Impacts to Socioeconomics 4.5-33
4.5.11	Public and Occupational Health and Safety Impacts 4.5-34
4.5.11.1	Construction Impacts to Public and Occupational Health and Safety 4.5-34
4.5.11.2	Operation Impacts to Public and Occupational Health and Safety 4.5-34
4.5.11.2.1	Radiological Impacts to Public and Occupational Health and Safety for Normal Operations 4.5-34
4.5.11.2.2	Radiological Impacts to Public and Occupational Health and Safety From Accidents 4.5-34
4.5.11.2.3	Non-radiological Impacts to Public and Occupational Health and Safety for Normal Operations 4.5-35
4.5.11.2.4	Non-radiological Impacts to Public and Occupational Health and Safety for Accidents 4.5-35
4.5.11.3	Aquifer Restoration Impacts to Public and Occupational Health and Safety 4.5-35
4.5.11.4	Decommissioning Impacts to Public and Occupational Health and Safety 4.5-35
4.5.12	Waste Management Impacts 4.5-36
4.5.12.1	Construction Impacts to Waste Management 4.5-36
4.5.12.2	Operation Impacts to Waste Management 4.5-36
4.5.12.3	Aquifer Restoration Impacts to Waste Management .. 4.5-36
4.5.12.4	Decommissioning Impacts to Waste Management 4.5-37
4.5.13	References 4.5-37
5	CUMULATIVE EFFECTS 5-1
5.1	Introduction 5-1
5.2	Other Past, Present, and Reasonably Foreseeable Future Actions in the Four Regions 5-3
5.2.1	Uranium Recovery Sites 5-3
5.2.2	EISs as Indicators of Present and RFFAs 5-3

CONTENTS (continued)

Section	Page
5.3	Concurrent Actions 5-12
5.3.1	Wyoming West Uranium Milling Region 5-12
5.3.2	Wyoming East Uranium Milling Region 5-12
5.3.3	Nebraska-South Dakota-Wyoming Uranium Milling Region 5-18
5.3.4	Northwestern New Mexico Uranium Milling Region 5-22
5.4	Approaches to Conducting a Site-Specific Cumulative Effects Analysis 5-22
5.5	References 5-25
6	ENVIRONMENTAL JUSTICE 6-1
6.1	Environmental Justice Analysis 6-2
6.1.1	Background and Guidance 6-2
6.1.2	Identifying Minority and Low-Income Populations in the Four Geographic Uranium Milling Regions Considered in This GEIS 6-5
6.2	Wyoming West Uranium Milling Region 6-13
6.3	Wyoming East Uranium Milling Region 6-14
6.4	Nebraska-South Dakota-Wyoming Uranium Milling Region 6-14
6.5	Northwestern New Mexico Uranium Milling Region 6-16
6.6	Summary 6-19
6.7	References 6-19
7	POTENTIAL BEST MANAGEMENT PRACTICES, MITIGATION MEASURES, AND MANAGEMENT ACTIONS TO MITIGATE ADVERSE ENVIRONMENTAL IMPACTS 7-1
7.1	Introduction 7-1
7.2	Best Management Practices 7-1
7.3	Management Actions 7-1
7.4	Potential Best Management Practices, Management Actions, and Mitigation Measures 7-2
7.5	References 7-6
8	ENVIRONMENTAL MONITORING ACTIVITIES 8-1
8.1	Introduction 8-1
8.2	Radiological Monitoring 8-1
8.2.1	Airborne Radiation Monitoring Program 8-2
8.2.2	Soils and Sediments Monitoring 8-2
8.2.3	Vegetation, Food, and Fish Monitoring 8-2
8.2.4	Surface Water Monitoring 8-2
8.2.5	Groundwater Monitoring 8-3
8.3	Physiochemical Monitoring 8-3
8.3.1	Groundwater Monitoring 8-3
8.3.1.1	Pre-Operational Groundwater Sampling 8-3
8.3.1.2	Groundwater Quality Monitoring 8-4
8.3.2	Well Field and Pipeline Flow and Pressure Monitoring 8-6
8.4	Ecological Monitoring 8-7
8.5	References 8-7
9	CONSULTATIONS 9-1

CONTENTS (continued)

Section	Page
10 SUMMARY OF ENVIRONMENTAL CONSEQUENCES	10-1
11 LIST OF PREPARERS	11-1
12 GLOSSARY	12-1
APPENDIX A SCOPING SUMMARY REPORT	A-1
APPENDIX B POTENTIALLY APPLICABLE FEDERAL STATUTES, REGULATIONS, AND EXECUTIVE ORDERS	B-1
APPENDIX C SUMMARY OF CONVENTIONAL URANIUM MILLING TECHNOLOGIES	C-1
APPENDIX D CULTURAL AND HISTORICAL RESOURCE MANAGEMENT PROCESSES	D-1
APPENDIX E HAZARDOUS CHEMICALS	E-1
APPENDIX F DESCRIPTION OF PROCESSES FOR REVIEW OF CUMULATIVE EFFECTS	F-1

FIGURES

Figure	Page
1.1-1	Four Geographic Regions Used as a Framework for the Analyses Presented in This GEIS.....
1.1-2	Major Uranium Reserves Within the United States.....
1.4-1	Structure of This GEIS
1.7-1	General Flow Diagram of the NRC Licensing Process for 10 CFR Part 40 Licenses
2.1-1	Simplified Cross-Section of Sandstone Uranium Roll-Front Deposits Formed by Regional Groundwater Migration.....
2.1-2	Schematic Diagram of the Different Types of Tabular Stratabound Uranium Deposits in the Grants Uranium District, New Mexico
2.1-3	Layout of the Crow Butte Uranium Project in Dawes County, Nebraska.....
2.1-4	Well Heads and a Header House at Smith Ranch, Converse County, Wyoming.....
2.3-1	Schematic Diagram of a Well Field Showing Typical Injection/Production Well Patterns, and Monitor Wells
2.3-2	Manifold Inside Well Field Header House at an ISL Facility
2.3-3	Computerized Meter for Monitoring Well Field Flow Rates.....
2.3-4	Cross Section of a Typical Injection, Production, or Monitoring Well Completed Using the Underreamed Method
2.4-1	Idealized Schematic Cross Section to Illustrate Ore-Zone Geology and Lixiviant Migration From an Injection Well to a Production Well
2.4-2	Flow Diagram of an ISL Uranium Recovery Process.....
2.4-3	Typical Ion Exchange Vessels in an ISL Facility.....
2.4-4	A Typical Thickener for an ISL Uranium Processing Facility
2.4-5	Typical Vacuum Dryer for Uranium Yellowcake Processing at an ISL Uranium Processing Facility.....
2.4-6	Labeled and Placarded 208-L [55-gal] Drum Used for Packaging and Shipping Yellowcake
2.5-1	Schematic Diagram of Groundwater Sweep During Aquifer Restoration
2.7-1	Downwind Distance Versus Relative Air Concentration (Which Is Proportional to Dose).....
3.1-1	Wyoming West Uranium Milling Region With Current and Potential ISL Milling Sites
3.1-2	Wyoming East Uranium Milling Region With Current and Potential ISL Milling Sites
3.1-3	Nebraska-South Dakota-Wyoming Uranium Milling Region With Current and Potential ISL Milling Sites.....
3.1-4	New Mexico Uranium Milling Region With Current and Potential ISL Milling Sites.....
3.1-5	Simplified Cross-Section of Sandstone Uranium Roll-Front Deposits Formed by Regional Groundwater Migration.....
3.2-1	Wyoming West Uranium Milling Region General Map With Current and Future Uranium Milling Site
3.2-2	Map Showing Outline of the Wyoming West Uranium Milling Region and Locations of the Crooks Gap Uranium District in the Great Divide Basin and the Gas Hills Uranium District in the Wind River Basin

FIGURES (continued)

Figure	Page
3.2-3	Wyoming West Uranium Milling Region Transportation Corridor..... 3.2-5
3.2-4	Index and Structure Map of Central Wyoming Showing Relation of Sweetwater Arch to the Great Divide Basin and the Wind River Basin..... 3.2-9
3.2-5	Stratigraphic Section of Tertiary Age Formations in the Great Divide Basin and Wind River Basin of Central Wyoming..... 3.2-10
3.2-6	Watersheds in the Wyoming West Uranium Milling Region..... 3.2-14
3.2-7	Ecoregions of the Wyoming West Uranium Milling Region..... 3.2-23
3.2-8	Antelope Wintering Areas for the Wyoming West Uranium Milling Region 3.2-28
3.2-9	Big Horn Wintering Areas for the Wyoming West Uranium Milling Region..... 3.2-29
3.2-10	Elk Wintering Areas for the Wyoming West Uranium Milling Region 3.2-30
3.2-11	Moose Wintering Areas for the Wyoming West Uranium Milling Region..... 3.2-31
3.2-12	Mule Deer Wintering Areas for the Wyoming West Uranium Milling Region.... 3.2-32
3.2-13	Sage-Grouse/Lek Nesting Areas for the Wyoming West Uranium Milling Region 3.2-33
3.2-14	White Tailed Deer Wintering Areas for the Wyoming West Uranium Milling Region 3.2-34
3.2-15	Air Quality Attainment Status for Wyoming and Surrounding Areas 3.2-48
3.2-16	Prevention of Significant Deterioration Class I Areas in Wyoming East Uranium Milling Region and Surrounding Areas..... 3.2-50
3.2-17	Comparison of Noise Levels Associated With Common Activities 3.2-52
3.2-18	General Location of Native American Plains Tribes 3.2-54
3.2-19	Regional Distribution of Native American Tribes in Wyoming, South Dakota, and Nebraska 3.2-60
3.2-20	BLM Visual Resource Classifications for the Wyoming West Uranium Milling Region 3.2-65
3.2-21	Wyoming West Uranium Milling Region With Population..... 3.2-70
3.2-22	Average Annual Background Radiation in the United States..... 3.2-82
3.3-1	Wyoming East Uranium Milling Region General Map With, Past, Current, and Future Uranium Milling Site Locations 3.3-2
3.3-2	Map Showing Outline of the Wyoming East Region and Locations of the Pumpkin Buttes and Monument Hill Districts in the Powder River Basin 3.3-3
3.3-3	Wyoming East Uranium Milling Region General Map With Current and Future Uranium Milling Site Transportation Corridor 3.3-5
3.3-4	Index and Structure Map of East-Central Wyoming Showing Relation of the Sweetwater Arch and Laramie Range to the Powder River Basin and the Shirley Basin 3.3-9
3.3-5	Stratigraphic Section of Tertiary Age Formations in the Powder River Basin and Shirley Basin of Central Wyoming 3.3-10
3.3-6	Watersheds Within the Wyoming East Uranium Milling Region..... 3.3-13
3.3-7	Ecoregions of the Wyoming East Uranium Milling Region..... 3.3-21
3.3-8	Antelope Wintering Area for the Wyoming East Uranium Milling Region 3.3-25
3.3-9	Big Horn Wintering Area for the Wyoming East Uranium Milling Region 3.3-26
3.3-10	Elk Wintering Area for the Wyoming East Uranium Milling Region 3.3-27
3.3-11	Moose Wintering Area for the Wyoming East Uranium Milling Region 3.3-28
3.3-12	Mule Deer Wintering Area for the Wyoming East Uranium Milling Region..... 3.3-29

FIGURES (continued)

Figure	Page
3.3-13 Sage-Grouse/Lek Nesting Areas for the Wyoming East Area Uranium Milling Region	3.3-30
3.3-14 White Tailed Deer Wintering Area for the Wyoming East Uranium Milling Region	3.3-31
3.3-15 Air Quality Attainment Status for Wyoming and Surrounding Areas	3.3-38
3.3-16 Prevention of Significant Deterioration Class I Areas in the Wyoming East Uranium Milling Region and Surrounding Areas	3.3-40
3.3-17 BLM Visual Resource Classifications for the Wyoming East Uranium Milling Region	3.3-44
3.3-18 Wyoming East Uranium Milling Region With Population	3.3-47
3.4-1 Nebraska-South Dakota-Wyoming Uranium Milling Region General Map With Current and Current (Crow Butte, Nebraska) and Potential Future Uranium Milling Site Locations	3.4-2
3.4-2 Map Showing the Nebraska-South Dakota-Wyoming Uranium Milling Region and Uranium Milling Sites in the Black Hills Uranium Districts in South Dakota and Wyoming and in the Crow Butte Uranium District in Nebraska	3.4-3
3.4-3 Nebraska-South Dakota-Wyoming Uranium Milling Region Transportation Corridor	3.4-5
3.4-4 Outcrop Map of the Inyan Kara Group in the Black Hills of Western South Dakota and Northeastern Wyoming Showing the Locations of Principal Uranium Mining Districts	3.4-9
3.4-5 Principal Stratigraphic Units in the Black Hills Area of Western South Dakota and Northeastern Wyoming	3.4-10
3.4-6 Schematic Cross Section Through a Typical Inyan Kara Roll-Front Deposit Showing Differences in U ₃ O ₈ Concentration Between "Fresh" (i.e., Unoxidized) and "Altered" Ground	3.4-12
3.4-7 Bedrock Geology and Major Structural Features of the Crawford Basin	3.4-13
3.4-8 Generalized Stratigraphic Units in the Crow Butte Area of Northwestern Nebraska	3.4-15
3.4-9 Location of Oxidation-Reduction Fronts Detected During Exploration Drilling Within the Chadron Sandstone in Northwestern Nebraska	3.4-16
3.4-10 Watersheds Within the Nebraska-South Dakota-Wyoming Uranium Milling Region	3.4-18
3.4-11 Ecoregions for the Nebraska-South Dakota-Wyoming Uranium Milling Region	3.4-28
3.4-12 Antelope Wintering Areas for the Nebraska-South Dakota-Wyoming Uranium Milling Region	3.4-32
3.4-13 Big Horn Wintering Areas for the Nebraska-South Dakota-Wyoming Uranium Milling Region	3.4-33
3.4-14 Elk Wintering Areas for the Nebraska-South Dakota-Wyoming Uranium Milling Region	3.4-34
3.4-15 Moose Wintering Areas for the Nebraska-South Dakota-Wyoming Uranium Milling Region	3.4-35
3.4-16 Mule Deer Wintering Areas for the Nebraska-South Dakota-Wyoming Uranium Milling Region	3.4-36
3.4-17 Sage-Grouse/Lek Nesting Areas for the Nebraska-South Dakota-Wyoming Uranium Milling Region	3.4-37

FIGURES (continued)

Figure	Page
3.4-18	White Tailed Deer Wintering Areas for the Nebraska-South Dakota-Wyoming Uranium Milling Region..... 3.4-38
3.4-19	Air Quality Attainment Status for Western South Dakota/Nebraska Uranium Milling Region and Surrounding Areas 3.4-50
3.4-20	Prevention of Significant Deterioration Class I Areas in the Western South Dakota/Nebraska Uranium Milling Region and Surrounding Areas 3.4-52
3.4-21	Nebraska-South Dakota-Wyoming Uranium Milling Region With Population... 3.4-73
3.5-1	Northwestern New Mexico Uranium Milling Region General Map with Current and Future Uranium Milling Site Locations 3.5-2
3.5-2	Map Showing Outline of the Northwestern New Mexico Region and the Location of the Grants Uranium District Along the Southern Margin of the San Juan Basin..... 3.5-3
3.5-3	Northwestern New Mexico Uranium Milling Region Transportation Corridor Locations 3.5-5
3.5-4	Index Map of the Grants Uranium District, San Juan Basin, New Mexico, Showing Eight Subdistricts 3.5-8
3.5-5	Generalized Stratigraphic Section of Upper Jurassic and Cretaceous Formations in the Grants Uranium District 3.5-9
3.5-6	Generalized Geologic Section Showing the Stratigraphic Relations of the Morrison Formation Between the Ambrosia Lake and Laguna Areas..... 3.5-10
3.5-7	Schematic Diagram of the Different Types of Uranium Deposits in the Morrison Formation, Grants Uranium District, New Mexico 3.5-12
3.5-8	Watersheds in the Northwestern New Mexico Uranium Milling Region 3.5-15
3.5-9	Ecoregions for the Northwestern New Mexico Uranium Milling Region..... 3.5-22
3.5-10	Elk Winter Habitat and Calving Areas for the Northwestern New Mexico Uranium Milling Region..... 3.5-25
3.5-11	Windrose for Gallup, New Mexico, Airport for 1991..... 3.5-35
3.5-12	Air Quality Attainment Status for the Northwestern New Mexico Uranium Milling Region and Surrounding Areas..... 3.5-37
3.5-13	Prevention of Significant Deterioration Class I Areas in the Northwestern New Mexico Uranium Milling Region and Surrounding Areas 3.5-38
3.5-14	Distribution of Recorded Archaeological Sites in McKinley and Cibola Counties, New Mexico 3.5-41
3.5-15	Paleoindian Sites 3.5-43
3.5-16	Distribution of Archaic-Period Sites..... 3.5-44
3.5-17	Distribution of Ancestral Puebloan Sites 3.5-46
3.5-18	Distribution of Historic Pueblo Sites 3.5-49
3.5-19	Distribution of Navajo Archaeological Sites..... 3.5-51
3.5-20	BLM Visual Resource Classifications for the Northwestern New Mexico Uranium Milling Region..... 3.5-60
3.5-21	Northwestern New Mexico Uranium Milling Region With Population..... 3.5-64
4.2-1	Sound Levels Around a Typical Water Well Work Site (From Reinke, 2005)... 4.2-38
4.2-2	Community Surveys of Noise Annoyance 4.2-39

FIGURES (continued)

Figure	Page
6.1-1 Environmental Justice Process Flow Chart.....	6-3
6.1-2 Affected Minority and Low-Income Population for the Wyoming West Uranium Milling Region.....	6-7
6.1-3 Affected Minority and Low-Income Population for the Wyoming East Uranium Milling Region.....	6-9
6.1-4 Affected Minority and Low-Income Population for the Nebraska-South Dakota- Wyoming Uranium Milling Region.....	6-10
6.1-5 Affected Minority and Low-Income Populations for the Northwestern New Mexico Uranium Milling Region.....	6-11

TABLES

Table	Page
1.7-1	New Source Review Permit Summary Information for Nebraska, New Mexico, South Dakota, and Wyoming..... 1-20
2.2-1	Typical Baseline Water Quality Parameters and Indicators 2-6
2.4-1	Typical Lixiviant Chemistry 2-16
2.6-1	Estimated Decommissioning and Reclamation Waste Volumes (yd ³) for Offsite Disposal, Smith Ranch <i>In-Situ</i> Leach Facility 2-31
2.7-1	Combustion Engine Exhaust Sources for the Gas Hills <i>In-Situ</i> Leach Satellite Facility During Construction, Operations, Reclamation, and Decommissioning .. 2-33
2.7-2	Estimated Particulate (Fugitive Dust) and Gaseous (Diesel Combustion Products) Emissions for the Crownpoint, New Mexico, <i>In-Situ</i> Leach Facility 2-34
2.7-3	Estimated Flow Rates and Constituents in Liquid Waste Steams for the Highland <i>In-Situ</i> Leach Facility 2-36
2.8-1	Estimated Annual Vehicle Trips for Phases of ISL Facility Lifecycle 2-39
2.11-1	Size of Permitted Areas for ISL Facilities..... 2-43
2.11-2	Common Bulk Chemicals Required at the Project Processing Sites 2-44
2.11-3	Onsite Quantities of Process Chemicals at ISL Facilities..... 2-45
2.11-4	Irigaray Post-Uranium Recovery Water Quality 2-48
2.11-5	Baseline Groundwater Conditions, Aquifer Restoration Goals, and Actual Final Restoration Values NRC Approved for the Q-Sand Pilot Well Field, Smith Ranch, Wyoming..... 2-50
3.2-1	Land Ownership and General Use in the Wyoming West Uranium Milling Region..... 3.2-4
3.2-2	Average Annual Daily Traffic Counts for Roads in the Wyoming West Uranium Milling Region 3.2-6
3.2-3	Representative Transportation Routes for Yellowcake Shipments From the Wyoming West Uranium Milling Region..... 3.2-7
3.2-4	Primary Watersheds in the Wyoming West Uranium Milling Region Range of Designated Uses of Water Bodies Within Each Watershed 3.2-13
3.2-5	Native Fish Species Found in Wyoming 3.2-35
3.2-6	Information on Two Climate Stations in the Wyoming West Uranium Milling Region..... 3.2-44
3.2-7	Climate Data for Stations in the Wyoming West Uranium Milling Region..... 3.2-44
3.2-8	National Ambient Air Quality Standards..... 3.2-46
3.2-9	Allowable Prevention of Significant Deterioration Class I and Class II Areas ... 3.2-47
3.2-10	U.S. Environmental Protection Agency Class I Prevention of Significant Deterioration Areas in Wyoming 3.2-49
3.2-11	National Register Listed Properties in Counties Included in the Wyoming West Uranium Milling Region 3.2-58
3.2-12	List of Tribal Contacts for Tribes With Interests in Nebraska, Montana, South Dakota, and Wyoming..... 3.2-62
3.2-13	Summary of the Affected Environment Within the Wyoming West Uranium Milling Region..... 3.2-67
3.2-14	2000 U.S. Bureau of Census Population and Race Categories of the Wyoming West Uranium Milling Region..... 3.2-68
3.2-15	U.S. Bureau of Census State Income Information for the Region Surrounding the Wyoming West Uranium Milling Region..... 3.2-72

TABLES (continued)

Table	Page
3.2-16 U.S. Bureau of Census County and Native American Income Information for the Wyoming West Uranium Milling Region.....	3.2-73
3.2-17 U. S. Bureau of Census Housing Information for Wyoming.....	3.2-75
3.2-18 Employment Structure of the Wind River Indian Reservation Within the Affected Area.....	3.2-78
3.2-19 2007 Sales and Use Tax Distribution of the Affected Counties Within Wyoming West Uranium Milling Region.....	3.2-78
3.2-20 Public Radiation Doses.....	3.2-81
3.3-1 Land Ownership and General Use in the Wyoming East Uranium Milling Region.....	3.3-4
3.3-2 Average Annual Daily Traffic Counts for Roads in the Wyoming East Uranium Milling Region.....	3.3-7
3.3-3 Representative Transportation Routes for Yellowcake Shipments From the Wyoming East Uranium Milling Region.....	3.3-7
3.3-4 Primary Watersheds in the Wyoming East Uranium Milling Region, Range of Designated Uses of Water Bodies Within Each Watershed	3.3-14
3.3-5 Information on Two Climate Stations in the Wyoming East Uranium Milling Region.....	3.3-35
3.3-6 Climate Data for Stations in the Wyoming East Uranium Milling Region	3.3-36
3.3-7 U.S. Environmental Protection Agency Class I Prevention of Significant Deterioration Areas in Wyoming	3.3-39
3.3-8 National Register Listed Properties in Counties Included in the Wyoming East Uranium Milling Region.....	3.3-41
3.3-9 Summary of Affected Environment Within the Wyoming East Uranium Milling Region.....	3.3-45
3.3-10 2000 U.S. Bureau of Census Population and Race Categories of the Wyoming East Uranium Milling Region.....	3.3-46
3.3-11 U.S. Bureau of Census State Income Information for Wyoming East Uranium Milling Region.....	3.3-49
3.3-12 U.S. Bureau of Census County Income Information for Wyoming East Uranium Milling Region.....	3.3-50
3.3-13 U.S. Bureau of Census Housing Information for the Wyoming East Uranium Milling Region.....	3.3-51
3.3-14 2007 Sales and Use Tax Distribution of the Affected Counties Within the Wyoming East Uranium Milling Region	3.3-55
3.4-1 Land Ownership and General Use in the Nebraska-South Dakota-Wyoming Uranium Milling Region.....	3.4-4
3.4-2 Average Annual Daily Traffic Counts for Roads in the Nebraska-South Dakota-Wyoming Uranium Milling Region.....	3.4-7
3.4-3 Representative Transportation Routes for Yellowcake Shipments From the Nebraska-South Dakota-Wyoming Uranium Milling Region	3.4-7
3.4-4 Primary Watersheds in the Nebraska-South Dakota-Wyoming Uranium District and Range of Generic Designated Uses of Water Bodies Within Each Watershed	3.4-19
3.4-5 Fishes of the Angostura Reservoir, Cheyenne River Watershed	3.4-40
3.4-6 Fishes of the Niobrara River Watershed	3.4-42
3.4-7 Information on Three Climate Stations in the Nebraska-South Dakota-Wyoming Uranium Milling Region.....	3.4-47

TABLES (continued)

Table	Page
3.4-8	Climate Data for Stations in the Nebraska-South Dakota- Wyoming Uranium Milling Region..... 3.4-47
3.4-9	U.S. Environmental Protection Agency Class I Prevention of Significant Deterioration Areas in Wyoming, South Dakota, Nebraska, and Montana 3.4-51
3.4-10	National Register Listed Properties in Counties Included in the Nebraska-South Dakota-Wyoming Uranium Milling Region 3.4-59
3.4-11	Summary of Affected Areas Within the Nebraska-South Dakota- Wyoming Uranium Milling Region..... 3.4-68
3.4-12	2000 U.S. Bureau of Census Population and Race Categories of Nebraska ... 3.4-69
3.4-13	2000 U.S. Bureau of Census Population and Race Categories of South Dakota 3.4-70
3.4-14	2000 U.S. Bureau of Census Population and Race Categories of Northwestern Wyoming 3.4-72
3.4-15	U.S. Bureau of Census State Income Information for the Nebraska-South Dakota-Wyoming Uranium Milling Region 3.4-74
3.4-16	U.S. Bureau of Census County and Native American Income Information for the Nebraska-South Dakota-Wyoming Uranium Milling Region 3.4-76
3.4-17	U.S. Bureau of Census Housing Information for the Nebraska Uranium Milling Region..... 3.4-79
3.4-18	U.S. Bureau of Census Housing Information for South Dakota..... 3.4-79
3.4-19	U.S. Bureau of Census Housing Information for the Nebraska-South Dakota- Wyoming Uranium Milling Region..... 3.4-80
3.4-20	Employment Structure of the Pine Ridge Indian Reservation Within the Affected Area 3.4-82
3.4-21	2007 Sales and Use Tax Distribution of the Affected Counties Within Wyoming 3.4-87
3.5-1	Land Ownership and General Use in the Northwestern New Mexico Uranium Milling Region..... 3.5-4
3.5-2	Average Annual Daily Traffic Counts for Roads in the Northwestern New Mexico Uranium Milling Region 3.5-6
3.5-3	Representative Transportation Routes for Yellowcake Shipments From the Northwestern New Mexico Uranium Milling Region 3.5-7
3.5-4	Named Uranium Deposits in New Mexico and Corresponding Watersheds..... 3.5-16
3.5-5	Primary Watersheds in New Mexico, Designated Uses and Known Impairments..... 3.5-16
3.5-6	Native Fish Species Found in New Mexico..... 3.5-26
3.5-7	Information on Two Climate Stations in the Northwestern New Mexico Uranium Milling Region 3.5-33
3.5-8	Climate Data for Stations in the Northwestern New Mexico Uranium Milling Region..... 3.5-33
3.5-9	U.S. Environmental Protection Agency Class I Prevention of Significant Deterioration Areas in New Mexico and Arizona..... 3.5-34
3.5-10	Number of Recorded Sites by Time Period and County..... 3.5-40
3.5-11	Cebolleta Mesa and Pecos Chronologies..... 3.5-45
3.5-12	National Register Listed Properties in Counties Included in the Northwestern New Mexico Uranium Milling Region 3.5-53
3.5-13	Known Culturally Significant Places in McKinley and Cibola Counties..... 3.5-56

TABLES (continued)

Table	Page
3.5-14	2008 Pueblo and Tribal Government Contacts for McKinley and Cibola Counties, New Mexico 3.5-57
3.5-15	Summary of Affected Environment Within the Northwestern New Mexico Uranium Milling Region 3.5-62
3.5-16	2000 U.S. Bureau of Census Population and Race Categories of the Northwestern New Mexico Uranium Milling Region 3.5-63
3.5-17	U.S. Bureau of Census State Income Information for the Northwestern New Mexico Uranium Milling Region 3.5-66
3.5-18	U.S. Bureau of Census County Income Information for the Northwestern New Mexico Uranium Milling Region 3.5-68
3.5-19	U.S. Bureau of Census Housing Information for the Northwestern New Mexico Uranium Milling Region 3.5-69
3.5-20	Employment Structure of Native American Communities Within the Affected Environment of the Northwestern New Mexico Uranium Milling Region..... 3.5-72
3.5-21	Net Taxable Values for Affected Counties Within New Mexico for 2006 3.5-73
3.5-22	Percent Change in Tax Values From 2005 to 2006 for the Affected Counties Within New Mexico 3.5-73
3.5-23	Percent Distribution of New Mexico Property Tax Obligations Within Affected Counties for 2006 3.5-73
4.2-1	Average Noise Levels at 15 m [50 ft] From Representative Construction Heavy Equipment 4.2-37
4.2-2	Dose to Offsite Receptors From ISL Facilities 4.2-53
4.2-3	Pertinent Regulations for Chemicals Used at ISL Facilities 4.2-58
5.2-1	Past, Existing, and Potential Uranium Recovery Sites in Wyoming, South Dakota, Nebraska, and New Mexico 5-4
5.2-2	Draft and Final Environmental Impact Statements (EISs) to the Wyoming West Uranium Milling Region 5-7
5.2-3	Draft and Final Environmental Impact Statements (EISs) Related to the Wyoming East Uranium Milling Region..... 5-8
5.2-4	Draft and Final Programmatic or Large-Scale Environmental Impact Statements (EISs) Related to One or Both of the Wyoming Regional Study Areas 5-8
5.2-5	Draft and Final Environmental Impact Statements (EISs) Related to Nebraska-South Dakota-Wyoming Uranium Milling Region..... 5-9
5.2-6	Draft and Final Environmental Impact Statements (EISs) Related to the Northwestern New Mexico Uranium Milling Region 5-11
5.3-1	Other Actions Concurrent With Uranium Recovery in the Wyoming West Uranium Milling Region 5-13
5.3-2	Coal Mining Project as Identified by the Wyoming Mining Association..... 5-15
5.3-3	Other Actions Relations to or Conflicting With Uranium Recovery in the Wyoming East Uranium Milling Region..... 5-16
5.3-4	Other Actions Concurrent With Uranium Recovery in Nebraska-South Dakota-Wyoming Uranium Milling Region..... 5-19
5.3-5	Other Actions Concurrent With Uranium Recovery in the Northwestern New Mexico Uranium Milling Region 5-22

TABLES (continued)

Table	Page
6.1-1	Minority and Low-Income Populations in the Four Geographic Uranium Milling Regions Considered in This Generic Environmental Impact Statement 6-6
7.4-1	Summary of Potential Best Management Practices and Management Actions 7-3
8.2-1	Typical Baseline Water Quality Parameters and Indicators for Groundwater 8-4
10-1	Summary of Impacts for the Wyoming West Uranium Milling Region 10-3
10-2	Summary of Impacts for the Wyoming East Uranium Milling Region 10-18
10-3	Summary of Impacts for the Nebraska-South Dakota-Wyoming Uranium Milling Region..... 10-33
10-4	Summary of Impacts for the Northwestern New Mexico Uranium Milling Region..... 10-48

EXECUTIVE SUMMARY

PURPOSE AND NEED

NRC prepared this Draft Generic Environmental Impact Statement (Draft GEIS) to identify and evaluate the potential environmental impacts associated with the construction, operation, aquifer restoration, and decommissioning of *in-situ* leach (ISL) uranium recovery facilities. Based on discussions between uranium mining companies and the NRC staff, these facilities potentially could be located in portions of Wyoming, Nebraska, South Dakota, and New Mexico, which are States where NRC has regulatory authority over the licensing of uranium recovery facilities. Given that the large majority of these potential license applications would involve use of the ISL process and would be submitted over a relatively short period of time, NRC decided to prepare a GEIS to support an efficient and consistent approach to reviewing site-specific license applications for ISL facilities. The NRC staff plans to use the GEIS as a starting point for its National Environmental Policy Act (NEPA) analyses for site-specific license applications for new ISL facilities. Additionally, the NRC staff plans to use the GEIS, along with applicable previous site-specific environmental review documents, in its NEPA analysis for the restart or expansions of existing facilities.

Uranium milling techniques are designed to recover the uranium from uranium-bearing ores. Various physical and chemical processes may be used, and selection of the uranium milling technique depends on the physical and chemical characteristics of the ore deposit and the attendant cost considerations. Generally, the ISL process is used to recover uranium from low-grade ores or deeper deposits that are not economically recoverable by conventional mining and milling techniques. In this process, a leaching agent, such as oxygen with sodium carbonate, is injected through wells into the subsurface ore body to dissolve the uranium. The leach solution is pumped from there to the surface processing plant and then ion exchange separates the uranium from the solution. After additional purification and drying, the uranium in the form of U_3O_8 (also known as "yellowcake") is placed in 55-gallon drums prior to shipment offsite.

THE PROPOSED FEDERAL ACTION AND ALTERNATIVES

In States where NRC is the regulatory authority over the licensing of uranium milling (including the ISL process), NRC has a statutory obligation to assess each site-specific license application to ensure it complies with NRC regulations before issuing a license. The proposed federal action is to prepare a GEIS that identifies and evaluates the potential environmental impacts associated with the construction, operation, aquifer restoration, and decommissioning of ISL milling facilities in portions of Wyoming, Nebraska, South Dakota, and New Mexico. As stated above, NRC intends to make use of the GEIS during subsequent site-specific ISL licensing actions.

A range of alternatives to the proposed action was evaluated for inclusion in the Draft GEIS. The No-Action alternative was included in the detailed impact analysis. In the No-Action Alternative, no ISL facilities would be licensed, and therefore constructed and operated, in the four uranium milling regions considered in this Draft GEIS. The environment in these regions would not be affected by uranium extraction, although other ongoing and future non-ISL activities would continue as planned.

Alternative methods for milling uranium were considered as possible alternatives to the ISL process. As stated previously, not all uranium deposits are suitable for ISL extraction. For example, if the uranium mineralization is above the saturated zone (i.e., all of the pore spaces in

the ore-bearing rock are not filled with water) ISL techniques may not be appropriate. Likewise, if the ore is not located in a porous and permeable rock unit, it will not be accessible to the leach solution used in the ISL process. Because ISL techniques may not be appropriate in these circumstances, conventional mining (underground or open-pit/surface mining) and milling techniques (e.g., heap leaching) are possible viable alternative technologies.

Inasmuch as the suitability and practicality of using alternative milling methodologies depends upon site-specific conditions, a generic discussion of alternative milling methodologies is not appropriate. Accordingly, this Draft GEIS does not contain a detailed analysis of alternative milling methodologies. A detailed analysis of alternative milling methodologies that can be applied at a specific site will be addressed in NRC's site-specific environmental review for individual ISL license applications.

In addition, it should be noted that previous analyses have indicated that the potential environmental impacts associated with conventional uranium milling operations are significant, because the mill tailings, or waste, are a significant source of radon and radon progeny. For this reason, NRC has made a policy decision to prepare site-specific EISs for applications for a new, or restart of a former, conventional or heap leach facility, as required under 10 CFR 51.20(b)(8).

APPROACH

NRC developed this Draft GEIS, based on NRC's experience in licensing and regulating ISL facilities gained during the past 30 years. In the Draft GEIS, NRC does not consider specific facilities, but rather provides an assessment of potential environmental impacts associated with ISL facilities that might be located in four regions of the western United States. These regions are used as a framework for discussions in this Draft GEIS, and were identified based on several considerations, including:

- Past and existing uranium milling sites are located within States where NRC has regulatory authority over uranium recovery;
- Potential new sites are identified based on NRC's understanding of where the uranium recovery industry has plans to develop uranium deposits using ISL technology; and
- Locations of historical uranium deposits within portions of Wyoming, Nebraska, South Dakota, and New Mexico.

Using these criteria, four geographic regions were identified (Figure ES-1). For the purpose of this Draft GEIS, these regions are titled

- Wyoming West Uranium Milling Region;
- Wyoming East Uranium Milling Region;
- Nebraska-South Dakota-Wyoming Uranium Milling Region; and
- Northwestern New Mexico Uranium Milling Region.

The foundation of the environmental impact assessment in the Draft GEIS is based on (1) the historical operations of NRC-licensed ISL facilities and (2) the affected environment in each of the four regions. The structure of the GEIS is presented in Figure ES-2.




Figure ES-1. Location of Four Geographic Regions Used as a Framework for the Analyses Presented in this GEIS




Figure ES-2. Structure of this GEIS

Chapter 2 of the Draft GEIS provides a description of the ISL process, addressing construction, operation, aquifer restoration, and decommissioning of an ISL facility. This section also discusses financial assurance, whereby the licensee or applicant establishes a bond or other financial mechanism prior to operations to ensure that sufficient funds are available to complete aquifer restoration, decommissioning, and reclamation activities.

Chapter 3 of the Draft GEIS describes the affected environment in each uranium milling region using the environmental resource areas and topics identified through public scoping comments on the GEIS and from NRC guidance to its staff found in NUREG-1748, "Environmental Review Guidance for Licensing Actions Associated With NMSS Programs," issued by NRC in 2003.

Chapter 4 of the GEIS provides an evaluation of the potential environmental impacts of constructing, operating, aquifer restoration, and decommissioning at an ISL facility in each of the four uranium milling regions. In essence, this involves placing an ISL facility with the characteristics described in Chapter 2 of the Draft GEIS within each of the four regional areas described in Chapter 3 and describing and evaluating the potential impacts in each region separately. The potential environmental impacts are evaluated for the different stages in the ISL process: construction, operation, aquifer restoration, and decommissioning. Impacts are examined for the resource areas identified in the description of the affected environment. These resource areas are:

- Land use
- Transportation
- Geology and soils
- Water resources
- Ecology
- Air Quality
- Noise
- Historical and cultural resource
- Visual and scenic resources
- Socioeconomic
- Public and occupational health

NRC identified a number of other issues that helped in the evaluation of the potential environmental impacts of an ISL facility. These issues include:

- **Applicable Statutes, Regulations and Agencies.** Various statutes, regulations, and implementing agencies at the federal, state, tribal and local levels that have a role in regulating ISL facilities are identified and discussed.
- **Waste Management.** Potential impacts from the generation, handling, treatment, and final disposal of chemical, radiological, and municipal wastes are addressed.
- **Accidents.** Potential accident conditions are assessed in the Draft GEIS. This includes consideration of a range of possible accidents and estimation of their consequences including: well field leaks and spills, excursions, processing chemical spills, and ion exchange resin and yellowcake transportation accidents.
- **Environmental Justice.** Although not required for a GEIS, to facilitate subsequent site-specific analyses, this Draft GEIS provides a first order definition of minority and low income populations. Early consultations will be initiated with some of these populations, and the potential for disproportionately high and adverse impacts from future ISL licensing in the uranium milling regions will be evaluated.
- **Cumulative Impacts.** The Draft GEIS addresses cumulative impacts from proposed ISL facility construction, operation, ground water restoration, and decommissioning on all

aspects of the affected environment, considering the impacts from past, present, and reasonably foreseeable future actions in the uranium milling regions.

- **Monitoring.** The Draft GEIS discusses various monitoring methodologies and techniques used to detect and mitigate the spread of radiological and non-radiological contaminants beyond ISL facility boundaries.

SIGNIFICANCE OF LEVELS

In the Draft GEIS, NRC has categorized the potential environmental impacts using significance levels. According to the Council on Environmental Quality, the significance of impacts is determined by examining both context and intensity (40 CFR 1508.27). Context is related to the affected region, the affected interests, and the locality, while intensity refers to the severity of the impact, which is based on a number of considerations. In this Draft GEIS, the NRC used the significance levels identified in NUREG-1748:

- **SMALL Impact:** The environmental effects are not detectable or are so minor that they will neither destabilize nor noticeably alter any important attribute of the resource considered.
- **MODERATE Impact:** The environmental effects are sufficient to alter noticeably, but not destabilize, important attributes of the resource considered.
- **LARGE Impact:** The environmental effects are clearly noticeable and are sufficient to destabilize important attributes of the resource considered.

SUMMARY OF IMPACTS

As discussed previously, Chapter 4 of the Draft GEIS provides NRC's evaluation of the potential environmental impacts of the construction, operation, aquifer restoration, and decommissioning at an ISL facility in each of the four uranium milling regions. A summary of this evaluation by environmental resource area and phase of the ISL facility lifecycle is provided below.

Land Use Impacts

CONSTRUCTION—Land use impacts could occur from land disturbances (including alterations of ecological cultural or historic resources) and access restrictions (including limitations of other mineral extraction activities, grazing activities, or recreational activities). The potential for land use conflicts could increase in areas with higher percentages of private land ownership and Native American land ownership or in areas with a complex patchwork of land ownership. Land disturbances during construction would be temporary and limited to small areas within permitted areas. Well sites, staging areas, and trenches would be reseeded and restored. Unpaved access roads would remain in use until decommissioning. Competing access to mineral rights could be either delayed for the duration of the in-situ leach (ISL) project or be intermixed with ISL operations (e.g., oil and gas exploration). Changes to land use access including grazing restrictions and impacts on recreational activities would be limited due to the small size of restricted areas, temporary nature of restrictions, and availability of other land for these activities. Ecological, historical, and cultural resources could be affected, but would be protected by careful planning and surveying to help identify resources and avoid or mitigate impacts. For all land use aspects except ecological, historical and cultural resources, the potential impacts would be SMALL. Due to the potential for unidentified resources to be altered

or destroyed during excavation, drilling, and grading, the potential impacts to ecological, historical or cultural resources would be SMALL to LARGE, depending on local conditions.

OPERATION—The types of land use impacts for operational activities would be similar to construction impacts regarding access restrictions because the infrastructure would be in place. Additional land disturbances would not occur from conducting operational activities. Because access restriction and land disturbance related impacts would be similar to, or less than, for construction, the overall potential impacts to land use from operational activities would be SMALL.

AQUIFER RESTORATION—Due to the use of the same infrastructure, land use impacts would be similar to operations during aquifer restoration, although some operational activities would diminish—SMALL.

DECOMMISSIONING—Land use impacts would be similar to those described for construction with a temporary increase in land-disturbing activities for dismantling, removing, and disposing of facilities, equipment, and excavated contaminated soils. Reclamation of land to preexisting conditions and uses would help mitigate potential impacts—SMALL to MODERATE during decommissioning, and SMALL once decommissioning is completed.

Transportation Impacts

CONSTRUCTION—Low magnitude traffic generated by ISL construction relative to local traffic counts would not significantly increase traffic or accidents on many of the roads in the region. Existing low traffic roads could be moderately impacted by the additional worker commuting traffic during periods of peak employment. This impact would be expected to be more pronounced in areas with relatively lower traffic counts. Moderate dust, noise, and incidental wildlife or livestock kill impacts would be possible on, or near, site access roads (dust in particular for unpaved access roads)—SMALL to MODERATE.

OPERATION— Low magnitude traffic relative to local traffic counts on most roads would not significantly increase traffic or accidents. Existing low traffic roads could be moderately impacted by commuting traffic during periods of peak employment including dust, noise, and possible incidental wildlife or livestock kill impacts on or near site access roads. High consequences would be possible for a severe accident involving transportation of hazardous chemicals in a populated area. However, the probability of such accidents occurring would be low owing to the small number of shipments, comprehensive regulatory controls, and use of best management practices. For radioactive material shipments (yellowcake product, ion exchange resins, waste materials), compliance with transportation regulations would limit radiological risk for normal operations. Low radiological risk is estimated for accident conditions. Emergency response protocols would help mitigate long-term consequences of severe accidents involving release of uranium—SMALL to MODERATE.

AQUIFER RESTORATION—The magnitude of transportation activities would be lower than for construction and operations, with the exception of workforce commuting which could have moderate impacts on, or in the vicinity of, existing low traffic roads—SMALL to MODERATE.

DECOMMISSIONING—The types of transportation activities and, therefore, the types of impacts would be similar to those discussed for construction and operations except the magnitude of transportation activities (e.g., number and types of waste and supply shipments, no yellowcake shipments) from decommissioning could be lower than for operations. Accident risks would be bounded by operations yellowcake transportation risk estimates—SMALL.

Geology and Soils Impacts

CONSTRUCTION—Disturbance to soil would occur from construction (clearing, excavation, drilling, trenching, road construction); however, such disturbances would be expected to be temporary, disturbed areas would be **SMALL** (approximately 10 percent of the total site area), and potential impacts would be mitigated by using best management practices. A large portion of the well fields, trenches, and access roads would be restored and reseeded after construction. Excavated soils would be stockpiled, seeded, and stored onsite until needed for reclamation fill. No impacts to subsurface geological strata would be likely—**SMALL**.

OPERATION—Temporary contamination or alteration of soils would be likely from operational leaks and spills and possible from transportation, use of evaporation ponds, or land application of treated waste water. However, detection and response to leaks and spills (e.g., soil cleanup), monitoring of treated waste water, and eventual survey and decommissioning of all potentially impacted soils would limit the magnitude of overall impacts to soils—**SMALL**.

AQUIFER RESTORATION—Impacts to geology and soils from aquifer restoration activities would be similar to impacts from operations due to use of the same infrastructure and similar activities conducted (e.g., well field operation, transfer lines, liquid effluent treatment and disposal)—**SMALL**.

DECOMMISSIONING—Impacts to geology and soils from decommissioning would be similar to impacts from construction. Activities to cleanup, re-contour and reclaim disturbed lands during decommissioning would mitigate long-term impacts to soils—**SMALL**.

Surface Water Impacts

CONSTRUCTION—Impacts to surface waters and related habitats from construction (road crossings, filling, erosion, runoff, spills or leaks of fuels and lubricants for construction equipment) would be mitigated through proper planning, design, construction methods, and best management practices. Some impacts directly related to the construction activities would be temporary and limited to the duration of the construction period. U.S. Army Corps of Engineers permits may be required when filling and crossing of wetlands. Temporary changes to spring and stream flow from grading and changes in topography and natural drainage patterns could be mitigated or restored after the construction phase. Impacts from incidental spills of drilling fluids into local streams could occur, but would be temporary, due to the use of mitigation measures. Impacts from roads, parking areas, buildings on recharge to shallow aquifers would be **SMALL**, owing to the limited area of impervious surfaces proposed. Impacts from infiltration of drilling fluids into the local aquifer would be localized, small, and temporary—**SMALL** to **MODERATE** depending on site-specific characteristics.

OPERATION—Through permitting processes, federal and state agencies regulate the discharge of storm water runoff and the discharge of process water. Impacts from these discharges would be mitigated as licensees would within the conditions of their permits. Expansion of facilities or pipelines during operations would generate impacts similar to construction—**SMALL** to **MODERATE** depending on site-specific characteristics.

AQUIFER RESTORATION—Impacts from aquifer restoration would be similar to impacts from operations due to use of the same (in-place) infrastructure and similar activities conducted (e.g., well field operation, transfer lines, water treatment, storm water runoff)—**SMALL** to **MODERATE** depending on site-specific characteristics.

DECOMMISSIONING—Impacts from decommissioning would be similar to impacts from construction. Activities to clean up, re-contour and reclaim disturbed lands during decommissioning would mitigate long-term impacts to surface waters—SMALL to MODERATE depending on site-specific characteristics.

Groundwater Impacts

CONSTRUCTION—Water use impacts would be limited by the small volumes of groundwater used for routine activities such as dust suppression, mixing cements, and drilling support over short and intermittent periods. Contamination of groundwater from construction activities would be mitigated by best management practices—SMALL to LARGE, depending on site-specific conditions.

OPERATION—Potential impacts to shallow aquifers can occur from leaks or spills from surface facilities and equipment. Shallow aquifers are important sources of drinking water in some areas of the four uranium milling regions. Potential impacts to the ore-bearing and surrounding aquifers include consumptive water use and degradation of water quality (from normal production activities, off-normal excursion events, and deep well injection disposal practices). Consumptive use impacts from withdrawal of groundwater would occur because approximately 1 to 3 percent of pumped groundwater is not returned to the aquifer (e.g., process bleed). That amount of water lost could be reduced substantially by available treatment methods (e.g., reverse osmosis, brine concentration). Effects of water withdrawal on surface water would be expected to be SMALL as the ore zone normally occurs in a confined aquifer. Estimated drawdown effects vary depending on site conditions and water treatment technology applied. Excursions of lixiviant and mobilized chemical constituents could occur from failure of well seals or other operational conditions that result in incomplete recovery of lixiviant. Well seal related excursions would be detected by the groundwater monitoring system and periodic well mechanical integrity testing and impacts would be expected to be mitigated during operation or aquifer restoration. Other excursions could result in plumes of mobilized uranium and heavy metals extending beyond the mineralization zone. The magnitude of potential impacts from vertical excursions would vary depending on site-specific conditions. To reduce the likelihood and consequences of potential excursions at ISL facilities, NRC requires licensees to take preventative measures prior to starting operations including well tests, monitoring, and development of procedures that include excursion response measures and reporting requirements. Alterations of ore body aquifer chemistry would be SMALL, because the aquifer would: (1) be confined, (2) not be a potential drinking water source, and (3) be expected to be restored within statistical range of preoperational baseline water quality during the restoration period. Potential environmental impacts to confined deep aquifers below the production aquifers from deep well injection of processing wastes would be addressed by the underground injection permitting process regulated by the states—SMALL to LARGE, depending on site-specific conditions.

AQUIFER RESTORATION—Potential impacts would be from consumptive use and potential deep disposal of brine slurries after reverse osmosis, if applicable. The volume of water removed from the aquifer and related impacts would be dependent on site-specific conditions and the type of water treatment technology the facility uses. In some cases, groundwater consumptive use for the aquifer restoration has been reported to be less than groundwater use during the ISL operation and drawdowns due to aquifer restorations have been smaller than drawdown caused by ISL operations. Potential environmental impacts associated with water consumption during aquifer restorations are determined by: (1) the restoration techniques chosen, (2) the volume of water to be used, (3) the severity and extent of the contamination,

and (4) the current and future use of the production and surrounding aquifers near the ISL facility or at the regional scale—SMALL to LARGE, depending on site-specific conditions.

DECOMMISSIONING—Potential impacts from decommissioning would be similar to construction (water use, spills) with an additional potential to mobilize contaminants during demolition and cleanup activities. Contamination of groundwater from decommissioning activities would be mitigated by implementation of an NRC-approved decommissioning plan and use of best management practices—SMALL.

Terrestrial Ecology Impacts

CONSTRUCTION—Potential terrestrial ecology impacts would include the removal of vegetation from the well fields, the milling site, the modification of existing vegetative communities, the loss of sensitive plants and habitats from clearing and grading, and the potential spread of invasive species and noxious weed populations. These impacts would be expected to be temporary because restoration and reseeding occur rapidly after the end of construction. Introduction of invasive species and noxious weeds would be mitigated by restoration and reseeding after construction. Shrub and tree removal and loss would take longer to restore. Construction noise could affect reproductive success of sage grouse leks by interfering with mating calls. Temporary displacement of some animal species would also occur. Critical wintering and year-long ranges are important to survival of both big game and sage grouse. Raptors breeding onsite may be impacted by construction activities or milling operations, depending on the time of year construction occurs. Wildlife habitat fragmentation, temporary displacement of animal species, and direct or indirect mortalities would be possible. Implementation of wildlife surveys and mitigation measures following established guidelines would limit impacts. The magnitude of impacts depends on whether a new facility is being licensed or an existing facility is being extended—SMALL to MODERATE, depending on site-specific habitat conditions.

OPERATION—Habitats could be altered by operations (fencing, traffic, noise), and individual takes could occur due to conflicts between species habitat and operations. Access to crucial wintering habitat and water could be limited by fencing. However, the State of Wyoming Game and Fish Department specifies fencing construction techniques to minimize impediments to big game movement. Migratory birds could be affected by exposure to constituents in evaporation ponds, but perimeter fencing, netting, and periodic wildlife surveys (e.g., raptor surveys) would limit impacts. Temporary contamination or alteration of soils would be likely from operational leaks and spills and possible from transportation or land application of treated waste water. However, detection and response to leaks and spills (e.g., soil cleanup) and eventual survey and decommissioning of all potentially impacted soil limits the magnitude of overall impacts to terrestrial ecology. Mitigation measures such as perimeter fencing, netting, alternative sites, and periodic wildlife surveys would reduce overall impacts—SMALL.

AQUIFER RESTORATION—Impacts include habitat disruption, but existing (in-place) infrastructure would be used during aquifer restoration, with little additional ground disturbance. Migratory birds could be affected by exposure to constituents in evaporation ponds, but perimeter fencing, netting, and periodic wildlife surveys (e.g., raptor surveys) would limit impacts. Contamination of soils could be result from leaks and spills, and land application of treated waste water. However, detection and response techniques, and eventual survey and decommissioning of all potentially impacted soils, would limit the magnitude of overall impacts to terrestrial ecology. Mitigation measures such as perimeter fencing, netting, alternative sites, and periodic wildlife surveys would reduce overall impacts—SMALL.

DECOMMISSIONING—During decommissioning and reclamation, there would be a temporary disturbance to land (e.g., excavating soils, buried piping, removal of structures). However, re-vegetation and re-contouring would restore habitat altered during construction and operations. Wildlife would be temporarily displaced, but are expected to return after decommissioning and reclamation are completed and vegetation and habitat reestablished—**SMALL**.

Aquatic Ecology Impacts

CONSTRUCTION—Clearing and grading activities associated with construction could result in a temporary increase in sediment load in local streams, but aquatic species would recover quickly as sediment load decreases. Clearing of riparian vegetation could affect light and temperature of water. Construction impacts to wetlands would be identified and managed through U.S. Army Corps of Engineers permits, as appropriate. Construction impacts to surface waters and aquatic species would be temporary and mitigated by best management practices—**SMALL**.

OPERATION—Impacts could result from spills or releases into surface water. Impacts would be minimized by spill prevention, identification and response programs, and National Pollutant Discharge Elimination System (NPDES) permit requirements—**SMALL**.

AQUIFER RESTORATION—Activities would use existing (in-place) infrastructure, and impacts could result from spills or releases of untreated groundwater. Impacts would be minimized by spill prevention, identification, and response programs, and NPDES permit requirements—**SMALL**.

DECOMMISSIONING—Decommissioning and reclamation activities could result in temporary increases in sediment load in local streams, but aquatic species would recover quickly as sediment load decreases. With completion of decommissioning, re-vegetation, and re-contouring, habitat would be reestablished and impacts would, therefore, be limited—**SMALL**.

Threatened and Endangered Species Impacts

CONSTRUCTION—Numerous threatened and endangered species and state species of concern are located in the four uranium milling regions. Small fragmentation of habitats would occur, but most species readapt quickly. The magnitude of impact would depend on the size of a new facility or extension to an existing facility and the amount of land disturbance. Inventory of threatened or endangered species would be developed during site-specific reviews to identify unique or special habitats, and Endangered Species Act consultations conducted with the U.S. Fish and Wildlife Service would reduce impacts—**SMALL to MODERATE to LARGE**—depending on site-specific habitat and presence of threatened or endangered species.

OPERATION—Impacts could result from individual takes due to conflicts with operations. Small fragmentation of habitats would occur, but most species readapt quickly. The magnitude of impact would depend on the size of a new facility or extension to an existing facility and the amount of land disturbance. Impacts could potentially result from spills or permitted effluents, but would be minimized through the use of spill prevention measures, identification and response programs, and NPDES permit requirements. Inventory of threatened or endangered species developed during site-specific reviews would identify unique or special habitats, and Endangered Species Act consultations conducted with the U.S. Fish and Wildlife Service would assist in reducing impacts—**SMALL to MODERATE**—depending on site-specific habitat and presence of threatened or endangered species.

AQUIFER RESTORATION—Impacts could result from individual takes due to conflicts with aquifer restoration activities (equipment, traffic). Existing (in-place) infrastructure would be used during aquifer restoration, so additional land-disturbing activities and habitat fragmentation would not be anticipated. Impacts may result from spills or releases of treated or untreated groundwater, but impacts would be minimized through the use of spill prevention measures, identification, and response programs, and NPDES permit requirements. Inventory of threatened or endangered species would be developed during site-specific reviews to identify unique or special habitats, and Endangered Species Act consultations with the U.S. Fish and Wildlife Service would assist in reducing impacts—SMALL.

DECOMMISSIONING—Impacts resulting from individual takes would occur due to conflicts with decommissioning activities (equipment, traffic). Temporary land disturbance would occur as structures are demolished and removed and the ground surface is re-contoured. Inventory of threatened or endangered species developed during site-specific environmental review of the decommissioning plan would identify unique or special habitats, and Endangered Species Act consultations with the U.S. Fish and Wildlife Service would assist in reducing impacts. With completion of decommissioning, re-vegetation, and re-contouring, habitat would be reestablished and impacts would, therefore, be limited—SMALL.

Air Quality Impacts

CONSTRUCTION—Fugitive dust and combustion (vehicle and diesel equipment) emissions during land-disturbing activities associated with construction would be small, short-term, and reduced through best management practices (e.g., dust suppression). For example, estimated fugitive dust emissions during ISL construction is less than 2 percent of the National Ambient Air Quality Standards (NAAQS) for $PM_{2.5}$ and less than 1 percent for PM_{10} . For NAAQS attainment areas, non-radiological air quality impacts would be SMALL. A Prevention of Significant Deterioration (PSD) Class I area exists in only one of the four regions (Wind Cave National Park in the Nebraska-South Dakota-Wyoming Region). Here, more stringent air quality standards would apply to a facility that impacts the air quality of that area. If impacts were initially assessed at a higher significance level, permit requirements would impose conditions or mitigation measures to reduce impacts—SMALL.

OPERATION—Radiological impacts can result from dust releases from drying of lixiviant pipeline spills, radon releases from well system relief valves, resin transfer, or elution, and gaseous/particulate emissions from yellowcake dryers. Only small amounts of low dose materials would be expected to be released based on operational controls and rapid response to spills. Required spill prevention, control, and response procedures would be used to minimize impacts from spills. HEPA filters and vacuum dryer designs reduce particulate emissions from operations and ventilation reduces radon buildup during operations. Compliance with the NRC-required radiation monitoring program would ensure releases are within regulatory limits. Other potential non-radiological emissions during operations include fugitive dust and fuel from equipment, maintenance, transport trucks, and other vehicles. For NAAQS attainment areas, non-radiological air quality impacts would be SMALL. A PSD Class I area is located in the Nebraska-South Dakota-Wyoming Region (Wind Cave National Park). More stringent air quality standards would apply to a facility that impacts the air quality of that area. If impacts were initially assessed at a higher significance level, permit requirements would impose conditions or mitigation measures to reduce impacts—SMALL.

AQUIFER RESTORATION—Because the same infrastructure is used, air quality impacts are expected to be similar to, or less than, during operations. For NAAQS attainment areas, non-radiological air quality impacts would be SMALL. Where a PSD Class I area exists, such as the

Wind Cave National Park in the Nebraska-South Dakota-Wyoming Region, more stringent air quality standards would apply to a facility that impacts the air quality of that area. If impacts were initially assessed at a higher significance level, permit requirements would impose conditions or mitigation measures to reduce impacts—SMALL.

DECOMMISSIONING—Fugitive dust, vehicle, and diesel emissions during land-disturbing activities associated with decommissioning would be similar to, or less than, those associated with construction, short-term, and reduced through best management practices (e.g., dust suppression). Potential impacts would decrease as decommissioning and reclamation of disturbed areas are completed. For NAAQS attainment areas, non-radiological air quality impacts would be SMALL. However, where a PSD Class I area exists (Wind Cave National Park, in the Nebraska-South Dakota-Wyoming Region), more stringent air quality standards would apply to a facility that impacts the air quality of that area. If impacts were initially assessed at a higher significance level, permit requirements would impose conditions or mitigation measures to reduce impacts—SMALL.

Noise Impacts

CONSTRUCTION—Noise generated during construction would be noticeable in proximity to operating equipment, but would be temporary (typically daytime only). Administrative and engineering controls would be used to maintain noise levels in work areas below Occupational Health and Safety Administration (OSHA) regulatory limits and mitigated by use of personal hearing protection. Traffic noise during construction (commuting workers, truck shipments to and from the facility, and construction equipment such as trucks, bulldozers, and compressors) would be localized, limited to highways in the vicinity of the site, access roads within the site, and roads in the well fields. Relative increases in traffic levels would be SMALL for the larger roads, but may be MODERATE for lightly traveled rural roads through smaller communities. Noise may also adversely affect wildlife habitat and reproductive success in immediate vicinity of construction activities. Noise levels decrease with distance, and at distances more than about 300 m [1,000 ft], ambient noise levels would return to background. Wildlife avoid construction areas because of noise and human activity. All of the uranium districts are located more than 300 m [1,000 ft] from the closest community. As a result, noise impacts would be—SMALL to MODERATE.

OPERATION—Noise-generating activities in the central uranium processing facility would be indoors, reducing offsite sound levels. Well field equipment (e.g., pumps, compressors) would be contained within structures (e.g., header houses, satellite facilities) also reducing sound levels to offsite receptors. Administrative and engineering controls would be used to maintain noise levels in work areas below OSHA regulatory limits and mitigated by use of personal hearing protection. Traffic noise from commuting workers, truck shipments to and from the facility, and facility equipment would be expected to be localized, limited to highways in the vicinity of the site, access roads within the site, and roads in well fields. Relative increases in traffic levels would be SMALL for the larger roads, but may be MODERATE for lightly traveled rural roads through smaller communities. Most noise would be generated indoors and mitigated by regulatory compliance and best management practices. Noise from trucks and other vehicles are typically of short duration. Also, noise usually is not discernable to offsite receptors at distances of more than 300 m [1,000 ft]. All the uranium districts are located more than 300 m [1,000 ft] from the closest community—SMALL to MODERATE.

AQUIFER RESTORATION—Noise generation is expected to be less than during construction and operations. Pumps and other well field equipment contained in buildings reduce sound levels to offsite receptors. Existing operational infrastructure would be used and traffic levels

would be expected to be less than during construction and operations. There are additional sensitive areas that should be considered within some of the regions, but because of decreasing noise levels with distance, construction activities would have only SMALL and temporary noise impacts for residences, communities, or sensitive areas, especially those located more than about 300 m [1,000 ft] from specific noise generating activities. Noise usually is not discernable to offsite receptors at distances more than 300 m [1,000 ft]. All the uranium districts are located more than 300 m [1,000 ft] from the closest community—SMALL to MODERATE.

DECOMMISSIONING—Noise generated during decommissioning would be noticeable only in proximity to equipment and temporary (typically daytime only). Administrative and engineering controls would be used to maintain noise levels in work areas below OSHA regulatory limits and mitigated by use of personal hearing protection. Noise levels during decommissioning would be less than during construction and would diminish as less and less equipment is used and truck traffic is reduced. Noise usually is not discernable to offsite receptors at distances more than 300 m [1,000 ft]. All the uranium districts are located more than 300 m [1,000 ft] from the closest community—SMALL.

Historical and Cultural Resources Impacts

CONSTRUCTION—Potential impacts during ISL facility construction could include loss of, or damage and temporary restrictions on access to, historical, cultural, and archaeological resources. The eligibility evaluation of cultural resources for listing in the National Register of Historic Places (NRHP) under criteria in 36 CFR 60.4(a)–(d), and/or as Traditional Cultural Properties (TCP) would be conducted as part of the site-specific review and NRC licensing procedures undertaken during the National Environmental Policy Act (NEPA) review process. The evaluation of impacts to any historic properties designated as TCPs and tribal consultations regarding cultural resources and TCPs also occurs during the site-specific licensing application and review process. To determine whether significant cultural resources would be avoided or mitigated, consultations with State Historic Preservation Offices (SHPO), other government agencies (e.g., U.S. Fish and Wildlife Service and State Environmental Departments), and Native American Tribes (THPO) occur as part of the site-specific review. Additionally, as needed, the NRC license applicant would be required, under conditions in its NRC license, to adhere to procedures regarding the discovery of previously undocumented cultural resources during initial construction. These procedures typically require the licensee to stop work and to notify the appropriate federal, tribal, and state agencies with regard to mitigation measures—SMALL or MODERATE to LARGE depending on site-specific conditions.

OPERATION—Because less land disturbance occurs during the operations phase, potential impacts to historical, cultural, and archaeological resources would be less than during construction. Conditions in the NRC license requiring adherence to procedures regarding the discovery of previously undocumented cultural resources would apply during operation. These procedures typically require the licensee to stop work and to notify the appropriate federal, tribal, and state agencies with regard to mitigation measures—SMALL, but depending on site-specific conditions.

AQUIFER RESTORATION—Because less land disturbance occurs during the aquifer restoration phase, potential impacts to historical, cultural, and archaeological resources would be less than during construction. Conditions in the NRC license requiring adherence to procedures regarding the discovery of previously undocumented cultural resources would apply during aquifer restoration. These procedures typically require the licensee to stop work and to

notify the appropriate federal, tribal, and state agencies with regard to mitigation measures—SMALL, but depending on site-specific conditions.

DECOMMISSIONING—Because less land disturbance occurs during the decommissioning phase and because decommissioning and reclamation activities would be focused on previously disturbed areas, potential impacts to historical, cultural, and archaeological resources would be less than during construction. Conditions in the NRC license requiring adherence to procedures regarding the discovery of previously undocumented cultural resources would apply during decommissioning and reclamation. These procedures typically require the licensee to stop work and to notify the appropriate federal, tribal, and state agencies with regard to mitigation measures—SMALL, depending on site-specific conditions.

Visual and Scenic Impacts

CONSTRUCTION—Visual impacts result from equipment (drill rig masts, cranes), dust/diesel emissions from construction equipment, and hillside and roadside cuts. Most of the four uranium milling regions are classified as Visual Resource Management (VRM) Class II through IV by the BLM. A number of VRM Class II areas surround national monuments (El Morro and El Malpais), the Chaco Culture National Historic Park, and sensitive areas managed within the Mt. Taylor district, in the Northwestern New Mexico Uranium Milling District, and would have the greatest potential for impacts to visual resources. Most of these areas, however, are located away from potential ISL facilities, at distances greater than 16 km [10 mi]. Most potential facilities are located in VRM Class III and IV areas. The general visual and scenic impacts associated with ISL facility construction would be temporary and SMALL, but from a Native American perspective, any construction activities would likely to result in adverse impacts to the landscape, particularly for facilities located in areas within view of tribal lands and areas of special significance such as Mt. Taylor. In addition, a PSD Class I area (Wind Cave National Park) is located in the Nebraska-South Dakota-Wyoming Uranium Milling Region. Nevertheless, most potential visual impacts during construction would be temporary as equipment is moved, and would be mitigated by best management practices (e.g., dust suppression). Because of the generally rolling topography of the region, most visual impacts during construction would not be visible from more than about 1 km [0.6 mi]. The visual impacts associated with ISL construction would be consistent with the predominant VRM Class III and IV—SMALL.

OPERATION—Visual impacts during operations would be less than those associated with construction. Most of the well field surface infrastructure has a low profile, and most piping and cables would be buried. The tallest structures include the central uranium processing facility {10 m [30 ft]} and power lines {6 m [20 ft]}. Because of the generally rolling topography of the regions, most visual impacts during operations would not be visible from more than about 1 km [0.6 mi]. Irregular layout of well field surface structures such as wellhead protection and header houses would further reduce visual contrast. Best management practices, design (e.g., painting buildings) and landscaping techniques would be used to mitigate potential visual impact. The uranium districts in the four regions are all located more than 16 km [10 mi] from the closest VRM Class II region, and the visual impacts associated with ISL construction would be consistent with the predominant VRM Class III and IV—SMALL.

AQUIFER RESTORATION—Aquifer restoration activities would use in-place infrastructure. As a result, potential visual impacts would be the same as, or less than, those during operations—SMALL.

DECOMMISSIONING SMALL—Because similar equipment would be used and activities conducted, potential visual impacts during decommissioning would be the same as, or less than, those during construction. Most potential visual impacts during decommissioning would be temporary as equipment is moved, and mitigated by best management practices (e.g., dust suppression). Visual impacts would be low, because these sites are in sparsely populated areas, and impacts would diminish as decommissioning activities decrease. An approved site reclamation plan is required prior to license termination, with the goal of returning the landscape to preconstruction condition (predominantly VRM Class III and IV). Some roadside cuts and hill slope modifications, however, may persist beyond decommissioning and reclamation—SMALL.

Socioeconomic Impacts

CONSTRUCTION—Potential impacts to socioeconomics would result predominantly from employment at an ISL facility and demands on the existing public and social services, tourism/recreation, housing, infrastructure (schools, utilities), and the local work force. Total peak employment would be about 200 people, including company employees and local contractors, depending on timing of construction with other stages of the ISL lifecycle. During construction of surface facilities and well fields, the general practice would be to use local contractors (drillers, construction), as available. A local multiplier of 0.7 (U.S. Bureau of the Census) is used to indicate how many ancillary jobs could be created (in this case about 140). For example, local building materials and building supplies would be used to the extent practical. Most employees would live in larger communities with access to more services. Some construction employees, however, would commute from outside the county to the ISL facility, and skilled employees (e.g., engineers, accountants, managers) would come from outside the local work force. Some of these employees would temporarily relocate to the project area and contribute to the local economy through purchasing goods and services and taxes. Because of the small relative size of the ISL workforce, net impacts would be SMALL to MODERATE.

OPERATION—Employment levels for ISL facility operations would be less than for construction, with total peak employment depending on timing and overlap with other stages of the ISL lifecycle. Use of local contract workers and local building materials would diminish, because drilling and facility construction would diminish. Revenues would be generated from federal, state, and local taxes on the facility and the uranium produced. Employment types would be similar to construction, but the socioeconomic impacts would be less due to fewer employees—SMALL to MODERATE.

AQUIFER RESTORATION—In-place infrastructure would be used for aquifer restoration, and employment levels would be similar to those for operations—SMALL to MODERATE.

DECOMMISSIONING—A skill set similar to the construction workforce would be involved in dismantling surface structures, removing pumps, plugging and abandoning wells, and reclaiming/re-contouring the ground surface. Employment levels and use of local contractor support during decommissioning would be similar to that required for construction. Employment would be temporary, however, as decommissioning activities are in duration. Because of similar employment levels, other socioeconomic impacts would be similar to construction—SMALL to MODERATE.

Public and Occupational Health and Safety Impacts

CONSTRUCTION—Worker safety would be addressed by standard construction safety practices. Fugitive dust would result from construction activities and vehicle traffic, but would likely be of short duration and would not result in a radiological dose. Diesel emissions would also be of short duration and readily dispersed into the atmosphere—SMALL to MODERATE.

OPERATION—Potential occupational radiological impacts from normal operations would result from: (1) exposure to radon gas from well field, (2) ion-exchange resin transfer operations, and (3) venting during processing activities. Workers would also be exposed to airborne uranium particulates from dryer operations and maintenance activities. Potential public exposures to radiation could occur from the same radon releases and uranium particulate releases (i.e., from facilities without vacuum dryer technology). Both worker and public radiological exposures are addressed in NRC regulations at 10 CFR Part 20, which require licensees to implement an NRC-approved radiation protection program. (Measured and calculated doses for workers and the public are commonly only a fraction of regulated limits.) Non-radiological worker safety matters are addressed through commonly-applied occupational health and safety regulations and practices. Radiological accident risks could involve processing equipment failures leading to yellowcake slurry spills, or radon gas or uranium particulate releases. Consequences of accidents to workers and the public are generally low, with the exception of a dryer explosion which could result in worker dose above NRC limits. The likelihood of such an accident would be low, and therefore the risk would also be low. Potential non-radiological accidents impacts include high consequence chemical release events (e.g., ammonia) for both workers and nearby populations. The likelihood, however, of such release events would be low based on historical operating experience at NRC-licensed facilities, primarily due to operators following commonly-applied chemical safety and handling protocols—SMALL to MODERATE.

AQUIFER RESTORATION—Activities involving aquifer restoration overlap with similar operational activities (e.g., operation of well fields, waste water treatment and disposal). The resultant types of impacts on public and occupational health and safety are similar to operational impacts. The absence of some operational activities (e.g., yellowcake production and drying, remote ion exchange) further limits the relative magnitude of potential worker and public health and safety hazards—SMALL.

DECOMMISSIONING—Worker and public health and safety would be addressed in a NRC-required decommissioning plan. This plan details how a 10 CFR Part 20 compliant radiation safety program would be implemented during decommissioning, ensuring the safety of workers and the public would be maintained and applicable safety regulations complied with—SMALL.

Waste Management Impacts

CONSTRUCTION—Relatively small scale construction activities (Section 2.3) and incremental well field development at ISL facilities would generate low volumes of construction waste—SMALL.

OPERATION—Operational wastes primarily result from liquid waste streams including process bleed, flushing of depleted eluant to limit impurities, resin transfer wash, filter washing, uranium precipitation process wastes (brine), and plant wash down water. State permit actions, NRC license conditions, and NRC inspections ensure the proper practices would be used to comply with safety requirements to protect workers and the public. Waste treatments such as reverse osmosis and radon settling would be used to segregate wastes and minimize disposal volumes. Potential impacts from surface discharge and deep well injection would be limited by the conditions specified in the applicable state permit. NRC regulations address constructing, operating, and monitoring for leakage of evaporation ponds used to store and reduce volumes of liquid wastes. Potential impacts from land application of treated wastewater would be addressed by NRC review of site-specific conditions prior to approval and routine monitoring in decommissioning surveys. Offsite waste disposal impacts would be SMALL for radioactive wastes as a result of required preoperational disposal agreements. Impacts for hazardous and

municipal waste would also be SMALL due to the volume of wastes generated. For remote areas with limited available disposal capacity, such wastes may need to be shipped greater distances to facilities that have capacity; however, the volume of wastes generated and magnitude of such shipments are estimated to be low—SMALL.

AQUIFER RESTORATION—Waste management activities during aquifer restoration would use the same treatment and disposal options implemented for operations. Therefore, impacts associated with aquifer restoration would be similar to operational impacts. While the amount of wastewater generated during aquifer restoration would be dependent on site-specific conditions, the potential exists for additional wastewater volume and associated treatment wastes during the restoration period. However, this would be offset to some degree by the reduction in production capacity from the removal of a well field. NRC review of future ISL facility applications would verify that sufficient water treatment and disposal capacity (and the associated agreement for disposal of byproduct material) are addressed. As a result, waste management impacts from aquifer restoration would be—SMALL.

DECOMMISSIONING—Radioactive wastes from decommissioning ISL facilities (including contaminated excavated soil, evaporation pond bottoms, process equipment) would be disposed of as byproduct material at an NRC-licensed facility. A preoperational agreement with a licensed disposal facility to accept radioactive wastes ensures sufficient disposal capacity would be available for byproduct wastes generated by decommissioning activities. Safe handling, storage, and disposal of decommissioning wastes would be addressed in a required decommissioning plan for NRC review prior to starting decommissioning activities. Such a plan would detail how a 10 CFR Part 20 compliant radiation safety program would be implemented during decommissioning to ensure how the safety of workers and the public would be maintained and applicable safety regulations complied with. Overall, volumes of decommissioning radioactive, chemical, and solid wastes would be—SMALL.

ABBREVIATIONS/ACRONYMS

BLM	U.S. Bureau of Land Management
CBSA	Core-Based Statistical Area
CEA	Cumulative Effects Assessment
CERCLA	Comprehensive Environmental Response, Compensation, and Liability Act of 1980
CEQ	Council on Environmental Quality
Dod	Department of Defense
EIS	Environmental Impact Statement
EPA	U.S. Environmental Protection Agency
FONSI	Finding of No Significant Impact
GEIS	Generic Environmental Impact Statement
ISL	<i>In-situ</i> Leaching
MIT	Mechanical Integrity Testing
NAAQS	National Ambient Air Quality Standards
NAGPRA	Native American Graves Protection and Repatriation Act
NDEQ	Nebraska Department of Environmental Quality
NEPA	National Environmental Policy Act
NHPA	National Historic Preservation Act
NPDES	National Pollutant Discharge Elimination System
NRC	U.S. Nuclear Regulatory Commission
NRCS	Natural Resources Conservation Service
NRHP	National Register of Historic Places
PVC	Polyvinyl Chloride
RFFA	Reasonably Foreseeable Future Action
SHPO	State Historic Preservation Officer
TDS	Total Dissolved Solids
THPO	Tribal Historic Preservation Officer
UCL	Upper Control Limit
UIC	Underground Injection Control
UMTRCA	Uranium Mill Tailings Radiation Control Act
USACE	U.S. Army Corps of Engineers
USDA	U.S. Department of Agriculture
USFS	U.S. Forest Service
VRM	Visual Resource Management
WDEQ	Wyoming Department of Environmental Quality

SI* (MODERN METRIC) CONVERSION FACTORS

Approximate Conversions From SI Units				
Symbol	When You Know	Multiply By	To Find	Symbol
Length				
mm	millimeters	0.039	inches	in
m	meters	3.28	feet	ft
m	meters	1.09	yards	yd
km	kilometers	0.621	miles	mi
Area				
mm ²	square millimeters	0.0016	square inches	in ²
m ²	square meters	10.764	square feet	ft ²
m ²	square meters	1.195	square yards	yd ²
ha	hectares	2.47	acres	ac
km ²	square kilometers	0.386	square miles	mi ²
Volume				
mL	milliliters	0.034	fluid ounces	fl oz
L	liters	0.264	gallons	gal
m ³	cubic meters	35.314	cubic feet	ft ³
m ³	cubic meters	1.307	cubic yards	yd ³
m ³	cubic meters	0.0008107	acre-feet	acre-feet
Mass				
g	grams	0.035	ounces	oz
kg	kilograms	2.202	pounds	lb
Mg (or "t")	megagrams (or "metric ton")	1.103	short tons (2000 lb)	T
Temperature (Exact Degrees)				
°C	Celsius	1.8C + 32	Fahrenheit	°F
*SI is the symbol for the International System of Units. Appropriate rounding should be performed to comply with Section 4 of ASTM E380 (ASTM International. "Standard for Metric Practice Guide." West Conshohocken, Pennsylvania: ASTM International. Revised 2003.).				

1 INTRODUCTION

The U.S. Nuclear Regulatory Commission (NRC) prepared this Draft Generic Environmental Impact Statement (GEIS) to identify and evaluate potential environmental impacts associated with the construction, operation, aquifer restoration, and decommissioning of *in-situ* leach (ISL) uranium recovery facilities on a programmatic basis for specific identified regions in the western United States. Based on discussions between uranium mining companies and the NRC staff, ISL facilities could be located in portions of Wyoming, Nebraska, South Dakota, and New Mexico. NRC is the regulatory authority that licenses ISL facilities in these States.

1.1 Purpose of the GEIS

The purpose behind developing the GEIS is to improve the efficiency of NRC's environmental reviews for ISL license applications required under the National Environmental Policy Act of 1969, as amended (NEPA). NRC regulations that implement NEPA and discuss environmental reviews are found in Title 10, "Energy," of the Code of Federal Regulations (10 CFR) Part 51. The NRC staff plans to use the GEIS as a starting point for its NEPA analyses for site-specific license applications for new ISL facilities. Additionally, the NRC staff plans to use the GEIS, along with applicable previous site-specific environmental review documents, in its NEPA analysis for the restart or expansions of existing facilities.

NRC developed this Draft GEIS using (1) knowledge gained during the past 30 years licensing and regulating these facilities, (2) the active participation of the State of Wyoming as a cooperating agency, and (3) public comments received during the scoping period for the GEIS. NRC's research indicates that the technology used for ISL uranium recovery is relatively standardized throughout the industry and therefore appropriate for a programmatic evaluation in a GEIS.

NRC has identified four regions (Figure 1.1-1) to use as a framework for discussions in this Draft GEIS based on several considerations, including:

- Past and existing uranium milling sites are located within States where NRC has regulatory authority over uranium recovery (see text box)
- Potential new sites are identified

The NRC Agreement State Program

In accordance with Section 274 of the Atomic Energy Act of 1954, as amended, NRC may relinquish certain portions of its regulatory authority to those States that express interest in establishing their own programs for regulating the use of certain nuclear materials and demonstrated the adequacy and compatibility of their programs. The areas of regulatory authority that NRC may relinquish include the regulation of byproduct materials as defined in section 11e.(1), (3), and (4); source materials (uranium and thorium), certain quantities of special nuclear materials, byproduct material as defined in section 11e.(2) and the facilities that generate this material (uranium milling), the commercial disposal of low-level waste, and the evaluation of sealed sources and devices. A signed agreement between the Chairman of NRC and the Governor of the State identifies and documents the specific authorities transferred to the State. NRC reviews the performance of each Agreement State on a periodic basis as part of its Integrated Materials Performance Evaluation Program (NRC, 2004). Agreement State reviews are coordinated with the individual State and typically happen once every 4 years (NRC, 2004). Starting with Kentucky in 1962, more than 30 States have entered into the NRC Agreement State program.

Wyoming and South Dakota are Non-Agreement States, and NRC has authority for regulating nuclear materials in these States, including ISL facilities. New Mexico and Nebraska are Agreement States; however, their Agreements do not include the authority for 11e.(2) byproduct material (uranium milling). Therefore, NRC maintains regulatory authority with respect to uranium recovery facilities (uranium milling) in these states. (NRC, 2007a). Utah, Colorado, and Texas are full Agreement States and have regulatory authority over ISL facilities within their boundaries.

based on NRC's understanding of where the uranium recovery industry has plans to develop uranium deposits using ISL technology (NRC, 2008a)

- Locations of historical uranium deposits within portions of Wyoming, Nebraska, South Dakota, and New Mexico (EPA, 2006, 2007a) (Figure 1.1-2).





Figure 1.1-1. Four Geographic Regions Used as a Framework for the Analyses Presented in This GEIS

In this Draft GEIS, NRC documents the potential environmental impacts that would be associated with the construction, operation, aquifer restoration, and decommissioning of an ISL facility in specified regions of the western U.S. and evaluates the significance of those impacts on a programmatic basis. In its review of individual ISL license applications, NRC would evaluate the site-specific data to determine whether relevant sections of the GEIS could be incorporated by reference into the site-specific environmental review. Additionally, NRC would

determine whether aspects of the site and/or the applicant's proposed activities are consistent with those evaluated in the GEIS or are such that additional analysis in specific topic areas would be required. Section 1.8 of the Draft GEIS provides a more detailed discussion of the use of the GEIS in the site-specific licensing review process



Legend

○ EPA-Identified Uranium Locations

Miles
0 75 150 300 450



Figure 1.1-2. Major Uranium Reserves Within the United States. (From Energy Information Administration, 2004).

1.2 The Proposed Federal Action

In States where NRC is the regulatory authority over the licensing of uranium milling (including the ISL process), NRC has a statutory obligation to assess each site-specific license application

to ensure it complies with NRC regulations before issuing a license. The proposed federal action is to prepare a GEIS that identifies and evaluates the potential environmental impacts associated with the construction, operation, aquifer restoration, and decommissioning of ISL milling facilities in portions of Wyoming, Nebraska, South Dakota, and New Mexico. NRC intends to make use of the GEIS during subsequent site-specific ISL licensing actions.

1.3 Purpose and Need for the Action

NRC is the regulatory authority responsible for licensing ISL facilities in Wyoming, Nebraska, South Dakota, and New Mexico. Commercial uranium recovery companies have approached NRC with their plans to submit as many as 21 license applications for new uranium recovery sites, as well as for potentially 10 applications for the restart or expansion of existing facilities in the next several years (NRC, 2008a). The companies have indicated that these new, restarted, and expanded facilities would be located in these States. Given that the large majority of these potential applications (perhaps 24 of the 31) would involve use of the ISL process and that such applications may be submitted over a relatively short period of time, NRC decided to prepare a GEIS to increase the efficiency of and support a consistent approach to NRC's site-specific environmental review of license applications for ISL facilities (NRC, 2007b).

This Draft GEIS, however, does not address the purpose and need of the primary Federal action of issuing licenses for ISL facilities. As discussed in Section 1.8, NRC plans to conduct a site-specific environmental analysis in support of its review of a license application for an ISL facility. Relevant sections of the GEIS can be incorporated by reference into the site-specific environmental review in a process known as tiering. It is not appropriate for NRC to determine in the Draft GEIS the purpose and need for individual ISL applications. The purpose and need for each ISL application will be addressed in the site-specific environmental review that NRC will conduct.

1.4 Structure of the GEIS

In this Draft GEIS, NRC systematically evaluated the potential environmental impacts of constructing, operating, restoring aquifers, and decommissioning an ISL uranium recovery facility in four separate geographic regions of the western United States. The regions represent areas in four western states: Wyoming, Nebraska, South Dakota, and New Mexico. As stated in Section 1.1, three criteria were used to identify these regions for the purpose of the Draft GEIS analysis. These regions are:

- **The Wyoming West Uranium Milling Region.** This region includes portions of four Wyoming counties (Carbon, Fremont, Natrona, and Sweetwater).
- **The Wyoming East Uranium Milling Region,** which includes portions of eight Wyoming counties (Albany, Campbell, Carbon, Converse, Johnson, Natrona, Platte, and Weston) east of the Bighorn Mountains.
- **The Nebraska-South Dakota-Wyoming Uranium Milling Region.** This region includes the portions of northwestern Nebraska (Dawes and Sioux Counties), western South Dakota (Custer, Fall River, Lawrence, and Pennington Counties), and the extreme eastern portion of Wyoming (Crook, Niobrara, and Weston Counties).
- **The Northwestern New Mexico Uranium Milling Region,** which includes McKinley County and portions of Cibola and Sandoval Counties.

1.4.1 Describing the ISL Process

Chapter 2 of this Draft GEIS describes the ISL process, addressing construction, operation, aquifer restoration, and decommissioning of an ISL facility. This description is based on historical operations information from ISL facilities NRC licenses and regulates. The construction stage includes well field development and the construction of surface facilities and supporting infrastructure. Operations includes injection and production of solutions from uranium mineralization in the subsurface, as well as the process to recover the uranium from these solutions. Aquifer restoration includes activities to restore the groundwater quality in the production zone after uranium recovery is completed within a well field. Decommissioning includes the final stages of removing surface and subsurface infrastructure and reclaiming the surface after uranium production activities at a site has been completed. Chapter 2 of the Draft GEIS also includes a section on financial surety arrangements, where the licensee or applicant establishes a bond or other financial mechanism prior to operations to ensure that sufficient funds are available to complete aquifer restoration, decommissioning, and reclamation activities.

Site-specific license applications may not include all stages of the ISL process. For example, an applicant may propose to limit activities to well field construction, uranium mobilization and ion exchange, and then ship the uranium-bearing resin to an existing processing plant for final processing. In this case, the applicant's license application would likely exclude the construction, operation, and decommissioning of a processing plant. NRC categorizes the ISL operations by various stages so that relevant portions of the GEIS can be incorporated by reference into the subsequent site-specific environmental reviews.

1.4.2 Describing the Affected Environment

Chapter 3 of the Draft GEIS describes the affected environment for each of the four geographic regions using the environmental resource areas identified in (NRC, 2003b), which provides guidance to the NRC staff in conducting environmental reviews. These resource areas are

- | | |
|---------------------|------------------------------------|
| • Land use | • Noise |
| • Transportation | • Historical and cultural resource |
| • Geology and soils | • Visual and scenic resources |
| • Water resources | • Socioeconomic |
| • Ecology | • Public and occupational health |
| • Air Quality | |

NRC staff will conduct independent, site-specific environmental reviews for each license application (see Section 1.8.3). Chapter 3 of this Draft GEIS is divided into regional area discussions to facilitate using the Draft GEIS in these site-specific reviews. Relevant sections of the regional discussions can be incorporated by reference in the site-specific environmental reviews.

1.4.3 Identifying Environmental Issues and Characterizing Significance

In Chapter 4, NRC evaluates the potential environmental impacts of construction, operation, aquifer restoration, and decommissioning of an ISL facility in each of the four regions. In essence, this involves placing an ISL facility with the characteristics described in Chapter 2 of the Draft GEIS within each of the four regional areas described in Chapter 3 and then describing and evaluating the significance of potential impacts in each region separately. The description

for each identified potential environmental impact includes the type and magnitude of the ISL activity that would affect the environment and the attributes of the resource area that would be potentially affected.

The assessment of impacts considers potential environmental consequences at each stage in an ISL facility's lifetime—construction, operation, aquifer restoration, and decommissioning/reclamation—and presents them for each of the resource areas identified in Chapter 3.

According to the Council on Environmental Quality (CEQ), the significance of impacts is determined by examining both context and intensity (40 CFR 1508.27). Context is related to the affected region, the affected interests, and the locality, while intensity refers to the severity of the impact, which is based on a number of considerations. In describing the significance of potential impacts in this Draft GEIS, the NRC used the significance levels identified in NUREG-1748 (NRC, 2003b) (see text box).

Considerations related to potential cumulative impacts are described in Chapter 5, and environmental justice is discussed in Chapter 6. Mitigation measures and best management practices that may reduce potential environmental impacts are identified and discussed in Chapter 7. Required monitoring programs are described in Chapter 8 and are included in the determination of significance. Chapter 9 discusses the process for NRC's consultation with federal, tribal, state, and local agencies. In Chapter 10, impacts are summarized in a table for each of the four geographic regions. The structure of this Draft GEIS is shown graphically in Figure 1.4-1.

1.5 Scope of the GEIS

The scoping process occurs early in the development of an EIS in accordance with NEPA. Scoping provides an opportunity for the public and other stakeholders to identify key issues and concerns that they believe should be addressed in the document. The NRC requirements for scoping are found at 10 CFR 51.26-29, while the general NRC approach to scoping is described in NUREG-1748 (NRC, 2003b, Section 4.2.3).

1.5.1 The GEIS Scoping Process

On July 24, 2007, NRC published in the *Federal Register* a notice of intent to prepare a GEIS to examine the potential impacts associated with ISL uranium recovery facilities (NRC, 2007b). In that notice, NRC described the scoping process for the GEIS and established a public comment period from July 24, 2007, to September 4, 2007. NRC also announced dates and times for two public scoping meetings to be held—one in Albuquerque, New Mexico, and the other in Casper, Wyoming. NRC published a revised notice of intent in the *Federal Register* on August 31, 2007, announcing a third public scoping meeting in Gallup, New Mexico, and extended the public comment period to October 8, 2007 (NRC, 2007c). Following the Gallup public meeting, NRC subsequently extended the comment period further to October 31, 2007, and finally to

Classifying Impact Significance (after NRC, 2003b)

- *Small Impact:* The environmental effects are not detectable or are so minor that they will neither destabilize nor noticeably alter any important attribute of the resource considered.
- *Moderate Impact:* The environmental effects are sufficient to alter noticeably, but not destabilize, important attributes of the resource considered.
- *Large Impact:* The environmental effects are clearly noticeable and are sufficient to destabilize important attributes of the resource considered.




Figure 1.4-1. Structure of This GEIS

November 30, 2007 (NRC, 2007c). At each of the three public scoping meetings, NRC described its role and mission and reviewed NRC procedures and responsibilities. Then tribal, state, and local government agencies; concerned local citizens; and other stakeholders were invited to identify scoping issues and concerns and ask questions. Transcripts (NRC, 2008b, 2007d,e) were prepared for all three meetings and are available online at the NRC Agencywide Documents Access and Management System (ADAMS), which is accessible at www.nrc.gov or through the NRC website for the GEIS at <http://www.nrc.gov/materials/fuel-cycle-fac/licensing/geis.html>.

In addition to the comments received at the public meetings, NRC also accepted written comments submitted either by regular mail or electronically. Using these varied methods, comments were received from approximately 1,600 entities (i.e., federal, state, and local agencies; industry organizations; public advocacy groups; and individual members of the public).

A summary of all comments NRC received during scoping is provided in a scoping summary report included as Appendix A to this Draft GEIS.

1.5.2 Issues To Be Studied in Detail

From the scoping process, NRC determined that the following issues identified by the public and other stakeholders will be addressed in the GEIS.

- **Proposed Action and Alternatives.** Scoping comments recommended clarifying the scope of the proposed action. Commenters also suggested a variety of alternatives for consideration. The proposed action is described in Section 1.2 and alternatives are described in Sections 2.12 and 2.13.
- **Applicable Statutes, Regulations, and Agencies.** Scoping comments expressed a need to clarify applicable regulations and the roles of government agencies in regulating ISL facilities. Various statutes, regulations, and implementing agencies at the federal, state, and local levels that have a role in regulating ISL facilities are identified and discussed in Section 1.6. The roles of these agencies are also described, as appropriate.
- **Purpose of the Draft GEIS and Use in Site-Specific Licensing Reviews.** A number of scoping comments conveyed various interpretations of the purpose and intended use of the GEIS, suggesting the purpose and intended use needed to be clarified. For example, some thought the GEIS was going to be the only NEPA analysis conducted for all ISL facilities while others thought the GEIS would eliminate or substantially degrade the rigor of NRC site-specific environmental reviews. A statement of purpose is included in Section 1.3, the NRC licensing process is described in Section 1.7.1, and the ways NRC intends to use the GEIS to evaluate environmental impacts in site-specific licensing reviews is provided in Section 1.8.
- **Opportunities for Public Involvement.** Many scoping comments reflected a perception that the GEIS would limit public involvement in ISL licensing. Some requested the opportunities for public involvement be described. Section 1.8.4 describes opportunities for public participation in the ISL licensing process.

- **Applicable Rulemaking Activities.** Some scoping comments recommended a discussion of ongoing rulemaking activities that are applicable to ISL licensing or the GEIS. The Draft GEIS is based on the existing regulations in effect at the time of writing.
- **Land Use.** Concerns regarding potential land use impacts on ranching operations and livestock were raised during the scoping process. Potential impacts to existing land uses in the ISL milling regions including potential impacts to ranching, grazing, recreation, industrial, and cultural activities are discussed in Sections 4.2.1, 4.3.1, 4.4.1, and 4.5.1.
- **Transportation.** Scoping comments addressed general concerns with the safety of shipping yellowcake, road construction, fugitive dust generation, infrastructure damage, and incidental livestock kills. Potential radiological and nonradiological impacts from ISL transportation activities are discussed in Sections 4.2.2, 4.3.2, 4.4.2, and 4.5.2. Impacts regarding shipment of supplies, yellowcake product, and wastes associated with each phase of the ISL facility lifecycle are discussed. Normal transportation and accident conditions are considered. Potential nonradiological impacts evaluated include dust and noise generation, impacts on infrastructure such as roads, incidental livestock and wildlife kills, and changes to local traffic conditions. Potential radiological impacts considered include direct radiation and potential release of radioactive material from accidents during shipment.
- **Geology.** Scoping comments were received regarding the extent of soil disturbance and questioning the usefulness of a generic analysis of geology. The Draft GEIS describes the geology of the ISL milling regions in sufficient detail to support the evaluation of impacts to geology and soils (Sections 4.2.3, 4.3.3, 4.4.3, and 4.5.3) and groundwater (Sections 4.2.4.2, 4.3.4.2, 4.4.4.2, and 4.5.4.2) from ISL activities. Chapter 2 of the Draft GEIS describes soil-disturbing activities (e.g., clearing, excavation, drilling, trenching, road construction, leaks, spills) and the magnitude of surface area disturbed at existing ISL facilities.
- **Water Resources.** A variety of water resource issues were raised in scoping comments including concerns about potential groundwater and surface water contamination, water availability and consumptive use, groundwater protection requirements, and aquifer restoration goals and techniques. The Draft GEIS addresses potential impacts to surface waters, groundwater, and wetlands from each phase of the ISL facility lifecycle in Sections 4.2.4, 4.3.4, 4.4.4, and 4.5.4. Specific topics addressed include permitted surface water discharges, leaks and spills, groundwater excursions, consumptive water use, aquifer restoration, deep well injection, and applicable regulations. Hydrologic conditions in uranium milling regions are considered, as well as available restoration technologies and methods. The restoration of the aquifer water quality in the production zone following operations is addressed in the Draft GEIS. Data from aquifer restoration efforts at ISL sites informs the analysis. Regulatory requirements and the roles of various federal, state, and local agencies regarding aquifer restoration are also discussed. Potential for groundwater impacts, in particular, is a key concern that has been historically an area of focus in NRC ISL licensing reviews.
- **Ecology.** Scoping comments on ecology raised topics regarding surface disturbance impacts on wildlife and vegetation, practices for isolating wildlife from exposure to

uranium and other metals, recommended construction guidelines, habitat loss and fragmentation, and avoiding establishment of invasive species. The Draft GEIS assesses the potential impacts to ecology in the uranium milling regions from all phases of the ISL facility lifecycle in Sections 4.2.5, 4.3.5, 4.4.5, and 4.5.5. This includes consideration of potential impacts to terrestrial, aquatic, and threatened and endangered species. Specific topics addressed include evaluating ecoregions and habitat for a variety of listed species and assessing potential impacts from surface disturbances, habitat loss and fragmentation, and incidental kills. Applicable regulations and various management practices designed to protect species or mitigate potential impacts are discussed.

- **Meteorology, Climatology, and Air Quality.** Scoping comments included general environmental and safety concerns about the potential for airborne contamination, the magnitude of airborne facility releases, and applicable regulations. Sections 4.2.6, 4.3.6, 4.4.6, and 4.5.6 of the Draft GEIS consider the potential impacts of all phases of the ISL facility lifecycle on local and regional air quality from both radiological and nonradiological emissions. The radiological air emissions addressed in the Draft GEIS include radon from well fields, processing, and waste treatment operations and the potential for uranium particulate emissions from yellowcake drying operations. Nonradiological emissions addressed in the Draft GEIS include combustion engine exhausts from trucking and well drilling operations and fugitive dusts from a variety of activities.

- **Noise.** Scoping comments on noise were limited to a statement regarding the low levels of noise ISL facilities generate. NRC recognizes that some activities in the ISL facility lifecycle can potentially generate additional noise, and impacts are evaluated in the Draft GEIS Sections 4.2.7, 4.3.7, 4.4.7, and 4.5.7. This includes noise from well field development, uranium processing activities, and trucking activities associated with all phases of the ISL facility lifecycle.

- **Historic and Cultural.** Scoping comments were provided on historic and cultural resources including recommendations for documenting compliance with the National Historic Preservation Act regarding protecting historic properties on tribal lands, concerns about the notification process when cultural artifacts are found at an ISL facility, and opportunities for public participation regarding historic and cultural concerns. A number of individuals and organizations, primarily in New Mexico, expressed concerns on topics regarding proximity of uranium facilities to Native American communities and requested government-to-government consultations and documentation of consultations in the GEIS. The Draft GEIS assesses potential impacts from all phases of the ISL facility lifecycle on historical and cultural resources in Sections 4.2.8, 4.3.8, 4.4.8, and 4.5.8. Local and regional historic and cultural properties and practices in ISL milling regions such as those involving Native American communities and governments are included. A description of NRC's process for consultation with Native American governments is provided in Chapter 9 of the Draft GEIS.

- **Visual Resources.** Scoping comments on visual resource impacts were varied. Potential impacts to visual resources in uranium milling regions from all phases of the ISL facility lifecycle are assessed in Draft GEIS Sections 4.2.9, 4.3.9, 4.4.9, and 3.5.9.

Assessments consider scenic vistas and sensitive viewsheds within uranium milling regions and ISL facility lifecycle impacts on these resources based on proximity.

- **Socioeconomics.** Scoping comments recommended evaluating social and economic impacts to local communities including job creation impacts; changes to tax base; and cumulative impacts on housing, roads, services, and labor to towns already overburdened by oil, gas, and coal development. The Draft GEIS assesses potential impacts to socioeconomic conditions in uranium milling regions from all phases of the ISL facility lifecycle in Sections 4.2.10, 4.3.10, 4.4.10, 4.5.10. Local and regional characteristics pertaining to demographics, income, tax structure and distribution, housing, employment, finances, education, and services are considered.
- **Public and Occupational Health.** A number of scoping comments expressed general public and worker safety concerns and more specific concerns about potential contamination of soils, surface water, air, and groundwater; risks from radon gas and spills and from processing chemicals and resins; and emergency response and reporting. Potential impacts to public and occupational health from all phases of the ISL facility lifecycle are assessed in Draft GEIS Sections 4.2.11, 4.3.11, 4.4.11, and 4.5.11. Both nonradiological (including chemical) and radiological effluents and releases under normal (routine) and accident conditions are assessed. Dose calculation results from previously licensed ISL facilities that include airborne uranium particulate and radon gas are provided. Hazards and risks for ISL processing chemicals are also considered. Potential soil contamination impacts from leaks and spills are discussed in Sections 4.2.3, 4.3.3, 4.4.3, and 4.5.3, and potential groundwater contamination is in 4.2.4, 4.3.4, 4.4.4, and 4.5.4.
- **Waste Management.** Scoping comments expressed concerns about waste management in general and also about handling and disposal practices, deep well injection and permitted discharges, land application, disposal capacity, annual waste volumes, transportation, and applicable regulations. The Draft GEIS considers impacts from waste management activities in all phases of the ISL facility lifecycle in Sections 4.2.12, 4.3.12, 4.4.12, and 4.5.12. Generation, handling, treatment, transportation, and final disposal of chemical, radiological, and municipal wastes are addressed. Constituents in various waste streams are identified and volume estimates are provided.
- **Decontamination, Decommissioning, Reclamation.** A number of scoping comments expressed concerns about the site cleanup after operations end. The Draft GEIS assesses impacts to the environment from terminating ISL operations, which includes removal of facilities and equipment, disposal of waste materials, cleanup of contaminated areas, and reclamation of lands to pre-milling conditions. Decommissioning impacts are assessed for each resource area discussed in Chapter 4. Waste volume estimates by type of waste are provided and applicable requirements are discussed.
- **Accidents.** Scoping comments requested consideration of credible accident scenarios. Potential accident conditions are assessed in various sections in the Draft GEIS. This includes considering a range of possible accidents and off-normal operating conditions and estimating and evaluating consequences including well field leaks and spills,

excursions, processing chemical spills, and ion exchange resin and yellowcake transportation accidents.

- **Environmental Justice.** A range of opinions was provided in scoping comments on environmental justice in the GEIS. Some commenters thought it should be included in the GEIS and others thought it should not be included. Still others provided various suggestions on how to do the analysis. The Draft GEIS (Chapter 6) discusses the potential for disproportionately high and adverse environmental and health impacts on minority and low income populations from future ISL licensing in the specified uranium milling regions.
- **Cumulative Impacts.** Scoping comments on cumulative impacts offered a number of suggestions for reasonably foreseeable future actions to be included in the GEIS, including coal bed methane operations, and oil and gas development. The Draft GEIS (Chapter 5) describes past, present, and reasonably foreseeable future actions in the uranium milling regions and evaluates which resource areas would be potentially impacted by both ISL facilities and the types of reasonably foreseeable future actions identified in the regions. Due to the complex and site-specific nature of a cumulative impact assessment, the Draft GEIS provides useful information for understanding the potential for cumulative impacts when licensing future ISL facilities in the milling regions, but does not make conclusions regarding cumulative impacts for specific sites.
- **Monitoring.** Scoping comments on monitoring recommended the GEIS discuss monitoring programs designed to assess impacts from operations and waste management practices. The Draft GEIS discusses various monitoring techniques and programs (Chapter 2, Chapter 8) used to detect radiological and nonradiological contaminants within and beyond ISL facility boundaries. This includes effluent monitoring, workplace radiological monitoring, groundwater monitoring to detect potential excursions, and environmental monitoring at the facility boundary.
- **Financial Assurance.** Scoping comments recommended the GEIS discuss bonding for complete restoration of groundwater and land. Requirements and practices designed to ensure companies engaged in ISL recovery have sufficient funds to close down operations, restore aquifers, decontaminate and decommission facilities, and reclaim lands are described in Draft GEIS Section 2.10.

1.5.3 Issues Eliminated From Detailed Study

The analyses presented in this Draft GEIS focus on potential impacts within the four geographic regions described in Section 1.1 and illustrated in Figure 1.1-1; they are not intended to provide a detailed assessment of any specific site. Yellowcake transportation from uranium mills to the uranium hexafluoride (UF₆) conversion facility in Metropolis, Illinois, is anticipated to be by truck over existing highways. Access roads may need to be constructed to bring the yellowcake from the mill to the state and national (interstate) highway system. The existing national transportation routes are not expected to be altered. Because the environmental impacts of national transportation of yellowcake uranium have been previously analyzed, they will not be studied in detail within this Draft GEIS (NRC, 1977, 1980).

1.5.4 Issues Outside of the Scope of the GEIS

NRC has determined that comments received on topics in the following areas are outside the scope of this Draft GEIS:

- NRC's licensing process and the decision to prepare the Draft GEIS.
- General support or opposition for GEIS or uranium milling.
- Requests for cooperation or agreements.
- Matters that are regulated by Agreement States.
- Impacts associated with conventional uranium milling past or present.
- Requests for compensation for past mining impacts.
- Resolution of dual regulation issues.
- Consideration of human-induced climate change.
- Analysis of all variations of ISL technology.
- Alternative sources of uranium.
- Cumulative Impact Analysis.
- Energy debate.
- NRC credibility.

A discussion of why NRC determined that comments in these topic areas were outside the scope of the GEIS is provided in the Scoping Summary Report (Appendix A of the Draft GEIS).

1.6 Agencies Involved in Uranium ISL Facility Licensing

Different federal, tribal, state, and local agencies potentially have a role in licensing and permitting a uranium ISL facility. Specific statutes and regulations that may be applicable for uranium ISL facilities are detailed in Appendix B.

1.6.1 Federal Agencies

1.6.1.1 NRC

NRC responsibilities include regulating the nuclear industry in a manner that

- Protects public health and safety;
- Protects the environment; and

- Protects and safeguards materials and nuclear facilities in the interest of national security.

NRC is the federal agency with lead responsibility in licensing and regulating uranium ISL facilities through the statutory requirements of the Uranium Mill Tailings Radiation Control Act (UMTRCA) of 1978 and the Atomic Energy Act of 1954, as amended. In part, these statutes require that NRC ensure byproduct material, as defined in Section 11e.(2) of the Atomic Energy Act, is managed to conform with applicable general standards the U.S. Environmental Protection Agency (EPA) promulgated under Section 275 of the Atomic Energy Act. EPA standards of general application for 11e.(2) byproduct material were established in 40 CFR Part 192. The UMTRCA and the Atomic Energy Act also require that the generally applicable standards EPA promulgates for nonradiological hazards under UMTRCA be consistent with the standards EPA promulgates under the Safe Drinking Water Act/Resources Conservation and Recovery Act for such hazards. NRC conforming regulations are in 10 CFR Part 40, Appendix A.

NRC is the regulatory authority for ISL facilities unless NRC relinquishes its authority to a State in a written agreement. The text box on page 1-1 provides additional information on NRC's Agreement State program.

1.6.1.2 EPA

EPA also has a role in permitting nonradiological emissions and effluents. Water quality issues are administered predominantly through underground injection control (UIC) programs and National Pollutant Discharge Elimination System (NPDES) permits. Air quality issues are addressed through National Ambient Air Quality Standards (NAAQS) and National Emission Standards for Hazardous Air Pollutants (NESHAPS) programs. These programs may be administered directly by EPA, by States and Tribes granted primacy, or by joint programs between EPA and the state (EPA, 2008a-f).

1.6.1.3 Occupational Safety and Health Administration

The mission of the Occupational Health and Safety Administration (OSHA) is to assure the safety and health of workers in the United States, and it is the lead federal agency with responsibility for regulating the industrial safety of the work force at uranium ISL facilities. Recognizing the different agency responsibilities, NRC and OSHA have entered into memorandum of understanding to coordinate their inspection programs and avoid duplication of effort (Occupational Safety and Health Administration, 1988). As part of this program, NRC inspectors do not perform the role of OSHA, but they may identify safety concerns or receive complaints from employees about working conditions within the areas of responsibility for OSHA, notifying the OSHA Regional Office as appropriate (Occupational Safety and Health Administration, 1988).

1.6.1.4 U.S. Department of Transportation

The U.S. Department of Transportation regulates the shipments of radiological and nonradiological hazardous materials and sets regulatory requirements for type and condition of hazardous material containers, the mechanical condition of the transportation vehicles, the

training of personnel, and the routing requirements, package labels, vehicle placards, and shipping papers associated with shipments of radioactive materials. The U.S. Department of Transportation also inspects containers, storage facilities, and carrier equipment (Office of Technology Assessment, 1986).

1.6.1.5 Other Federal Agencies

For individual new uranium ISL facilities proposed near or on federally managed lands, agencies such as the Bureau of Land Management (BLM), the U.S. Forest Service, or National Park Service may have jurisdiction or special expertise that leads to a role in reviewing applications for these facilities. The Bureau of Indian Affairs has responsibilities under 25 CFR Part 216 to evaluate mineral leases involving lands held in trust for Native American tribes. Other federal agencies that may be consulted on specific resource areas include the U.S. Army Corps of Engineers (wetlands) and the U.S. Fish and Wildlife Service (endangered and threatened species).

1.6.2 Tribal Agencies

Native American tribes do not formally have licensing authority over uranium ISL facilities. Consultations with Native American tribes would be conducted in a government-to-government relationship that exists based on applicable federal law and treaties (NRC, 2003a) during the ISL licensing process. EPA can authorize tribes to implement specific environmental permitting programs. Tribes may also have their own local laws that impact ISL facilities. Additionally, tribes may have a tribal historic preservation officer (THPO) that would coordinate with NRC to support cultural resource inventories for ISL facility applications.

1.6.3 State Agencies

Individual states have regulatory authority over construction, operation, aquifer restoration, and decommissioning and reclamation at uranium ISL facilities through state-administered permitting processes. For the purposes of the Draft GEIS, specific agencies within each state that have regulatory authority over uranium ISL facilities are identified in the following sections.

1.6.3.1 Wyoming Department of Environmental Quality

The lead agency for permitting uranium ISL facilities in Wyoming is the Wyoming Department of Environmental Quality (WDEQ). With statutory authority from the Federal Surface Mining Reclamation and Control Act and the Wyoming Environmental Quality Act, the Land Quality Division within WDEQ administers and enforces permits and licensing requirements for all operators engaged in land-disturbing activities related to mining and reclamation within Wyoming. In the context of Wyoming regulations, uranium ISL facilities are considered to be noncoal mining activities that are subject to Land Quality Division permits. Each operation must be covered by a reclamation bond to provide financial surety that reclamation requirements can be met. Through its review and consultation program, the Wyoming State Historic Preservation Office (SHPO) coordinates with NRC and WDEQ to support cultural resource inventories for uranium ISL facilities.

1.6.3.2 Nebraska Department of Environmental Quality

The Nebraska Department of Environmental Quality (NDEQ) regulates air and water quality, with statutory authority from the Nebraska Environmental Protection Act. General water quality standards and use classifications are established in Title 117 (surface water) and Title 118 (groundwater) of the Nebraska Administrative Code (NDEQ, 2006a,b). The Nebraska NPDES program is described in Title 119 (NDEQ, 2005), and the regulatory requirements for underground injection, mineral production wells, and waste disposal wells related to ISL uranium recovery are governed by UIC requirements in Title 122 of the Nebraska Administrative Code (NDEQ, 2002a). The Nebraska SHPO is a division of the Nebraska State Historical Society. The Nebraska SHPO manages historic preservation programs within the state, which includes developing and maintaining a statewide historic preservation plan and providing supporting planning programs for other state agencies.

1.6.3.3 South Dakota Department of Environment and Natural Resources

With renewed interest in uranium resources in South Dakota, the 2006 State Legislature passed legislation to fill gaps in the existing state laws that govern uranium exploration and recovery. This legislation authorized the South Dakota Board of Minerals and Environment to develop rules to govern the construction, operation, monitoring, and closure of uranium and other ISL facilities under the South Dakota Mined Land Reclamation Act (South Dakota Codified Law 45–6B). The final rules were adopted in April 2007 (South Dakota Department of Environment and Natural Resources, 2007a). The South Dakota SHPO is a program of the South Dakota State Historical Society within the Department of Tourism and State Development. The South Dakota SHPO manages historic preservation programs within the state and coordinates and plans historic preservation efforts across the state.

1.6.3.4 New Mexico Environmental Department

The New Mexico Environmental Department was established under the provisions set forth in the Department of the Environment Act by the 40th State Legislature, enacted July 1, 1991 (Laws of 1991, Chapter 25). With the exception of potential facilities in the Navajo Nation and other Native American tribal lands, the New Mexico Environmental Department, with statutory authority from the New Mexico Oil and Gas Act and the New Mexico Water Quality Act, has permitting authority over uranium ISL facilities through its state-administered UIC program. The New Mexico SHPO is part of the Historic Preservation Division within the New Mexico Department of Cultural Affairs. The New Mexico SHPO administers historic preservation programs within the state and provides information and technical assistance to state agencies, local governments, and private owners.

1.7 Licensing and Permitting Process for a Uranium ISL Facility

As noted in Section 1.6, NRC has statutory authority through the Atomic Energy Act and UMTRCA to regulate uranium ISL facilities. In addition to obtaining an NRC license, uranium ISL facilities also must obtain the necessary permits from the appropriate federal, tribal, and state agencies. The NRC licensing process and other potential federal, tribal, and state permitting processes are briefly discussed in this section to provide a basic understanding of potential permitting requirements for uranium ISL facilities in the four geographic regions

identified previously in Figure 1.1-1. This is not intended to be an exhaustive description of all permits that may be necessary for a specific facility.

1.7.1 The NRC Licensing Process

The NRC process for licensing ISL uranium recovery facilities is described in NRC (2003b) and illustrated in Figure 1.7-1. After receiving a license application for either a new facility or a restart/expansion of an existing facility, NRC conducts an acceptance review to determine whether the application is complete enough to support more detailed technical review. If NRC determines that a new license application is acceptable for detailed review, NRC will formally docket the application and publish a Notice of Availability of the application in the *Federal Register*. NRC's detailed technical review of a site-specific license application is composed of a safety review and an environmental review. NRC conducts the safety review to assess compliance with the regulatory requirements of 10 CFR Part 20 and 10 CFR Part 40, Appendix A. In parallel with the safety review, the NRC staff is required under NEPA to conduct an environmental review for each license application. The NRC environmental protection regulations applicable to licensing actions are found in 10 CFR Part 51. The NRC hearing process (10 CFR Part 2) applies to NRC licensing actions and offers stakeholders a separate opportunity to raise concerns with the proposed action during the licensing process.

If a license is issued or a license amendment granted for expansion or restart of a facility, NRC ensures that the licensee complies with the conditions of its NRC license and the applicable regulations through an inspection program managed out of one of its four regional offices. The NRC Region IV office in Arlington, Texas, would manage inspection programs for ISL uranium recovery facilities located in each of the four regions analyzed in this Draft GEIS.

1.7.2 EPA Permitting

Under different environmental laws such as the Clean Water Act, the Safe Drinking Water Act, and the Clean Air Act, EPA has statutory authority to regulate activities that may affect the environment. EPA permitting that is most relevant for uranium ISL facilities is related to underground injection of the leaching solution (i.e., the lixiviant) and liquid effluents, surface discharge of treated waters and industrial and construction stormwaters, and air quality.

1.7.2.1 Water Resources

Under the Safe Drinking Water Act, EPA was granted primary authority to regulate underground injection and protect current and future sources of drinking water. Underground injection is broadly defined as the process of placing fluids underground through wells or other similar conveyance systems. EPA implements this responsibility through its UIC program (EPA, 2008a). EPA may administer the programs directly for states or tribal lands or jointly with the state government. Alternatively, EPA may also authorize individual states or tribes the opportunity to administer the UIC programs in accordance with EPA regulations. Currently, Wyoming, Nebraska, and New Mexico are authorized states. South Dakota administers the UIC program jointly with EPA, with the state administering the program for UIC Class II permits (EPA, 2008b).

1




Figure 1.7-1. General Flow Diagram of the NRC Licensing Process for 10 CFR Part 40 Licenses (From NRC, 2003a). ASLBP–Atomic Safety Licensing Board Panel; EA–Environmental Assessment; EIS–Environmental Impact Statement; FONSI–Finding of No Significant Impact; NEPA–National Environmental Policy Act; SER–Safety Evaluation Report.

Native American tribes can follow the same rules as states for obtaining authorization (40 CFR Part 145) if they are considered a "Federally Recognized Tribe" and have been designated for "Treatment Similar to a State." As of this writing (March 2008), no tribes have been granted authorization with respect to administering UIC programs. Tribes that want to enforce the federal UIC requirements must submit an application to EPA. If the application meets the minimum federal requirements for an authorized program, EPA will authorize the tribe to implement the UIC program. Two tribes currently are developing applications, but no tribal programs have been authorized yet (EPA, 2008c). The primacy application of the Ft. Peck Tribe in Montana is currently in hearings. The Navajo Nation has applied for authorization over all but Class III wells, which would include injection and production wells at uranium ISL facilities. In the absence of tribal authorization, EPA directly administers the UIC program on Indian Country lands, although tribes retain an option to establish additional requirements.

Unless authorized by rule or by permit, any underground injection is unlawful and violates the Safe Drinking Water Act and UIC regulations. Before an NRC-licensed uranium ISL facility can begin operations at any project site, the licensee must obtain the necessary UIC authorizations. These will include (1) an aquifer exemption (also called exempting the aquifer) as an underground source of drinking water or for the aquifer or portion of the aquifer where the uranium mobilization and recovery will occur and (2) a Class III UIC permit to operate injection wells. In addition, if deep well injection will be used to dispose of certain liquid wastes, the licensee will need to obtain a Class I UIC permit.

Under the provisions of the Clean Water Act, the NPDES program regulates discharges of pollutants from a point source into surface water of the United States. Operators of a point source discharge must obtain an NPDES discharge permit (EPA, 2008d). The permits contain limitations and conditions that are intended to protect surface water quality. Permits can cover either operational (industrial stormwater) discharges or construction phases. Construction stormwater NPDES authorizations are applied for and issued annually under a general permit based on projected construction activities. For a construction stormwater authorization, a notice of intent is filed before construction activities begin.

As with the UIC program, EPA either directly administers the NPDES permitting program or may authorize the permitting authority to a state or tribe (EPA, 2008e). State-implemented NPDES

UIC Permitting (from EPA, 2008a)

In the four regions covered in this Draft GEIS, the state implements UIC permitting for all five UIC permit classes for Wyoming, Nebraska, and New Mexico and for UIC Class II for South Dakota. Classes I and III are most applicable to uranium ISL facility operations.

- *Aquifer Exemption.* UIC criteria for exemption of an aquifer that might otherwise be defined as an underground source of drinking water are found at 40 CFR 146.4. These criteria include whether the aquifer is currently a source of drinking water and whether the water quality is such that it would be economically or technologically impractical to use the water to supply a public water system.
- *Industrial and Municipal Waste Disposal Wells (UIC Class I).* This permit class governs deep disposal of industrial, commercial, or municipal waste below the deepest usable aquifer. This type of injection uses wells and requires applied pressure. It includes all wells that dispose of waste on a commercial basis, even if the waste would be otherwise eligible for disposal into a Class II well (e.g., WDEQ, 2005, 1993). For uranium ISL facilities, this type of UIC permit is necessary to use deep well injection for waste disposal.
- *Mining Wells (UIC Class III).* These permits govern injection wells drilled to recover minerals. They include experimental technology wells; underground coal gasification wells; and wells for the *in-situ* recovery of materials such as copper, uranium, and trona. For uranium ISL facilities, this type of UIC permit covers wells that inject the lixiviant into the uranium mineralization.

programs (covering commercial industrial facilities like uranium ISL mills) are authorized in Wyoming, Nebraska, and South Dakota. EPA directly administers the NPDES program in New Mexico (EPA, 2008f).

1.7.2.2 Air Quality

EPA was given the primary responsibility to set standards and oversee the Clean Air Act. Similar to water protection programs, EPA may authorize the states, tribes, and local agencies to prevent and control air pollution. Under the Clean Air Act, EPA developed the following standards:

- National Primary and Secondary Ambient Air Quality Standards in 40 CFR Part 50
- National Emission Standards for Hazardous Air Pollutants in 40 CFR Part 40
- Prevention of Significant Deterioration in 40 CFR Part 52

As described in 40 CFR Part 51, Requirements for Preparation, Adoption, and Submittal of Implementation Plans, states must develop State Implementation Plans consisting of regulations, programs, and policies that describe how each state will control air pollution under the Clean Air Act. Agencies must obtain EPA approval for these implementation plans. The permitting process is a mechanism agencies use to put the implementation plans into effect. EPA's Tribal Authority Rule gives tribes the ability to: (1) develop air quality management programs, (2) write air pollution reduction rules, and (3) implement and enforce these rules. Similar to the states, tribes must obtain EPA approval for these implementation plans. The Clean Air Act permitting process is divided into two programs: the New Source Review program (pre-construction) and the Title V program (operation). Before any construction of or major modification to an ISL facility begins, a New Source Review permit scrutinizes the site-specific air quality impacts. The operation of the New Source Review permitting system varies by state (see Table 1.7-1).

Table 1.7-1. New Source Review Permit Summary Information for Nebraska, New Mexico, South Dakota, and Wyoming*

Area	Permitting Authority	Regulations
Nebraska†	State and local agencies	State Implementation Plan
New Mexico†	State and local agencies	State Implementation Plan
South Dakota†	State agency	State Implementation Plan‡
Wyoming†	State agency	State Implementation Plan
Indian Country (all four states)	Appropriate U.S. Environmental Protection Agency regional office	40 CFR 52.21
*Modified from U.S. Environmental Protection Agency. "Prevention of Significant Deterioration (PSD) Permit Program Status: May 2007." 2007. < http://www.epa.gov/nsr/where.html > (26 September 2007). †Except for Indian country. ‡Except for Prevention of Significant Deterioration permitting that is regulated by 40 CFR 52.21.		

Three types of New Source Review permits exist: (1) Prevention of Significant Deterioration, (2) nonattainment New Source Review, and (3) minor New Source Review. In attainment areas, Prevention of Significant Deterioration permits are required for major stationary pollutant sources that are new or making major modifications. In nonattainment areas, the nonattainment New Source Review permits are required for major stationary pollutant sources that are new or making major modifications. The minor New Source Review permits are for sources that do not require Prevention of Significant Deterioration or nonattainment New Source Review permits. A minor New Source Review permit is intended to support the Prevention of Significant Deterioration and nonattainment New Source Review programs by implementing permit conditions as needed that limit emissions from sources not covered by those two programs. For ISL facilities, NAAQS compliance status and emission levels determine which permit applies to a particular proposed facility.

Operating permits, called Title V permits, are required for most large sources and some smaller sources of air pollution. State or local agencies issue most Title V permits. In general, ISL facilities do not meet the emissions thresholds that invoke Title V requirements or require operating permits. However, to the extent that an ISL facility would meet the general requirements identified for EPA regulations at 40 CFR Part 70 and 71 (e.g., by exceeding either a general emissions threshold of 90.7 metric tons [100 short tons] for any air pollutant, lower thresholds for areas that are in nonattainment with air quality standards, or major source thresholds for hazardous air pollutants), the licensee or applicant would need to obtain the necessary Title V permit before beginning operations.

1.7.3 Other Federal Agencies

NRC and the Department of Transportation jointly regulate the safety of radioactive material shipments. The NRC regulations to transport radiological materials such as yellowcake and uranium-loaded resins are established in 10 CFR Part 71. For example, refined yellowcake is packaged and shipped in 208-L [55-gal], 18-gauge steel drums holding an average of 430 kg [950 lb]. The Department of Transportation classifies this as Type A packaging (49 CFR Part 171–189 and 10 CFR Part 71).

Because the federal government manages a portion of the land in the four geographic regions discussed in this Draft GEIS, BLM may control surface access at uranium ISL sites proposed for federal lands. BLM administers grazing on public ranchlands through field offices located in each state. The licensee must obtain the necessary mineral rights and environmental clearances from BLM for surface disturbances and approval for temporary occupancy. BLM requires (per 43 CFR 3809) the ISL licensee or applicant to submit a Plan of Operations. The BLM-required information can be (and usually is) included as part of the applicant's state-required forms/applications. Unlike NRC, BLM considers all mineral recovery to be mining. BLM regulates land use for operations proposed on BLM land and where the surface rights are privately owned and the mineral rights are under federal jurisdiction.

1.7.4 Tribal Agencies

Like States, Native American tribes can be authorized to implement the EPA Clean Water Act and Clean Air Act programs and can have their own permitting authority (e.g., Navajo Nation Environmental Protection Agency). This is discussed further in sections 1.7.2.1 and 1.7.2.2. Additionally, NRC has a responsibility to consult with tribes; the process for doing so is discussed in Chapter 9 of the Draft GEIS.

At least one tribe, the Navajo Nation, has enacted tribal legislation that prohibits all uranium processing activities. On April 29, 2005, Navajo Nation President Joe Shirley, Jr. signed the Diné Natural Resources Protection Act of 2005. The Navajo ban on uranium milling and processing presents a number of complex legal and policy issues, including whether a particular site falls under the definition of "Navajo land" in the Diné Natural Resources Protection Act of 2005. This latter issue is currently being litigated in the U.S. Court of Appeals for the 10th Circuit in a case brought against EPA with respect to certain proposed uranium processing sites in New Mexico. However, the fundamental question the Navajo ban poses is the relationship between the laws of the Navajo Nation and the laws and regulations of other governmental organizations, such as the NRC.

The NRC Commission's approach to these types of jurisdictional issues has been to fulfill NRC's statutory mandate to evaluate license applications and determine whether a particular application complies with the Atomic Energy Act and NRC regulations. At the same time, NRC recognizes that other governmental entities, in this case the Navajo Nation, may also have jurisdiction over some issues. The Commission acknowledges and recognizes that the Navajo Nation has certain sovereign powers under federal law. In general, although a license applicant may demonstrate that it meets the Atomic Energy Act and NRC regulations and thereby receives an NRC license, the applicant may nonetheless need to address other applicable requirements and obtain other necessary permits from appropriate regulatory authorities to go forward with its project.

1.7.5 State Agencies

The following sections briefly describe relevant state permitting requirements for Wyoming, Nebraska, South Dakota, and New Mexico.

1.7.5.1 Wyoming

WDEQ provides general guidance on Wyoming regulatory requirements for ISL operations in several reports (WDEQ, 2000a, 2005). WDEQ issues state permits relevant to ISL uranium recovery operations under Title 35, Chapter 11, of the Wyoming Environmental Quality Act. Most of these permits are related to water supply and air and water quality issues and include aquifer exemption; UIC Class I, III, and V permits; and NPDES permits (WDEQ, 2007, 2005, 2001, 2000b, 1993, 1984). Wyoming requires UIC Class III permits for injection wells in areas not previously mined using conventional mining and milling. UIC Class V permits are required for injection wells leaching from older conventional operations. In addition, the WDEQ Land Quality Division issues permits to mine for noncoal resources and *in-situ* recovery operations (WDEQ, 2003, 2000a). These permits identify site-specific requirements related to establishing baseline conditions (e.g., water, soils, vegetation, cultural values) and establishing reclamation bonds based on estimated site-specific costs. Wyoming also implements the NPDES program

regarding discharges to surface waters. With regard to air quality permitting, WDEQ establishes the NAAQS requirements (WDEQ, 2006) (see Table 1.7-1). In addition, the Wyoming State Land Use Planning Act established a State Land Use Commission to govern leases, easements, and temporary uses of state lands. The state also regulates drilling and well spacing and requires drilling permits for wells regardless of land ownership.

1.7.5.2 Nebraska

The regulations established in Title 122 of the Nebraska Administrative Code ensure proper well construction and regulate the injection of fluids containing potential contaminants into, above, or below underground sources of drinking water. NDEQ must approve injection wells, which must be operated and managed in accordance with the applicable NDEQ regulations. NDEQ issues and reviews UIC permits, conducts inspections, and performs compliance reviews for wells that inject fluids into the subsurface to ensure that injection activities comply with state and federal regulations and that groundwater is protected from potential contamination sources. Similar to WDEQ in Wyoming, NDEQ has authority over and manages Class I, III, and V wells in Nebraska. Injection wells not included in the other specific classes are considered Class V wells. In Nebraska, regulations adopted in 2002 prohibit a number of Class V wells types, including radioactive waste disposal wells. The NDEQ UIC program is currently closing existing waste disposal systems that fall into these prohibited types. EPA reviews and approves the aquifer exemption portion of the NDEQ UIC program (40 CFR 146.4). Nebraska also implements the NPDES program regarding discharges to surface waters. With regard to air quality permitting, NDEQ establishes the ambient air quality standards through a state-administered NAAQS program described in Title 129 of the Nebraska administrative code (NDEQ, 2002b).

1.7.5.3 South Dakota

As described in Section 1.7.3.3, recent legislation passed in South Dakota establishes permitting requirements for uranium recovery activities. Activities covered under these permits include sinking shafts; tunneling; and drilling test holes, cuts, or other works to extract samples (including bulk samples) to confirm the commercial grade of a uranium deposit before mining operations or test facility development begins. Uranium milling, including ISL uranium recovery, requires a state mine permit issued under South Dakota Codified Law 45-6B and South Dakota Administrative Rule Chapter 74:29. The Board of Minerals and Environment evaluates permit applications for uranium exploration in South Dakota (South Dakota Department of Environment and Natural Resources, 2007a, 2006). South Dakota implements the NPDES program regarding discharges to surface waters. The South Dakota Department of Environmental and Natural Resources is the air quality permitting authority through its NAAQS program (South Dakota Department of Environment and Natural Resources, 2007b).

1.7.5.4 New Mexico

Water quality standards in New Mexico are established in accordance with Water Quality Control Commission regulations in Title 20, Chapter 6, of the New Mexico Administrative Code. The New Mexico Environmental Department administers the state's UIC programs, excluding Native American tribal lands. The state's authority does not extend to any parts of the proposed project that would be on Native American tribal lands, such as allotments, land held in trust for the Navajo Nation, and land within a dependent Indian community, whereas EPA retains authority over UIC permitting. EPA Region IX administers the federal UIC program for all Navajo Indian country. For ISL uranium milling operations in Indian country (including Navajo

Indian lands) in New Mexico, an operator must obtain a Class III injection well permit and an aquifer exemption from EPA requiring aquifer cleanup and monitoring to protect surrounding underground sources of drinking water. For operations outside Indian lands in New Mexico, operators need to obtain the Class III injection well permit and a temporary aquifer designation from New Mexico Environmental Department, subject to EPA review and approval. EPA directly administers the NPDES program for surface water discharges in New Mexico. With regard to air quality permitting, the New Mexico Environmental Department is the permitting authority through its NAAQS program (New Mexico Environmental Department, 2002).

1.8 Use of the GEIS in the NRC Licensing Process

NRC plans to use the GEIS to fulfill its requirement at 10 CFR 51.20(b)(8) to prepare an environmental impact statement or supplement to an environmental impact statement, for site-specific ISL license applications. NRC environmental regulations in Appendix A to Subpart A of Part 51 discuss the format for presentation of material in environmental impact statements. In particular, Section 1(b) states "[T]he techniques of tiering and incorporation by reference described respectively in CEQ's NEPA regulations 40 CFR 1502.20 and 1508.28 and 40 CFR 1502.21 may be used as appropriate to aid in the presentation of issues, eliminate repetition or reduce the size of the environmental impact statement."

NRC also uses other CEQ regulations as guidance. In this case, CEQ's regulation 40 CFR 1502.4 allows, and in some cases requires, preparation of EISs for "broad federal actions." In preparing EISs on broad actions, the CEQ offers different approaches for agencies to take in their evaluations. These include evaluating proposals: (1) geographically (i.e., those actions occurring in the same general location) and (2) generically (i.e., those actions which have relevant similarities, such as common timing, impacts, alternative, methods or implementation, media or subject matter).

NRC plans to use tiering and incorporation by reference for environmental reviews of site-specific ISL license applications. Tiering (defined in 40 CFR 1508.28) is a procedure by which more specific or more narrowly focused environmental documents can be prepared without duplicating relevant parts of previously prepared, more general, or broader documents. The more specific environmental document incorporates by reference the general discussions and analyses from the existing broader document and concentrates on the issues and impacts of the project which are not specifically covered in the broader document. Often, other federal agencies refer to this broader document as a Programmatic EIS (or PEIS). The NRC uses the term Generic Environmental Impact Statement (GEIS) to refer these broader environmental documents.

The NRC Safety Review

In addition to meeting its responsibilities under the Atomic Energy Act of 1954, as amended, NRC prepares a Safety Evaluation Report to analyze the safety of the proposed action and assess its compliance with applicable NRC regulations.

The safety and environmental reviews are conducted in parallel (Figure 1.7-1). Although there is some overlap between the content of a Safety Evaluation Report and the environmental review document, the intent of the documents is different.

To aid in the decision process, the environmental review document summarizes the more detailed analyses included in the Safety Evaluation Report. For example, the environmental review document would not address how accidents are prevented but the environmental impacts that would result if an accident occurred.

Much of the information describing the affected environment in the environmental review document also is applicable to the Safety Evaluation Report (e.g., demographics, geology, and meteorology) (NRC, 2003b).

In this GEIS, NRC evaluates the potential environmental impacts of the relatively standard technology used in ISL facilities as operated in specified geographic areas. Relevant portions of this GEIS can then be incorporated by reference into the NRC's site-specific environmental review. In some cases, the site-specific environmental review will be an environmental assessment (EA) that supports a Finding of No Significant Impact (FONSI). In other cases, a site-specific EIS will be developed to analyze topic areas where a FONSI cannot be supported.

Section 1.7.1 summarizes NRC's licensing process. The following discussion provides a more detailed description of how the NRC staff will use the GEIS as part of the staff's environmental reviews for new ISL license applications.

1.8.1 Applicant's or Licensee's Environmental Report

License applicants must submit an environmental report to support their application for an NRC license to possess and use source material for ISL uranium milling. NRC regulations at 10 CFR 51.45 list the general content of the environmental report to include, among other things:

- A description of the proposed action
- A statement of its purposes
- A description of the environment affected
- Consideration of the impact of the proposed action on the environment
- Identification of any adverse environmental effects that cannot be avoided
- Discussion of alternatives to the proposed action

To help potential uranium milling license applicants develop their environmental reports, NRC provides additional guidance in

- Regulatory Guide 3.46, "Standard Format and Content of License Applications, Including Environmental Reports, for *In-Situ* Uranium Solution Mining" (NRC, 1982)
- NUREG-1569, "Standard Review Plan for *In-Situ* Leach Uranium Extraction License Applications" (NRC, 2003a)
- NUREG-1748, "Environmental Review Guidance for Licensing Actions Associated with NMSS Programs" (NRC, 2003b)

1.8.2 Acceptance Review of the License Application and Environmental Report

After receiving the license application and accompanying environmental report, the NRC staff first reviews the application and environmental report for completeness. This initial "acceptance review" ensures that the application and environmental report are comprehensive and address all relevant aspects of the applicant's proposed actions. When the NRC staff determine that the application is acceptable for detailed technical review, the application is officially docketed in accordance with NRC's regulations at 10 CFR Part 2. Then NRC publishes in the *Federal Register* notice of the public availability of the application and accompanying notice of opportunity for hearing on the application.

1 In their subsequent detailed technical review of an ISL license application, the NRC staff
2 analyzes both the health and safety impacts (documented in a Safety Evaluation Report) and
3 the potential environmental impacts of the proposed action (discussed in a separate
4 environmental review document—either an EA or an EIS).

5 6 **1.8.3 NRC's Site-Specific Environmental Review**

7
8 To meet its NEPA obligations for a site-specific license application, the NRC staff will conduct
9 an independent, detailed evaluation of the potential environmental impacts of the applicant's
10 proposed action to construct, operate, and decommission an ISL facility. This evaluation will
11 use the conclusions reached in the GEIS to the extent applicable to the specific site.

12
13 In their environmental review, the NRC staff can request additional information from the
14 applicant. These requests require the applicant to provide the information and data the NRC
15 staff consider necessary to conduct their review and reach their environmental conclusions.

16
17 As the basis for their independent evaluation, the NRC staff relies initially on the applicant's
18 environmental report for background information on the proposed action, including the potential
19 ISL facility's location, the extent of proposed operations and schedule, and the surrounding local
20 and regional affected environment. The NRC staff confirms important attributes of these
21 descriptions through visits to the proposed site location and vicinity, independent research
22 activities, and consultations with appropriate federal, tribal, state, and/or local agencies. The
23 NRC staff compares relevant aspects of the applicant's description of its proposed facility, its
24 use of the ISL process, and the affected environment to the descriptions of these aspects in the
25 GEIS. To the extent applicable, the NRC staff incorporates by reference the GEIS descriptions
26 into the site-specific environmental document.

27
28 The NRC staff will focus on the applicant's assessment of potential environmental impacts from
29 the proposed action and the identified alternatives. In its site-specific environmental review
30 document, NRC will evaluate a reasonable range of alternatives that may include alternatives
31 not identified by the applicant. NRC's independent evaluation of potential environmental
32 impacts will be conducted for each of the environmental resource areas identified in NRC
33 (2003b) (e.g., air quality, transportation, groundwater). In the specific assessment, the NRC
34 staff will evaluate the applicant's analysis of the potential impacts to each resource area, and to
35 the extent needed, independently confirm and verify essential aspects of the analysis. The
36 NRC staff may use computer codes and other verification techniques for these
37 confirmatory assessments.

38
39 With respect to the GEIS, the purpose of the NRC staff's site-specific impacts assessment is to
40 evaluate whether the conclusions concerning the potential environmental impacts identified in
41 the GEIS for that resource area can be adopted in the site-specific document. The NRC staff
42 may find that the GEIS conclusions for a specific resource area can be adopted in full, only in
43 part, or not at all. For those cases in which the GEIS conclusions can be adopted only in part or
44 not at all, the NRC staff will determine whether development of a site-specific EA or EIS is
45 appropriate due to the significance of the differing environmental impacts. The NRC staff will
46 document its decision regarding the adoption of the GEIS conclusions in the site-specific
47 environmental review document.

1.8.4 Public Participation Activities

As discussed previously, the NRC staff may prepare either an EA or an EIS for the site-specific license application (see Figure 1.7-1). If the NRC staff concludes that it needs to prepare a site-specific EIS, a notice of intent will be published in the *Federal Register*. Then, the NRC staff will follow the public participation procedures outlined in 10 CFR Part 51, which include requests for public input on the scope of the EIS and for public comment on the draft EIS for ISL applications. However, if the NRC staff determines that an EA is appropriate, the staff will make a draft of the EA and accompanying draft FONSI available for public comment before taking any licensing action on the applicant's proposal. The NRC staff will address public comments received on the draft EA/FONSI in the staff's final environmental review document. This approach is consistent with NRC regulations at 10 CFR 51.33 and was noticed in the *Federal Register* on September 27, 2007 (72 FR 54947).

As stated in Section 1.8.2, upon acceptance of a license application for detailed technical review, NRC publishes in the *Federal Register* a notice of opportunity for hearing on the application. Individuals or entities that may be affected by the potential issuance of the site-specific ISL license may request a hearing under NRC's formal hearing process. 10 CFR Part 2 provides the requirements needed to be granted a hearing.

1.8.5 NRC's Final Environmental Review Document and Findings

The NRC staff will issue a final EIS or final EA/FONSI as part of the licensing review for each site-specific license application. These final documents will provide the NRC staff's site-specific environmental review determinations that consider public input and the evaluations in the GEIS, to the extent applicable. The final environmental document and the site-specific Safety Evaluation Report together form the basis for the NRC's decision on whether to issue a 10 CFR Part 40 source material license to the applicant.

NRC's final action to issue a license may also be subject to a formal NRC hearing. As discussed in Section 1.8.4, 10 CFR Part 2 provides NRC's requirements concerning hearings.

1.9 References

Energy Information Administration. "Domestic Uranium Production Report—Quarterly." Data for 3rd Quarter 2007. <www.eia.doe.gov/cneaf/nuclear/dupr/qupd.html> (30 November 2007).

Energy Information Administration. "U.S. Uranium Reserve Estimates." Washington, DC: Energy Information Administration. 2004. <www.eia.doe.gov/cneaf/nuclear/page/reserves/ures.html> (14 September 2007).

EPA. "Underground Injection Control Program." 2008a. <<http://www.epa.gov/safewater/uic/index.html>> (12 February 2008).

EPA. "States' and Territories' Responsibility for the UIC Program." 2008b. <<http://www.epa.gov/safewater/uic/pdfs/Delegation%20status.pdf>> (12 February 2008).

EPA. "UIC Programs on Tribal Lands." 2008c. <<http://www.epa.gov/safewater/uic/tribal.html>> (12 February 2008).

1
2 EPA. "National Pollutant Discharge Elimination System (NPDES)." 2008d.
3 <<http://cfpub.epa.gov/npdes/index.cfm>> (12 February 2008).

4
5 EPA. "National Pollutant Discharge Elimination System (NPDES): State and Tribal Program
6 Authorization Status." 2008e. <<http://cfpub.epa.gov/npdes/statestribes/astatus.cfm>>.
7 (12 February 2008).

8
9 EPA. "National Pollutant Discharge Elimination System (NPDES): Specific State Program
10 Status." 2008f. <<http://cfpub.epa.gov/npdes/statestats.cfm?view=specific>>
11 (12 February 2008).

12
13 EPA. "Technical Report on Technologically Enhanced Naturally Occurring Radioactive
14 Materials From Uranium Mining: Volume 1—Mining and Reclamation Background."
15 EPA 402-R-05-007. Washington, DC: EPA, Office of Radiation and Indoor Air. 2007a.
16 <www.epa.gov/radiation/docs/tenorm/402-r-05-007-rev0607.pdf> (20 November 2007).

17
18 EPA. "Counties Designate Nonattainment or Maintenance for Clean Air Act's National Ambient
19 Air Quality Standards (NAAQS)." 2007b. <[http://www.epa.gov/oar/oaqps/greenbk/](http://www.epa.gov/oar/oaqps/greenbk/mapnmpoll.html)
20 [mapnmpoll.html](http://www.epa.gov/oar/oaqps/greenbk/mapnmpoll.html)> (29 September 2007).

21
22 EPA. "Uranium Location Database Compilation." EPA 402-R-05-009. Washington, DC:
23 EPA, Office of Radiation and Indoor Air. 2006. <[www.epa.gov/radiation/docs/tenorm/402-r-05-](http://www.epa.gov/radiation/docs/tenorm/402-r-05-009.pdf)
24 [009.pdf](http://www.epa.gov/radiation/docs/tenorm/402-r-05-009.pdf)> (20 November 2007).

25
26 NDEQ. "Nebraska Administrative Code, Title 117—Nebraska Surface Water Quality
27 Standards." Lincoln, Nebraska: NDEQ. July 2006a.

28
29 NDEQ. "Nebraska Administrative Code, Title 118—Groundwater Quality Standards and Use
30 Classification." Lincoln, Nebraska: NDEQ. March 2006b.

31
32 NDEQ. "Nebraska Administrative Code, Title 119—Rules and Regulations Pertaining to the
33 Issuance of Permits Under the National Pollutant Discharge Elimination System." Lincoln,
34 Nebraska: NDEQ. May 2005.

35
36 NDEQ. "Nebraska Administrative Code, Title 122—Rules and Regulations for Underground
37 Injections and Mineral Production Wells." Lincoln, Nebraska: NDEQ. April 2002a.

38
39 NDEQ. "Nebraska Administrative Code, Title 129—Ambient Air Quality Standards." Lincoln,
40 Nebraska: NDEQ. April 2002b.

41
42 New Mexico Environmental Department. "Title 20—Environmental Protection; Chapter 2—Air
43 Quality (Statewide); Part 3—Ambient Air Quality Standards." Albuquerque, New Mexico:
44 New Mexico Environmental Department. October 2002.

45
46 NRC. "Expected New Uranium Recovery Facility Applications/Restarts/Expansions: Updated
47 1/24/2008." 2008a. <[http://www.nrc.gov/info-finder/materials/uranium/2008-ur-projects-list-](http://www.nrc.gov/info-finder/materials/uranium/2008-ur-projects-list-public-012408.pdf)
48 [public-012408.pdf](http://www.nrc.gov/info-finder/materials/uranium/2008-ur-projects-list-public-012408.pdf)> (08 February 2008).

- 1 NRC. "Generic Environmental Impact Statement for Uranium Milling Facilities." 2008b.
2 <<http://www.nrc.gov/materials/fuel-cycle-fac/licensing/geis.html>> (08 February 2008).
3
- 4 NRC. "Agreement State Program." Washington, DC: NRC. 2007a.
5 <<http://www.nrc.gov/about-nrc/state-tribal/agreement-states.html>> (03 December 2007).
6
- 7 NRC. "Notice of Intent To Prepare a Generic Environmental Impact Statement for Uranium
8 Milling Facilities." *Federal Register*. Vol. 72. pp. 40344–40346. July 24, 2007b.
9
- 10 NRC. "Revised Notice of Intent to Prepare a Generic Environmental Impact Statement for
11 Uranium Milling Facilities." *Federal Register*. Vol. 72. pp. 50414–50416. August 31, 2007c.
12
- 13 NRC. "Public Scoping Meeting Re: General Environmental Impact Statement (GEIS) for
14 Uranium Recovery." ML072670251. <<http://www.nrc.gov/reading-rm/adams.html>>.
15 Washington, DC: NRC. 2007d.
16
- 17 NRC. "Uranium Recovery GEIS—Transcript for 09/27/2007 Gallup, NM Public Scoping
18 Meeting." ML073090104. <<http://www.nrc.gov/reading-rm/adams.html>>. Washington, DC:
19 NRC. 2007e.
20
- 21 NRC. "Integrated Materials Performance Evaluation Program (IMPEP): Vol. 5—Governmental
22 Relations and Public Affairs." NRC Management Directive 5.6. Washington, DC: NRC.
23 February 2004.
24
- 25 NRC. NUREG–1569, "Standard Review Plan for *In-Situ* Leach Uranium Extraction License
26 Applications-Final Report." Washington, DC: NRC. June 2003a.
27
- 28 NRC. NUREG–1748, "Environmental Review Guidance for Licensing Actions Associated With
29 NMSS Programs—Final Report." Washington, DC: NRC. August 2003b.
30
- 31 NRC. Regulatory Guide 3.46, "Standard Format and Content of License Applications, Including
32 Environmental Reports, for *In-Situ* Uranium Solution Mining." Washington, DC: NRC. 1982.
33
- 34 NRC. NUREG–0706, "Final Generic Environmental Impact Statement on Uranium Milling."
35 Washington, DC: NRC. September 1980.
36
- 37 NRC. NUREG–0170, "Final Environmental Report on the Transportation of Radioactive
38 Materials by Air and Other Modes." Washington, DC: NRC. 1977.
39
- 40 Occupational Safety and Health Administration. "Memorandum of Understanding Between the
41 U.S. Nuclear Regulatory Commission and the Occupational Safety And Health Administration."
42 November 16, 1988. <[http://www.osha.gov/pls/oshaweb/](http://www.osha.gov/pls/oshaweb/owadisp.show_document?p_table=MOU&p_id=233)
43 [owadisp.show_document?p_table=MOU&p_id=233](http://www.osha.gov/pls/oshaweb/owadisp.show_document?p_table=MOU&p_id=233)> (12 February 2008).
44
- 45 Office of Technology Assessment. "Transportation of Hazardous Materials." Washington, DC:
46 Office of Technology Assessment. 1986. <<http://ntl.bts.gov/lib/000/600/652/8636.pdf>>
47 (12 February 2008).
48
- 49 South Dakota Department of Environment and Natural Resources. "Chapter 74:29:11—*In Situ*
50 Leach Mining." Pierre, South Dakota: South Dakota Department of Environment and Natural
51 Resources. April 2007a.

1
2 South Dakota Department of Environment and Natural Resources. "Air Quality Program."
3 2007b. Pierre, South Dakota: South Dakota Department of Environment and Natural
4 Resources. <<http://www.state.sd.us/denr/DES/AirQuality/airprogr.htm>> (25 October 2007).
5

6 South Dakota Department of Environment and Natural Resources. "Question and Answer
7 Sheet for the Preliminary Draft Rules Chapter 74:29:11 *In Situ* Leach Mining." Pierre,
8 South Dakota: South Dakota Department of Environment and Natural Resources. November
9 2006.
10

11 WDEQ. "Water Quality Rules and Regulations, Chapter 1, Wyoming Surface Water Quality
12 Standards." Cheyenne, Wyoming: WDEQ, Water Quality Division. April 2007.
13

14 WDEQ. "Wyoming Department Of Environmental Quality Air Quality Division Standards and
15 Regulations Chapter 2 Ambient Standards." Cheyenne, Wyoming: WDEQ, Water Quality
16 Division. January 2006.
17

18 WDEQ. "Water Quality Rules and Regulations, Chapter 8, Quality Standards for Wyoming
19 Groundwaters." Cheyenne, Wyoming: WDEQ, Water Quality Division. April 2005.
20

21 WDEQ. "Guideline No. 6, Noncoal; Application for a 'Permit To Mine' or an 'Amendment.'"
22 Cheyenne, Wyoming: WDEQ, Land Quality Division. January 2003.
23

24 WDEQ. "Water Quality Rules and Regulations, Chapter 16, Class V Injection Wells and
25 Facilities Underground Injection Control Program." Cheyenne, Wyoming: WDEQ, Water
26 Quality Division. July 2001.
27

28 WDEQ. "Guideline No. 4, *In-Situ* Mining." Cheyenne, Wyoming: WDEQ, Land Quality Division.
29 March 2000a.
30

31 WDEQ. "Water Quality Rules and Regulations, Chapter 3, Regulations for Permit To Construct,
32 Install or Modify Public Water Supplies, Wastewater Facilities, Disposal Systems, Biosolids
33 Management Facilities, Treated Wastewater Reuse Systems and Other Facilities Capable of
34 Causing or Contributing to Pollution." Cheyenne, Wyoming: WDEQ, Water Quality Division.
35 January 2000b.
36

37 WDEQ. "Water Quality Rules and Regulations, Chapter 13, Class I Hazardous Waste and
38 Non-Hazardous Waste Wells Underground Injection Control Program." Cheyenne, Wyoming:
39 WDEQ, Water Quality Division. March 1993.
40

41 WDEQ. "Water Quality Rules and Regulations, Chapter 11, Design and Construction Standards
42 for Sewerage Systems, Treatment Works, Disposal Systems or Other Facilities Capable of
43 Causing or Contributing to Pollution and Mobile Home Park and Campground Sewerage and
44 Public Water Supply Distribution Systems." Cheyenne, Wyoming: WDEQ, Water Quality
45 Division. May 1984.

2 IN-SITU LEACH URANIUM RECOVERY AND ALTERNATIVES

Chapter 2 provides information on uranium recovery using the *in-situ* leach (ISL) process. The first part of the chapter gives basic information on the type of uranium deposits that are amenable to ISL technology and an overview description of the different parts of an ISL facility. Sections 2.2 through 2.6 describe different stages of an ISL facility's lifecycle, including pre-construction, construction, operation, aquifer restoration, and decommissioning. Sections 2.7 through 2.10 include discussions of aspects such as occupational health radiation monitoring, waste management, transportation, and financial assurance that are common to all ISL uranium facilities and not confined to any one stage. Section 2.11 summarizes operational experience of ISL facilities regulated by the U.S. Nuclear Regulatory Commission (NRC). Sections 2.12 and 2.13 discuss the alternatives considered in this Draft GEIS.

As stated, this chapter is organized by different stages in the life of an ISL facility. NRC recognizes that other than the pre-construction phase, aspects of the other four phases could be performed concurrently. However, by describing the ISL process in terms of these stages, NRC considers that this aids in the discussion of the ISL process and in the evaluation of potential environmental impacts during the lifecycle of an ISL facility.

2.1 Overview of ISL Uranium Recovery

Only certain uranium deposits are amenable to the ISL recovery process. To understand why the ISL recovery process is an effective recovery method for certain uranium deposits, it is necessary to understand the chemical and physical characteristics of uranium ore. This section will describe the geochemistry of uranium, provide a brief geologic overview of uranium ore bodies in the four Draft GEIS regions, and a general description of ISL facilities.

2.1.1 Geochemistry of Uranium

Natural uranium occurs in minerals as each of these isotopes: U-238 (99.274 percent), U-235 (0.720 percent), and U-234 (0.0055 percent) (EPA, 2007a) and predominantly exists in one of two ionic states: U^{6+} (the uranyl oxidized ion) and U^{4+} (the uranous reduced ion) (EPA, 1995). In the oxidized (uranyl) state, uranium is more readily dissolved. In the uranous (U^{4+}) state, uranium solubility is very low (i.e., it does not readily dissolve in water). Common uranous minerals include uraninite (UO_2), pitchblende (a crystalline variant of uraninite), and coffinite [$U(SiO_4)(OH)_4$] (EPA, 1995; Nash et al., 1981).

Characteristics of Uranium Deposits That Are Amenable to ISL Extraction

Certain geologic and hydrological features make a uranium deposit suitable for ISL technologies (based on Holen and Hatchell, 1986):

- **Deposit geometry.** The operator defines well field boundaries based on the geometry of the specific uranium mineralization. The deposit should generally be horizontal and have sufficient size and lateral continuity to economically extract uranium.
- **Permeable host rock.** The host rock must be permeable enough to allow the mining solutions to access and interact with the uranium mineralization. Preferred flow pathways such as fractures may short circuit portions of the mineralization and reduce the recovery efficiency. The most common host units are sandstones.
- **Confining layers.** Hydrogeologic (formation) geometry must prevent uranium-bearing fluids (i.e., lixiviant) from vertically migrating. Typically, low permeability layers such as shales or clays confine the uranium-bearing sandstone both above and below. This isolates the uranium-producing horizon from overlying and underlying aquifers.
- **Saturated conditions.** For ISL extraction techniques to work, the mineralization should be located in a hydrologically saturated zone.

2.1.2 Physical Characteristics of Uranium Deposits

Uranium deposits subject to recovery in the United States are primarily found in four types of deposits: stratabound, breccia pipes, vein, and phosphatic (EPA, 1995). Deposits that are generally amenable to ISL recovery in the four Draft GEIS regions are stratabound deposits. These deposits are contained within a single layer (strata) of sedimentary rock. It is believed that these deposits were formed through the transport of uranium (and associated elements) by oxidizing groundwater (i.e., groundwater with chemical properties that cause the uranium ion to lose electrons) (EPA, 1995; Nash et al., 1981). The groundwater flowed through the uranium-containing rocks, causing the uranium to dissolve and leach from the rock. The uranium remained soluble in the groundwater until it encountered a reducing environment (i.e., an environment with chemical properties that caused the uranium ion to gain electrons), became less soluble in water and precipitated.

Depending upon the environmental conditions, stratabound deposits can take different physical forms and are typically described as either roll-front deposits or tabular deposits. Roll-front deposits (Figure 2.1-1) are found in basins in Wyoming, southwestern South Dakota and northwestern Nebraska. Tabular deposits (see Figure 2.1-2) are found in the Colorado Plateau, including northwestern New Mexico.




Figure 2.1-1. Simplified Cross-Section of Sandstone Uranium Roll-Front Deposits Formed by Regional Groundwater Migration (NRC, 1997a)

A roll-front deposit is a uranium ore-body deposited at the interface of oxidizing and reducing groundwater (EPA, 1995; Nash et al., 1981). In basins in Wyoming, oxidized groundwater containing uranium flowed through permeable sandstone beds until reducing groundwater was

reached, and the uranium precipitated out at this interface. The sandstone beds are generally confined by low- or semi-permeable units such as claystones, siltstones, mudstones, or shales. As the oxidizing and reducing environments migrated within the sandstone beds, the uranium ore deposited over a laterally extended area (EPA, 1995). These roll-front deposits have a crescent shape and may extend hundreds of meters [feet] in length, but may only be a few meters [feet] thick.

The tabular deposits of the Colorado Plateau were formed when oxidized groundwater with higher concentrations of uranium and vanadium flowed through zones of highly permeable organic matter (humates), gases (hydrogen sulfide), or liquids capable of reducing the uranyl ion (EPA, 1995). The uranium deposited in the areas where the reducing conditions were created. The deposits are typically tabular in shape and can be found in sandstones, limestones, siltstones, and conglomerates scattered throughout various portions of the Colorado Plateau, including northwestern New Mexico. The tabular deposits found in northwestern New Mexico result from organic matter and occur in sandstones and siltstones. These deposits can range from about 0.5 to 2 m [2 to 6 ft] thick and hundreds of meters [feet] wide. These deposits have provided over 50% of the total uranium production in the United States (EPA, 1995).




Figure 2.1-2. Schematic Diagram of the Different Types of Stratabound Uranium Deposits in the Grants Uranium District, New Mexico (Modified from Holen and Hatchell, 1986)

Uranium concentrations in the ore deposit vary depending on system geochemistry and hydrology. For example, in New Mexico, uranium deposits typically contain about 0.2 to 0.3 percent U_3O_8 by weight, while deposits in Wyoming contain about less (about 0.1 to 0.25 percent) (Energy Information Administration, 2004; McLemore, 2007). The depth to the uranium mineralization ranges from about 100–300 m [328 to 984 ft] (e.g., Church Rock,

1 New Mexico; Gas Hills, Wyoming; Smith Ranch, Wyoming, and Crow Butte, Nebraska) to
2 greater than 560 m [1,840 ft] at Crownpoint, New Mexico. The most common uranium minerals
3 in roll-front deposits are uraninite (UO_2), pitchblende, and coffinite [$\text{U}(\text{SiO}_4)(\text{OH})_4$]. Minor
4 quantities of the uranium-vanadium mineral tyuyamunite [$\text{Ca}(\text{UO}_2)_2(\text{VO}_4)_2 \cdot \text{H}_2\text{O}$] are also typically
5 present (Nash, et al., 1981).
6

7 **2.1.3 General Description of ISL Facilities**

8
9 This section briefly describes the layout of an ISL facility. More detailed descriptions of the
10 individual stages of ISL uranium recovery (construction, operations, aquifer restoration,
11 decommissioning/reclamation) are included in Sections 2.3 through 2.6. A commercial ISL
12 facility consists of both an underground and a surface infrastructure. The underground
13 infrastructure includes injection and production wells drilled to the uranium mineralization zone,
14 monitoring wells drilled to the adjacent overlying and underlying aquifers, and perhaps deep
15 injection wells to dispose of liquid wastes. Pipelines to transfer groundwater extracted from the
16 well fields to the uranium processing circuit are buried to avoid freezing and thus are also
17 considered in this Draft GEIS to be part of the underground infrastructure.
18

19 ISL facilities also include a surface infrastructure that supports uranium processing. The
20 surface facilities can include a central uranium processing facility, header houses to control flow
21 to and from the well fields, satellite facilities that house ion exchange columns and reverse
22 osmosis for ground water restoration, and ancillary buildings that house administrative and
23 support personnel. Surface impoundments such as solar evaporation ponds may be
24 constructed to manage liquid effluents from the central processing plant and the ground water
25 restoration circuit (Figure 2.1-3).
26

27 The surface extent of a full-scale (i.e., commercial) ISL
28 facility includes a central processing facility and
29 supporting surface infrastructure for one or more well
30 fields (sometimes called mine units) encompasses
31 about 1,000 to 6,000 ha [2,500 to 16,000 acres] (NRC,
32 1992, 1997a) (see Section 2.11). However, the total
33 amount of land disturbed by such infrastructure and
34 ongoing activities at any one time is much smaller, and
35 only a small portion around surface facilities is fenced
36 to limit access (Figures 2.1-3 and 2.1-4). Using license
37 conditions, NRC establishes the total flow rates and the
38 maximum amount of uranium that can be produced
39 annually at a commercial ISL facility. NRC-licensed
40 flow rates typically range from about 15,100 to 34,000
41 L/min [4,000 to 9,000 gal/min], and licensed maximum
42 limits on annual uranium production range from about
43 860,000 to 2.5 million kg/yr [1.9 million to 5.5 million
44 lb/yr] of yellowcake (NRC, 1995, 1998a,b, 2006, 2007).
45 Actual production rates are somewhat lower (Energy Information Administration, 2008).
46
47

What is Yellowcake?

Yellowcake is the common name given to the uranium concentrate produced by milling and chemical processing. The yellowcake produced by most modern mills is a coarse, insoluble (does not dissolve in water) powder that is actually brown or black, not yellow. The name comes from the color and texture of the concentrates produced by early uranium milling production methods.

U_3O_8 depends on the processes used, but modern yellowcake typically contains 70 to 90 percent U_3O_8 by weight. Yellowcake is produced by all countries in which uranium is milled.




Figure 2.1-3. Layout of the Crow Butte Uranium Project in Dawes County, Nebraska (From Crow Butte Resources, Inc., 2007)




Figure 2.1-4. Well Heads and a Header House at Smith Ranch, Converse County, Wyoming

2.2 Pre-Construction

The applicant must characterize the potential site to support an application for a license to construct and operate an ISL facility (NRC, 2003a, Chapters 2 and 7). During the initial licensing review for a new ISL facility, NRC does not require a comprehensive discussion of all aspects of the site and of planned operations (NRC, 2003a). Instead, at this stage, the applicant needs to provide enough information to generally locate the uranium mineralization, understand the natural systems involved, and establish baseline conditions prior to operation. If a license is granted, the licensee would collect more detailed information as each well field is developed and brought into production (NRC, 2003a).

A number of general types of site baseline information to be provided by the license applicant are described in NRC guidance (NRC, 2003a, Chapter 2; 1982). Specific features of the site or its environs may also be identified and used by the applicant to support the proposed facility description. The applicant would provide maps to locate the proposed site, and identify proposed surface facilities, well fields, and other features of the ISL facility. In addition to providing information about the proposed site location and the environment in the vicinity of that location (e.g., water use, subsurface geology, hydrology, ecology, historical and cultural resources), the applicant also provides required population data and assessments of trends in population and industry patterns (NRC, 2003b, Appendix C).

Given the nature of the ISL uranium recovery process, hydrologic characterization of the site is a critical component of the applicant's pre-construction activities. This characterization describes surface-water features in the site area and the specific groundwater hydrogeologic setting, including detailed hydrogeologic and hydraulic descriptions of the proposed uranium production zone, adjacent aquifers, and low-permeability units that isolate the production zone.

Applicants are to determine baseline water quality for both the production zone and for adjacent un-mineralized zones (NRC, 2003a). An NRC-accepted list of constituents to be sampled is shown in Table 2.2-1, although an applicant can propose a list of constituents that is tailored to a particular location. To establish appropriate groundwater restoration standards, NRC requires that applicants and licensees establish pre-operational nonradiological and radiological groundwater quality baselines within the proposed permit boundaries and adjacent properties. These baseline conditions are based on samples collected over a period of at least 1 year, with a distribution that is sufficient to characterize the different aquifers and surface water bodies (NRC, 2003a).

Table 2.2-1. Typical Baseline Water Quality Parameters and Indicators*

Physical Indicators		
Specific Conductivity	Total Dissolved Solids†	pH‡
Major Elements and Ions		
Alkalinity	Chloride	Sodium
Bicarbonate	Magnesium	Sulfate
Calcium	Nitrate	
Carbonate	Potassium	
Trace and Minor Elements		
Arsenic	Iron	Selenium
Barium	Lead	Silver

**Table 2.2-1. Typical Baseline Water Quality Parameters and Indicators*
(continued)**

Trace and Minor Elements (continued)		
Boron	Manganese	Uranium
Cadmium	Mercury	Vanadium
Chromium	Molybdenum	Zinc
Copper	Nickel	
Fluoride	Radium-226§	
Radiological Parameters		
Gross Alpha [@]	Gross Beta	
*Based on U.S. Nuclear Regulatory Commission (NRC). NUREG-1569, "Standard Review Plan for <i>In-Situ</i> Leach Uranium Extraction License Applications—Final Report." Table 2.7.3-1. Washington, DC: NRC. June 2003. † Laboratory only. ‡ Field and laboratory determination. § If site initial sampling indicates the presence of thorium-232, then radium-228 should be considered in the baseline sampling, or an alternative may be proposed. @ Excluding radon, radium, and uranium.		

License applicants also collect site-specific data to establish background radiological characteristics of the site. These data may include measurements of radionuclides occurring in important flora and fauna species, soil, air, and surface and groundwaters that ISL operations could affect.

2.3 Construction

General construction activities associated with ISL facilities include drilling wells, clearing and grading associated with road construction and building foundations, building construction, trenching and laying pipelines, and building evaporation pond impoundments. Construction-related activities continue throughout much of the life of the project as different well fields are developed and additional wells and surface structures are added. For a satellite facility, the initial construction of the surface facilities would take about 2–3 months (NRC, 2004). Construction and testing of a well field may take about a year and a half (NRC, 2006), with about four to eight drill rigs and support vehicles operating in the field (NRC, 2004, 1997a). Well field construction would require about 50 to 75 contractors and full-time employees (NRC, 2004).

2.3.1 Underground Infrastructure

The underground infrastructure at an ISL facility is established to inject, produce, and monitor groundwater, and to transfer fluids between the wells and other production facilities.

2.3.1.1 Well Fields

Well Field Design. The licensee establishes the injection and production well patterns to recover uranium. The well patterns are developed for a specific site, and installation for a given well field is based on the subsurface geometry of the ore deposit. Various pattern shapes are used, although five-spot and seven-spot patterns are common (NRC, 2003a). A typical well arrangement using five- and seven-spot patterns is shown in Figure 2.3-1. Because roll-front uranium deposits normally have irregular shapes, some of the well patterns in a given well field are also irregular, and the licensee may alter well patterns to fit the size, shape, and boundaries of individual ore bodies.

1




Figure 2.3-1. Schematic Diagram of a Well Field Showing Typical Injection/Production Well Patterns, Monitor Wells, Manifold Buildings, and Pipelines (From NRC, 1997a)

2
3

These characteristics will also influence the number of wells in a well field. For example, at the Crow Butte ISL facilities in Dawes County, Nebraska, the number of injection and production wells varied from about 190 in the first well field (MU-1) to about 900 wells in later well fields (MU-5 and MU-6) (NRC, 1998b).

Three types of wells are predominant at uranium ISL facilities:

- Injection wells for introducing solutions into the uranium mineralization
- Production wells for uranium production
- Monitoring wells for assessing ongoing operations

In addition, the licensee or applicant may also drill deep injection wells permitted by the EPA or state for liquid waste disposal. Injection and production wells are connected to manifolds in a nearby header house (Figure 2.3-2). The manifolds connect to a series of pipelines that carry solutions to and from the recovery plant or satellite facility. Meters and control valves (usually computerized) in individual well lines monitor and control flow rates and pressures for each well to maintain water balance and to aid in identifying leaks in the system (Figure 2.3-3). The well field piping is typically high-density polyethylene pipe, polyvinyl chloride (PVC), and/or steel. Individual well lines and larger trunk lines to the recovery plant are buried below the frost line {e.g., 2 m [6 ft] in Wyoming} to prevent transferred solutions from freezing (NRC, 2006).




Figure 2.3-2. Manifold Inside Well Field Header House at an ISL Facility

Commercial-scale uranium ISL facilities usually have more than one well field. For example, the Crow Butte facility in Dawes County, Nebraska, has constructed 10 well fields since 1991 and has plans for an eleventh (Crow Butte Resources, Inc., 2007). The Reynolds Ranch satellite facility in Converse County, Wyoming, plans to include eight well fields (NRC, 2006). As described in Section 2.1.1, the well fields are developed in sequence, and at any one time, different well fields are likely to be in different stages of construction, operation, aquifer restoration, and decommissioning/reclamation (Crow Butte Resources, Inc., 2007). Construction and testing for each well field may take up to a year and a half before production begins (NRC, 2006). The locations and boundaries for each well field are adjusted as more detailed data on the subsurface stratigraphy and uranium mineralization distribution are collected during well field construction.




Figure 2.3-3. Computerized Meter for Monitoring Well Field Flow Rates

Well Drilling. Standard drilling techniques are used to develop ISL well fields. Temporary access roads for drilling rig trucks, support vehicles, and excavators lead to each well location. At the drilling location, a flat drill pad may be graded. At most ISL well fields, injection, production, and monitoring wells are drilled to the desired depth {e.g., 100–300 m [328–984 ft] for a target uranium production zone} by a standard method such as mud rotary drilling. In this method, a string of drill pipe and a drill bit is rotated against the formation. A water-based

drilling fluid (mud) is circulated through the hole to lubricate the bit and to carry the drilled material to the surface. A temporary mud pit is excavated directly in the ground next to the drill site to contain the drilling mud. Depending on the depth to the uranium mineralization and site-specific hydrogeological characteristics, other drilling methods may be used. While a well field is being drilled, detailed stratigraphic information and uranium ore occurrence data are collected. The locations and boundaries of a well field are then adapted to the subsurface geometry of a specific ore body. As the driller reaches the final depth of a well, it is usually logged with a variety of downhole geophysical tools (e.g., natural gamma ray logging, electrical resistivity) to characterize the well stratigraphy and reamed out to adjust the borehole diameter to construct a well. Residual cuttings and drilling fluids are typically held in the mud pit after drilling and construction activities are completed. Depending on state and local regulations, such pits are backfilled and graded or are alternatively emptied and cleaned, and residual solids and liquids are transported and disposed of offsite (NRC, 2006).

Well Construction. The geologic units above the aquifer of interest typically are sealed with steel or PVC casing grouted in place (Figure 2.3-4). This firmly sets the casing and prevents groundwater leakage from or to overlying aquifer(s). Grouts and casing materials are selected by the licensee or applicant to be inert with respect to the lixiviant and based on the depth of the well and anticipated well pressures. Depending on local hydrogeologic conditions, these well construction steps generally are followed:

- Sections of the uranium mineralized aquifers are left as open holes and screened with either steel or PVC screen material.
- Screens are then connected to the ground surface with steel or PVC riser pipes.
- The space between the casing and the borehole (i.e., the annulus) is filled with properly graded sand or gravel pack material, or the formation is simply left to collapse around the screen.
- A seal of bentonite clay is installed above the top of the screen.
- The annulus above the bentonite seal between the screen/riser pipe assembly and the borehole is typically grouted to the ground surface with a mixture of cement, bentonite, and water.

To make access and maintenance easier, well heads are completed above ground. Depending on local weather and land conditions, a variety of protective enclosures is used around the well head to protect it from the elements. Before the well head construction of an injection or production well is completed, the well is connected by underground piping to an injection or production manifold of a nearby header house.

Monitoring wells are not usually connected to any other structure but can have cables connected to different sensors in the well (NRC, 2006).




Figure 2.3-4. Cross Section of a Typical Injection, Production, or Monitoring Well Completed Using the Underreamed Method (Modified From NRC, 1997)

[1 in = 2.54 cm; 1 ft = 0.305 m]

Well Development and Integrity Testing.

Wells are usually developed using an air lift method or other pumping method appropriate for local conditions. Well development removes remaining drilling mud, cuttings, and fine particles (i.e., silt and clay) from inside the well, the screen, and surrounding gravel/sand pack. Development improves well yield by enhancing hydraulic communication between the undisturbed aquifer and the well. The licensee also performs a mechanical integrity test (MIT) to verify that the well casing does not fail, causing water loss during injection or recovery operations. In an MIT, the bottom and top of the casing are plugged (sealed) with an inflated downhole packer or similar sealing device. The well is pressurized, and pressure gauges monitor pressure changes inside the casing. Based on site-specific conditions, after maintaining a specified pressure for a specified period without a measurable decrease, the well casing is considered to have passed an MIT and the well is fit for injection or production operations (NRC, 2006).

Mechanical Integrity Testing

After completion and before brining into service, injection and recovery wells are tested for mechanical integrity. As described in NRC (2003a, Section 3.1.3), a packer is set above the well screen and the well casing is filled with water. At the surface, the well is pressurized with either air or water to 125 percent of the maximum operating pressure, which is calculated based on the strength of the casing material and depth. The well pressure is monitored to ensure significant pressure drops do not occur through borehole leaks. A pressure drop of no more than 10 percent in a period of 10 to 20 minutes indicates the casing and grout are sound and the well is fit for service. Well integrity tests are also performed if a well has been serviced with equipment or procedures that could damage the well casing. Additionally, each well is retested periodically (once each 5 years or less) to ensure its continued integrity.

2.3.1.2 Pipelines

The following piping systems are typically installed as part of the underground infrastructure:

- Between the central uranium processing facility or the satellite facility and the pump house for transporting lixiviant
- Between the pump house and well field for injecting and recovering lixiviant
- Between processing facilities and wastewater disposal sites (e.g., deep injection wells, evaporation ponds)

The network of process pipelines and cables required in ISL operations would be buried because of freezing temperatures that are common in the regions considered in this Draft GEIS and because of safety and land imprint issues. This network of pipelines and cables connects

- Injection and recovery wells to manifolds inside pumping/injection header houses
- Header houses to a central uranium processing facility or to satellite resin facilities (if present)
- Header houses to a central uranium processing facility or the central facility to deep injection wells used for liquid waste disposal

Depending on local winter conditions, burial trenches can be excavated as deep as 2 m [6 ft] below the ground surface to avoid any potential freezing problem (e.g., NRC, 2006).

High-density polyethylene, PVC, or steel pipes used to convey water, lixiviant, resin, and

wastewater are placed in these unlined trenches along with numerous electrical, communication, and sensor cables. Trenches are typically backfilled with native soil and graded to surrounding ground topography. Pipeline pressures are instrumented and recorded to monitor for potential leaks and spills that might result from the failure of pipeline fittings and valves.

2.3.2 Surface Facilities

ISL facilities require construction of different surface facilities, ranging from standard industrial buildings with associated power, water, and heating, ventilation, and air conditioning to specialized structures such as evaporation ponds (NRC, 2003a). Examples of surface facilities may include

- Central uranium processing facilities, with a typical footprint of about 3,060 m² [33,000 ft²] (NRC, 1998b)
- Satellite facilities {about 1,200 m² [13,000 ft²] (NRC, 2006)} that contain remote ion exchange columns
- Administration, operation, and field office or other support facilities
- Pump and header houses that house equipment to transfer lixiviant between the wells and pipelines
- Liquid effluent handling facilities, such as solar evaporation ponds. Typical evaporation ponds have surface areas ranging from 0.04 to 2.5 ha [0.1 to 6.2 acres] (NRC, 1998a; Crow Butte Resources, Inc., 2007)

In addition, to provide access between the well field and various surface facilities, the applicant or licensee would construct roads (dirt and/or paved) for

- Access to well fields and pump houses
- Access between the well fields/pump houses and the satellite facilities
- Access between the satellite facilities and the central processing facility
- Access between the processing plant and main transportation routes

The surface facilities and access roads are designed and built using standard construction techniques. Specific building codes are used as appropriate. Construction vehicles may include bulldozers, drilling rigs, water trucks, forklifts, pump hoist trucks, coil tubing trucks, pickup trucks, portable air compressors, and other support vehicles.

Evaporation ponds may be constructed to dispose of effluent from the processing circuit or from aquifer restoration activities. These impoundments are designed and constructed with liners and leak detection systems installed in accordance with applicable NRC guidance (NRC, 1977, 2003a, 2008). Embankments for these evaporation ponds are constructed to resist erosion from wave action in the pond. The size and shape of the ponds are designed based on the amount of water that must be managed and the evaporation rates for the region. Sufficient

space is conserved so that the contents of one pond may be transferred to another to allow any identified pond system leaks to be repaired and also to meet freeboard requirements from possible wave action.

2.4 Operations

Although specific operations will vary depending on the individual operator and site-specific characteristics, the ISL uranium recovery process generally involves two primary operations: (1) injection of barren lixiviant to mobilize uranium in underground aquifers and (2) extracting and processing the pregnant lixiviant in surface facilities to recover the uranium and prepare it for shipment.

2.4.1 Uranium Mobilization

During ISL operations, chemicals are added to the groundwater to produce a leaching solution or lixiviant. The lixiviant is injected into the production zone to mobilize (dissolve) uranium from the underground formation and subsequently remove uranium from the deposit.

2.4.1.1 Lixiviant Chemistry

The lixiviant that is selected must leach uranium from the host rock and keep it in solution during groundwater pumping from the host aquifer. Based on experience with conventional uranium milling, early ISL facilities tended to use aggressive acid-based lixiviants, such as sulfuric acid (International Atomic Energy Agency, 2001). These acid-based systems generally achieved high yield and efficient, rapid uranium recovery, but they also dissolved other heavy metals associated with uranium in the host rock and other chemical constituents that required additional remediation. In the United States, acid-based lixiviants have been used only for small-scale research and development operations [e.g., Nine Mile Lake and Reno Ranch in Wyoming (Mudd, 2001)], but have not been used in commercial operations (Davis and Curtis, 2007; International Atomic Energy Agency, 2005). Licensees or applicants may propose the use of acid-based lixiviants in the future. Other technologies that used ammonia-based lixiviants experienced difficulties: the ammonia tended to adsorb onto clay minerals in the subsurface. The ammonia desorbs slowly from the clay during restoration, and therefore the system requires that much larger amounts of groundwater be removed and processed during aquifer restoration (Energy Information Administration, 1995; Davis and Curtis, 2007). Although applicants or licensees may decide to use different lixiviants for a given deposit (see text box "Lixiviant Selection" in Section 2.4.1.2), ISL operations in the United States are expected to use alkaline lixiviants that are based on sodium carbonate-bicarbonate as the complexing agent and gaseous oxygen or hydrogen peroxide as the oxidizing agents (Table 2.4-1). For the purposes of the analyses presented in this Draft GEIS, it is assumed that alkaline lixiviants will be used in uranium recovery operations.

Basic Steps in Uranium Mobilization

- **Groundwater Injection.** The operator injects a nonuranium-bearing (barren) extraction solution or lixiviant through wells into the mineralized zone. The lixiviant moves through pores in the production zone, dissolving uranium and other metals.
- **Groundwater Extraction.** Production wells withdraw the resulting "pregnant" lixiviant, which now contains uranium and other dissolved metals, and pump it to a central processing plant or to a satellite processing facility for further uranium recovery and purification.

1

Table 2.4-1. Typical Lixiviant Chemistry (From NRC, 1998b)

Species	Range (in mg/L)*	
	Low	High
Sodium (Na)	≤400	6,000
Calcium (Ca)	≤20	500
Magnesium (Mg)	≤3	100
Potassium (K)	≤15	300
Carbonate (CO ₃)	≤0.5	2,500
Bicarbonate (HCO ₃)	≤400	5,000
Chloride (Cl)	≤200	5,000
Sulfate (SO ₄)	≤400	5,000
Uranium (as U ₃ O ₈)	≤0.01	500
Vanadium (as V ₂ O ₅)	≤0.01	100
Total Dissolved Solids	≤1,650	12,000
pH (in std unit)	≤6.5	10.5

* 1 mg/L is approximately equal to 1 part per million (ppm)

The principal geochemical reactions caused by the lixiviant are the oxidation and subsequent dissolution of uranium and other metals from the ore body (Davis and Curtis, 2007). These reactions are effectively the reverse of those that initially caused the uranium deposition. The oxidant (oxygen or hydrogen peroxide) in the lixiviant oxidizes uranium from the relatively insoluble tetravalent state (U⁴⁺) to the more soluble hexavalent state (U⁶⁺). Once the uranium is in the 6+ oxidation state, the dissolved carbonate/bicarbonate causes the formation of aqueous uranyl-carbonate complexes that maintain oxidized uranium in solution as uranyl ion (UO₂²⁺).

2.4.1.2 Lixiviant Injection and Production

Dissolved carbonate/bicarbonate lixiviants are created by introducing reagents such as sodium carbonate/bicarbonate or by injecting carbon dioxide gas (CO₂) into the groundwater. Carbon dioxide can also be added for pH control (Table 2.4-1). Lixiviant is pumped down injection wells to the mineralized zones, where it oxidizes and dissolves uranium from the sandstone formation (Figure 2.4-1). The uranium-bearing solution migrates through the pore spaces in the sandstone and is recovered by production wells. This uranium-rich (pregnant) lixiviant is pumped to the processing plant or satellite ion exchange facility, where the

Lixiviant Selection

The geology and groundwater chemistry determine the proper leaching techniques and chemical reagents ISL milling uses for uranium recovery. For example, if the ore-bearing aquifer is rich in calcium (e.g., limestone or gypsum), alkaline (carbonate) leaching might be used [e.g., as discussed by Hunkin (1977), acid systems were generally considered unsuitable for Texas deposits because of higher carbonate]. Otherwise, acid (sulfate) leaching might be preferable. The leaching agent chosen for the ISL operation may affect the type of potential contamination and vulnerability of aquifers during and after ISL operations.

For example, acid leaching ISL uranium recovery at Nine Mile Lake and Reno Ranch, Wyoming, presented two major problems: (1) gypsum precipitated on well screens and within the aquifer during uranium recovery, plugging wells and reducing the formation permeability (critical for economic operation) and (2) the precipitated gypsum gradually dissolved after restoration, increasing salinity and sulfate levels in groundwater (Mudd, 2001).

Typical ISL uranium recovery operations in the United States use an alkaline sodium bicarbonate system to remove the uranium from ore-bearing aquifers. Alkaline lixiviants are used in all currently active and proposed ISL facilities in Wyoming, Nebraska, and New Mexico (NRC, 2006, 2004, 1998a, 1997a; Energy Metals Corporation, U.S., 2007) (see Table 2.4-1). Alkaline-based ISL operations are considered to be easier to restore than acid mine sites (Tweeton and Peterson, 1981; Mudd, 1998).

uranium is extracted through a series of chemical processes. Stripped of its uranium, the now-barren lixiviant is recharged with carbonate/bicarbonate and oxidant and the solution is returned through the injection wells to dissolve additional uranium. This process continues until the operator determines that further uranium recovery is uneconomical.

During the uranium recovery process, the groundwater in the production zone becomes progressively enriched in uranium and other metals that are typically associated with uranium in nature. The most common metals are arsenic, selenium, vanadium, iron, manganese, and radium. These and other constituents such as chloride, which is introduced by the ion exchange resin system, are removed or precipitated from the groundwater during aquifer restoration after uranium recovery is completed. Aquifer restoration will be detailed in Section 2.5.

The production wells are normally positioned to pump pregnant lixiviant from a number of injection wells. After processing but before reinjection, about 1–3 percent of the lixiviant, called the production bleed, is removed from the circuit and disposed of (see Section 2.7.2). The purpose of the production bleed is to ensure that more groundwater is extracted than re-injected. Maintaining this negative water balance helps to ensure that there is a net inflow of groundwater into the well field to minimize the potential movement of lixiviant and its associated contaminants out of the well field.




Figure 2.4-1. Idealized Schematic Cross Section To Illustrate Ore-Zone Geology and Lixiviant Migration From an Injection Well to a Production Well (From NRC, 1997a)

Pregnant lixiviant is pumped from the well fields by submersible pumps located in each production well. In some cases, booster pumps are installed in the lines to the processing plants or satellite facilities. Given the seasonal temperature variation in the four regions considered in this Draft GEIS, the main injection and production lines to and from the processing plants will be buried up to several meters [feet] to prevent freezing. These lines are usually 10.2- to 35.6-cm [4- to 14-in] diameter high density polyethylene or PVC pipes. The pregnant lixiviant is enriched in uranium relative to groundwater {typically about 60 mg/L [0.0005 lb/gal]} and is also likely to contain the trace elements and contaminants as discussed previously. The pipeline pressures are monitored continuously for spills and leaks.

2.4.1.3 Excursions

As described previously, ISL operations may affect the groundwater quality near the well fields when lixiviant moves from the production zone and beyond the boundaries of the well field. These occurrences are known as excursions. These excursions can be caused by

- Improper water balance between injection and recovery rates
- Undetected high permeability strata or geologic faults
- Improperly abandoned exploration drill holes
- Discontinuity within the confining layers
- Poor well integrity, such as a cracked well casing or leaking joints between casing sections
- Hydrofracturing of the ore zone or surrounding units

NRC license and underground injection control (UIC) permit conditions require that licensees conduct periodic tests to protect against excursions. These include but are not limited to

- Conducting pump tests for each well field prior to operations within the well field to evaluate the confinement of the production horizon
- Continued well field characterization to identify geologic features (e.g., thinning confining layers, fractures, high flow zones) that might result in excursions
- Mechanical integrity testing of each well to check for leaks or cracks in the casing

An excursion that moves laterally away from the production zone is a horizontal excursion. Vertical excursions occur where barren or pregnant lixiviant migrates into other aquifers above or below the production zone.

2.4.1.4 Excursion Monitoring

Licensees must maintain groundwater monitoring programs (see Chapter 8) to detect both vertical and horizontal excursions and must have operating procedures to analyze an excursion and determine how to remediate it. Geochemical excursion indicators are identified based on

the well fields' pre-operational baseline water quality (see text box "Identifying Excursion Indicators and UCLs").

The spacing of horizontal excursion monitoring wells is based on site-specific conditions, but typically they are spaced about 90–150 m [300–500 ft] apart and screened in the production zone (NRC, 2003a, 1997a; Mackin, et al., 2001a; Energy Information Administration, 1995). The specific location and spacing of the monitoring wells is established on a site-by-site basis by license condition. It is often modified according to site-specific, hydrogeologic characteristics of the uranium deposit and as the licensee gains experience detecting, recovering, and cleaning up these excursions.

NRC licenses also include requirements to establish monitoring wells in overlying and, as appropriate, in underlying aquifers to detect vertical excursions.

Although uranium deposits are typically located in hydrogeologic units bounded above and below by adequately confining units, the possibility of vertical contaminant transport must be considered. Historically, these monitoring wells are more widely spaced than those within the host aquifer, although underlying aquifer monitoring wells may not be required under some circumstances (Mackin, et al., 2001a). There are general guidelines for monitoring well placement: (1) one monitoring well per 1.6 ha [4 acres] of well field in the first overlying aquifer, (2) one monitoring well per 3.2 ha [8 acres] in each higher aquifer, and (3) one monitoring well per 1.6 to 3.2 ha [4 to 8 acres] in the underlying aquifer. These monitoring wells are typically sampled every 2 weeks during operations.

An excursion is defined to occur when two or more excursion indicators in a monitoring well exceed their UCLs (NRC, 2003a). If an excursion is detected, the licensee takes several steps to notify NRC and confirm the excursion through additional and more frequent sampling (NRC, 2003a) (see Chapter 8). As described in NRC guidance (NRC, 2003a, Section 5.7.8.3), licensees typically retrieve horizontal and vertical excursions back into the production zone by adjusting the flow rates of the nearby injection and production wells to increase

Identifying Excursion Indicators and UCLs

The applicant or licensee proposes excursion indicators and upper control limits (UCLs) based on lixiviant content and baseline groundwater quality (see Section 2.2.7). NRC staff review and approve the excursion indicators and proposed UCLs. UCLs are set on a well field basis and are concentrations for excursion indicators that provide early warning if leaching solutions are moving away from the well fields. As described in NRC (2003a, Section 5.7.8.3), the best excursion indicators are easily measurable parameters that are found in higher concentrations during ISL operations than in the natural waters. For example, at most ISL uranium recovery operations, chloride is selected because it does not interact strongly with minerals in the subsurface, it is easily measured, and chloride concentrations are significantly increased during ISL operations. Conductivity, which is correlated to total dissolved solids, is also considered to be a good excursion indicator because of the high concentrations of different dissolved constituents in the lixiviant as compared to the surrounding aquifers (Staub, et al., 1986; Deutsch, et al., 1985). Total alkalinity (carbonate plus bicarbonate plus hydroxide) is used as an indicator in well fields where sodium bicarbonate or carbon dioxide is used in the lixiviant.

A minimum of three excursion indicators are selected, and the UCLs are determined using statistical analyses of the preoperational baseline water quality in the well field. The NRC staff has identified several statistical methods that can be used to establish UCLs. For example, in areas with good water quality (total dissolved solids less than 500 mg/L), the UCL may be set at a value of 5 standard deviations above the mean of the measured concentrations. Conversely, if the chemistry or a particular excursion indicator is very consistent, a concentration may be specified as the UCL. If baseline data indicate that the groundwater is homogeneous across the well field, the same UCLs may be used for all monitoring wells. Alternatively, if the water chemistry in the well field is highly variable, UCLs may be set for individual wells. An excursion is defined to occur when two or more excursion indicators in a monitoring well exceed their UCLs (NRC, 2003a).

process bleed in the area of the excursion. Vertical excursions are more difficult to retrieve, persisting for years in some cases (see Section 2.11.4). If an excursion cannot be recovered, the licensee may be required to stop injection of lixiviant into a well field (NRC, 2003a, Section 5.7.8.3).

2.4.2 Uranium Processing

Uranium is recovered from the pregnant lixiviant and processed as yellowcake in a multistep process (Figure 2.4-2). The following sections briefly describe key aspects of the uranium process circuit.

2.4.2.1 Ion Exchange

As pregnant lixiviant from the production wells enters the ion exchange circuit, it may either be stored in a surge tank or sent directly to the ion exchange columns (Figure 2.4-3). The number and size of ion exchange columns in the circuit may vary, depending on facility design. For example, at the Smith Ranch Uranium Project in Converse County, Wyoming, the ion exchange circuit consists of six pressurized downflow vessels, each with a volume of 14.2 m³ [501.5 ft³] (Stout and Stover, 1997). At the Crow Butte facility in Dawes County, Nebraska, the ion exchange circuit consists of eight upflow columns, with a recent addition of six downflow columns, each about 3.5 m [11.5 ft] in diameter and 4.6 m [15 ft] tall and a volume of about 44 m³ [1,554 ft³] (NRC, 2007; Crow Butte Resources, Inc., 2007). In the ion exchange columns, the uranium is adsorbed onto resin beads that selectively remove uranium from solution. The primary reaction is the exchange of the uranium carbonate complexes for chloride. The (now barren) lixiviant exits the ion exchange columns, is recharged with oxidant and bicarbonate, and is returned to the well field for reinjection and further uranium recovery. It carries chloride that was exchanged for uranium on the resin. The chloride content of the water in the ore-bearing aquifer builds up with time as the lixiviant is circulated and the resin is recharged. The production bleed discussed previously in Section 2.4.1 is removed downstream of the ion exchange columns, before re-injecting the barren lixiviant into the well field (see Figure 2.4-2).

When the resin beads in the ion exchange columns become saturated with uranium, the columns are taken offline and other columns are brought online. Some facilities may not process the ion exchange resins further (NRC, 2004, 2006). In these facilities (called satellite facilities), the resin is discharged to a truck and then transported to a facility that has the capacity for further processing of the uranium-loaded resin. Later sections of this Draft GEIS assess the hazards associated with transferring and transporting loaded ion exchange resin.

2.4.2.2 Elution

At ISL facilities that can process resin, after the resin is loaded with uranium, it enters the elution circuit. In addition, uranium-loaded resins transported from satellite plants in a remote ion exchange operation enter the processing circuit at this point. In the elution circuit, the uranium is washed (eluted) from the resin and the resin is made available for further cycles of uranium absorption. The resin may be eluted directly in the ion exchange column, or it may be transferred to a separate elution tank. In the elution process, the uranium is removed from the resin by flushing with a concentrated brine solution. This process returns chloride ions to the resin exchange sites, regenerating the resin at the same time that the uranium is released for further processing. A sodium carbonate or bicarbonate rinse is also used during this phase to keep the stripped uranium from precipitating in the elution vessel. The resulting uranium-rich




Figure 2.4-2. Flow Diagram of an ISL Uranium Recovery Process (Mackin, et al., 2001a)

solution is termed pregnant or rich eluant and typically contains 8 to 20 g/L [0.067 to 0.17 lb/gal] of uranium (Mackin, et al., 2001a). It is normally discharged to a holding tank. After enough pregnant eluant is obtained, it is moved to the precipitation, drying, and packaging circuit (Mackin, et al., 2001a).




Figure 2.4-3. Typical Ion Exchange Vessels in an ISL Facility

2.4.2.3 Precipitation, Drying, and Packaging

In the precipitation and drying circuit, the pregnant eluant is typically acidified using hydrochloric or sulfuric acid to destroy the uranyl carbonate complex. Hydrogen peroxide (H_2O_2) is then

1 added to precipitate the uranium as uranyl peroxide (UO_2O_2). Caustic soda (NaOH) or
2 ammonia (NH_3) is also normally added at this stage to neutralize the acid remaining in the
3 eluate. The (now barren) eluant is typically recycled. Water left over from these processes may
4 be reused in the eluant circuit or may be disposed as 11e.(2) byproduct material. Effluent
5 management is discussed in Section 2.7.2.

6
7 After the precipitation process, the resulting slurry is sent to a thickener where it is settled,
8 washed, filtered, and dewatered (Figure 2.4-4). At this point, the slurry is 30 to 50 percent
9 solids. This thickened slurry may be transported offsite to a uranium processing plant to
10 produce yellowcake (U_3O_8), or it may be filter pressed to remove additional water, dried and
11 packaged onsite.




Figure 2.4-4. A Typical Thickener for an ISL Uranium Processing Facility

12
13 For onsite processing, the slurry is next dried in the yellowcake dryer. Two kinds of yellowcake
14 dryers have been used: multihearth dryers and vacuum dryers. Older uranium ISL facilities
15 used gas-fired multi-hearth dryers. These dryers typically dry the yellowcake at about 400 to
16 620 °C [750 to 1,150 °F]. Because of the high temperatures involved, any organic contaminants
17 in the yellowcake (e.g., grease from bearings) will be completely burned and will exit the system
18 with the dryer offgas. This is advantageous because leftover organic residues in the packaged
19 yellowcake product may oxidize while in the drum, causing the drum to pressurize and burst due
20 to the evolution of gases (primarily CO_2) inside it (NRC, 1999). The offgas discharge from the
21 dryer is scrubbed with a high intensity venturi scrubber that is 95 to 99 percent efficient at

1 removing uranium particulates before they are released to the atmosphere. Solutions from the
2 scrubber are normally returned to the precipitation circuit and are processed to recover any
3 uranium particulates. As a result, the stack discharge normally contains only water vapor and
4 quantities of uranium fines that are managed to be below regulatory limits (see Sections 2.7.1
5 and Chapter 8).

6
7 Newer ISL facilities usually use vacuum yellowcake dryers. In a vacuum dryer (Figure 2.4-5),
8 the heating system is isolated from the yellowcake so that no radioactive materials are entrained
9 in the heating system or its exhaust. The drying chamber that contains the yellowcake slurry is
10 under vacuum. Therefore, any potential leak would cause air to flow into the chamber, and the
11 drying can take place at relatively low temperature {e.g., 149 °C [250 °F]}. Moisture in the
12 yellowcake is the only source of vapor. Emissions from the drying chamber are normally treated
13 in two ways. First, vapor passes through a bag filter to remove yellowcake particulates with an
14 efficiency exceeding 99 percent. Any captured particulates are returned to the drying chamber.
15 Then, any water vapor exiting the drying chamber is cooled and condensed. This process is
16 designed to capture virtually all escaping particles (Mackin, et al., 2001a).




Figure 2.4-5. Typical Vacuum Dryer for Uranium Yellowcake Processing at an ISL Uranium Processing Facility

19 The dried product (yellowcake) is removed from the bottom of the dryer and packaged in drums
20 for eventual shipping offsite. The packaging area normally has a baghouse dust collection
21 system to protect personnel and to minimize yellowcake release. Air from the baghouse dust
22

collection system is typically routed to the dryer offgas line and scrubber. During drum loading, the drum is normally kept under negative pressure via a drum hood with a suction line. The drum hood transports any released particulates to a baghouse dust collector. The filtered air from this baghouse joins the dryer offgas and is passed through the scrubber. Parameters important to the effective operation of the dryer must be monitored, and existing NRC regulations at 10 CFR Part 40, Appendix A, Criterion (8), prohibit dryer operations when these parameters are outside prescribed ranges. After the dried product is cooled, it is packaged and shipped in 208-L [55-gal] drums (Figure 2.4-6).



Figure 2.4-6. Labeled and Placarded 208-L [55-gal] Drum Used for Packaging and Shipping Yellowcake

2.4.3 Management of Production Bleed and Other Liquid Effluents

Uranium mobilization and processing produce excess water that must be properly managed. The production wells extract slightly more water than is re-injected into the host aquifer, which creates a net inward flow of groundwater in the well field. This production bleed is about 1 to 3 percent of the circulation rate, which can amount to an excess production of several tens to a hundred liters per minute (several tens of gallons per minute). As described in Section 2.4.1,

the production bleed is diverted from the ISL circuit after the uranium is removed in the ion exchange resin system, but before the lixiviant is recharged. This water still contains lixiviant and minerals leached from the aquifer. The excess water can be discharged to an evaporation pond or a deep well injection for disposal, or treated further for discharge to the environment (Section 2.7.2). Other liquid waste streams produced during ISL operation can include spent eluant from the ion exchange system, and liquids from process drains. These are handled in the same manner as the production bleed.

2.5 Aquifer Restoration

Aquifer restoration within the well field ensures that the water quality and groundwater use in surrounding sources of drinking water will not be adversely affected by the uranium recovery operation. Before ISL operations can begin, the portion of the aquifer designated for uranium recovery must be exempted from U.S. Environmental Protection Agency (EPA) regulatory protection, in accordance with the Safe Drinking Water Act (see Section 1.7.2.1). Groundwater adjacent to the exempted portion of the aquifer, however, must still be protected. The states authorized to implement the EPA groundwater protection program as well as the NRC require well field restoration to protect human health and the environment.

After uranium is recovered, the groundwater in the well field contains constituents that were mobilized by the lixiviant. Licensees usually begin aquifer restoration in each well field as the uranium recovery operations end. Aquifer restoration criteria are determined on a site-specific, well field-by-well field basis. NRC's restoration standards are found in Appendix A to 10 CFR Part 40, and NRC historically has supplemented these regulatory standards through the use of guidance documents and conditions in NRC-issued licenses for ISL facilities. [NRC is currently engaged in a rulemaking that would clarify the requirements for groundwater protection at ISL facilities.]

Aquifer restoration programs typically use a combination of methods including (1) groundwater transfer, (2) groundwater sweep, (3) reverse osmosis with permeate injection, (4) groundwater recirculation, and (5) stabilization monitoring (Energy Information Administration, 1995; Mackin, et al., 2001a; Davis and Curtis, 2007).

2.5.1 Groundwater Transfer

Groundwater transfer involves moving groundwater between the well field entering restoration and another well field where uranium leach operations are beginning, or alternately, within the same well field, if one area is in a more advanced state of restoration than another (NRC, 2006). This technique displaces mining-affected waters in the restoration well field with baseline quality waters from the well field beginning leach operations. As a result, the groundwater in the two well fields becomes blended until the waters are similar in conductivity and therefore similar in the amount of dissolved constituents. Because water is transferred from one well field to another, groundwater transfer typically does not generate liquid effluents.

2.5.2 Groundwater Sweep

During the groundwater sweep phase, contaminated groundwater in the well field is removed by pumping. This pumping causes uncontaminated, native groundwater to flow into the ore body. The groundwater sweep process is depicted in Figure 2.5-1. During groundwater sweep, the licensee pumps water from the well field to the processing plant through all production and injection wells without reinjection. This draws native groundwater inward, flushing the contaminants from areas that have been affected by the horizontal spreading of the lixiviant in the affected zone during uranium recovery. Groundwater produced by the onsite wells will contain uranium and other contaminants released during uranium recovery and residual lixiviant. The initial concentrations of these substances would be similar to those during the uranium recovery operation phase, but would decline gradually with time (Davis and Curtis, 2007). The water removed from the aquifer during the sweep first is passed through the processing plant ion exchange system to recover the uranium and then disposed either in evaporation ponds or via deep well injection in accordance with the limits in the UIC permit.

The duration of the aquifer sweep and volume of water removed depend on the volume of the aquifer affected by the ISL process. The aquifer volume typically is described in terms of "pore volumes" (see text box). Based on operational data (see Section 2.11.5), it is likely that more than one pore volume would be removed during the sweep. At the Crow Butte ISL facility in Dawes County, Nebraska, the pore volumes for the first six well fields {3.8 to 16.3 ha [9.3 to 40.2 acres]} were estimated to range from 58.3 to 298.7 million L [15.4 to 78.9 million gal] (NRC, 1998b). In comparison, the total pore volume for the nine well fields at the Irigaray Project was estimated to be 232.8 million L [61.5 million gal] (Cogema Mining, 2005).

Pore Volume and Flare

Pore volume is a term of convenience used by the *in situ* leach industry to describe the quantity of free water in the pores of a given volume of aquifer material. It provides a unit reference that an operator can use to describe the amount of lixiviant circulation needed to leach an ore body, or describe the unit number of treated water circulations needed to flow through a depleted ore body to achieve restoration. A pore volume provides a way for an operator to use relatively small-scale studies and scale the results to field-level pilot tests or to commercial well field scales. Typically, a "pore volume" is calculated by multiplying the surficial area of a well field (the area covered by injection and recovery wells) by the thickness of the production zone being exploited and the estimated or measured porosity of the aquifer material (NRC, 2003a).

A proportionality factor, known as "flare," is designed to estimate the amount of aquifer water outside of the pore volume that has been impacted by lixiviant flow during the extraction phase. The flare is usually expressed as a horizontal and vertical component to account for differences between the horizontal and vertical hydraulic conductivity of an aquifer material (NRC, 2003a).




Figure 2.5-1. Schematic Diagram of Groundwater Sweep During Aquifer Restoration (after Energy Information Administration, 1995)

2.5.3 Reverse Osmosis, Permeate Injection, and Recirculation

Reverse osmosis and permeate injection are used after groundwater sweep operations. This phase returns total dissolved solids, trace metal concentrations, and aquifer pH to baseline values (Davis and Curtis, 2007; NRC, 2003a). During permeate injection and recirculation, uranium in the groundwater is removed by passing the water through the ion exchange circuit, as during operations. After that, other chemical constituents in the groundwater are removed by passing the groundwater through a reverse osmosis system consisting of pressurized, semi-permeable membranes.

The reverse osmosis process yields two fluids: clean water (permeate: about 70 percent) and water with concentrated ions (brine: about 30 percent). Water sent to the reverse osmosis system must be pre-treated so the semipermeable membranes used in the system are not fouled. The pH is lowered, and additives called antiscalants are added to the groundwater upstream of the reverse osmosis unit to prevent precipitation of minerals (particularly calcium carbonate). Typically, sodium hexametaphosphate or polycarboxylic acid are used as antiscalants and sulfuric acid is used for pH adjustment. After reverse osmosis, sodium hydroxide is added to readjust the pH of the groundwater to baseline levels.

The pumping and injection rates during the recirculation phase are likely to be similar to those during the sweep phase (hundreds of gallons per minute), but many pore volumes (often more than 10) must be circulated to achieve aquifer restoration goals (Davis and Curtis, 2007; Mackin, et al., 2001b). The net withdrawal from the aquifer depends on how the rejected liquid (reject) from the reverse osmosis system, which is about 30 percent of the pumping rate, is handled. Because the reject is a brine solution, it cannot be directly injected into the aquifer or discharged to the environment. The reject can be disposed directly in an evaporation pond or via a deep well injection in accordance with the discharge limits in the UIC permit. If the reject is sent directly to an evaporation pond or a deep disposal well, the net withdrawal from the aquifer could be about 30 percent of the pumping rate (tens of gallons per minute).

Alternatively, a brine concentrator can be used to treat the reject. The brine concentrator heats and evaporates the water, concentrating the brine, which then contains precipitated solids in the form of common salts. The brine concentration process typically results in about one part briny slurry and salts to 300 parts purified water. The purified water can be reintroduced into the aquifer and thus the net withdrawal from the aquifer would be only a small percentage of the recirculation rate. The briny slurry is disposed in an evaporation pond or via deep well injection (Section 2.7.2).

After completing the reverse osmosis/permeate injection phase, the well field water will have characteristics similar to the permeate, and the recirculation phase takes place. To homogenize the groundwater, well field water may be circulated using the original injection and production wells. The quantity of water that is recirculated depends on site-specific baseline parameters and contaminant levels.

2.5.4 Stabilization

The purpose of the stabilization phase of aquifer restoration is to establish a chemical environment that reduces the solubility of dissolved constituents such as uranium, arsenic, and selenium. An important part of stabilization during aquifer restoration is metals reduction (Davis and Curtis, 2007). During uranium recovery, if the oxidized (more soluble) state is allowed to persist after uranium recovery is complete, metals and other constituents such as arsenic, selenium, molybdenum, uranium, and vanadium may continue to leach and will remain at elevated levels. To stabilize metals concentrations, the pre-operational oxidation state in the ore production zone should be reestablished as much as is possible. This is achieved by adding an oxygen scavenger or reducing agent such as hydrogen sulfide (H_2S) or a biodegradable organic compound (such as ethanol) into the uranium production zone during the later stages of recirculation (Davis and Curtis, 2007). The need for an aquifer stabilization phase depends on how effectively the sweep and recirculation phases restore the affected aquifer to background water quality. The total volume and rate of net groundwater recovery during the stabilization phase will be similar to that during the restoration recirculation phase.

Following stabilization, the licensee monitors the groundwater by quarterly sampling to ensure that baseline or pre-operational class-of-use conditions have been permanently restored and that any adjacent nonexempt aquifers are unaffected. The licensee would reinitiate aquifer restoration if stabilization monitoring determines it is necessary. Both the state permitting agency and the NRC must review and approve the monitoring results before aquifer restoration is considered to be complete.

2.6 Decontamination, Decommissioning, and Reclamation

Decommissioning an ISL facility is based on an NRC-approved decommissioning plan. This section discusses activities based on previous summaries (Energy Information Administration, 1995; Mackin, et al., 2001a). The primary steps involved in decommissioning an ISL facility include

- Conducting radiological surveys of facilities, process equipment, and materials to evaluate the potential for exposure during decommissioning
- Removing contaminated equipment and materials for disposal at an approved facility or for reuse
- Decontaminating items to be released for unrestricted use
- Cleaning up areas used for contaminated equipment and materials
- Cleaning up evaporation ponds
- Plugging and abandoning wells
- Surveying excavated areas for contamination and removing contamination to meet cleanup limits
- Backfilling and recontouring disturbed areas
- Performing final site soil radiation background surveys
- Revegetating and reclaiming disturbed areas
- Monitoring the environment

Process buildings and equipment are surveyed to identify any radiation hazards. Alternatives for handling process buildings and equipment include reuse, removal, or disposal. Contaminated items are decontaminated if they are to be released for offsite unrestricted use; otherwise, they are disposed of as 11e.(2) byproduct material in a licensed disposal facility. Estimated volumes of building demolition and removed equipment wastes for an ISL facility are provided in Table 2.6-1.

Pond liners and leak detection systems are surveyed. If radiological contamination is found, the liners and detection systems are typically removed and disposed in a licensed disposal facility. Estimated volumes of pond reclamation wastes for an ISL facility are provided in Table 2.6-1.

Well fields are decommissioned after groundwater restoration has been completed. Proper well field decommissioning protects the groundwater supply and eliminates physical hazards. First, surface equipment (such as injection and production lines), electrical components, and well head equipment (such as valves, meters, or fixtures) are salvaged. Then buried piping is removed, and the wells are plugged and abandoned using accepted practices identified as part

Table 2.6-1. Estimated Decommissioning and Reclamation Waste Volumes (yd³)* for Offsite Disposal, Smith Ranch *In-Situ* Leach Facility†

ISL Decommissioning Activity	Byproduct Radioactive Waste	Other Solid Waste
Processing Equipment Removal	342	0
Building Demolition	546	531
Well Field Equipment	1,361	404
Trunk Line Removal	2,263	0
Contaminated Soil Removed	1,428	0
Evaporation Pond Reclamation	68	0

*To convert yd³ to m³, multiply by 0.7646.

†Volumes were compiled and summed from an annual surety report. McCarthy, J. "Smith Ranch: 2007–2008 Surety Estimate Revision." Letter (June 29) to G. Janosko, NRC. Glenrock, Wyoming: Power Resources International. 2007.

of the EPA- or state-administered UIC program. Based on past experience, about 90 percent of the materials will be suitable for unrestricted release or disposal at an unrestricted area landfill. Estimated volumes of well field decommissioning wastes for an ISL facility are provided in Table 2.6-1. The well field area is decontaminated in accordance with NRC regulatory limits at 10 CFR Part 40, Appendix A, and surveys are performed to ensure compliance with standards. Surface reclamation is completed using an NRC-approved plan.

Contaminated soils are cleaned up as necessary for decommissioning. A gamma radiation survey is conducted to determine whether any contaminated areas exist. Criteria at 10 CFR Part 40, Appendix A, are used for identifying contaminated soils and for determining when cleanup is complete. The NRC reviews and approves survey and sampling results. In the well fields where gamma radiation surveys correlate strongly with actual radiation concentrations in soil, gamma surveys are conducted as each well field unit is decommissioned. Soil samples are obtained from any areas that have elevated gamma readings. Areas contaminated with Ra-226, Ra-228, or other radionuclides exceeding the limits specified at 10 CFR Part 40, Appendix A, Criterion 6-(6), are cleaned up. Contaminated soil is removed and disposed as 11e.(2) byproduct material at a licensed disposal facility. The estimated volume of contaminated soil removal for an ISL facility is provided in Table 2.6-1. The most likely areas for contaminated soils are well field surfaces, evaporation pond bottoms and berms, process building areas, storage yards, transportation routes for uranium recovery products or contaminated materials, and pipeline runs. Areas used for land application of treated water are also surveyed and decontaminated as necessary.

All radioactive wastes generated during ISL facility decommissioning (as well as radioactive wastes generated during construction, operation, and aquifer restoration) are considered 11e.(2) byproduct material that must be disposed at a licensed facility (Section 2.7).

An NRC-approved surface reclamation plan ensures disturbed lands are returned to production or to planned post-operational land use. Baseline data on soils, vegetation, wildlife, and radiation are used as guidelines for the surface reclamation. Areas disturbed by the uranium recovery operations are restored as closely as possible to pre-operational conditions. Reclamation activities include replacing excavated soils, recontouring affected areas, reestablishing original drainage, and revegetation. The magnitude of reclamation activities vary, in part, with the size of the ISL facility. A large ISL facility, Smith Ranch (see Table 2.11-1) has estimated applying approximately 43,748 m³ [57,221 yd³] of topsoil to the ground surface during

site reclamation (McCarthy, 2007). Because topsoil excavated during construction was stockpiled and reseeded to limit erosion (NRC, 1992), the net amount of topsoil needed to replace topsoil removed during decommissioning is approximated by the estimated volume of excavated soil destined for offsite disposal shown in Table 2.6-1 (1,092 m³ [1,428 yd³]). After reclamation is complete, lands are normally capable of supporting wildlife and land uses such as livestock grazing.

A financial surety (Section 2.10), established when an NRC license is granted, provides assurance that the costs of aquifer restoration and site decommissioning are covered when facility operations end. The surety also covers costs to close the site at any point during operations.

2.7 Effluents and Waste Management

ISL facilities generate airborne effluents, liquid wastes, and solid wastes that must be handled and disposed of properly. Effluents, waste streams, and waste management practices applicable to ISL facilities are described in this section.

2.7.1 Gaseous or Airborne Particulate Emissions

During construction, operations, aquifer restoration, and decommissioning, ISL facilities can produce airborne emissions including

- Fugitive dusts
- Combustion engine exhausts
- Radon gas emissions from lixiviant circulation and evaporation ponds
- Uranium particulate emissions from yellowcake drying

Fugitive dusts and engine exhausts are generated primarily during construction, transportation, and decommissioning activities. The fugitive dust is generated by travel on unpaved roads and from disturbed land associated with the construction of well fields, roads, and support facilities. Vehicles workers use to commute to the facility, to support onsite activities, or to transport supplies to the site emit fuel combustion products. Diesel emissions originate from drill rigs, diesel-powered water trucks, and other equipment used during the construction phase. Table 2.7-1 provides information from a previously licensed ISL satellite facility on the nature and duration of nonradiological emission-generating activities during construction, operation, and decommissioning. Table 2.7-2 contains the annual total releases and average air concentrations of particulate (fugitive dust) and gaseous (diesel combustion products) emissions estimated for the construction phase of the ISL facility near Crownpoint, New Mexico.

Table 2.7-1. Combustion Engine Exhaust Sources for the Gas Hills In-Situ Leach Satellite Facility During Construction, Operations, Reclamation, and Decommissioning*

Period	Activity	Equipment Type	Number of Units	Frequency of Operation	Duration of Operation
Construction	Initial Construction/ Well Field Road Construction	Scraper	1	8 hr/day, 5 day/wk	2 months
		Bulldozer	1	"	"
		Motor Grader	1	"	"
	Well Preparation	Truck Mount Rotary Drill Rig, Diesel Truck	4-8	8 hr/day, 5 day/wk	12 mo/yr
		Pump Pulling Vehicle 1-ton gas or diesel	2	"	"
		Motor Grader	1	"	3 mo/yr
		Backhoe	3	"	12 mo/yr
		Forklift	2	"	"
		Cementer (gas)	4	"	"
		Light Duty Truck	8-10	8 hr/day, 7 day/wk	"
	Construction Material Transport	Heavy Duty Water Truck (1,500 gal)	4-8	"	"
		Heavy Duty Diesel Truck	1	1 trip/day	2 mo/yr
	Commuting	Light Duty Vehicles	30	"	6 mo/yr
Operation	Satellite Facility	Gas or Propane Heater	6	24 hr/day	6 mo/yr
	Product Transport	Truck to Highland Site Diesel Semi with Trailer	2	1 trip/day	12 mo/yr
	Commuting	Light Duty Vehicles	30	"	"
Decommissioning	Reclamation	Scraper	1	2 × 8 hr shift/day*	2-3 yr
		Motor Grader	1	"	"
		Backhoe	2	"	"
		Heavy Duty Truck (Diesel)	3	"	"
		Light Duty Truck	15	"	"
		Light Duty Vehicles	20	1 trip/day	"

*NRC. "Environmental Assessment for the Operation of the Gas Hills Project Satellite In-Situ Leach Uranium Recovery Facility." Docket No. 40-8857. Washington, DC: NRC. January 2004.

Table 2.7-2. Estimated Particulate (Fugitive Dust) and Gaseous (Diesel Combustion Products) Emissions for the Crownpoint, New Mexico, In-Situ Leach Facility Construction Phase*

Emission Type	Annual Total (metric tons)†	Annual Average Concentration ($\mu\text{g}/\text{m}^3$)‡
Particulates	10.0	0.28
Sulfur dioxides (SO_x)	6.4	0.18
Nitrous oxides (NO_x)	76.2	2.1
Hydrocarbons	9.8	0.27
Carbon monoxide	63.7	1.8
Aldehyde	1.4	0.04

*Modified from U.S. Nuclear Regulatory Commission. NUREG-1508, "Final Environmental Impact Statement To Construct and Operate the Crownpoint Uranium Solution Mining Project, Crownpoint, New Mexico." Washington, DC: U.S. Nuclear Regulatory Commission. February 1997.

†Multiply metric ton value by 1.1023 to convert units to short ton.

‡Multiply $\mu\text{g}/\text{m}^3$ value by 2.74×10^{-8} to convert units to oz/yd^3 .

Radon gas is released during operation and aquifer restoration. Pressurized processing systems may contain most of the radon in solution; however, radon may escape from the processing circuit in the central uranium processing facility through vents or leaks, during well field operations, or during resin transfer when remote ion exchange is used. For open air activities, the gas quickly disperses into the air. In closed processing areas, the building ventilation systems are designed to limit indoor radon concentrations. Radon detectors are placed in appropriate locations to ensure compliance with worker protection regulations in 10 CFR Part 20. Airborne particulate emissions from yellowcake drying and packaging and the filling of sodium bicarbonate storage containers are controlled by using vacuum drying equipment and baghouse dust collection systems.

Both radon releases and uranium particulate emissions can migrate downwind from processing facilities and well fields. Downwind radiation dose from such ISL facility emissions varies due to the effects of dispersion as a function of distance. Particulate emissions are further reduced by the effect of dry deposition during airborne transport. Calculations of downwind dose are based on estimating the relative air concentration of released radionuclides (which is proportional to dose). Figure 2.7-1 shows relative air concentration for particulate matter as a function of distance estimated for the Bison Basin ISL facility (NRC, 1981, Table D.3). These results apply to the downwind area with the highest relative air concentrations. As shown, relative air concentration of uranium particulates, and therefore dose, drops by about a factor of 10 from the first data point {500 m [1,640 ft]} to the second {1,500 m [4,920 ft]}. The reduction in relative air concentration, and therefore dose, becomes less significant as downwind distance increases. The effect of distance on air concentration estimates is less pronounced for transport of gases (e.g., radon) due to the absence of dry deposition, which does not apply to gaseous transport. Airborne transport and dose modeling results for ISL facility releases to air (including both radon and uranium particulate releases, where applicable) are provided in Sections 4.2.11.2, 4.3.11.2, 4.4.11.2, and 4.5.11.2.




Figure 2.7-1. Downwind Distance Versus Relative Air Concentration (Which Is Proportional to Dose) [Bison Basin ISL Facility (NRC, 1981, Table D.3)]

2.7.2 Liquid Wastes

Liquid wastes from ISL facilities are generated during all phases of uranium recovery; construction, operations, aquifer restoration, and decommissioning. Liquid wastes may contain elevated concentrations of radioactive and chemical constituents. Table 2.7-3 shows estimated flow rates and constituents in liquid waste streams for the Highland ISL facility (NRC, 1978). Liquid waste streams are predominantly production bleed (1 to 3 percent of the process flow rate) and aquifer restoration water (NRC, 1997a). Additional liquid waste streams are generated from well development, flushing of depleted eluant to limit impurities, resin transfer wash, filter washing, uranium precipitation process wastes (brine), and plant wash down water. ISL facilities have concrete curbed floors with drains and a sump to control and retain water from spills and wash downs. Sumps direct water to treatment facilities, to evaporation ponds, or back to the process circuit. Chemical tanks have berms that can hold tank contents if tanks rupture.

Table 2.7-3. Estimated Flow Rates and Constituents in Liquid Waste Streams for the Highland In-Situ Leach Facility*

	Water Softener Brine	Resin Rinse	Elution Bleed	Yellowcake Wash Water	Restoration Wastes
Flow Rate, gal/min	1	<3	3	7	450
As, ppm					0.1–0.3
Ca, ppm	3,000–5,000				
Cl, ppm	15,000–20,000	10,000–15,000	12,000–15,000	4,000–6,000	
CO ₃ , ppm		500–800			300–600
HCO ₃ , ppm		600–900			400–700
Mg, ppm	1,000–2,000				
Na, ppm	10,000–15,000	6,000–11,000	6,000–8,000	3,000–,000	380–720
NH ₄ , ppm			640–180		
Se, ppm					0.05–0.15
Ra-226, pCi/L	<5	100–00	100–300	20–50	50–100
SO ₄ , ppm					100–200
Th-230, pCi/L	<5	50–100	10–30	10–20	50–150
U, ppm	<1	1–3	5–10	3–5	<1
Gross Alpha, pCi/L					2,000–3,000
Gross Beta, pCi/L					2,500–3,500

*NRC. NUREG-0489, "Final Environmental Statement Related to Operation of Highland Uranium Solution Mining Project, Exxon Minerals Company, USA." Washington, DC: NRC. November 1978.

Some liquid wastes are treated at the processing facility to remove or reduce contaminants prior to disposal. Reverse osmosis is commonly used to segregate contaminants from liquid waste streams (e.g., Section 2.5.3). Radium concentrations are also selectively reduced when water is treated with barium chloride. The barium chloride chemically binds to radium in solution and deposits as a sludge that is sent to a licensed disposal facility. Results from Hydro Resources, Inc. reported in NRC (1997a) show radium concentrations of 74 pCi/l were reduced to less than 1 pCi/L following treatment with barium chloride.

Byproduct Material

11e.(2) byproduct materials are tailings or waste generated by extraction or concentration of uranium or thorium processed ores, as defined under Section 11e.(2) of the Atomic Energy Act.

Liquid effluent disposal practices that NRC previously has approved for use at specific sites include evaporation ponds, land application, deep well injection, and surface water discharge.

Evaporation ponds are used to retain the process-related liquid effluents that cannot be discharged directly to the environment. These effluents are 11e.(2) byproduct material. The residual solid waste materials normally remain in ponds until the ponds are decommissioned, when sludges are disposed of as 11e(2) material at a licensed disposal facility (Section 2.6). Guidance for the construction, operation, and monitoring of evaporation ponds is found in NRC

Regulatory Guide 3.11 (NRC, 1977, 2008). Typical evaporation ponds have surface areas ranging from 0.04 to 2.5 ha [0.1 to 6.2 acres] (NRC, 1998a; Crow Butte Resources, 2007). Evaporation ponds at NRC-licensed ISL facilities are designed with leak detection systems to detect liner failures. The licensee also must maintain sufficient reserve capacity in the retention pond system so that the contents of a pond can be transferred to other ponds in the event of a leak and subsequent corrective action and liner repair. Licensee and applicants can minimize the likelihood of impoundment failure by designing the pond embankments in accordance with the criteria found in NRC Regulatory Guide 3.11 (NRC, 1977, 2008). Sufficient freeboard height above the liquid level ensures containment during wind and rain events.

Land application uses agricultural irrigation equipment to apply treated water to land where the water can evaporate directly or be transpired by plants. Uranium and radium levels are reduced in the effluents disposed of by land application so as to limit contamination of surface soils and plants. Areas of a site where land application of treated water has been used are included in decommissioning surveys to ensure soil concentration limits are not exceeded. Land application may also require approval and permitting by other applicable State agencies.

NRC staff may also review and approve deep well injection for a specific ISL site as a method to dispose of particular process fluids such as reverse osmosis brine. [EPA or the state give the final approval, though, for the use of this method of waste disposal.] Deep well injection involves pumping the waste fluids into a deep confined aquifer at depths typically greater than 1,524 m [5,000 ft] below the ground surface (NRC, 1997a). Aquifer water quality in the deep confined aquifer is often poor (e.g., high salinity or total dissolved solids) and below drinking water standards. The approval process verifies that site-specific and regional characteristics limit the potential for contamination of local drinking water sources. Licensees must obtain an UIC permit from EPA or the appropriate state agency (Section 1.7).

The National Pollutant Discharge Elimination System (NPDES) permitting process (Section 1.8) allow for surface discharge of treated liquid effluents to local waterways including ephemeral stream channels. Water discharged in this way must be treated to remove contaminants to meet state and federal water quality standards.

2.7.3 Solid Wastes

All phases of the ISL facilities lifecycle generate solid wastes. These wastes include spent resin, empty chemical containers, pipes and fittings, pond sludge, tank sediments, contaminated soil from leaks and spills, and municipal waste. Solid wastes are classified as radioactive or nonradioactive prior to disposition. Radioactive wastes are disposed of as 11e(2) byproduct material at a licensed facility. Contaminated equipment and buildings may be similarly disposed or decontaminated and released according to NRC requirements. Nonradioactive hazardous wastes are segregated and disposed of at a hazardous waste disposal facility. Nonradiological uncontaminated wastes are disposed of as ordinary solid waste at a municipal solid waste facility. The largest volumes of solid wastes requiring disposal are generated during facility decommissioning (EPA, 2007a,b). Table 2.6-1 provides estimated volumes of radioactive and noncontaminated ISL facility decommissioning wastes designated for offsite disposal.

2.8 Transportation

Trucks transport construction equipment and materials, operational processing supplies, ion exchange resins, yellowcake product, and waste materials during all phases of the ISL facility lifecycle.

Trucks transport construction equipment and materials to the site to support facility and well field construction activities along local roads. Because ISL facilities are small magnitude construction projects and well field construction is phased over a period of years, the magnitude of trucking activity to support construction is small relative to other industrial activities. The estimated frequency of truck shipments for construction of an ISL facility is provided in Table 2.8-1.

During the operational period, trucks supply an ISL facility with materials needed to support processing operations. Shipments involve hazardous chemicals such as ammonia, sulfuric acid, liquid and gaseous oxygen, hydrogen peroxide, sodium hydroxide, barium chloride, carbon dioxide, hydrochloric acid, sodium carbonate, sodium chloride, hydrogen sulfide, and sodium sulfide. These chemicals are commonly used in a variety of industrial applications, and the U.S. Department of Transportation regulates their transport. The estimated frequency of truck shipments to support ISL facility operation is provided in Table 2.8-1.

In areas where ore deposits are smaller and more spread out, a producer may construct a series of small satellite plants at the well field where ion exchange processing is conducted remotely rather than at the central uranium processing facility (NRC, 2004, 2006). The products of ion exchange processing are then transported by truck to a central uranium processing facility (Section 2.4). Uranium production using these types of satellite facilities is sometimes known as satellite remote ion exchange (Finch, 2007). Facilities that incorporate remote ion exchange operations will transport loaded ion exchange resins or uranium slurry from well fields to centralized processing facilities by truck. These trucks are typically modified three-compartment cement trailers. The carbon steel compartments are pressurized and rubber lined. The first compartment carries the uranium-loaded resin, the second is empty, and the third compartment holds unloaded resins (Finch, 2007). Each shipment can contain about 900–1,350 kg [2,000–3,000 lb] of uranium-loaded resin, although the actual amount depends on the size of the trailer. These trucks are generally sole-use vehicles that are labeled for this purpose in accordance with U.S. Department of Transportation requirements at 49 CFR 171–189 and NRC regulations at 10 CFR Part 71. In accordance with these regulations, no liquids are permitted in the truck during transport of uranium resins. The estimated frequency of remote ion exchange truck shipments to support ISL facility operation is provided in Table 2.8-1.

The refined yellowcake product is packed in 208-L [55-gal], 18-gauge drums holding an average of 430 kg [950 lb] and classified by the U.S. Department of Transportation as Type A packaging (49 CFR Parts 171–189 and 10 CFR Part 71). The yellowcake is shipped by truck to a remote conversion plant that transforms the yellowcake to uranium hexafluoride (UF_6) for the enrichment step of the reactor fuel cycle. An average truck shipment contains approximately 40 drums or 17 metric tons [19 short tons] of yellowcake (NRC, 1980). The annual number of shipments from a given ISL facility depends on the yellowcake production rate of the facility. A range of estimated annual shipment totals based on prior ISL facility production limits is provided in Table 2.8-1.

1

Table 2.8-1. Estimated Annual Vehicle Trips for Phases of ISL Facility Lifecycle		
Cargo	Estimated Number of Truck Shipments	Remarks
Construction Equipment/Supplies	62*	1 per day for 2 months
Remote IX Shipments	365*	1 per day annually
Processing Chemicals	272†	Less than 1 per day annually
Processing Wastes	Range: 2.5–15*	Less than 1 per month annually
Yellowcake	Range: 21–145‡§¶¶#	Maximum is based on production assumed at the permitted limit at the largest facility
Decommissioning Nonhazardous Solid Waste	44**	Based on waste volumes from Smith Ranch (Table 2.6-1) and truck volume of 20 yd ³ /shipment
Decommissioning Byproduct Waste	100**	Based on waste volumes from Smith Ranch (Table 2.6-1) and truck volume of 20 yd ³ /shipment
Decommissioning Hazardous Waste	To be determined	To be determined
Employee Commuting	5,200–52,000 trips*	20 to 200 employees per day assumed for 12 months/yr. Maximum in range is expected to depend on timing of construction, drilling, and operational activities (Section 2.11.6)
<p>*NRC. "Environmental Assessment for the Operation of the Gas Hills Project Satellite <i>In-Situ</i> Leach Uranium Recovery Facility." Docket No. 40-8857. Washington, DC: NRC. January 2004.</p> <p>†NRC. "Environmental Assessment for Renewal of Source Material License No. SUA-1534—Crow Butte Resources Inc., Crow Butte Uranium Project Dawes County, Nebraska." Docket No. 40-8943. Washington, DC: NRC. 1998.</p> <p>‡NRC. NUREG-0489, "Final Environmental Statement Related to Operation of Highland Uranium Solution Mining Project, Exxon Minerals Company, USA." Washington DC: NRC. November 1978.</p> <p>§NRC. "Final Environmental Statement Related to the Operation of Bison Basin Project." Docket No. 40-8745. Washington, DC: NRC. 1981.</p> <p>¶NRC. NUREG-1508, "Final Environmental Impact Statement To Construct and Operate the Crownpoint Uranium Solution Mining Project, Crownpoint, New Mexico." Washington, DC: NRC. February 1997.</p> <p>¶¶NRC. "Environmental Assessment for Renewal of Source Material License No. SUA-1534—Crow Butte Resources Inc., Crow Butte Uranium Project Dawes County, Nebraska." Docket No. 40-8943. Washington, DC: NRC. 1998.</p> <p>#NRC. "Environmental Assessment Construction and Operation of In Situ Leach Satellite SR-2 Amendment No. 12 to Source Material License No. SUA-1548—Power Resources, Inc., Smith Ranch-Highland Uranium Project (SR-HUP) Converse County, Wyoming." Docket No. 40-8964. Washington DC: NRC. December 2007.</p> <p>**Waste volumes compiled and summed from estimates reported in McCarthy, J. "Smith Ranch: 2007–2008 Surety Estimate Revision." Letter (June 29) to G. Janosko, NRC. Glenrock, Wyoming: Power Resources International. 2007.</p>		

2

3

4

5

6

7

Waste materials generated by construction, operation, aquifer restoration, and decommissioning activities including hazardous chemical, radioactive, and ordinary municipal waste streams are segregated by waste type and transported by truck to approved disposal facilities. The estimated frequency of waste shipments for operation and decommissioning an ISL facility is

provided in Table 2.8-1. Section 2.7 provides additional information on waste streams and waste management activities.

2.9 Radiological Health and Safety

NRC regulations at 10 CFR Part 20 address the health and safety of workers and the public in the event of exposure to radiation from all phases of the ISL facility lifecycle. These regulations require ISL facility operators to develop and implement an NRC-approved radiation protection program. During NRC inspections and other oversight activities, including reviews of monitoring and incident reports, NRC checks compliance with this program. This section briefly summarizes basic elements of a 10 CFR Part 20 radiation protection program. More detailed descriptions of radiological safety requirements and programs are found in the regulations at 10 CFR Part 20 and applicable NRC guidance documents summarized in the NRC Standard Review Plan for ISL facilities (NRC, 2003a).

A radiological protection program includes plans and procedures addressing the following topics:

- **Effluent Control.** Effluents to air (e.g., radon, uranium particulates) and surface water (e.g., permitted wastewater discharges) must meet NRC limits in 10 CFR Part 20 for radioactive effluents and worker and public doses. To ensure proper performance to specifications, plans and procedures include minimum performance specifications for control technologies (e.g., yellowcake dryer emission controls) and frequencies of tests and inspections.
- **External Radiation Exposure Monitoring Program.** This program specifies survey methods (including monitoring locations), instrumentation, and equipment for measuring worker exposures to external radiation during routine and nonroutine operations, maintenance, and cleanup activities. The program is designed to ensure worker dose levels are as low as reasonably achievable and comply with NRC requirements in 10 CFR Part 20.
- **Airborne Radiation Monitoring Program.** This program determines concentrations of airborne radioactive materials (including radon) in the workplace during routine and nonroutine operations, maintenance, and cleanup. This program is designed to ensure airborne radiation releases and worker exposures are as low as reasonably achievable and meet requirements specified in 10 CFR Part 20.
- **Exposure Calculations.** Procedures document the methodologies used to calculate intake of airborne radioactive materials in the workplace during routine and nonroutine operations, maintenance, and cleanup activities.
- **Bioassay Program.** A bioassay program assesses biological intake of uranium by workers routinely involved in operations where radioactive material can be inhaled (e.g., yellowcake dust from dryer operations or baghouse maintenance). Programs include collection and analysis of urine samples that are assessed for the presence of uranium. Action levels are set to maintain exposures as low as reasonably achievable and within worker requirements in 10 CFR Part 20.

- 1 • **Contamination Control Program.** A contamination control program includes standard
2 operating procedures to prevent employees from entering clean areas or leaving the site
3 while contaminated with radioactive materials. Such programs involve radiation
4 surveys of personnel and surfaces, housekeeping requirements, specifications to
5 control contamination in processing areas, and controls for the release of
6 contaminated equipment.
7
- 8 • **Airborne Effluent and Environmental Monitoring Program.** This program measures
9 concentrations and quantities of radioactive and nonradioactive materials released to the
10 environment surrounding the facility. Such programs measure concentrations of
11 constituents in stack effluents at the facility and in the environment near and beyond the
12 site boundary emphasizing surface water, groundwater, vegetation, food and fish, and
13 soil and sediment. Direct radiation and radon flux are also measured. Offsite
14 radiological and environmental monitoring is detailed in Chapter 8.
15

16 **2.10 Financial Surety**

17
18 NRC regulations [10 CFR Part 40, Appendix A, Criterion (9)] require that applicants or licensees
19 cover the costs for a third party to conduct decommissioning, reclamation of disturbed areas,
20 waste disposal, and groundwater restoration (Mackin, et al., 2001b). NRC annually reviews a
21 licensee's financial surety to assess expansions in operations, changes in engineering design,
22 completion of decommissioning activities, actual experience in aquifer restoration, and inflation.
23 Specific considerations for estimating these costs are detailed in Appendix C of NRC, 2003a,
24 and financial surety arrangements are discussed only briefly here.
25

26 Each licensee establishes financial surety arrangements before uranium recovery operations
27 begin to assure there will be sufficient funds to carry out the activities described in Sections 2.5
28 and 2.6. The surety funds also must be sufficient for monitoring and control required as part of
29 the license termination. Acceptable financial surety arrangements include surety bonds, cash
30 deposits, certificates of deposit, deposits of government securities, parent company guarantees
31 (subject to specific NRC criteria), trusts and standby trusts, irrevocable letters or lines of credit,
32 and combinations of these instruments. Self-insurance is not an acceptable form of surety for
33 NRC, although it may be accepted by individual states. The term of the surety mechanism must
34 be open ended so that it will not expire before cleanup is complete. [NRC is currently engaged
35 in a rulemaking that may change the list of NRC-approved surety instruments and conditions for
36 other approved forms of financial assurance. The final rule may be issued in late 2008 or early
37 2009.]
38

39 As required under 10 CFR Part 40, Appendix A, Criterion 9, the licensee must supply
40 enough information for NRC to verify that the amount of financial coverage will allow all
41 decontamination and decommissioning and reclamation of sites, structures, and equipment
42 used in conjunction with facility operation to be completed. Cost estimates for the following
43 activities (where applicable) should be submitted to NRC with the initial license application or
44 reclamation plan and should be updated annually as specified in the operator's NRC license.
45 A third party (an independent contractor or operator who is not financially affiliated with the
46 licensee) must calculate cost estimates based on completion of all activities. Unit costs,
47 calculations, references, assumptions, equipment and operator efficiencies, and other
48 breakdown details must be provided.
49

In the required annual surety estimate, the licensee must provide estimated costs for all decommissioning, reclamation, and groundwater restoration work remaining to be performed at the site—not simply deduct the cost of work already performed from the previous surety estimate (see NRC, 1997b). For each activity, estimates should include costs for equipment; materials; labor and overhead; licenses, permits, and miscellaneous site-specific costs; and any other activity or resource that will require spending funds. The licensee should add a contingency amount to the total cost estimate for the final site closure. NRC typically considers a 15 percent contingency to be an acceptable minimum amount (NRC, 2003a, Appendix C). The licensee is required by 10 CFR Part 40, Appendix A, Criterion 9, to adjust cost estimates annually to account for inflation and changes in reclamation plans. In addition, all costs are to be estimated based on third party, independent contractor costs (including overhead and profit in unit costs or as a percentage of the total). Licensee-owned equipment and the availability of licensee staff should not be considered in the financial surety estimate, because this can reduce cost calculations.

To avoid unnecessary duplication and expense, NRC also takes into account surety arrangements that other federal, state, or other local agencies may require. However, NRC is not required to accept such sureties if they are insufficient. NRC reviews the licensee's surety analysis annually to ensure that the funding reflects ongoing aquifer restoration and decommissioning/reclamation activities. The surety remains in place until the final NRC decommissioning surveys are complete and the license is terminated.

2.11 Information From Historical Operation of ISL Uranium Milling Facilities

2.11.1 Area of ISL Uranium Milling Facilities

The permitted areas for past and current ISL uranium recovery operations have varied in size. As shown in Table 2.11-1 facilities range from about 1,034 ha [2,552 acres] for the proposed Crownpoint facility in McKinley County, New Mexico, to over 6,480 ha [16,000 acres] for the Smith Ranch property in Converse County, Wyoming. However, much of the permitted area of a site is undisturbed, and surface operations (wells, processing facilities) affect only a small portion of it. For example, the well fields and excursion monitoring wells that go along with them occupy between 40 and 2,500 ha [100 and 6,000 acres], although most occupy less than about 1,000 ha [2,500 acres]. The central processing facility may occupy only 1 to 6 ha [2.5 to 15 acres], and satellite plants would be even smaller (NRC, 2006).

Surface facilities are considered controlled areas where security fencing limits access. The well fields, which consist of injection and recovery (production) wells, are the areas where most activities that disturb the surface and subsurface take place. Select areas around header houses and well heads are fenced to prevent livestock grazing. Lands near surface operations and in active uranium recovery are excluded from agricultural production for the duration of the project. Despite the large permitted area of a typical ISL facility, the amount of land that is disturbed by earthmoving activities at any one time is relatively small. For example, while the total area disturbed by construction activities between 1987 and 2007 is about 530 ha [1,310 acres] for the Crow Butte ISL facility in Dawes County, Nebraska, only about 50 ha [120 acres] is estimated to be the total disturbed area at any one time (Crow Butte Resources, Inc., 2007). After the surface operations are complete and well fields are restored, the final steps of decommissioning and surface reclamation are intended to return the land to its pre-operational conditions.

1

Table 2.11-1. Size of Permitted Areas for ISL Facilities		
Name	Permitted Area in Hectares [acres]	Status of Facility as of February 2008
Crownpoint, New Mexico	1,034 [2,552]†	Partially permitted and licensed
Crow Butte, Nebraska	1134 [2,800] ‡	Operating
Gas Hills, Wyoming (Satellite)	3,442 [8,500]*	Under development as a satellite of Smith Ranch/Highland, intend to expand
Reynolds Ranch, Wyoming (Satellite)	3,525 [8,704]§x	Under development as satellite of Smith Ranch/Highland
Highland, Wyoming	6,075 [15,000] ‡	Operating, combined with Smith Ranch
Irigaray, Christensen Ranch	6,075 [15,000]¶	Previously issued license, intend to restart
Smith Ranch, Wyoming	6,480 [16,000]#	Operating, combined with Highland, Gas Hills, North Butte, and Ruth, intend to expand
<p>*NRC. NUREG-1508, "Final Environmental Impact Statement To Construct and Operate the Crownpoint Uranium Solution Mining Project, Crownpoint, New Mexico." Washington, DC: NRC. February 1997.</p> <p>†NRC. "Environmental Assessment for Renewal of Source Material License No. SUA-1534—Crow Butte Resources Inc., Crow Butte Uranium Project Dawes County, Nebraska." Docket No. 40-8943. Washington, DC: NRC. 1998.</p> <p>‡NRC. "Environmental Assessment for the Operation of the Gas Hills Project Satellite <i>In-Situ</i> Leach Uranium Recovery Facility." Docket No. 40-8857. Washington, DC: NRC. January 2004.</p> <p>§NRC. "Environmental Assessment for the Addition of the Reynolds Ranch Mining Area to Power Resources Inc., Smith Ranch/Highlands Uranium Project Converse County Wyoming, Source Material License No SUA-1548." Docket No. 40-8964. Washington, DC: NRC. November 2006.</p> <p>¶NRC. "Environmental Assessment for Renewal of Source Material License No. SUA-1511 Power Resources Inc., Highland Uranium Project Converse County, Wyoming." Docket No. 40-8857. Washington DC: NRC. August 18, 1995.</p> <p>¶NRC. "Environmental Assessment for Renewal of Source Material License No. SUA-1341, Cogema Mining, Inc. Irigaray and Christensen Ranch Projects, Campbell and Johnson Counties, Wyoming." Docket No. 40-8502. Washington, DC: NRC. June 1998.</p> <p>#NRC. "Environmental Assessment for Rio Algom Mining Corporation Smith Ranch <i>In-Situ</i> Leach Mining Project, Converse County, Wyoming in Consideration of a Source and Byproduct Material License Application." Docket No. 40-8964. Washington, DC: NRC. January 1992.</p>		

2.11.2 Spills and Leaks

During ISL operations and aquifer restoration, barren and pregnant uranium-bearing process solutions are moved through pipelines to and from the well field and among different surface facilities (e.g., processing circuit, evaporation ponds). If a pipeline ruptures or fails, process solutions can be released and (1) pond on the surface, (2) run off into surface water bodies, (3) infiltrate and adsorb in overlying soil or rock, or (4) infiltrate and percolate to groundwater. For example, from 2001 to 2005, the operators of the Smith Ranch-Highland uranium ISL facility

in Converse County, Wyoming, reported 24 spills of uranium recovery solutions, and the WDEQ identified more than 80 spills during commercial operations (WDEQ, 2008). This is the largest NRC-licensed ISL uranium recovery facility. The size of the spills at Smith Ranch-Highland has ranged from a 190- to 380-liter [50- to 100-gallon] spill in February 2004 to a 751,400-L [198,500-gal] spill of injection fluid in June 2007 (WDEQ, 2007; NRC, 2006). The spills most commonly involved injection fluids {0.5 to 3.0 mg/l uranium [0.5 to 3.0 parts per million]}, although spills of production fluids {10.0 to 152 mg/l uranium [10.0 to 152 parts per million]} also have occurred (NRC, 2007). These spills have been predominantly caused by the failure of joints, flanges, and unions of pipelines and at wellheads (NRC, 2006, 2007). The large June 2007 spill at Highland was the apparent result of a failed fitting. The spilled fluids flowed into a drainage and continued downstream for about 700 m [2,300 ft]. The WDEQ Land Quality Division estimated the affected area at 0.44 ha [1.08 acres] (WDEQ, 2007).

Reporting requirements for spills differ from State to State. NRC's requirements for spill reporting are found in Subpart M of 10 CFR Part 20 and at 10 CFR 40.60. Additionally, NRC may incorporate reporting requirements as conditions in the issued operating license. Generally, such NRC and State requirements include a more immediate report (e.g., notifications within 24 to 48 hours of the spill) followed by a later written report addressing items such as, the conditions leading to the spill, the corrective actions taken, and the results achieved. A licensee's documentation of its spills helps in final site decommissioning activities.

For hazardous chemicals stored at the processing facility, spill responses would be similar to those described previously for yellowcake transportation, although nonradiological material spills are primarily reportable to the appropriate state agency and EPA. Concrete berms with at least the volume of the tank are used to contain spills from process chemical storage tanks and simplify cleanup (e.g., NRC, 1998a,b). The Occupational Safety and Health Administration sets worker exposure limits to process chemicals at the ISL surface facilities. Typical onsite quantities of process chemicals used at ISL facilities are included in Tables 2.11-2 and 2.11-3.

Table 2.11-2. Common Bulk Chemicals Required at the Project Processing Sites*†

Shipped as Dry Bulk Solids	Shipped as Liquids and Gases
Salt (NaCl)	Hydrochloric acid (HCl)
Sodium bicarbonate (NaHCO ₃)	Sulfuric acid (H ₂ SO ₄)
Sodium carbonate (Na ₂ CO ₃)	Hydrogen peroxide (H ₂ O ₂)
Sodium hydroxide (NaOH)	Oxygen (O ₂)
—	Carbon dioxide (CO ₂)
—	Anhydrous ammonia (NH ₃)
—	Diesel oil
—	Bottled gases
—	Liquified petroleum gas (LPG)

*NRC. NUREG-1508, "Final Environmental Impact Statement to Construct and Operate the Crownpoint Uranium Solution Mining Project, Crownpoint, New Mexico." Washington, DC: NRC. February 1997.

†Energy Metals Corporation, U.S. "Application for USNRC Source Material License Moore Ranch Uranium Project, Campbell County, Wyoming: Environmental Report." ML072851249. Casper, Wyoming. Energy Metals Corporation, U.S. September 2007.

Table 2.11-3. Onsite Quantities of Process Chemicals at ISL Facilities*

Chemical	Typical Onsite Quantity	Use in Uranium ISL Process
Ammonia (NH ₃)	40,820 kg [90,000 lb]	pH adjustment
Sulfuric acid (H ₂ SO ₄)	37,850 L [10,000 gal]	pH control during lixiviant processing, and splitting uranyl carbonate complex into CO ₂ gas and uranyl ions in preparation for their precipitation
Liquid and gaseous oxygen	No specific typical quantities available	Oxidant in lixiviant, and precipitation of uranium as an insoluble uranyl peroxide compound
Hydrogen peroxide (H ₂ O ₂)	26,500 L [7,000 gal]	Uranium precipitation and oxidant in lixiviant
Sodium hydroxide (NaOH)	Typically stored in 208-L [55-gal] drums	pH adjustment
Barium chloride (BaCl ₂)	No specific typical quantities available	Precipitation of radium during groundwater restoration, and wastewater treatment
Carbon dioxide (CO ₂)	No specific typical quantities available	Carbonate complexing
Hydrochloric acid (HCl)	37,850 L [10,000 gal]	pH adjustment
Sodium carbonate (Na ₂ CO ₃)	64,350 L [17,000 gal]	Carbonate complexing and resin regeneration
Sodium chloride (NaCl)	127,000 kg [280,000 lb]	Resin regeneration
Hydrogen sulfide (H ₂ S)	No specific typical quantities available	Groundwater restoration
Sodium sulfide (Na ₂ S)	No specific typical quantities available	Groundwater restoration

*Mackin, P.C., D. Daruwalla, J. Winterle, M. Smith, and D.A. Pickett. NUREG/CR-6733, "A Baseline Risk-Informed Performance-Based Approach for *In-Situ* Leach Uranium Extraction Licensees." Washington, DC: NRC. September 2001.

Evaporation ponds are typically constructed in accordance with NRC staff guidance in NRC (1977, 2008), and license conditions require that these ponds be periodically monitored. Pond leaks have, however, occurred at active ISL facilities. For example, at the Crow Butte ISL facility in Dawes County, Nebraska, seven leaks were identified for three different commercial evaporation ponds from 1991 through 1997 (NRC, 1998b). The volumes of the leaks ranged from about 257.4 to 1,135.6 L [68 to 300 gal], but in all cases, the leaks involved only the upper liner of the double-lined system. To repair the leaks, the licensee exposed the liner by transferring water to other ponds to lower the water level, patched the holes, and pumped the water from the underdrain system (NRC, 1998b). Since, 1997, the Crow Butte facility has reported and repaired an additional eight pond leaks, with the most recent leak identified and

the pond liner repaired in May 2006 (Teahon, 2006). From 1988 to 1997, one pond leak was reported in 1992 at the Irigary/Christensen Ranch ISL facility in Campbell and Johnson Counties, Wyoming (NRC, 1998a). The licensee's corrective actions included temporarily transferring water to expose the liner and repair the leak.

The EPA- or state-issued UIC permit requires monitoring and testing the mechanical integrity of production and injection wells, reducing the potential for these types of failures. At the proposed Reynolds Ranch expansion of the Smith Ranch-Highland ISL facility in Converse County, Wyoming, the applicant established immediate spill responses through onsite standard operating procedures. These include shutting down the affected well or pipeline; recovering as much of the spilled fluid as possible; collecting samples of the affected soil so it can be compared to background values for uranium, radium-228, and selenium; and cleaning it up if necessary (NRC, 2006).

2.11.3 Groundwater Use

During construction, groundwater use is limited to routine activities such as dust suppression, mixing cements, and drilling support. Although large amounts of groundwater are moved and processed during ISL facility operations, most of the water is reinjected maintaining the overall water balance. A production bleed of about 1–3 percent, as discussed earlier, means that about 97–99 percent of the water produced from a well field is reinjected for additional uranium recovery. For example, for the proposed Reynolds Ranch addition to the Smith Ranch ISL facility in Converse County, Wyoming, the NRC staff estimated that the amount of water used in the ion exchange columns at the satellite facilities or discharged to a deep disposal well could be as much as 1,480,000,000 L [391 million gal] over the course of an assumed operating period of 15 years (NRC, 2006). For the Crow Butte ISL facility in Dawes County, Nebraska, the average operating flow rate in 2007 was about 16,200 L/min [4,279 gal/min] (Cameco Resources, Inc., 2008). The total net volume of groundwater produced for 2007 (volume produced–volume injected) was 346,900,000 L [91,640,000 gal], and the production bleed ranged from about 1.1 to 1.6 percent. During the last six months of 2007, about 76,200,000 L [20,130,000 gal] was disposed in the licensed Class I UIC deep disposal well and about 14,370,000 L [3,800,000 gal] was discharged to the evaporation pond system (Cameco Resources, 2008).

2.11.4 Excursions

As discussed in Section 2.4, ISL operations may affect the groundwater quality near the well fields or in over- or underlying aquifers when lixiviant travels from the production zone and beyond the well field boundaries. Monitoring wells are designed and placed to capture any lixiviant that moves out of the production zone. A monitoring well is placed on excursion status when two or more excursion indicators exceed their respective upper control limits (UCLs) (NRC, 2003a). NRC licensees are required by license conditions to identify reporting, monitoring, and response measures to be taken to determine the extent and cause of the excursion, as well as measures to recover the excursion into the well field and remove the well from excursion status.

Historical information for several facilities indicates that excursions can and do occur at ISL operations (NRC, 2006, 1998a,b, 1995; Crow Butte Resources, Inc., 2007; Cameco Resources, 2008; Arbogast, 2008). For example, from 1987 to 1998, 49 different wells were placed on excursion status at the Irigary and Christensen Ranch uranium recovery facility in Campbell and

Johnson Counties in the Wyoming East Uranium Milling Region (NRC, 1998a). Most of these excursions were recovered within a period of weeks to months, but six vertical excursions proved more difficult to return to baseline, with two wells remaining on excursion status for at least 8 years. These excursions were believed to be due to improperly abandoned wells from earlier exploratory programs prior to regulation by a UIC program. In 2007, three wells were on excursion status at the Christensen Ranch project, with only one, originally identified in 2004, remaining on excursion status at the end of 2007 (Arbogast, 2008a). None of the earlier wells identified in NRC (1998a) were still on excursion status. An additional well at the Christensen Ranch project was placed on excursion status in 2008 (Arbogast, 2008b).

From 1988 through 1995, 22 monitoring wells (11 vertical and 11 horizontal) were placed on excursion status for the Highland Uranium Project located in Converse County in the Wyoming East Uranium Milling Region (NRC, 1995). Most of the excursions were recovered within less than 1 year, but four horizontal excursions lasted up to at least five years. In two of these wells, the excursions were due to a thinning of the confining layer that separated two different production zones. Groundwater pumping during restoration of the underlying production zone resulted in establishing a hydraulic gradient that brought production fluids down from the overlying aquifer. One of the other excursions was believed to be the result of fluids migrating from an upgradient abandoned uranium mine (NRC, 1995). No cause was identified for the final long-term excursion at the Highland Uranium Project. Only one horizontal excursion was reported between 2001 and 2005 at the Smith Ranch-Highland uranium recovery facility, and corrective action brought the well back below the UCLs within less than one month (NRC, 2006).

At the Crow Butte ISL facility located in Dawes County, Nebraska (Nebraska-South Dakota-Wyoming Uranium Milling Region), the operator reported five vertical excursions into the overlying aquifer from the start of commercial operations in 1989 through the license renewal in 1998 (NRC, 1998b). In two cases, these excursions resulted from well integrity problems (borehole cement contamination and a failed casing coupling). One excursion resulted from a leak in a plugged and abandoned injection well, and the remaining two were believed to result from natural fluctuations in the groundwater quality (NRC, 1998b). Between 1999 and 2006, 17 wells at the Crow Butte facility were placed on excursion status (7 vertical and 10 horizontal). Most of these wells were restored below the UCLs within 1 to 6 months, although one vertical well took almost four years to restore (Crow Butte Resources, Inc., 2007). In the second half of 2007, three horizontal monitoring wells were on excursion status (Cameco Resources, 2008). These excursions were first identified in April 2000, December 2003, and September 2006 (Crow Butte Resources, Inc., 2007). The licensee believes that these longer term excursions resulted from well field geometry and well field flare as a result of ongoing groundwater transfer and well field restoration activities.

Operational experience at these facilities indicates that lixiviant excursions can result from

- Thinning or discontinuous confinement
- Improperly abandoned wells that may provide vertical flow pathways
- Casing failure or other well leaks
- Natural fluctuations in groundwater quality
- Improper balance of well field hydrologic gradients

Most horizontal excursions could be recovered quickly (weeks to months) by fixing and reconditioning wells and adjusting pumping rates in the well field, consistent with the findings of

Mackin, et al. (2001a). Vertical excursions tended to be more difficult to recover than horizontal excursions, and in a few cases, a well could remain on excursion status for a period of as much as 8 years.

2.11.5 Aquifer Restoration

Operational history at NRC-licensed ISL facilities is available to examine aquifer restoration at the well-field scale. In preparing the environmental report for the proposed Moore Ranch facility in Campbell County, Wyoming, Energy Metals Corporation, U.S., (2007) summarized mean groundwater quality conditions at the end of uranium recovery operations for a 12-ha [30-acre] area covered by Production Units 1–9 at the nearby COGEMA Irigaray ISL facility (Table 2.11-4). Before May 1980, the uranium recovery operations at Irigaray used an ammonium bicarbonate-hydrogen peroxide lixiviant. In May 1980, the facility was converted to a sodium bicarbonate-gaseous oxygen lixiviant. A comparison of the baseline and past recovery groundwater analytical data indicates that the water quality in the production zone is degraded for elements that make up part of the lixiviant (e.g., ammonia, bicarbonate, sodium) and for other elements (e.g., calcium and chloride).

Table 2.11-4. Irigaray Post-Uranium Recovery Water Quality*

Parameters (units)	Irigaray Baseline Range	Irigaray Post-Uranium Recovery Mean
Dissolved aluminum (mg/L)†	<0.05–4.25	<1.037
Ammonia nitrogen as N (mg/L)‡	<0.05–1.88	23
Dissolved arsenic (mg/L)	<0.001–0.105	<0.601
Dissolved barium (mg/L)	<0.01–0.12	<1.067
Boron (mg/L)	<0.01–0.225	<0.442
Dissolved cadmium (mg/L)	<0.002–0.013	<0.979
Dissolved chloride (mg/L)†	5.3–15.1	277
Dissolved chromium (mg/L)	<0.002–0.063	<1.018
Dissolved copper (mg/L)	<0.002–0.04	<0.828
Fluoride (mg/L)	0.11–0.66	<1
Total and dissolved iron (mg/L)	0.02–11.8	<1.098
Dissolved mercury (mg/L)	<0.0002–<0.001	<0.971
Dissolved magnesium (mg/L)	0.02–9.0	45.7
Total manganese (mg/L)	<0.005–0.190	1.249
Dissolved molybdenum (mg/L)	<0.02–<0.1	<1.067
Dissolved nickel (mg/L)	<0.01–<0.2	<1.018
Nitrate + nitrite as N (mg/L)	<0.2–1.0	<3
Dissolved lead (mg/L)	<0.002–<0.050	<1.018
Radium-226 (pCi/L)	0–247.7	200.5
Dissolved selenium (mg/L)	<0.001–0.416	0.247
Dissolved sodium (mg/L)	95–280	827
Sulfate (mg/L)	136–824	639

Table 2.11-4. Irigaray Post-Uranium Recovery Water Quality*
(continued)

Parameters (units)	Irigaray Baseline Range	Irigaray Post-Uranium Recovery Mean
Uranium (mg/L)	<0.0003–18.8	7.411
Vanadium (mg/L)	<0.05–0.55	<1.067
Dissolved zinc (mg/L)	<0.01–0.200	<0.065
Dissolved calcium (mg/L)†	1.6–33.5	199.2
Bicarbonate (mg/L)†	5–144	1,343
Carbonate (mg/L)	0–96	<2
Dissolved potassium (mg/L)	0.4–17.5	9
Total dissolved solids at 180 °F (mg/L)	308–1,054	2,451
*Energy Metals Corporation, U.S. "Application for USNRC Source Material License Moore Ranch Uranium Project, Campbell County, Wyoming: Environmental Report." ADAMS ML072851249. Casper, Wyoming: Energy Metals Corporation U.S. 2007.		
†1 mg/L = 1 ppm		
‡Parameters with restoration value other than baseline.		

Catchpole, et al. (1992a,b) provide an early discussion of small-scale restoration efforts for research and development (R&D) of ISL uranium recovery facilities in Wyoming. These include the Bison Basin facility in Fremont County (described in NRC, 1981), the Reno Creek project in Campbell County, and the Leuenberger Project in Converse County. Restoration activities required treatment of water from nine pore volumes at Bison Basin and five pore volumes at Reno Creek. In all cases, most water quality parameters were returned to within a statistical range of baseline values with the exception of uranium (Bison Basin and Reno Creek) and radium-226 (Leuenberger). For these parameters, Catchpole, et al. (1992a,b) report that water in the well field was returned to the same class of use.

Davis and Curtis (2007) detailed available information on aquifer restoration at ISL uranium recovery facilities. These include a pilot scale study by Rio Algom for the Smith Ranch facility in Converse County, Wyoming (Rio Algom Mining Corporation, 2001); the proposed Crownpoint ISL facility near Crownpoint, New Mexico (NRC, 1997); the A-Well Field at the Highland Uranium Project in Converse County, Wyoming (Power Resources, Inc., 2004a); and the Crow Butte Mine Unit No. 1 in Dawes County, Nebraska (NRC, 2002, 2003c). Rock core laboratory studies that Hydro Resources Inc. conducted for the Crownpoint facility (NRC, 1997a) also provide useful insights to water quality parameters that may present challenges for aquifer restorations.

Davis and Curtis (2007) generally concluded that for the sites and data they examined, aquifer restoration took longer and required more pore volumes than originally planned. For example, at the A-Well Field at the Highland Uranium Project, the licensee's original plan anticipated that restoration would last from four to seven years and require treating 5–7 pore volumes of groundwater. When uranium recovery in the well field ended in 1991, the baseline and class of use were not restored in the well field until 2004 (Table 2.11-5), and more than 15 pore volumes of water were involved (NRC, 2006, 2004). Similarly, WDEQ has noted that the C-Well field at Smith-Ranch-Highland has been undergoing restoration for 10 years (WDEQ, 2008). At the Crow Butte Mine Unit No. 1, more than 9.85 pore volumes of groundwater were used in all the stages of aquifer restoration over approximately 5 years as compared to the 8 pore volumes

estimated before restoration (NRC, 2002, 2003c). CBR extracted uranium from an additional 26 pore volumes using ion exchange, without lixiviant injection, prior to active restoration.

Table 2.11-5. Baseline Groundwater Conditions, Aquifer Restoration Goals, and Actual Final Restoration Values NRC Approved for the Q-Sand Pilot Well Field, Smith Ranch, Wyoming*†

Parameter (units)	Range	Mean	Restoration Goal	Actual Restoration
Arsenic (mg/L‡)	0.001–0.0013	0.004	0.05	0.008
Boron (mg/L)	0.002–0.70	0.15	0.54	0.14
Calcium (mg/L)	24–171	72	120	78
Iron (mg/L)	0.01–0.27	0.025	0.3	0.24
Magnesium (mg/L)	3–22	16	0.092	0.06
Manganese (mg/L)	0.01–0.077	0.023	Not applicable	0.1
Selenium (mg/L)	0.001–0.024	0.004	0.029	0.003
Uranium (mg/L)	0.001–3.1	0.28	3.7	1.45
Chloride (mg/L)	4–65	18	250	15
Bicarbonate (HCO ₃) (mg/L)	129–245	199	294	254
Carbonate (CO ₃) (mg/L)	Nondetectible–75	18	15	Nondetectible
Nitrate (mg/L)	0.1–1.0	0.4	Not applicable	0.13
Potassium (mg/L)	7–34	12	23	8
Sodium (mg/L)	19–87	28	41	38
Sulfate (mg/L)	100–200	124	250	128
Total dissolved solids (mg/L)	155–673	388	571	443
Specific conductivity (µmhos/cm)	518–689	582	827	642
pH (standard units)	7.5–9.4	8.0	6.5–8.6	7.0
Radium-226 (pCi/l)	6–1132	340	923	477
Thorium-230 (pCi/l)	0.027–4.65	1.03	5.62	3.4

*NRC. "Environmental Assessment for the Addition of the Reynolds Ranch Mining Area to Power Resources, Inc.'s Smith Ranch/Highlands Uranium Project Converse County, Wyoming." Source Material License No. SUA-1548. Docket No. 40-8964. Washington, DC: NRC. 2006.

†Sequoyah Fuels Corporation. "Re: License Application, Smith Ranch Project, Converse County, Wyoming." ML8805160068. Glenrock, Wyoming: Sequoyah Fuels Corporation. 1988.

‡1 mg/L = 1 ppm

As a field test of groundwater stabilization during aquifer restoration, hydrogen sulfide gas was injected as a reductant into the Ruth ISL research and development facility in Campbell County, Wyoming. After 6 weeks of hydrogen sulfide injection, pH dropped relatively quickly from 8.6 to 6.3, and sulfate concentration increased from 28 ppm to 91 ppm indicating a more reducing environment (Schmidt, 1989; Davis and Curtis, 2007). Concentrations of dissolved uranium, selenium, arsenic, and vanadium decreased by at least one order of magnitude. After one year

of monitoring, however, reducing conditions were not maintained, and uranium, arsenic, and radium concentrations began to increase.

Based on the available field data from aquifer restoration, Davis and Curtis (2007) concluded that aquifer restoration is complex and results could be influenced by a number of site-specific hydrological and geochemical characteristics. As discussed previously, in some cases, such as at Bison Basin and Reno Creek, the aquifer was restored in a relatively short time. In other cases, restoration required much more time and treatment than was initially estimated (e.g., the A- and C- Well Fields at the Highland ISL facility).

2.11.6 Socioeconomic Information

Because they are generally located in remote areas, uranium ISL facilities tend to be important employers in the local economy. The total number of full-time, permanent employees and local contractors varies during an operational life that may span several decades. Based on employment levels at existing operations and projected employment for proposed projects, staff levels at ISL facilities range from about 20 to 200, with peak employment depending on the scheduling of construction, drilling, and operational activities (Crow Butte Resources, Inc., 2007; Power Resources, Inc., 2004a; NRC, 1997a).

Another economic effect from ISL facilities is contributions to the local economy through purchases and through tax revenues from the uranium produced at the facility. For example, at the Crow Butte ISL facility in Dawes County, Nebraska, local purchases of goods and services in 2006 were estimated at about \$5,000,000 (Crow Butte Resources, Inc., 2007). Annual tax revenues depend on uranium prices and the amount of uranium produced at a given facility. For example, for a 272,155-kg [600,000-lb] increase in annual yellowcake production at the Crow Butte facility at a price of \$80/lb, an incremental contribution to federal, state, and local taxes on the order of \$1 million to \$1.4 million would result (Crow Butte Resources, Inc., 2007).

2.12 Alternatives Considered and Included in the Impact Analysis

The NRC's environmental review regulations in 10 CFR Part 51 that implement the National Environmental Policy Act (NEPA) require the NRC to consider reasonable alternatives, including the no-action alternative, to a proposed action before acting on a proposal. The intent is to enable the agency to consider the relative environmental consequences of an action given the environmental consequences of other activities that also meet the need for the action, as well as the environmental consequence of taking no action at all. The information in this section does not constitute NRC's final consideration of reasonable alternatives for the site-specific environmental reviews of ISL license applications.

2.12.1 The No-Action Alternative

As defined in Chapter 1, the proposed action is to identify and evaluate the potential environmental impacts associated with the construction, operation, aquifer restoration, and decommissioning of ISL facilities in designated regions of the western U.S. In the No-Action Alternative, no additional ISL activity would take place in the four geographic regions considered in this Draft GEIS. As a result, the regions would not see additional ISL activities as described in Chapter 2 nor the associated potential environmental impacts discussed in Chapter 4. Ongoing and reasonably foreseeable future activities as described in Chapter 5 would still impact the regions.

2.13 Alternatives Considered and Excluded From the Impact Analysis

Alternative methods for uranium recovery include conventional mining/milling methods and heap leaching. Heap leaching (i.e., use of chemical solutions to leach uranium from a pile of crushed ore) may be used for low grade or small ore bodies, but mining and some crushing and grading is necessary to build up the ore pile (EPA, 2007a; NRC, 1980). The heap leach process is a technology that is considered to be part of the conventional mining and milling industry; NRC regulates this technology using the criteria in 10 CFR 40, Appendix A, that are deemed applicable to such operations (NRC, 1980, Appendix B). These two alternative uranium recovery technologies are discussed further in Appendix C.

Because the Draft GEIS focuses on the future licensing of ISL facilities and does not evaluate available technologies for uranium recovery, conventional mining/milling and heap leaching were not included in the impact analysis. However, such uranium recovery methods may be among the reasonable alternatives evaluated in a site-specific review of an ISL license application. As described in Section 2.1, there are particular types of uranium deposits that are amenable to ISL uranium recovery technology. In certain cases (e.g., the ore body is located near the surface), these deposits may also be accessible by conventional mining techniques, with the uranium in the mined ore recovered by conventional milling methods or by heap leaching. Therefore, the alternatives to be considered will be addressed in the site-specific environmental reviews.

2.14 References

- Arbogast, L. "Letter and Introduction—Submittal of 2007 Annual Effluent and Monitoring Report for COGEMA Mining, Inc." Letter (February 15) to K. McConnell, NRC. Mills, Wyoming: Cogema Mining, Inc. 2008a.
- Arbogast, L. "Monitor Well 5MW 48 on Excursion Status." April 21, 2008. Mills, Wyoming: Cogema Mining, Inc. 2008b.
- Cameco Resources. "Crow Butte Uranium Project Radiological Effluent and Environmental Monitoring Report for Third and Fourth Quarters 2007." Crawford, Nebraska: Cameco Resources, Crow Butte Operations. February 2008.
- Catchpole, G., R. Garling, and M. Neumann. "Groundwater Restoration at Wyoming Uranium Solution Mining Sites." *The Mining Claim*. Cheyenne, Wyoming: Wyoming Mining Association. pp. 6–9. 1992a.
- Catchpole, G., R. Garling, and M. Neumann. "Groundwater Restoration at Wyoming Uranium Solution Mining Sites. Part II." *The Mining Claim*. Cheyenne, Wyoming: Wyoming Mining Association. pp. 14–19. 1992b.
- Cogema Mining, Inc. "Request for NRC Concurrence: Irigaray Restoration Approval." Letter (November 7) to G. Janosko, NRC. Docket No. 40-8502. 2005.
- Crow Butte Resources, Inc. "SUA-1535 License Renewal Application." Crawford, Nebraska: Crow Butte Resources, Inc. November 2007.

Davis, J.A. and G.P. Curtis. NUREG/CR-6870, "Consideration of Geochemical Issues in Groundwater Restoration at Uranium *In-Situ* Leaching Mining Facilities." Washington, DC: NRC. January 2007.

Deutsch, W.J., W.J. Martin, L.E. Eary, and R.J. Serne. NUREG/CR-3709, "Method of Minimizing Ground-Water Contamination From In Situ Leach Uranium Mining." Washington, DC: NRC. 1985.

Energy Information Administration. "Domestic Uranium Production Report—Quarterly: Data for 4th Quarter 2007." Release Date February 12, 2008. Washington, DC: Energy Information Administration. <<http://www.eia.doe.gov/cneaf/nuclear/dupr/qupd.html>> (11 April 2008).
Energy Information Administration. "Uranium Reserve Estimates." 2004. <www.eia.doe.gov/cneaf/nuclear/page/reserves/ures.html> (14 September 2007).

Energy Information Administration. DOE/EIA-0592, "Decommissioning of U.S. Uranium Production Facilities." Washington, DC: Energy Information Administration, Office of Coal, Nuclear, Electric, and Alternate Fuels. February 1995.

Energy Metals Corporation, U.S.. "Application for USNRC Source Material License, Moore Ranch Uranium Project, Campbell County, Wyoming, Environmental Report." Casper, Wyoming: Energy Metals Corporation. September 2007.

EPA. "Technical Report on Technologically Enhanced Naturally Occurring Radioactive Materials From Uranium Mining. Volume 1: Mining and Reclamation Background." EPA 402-R-05-007. Washington, DC: EPA, Office of Radiation and Indoor Air. 2007a. <www.epa.gov/radiation/docs/tenorm/402-r-05-007-rev0607.pdf> (20 November 2007).

EPA. "Technical Report on Technologically Enhanced Naturally Occurring Radioactive Materials From Uranium Mining. Volume 2: Investigation of Potential Health, Geographic, and Environmental Issues of Abandoned Uranium Mines." EPA 402-R-05-007. Washington, DC: EPA, Office of Radiation and Indoor Air. 2007b. <<http://www.epa.gov/radiation/docs/tenorm/volume-ii/402-r-05-007.pdf>> (20 November 2007).

EPA. Technical Resource Document. Extraction and Beneficiation of Ores and Minerals, Volume 5, Uranium." EPA 530-R-94-032. Washington, DC: EPA. January 1995.
Finch, J. "New ISR Processing Technology Could Help Boost U.S. Uranium Mining." 2007. <www.stockinterview.com/News/03192007/ISR-Uranium-Remote-Ion-Exchange.html> (14 September 2007).

Holen, H.K. and W.O Hatchell. "Geological Characterization of New Mexico Uranium Deposits for Extraction by *In-Situ* Leach Recovery." Open-File Report No. 251. Socorro, New Mexico: New Mexico Bureau of Mines and Mineral Resources. August 1986.

Hunkin, G.G. Uranium In-Situ Leaching in the Tertiary Deposits of South Texas, Chapter 3: Resource Development / Utilization in M.D. Campbell ed., "Geology of Alternate Energy Resources in the South-Central United States" Houston Geological Society, pp. 67-82. 1977.

International Atomic Energy Agency. *Guidebook on Environmental Impact Assessment for In-Situ Leach Mining Project*. TECDOC-1428. Vienna, Austria: International Atomic Energy Agency. 2005.

- 1
2 International Atomic Energy Agency. "Manual of Acid In Situ Leach Uranium Mining
3 Technology." IAEA-TECDOC-1239. Vienna, Austria: International Atomic Energy Agency.
4 August 2001.
- 5
6 Mackin, P.C., D. Daruwalla, J. Winterle, M. Smith, and D.A. Pickett. NUREG/CR-6733, "A
7 Baseline Risk-Informed Performance-Based Approach for *In-Situ* Leach Uranium Extraction
8 Licensees." Washington, DC: NRC. September 2001a.
- 9
10 Mackin, P.C., W.A. Illman, D.L. Hughson, and D.A. Pickett. "Surety Estimation Methodology for
11 Groundwater Corrective Actions at Title II Conventional Mills—Final Report." San Antonio,
12 Texas: Center for Nuclear Waste Regulatory Analyses. August 2001b.
- 13
14 McCarthy, J. "Smith Ranch: 2007–2008 Surety Estimate Revision." Letter (June 29) to
15 G. Janosko, NRC. Glenrock, Wyoming: Power Resources International. 2007.
- 16
17 McLemore, V.T. "Uranium Resources in New Mexico." Society of Mining and Metallurgical
18 Engineering Annual Meeting, Denver, Colorado, February 25–28, 2007. Littleton, Colorado:
19 Society of Mining and Metallurgical Engineering. 2007.
- 20
21 Mudd, G.M. "Critical Review of Acid *In-Situ* Leach Uranium Mining: 1—USA and Australia."
22 *Environmental Geology*. Vol. 41. pp. 390–403. 2001.
- 23
24 Mudd, G.M. "An Environmental Critique of *In-Situ* Leach Uranium Mining; the Case Against
25 Uranium Solution Mining." Research report. Melbourne, Australia: The Sustainable Energy
26 and Anti-Uranium Service, Inc. 1998. <<http://www.sea-us.org.au/isl>> (15 November 2007).
- 27
28 Nash, J.T., H.C. Granger, and S. Adams. "Geology and Concepts of Important Types of
29 Uranium Deposits." B.J. Skinner, ed. *Economic Geology*. 75th Anniversary Volume
30 1905–1980. pp. 63–116. 1981.
- 31
32 NRC. NRC Draft Regulatory Guide DG-3032. "Design, Construction, and Inspection of
33 Embankment Retention Systems at Uranium Recovery Facilities." Proposed Revision 3 of
34 Regulatory Guide 3.11, dated December 1977. Washington, DC: NRC. February 2008.
- 35
36 NRC. "Environmental Assessment for Amendment to Source Materials License SUA-1534 for a
37 Central Processing Plant Upgrade. Crow Butte Resources, Inc. *In-Situ* Leach Uranium
38 Recovery Facility Crawford, Dawes County, Nebraska." Washington, DC: NRC. October
39 2007a.
- 40
41 NRC. "Environmental Assessment for Construction and Operation of In Situ Leach Satellite
42 SR-2, Amendment No. 12 to Source Materials License No. SUA-1548." Docket No. 40-8964.
43 Washington, DC: NRC. 2007.
- 44
45 NRC. "Environmental Assessment for the Addition of the Reynolds Ranch Mining Area to
46 Power Resources, Inc.'s Smith Ranch/Highlands Uranium Project Converse County, Wyoming."
47 Source Material License No. SUA-1548. Docket No. 40-8964. Washington, DC: NRC. 2006.
- 48
49 NRC. "Review of Power Resources, Inc.'s A-Well Field Ground Water Restoration Report for
50 the Smith Ranch-Highland Uranium Project." ML041840470. Washington, DC: NRC. 2004.
- 51

- 1 NRC. NUREG-1569, "Standard Review Plan for *In-Situ* Leach Uranium Extraction License
2 Applications—Final Report." Washington, DC: NRC. June 2003a.
- 3
- 4 NRC. NUREG-1748, "Environmental Review Guidance for Licensing Actions Associated With
5 NMSS Programs. Final Report." Washington, DC: NRC. August 2003b.
- 6
- 7 NRC. "License Amendment 15, Crow Butte Resources In Situ Leach Facility, License
8 No. SUA-1534, Well Field #1 Restoration Acceptance." ML030440055. Washington, DC:
9 NRC. 2003c.
- 10
- 11 NRC. "Denial, Well Field Unit 1 Ground-Water Restoration Approval, Crow Butte Resources In
12 Situ Leach Facility, License No. SUA-1534." ML020930087. Washington, DC: NRC. 2002.
- 13
- 14 NRC. "Exothermic Reactions Involving Dried Uranium Oxide Powder (Yellowcake)."
15 Notice 99-03. Washington, DC: NRC. 1999.
- 16
- 17 NRC. "Environmental Assessment for Renewal of Source Material License No. SUA-1341,
18 Cogema Mining, Inc. Irigaray and Christensen Ranch Projects, Campbell and Johnson
19 Counties, Wyoming." Docket No. 40-8502. Washington, DC: NRC. June 1998a.
- 20
- 21 NRC. "Environmental Assessment for Renewal of Source Material License
22 No. SUA-1534—Crow Butte Resources Incorporated Crow Butte Uranium Project Dawes
23 County, Nebraska." Docket No. 40-8943. Washington, DC: NRC. 1998b.
- 24
- 25 NRC. NUREG-1508, "Final Environmental Impact Statement To Construct and Operate the
26 Crownpoint Uranium Solution Mining Project, Crownpoint, New Mexico." Washington, DC:
27 NRC. February 1997a.
- 28
- 29 NRC. "Annual Financial Surety Update Requirements for Uranium Recovery Licensees."
30 Generic Letter 97-03. Washington, DC: NRC. July 1997b. <[http://www.nrc.gov/reading-](http://www.nrc.gov/reading-rm/doc-collections/gen-comm/gen-letters/_1997/gl97003.html)
31 [rm/doc-collections/gen-comm/gen-letters/_1997/gl97003.html](http://www.nrc.gov/reading-rm/doc-collections/gen-comm/gen-letters/_1997/gl97003.html)> (23 October 2007).
- 32
- 33 NRC. "Environmental Assessment for Renewal of Source Materials License No. SUA-1511.
34 Power Resources Incorporated Highland Uranium Project, Converse County, Wyoming."
35 Washington, DC: NRC. August 1995.
- 36
- 37 NRC. "Environmental Assessment for Rio Algom Mining Corporation Smith Ranch *In-Situ*
38 Leach Mining Project, Converse County, Wyoming In Consideration of a Source and Byproduct
39 Material License Application." Docket No. 40-8964. Washington DC: NRC. January 1992.
- 40
- 41 NRC. Regulatory Guide 3.46, "Standard Format and Content of License Applications, Including
42 Environmental Reports, for *In-Situ* Uranium Solution Mining." Washington, DC: NRC. 1982.
- 43
- 44 NRC. "Final Environmental Statement Related to the Operation of Bison Basin Project." Docket
45 No. 40-8745. Washington, DC: NRC. 1981.
- 46
- 47 NRC. NUREG-0706, "Final Generic Environmental Impact Statement on Uranium Milling
48 Project M-25." Washington, DC: NRC. September 1980.
- 49
- 50 NRC. NRC Regulatory Guide 3.11, Rev. 2, "Design, Construction, and Inspection of
51 Embankment Retention Systems for Uranium Mills." Washington, DC: NRC. December 1977.

- 1
2 NRC. NUREG-0489, "Final Environmental Statement Related to Operation of Highland
3 Uranium Solution Mining Project, Exxon Minerals Company, USA." Washington DC: NRC.
4 November 1978.
5
- 6 Power Resources, Inc. "Smith Ranch--Highland Uranium Project, A-Well Field Groundwater
7 Restoration Information." ML040300369. Glenrock, Wyoming: Power Resources, Inc. 2004a.
8
- 9 Power Resources, Inc. "Reynolds Ranch Amendment—Permit to Mine No 1548—
10 Smith Ranch-Highland Uranium Project—Volume I, Chapters 1–10." ML050390095. Glenrock,
11 Wyoming: Power Resources, Inc. 2004b.
12
- 13 Rio Algom Mining Corporation. "Amendment 1 to Source Material License SUA-1548."
14 ML020220040. Washington, DC: NRC. 2001.
15
- 16 Schmidt, C. "Groundwater Restoration and Stabilization at the Ruth ISL Test Site in Wyoming,
17 USA." *In-Situ Leaching of Uranium: Technical, Environmental, and Economic Aspects*.
18 IAEA-TECDOC-492. Vienna, Austria: International Atomic Energy Agency. pp. 97–126.
19 1989.
20
- 21 Staub, W.P., N.E. Hinkle, R.E. Williams, F. Anastasi, J. Osiensky, and D. Rogness.
22 NUREG/CR-3967, "An Analysis of Excursions at Selected *In Situ* Uranium Mines in Wyoming
23 and Texas." Washington, DC: NRC. July 1986.
24
- 25 Stout, R.M. and D.E. Stover. "The Smith Ranch Uranium Project." The Uranium Institute 22nd
26 Annual Symposium, September 3–5, 1997, London, England. London, United Kingdom: World
27 Nuclear Association. 1997.
28
- 29 Teahon, L. "Evaporation Pond 4 Liner Leak." Letter (June 1) to G. Janosko, NRC. Crawford,
30 Nebraska, Crow Butte Resources, Inc. 2006.
31
- 32 Tweeton, D.R. and K.A. Peterson. "Selection of Lixivants for *In-Situ* Leach Mining." *In-Situ*
33 Mining Research. U.S. Bureau of Mines Technology Transfer Seminar. Information
34 Circular 8852. 1981.
35
- 36 WDEQ. "*In-Situ* Uranium Permits 603 and 633, Notice of Violation, Docket No. 4231-08."
37 Cheyenne, Wyoming: WDEQ, Land Quality Division. March 10, 2008.
38
- 39 WDEQ. "Notice of Violation—Re: *In-Situ* Uranium Operation Permit #603
40 (Docket No. 4122-07)." Cheyenne, Wyoming: WDEQ, Land Quality Division. 2007.
41 <<http://deq.state.wy.us/out/downloads/LQ%20NOV%209.12.07a.pdf>> (27 February 2008)
42

3 DESCRIPTION OF THE AFFECTED ENVIRONMENT

3.1 Introduction

This chapter of the Draft GEIS provides a description of the environmental conditions and resources in four regions of Wyoming, South Dakota, Nebraska, and New Mexico where previous and existing ISL uranium recovery operations have been licensed by NRC and where new ISL facilities may be proposed for NRC review. These uranium milling regions are defined in Section 3.1.1 and provide the basis for the structure of Chapter 3, which describes the affected environments for each region. Section 3.1.2 includes general information that applies to each of the four regions.

3.1.1 Geographic Scope—Defining Uranium Milling Regions

For the purpose of analysis in this Draft GEIS, NRC assumptions about potential future ISL facility locations were based on:

- The locations of past and existing uranium milling operations in States where NRC has the regulatory authority over uranium recovery;
- The locations where uranium milling companies have expressed interest in future uranium recovery using the ISL process; and
- The locations of historical uranium ore deposits in Wyoming, South Dakota, Nebraska, and New Mexico.

In the United States, uranium ore deposits have been studied and developed in a number of western states: Arizona, Colorado, Montana, Nebraska, New Mexico, South Dakota, Utah, Washington, Wyoming, and Texas (see Figure 1.1-2). Regional ore deposits found in those states can encompass portions of several contiguous states.

The affected environment described in this chapter is further limited to states where NRC has authority to license ISL facilities. NRC does not have regulatory authority in all states because at the state's request, NRC may relinquish its regulatory authority to the state. Therefore, in certain states, known as Agreement States, NRC has relinquished its regulatory authority to license uranium milling facilities. Colorado, Utah, and Texas are Agreement States with state, not NRC, regulation of uranium milling. NRC has retained its regulatory authority over uranium milling activities in non-Agreement States. Western non-Agreement States where NRC regulates uranium milling activities include Wyoming, South Dakota, Nebraska, and New Mexico. Montana and Arizona are also non-Agreement States with respect to uranium milling. One uranium milling company has indicated to NRC its plans for an ISL facility in Montana near its border with Wyoming, but no companies have indicated to NRC their plans to construct and operate ISL facilities in Arizona (NRC, 2008).

Locations within Wyoming, South Dakota, Nebraska, and New Mexico that include ore deposits and where past, existing, or future uranium milling activities or interest has been identified are shown in Figures 3.1-1, 3.1-2, 3.1-3, and 3.1-4.




Figure 3.1-1. Wyoming West Uranium Milling Region With Current and Potential ISL Milling Sites





Figure 3.1-2. Wyoming East Uranium Milling Region With Current and Potential ISL Milling Sites

1
2



SOUTH DAKOTA - NEBRASKA REGION




Figure 3.1-3. Nebraska-South Dakota-Wyoming Uranium Milling Region With Current and Potential ISL Milling Sites

3
4




Figure 3.1-4. New Mexico Uranium Milling Region With Current and Potential ISL Milling Sites

1




Figure 3.1-5. Simplified Cross-Section of Sandstone Uranium Roll-Front Deposits Formed by Regional Groundwater Migration (NRC, 1997)

As shown in the figures, NRC has delineated separate uranium milling regions where the boundaries of each milling region encompass past, existing, and potential future ISL milling sites. In defining these regions, NRC also considered aspects of the affected environment (e.g., regional ground water characteristics, regional demographics) such that potential future ISL milling sites within each region would more likely share those aspects for the purpose of evaluating potential environmental impacts. Therefore, NRC considers that these regions reasonably bound the geographic scope of the Draft GEIS for describing the affected environment and for assessing potential environmental impacts within each region.

For the purposes of the Draft GEIS, the regions have been named (see Section 1.4)

- Wyoming West Uranium Milling Region (Section 3.2)
- Wyoming East Uranium Milling Region (Section 3.3)
- Nebraska-South Dakota-Wyoming Uranium Milling Region (Section 3.4)
- Northwestern New Mexico Uranium Milling Region (Section 3.5)

Using this regional approach, the assessments of impacts in the Draft GEIS may or may not be applicable or informative to reviews of ISL facilities proposed outside of the designated uranium milling regions. In such cases, the applicability of the Draft GEIS would depend on the similarities of the proposed site and regional conditions with those described in the Draft GEIS.

Identifying regions based on the locations of past, existing, and potential future uranium recovery operations as is done in the Draft GEIS does not mean NRC prefers these locations or would prevent uranium recovery in other areas. It is the applicant or licensee that proposes the location of an ISL facility in the license application submitted to NRC, and NRC reviews such applications to fulfill its regulatory responsibilities.

3.1.2 General Information for All Uranium Milling Regions

To limit redundancies in discussing general information applicable to all four uranium milling regions addressed by the Draft GEIS, that information is provided in this section.

Sandstone-hosted uranium deposits account for the vast majority of the uranium ore produced in Wyoming, South Dakota, Nebraska, and New Mexico (Chenoweth, 1988, 1991; Collings and Knode, 1984; McLemore and Chenoweth, 1989, 2003). Uranium mineralization in these sandstone deposits occurs primarily in what have been termed stratabound or roll-front deposits (Rackley, 1972; Renfro, 1969; Collings and Knode, 1984; McLemore, 2007). A conceptual model of a roll-front uranium deposit is illustrated in Figure 3.1-5. Roll fronts occur where water infiltrates from the surface and flows through an aquifer with slight amounts of uranium. Near the surface, oxidizing conditions cause the minerals and volcanic ash to weather (or dissolve) and release minute quantities of uranium into the groundwater. As groundwater continues to flow, it can encounter reducing conditions where the uranium is no longer stable in solution. In an aquifer, a reducing environment is characterized by the presence of hydrogen sulfide (H_2S), iron sulfides, or organic material. As a result, uranium precipitates from the groundwater and forms mineral coatings on the sediment grains in the formation. Principal uranium ore mineral coatings found in the roll-front deposits include uraninite (UO_2) and coffinite (USiO_4). Roll-front deposits are ideally crescent- or C-shaped when viewed in cross section, with thin mineralization forming the tips of the crescents. Thick mineralization occurs in the center of the concave C-shaped ore body in the direction of groundwater flow. Individual mineralization fronts are typically from 0.6 m [2 ft] to more than 7.5 m [25 ft] thick and may be several hundred meters [feet] long. Fronts may coalesce to form ore bodies kilometers [miles] in length. Thin mineralized trails and more finely disseminated minerals branch off the main front and are located between fronts. High grade uranium roll-front deposits average about 0.2 percent U_3O_8 . Lower grade ore (0.05–0.10 percent U_3O_8) is commonly present on the unaltered side of the higher grade roll-front.

Several features are common to most major sandstone roll-front uranium deposits and their host rocks in Wyoming, South Dakota, Nebraska, and New Mexico (Rackley, 1972; McLemore, 2007). These features are: (1) sandstones of fluvial origin (i.e., produced by the action of a stream or river); (2) common association with arkosic (i.e., sediments with a considerable amount of the mineral feldspar) or micaceous sediment; (3) siltstones and mudstones interbedded with sandstones; (4) association with organic materials; (5) presence of pyrite in unweathered deposits; (6) gray color of the sandstones and light-gray or green color of the mudstones in unweathered deposits; (7) association with volcanic debris in the host formation or in overlying formations; (8) the discordant roll front features or solution fronts; and (9) the sharp contact between mineralized zones and adjacent carbonaceous-free or oxidized zones. The first seven features are related directly to the source rock, sedimentation, and the sedimentary environment; the last two features are related to the mineralizing process.

3.1.3 References

- Chenoweth, W.L. "A Summary of Uranium Production in Wyoming." Mineral Resources of Wyoming, Wyoming Geological Association, 42nd Annual Field Conference Guidebook. Casper, Wyoming: Wyoming Geological Association. pp. 169–179. 1991.
- Chenoweth, W.L. "Geology and Production History of the Uranium Deposits in the Northern Black Hills, Wyoming—South Dakota." Eastern Powder River Basin, Wyoming Geological Association, 39th Annual Field Conference Guidebook. Casper, Wyoming: Wyoming Geological Association. pp. 263–270. 1988.
- Collings, S.F. and R.H. Knode. "Geology and Discovery of the Crow Butte Uranium Deposit, Dawes County, Nebraska." Practical Hydromet '83, 7th Annual Symposium on Uranium and Precious Metals. Littleton, Colorado: American Institute of Mining, Metallurgical, and Petroleum Engineering. 1984.
- Energy Information Administration. "Uranium Reserve Estimates." 2004. <www.eia.doe.gov/cneaf/nuclear/page/reserves/ures.html> (14 September 2007).
- McLemore, V.T. "Uranium Resources in New Mexico." Society of Mining and Metallurgical Engineering Annual Meeting, Denver, Colorado, February 25–28, 2007. Littleton, Colorado: Society of Mining and Metallurgical Engineering. 2007.
- McLemore, V.T. and W.L. Chenoweth. "Uranium Resources in the San Juan Basin, New Mexico." Geology of the Zuni Plateau: New Mexico Geological Society, Guidebook 54. pp. 165–178. 2003.
- McLemore, V.T. and W.L. Chenoweth. "Uranium Resources in New Mexico." New Mexico Bureau of Mines and Mineral Resources Map 18. Socorro, New Mexico: New Mexico Bureau of Mines and Mineral Resources. 1989.
- NRC. "Expected New Uranium Recovery Facility Applications/Restarts/Expansions: Updated 1/24/2008." 2008. <<http://www.nrc.gov/info-finder/materials/uranium/2008-ur-projects-list-public-012408.pdf>> (08 February 2008).
- NRC. NUREG–1508, "Final Environmental Impact Statement to Construct and Operate the Crown Point Uranium Solution Mining Project, Crown Point, New Mexico." Washington, DC: NRC. February 1997.
- Rackley, R.I. "Environment of Wyoming Tertiary Uranium Deposits." American Association Petroleum Geologists Bulletin. Vol. 56, No. 4. 1972.
- Renfro, A.R. "Uranium Deposits in the Lower Cretaceous of the Black Hills." Contributions to Geology. Laramie, Wyoming: University of Wyoming. Vol. 8, No. 2-1. pp. 87–92. 1969.

3.2 Wyoming West Uranium Milling Region

3.2.1 Land Use

Approximately 53.3 percent of the land in the State of Wyoming is public land (47 percent federal ownership and 6.3 percent state ownership). Most of these federal lands are located in the western and northwestern parts of Wyoming and the vast majority of private lands are located in the eastern half of the state. The U.S. Bureau of Land Management (BLM) administers the largest amount of public land in the state (28 percent). BLM lands are mixed with private and state lands. Private lands, including Native American lands, which are administered by the Bureau of Indian Affairs (BIA), represent 45.9 percent of Wyoming land. In terms of general landscape, Wyoming big sagebrush (30.8 percent) and mixed grass (20.2 percent) occupy about half of the land in Wyoming, while irrigated agriculture occupies only 4.2 percent of the land (Wyoming Geographic Information Science Center, 2008).

For the purpose of this Draft GEIS, the Wyoming West Uranium Milling Region encompasses parts of Carbon, Fremont, Natrona and Sweetwater Counties (Figure 3.2-1). This region, which is a part of the Rocky Mountain System, straddles the Wyoming Basin to the east and the Middle Rocky Mountains to the west (U.S. Geological Survey, 2004). Based on known past, current, and planned uranium milling operations, Figure 3.2-2 shows that these operations are concentrated in two major uranium districts known as the Crooks Gap area in the Great Divide Basin straddling northeastern Sweetwater County and southeastern Fremont County and the Gas Hills area in the Wind River Basin located in eastern Fremont County (see details in the Geology and Soils Section 3.2.3).

The land ownership and use statistics for the Wyoming West Uranium Milling Region shown in Table 3.2-1, were calculated using the Geographic Information System (GIS) used to prepare the map shown in Figure 3.2-1. The majority of the land of the four counties of this region is composed of federal land (66 percent) and Native American land (9 percent) (Table 3.2-1). Private lands, intermixed with BLM land, occupy approximately 25 percent of the region. The eastern tips of the Shoshone and Bridger National Forests form a very small part on the western edge of this region (1 percent). A portion of the Wind River Indian Reservation and land administered by the United States Bureau of Reclamation represent approximately 13 percent of the land at the northwestern corner of the Wyoming West Uranium Milling Region. Riverton, located in this corner, is the largest town of the region with almost 10,000 inhabitants (Figure 3.2-1). Riverton is located more than 80 km [50 mi] from the Crooks Gap area and the Gas Hills area. Towns in the vicinity of these two uranium districts include Jeffrey City, Sand Draw, and Bairoil, each of which has a population of a few hundred or less (Figure 3.2-2).


As shown on Figure 3.2-1, BLM manages the vast majority of the land in the Crooks Gap and the Gas Hills areas. The land is mostly used as rangeland for cattle and sheep grazing under the BLM permit system.

BLM Grazing Permit/ License/Lease

BLM grants official written permission to private permittees or lessees to allow a certain number, type and class of their livestock graze on public lands for a specified time period and on a defined rangeland.



WYOMING WEST REGION



- ▲ Ur Milling Site (NRC)
- ▭ Wyoming West Milling Region
- ▭ Wyoming East Milling Region
- Major City
- Interstate Highway
- US Highway

- State Highway
- Railroad
- Water bodies (Lakes, Bays, ...)
- Rivers and Streams
- ▭ Counties



Federal Lands

- ▨ Forest Service
- ▨ Bureau of Land Management
- ▨ Fish and Wildlife Service
- ▨ Bureau of Indian Affairs
- ▨ Bureau of Reclamation

Figure 3.2-1. Wyoming West Uranium Milling Region General Map With Current and Future Uranium Milling Site



WYOMING WEST REGION



Cities by population




Figure 3.2-2. Map Showing Outline of the Wyoming West Uranium Milling Region and Locations of the Crooks Gap Uranium District in the Great Divide Basin and the Gas Hills Uranium District in the Wind River Basin

Table 3.2-1. Land Ownership and General Use in the Wyoming West Uranium Milling Region

Land Ownership and General Use	Area (mi²)	Area (km²)	Percent
U.S. Bureau of Land Management, Public Domain Land	5,476	14,184	61.4
Private Lands	2,191	5,675	24.6
Bureau of Indian Affairs, Indian Reservatons	809	2,095	9.1
Bureau of Reclamation	352	911	3.9
U.S. Forest Service, National Forest	87	226	1
Totals	8,915	23,090	100.0

Most of the private land in the eastern and southern part of the region is intermixed with BLM grazing land, and is used to produce hay for feeding cattle in winter. Other scattered land uses in this region include wildlife habitat, wilderness areas, hunting, dispersed recreation and off-road vehicle (ORV) use, oil and gas recovery, gas and carbon dioxide pipelines and transmission lines, and cultural and historical sites, such as the Oregon/Mormon Pioneer National Historic Trail (BLM, 1987, 2007e). The presence and extent of these land uses will have to be addressed on a site-specific basis at, and in the vicinity of, any new potential uranium milling facility.

3.2.2 Transportation

Past experience at NRC licensed ISL facilities indicate these facilities rely on roads for transportation of goods and personnel (Section 2.8). As shown on Figure 3.2-3, the Wyoming West Uranium Milling Region is accessible by Interstate 80, which borders the south of the region between Rock Springs and Rawlins. The Wyoming West Uranium Milling Region is also accessed from the west by State Highway 28, from the northwest by U.S. Highway 26, from the north by U.S. Highway 20, and from the east by U.S. Highways 20 and State Route 220. Rail lines traverse the northern and southern portions of the region.

Areas of past, present, or future interest in uranium milling in the region are also shown in Figure 3.2-3. These areas are located in four main subregions when considering site access by local roads. Areas of milling interest that are located in the northeastern part of the region near the Natrona County and Fremont County border are accessible by State Route 136 from Riverton or by a local access road that travels south from Waltman until intersection with State Route 136. Another area of milling interest is in the central portion of the milling region adjacent to State Route 135, which is accessed from the north from Riverton or from the south from U.S. Highway 789. Traveling east from that point on U.S. Highway 789 to Jeffrey City is another area of milling interest. Other sites of interest in the southeastern portion of the Wyoming West Uranium Milling area (Great Divide Basin Area in Sweetwater County) are accessible by unpaved local access roads that extend west from U.S. Highway 287 at Bairoil and a location further south between Bairoil and Rawlins. These west trending roads intersect a north and south trending unpaved road that connects Wamsutter on the southern border of the region at Interstate 80 to Jeffrey City and Moneta to the north. U.S. Highway 287 continues south to Interstate 80.




Figure 3.2-3. Wyoming West Uranium Milling Region Transportation Corridor

Description of the Affected Environment

Table 3.2-2 provides available traffic count data for roads that support areas of past or future milling interest in the Wyoming West Uranium Milling Region. Counts are variable with the minimum all vehicle count at 130 vehicles per day on State Route 136 to Riverton and the maximum on U.S. Highway 20 from Riverton to Shoshoni at 19,620 vehicles per day. Most all vehicle counts in the Wyoming West Uranium Milling Region are above 800 vehicles per day.

Yellowcake product shipments are expected to go from the milling facility to a uranium hexafluoride production (conversion) facility in Metropolis, Illinois (the only facility currently licensed by NRC in the U.S. for this purpose). Major interstate transportation routes are expected to be used for these shipments, which are required to follow NRC packaging and transportation regulations in 10 CFR Part 71 and U.S. Department of Transportation hazardous material transportation regulations at 49 CFR Parts 171—189.

Table 3.2-2. Average Annual Daily Traffic Counts for Roads in the Wyoming West Uranium Milling Region*

Road Segment	Distance (mi)	Trucks		All Vehicles	
		2005	2006	2005	2006
State Route 136 to Riverton	44	10–20	20–30	130–260	200–270
State Route 135 from State Route 136 to State Route 789	1.04	170	210	840	1,090
State Route 789 from State Route 135 to U.S. Highway 26	1	570–650	570–650	11,500–17,000	11,650–17,100
U.S. Highway 20/26 from Riverton to Shoshoni	22	520–650	520–650	3,340–19,580	5,100–19,620
U.S. Highway 20/26 from Shoshoni to Waltman	51	270–580	470–550	2,350–3,090	2,190–3,060
U.S. Highway 20/26 from Waltman to Casper	49	470–670	480–650	2,480–13,740	2,450–13,580
Interstate 25 from Casper to State Route 95	21	570–1,030	610–1,030	2,610–10,220	2,710–10,220
U.S. Highway 287 (State Route 789) at Lander South	-	390	400	5,080	4,550
U.S. Highway 287 (State Route 789) at Jeffrey City	-	140	140	850	890
U.S. Highway 287 at Muddy Gap	-	140	140	910	910
State Route 220 at Muddy Gap North	-	620	620	1910	1910
State Route 73 from Bairoil to Lamont	4.64	30	30	230	230
U.S. Highway 287 from Lamont to Muddy Gap	11	700	690	2,400	2,400
*Wyoming Department of Transportation. "Wyoming Department of Transportation Vehicle Miles." Data for Calendar Year 2005 and 2006 Provided on Request. District 2 Office, Casper, Wyoming: Wyoming Department of Transportation. April 18, 2008.					

Table 3.2-3 describes representative routes and distances for shipments of Yellowcake from locations of Uranium milling interest in the Wyoming West Uranium Milling Region. Representative routes are considered owing to the number of routing options available that could be used by a future ISL facility.

Table 3.2-3. Representative Transportation Routes for Yellowcake Shipments From the Wyoming West Uranium Milling Region*

Origin	Destination	Major Links	Distance (mi)
South of Moneta, Wyoming	Metropolis, Illinois	Local access road to Waltman, Wyoming U.S. Highway 20 east to Casper, Wyoming Interstate 25 south to Denver, Colorado Interstate 70 east to St. Louis, Missouri Interstate 64 east to Interstate 57 Interstate 57 south to Interstate 24 Interstate 24 south to U.S. Highway 45 U.S. Highway 45 west to Metropolis, Illinois	1,390
Sand Draw, Wyoming	Metropolis, Illinois	Local access roads to State Route 135 State Route 135 south to U.S. Highway 287 U.S. Highway 287 south to Interstate 80 Interstate 80 east to Cheyenne, Wyoming Interstate 25 south to Metropolis, Illinois (as above)	1,400
Jeffrey City, Wyoming	Metropolis, Illinois	Local access roads to U.S. Highway 287 U.S. Highway 287 to Interstate 80 Interstate 80 east to Cheyenne, Wyoming Interstate 25 south to Metropolis, Illinois (as above)	1,360
Great Divide Basin Area, Wyoming	Metropolis, Illinois	Local access road south to Wamsutter Interstate 80 east to Cheyenne, Wyoming Interstate 25 south to Metropolis, Illinois (as above)	1,360

*American Map Corporation. "Road Atlas of the United States, Canada, and Mexico." Long Island City, New York: American Map Corporation. p. 144. 2006.

3.2.3 Geology and Soils

Wyoming contains the largest known reserves of uranium in the United States and has been the nation's leading producer of uranium ore since 1995 (Wyoming State Geological Survey, 2005). Sandstone-hosted uranium deposits account for the vast majority of the ore produced in Wyoming (Chenoweth, 1991). In the Wyoming West Uranium Milling Region, uranium mineralization is found in fluvial sandstones in two major uranium districts: the Crooks Gap area of the Great Divide Basin and the Gas Hills area of the Wind River Basin (Figure 3.2-2). The uranium mineralization in the sandstone-hosted deposits in the Crooks Gap and Gas Hills areas is amenable to recovery by ISL milling. Since 1991, all uranium produced from sandstones in these two districts has been by the ISL method (Wyoming State Geological Survey, 2005).

The Crooks Gap area is located in Fremont and Sweetwater Counties and encompasses approximately 9,100 km² [3,500 mi²] in south-central Wyoming (Bailey, 1969; Rackley, 1972;

Description of the Affected Environment

Boberg, 1981). In 1954, ore-grade mineralization was found at Crooks Gap, and by late 1957, 3,800 metric tons [4,200 tons] of ore had been mined, mostly from shallow workings (Bailey, 1969). Production plus minable reserves at Crooks Gap are estimated to be between 5,000 and 5,400 metric tons [5,500 and 6,000 tons] U_3O_8 .

The Gas Hills uranium district is located along the southeastern margin of the Wind River Basin in central Wyoming (Anderson, 1969; Rackley, 1972; Boberg, 1981). Uranium in the Gas Hills district was discovered in 1953, and ore production began in 1955 (Anderson, 1969). The mineralized ground encompasses an area of about 160 km² [100 mi²]. Prior to 1968, the Gas Hills uranium district produced approximately 26 million metric tons [29 million tons] of U_3O_8 , which accounted for about 12 percent of total uranium production in the United States (Chenoweth, 1991).

The dominant source of sediment in the Great Divide Basin and the Wind River Basin was Precambrian (greater than 453 million-year-old) granitic rock of the Sweetwater Arch (Rackley, 1972) (Figure 3.2-4). The Sweetwater Arch is also referred to as the Granite Mountains (Bailey, 1969; Anderson, 1969; Lageson and Spearing, 1988). The Sweetwater Arch is a large mass of granitic rock 140 km [87 mi] long, with a maximum width of 50 km [31 mi]. Uplift of the Sweetwater Arch began to affect sedimentation in the adjacent Great Divide Basin and Wind River Basin in Late Cretaceous time (65 to 99 million years ago). Rapidly subsiding portions of these basins received thick clastic wedges (i.e., wedges made up of fragments of other rock) of predominantly arkosic sediments (i.e., sediments containing a significant fraction of feldspar), while larger, more slowly subsiding portions of the basins received a greater proportion of paludal (marsh) and lacustrine (lake) sediments.

Sediment transported southward into the Great Divide Basin was deposited on an apron of alluvial fans (Rackley, 1972). One of the major fans is centered near the Crooks Gap milling district, and another is northwest of the Lost Soldier anticline. Sedimentation in the Gas Hills area of the Wind River Basin was on an alluvial (i.e., deposited by running water) fan in which ridges of older resistant rock protruded through the fan and controlled the movement of the streams and their pattern of deposition. Beginning in the middle Eocene (41 to 49 million years ago) and increasing in the Oligocene (23.8 to 33.7 million years ago), regional volcanic activity contributed a significant amount of tuffaceous materials (i.e., materials made from volcanic rock and mineral fragments in a volcanic ash matrix) to local sediments. Deposition within the basins probably continued through the Miocene (5.3 to 23.8 million years ago), but post-Miocene erosion has completely removed Oligocene and Miocene units.

A generalized stratigraphic section of Tertiary (1.8 to 65-million-year-old) formations in the Wyoming West Uranium Milling Region is shown in Figure 3.2-5. Stratigraphic descriptions presented here are limited to formations that may be involved in potential milling operations or formations that may have environmental significance, such as important aquifers and confining units above and below potential milling zones.

Formations hosting major sandstone-type uranium deposits in the Wyoming West Uranium Milling Region are the Wind River Formation in the Wind River Basin and the Bottle Springs Formation in the Great Divide Basin. Both the Wind River and Bottle Springs are lower Eocene (49 to 54.8 million years old) in age (Houston, 1969) and consist of interbedded, arkosic

1




Figure 3.2-4. Index and Structure Map of Central Wyoming Showing Relation of Sweetwater Arch to the Great Divide Basin and the Wind River Basin. The Distribution of Arkosic, Carbonaceous Sediments and Mineralized Areas in the Crooks Gap and Gas Hills Uranium Districts Are also Shown (Modified From Rackley, 1972).

1

Central Wyoming			
System	Series		Formation
Tertiary	Pliocene		Moonstone Formation
	Miocene		Browns Park Formation Split Rock Formation
	Oligocene		White River Formation
	Eocene	Upper	Wagon Bed Formation
		Middle	
		Lower	Battle Springs Formation Wind River Formation
	Paleocene		Fort Union Formation
Cretaceous	Upper		Lance Formation

Figure 3.2-5. Stratigraphic Section of Tertiary Age Formations in the Great Divide Basin and Wind River Basin of Central Wyoming. Major Sandstone-Type Uranium Deposits Are Hosted in the Battle Springs Formation in the Great Divide Basin and the Wind River Formation in the Wind River Basin (Modified From Harshman, 1968).

sandstone; conglomerate; siltstone; mudstone; and carbonaceous shale—all compacted but poorly cemented (Harshman, 1968). The source beds for uranium deposits are sandstones interstratified with lensing mudstones and shales (Anderson, 1969). The mineralized zone in the Battle Springs Formation at Crooks Gap occurs in a stratigraphic range of as much as 460 m [1,500 ft] {i.e., occurs in a zone up to 460 m [1,500 ft] thick} (Stephens, 1964). In the Gas Hills district, mineralization in the Wind River Formation occurs in a stratigraphic range of perhaps 150 m [500 ft] (Bailey, 1969).

The Wagon Bed Formation conformably overlies the Wind River and Bottle Springs formations. The Wagon Bed is composed of a series of interbedded arkosic sandstones and silicified claystones. Regionally, the Wagon Bed Formation may not be present in the central parts of the basins, having been removed by erosion. The White River Formation unconformably overlies the Wagon Bed Formation or the Wind River and Bottle Springs formations where the Wagon Bed has been removed by erosion. The White River consists of tuffaceous siltstone, claystone, and conglomerate with subordinate amounts of tuff. The White River overlaps older Tertiary formations and wedges out against pre-Tertiary rocks on the flanks of the basins. The White River Formation is overlain by the Browns Park Formation in the Great Divide Basin and the Split Rock Formation in the Wind River Basin. The Browns Park and Split Rock consist of tuffaceous siltstone and sandstone beds that sometimes cap prominent ridges (Harshman, 1968).

The Fort Union Formation underlies the Wind River and Bottle Springs formations and, to a limited extent, is also a host of sandstone-type uranium deposits (Davis, 1969; Langden, 1973). The Fort Union is a fluvial deposit consisting of alternating and discontinuous mudstones, siltstones, carbonaceous shales, and coarser arkosic sandstone. The Fort Union is unconformably underlain by sediments of the Lance Formation, which is in turn underlain by a thick sequence of older sandstones, mudstones, and shales.

The uranium deposits in the Wyoming West Uranium Milling Region are genetically related to geochemical interfaces or roll-fronts (see Section 3.1.1). Principal ore minerals at Crooks Gap are meta-autunite, uraninite, and coffinite. The uranium minerals occur as earthy brown to black coatings on and interstitial fillings between quartz sand grains. In the Gas Hills district, roll-fronts can be followed for long distances and individual ore bodies are found along them that may reach thousands of feet in length.

The source of uranium in sandstone roll-front deposits in central Wyoming is a topic of conjecture. Four theories on the source of uranium in these occurrences have been suggested: (1) leached uranium from overlying ash-fall tuffs; (2) leached uranium from igneous and metamorphic rocks in the highlands surrounding the basins; (3) leached uranium from the host sandstones themselves; and (4) hydrothermal uranium from a magma source at depth (Harris and King, 1993). Combinations of these theories have been proposed as well (Boberg, 1981). The most popular theories are the tuff leach (1) and the highland leach (2). The tuff leach theory is supported by extensive geochemical studies on uranium removal from tuff (Zielinski, 1983, 1984; Trentham and Orajaka, 1986). Further, it was the tuff leach theory that led to the discovery of most of the large uranium deposits in Wyoming (Love, 1952). On the other hand, many sandstone-hosted uranium deposits in Wyoming are found adjacent to crystalline rocks, especially the uraniferous granites of the northern Laramie and Granite mountains (Harris and King, 1993). Oxidized uranium leached from these crystalline terrains could have been transported to the sites of present mineralization.

Soils within the Wyoming West Uranium Milling Region are diverse and can vary substantially over relatively short distances. The distribution and occurrence of soils in central Wyoming can vary both on a regional basis (mountains, foothills, basins) and locally with changes in slope, geology, vegetation, climate, and time. The Great Divide Basin and the Wind River Basin present a mixture of old, tilted sedimentary rocks that often occur in bands along the basin margins and younger sediments showing varying degrees of incision by erosion in basin centers.

The topographic position and texture of typical soils in the Great Divide Basin and Wind River Basin areas of central Wyoming were obtained from the Soils Map of Wyoming (Munn and Arneson, 1998). This map was designed primarily for statewide study of groundwater vulnerability to contamination and would not be expected to be used for site-specific soil interpretations at proposed ISL milling facilities. For site-specific evaluations, detailed soils information would be expected to be obtained from published county soil surveys or the U.S. Department of Agriculture Natural Resource Conservation Service.

In the Great Divide and Wind River basin areas, loamy-skeletal soils (rocky soils) with little or no subsoil development occur along bedrock outcrops that form ridges along the flanks of the basins. On gently sloping to moderately steep slopes associated with ridge flanks, alluvial fans, and alluvial terraces, fine to fine-loamy soils with well-developed horizons of clay accumulation are found. These soils are generally light-colored and depleted in moisture. Moderately deep fine-loamy over sandy and coarse loamy soils with well-developed soil horizons occur on

terraces along major streams. Soils found on floodplains and drainageways include clay loams and fine sand loams. Dark-colored, base-rich soils formed under grass are generally associated with floodplains along streams with permanent high-water tables. These soils are generally very deep and have well-developed soil horizons.

3.2.4 Water Resources

Water resources of the Wyoming West Uranium Milling Region are described in terms of surface waters, wetlands and waters of the United States, and groundwater.

3.2.4.1 Surface Waters

The Wyoming West Uranium Milling Region (Figure 3.2.-1) includes major portions of Fremont and Sweetwater counties and small portions of Carbon and Natrona Counties. The watersheds within the Wyoming West Uranium Milling Region are listed in Table 3.2-4 along with the range of designated uses of surface water bodies assigned by the State of Wyoming (WDEQ, 2001). Because surface water uses are designated for specific water bodies, such as stream segments and lakes, within a watershed and the specific locations of future uranium milling activities are not known at this time, the range of designated uses is provided rather than a listing of designated uses for each water body within a watershed. Not all water bodies within a watershed may have all of the designated uses listed in Table 3.2-4. For information regarding specific water bodies, the reader is referred to the Wyoming Department of Environmental Quality Surface Water Standards webpage deq.state.wy.us/wqd/watershed/surfacestandards.

The historical uranium milling districts included in the Wyoming West Uranium Milling Region are called Gas Hills in the east-central portion of the Wyoming West Uranium Milling Region, and Crooks Gap near the Fremont-Sweetwater county line (Figure 3.2-2). Watersheds in the Wyoming West Uranium Milling Region are: Great Divide Closed Basin, Sweetwater River, Muskrat Creek, Little Wind River, Popo Agie River, Lower Wind River, Badwater Creek, and their associated tributaries. Historical or potential uranium milling sites are present in the Great Divide, Sweetwater River, Muskrat Creek, Littlewind River, and Lower Wind River watersheds (Figure 3.2-6).

The Great Divide Closed Basin is an area with internal drainage and no outlet to either the Atlantic or Pacific oceans located in northeastern Sweetwater County and western Carbon County (Figure 3.2-6). Surface water flows from the upland areas on the perimeter of the basin toward playa lakes near the center of the basin. The State of Wyoming has assigned surface classifications to streams in this watershed ranging from 2AB to 4C (WDEQ, 2001). Most of the streams are classified as 3A or 3B. The attainment status of these streams has not been assessed. The Crooks Gap Uranium District is partly located within the Great Divide Closed Basin.

The Sweetwater River watershed is located north of the Great Divide Closed Basin watershed in Sweetwater County. The Sweetwater River is a Class 1 water above Alkali Creek and Class 2AB water below Alkali Creek (Table 3.2-4). Crooks Creek is reported to be impaired due to oil and

Attainment Status

The attainment status of a water body refers to whether or not its water quality meets the standards for its designated use. The designated use of a water body is assigned by the state, such as swimming, drinking, and protection and propagation of aquatic life. If the chemical pollutants or other water quality parameters, such as temperature or turbidity, exceed the standards for its designated use, the attainment status of the water body is described as impaired.

grease from oil and natural gas production (WDEQ, 2006). The average flow in the Sweetwater River near Alcova, Wyoming is 1.1 m³/s [40 ft³/s] (U.S. Geological Survey, 2008). The Crooks Gap uranium district is within the Sweetwater River watershed and is drained primarily by Crooks Creek and its tributaries. Topographic maps of the area show a number of unnamed springs and small impoundments on the ephemeral streams within the district.

**Table 3.2-4. Primary Watersheds in the Wyoming West Uranium Milling Region
Range of Designated Uses of Water Bodies Within Each Watershed**

Watershed	Range of State Classification of Designated Uses*
Great Divide Closed Basin	2AB to 4C
Sweetwater River and Tributaries	1 (above Alkali Creek), 2AB (below Alkali Creek)
Muskrat Creek	2AB, 2C
Little Wind River	2AB
Popo Agie River	2AB
Lower Wind River	Generally 2AB with some tributaries 3B
Badwater Creek	Generally 2AB with some tributaries 3B and 4B
*Class 1 waters have designated uses including: Drinking Water, Game Fish, Non-Game Fish, Fish Consumption, Other Aquatic Life, Recreation, Wildlife Agriculture, Industry, Scenic Value. Class 2AB waters have designated uses including: Drinking Water, Game Fish, Non-Game Fish, Fish Consumption, Other Aquatic Life, Recreation, Wildlife Agriculture, Industry, Scenic Value. Class 2A waters have designated uses including: Drinking Water, Other Aquatic Life, Recreation, Wildlife Agriculture, Industry, Scenic Value. Class 2B waters exclude drinking water from the Class 2AB uses. Class 2C waters exclude drinking water and game fish from the Class 2AB uses. Class 3A, 3B and 3C waters have designated uses including: Other Aquatic Life, Recreation, Wildlife Agriculture, Industry, Scenic Value. Class 4A, 4B and 4C waters have designated uses include: Recreation, Wildlife Agriculture, Industry, Scenic Value.	

The Muskrat Creek watershed is located north of the Sweetwater River watershed in Fremont County. Classifications of water bodies in the Muskrat Creek watershed range from 2AB to 2C (Table 3.2-4). No data are available on average flow in Muskrat Creek. The Gas Hills uranium district is within the Muskrat Creek watershed which drains to the Wind River and ultimately to the Powder River (Figure 3.2-5). Muskrat Creek is ephemeral within the Gas Hills uranium district. The Gas Hills district is also drained by a number of other ephemeral stream channels with small surface water impoundments. Mapped springs in the district are Puddle Spring and Willow Spring.

The Little Wind River watershed is located west of the Muskrat Creek watershed and roughly centered on Riverton, Wyoming. The Little Wind River is classified as 2AB (Table 3.2-4). The average flow of the Little Wind River at Riverton is 6 m³/s [215 ft³/s] (U.S. Geological Survey, 2008).

The Popo Agie River watershed is located west of the Little Wind River watershed on the eastern flank of the Wind River Mountains in Fremont County. The Popo Agie River is classified as 2AB (Table 3.2-4). The average flow of the Popo Agie River between 1947 and 1971 was 2.3 m³/s [80 ft³/s] (U.S. Geological Survey, 2008). No historical uranium mining or milling has occurred within the Popo Agie watershed.




Figure 3.2-6. Watersheds in the Wyoming West Uranium Milling Region

The Lower Wind River watershed is located north and downstream of the Little Wind River water shed. Water bodies in the Lower Wind River watershed are generally classified as 2AB with some tributaries classified as 3B, the difference being that 3B waters are not designated as sources of drinking water or for fishing or fish consumption (Table 3.2-4). Lower Muddy Creek and Lower Poison Creek are described as impaired due to fecal coliform (WDEQ, 2006). The average flow of the Wind River below Boysen Reservoir is 29.5 m³/s [1,040 ft³/s] (U.S. Geological Survey, 2008).

The Badwater Creek watershed is located on the northern edge of the Wyoming West Uranium Milling Region northeast of the Muskrat Creek watershed. Water bodies in the Badwater Creek watershed are generally classified as 2AB with some tributaries classed as 3B and 4B. The difference between 3B and 4B waters is that 4B waters do not have "other aquatic life" as a designated use (Table 3.2-4). No data are available on average flow in Badwater Creek.

3.2.4.2 Wetlands and Waters of the United States

The regulatory program of the U.S. Army Corps of Engineers (USACE) plays a critical role in the protection of the aquatic ecosystem and navigation. Under Section 404 of the Clean Water Act and Section 10 of the Rivers and Harbors Act of 1899, the USACE performs the following services:

- Conducts jurisdictional determinations for wetlands and other waters of the United States and navigable waters of the United States
- Authorizes activities in these jurisdictional areas through individual and general permits
- Ensures compliance of issued permits
- Enforces requirements of the law for unpermitted activities

Under Section 404 of the Clean Water Act, the Secretary of the Army is responsible for administering a regulatory program that requires permits to discharge dredged or fill material into waters of the United States, including wetlands.

Areas regulated under Section 404 are collectively referred to as "Waters of the United States." Included are parts of the surface water tributary system down to the smallest streams; lakes, ponds, or other water bodies on those streams; and adjacent wetlands.

Isolated waters such as playa lakes, prairie potholes, old river scars, cutoff sloughs, and abandoned construction and milling pits may also be waters of the United States if they meet certain criteria. Wetlands are found in many different forms including bottomland hardwood forests, wooded swamps, marshes, wet meadows, bogs, and playa lakes. Wetlands are of particular concern because they are valuable to restoring and maintaining the quality of the waters of the United States. Their functions include sediment trapping, nutrient removal, chemical detoxification, shoreline stabilization, aquatic food chain support, fish and wildlife habitat, floodwater storage, and groundwater recharge.

According to the USACE Wetland Delineation Manual (USACE, 1987), wetlands are defined as "those areas that are inundated or saturated by surface or groundwater at a frequency and duration sufficient to support, and that under normal circumstances do support, a prevalence of

Description of the Affected Environment

vegetation typically adapted for life in saturated soil conditions. Wetlands generally include swamps, marshes, bogs, and similar areas.” A minimum of one positive indicator from each parameter (vegetation, hydrology, and soils) must be found to make a positive wetland determination.

- **Vegetation**—Under normal circumstances, an area is considered to have hydrophytic vegetation when more than 50 percent of dominant species, from all plant strata, are classified as Obligate (OBL), Facultative wet (FACW), or Facultative (FAC). Plants listed as Facultative Upland (FACUP), Not Listed (NL), or No Indicator (NI) are considered nonwetland plants for the purposes of wetland delineations.
- **Hydrology**—USACE (1987) requires that wetland soils must be continually saturated for a prolonged period (at least 5 percent) during the growing season.
- **Soils**—Hydric soils are those that are saturated, flooded, or ponded long enough during the growing season to develop anaerobic conditions in their upper parts. Typical field indicators of hydric soils are the presence of thick organic layers, or in the case of predominantly mineral soils, a low chroma matrix (gray color) and/or bright mottling.

Man-made ponds and other surface features not immediately adjacent to traditional navigable waters do not fall under the jurisdiction of the USACE. The landward regulatory limit for waters (in the absence of adjacent wetlands) is the ordinary high water mark. The ordinary high water mark is the line on the shores established by the fluctuations of water and indicated by physical characteristics such as

- A clear natural line impressed on the bank
- Shelving
- Changes in the character of the soil
- Destruction of terrestrial vegetation
- The presence of litter and debris
- Other appropriate means that consider the characteristics of the surrounding areas

Waters of the United States and special aquatic sites that include wetlands would need to be identified and the impact delineated upon individual site selection for a potential ISL facility. Based on impacts and consultation with each area, appropriate permit would need to be obtained from the local USACE district. Under Section 401 of the Clean Water Act, state water quality certification is required for work

According to the U.S. Fish and Wildlife Wetland Mapper (2007), numerous types of wetlands and waters located within the region:

- **Perennial Streams**—A perennial stream has flowing water year-round during a typical year. The water table is located above the stream bed for most of the year. Groundwater is the primary source of water for stream flow. Runoff from rainfall is a supplemental source of water for stream flow (USACE, 2000).
- **Intermittent Streams**—An intermittent stream has flowing water during certain times of the year, when groundwater provides water for stream flow. During dry periods, intermittent streams may not have flowing water. Runoff from rainfall is a supplemental source of water for stream flow (USACE, 2000).
- **Ephemeral Streams/Arroyos** (term used in arid regions)—An ephemeral stream has flowing water only during, and for a short duration after, precipitation events in a typical year. Ephemeral stream beds are located above the water table year round. Groundwater is not a source of water for the stream. Runoff from rainfall is the primary source of water for stream flow (USACE, 2000).

in waters of the United States. Within this region, the state of Wyoming regulates isolated wetlands and waters. Cumulative total project impacts greater than 0.04 ha [1 acre] require a general permit for wetlands mitigation by the Wyoming Department of Environmental Quality (WDEQ).

The majority of wetland areas located within the region consist of fresh water, ponds, emergent, or ponds with floating or submerged aquatic vegetation. These wetland areas are typically temporarily flooded on a seasonal basis. Numerous intermittent streams that are temporarily flooded are also found in the Wyoming West Uranium Milling Region.

3.2.4.3 Groundwater

Groundwater resources in the Wyoming West Uranium Milling Region are part of regional aquifer systems that extend well beyond the areas of uranium milling interest in this part of Wyoming. Uranium bearing aquifers exist within these regional aquifer systems in the Wyoming West Uranium Milling Region. This section provides a general overview of the regional aquifer systems to provide context for a more focused discussion of the uranium bearing aquifers in the Wyoming West Uranium Milling Region, including hydrologic characteristics, level of confinement, groundwater quality, water uses, and important surrounding aquifers.

3.2.4.3.1 Regional Aquifer Systems

The location of the Wyoming West Uranium Milling Region is shown in Figures 3.2-1 and 3.2-2. The Upper Colorado River Basin aquifer system is the major regional aquifer system (large-scale underground layer of water-bearing permeable rock or unconsolidated materials) in the Wyoming West Uranium Milling region. The Upper Colorado River Basin aquifer system extends over 51,800 km² [20,000 mi²] in the Green River, the Great Divide, and the Washakie structural basins in the southwestern parts of Wyoming (Whitehead, 1996).

Groundwater in the Upper Colorado River Basin aquifer system flows from aquifer recharge areas toward the centers of the structural basins. Discharge from the aquifers is by upward leakage to shallower aquifers and to major streams. Groundwater is less than 61 m [200 ft] below the land surface in most parts of the aquifer system and is nearest the land surface near the major streams. In and near mountainous areas, depth to groundwater ranges from 152 to 305 m [500 to 1,000 ft].

The Upper Colorado River Basin aquifer system in southwestern Wyoming consists of layered sedimentary formations. Whitehead (1996) grouped the sedimentary formations into five principal aquifers. From shallowest to deepest, they are the Laney aquifer, the Wasatch-Fort Union aquifer, the Mesaverde aquifer, a series of sandstone aquifers from the Dakota Sandstone through the Nugget Sandstone aquifers, and the Paleozoic aquifers.

The uppermost aquifer in the Wyoming part of the Upper Colorado River Basin aquifer system is the Laney aquifer. It is the highest permeability member of the Green River Formation. This aquifer consists of fractured sandstone beds and yield sufficient water for domestic and livestock watering supplies. Water in the Laney aquifer is fresh to slightly saline.

The Wasatch-Fort Union aquifer (that includes the Wasatch Formation and the Fort Union Formation) is composed of the major water-yielding sandstones interbedded with shale, mudstone, and some coal beds. The thickness of the Wasatch-Fort Union aquifer is notable and reported to be about 3,350 m [11,000 ft] thick in Sublette County and about 2,135 m

Description of the Affected Environment

[7,000 ft] thick near the center of the Great Divide Basin in south-central Wyoming. The regional groundwater flow direction in the eastern part of the aquifer is from recharge areas at basin margins toward the Great Divide Basin and southward into Colorado toward the center of the Washakie Basin. In the western part of the aquifer, water flows from recharge areas toward the Green River and its tributaries and toward the Flaming Gorge Reservoir in South Wyoming. Most of the fresh water in the Upper Colorado River Basin aquifer system is in the Wasatch-Fort Union aquifer, but the aquifer locally, where it is deeply buried, and contains saline water. The Green River Formation overlies the Wasatch-Fort Union aquifer and forms an effective confining unit in most places.

The Mesaverde aquifer is composed of sandstone beds. In most places, the Mesaverde aquifer and the Wasatch-Fort Union aquifer are hydraulically connected. However, the Lewis Shale locally overlies the Mesaverde aquifer in the Great Divide and the Washakie Basins. The Mesaverde aquifer crops out at the land surface surrounding the Rock Springs Uplift. The groundwater flow direction in the Mesaverde aquifer is from recharge areas at the Rock Springs Uplift and near the eastern limit of the aquifer system toward the centers of structural basins. The aquifer contains fresh water locally at outcrop (recharge) areas, but it contains saline or brine water where the aquifer is deeply buried (e.g., in the Washakie Basin in southwestern Wyoming). The Mesaverde aquifer is hydraulically separated from deeper aquifers in Mesozoic rocks through thick confining layers that consist primarily of shale.

The Dakota and the Nugget aquifers consist of several sandstone formations separated by confining units. These aquifers crop out only locally in southwestern Wyoming and contain very saline water or brine in most places. A thick confining unit of Triassic- and Permian-aged rocks hydraulically separates them from the deeper Paleozoic aquifers.

The Tensleep Sandstone and the Madison Limestone are the principal aquifers in Paleozoic rocks. Groundwater in these aquifers flows toward the centers of the structural basins from adjacent topographically high areas. Groundwater discharges from the Tensleep Sandstone to the shallower aquifers occur by upward leakage. Much of the discharge from the Madison Limestone occurs by lateral movement of the ground water into adjacent structural basins to the southeast and northeast. Because the Paleozoic aquifers are mostly deeply buried and contain saline water, they are not extensively used for water supply in southwestern Wyoming.

Recharge to the aquifers in most of the area is likely small, due to low annual precipitation and high evaporation rates (AATA International Inc., 2005). The mean annual precipitation in the Wyoming West Uranium Milling Region is typically in the range of 15-28 cm/yr [6-11 in/yr], but at high elevations, it locally exceeds 50 cm/yr [20 in/yr] based on precipitation data from 1971 to 2000. The evaporation rate was estimated to be 105.9 ± 7.1 cm/yr [41.7 ± 2.8 in/yr] using the Kohler-Nordenson-Fox equation at the station in Lander, Wyoming (Curtis and Grimes, 2004).

3.2.4.3.2 Aquifer Systems In The Vicinity Of Uranium Milling Sites

An underlying hydrogeological system in past and current areas of uranium milling interest in the Wyoming West Uranium Milling Region consists of a thick sequence of primarily sandstone aquifers and shale aquitards. Uranium-bearing sandstone aquifers in the Wind River Formation (equivalent to the Battle Springs Formation at the proposed Lost Creek site and to the Green River Formation at the regional scale) are important sources for water supplies in the milling region.

Areas of uranium milling interest in the southern parts of the Wyoming West Uranium Milling Region near the Great Divide Basin (Crooks Gap) are underlain, from shallowest to deepest, by sedimentary deposits and sandstone layers (Quaternary-aged), the Green River Formation, the Wasatch/Battle Springs Formation, the Fort Union Formation, and the Lance/Fox Hills Formation. This hydrogeological sequence is separated from the underlying Mesaverde Formation by the regionally continuous and low permeable Lewis Shale aquitard (AATA International Inc., 2005; Lost Creek ISR, LLC, 2007). All these Formations host sandstone aquifers.

Areas of uranium milling interest in the northern parts of the Wyoming West Uranium Milling Region near the Gas Hills is underlain by the Late Tertiary-aged Formation and deposits including the Split Rock, White River, and Wagon Bed Formations. Among these formations, the Split Rock Formation is the primary aquifer. This system is underlain by the Wind River Formation, the Fort Union Formation, and the Lance Formation. This sequence is underlain by a thick sequence of confined aquifers and aquitards. The most important underlying water supply aquifers involve the Cloverly aquifer, the Nugget Sandstone, and the Tensleep Sandstone (NRC, 2004).

3.2.4.3.3 Uranium-Bearing Aquifers

Uranium mineralization at locations of milling interest is typically hosted by the Early Tertiary-age confined sandstone aquifers in the Wyoming West Uranium Milling Region.

Confined sandstone beds in the Battle Springs Formation are the uranium bearing aquifers in the Great Divide Basin (south-central Wyoming) within the southern portion of the Wyoming West Uranium Milling Region (AATA International Inc., 2005). Similarly, the Wind River Formation in the northern parts of the Wyoming West Uranium Milling Region near the Gas Hills is the uranium-bearing aquifer. Uranium mineralization in the Gas Hills has been identified in six different sandstone layers in the Wind River Formation, which are named as 30, 40, 60, 70, and 80 Sands. In some areas, these sand layers are hydraulically separated by confining units including siltstone, clay, and shale beds, while in other areas they are hydraulically and stratigraphically connected (NRC, 2004).

For ISL operations to begin, portions of the uranium-bearing sandstone aquifers in the Battle Springs Formation and in the Wind River Formation in the Wyoming West Uranium Milling Region would need to be exempted by the Underground Injection Control (UIC) Program administered by WDEQ (Section 1.7.2.1) for the purposes of uranium recovery (NRC, 2004).

Hydrogeological characteristics: In the Wyoming West Uranium Milling Region, the production aquifer system typically consists of confined sandstone aquifers. Aquifer properties (e.g., transmissivity, thickness, storage coefficient) vary spatially in the region. Based on field test data at the Gas Hills and in the Great Divide Basin, transmissivity of the ore-bearing aquifers range from 0.01–90 m²/day [0.1 to

Hydrologic Terminology

Transmissivity: It is used to define the flow rate through the vertical section of an aquifer unit considering width and extending the full saturated height of an aquifer under unit hydraulic gradient. Transmissivity is a function of the aquifer's saturated thickness and hydraulic conductivity.

Storage Coefficient: It is used to characterize the capacity of an aquifer to release groundwater from storage in response to a decline in hydraulic head.

Hydraulic Conductivity: It is a measure of the capacity of a porous medium to transmit water. It is used to define the flow rate per unit cross sectional area of an aquifer under unit hydraulic gradient.

Description of the Affected Environment

1,000 ft²/day] in the region. For ISL operations to be practical, the hydraulic conductivity of the production aquifer must be large enough to allow reasonable water flow from injection to production wells. Hence, portions of the production aquifers with low hydraulic conductivities may not be amenable to uranium recovery using ISL techniques. The average storage coefficient of the ore-bearing aquifer is on the order of 10^{-4} , indicating the confined nature of the production aquifer (typical storage coefficients for confined aquifers range from 10^{-5} – 10^{-3} (Driscoll, 1986; p.68).

Sandstone aquifers in the Battle Springs Formation are typically confined at the Lost Soldier and Lost Creek areas. However, the Battle Springs Formation locally crops out in the region, and hence the formation becomes locally unconfined. The transmissivity of the aquifer ranges from 8,690 L/day/m to 24,800 L/day/m [700 gal/day/ft to 2,000 gal/day/ft] {9 – 25 m²/day [95 ft²/day to 270 ft²/day]} and the aquifer storage coefficient ranges from 3.0×10^{-4} to 8.0×10^{-4} (AATA International Inc., 2005; Lost Creek ISR, LLC, 2007). Lateral hydraulic gradients range from 0.05 at the Lost Soldier area to 0.0125 at the Lost Creek area, and range from 0.002 to 0.006 between these two sites (AATA International Inc., 2005). Hence, the lateral hydraulic gradients are an order of magnitude larger within the Lost Creek area and the Lost Soldier area than between these two sites. The maximum well yields from the uranium-bearing aquifers range from 760 to 3,780 L/day [200 to 1,000 gal/day] (AATA International Inc., 2005).

Groundwater levels in the shallow, intermediate, and deep monitoring wells in the uranium-bearing aquifer were 55 m [180 ft], 58 m [190 ft], and 64 m [210 ft] below the ground surface (AATA International Inc., 2005). These measurements indicate potential upward vertical flow within the Battle Springs Formation.

In the northern parts of the Wyoming West Uranium Milling Region, the uranium-bearing sandstone aquifers are typically confined as in the southern parts of the Wyoming West Uranium Milling Region. Transmissivity values in the uranium-bearing aquifers vary from 0.07 to 90 m²/day [0.7 to 965 ft²/day]. Aquifer storage coefficients vary in the range of 8.5×10^{-5} to 8.0×10^{-3} , with an average storage coefficient of 3.0×10^{-4} (NRC, 2004).

Level of confinement: The production aquifer is typically confined in the Wyoming West Uranium Milling Region; however, local unconfined conditions exist. The thickness of the confinement varies spatially.

At the regional scale, the thickness of the upper confinement of the Battle Springs Formation spatially varies. At the Lost Soldier and Lost Creek areas, the Battle Springs Formation is confined above by a 3–6 m [10–20 ft] thick Claystone unit (AATA International Inc., 2005). But, as noted previously, the Battle Springs Formation crops out over the northeastern portion of the Great Divide Basin, and hence locally unconfined conditions exist (Lost Creek ISR, LLC, 2007). The Battle Springs Formation is confined below by the continuous Lewis Shale at the local and regional scales. At the Lost Creek area, the Lewis Shale is up to 820 m [2,700 ft] thick (Lost Creek ISR, LLC, 2007). Thus, the sandstone aquifers in the Battle Springs Formation are confined at the Lost Soldier and Lost Creek areas. Aquitard vertical conductivity ranges from 1.2×10^{-3} to 2.2×10^{-3} m/day [4.0×10^{-3} to 7.3×10^{-3} ft/day] (AATA International Inc., 2005).

At the Gas Hills site, the production aquifers are typically confined. Five potential ISL sites are identified and the thickness of the confinement spatially varies with the location of the potential ISL sites. For example, at Mine Unit 1, the uranium-bearing 70 Sand is confined above and below by relatively thick, continuous, low permeability units of the Wind River Formation. At

Mine Unit 2, the 30, 50, 60, 70, and 80 Sands are typically separated by up to 6 m [20 ft] thick confining layers. At Mine Unit 3, the 30, 40, and 50 sands are separated by relatively thin (1.5 to 9 m [5 to 30 ft] thick) confining layers. At Mine Unit 4, a 3–12 m [10–40 ft] thick confining layer overlies the 80 Sand locally in some parts of the region while the 70 and 80 Sands are unconfined in other parts. The 60 Sand is locally confined above by a 3 to 6 m [10 to 20 ft] thick confining layer and the 50 Sand is typically underlain by a 1.5 to 9 m [5 to 30 ft] thick confining layer in the region. The 50 Sand at Mine 5 is confined above by a 4.5 to 12 m [15 to 40 ft] thick confining unit and confined below by a 6 to 12 m [20 to 40 ft] thick confining layer (NRC, 2004).

Groundwater quality: In some parts of the Wyoming West Uranium Milling Region, the total dissolved solids (TDS) levels in the uranium-bearing aquifers exceed the EPA's drinking water standards. The uranium and radium-226 concentrations in the uranium-bearing aquifers typically exceed their respective EPA Maximum Contaminant Levels.

Groundwater of the Battle Springs Formation is of bicarbonate-sulfate-calcium type or bicarbonate-calcium type. The TDS level ranges from 200 to 400 mg/L [200 to 400 ppm], which is below the EPA's Secondary Drinking Water Standard of 500 mg/L [500 ppm]. The quality of groundwater near the town of Bairoil meets drinking water quality standards for all chemical constituents except for the elevated uranium and radium-226 concentrations associated with the rollfront uranium deposits (AATA International Inc., 2005). Uranium and radium-226 concentrations typically exceed their respective EPA Maximum Contaminant Levels of 0.03 mg/L [0.03 ppm] and 5 pCi/L.

Groundwater from the Wind River Formation in the Gas Hills area is of calcium-sulfate and calcium-sodium-bicarbonate-sulfate type. The TDS level in the Wind River Formation is commonly higher (623 to 1,887 mg/L [623 to 1,887 ppm]) than in the Battle Springs Formation and exceeds the EPA's Secondary Drinking Water Standard. Similar to the Battle Springs Formation, both the uranium (0.04 mg/L [0.04 ppm on the average]) and radium-226 (5–50 pCi/L away from the ore zone) exceeds respective EPA Maximum Contaminant Levels (NRC, 2004).

Current groundwater uses: Groundwater withdrawn from the Battle Springs Formation is primarily used for public water supply and agricultural purposes of the Town of Bairoil (AATA International Inc., 2005). Groundwater use in the Gas Hills area is typically limited to livestock, wildlife watering and, to a lesser extent, industrial uses. In vicinity of the Gas Hills area, groundwater is not used for domestic and irrigation supplies (NRC, 2004). At the regional scale, the Laney aquifer also yields sufficient water for domestic and livestock watering (Whitehead, 1996).

3.2.4.3.4 Other Important Surrounding Aquifers for Water Supply

At the regional scale, the Laney aquifer, the Wasatch-Fort Union aquifer, the Mesaverde aquifer, the Dakota and the Nugget aquifers, and the Paleozoic aquifers are the important aquifers for water supply in the region (Whitehead, 1996). Among these aquifers, the Paleozoic aquifers are used less extensively, because they are mostly deeply buried and contain saline water. The Laney and the Wasatch-Fort Union aquifers are locally hydraulically connected. The Mesaverde aquifer is also locally hydraulically connected to the overlying Wasatch-Fort Union aquifer. However, in most places, these two aquifers are separated by the Lewis Shale at the regional scale.

At the Great Divide, the Battle Springs Formation interfingers with sandstone aquifers in the Wasatch Formation and the Green River Formation, and it is underlain by sandstone aquifers in the Fort Union Formation and Lance/Fox Hills Formation. The Fox Hill Formation is considered to be a minor aquifer, but the others are usually considered to be relatively important aquifers in the region (AATA International Inc., 2005). The Fort Union aquifer is largely undeveloped in the Lost Creek area, and the reported transmissivity values are typically less than 30 m²/day [325 ft²/day] (Collentine et al., 1981). The TDS levels in the Wasatch Formation in the west and south parts of the Great Divide Basin is typically higher than the U.S. EPA drinking water standards of 500 mg/L [500 ppm]. However, the TDS levels in the Battle Springs/Wasatch aquifers are generally less than 500 mg/L [500 ppm] along the northern side of the region (Lost Creek ISR, LLC, 2007).

In most parts of the Gas Hills area, the Wind River Formation is underlain by an aquitard that consists of the Chugwater (between the Nugget Sandstone and the Tensleep Sandstone) and Sundance Formations (between the Cloverly Formation and the Tensleep Sandstone). The other important aquifers, including the Cloverly Formation (equivalent to the Dakota Sandstone), Nugget Sandstone and Pennsylvanian Tensleep Sandstone, are separated from the Wind River Formation by a series of thick aquitards.

3.2.5 Ecology

3.2.5.1 Terrestrial

A generalized overview and description of the habitat types and terrestrial species that may be found in areas used for milling operation are discussed in this section. These areas are broad and contain many subregions. For specific future locations of new milling sites, potential license applicants and the NRC review would be expected to address sitespecific habitat types and terrestrial species.

Wyoming West Uranium Milling Region Flora

According to the EPA, the identified ecoregions in the Wyoming West Uranium Milling Region primarily consist of Wyoming Basin and the Middle Rockies ecoregions (Chapman, et al., 2004). Figure 3.2-7 depicts the various ecoregions found within the Wyoming West Uranium Milling Region. Uranium milling districts within the uranium districts in the region are located within the Rolling Sagebrush Steppe and the Salt Desert Shrub Basin ecoregions of the Wyoming Basin.

The Wyoming Basin ecoregion is a broad, arid, intermontane basin interrupted by hills and low mountains and dominated by grasslands and shrublands. Nearly surrounded by forest-covered mountains, the region is drier than the Northwestern Great Plains to the northeast and does not have the extensive cover of pinyon-juniper woodland found in the Colorado Plateaus to the south. Much of the region is used for livestock grazing, although many areas lack sufficient forage to support this activity (Chapman, et al., 2004). Within the Wyoming Basin, the Wyoming West Uranium Milling Region contains several subecoregions that are described below, based on the descriptions of Chapman, et al. (2004).

The Rolling Sagebrush Steppe area of the Wyoming basin is composed of rolling plains with hills, mesas, and terraces. Areas near the mountains may contain footslopes, ridges, alluvial fans, and outwash fans (Chapman, et al., 2004). The most abundant shrub vegetation in the region is Wyoming big sagebrush (*Artemisia tridentata* ssp. *wyomingensis*), with silver



2

Description of the Affected Environment

sagebrush (*Artemisia cana*) and black sagebrush (*Artemisia nova*) occurring in the lowlands and mountain big sagebrush (*Artemisia tridentata* ssp. *vaseyana*) in the higher elevations. Grass species include western wheatgrass (*Pascopyrum smithii*), needle-and-thread grass (*Stipa comata*), blue gramma (*Bouteloua gracilis*), Sandberg bluegrass (*Poa secunda*), junegrass (*Koeleria macrantha*), rabbitbrush (*Chrysothamnus nauseosus*), and fringed sage (*Artemisia frigida*) (Chapman, et al., 2004).

The Bighorn Basin is primarily an arid region influenced by the rainshadow effect of the Beartooth Mountains, Absaroka Range, and Pryor Mountains. This higher portion of the greater Bighorn Basin forms a transition from arid desert shrubland to semiarid shrubland. Sage steppe vegetation dominates this region and is composed of species such as Wyoming big sagebrush, western wheat grass, blue wheatgrass (*Elymus magellanicus*), needle-and thread grass, blue gramma, Sandberg bluegrass, junegrass, rabbitbrush, and fringed sage. (Chapman, et al., 2004).

The Foothill Shrublands ecoregion serves as a transitional zone between the forested Dry Mid-Elevation Sedimentary Mountains ecoregion to the arid grassland and sagebrush regions in the Wyoming Basin and the High Plains (Chapman, et al., 2004).

Vegetation found within this region include Sagebrush steppe communities, mountain mahogany woodlands that are often interspersed with mountain big sagebrush, blue grama, prairie junegrass, western wheatgrass, and ponderosa pine (*Pinus ponderosa*) savanna in the Laramie foothills (Chapman, et al., 2004).

The Sub-Irrigated High Valleys are wet meadow systems located in areas of high drainage density beneath surrounding mountain ranges. Soil in this region remains moist due to the presence of a high water table. This region is abundant with floodplains, low terraces, riparian wetlands, and alluvial fans. As a result, the riparian areas and wet meadows are dominated by willows, alders, cottonwoods and wetland plants, such as horsetail (*Equisetum* sp.), spikerush (*Eleocharis* sp.), sedges (*Cyperaceae* sp.), and tufted hairgrass (*Deschampsia cespitosa*) found in low drainage areas. Shrubland areas may include Wyoming big sagebrush, western Wheatgrass, needle-and-thread grass, blue gramma, Sandberg blue grass, junegrass, rabbitbrush, and fringe sage (Chapman, et al., 2004)

The Salt Desert Shrub Basins ecoregion is an arid environment that includes isolated playa lakes and sand dunes scattered throughout the Wyoming Basin. Vegetation in this area consists of arid land alkaline tolerant shrubs such as shadscale (*Atriplex confertifolia*), greasewood (*Sarcobatus vermiculatus*), and Gardner saltbush (*Atriplex gardneri*) low in abundance. Plant life is more diverse on sand dunes, which provide greater moisture, higher permeability, and lower alkalinity than the basin floor. Vegetation found on stable sand dune areas includes alkali cordgrass (*Spartina gracilis*), Indian grass (*Sorghastrum nutans*), blowout grass (*Redfieldia flexuosa*), alkali wildrye (*Leymus simplex*), and needle-and-thread grass (Chapman, et al., 2004).

The Bighorn Salt Desert Shrub Basins are composed of two large, arid, alkaline depressions surrounded by mountains. This region is geographically isolated from the other salt desert shrub basins in southern Wyoming. This region has a greater human influence due to the proximity to major rivers (Bighorn, Shoshone, and Greybull rivers), which provide water for irrigation. This region receives approximately 15 cm [6 in] of precipitation per year and supports desert shrubs and grasses. Vegetation found in this region may consist of greasewood,

1 Gardner saltbush, shadscale, alkali sacaton, and saltgrass (Chapman, et al., 2004). The
2 vegetation around major rivers consist of open woodland of plains cottonwood (*Populus*
3 *deltoides*), narrowleaf cottonwood (*Populus angustifolia*), peachleaf willow (*Salix amygdaloides*),
4 and wild plum (*Prunus americana*).
5

6 The Middle Rockies ecoregion is composed of steep-crested, high mountains that are largely
7 covered by coniferous forests.
8

9 The Bighorn, Beartooth mountains, and the Wind River and Teton ranges, comprise the Granitic
10 Subalpine Zone. Snow melt moisture, absorbed and released throughout the spring and
11 summer, provides water for humans and wildlife living at lower elevations in the droughty,
12 sedimentary fringes of these mountains. Subalpine forests are dominated by Lodgepole pine
13 (*Pinus contorta*) at the lower elevations with subalpine fir (*Abies lasiocarpa*) and Engelmann
14 spruce (*Picea engelmannii*) found in the higher elevations. The diversity of the understory is low
15 and consists mostly of grouse whortleberry (*Vaccinium scoparium*), Oregon grape (*Mahonia*
16 *aquifolium*), and birchleaf spirea (*Spiraea betulifolia*). The subalpine spruce-fir zone is not as
17 heavily grazed by livestock as mid-elevation areas; it serves as summer range for mule deer
18 and elk (Chapman, et al., 2004).
19

20 The Dry Mid-Elevation Sedimentary Mountains ecoregion includes the mid-elevation Bighorn
21 Mountains and the drier northeastern portion of the Wind River Range that are underlain by
22 sedimentary rocks. The lack of moisture in the soil is enhanced by the rainshadow effects of the
23 two mountain ranges. Upland forest cover is open and patchy due to arid conditions. Forests of
24 the Wind River Range are dominated by Douglas firs with an understory of grasses, forbs, and
25 shrubs. Forest cover is more extensive on the east slopes of the Bighorns where there is more
26 summer precipitation. A ponderosa pine/juniper/mountain mahogany association exists here
27 similar to one in the Black Hills region to the east, but it is of limited extent. The forest of the
28 eastern Bighorn Mountains lacks enough precipitation to support the eastern deciduous species
29 and boreal vegetation present in the Black Hills. Some quaking aspen groves occur in this
30 region, particularly in the Wind River Range (Chapman, et al., 2004).
31

32 A comprehensive listing of habitat types and species found in the aforementioned ecoregions
33 has been surveyed and compiled as part of the Wyoming Gap Analysis project (Wyoming
34 Geographic Information Science Center, 2007a,b).
35

36 The Wyoming Gap Analysis project is part of the National Gap Analysis Program. It began in
37 1991 and was officially completed in November 1996. The program's main goal was to analyze
38 the status of biodiversity within Wyoming, focusing on two biodiversity elements: land cover
39 types and terrestrial vertebrate species. Land ownership and management for the state of
40 Wyoming was combined with the data on land cover and species distributions in a geographic
41 overlay. A Geographical Information System was used to determine which biodiversity
42 elements were inadequately protected within the current system of areas managed for
43 conservation (Wyoming Geographic Information Science Center, 2007a,b).
44

Wyoming West Uranium Milling Region Fauna

According to the official state list of birds, mammals, amphibians, and reptiles in Wyoming compiled by the Wyoming Game and Fish Department, approximately 246 bird, 127 mammal, 12 amphibian, and 27 reptile species are found in Wyoming. The official state list of the common and scientific names of the birds, mammals, amphibians, and reptiles in Wyoming can be obtained from the Wyoming Game and Fish Department (2007a).

According to the World Wildlife Fund's species database (World Wildlife Fund, 2007a,b), approximately 285 different species are found within the Wyoming Basin. Common animals found in this region include large game mammals such as moose (*Alces Alce*), pronghorn (*Antilocapra Americana*), elk (*Cervus elaphus*), mule deer (*Odocoileus hemious*), white tailed deer (*Odocoileus virginianus*), bighorn sheep (*Ovis Canadensis*), and American black bear (*Ursus americanus*). Numerous rodents such as chipmunks (*Tamias spp.*), squirrels (*Speermophilus spp.*), shrews (*Sorex spp.*), and rabbits (*Sylvilagus spp.*) and numerous myotic bat species are found within this region. Reptiles and amphibians found in the region include species such as the western rattlesnake (*Crotalus viridis*), gopher snake (*Pituophis caterifer*), garter snake (*Thamnophis elegans*), tiger salamander (*Ambystoma tigrium*), Woodhouse's toad (*Bufo woodhouii*), and spadefoot toad (*Scaphiopus spp.*). A diverse number of birds also inhabit this region, including hawks like the Cooper's hawk (*Accipter cooperii*), goshawk (*Accipiter gentilis*), and red-tailed hawk (*Buteo jamaicensis*) and the golden eagle (*Aquila chrysaetos*). Common birds in the region include finches (*Leucosticte spp.*), sparrows (*Melospiza spp.*), owls (*Otus spp.*), swallows (*Tachycinets spp.*), and vireos (*Vireo spp.*) in addition to other songbirds. A noted species within this region is the white-tailed prairie dog (*Cynomys leucurus*). The white-tailed prairie dog towns in this region provide food for predators such as the coyote (*Canis latrans*), the swift fox (*Vulpes velox*), and the black-footed ferret (*Mustela nigripes*)—a federally recognized endangered species (World Wildlife Fund, 2007a,b).

The Foothill Shrublands ecoregion is a transition region between the Dry Mid-Elevation Sedimentary Mountains ecoregion, Wyoming Basin Shrublands, the Northwest Great Plains, and the South Central Rockies forest, species found in this region will overlap all regions. Again, large mammal species such as bighorn sheep, cougar, American bison, pronghorn, moose, elk, and coyotes can be found in this region. Shrews, voles, rabbits, squirrels, and prairie dogs common to the other ecoregions can also be found in this transition area. Raptors such as Cooper's hawk, goshawk, red-tailed hawk, golden eagles, and numerous owl species are bird predators in this area. Common bird species in the region include finches, sparrows, swallows, vireos, warblers, and kingbirds in addition to other songbirds (World Wildlife Fund, 2007a–e).

The Middle Rockies ecoregion contains over 300 different species. This region features large, important herds of elk and mule deer, which are the main game species in this region. Large predators such as cougar (*Puma concolor*) and black bear (*Ursus americanus*) are also abundant. Other mammals found in this region include the wolverine (*Gulo gulo*), lynx (*Lynx canadensis*), pronghorn, beaver (*Castor canadensis*), coyote, Gunnison's prairie dog, black-tailed prairie dog, porcupine (*Eremophila dorsatum*), bat, and American marten (*Martes americana*). Numerous rodents such as squirrels, voles, rabbits, rats, and mice occur in this region. Common birds in the region include many of the species found throughout Wyoming like bluebirds, sparrows, ducks, woodpeckers, owls, hawks, and eagles. Reptile and amphibian species include the soft-shelled turtle, plateau striped whiptail (*Cnemidophorus velox*), western rattlesnake, many-lined skink (*Eumeces multivirgatus*), fence lizard, tiger salamander, western

1 toad (*Bufo boreas*), and the Baird's spotted toad (*Bufo punctatus*) (World Wildlife Fund,
2 2007a–e).

3
4 According to the Wyoming Game and Fish Department, crucial wintering habitats are found
5 within this region for large game mammals and nesting leks for the sage grouse (Wyoming
6 Game and Fish Department, 2007b). Figures 3.2-8 through 3.2-14 depict the crucial winter and
7 yearlong areas ranges for large mammals and game birds found in this region. Most of the
8 crucial areas for big game animals in the Wyoming West Uranium Milling Region are located in
9 the Rattlesnake Hills and Granite Mountains in the central and northwestern parts of the region,
10 or along the Sweetwater River and its tributaries. Sites identified within Crook's Gap and Gas
11 Hills Uranium Districts are located in or near crucial winter/yearlong habitat for antelope, moose,
12 and mule deer. Numerous sage grouse leks nesting areas are located near sites in both
13 uranium districts, articularly in the southeastern portion of the study region (i.e., Crook's Gap
14 Uranium District).

15 16 **3.2.5.2 Aquatic**

17
18 Within the Wyoming West Uranium Milling Region, several watersheds have been listed as
19 aquatic habitat areas. These areas include the Lower Wind River/Boysen Reservoir watershed,
20 Upper Sweetwater River Watershed, lower Sweetwater watershed, Middle Fork Popo Agie,
21 Middle North Platte River Corridor, and the South Fork Powder River watersheds. These
22 watersheds are part of the larger Lower Wind River, Sweetwater, South Fork Powder River, and
23 Middle North Platte-Casper watersheds previously discussed in Section 3.2.4.1 (Wyoming
24 Game and Fish Department, 2007b). The two uranium districts within the Wyoming West
25 Uranium Milling Region are located in the Sweetwater (Crooks Gap) and Wind River (Gas Hills)
26 watersheds.

27
28 According to the Wyoming Fish and Game Department (Wyoming Game and Fish Department,
29 2007a), there are approximately 49 native fish species found in the watersheds throughout the
30 state. These species are identified in Table 3.2-5. Current conditions of these watersheds have
31 been evaluated, and fish species that would benefit from conservation measures within the
32 watersheds have been identified.

33
34 The Lower Wind River discharges into the Boysen Reservoir. Additional waterways which are
35 included in the basin are the Stagner Creek, Gold Creek, Cottonwood Creek, Birdseye Creek,
36 Reservoir Creek, Muddy Creek, Poison Creek, and Cottonwood Drain. Aquatic species found in
37 this system include Sauger (*Stizostedion canadense*), burbot (*Lota lota*), mountain whitefish
38 (*Prosopium williamsoni*), stonecat (*Noturus flavus*), channel catfish (*Ictalurus punctatus*),
39 longnose dace (*Rhinichthys cataractae*), Northern Redhorse (*Moxostoma aureouim*), and
40 Flathead chub (*Platygobio gracilis*). Sport fish that occur in the watershed include rainbow trout
41 (*Oncorhynchus mykiss*), brown trout (*Salmo trutta*), walleye (*Sander vitreus*), brook trout
42 (*Salvelinus fontinalis*), lake trout (*Salvelinus namaycush*), largemouth bass (*Micropterus*
43 *salmoides*), black crappie (*Pomoxis nigromaculatus*), bluegill (*Lepomis macrochirus*), yellow
44 perch (*Perca flavescens*), and black bullhead (*Ameiurus melas*) (Wyoming Game and Fish
45 Department, 2007b).

1
2




Figure 3.2-8. Antelope Wintering Areas for the Wyoming West Uranium Milling Region

3

1




Figure 3.2-9. Big Horn Wintering Areas for the Wyoming West Uranium Milling Region

2

1




Figure 3.2-10. Elk Wintering Areas for the Wyoming West Uranium Milling Region

2

1




Figure 3.2-11. Moose Wintering Areas for the Wyoming West Uranium Milling Region

2

1




Figure 3.2-12. Mule Deer Wintering Areas for the Wyoming West Uranium Milling Region

2

1




Figure 3.2-13. Sage-Grouse/Lek Nesting Areas for the Wyoming West Uranium Milling Region

2

1




Figure 3.2-14. White Tailed Deer Wintering Areas for the Wyoming West Uranium Milling Region

2

The Middle Fork Popo Agie watershed is found in the western and southern portion of the Wyoming West Uranium Milling Region. Contributing waterways include Saw creek and Sawmill Creeks. Species in this watershed have been impacted by erosion and sediment processes which have been accelerated by human activities such as prolonged annual herbivory, increased drainage from roads and trails, removal of water for irrigation, dewatering of wetlands, and rural subdivision development. Native species found within this watershed include the lakechub (*Couesius plumbeus*), longnose dace, longnose sucker (*Catostomus catostomus*), white sucker (*Catostomus commersonii*), mountain sucker (*Catostomus platyrhynchus*), mountain whitefish, and flathead minnow (*Pimephales promelas*). Sport fish found in this watershed include rainbow trout, brown trout, brook trout, Yellowstone trout (*Oncorhynchus clarki bouvieri*), Snake River cutthroat trout (*Oncorhynchus clarki ssp.*), and grayling (*Thymallus thymallus*) (Wyoming Game and Fish Department, 2007b).

Table 3.2-5. Native Fish Species Found in Wyoming

Common Name	Scientific Name
Arctic Grayling	<i>Thymallus arcticus</i>
Bigmouth Shiner	<i>Notropis dorsalis</i>
Black Bullhead	<i>Ameiurus melas</i>
Bluehead Sucker	<i>Catostomus discobolus</i>
Brassy Minnow	<i>Hybognathus hankinsoni</i>
Burbot	<i>Lota lota</i>
Central Stoneroller	<i>Campostoma anomalum</i>
Channel Catfish	<i>Ictalurus punctatus</i>
Common Shiner	<i>Luxilus cornutus</i>
Creek Chub	<i>Semotilus atromaculatus</i>
Cutthroat Trout	<i>Oncorhynchus clarki</i>
Fathead Minnow	<i>Pimephales promelas</i>
Finescale Dace	<i>Phoxinus neogaeus</i>
Flannelmouth Sucker	<i>Catostomus latipinnis</i>
Flathead Chub	<i>Platygobio gracilis</i>
Goldeye	<i>Hiodon alosoides</i>
Hornyhead Chub	<i>Nocomis biguttatus</i>
Iowa Darter	<i>Etheostoma exile</i>
Johnny Darter	<i>Etheostoma nigrum</i>
Lake Chub	<i>Couesius plumbeus</i>
Leatherside Chub	<i>Gila copei</i>
Longnose Dace	<i>Rhinichthys cataractae</i>
Longnose Sucker	<i>Catostomus catostomus</i>
Mottled Sculpin	<i>Cottus bairdi</i>
Mountain Sucker	<i>Catostomus platyrhynchus</i>
Mountain Whitefish	<i>Prosopium williamsoni</i>
Orangethroat Darter	<i>Etheostoma spectabile</i>
Paiute Sculpin	<i>Cottus beldingi</i>
Pearl Dace	<i>Margariscus margarita</i>
Plains Killifish	<i>Fundulus zebrinus</i>
Plains Minnow	<i>Hybognathus placitus</i>
Plains Topminnow	<i>Fundulus sciadicus</i>
Quillback	<i>Carpionodes cyprinus</i>

1

Table 3.2-5. Native Fish Species Found in Wyoming (continued)	
Common Name	Scientific Name
Red Shiner	<i>Cyprinella lutrensis</i>
Redside Shiner	<i>Richardsonius balteatus</i>
River Carpsucker	<i>Carpionodes carpio</i>
Roundtail Chub	<i>Gila robusta</i>
Sand Shiner	<i>Notropis stramineus</i>
Sauger	<i>Stizostedion canadense</i>
Shorthead Redhorse	<i>Moxostoma macrolepidotum</i>
Shovelnose Sturgeon	<i>Scaphirhynchus platyrhynchus</i>
Speckled Dace	<i>Rhinichthys osculus</i>
Stonecat	<i>Noturus flavus</i>
Sturgeon Chub	<i>Macrhybopsis gelida</i>
Suckermouth Minnow	<i>Phenacobius mirabilis</i>
Utah Chub	<i>Gila atraria</i>
Utah Sucker	<i>Catostomus ardens</i>
Western Silvery Minnow	<i>Hybognathus argyritis</i>
White Sucker	<i>Catostomus commersoni</i>

2

3

4

5

6

7

8

9

10

11

The Upper Sweetwater River headwaters in the Wind River Mountains and flows across the South Pass uplift area. Native species found within this watershed include the lake chub, creek chub (*Semotilus atromaculatus*), longnose dace, longnose sucker, white sucker, mountain whitefish, flathead minnow, Iowa darter (*Etheostoma exile*), and mountain sucker. Sport fish found in this watershed include rainbow trout, brown trout, brook trout, fallriver rainbow, Yellowstone cutthroat trout, Snake River cutthroat, and Bear River cutthroat (Wyoming Game and Fish Department, 2007b).

12

13

14

15

16

17

18

19

20

21

22

The Lower Sweetwater River watershed is found in the south central portion of the Wyoming West Uranium Milling Region. Contributing waterways include Crook Creek and Willow Creek. Species in this watershed have been impacted by erosion and sediment processes which have been accelerated by human activities such as prolonged annual herbivory, increase drainage from roads and trails as a result of previous uranium milling operations in the Green Mountain Area. Native species found within this watershed include the lake chub, creek chub, longnose dace, longnose sucker, white sucker, mountain sucker, flathead minnow, bigmouth sucker (*Ictiobus cyprinellus*) and Iowa darter. Sport fish found in this watershed include rainbow trout, brown trout, brook trout, fallriver rainbow, and bear river cutthroat (Wyoming Game and Fish Department, 2007b).

23

24

25

26

27

28

29

The South Fork Powder River-Murphy Creek basin is relatively dry and sparsely vegetated. Most of the streams are ephemeral or intermittent with few perennial streams. Many of these stream channels are degraded or actively degrading. Native fish species that can be found in this watershed include the creek chub, fathead minnow, flathead chub, longnose dace, plains minnow, sand shiner, mountain sucker, and the plains killifish (Wyoming Game and Fish Department, 2007b).

30

31

32

Middle North Platte River Corridor portion of the watershed is located on the eastern side of the Wyoming West Uranium Milling Region. Species found within this watershed include the brassy minnow (*Hybognathus hankinsoni*), common shiner (*Notropis cornutus*), creek chub, fathead

minnow, longnose dace, sand shiner (*Notropis stramineus*), stoneroller (*Camptostoma anomalum*), longnose sucker, white sucker with the rainbow trout, brown trout, cutthroat trout and channel catfish being sport fish (Wyoming Game and Fish Department, 2007b).

The Sweetwater River Muddy Creek and Horse Creek watersheds are located in the southern portion of the Wyoming West Uranium Milling Region. This watershed region has been impacted by intense herbivory, the successional advance of big sagebrush steppe and absence of beaver dams are the perceived bottlenecks limiting watershed function. Native species found within this watershed include the bigmouth shiner, creek chub, fathead minnow, longnose dace, sand shiner, longnose sucker, white sucker, and Iowa darter. Sport fish in the watershed include rainbow trout, brown trout, cutthroat trout, and brook trout.

3.2.5.3 Threatened and Endangered Species

Federally listed threatened and endangered species known to exist in habitats in the West Wyoming Uranium Milling Region include the following:

- Black-Footed Ferret (*Mustela nigripes*)—Ferrets were once found throughout the Great Plains, from Texas, New Mexico, and Arizona to southern Saskatchewan, Canada. Ferrets eat prairie dogs and live in prairie dog burrows. Typical wild ferret behavior revolves around prairie dog towns. Wild ferrets hunt prairie dogs at night, but occasionally they are active above ground during the day. This is especially true of female ferrets hunting to feed their young. In search of prey, they move from one prairie dog burrow to the next (U.S. Fish and Wildlife Service, 2008).
- Blowout Penstemon (*Penstemon haydenii*)—Limited to the sandhills region of west-central Nebraska, and sand dune habitat in the northeastern Great Divide Basin in Wyoming. In Nebraska this plant typically occurs in "blowouts"—sparsely vegetated depressions in active sand dunes created by wind erosion. In Wyoming it occurs on sandy aprons or the lower half of steep sandy slopes deposited at the base of granitic or sedimentary mountains or ridges. It occurs at elevations ranging from 850–1,150 m [2,800–3,800 ft] in Nebraska to 2,030–2,270 m [6,680–7,440 ft] in Wyoming. This species can be found in west-central Nebraska in Box Butte, Cherry, Garden, Morrill and Thomas counties, and in the Wyoming West Uranium Milling Region in northwestern Carbon County (Center for Plant Conservation, 2008).
- Bonytail Chub (*Gila elegans*)—Found in slower water habitats in the main stream such as eddies, pools, sidechannels, and coves. They are found in streams below 1,220 m [4,000 ft] elevation. Endemic to the Colorado River basin and found throughout the mainstem rivers and backwaters of the Upper and Lower Basins. This species is one of the rarest of the Colorado River fishes and is close to extinction (U.S. Fish and Wildlife Service, 2008).
- Canada Lynx (*Lynx canadensis*)—The Canada lynx inhabits mountain regions, primarily at elevations between 2,356 and 2,869 m [7,730 to 9,410 ft] and on slopes of 8 to 12 percent. It usually occurs in extensive tracts of dense coniferous forest, primarily Engelmann spruce and subalpine fir. It feeds primarily on snowshoe hares, especially during winter (thereby making habitat for snowshoe hares a key consideration for lynx habitat). Older forests with a substantial understory of conifers or small patches of shrubs and young trees provide good quality lynx foraging habitat. The most important

Description of the Affected Environment

component of denning habitat is large woody debris, especially dense tangles of fallen trees and root wads. Such preferred habitat is relatively limited in Wyoming and occurs primarily in multiple use areas of the Shoshone and Bridger-Teton National Forests along the western boundary of the Wyoming West Uranium Milling Region. The National Parks and designated wilderness areas in Wyoming tend to be marginal lynx habitat as they are either dominated by dry even aged lodgepole pine forests, or too steep and high elevation (Wyoming Game and Fish Department, 2008).

- Colorado Pikeminnow (*Ptychocheilus lucius*)—Colorado pikeminnow were once abundant in the main reach of the Colorado River and most of its major tributaries in Colorado, Wyoming, Utah, New Mexico, Arizona, Nevada, California and Mexico. Now, they exist primarily in the Green River below the confluence with the Yampa River, the lower Duchesne River in Utah, the Yampa River below Craig, Colorado, the White River from Taylor Draw Dam near Rangely, Colorado downstream to the confluence with the Green River, the Gunnison River in Colorado, and the Colorado River from Palisade, Colorado, downstream to Lake Powell. It is believed that the Colorado pikeminnow populations in the upper Colorado River basin are now relatively stable and in some areas may even be growing (U.S. Fish and Wildlife Service, 2008).
- Humpback Chub (*Gila cypha*)—The humpback chub lives primarily in canyons with swift currents and white water. Historically, it inhabited canyons of the Colorado River and four of its tributaries: the Green, Yampa, White and Little Colorado rivers. Now, there are two populations near the Colorado/Utah border—one at Westwater Canyon in Utah and one in an area called Black Rocks, in Colorado. Though now smaller in number than they were historically, the two populations seem to be fairly stable in these two areas (U.S. Fish and Wildlife Service, 2008).
- Interior Least Tern (*Sterna antillarum athalassos*)—Nesting habitat of the Interior Least Tern includes bare or sparsely vegetated sand, shell, and gravel beaches, sandbars, islands, and salt flats associated with rivers and reservoirs. The birds prefer open habitat, and tend to avoid thick vegetation and narrow beaches. Sand and gravel bars within a wide unobstructed river channel, or open flats along shorelines of lakes and reservoirs, provide favorable nesting habitat. Nesting locations are often at the higher elevations away from the water's edge, since nesting usually starts when river levels are high and relatively small amounts of sand are exposed. The size of nesting areas depends on water levels and the extent of associated sandbars and beaches. Highly adapted to nesting in disturbed sites, terns may move colony sites annually, depending on landscape disturbance and vegetation growth at established colonies (Texas Parks and Wildlife Department, 2007).
- Pallid Sturgeon (*Scaphirhynchus albus*)—This species is a bottom dweller, found in areas of strong current and firm sand bottom in the main channel of large turbid rivers such as the Missouri and Platte River. The pallid sturgeon is a member of a primitive family that, like other sturgeon, has lengthwise rows of bony plates covering its body, rather than scales. Pallids are slow growing, late-maturing fish that feed on small fishes and immature aquatic insects. Spawning occurs from June through August (Platte River Endangered Partnership, 2008).
- Piping Plover (*Charadrius melodus*)—Piping plovers breed only in North America in three geographic regions: the Atlantic Coast, the Northern Great Plains, and the Great

Lakes. Plovers in the Great Plains make their nests on open, sparsely vegetated sand or gravel beaches adjacent to alkali wetlands, and on beaches, sand bars, and dredged material islands of major river systems (U.S. Fish and Wildlife Service, 2008).

- Preble's Meadow Jumping Mouse (*Zapus hudsonius preblei*)—This species lives primarily in heavily vegetated, shrub-dominated riparian (streamside) habitats and immediately adjacent upland habitats along the foothills of southeastern Wyoming south to Colorado Springs along the eastern edge of the Front Range of Colorado. Documented distribution includes Albany, Laramie, Platte Goshen, and Converse counties in Wyoming (U.S. Fish and Wildlife Service, 2008)
- Razorback Sucker (*Xyrauchen texanus*)—This is a large river species not found in smaller tributaries and headwater streams. Found in water from 1–3 m [4–10 ft] in depth, adults are associated with areas of strong current and backwaters (Colorado Division of Wildlife, 2008). This species has been extirpated from Wyoming however it can be occasionally found in Sweetwater County (University of Wyoming, 2008).
- Ute Ladies' Tresses Orchid (*Spiranthes diluvialis*)—Populations of Ute ladies'-tresses orchids are known from three broad general areas of the interior western United States—near the base of the eastern slope of the Rocky Mountains in southwestern Wyoming and adjacent Nebraska and north-central and central Colorado; in the upper Colorado River basin, particularly in the Uinta Basin; and in the Bonneville Basin along the Wasatch Front and westward in the eastern Great Basin, in north-central and western Utah, extreme eastern Nevada, and southeastern Idaho. The orchid also has been discovered in southwestern Montana and in the Okanogan area and along the Columbia River in north-central Washington. The orchid occurs along riparian edges, gravel bars, old oxbows, high flow channels, and moist to wet meadows along perennial streams. It typically occurs in stable wetland and seepy areas associated with old landscape features within historical floodplains of major rivers. It also is found in wetland and seepy areas near fresh water lakes or springs (U.S. Fish and Wildlife Service, 2008).
- Western Prairie Fringed Orchid (*Platanthera praeclara*)—The western prairie fringed orchid is a plant of the tallgrass prairie and requires direct sunlight for growth. It is most often found in moist habitats or sedge meadows. (U.S. Fish and Wildlife Service, 2008).
- Whooping Crane (*Grus americana*)—The whooping crane prefers fresh water marshes, wet prairies, shallow portions of rivers and reservoirs, grain and stubble fields, shallow lakes and lagoons for feeding and loafing during migration. The whooping crane formerly nested from central Illinois west to eastern North Dakota and north through the Canadian prairie provinces. It presently breeds in Wood Buffalo National Park in the Northwest Territories, Canada. It overwinters on the Texas Gulf Coast on and in the vicinity of the Aransas National Wildlife Refuge. A second foster population migrates from Grays Lake National Wildlife Refuge in Idaho to the Bosque del Apache National Wildlife Refuge on the Rio Grande River in New Mexico. In South Dakota, the whooping crane is a predictable spring and fall migrant in the Missouri River drainage and in western South Dakota (Platte River Endangered Partnership, 2008).
- Yellow Billed Cuckoo (*Coccyzus americanus*)—(candidate)—Throughout their range, preferred breeding habitat includes open woodland (especially where undergrowth is

Description of the Affected Environment

thick), parks, and deciduous riparian woodland. In the West, they nest in tall cottonwood and willow riparian woodlands. Nests are found in trees, shrubs, or vines an average of 1 to 3 m [3–10 ft] above ground (Harrison, 1979). Western subspecies require patches of at least 10 hectares [25 acres] of dense, riparian forest with a canopy cover of at least 50 percent in both the understory and overstory (Montana Natural Heritage Program, 2008).

The state of Wyoming does not maintain a list of threatened or endangered plant or animal species, but has established a non-game bird and mammal plan that includes a list of species of special concern. All of the federally listed animal species are considered by the state species of special concern. Wyoming Species of Concern are described as special status Wyoming Native Species Status matrix 1 (populations are greatly restricted or declining—extirpation appears possible), and 2 (populations are declining or restricted in numbers and or distribution—extirpation is not imminent. Wyoming Species of Concern which may be found in the Wyoming West Uranium Milling Region include the following:

- Flannelmouth Sucker (*Catostomus latipinnis*) Native Species Status 1—This species prefers large rivers with deep riffles and runs, they can also be found in smaller streams and sometimes in lakes. Native to the Colorado River drainage basin, in Wyoming it is found in the Green and Little Snake river drainages. In the spring they leave the large rivers and ascend small tributary streams to spawn; migrations of over 225 km [140 mi] have been documented (Wyoming Game and Fish Department, 2008).
- Boreal Toad (*Bufo boreas*) Native Species Status 1—The southern Rocky Mountain population occurs from south-central Wyoming southward through the mountainous regions of Colorado to extreme north-central New Mexico. The toads inhabit a variety of wet habitats (i.e., marshes, wet meadows, streams, beaver ponds, glacial kettle ponds, and lakes interspersed in subalpine forest) at altitudes primarily between 2,400–3,400 m [8,000–11,500 ft] (U.S. Fish and Wildlife Service, 2008).
- Common Loon (*Gavia immer*) Native Species Status 1—Lakes that are suitable for breeding are extremely limited in Wyoming and must have the following characteristics: At least 4 ha (10 ac), although reproductive success is better on lakes that are greater than 10 hectares (25 acres); Free of human disturbance or have areas that are secluded from human activity; Between 1,800 and 2,400 m [1,000 and 8,000 ft] in elevation; Have clear water with a minimum visibility of 3 to 4 m [10 to 13 ft], as loons are visual predators; Islands or protected shore areas for nesting and raising young; Abundant populations of small to mid-sized fish; Greater than 2 m [6 ft] deep to prevent winter kill of fish; remain ice free for at least 4 months to allow young to fledge; and nesting, lakes with partially forested, rocky shorelines; an area of shallow water with emergent vegetation; and a steep slope adjacent to the shoreline for an underwater approach to the nest (Wyoming Game and Fish Department, 2008).
- Burbot (*Lota lota*) Native Species Status 1—The burbot lives in cold, deep lakes and large rivers. Immature fish prefer rubble substrate, while adults remain in deep water to prey on other fish. In Wyoming, the burbot is native to the Big Horn and Tongue River systems. It is found in larger lakes in the Lander and Dubois area, including Boysen Reservoir and Ocean Lake. It also occurs south to Missouri and Kansas and east to New England, as well as throughout Canada (Wyoming Game and Fish Department, 2008).

- 1
2 • Sauger (*Sander canadensis*) Native Species Status—The sauger prefers large rivers but
3 may also be found in reservoirs. The fish is tolerant of turbid waters. In rivers the key
4 component of sauger habitat is velocity. In the summer and spring they select low
5 velocity areas having sand or silt substrates. Pool habitats are preferred by sauger
6 especially in winter where they tend to select low velocity pools greater than 2 m [6 ft]
7 deep. Native to streams east of the Continental Divide, the sauger occurs in Wyoming
8 today in the Wind Big Horn River drainage and in the Tongue and Powder River
9 drainages. It has apparently been extirpated from the North Platte River, where it had
10 once been common (Wyoming Game and Fish Department, 2008).
- 11
12 • Yellowstone Cutthroat (*Oncorhynchus clarki bouvieri*) Native Species Status 1—The
13 Yellowstone cutthroat lives in lakes, large rivers, and small tributary streams. Native to
14 the Yellowstone River drainage downstream to the Tongue River, including the Big Horn
15 and Clarks Fork River drainages, this trout is also found in Pacific Creek and other
16 Snake River tributaries. All other occupation by this species east of the Continental
17 Divide is from introductions (Wyoming Game and Fish Department, 2008).
- 18
19 • Cliff Tree Lizard (*Urosaurus ornata wrightii*) Native Species Status 1—This lizard prefers
20 cliffs and rocky canyon slopes in sagebrush desert habitats. It is often found on the
21 vertical surfaces of large boulders or rock cliffs. In Wyoming, the cliff tree lizard occurs
22 in the extreme southwestern part of the state. It also ranges south through Utah and
23 western Colorado to northern Arizona and northern New Mexico (Wyoming Game and
24 Fish Department, 2008).
- 25
26 • Great Basin Gopher Snake (*Pituophis melanoleucas deserticola*) Native Species
27 Status 1—This snake prefers sagebrush communities and deserts in the plains zone. In
28 Wyoming, it can be found in the south-central counties at lower elevations, and west of
29 the Continental Divide in the Wyoming Basin. Elsewhere, it is distributed from the Great
30 Basin to eastern California, Oregon, and Washington (Wyoming Game and Fish
31 Department, 2008).
- 32
33 • Rubber Boa (*Charina bottae*) Native Species Status 1—The rubber boa prefers areas
34 with an abundance of flat rocks and water nearby. It does not inhabit Wyoming's arid
35 regions, but may be found in the foothills and lower mountain zones of the northwestern
36 corner of the state, south into Star Valley and east to the Big Horn Mountains. It is also
37 distributed west of Wyoming to the Pacific Coast from British Columbia to northern
38 California (Wyoming Game and Fish Department, 2008).
- 39
40 • Canada Lynx (*Lynx canadensis*) Native Species Status 1—The Canada lynx inhabits
41 mountain regions, primarily at elevations between 2,356 and 2,869 m [7,730 to 9,413 ft]
42 and on slopes of 8 to 12 percent. It usually occurs in extensive tracts of dense
43 coniferous forest, primarily Engelmann spruce and subalpine fir. It feeds primarily on
44 snowshoe hares, especially during winter, and the prime consideration for lynx is habitat
45 for snowshoe hares. Older forests with a substantial understory of conifers or small
46 patches of shrubs and young trees provide good quality lynx foraging habitat. The most
47 important component of denning habitat is large woody debris, especially dense tangles
48 of fallen trees and root wads. Such preferred habitat is relatively limited in Wyoming and
49 occurs primarily in multiple use areas of the Shoshone and Bridger-Teton National
50 Forests. The National Parks and designated wilderness areas in Wyoming tend to be

Description of the Affected Environment

marginal lynx habitat as they are either dominated by dry even-aged lodgepole pine forests, or too steep and high elevation (Wyoming Game and Fish Department, 2008).

- Pale Milk Snake (*Lampropeltis triangulum multistrata*) Native Species Status 2—The pale milk snake prefers grasslands, sandhills, and scarp woodlands below 1,800 m [6,000 ft] in elevation. It is distributed throughout the northern Great Plains. In Wyoming, it can be found in the eastern counties and the Big Horn Basin (Wyoming Game and Fish Department, 2008).
- Smooth Green Snake (*Opheodrys vernalis*) Native Species Status 2—This snake occupies forested areas of the foothills and montane zones, preferring to spend much of its time under rocks, logs, and other objects. It is usually associated with lush vegetation. Two subspecies occur in Wyoming. *O. vernalis vernalis*, the eastern smooth green snake, is a relict population that occurs only in the Black Hills of Wyoming and South Dakota. *O. vernalis blanchardi* is the western subspecies, and can be found in southeast and south-central Wyoming. Additionally, the smooth green snake occurs in parts of Canada, the northeastern and north-central United States, and as far west as Utah, Idaho and New Mexico. In the west, the snake's distribution is highly disjointed (Wyoming Game and Fish Department, 2008).
- Yellow-Billed Cuckoo Native Species Status 2—The Yellow-billed cuckoo nests primarily in large stands of cottonwood-riparian habitat below 2,100 m [7,000 ft], including such habitats that occur in urban areas. It is a riparian obligate species that prefers extensive areas of dense thickets and mature deciduous forests near water, and requires low, dense, shrubby vegetation for nest sites (Wyoming Game and Fish Department, 2008).
- Greater Sage Grouse (*Centrocercus urophasianus*) Native Species Status 2—Sage grouse depend on a variety of sagebrush community types and associated habitats, including basin-prairie and mountain foothills shrub lands, wet-moist meadows. Alfalfa and irrigated meadows also serve as habitat when immediately adjacent to sagebrush. Sage grouse use different habitats during different times of the year (Wyoming Game and Fish Department, 2008).
- Bald Eagle (*Haliaeetus leucocephalus*) Native Species Status 2—The Bald Eagle nests near large lakes and rivers in forested habitat where adequate prey and old, large-diameter cottonwood or conifer trees are available for nesting. Highly productive nesting areas in the Greater Yellowstone Area were found to have open water available in winter, low severity of early spring weather, limited human activity, and high sinuosity and an abundance of islands, riffles, runs, and pools in the river. Migrating and wintering eagles congregate near open water areas where concentrations of prey are available, such as carcasses of game animals, and spawning areas for kokanee, trout, and other fish (Wyoming Game and Fish Department, 2008).
- Trumpeter Swan (*Cygnus buccinator*) Native Species Status 2—The Trumpeter Swan inhabits shallow marshes, ponds, lakes, and river oxbows. It prefers stable, quiet, and shallow waters where small islands, muskrat houses, or dense emergent vegetation provide nesting and loafing sites. Nutrient-rich waters, with dense aquatic plant and invertebrate growth, provide the most suitable habitat. Adequate forage in the prenesting period (April to May) is critical for nesting success. Winter habitat must provide extensive beds of aquatic plants that remain ice free. In Wyoming, cold

temperatures and ice restrict trumpeters to sites where geothermal waters, springs, or outflow from dams maintain ice-free areas (Wyoming Game and Fish Department, 2008).

- **Fringed Myotis (*Myotis thysanodes*) Native Species Status 2**—The fringed myotis is found in a wide range of habitats, including coniferous forests, woodlands, grasslands, and shrublands, although it is probably most common in xeric woodlands, such as juniper, ponderosa pine, and Douglas fir. It typically forages over water, along forest edges, or within forests and woodlands. During summer, it uses a variety of roosts, including rock crevices, tree cavities, caves, abandoned mines, and buildings. During winter, it hibernates in caves, abandoned mines, and buildings (Wyoming Game and Fish Department, 2008).
- **Long-Eared Myotis (*Myotis evotis*) Native Species Status 2**—The long-eared myotis primarily inhabits coniferous forest and woodland, including juniper, ponderosa pine, and spruce fir. It typically forages over rivers, streams, and ponds within the forest-woodland environment. During summer, it roosts in a wide variety of structures, including cavities in snags, under loose bark, stumps, buildings, rock crevices, caves, and abandoned mines. During winter, it is thought to hibernate primarily in caves and abandoned mines (Wyoming Game and Fish Department, 2008).
- **Long-Legged Myotis (*Myotis volans*) Native Species Status 2**—The long-legged myotis inhabits open, mature forest with standing dead trees, including montane and subalpine forest and ponderosa pine and juniper woodlands, primarily from 1,500 m to more than 3,300 m [5,000 to more than 11,000 ft]. It usually forages over open areas such as campgrounds and small forest clearings; over vegetated riparian areas; and within, above, and under the forest canopy. During summer, it roosts in tree cavities, buildings, rock crevices, caves, abandoned mines, and under loose bark. During winter, it hibernates primarily in caves and abandoned mines (Wyoming Game and Fish Department, 2008).
- **Pallid Bat (*Antrozous pallidus*) Native Species Status 2**—The pallid bat generally inhabits low desert shrublands, juniper woodlands, and grasslands and occasionally cottonwood riparian zones in those habitats. It is most common in low, arid regions with rocky outcroppings, particularly near water. During summer, it usually roosts in rock crevices and buildings, but also uses rock piles, tree cavities, shallow caves, and abandoned mines (Wyoming Game and Fish Department, 2008).
- **Spotted Bat (*Euderma maculatum*) Native Species Status 2**—The spotted bat occupies a wide variety of habitats, from desert scrub to coniferous forest, although it is most often observed in low deserts and basins and juniper woodlands. It roosts in cracks and crevices in high cliffs and canyons. It also may occasionally roost in buildings, caves, or abandoned mines, although cliffs are the only roosting habitat in which reproductive females have been documented (Wyoming Game and Fish Department, 2008).
- **Townsend's Big-Eared Bat (*Plecotus townsendii*) Native Species Status 2**—The Townsend's big-eared bat occupies a variety of xeric to mesic habitats, including coniferous forests, juniper woodlands, deciduous forests, basins, and desert shrublands, and is absent only from the most extreme deserts and highest elevations. However, this species requires caves or abandoned mines for roost sites during all seasons and

stages of its life cycle, and its distribution is strongly correlated with the availability of these features (Wyoming Game and Fish Department, 2008).

3.2.6 Meteorology, Climatology, and Air Quality

3.2.6.1 Meteorology and Climatology

Wyoming's elevation results in relatively cool temperatures. Much of the temperature variations within the state can be attributed to elevation with average values dropping 1 to 2 °C [1.8 to 3.6 °F] per 300 m [1,000 ft] (National Climatic Data Center, 2005). Summer nights are normally cool although daytime temperatures may be quite high. The fall, winter, and spring can experience rapid changes with frequent variations from cold to mild periods. Freezes in early fall and late spring are typical and result in long winters and a short growing season. In the mountains and high valleys, freezes can occur any time in the summer. During winter warm spells, nighttime temperatures can remain above freezing. Valleys protected from the wind by mountain ranges can provide ideal pockets for cold air to settle and temperatures in the valley can be considerably lower than on nearby mountainsides. Table 3.2-6 identifies two climate stations located in the Wyoming West Uranium Milling Region. Climate data for these stations are found in the National Climatic Data Center's Climatology of the United States No. 20 Monthly Station Climate Summaries for 1971–2000 (National Climatic Data Center, 2004). This summary contains climate data for 4,273 stations throughout the United States and some territories. Table 3.2-7 contains temperature data for two stations in the Wyoming West Uranium Milling Region.

Table 3.2-6. Information on Two Climate Stations in the Wyoming West Uranium Milling Region*

Station (Map Number)	County	State	Longitude	Latitude
Gas Hills 4 E (042)	Fremont	Wyoming	107°31W	42°50N
Jeffrey City (049)	Fremont	Wyoming	107°50W	42°30N

*National Climatic Data Center. "Climatology of the United States No. 20: Monthly Station Climate Summaries, 1971–2000." Asheville, North Carolina: National Oceanic and Atmospheric Administration. 2004.

Table 3.2-7. Climate Data for Stations in the Wyoming West Uranium Milling Region*

		Gas Hills 4 E	Jeffrey City
Temperature (°C)†	Mean—Annual	5.5	5.3
	Low—Monthly Mean	-7.0	-7.0
	High—Monthly Mean	19.5	19.0
Precipitation (cm)‡	Mean—Annual	24.9	27.1
	Low—Monthly Mean	0.86	0.89
	High—Monthly Mean	3.33	5.71
Snowfall (cm)	Mean—Annual	154	143
	Low—Monthly Mean	0	0
	High—Monthly Mean	34.3	26.9

*National Climatic Data Center. "Climatology of the United States No. 20: Monthly Station Climate Summaries, 1971–2000." Asheville, North Carolina: National Oceanic and Atmospheric Administration. 2004.

†To convert Celsius (°C) to Fahrenheit (°F), multiply by 1.8 and add 32.

‡To convert centimeters (cm) to inches (in), multiply by 0.3937

Precipitation within Wyoming varies with spring and early summer being the wettest time for much of the state. Mountain ranges are generally oriented in a north-south direction. This is perpendicular to the prevailing westerlies. Therefore, these mountains often act as moisture barriers. Air currents for the Pacific Ocean rise and drop much of their moisture along the western slopes of the mountains. Summer showers are frequent but typically result in rainfall amounts of a few hundredths of an inch. Usually several times a year in the state, local thunderstorms will result in 2.5 to 5 cm [1 to 2 in] of rain in a 24-hour period. On rare occasions, rainfall in a 24-hour period can reach 7.5 to 12.5 cm [3 to 5 in] (National Climatic Data Center, 2005). Heavy rains can create flash flooding in headwater streams, and this flooding intensifies if these storms coincide with snow pack melting. Table 3.2-7 contains precipitation data for two stations in the Wyoming West Uranium Milling Region. The wettest month for both stations identified in Table 3.2-7 is May, which based on the snow depth data, coincides with snow pack melting (National Climatic Data Center, 2004). Both of these stations are in Fremont County. Data from National Climatic Data Center's Storm Events Database from 1950 to 2007 indicate that the vast majority of thunderstorms in Fremont County occur between June and September with the most occurring in July (National Climatic Data Center, 2007).

Hailstorms are the most destructive storm event for Wyoming. Most hailstorms pass over open rangeland with minimal impact. When a hailstorm passes over a city or farmland, the property and crop damage can be severe. Most of the severe hailstorms occur in the southeast corner of the state.

Low elevations typically experience light to moderate snowfall from November to May. Snowfall within Wyoming varies by location with the mountain ranges typically receiving the most. Significant storms of 25 to 40 cm [10 to 16 in] of snowfall are infrequent outside of the mountains. Wind often coincides or follows snowstorms and can form snow drifts several meters deep. Snow can accumulate to considerable depths in the high mountains. Blizzards that last more than 2 days are uncommon. Table 3.2-7 contains snowfall data for two stations in the Wyoming West Uranium Milling Region.

Wyoming is windy and ranks first in the US with an annual average speed of 6 m/s [12.9 mph]. During winter Wyoming frequently experiences periods where wind speed reaches 13 to 18 m/s [30 to 40 mph] with gusts to 22 to 27 m/s [50 or 60 mph] (National Climatic Data Center, 2005). Prevailing wind direction varies by location but usually ranges between west-southwest through west to northwest. Because the wind is normally strong and constant from those directions, trees often lean to the east or southeast.

The pan evaporation rates for the Wyoming West Uranium Milling Region range from about 76 to 127 cm [30 to 50 in] (National Weather Service, 1982). Pan evaporation is a technique that measures the evaporation from a metal pan typically 121 cm [48 in] in diameter and 25 cm [10 in] tall. Pan evaporation rates can be used to estimate the evaporation rates of other bodies of water such as lakes or ponds. Pan evaporation rate data is typically available only from May to October. Freezing conditions often prevent collection of quality data during the other parts of the year.

3.2.6.2 Air Quality

As described in Section 1.7.2.2, the permitting process is the mechanism used to address air quality. If warranted, permits may set facility air pollutant emission levels, require mitigation measures, or require additional air quality analyses. Except for Indian Country, New Source

Description of the Affected Environment

Review permits in Wyoming are regulated under the EPA-approved State Implementation Plan. For Indian Country in Wyoming, the New Source Review permits are regulated under 40 CFR 52.21 (EPA, 2007a).

State Implementation Plans and permit conditions are based in part on federal regulations developed by the EPA. As promulgated in 40 CFR Part 50, National Primary and Secondary Ambient Air Quality Standards (NAAQS), the NAAQS define acceptable ambient air concentrations for six common nonradiological air pollutants: nitrogen oxides, ozone, sulfur oxides, carbon monoxide, lead, and particulates. Primary NAAQS are established to protect public health, and secondary NAAQS are established to protect public welfare by safeguarding against environmental and property damage. Primary and secondary NAAQS are presented in Table 3.2-8. Some pollutants have multiple standards. Particulates are divided into two categories: PM₁₀ defined as particulate matter smaller than 10 micrometers [3.9×10^{-4} in] and PM_{2.5} defined as particulate matter smaller than 2.5 micrometers [9.8×10^{-5} in]. In June 2005,

Table 3.2-8. National Ambient Air Quality Standards*

Pollutant	Primary Standards	Averaging Times	Secondary Standards
Carbon Monoxide	9 ppm (10,000 $\mu\text{g}/\text{m}^3$)†	8 hours‡	None
	35 ppm (40,000 $\mu\text{g}/\text{m}^3$)†	1 hour‡	None
Lead	1.5 $\mu\text{g}/\text{m}^3$ †	Quarterly average	Same as primary
Nitrogen Dioxide	0.053 ppm (100 $\mu\text{g}/\text{m}^3$)†	Annual (arithmetic mean)	Same as primary
Particulate Matter 10- μm diameter (PM ₁₀)	150 $\mu\text{g}/\text{m}^3$ †	24 hours§	Same as primary
Particulate Matter 2.5- μm diameter (PM _{2.5})	15.0 $\mu\text{g}/\text{m}^3$ †	Annual (arithmetic mean)	Same as primary
	35 $\mu\text{g}/\text{m}^3$ †	24 hours¶	Same as primary
Ozone	0.08 ppm	8 hours#	Same as primary
	0.12 ppm	1 hour**	Same as primary
Sulfur Oxides	0.03 ppm	Annual (arithmetic mean)	Not applicable
	0.14 ppm	24 hours‡	Not applicable
	Not applicable	3 hours‡	0.5 ppm (1,300 $\mu\text{g}/\text{m}^3$)†

*Modified from U.S. Environmental Protection Agency. "National Ambient Air Quality Standards (NAAQS)." 2007. <<http://www.epa.gov/air/criteria.html>> (15 October 2007).

†Multiply $\mu\text{g}/\text{m}^3$ value by 2.7×10^{-8} to convert units to oz/yd³

‡Not to be exceeded more than once per year

§Not to be exceeded more than once per year on average over 3 years.

|| To attain this standard, the 3-year average of the weighted annual mean PM_{2.5} concentrations from single or multiple community-oriented monitors must not exceed 15.0 $\mu\text{g}/\text{m}^3$.

¶To attain this standard, the 3-year average of the 98th percentile of 24-hour concentrations at each population-oriented monitor within an area must not exceed 35.0 $\mu\text{g}/\text{m}^3$ (effective December 17, 2006).

#To attain this standard, the 3-year average of the fourth highest daily maximum 8-hour average ozone concentrations measured at each monitor within an area over each year must not exceed 0.08 ppm.

** (a) The standard is attained when the expected number of days per calendar year with maximum hourly average concentrations above 0.12 ppm is ≤ 1 , as determined by Appendix H. (b) As of June 15, 2005, the U.S. Environmental Protection Agency revoked the 1-hour ozone standard in all areas except the fourteen 8-hour ozone nonattainment Early Action Compact Areas.

EPA revoked the 1-hour ozone standard nationwide in all locations except certain Early Action Compact Areas. None of the 1-hour ozone Early Action Compact Areas are in Wyoming. States may develop standards that are stricter or supplement the NAAQS. Wyoming has a more restrictive annual average standard for sulfur dioxide at $60 \mu\text{g}/\text{m}^3$ [$1.6 \times 10^{-6} \text{ oz}/\text{yd}^3$] and a supplemental $50 \mu\text{g}/\text{m}^3$ [$1.3 \times 10^{-6} \text{ oz}/\text{yd}^3$] PM_{10} standard with an annual averaging time (Wyoming Department of Environmental Quality, 2006).

As promulgated in 40 CFR Part 52, Prevention of Significant Deterioration requirements identify maximum allowable increases in concentrations for particulate matter, sulfur dioxide, and nitrogen dioxide for areas designated as attainment. Different increment levels are identified for different classes of areas. Table 3.2-9 contains the maximum allowable Prevention of Significant Deterioration increments for Class I and Class II areas. Class I areas are locations with special natural, recreational, scenic, or historic value such as national parks or wilderness

Table 3.2-9. Allowable Prevention of Significant Deterioration Class I and Class II Areas*

Pollutant	Class I ($\mu\text{g}/\text{m}^3$)†	Class II ($\mu\text{g}/\text{m}^3$)†	Measurement
Nitrogen Dioxide (NO_2)	2.5	25	Annual average
$\text{PM}_{10}\ddagger$	4	17	Annual average
	8	30	24 hours‡
Sulfur Dioxide (SO_2)	2	20	Annual average
	5	91	24 hours§
	25	512	3 hours§

*Modified from Code of Federal Regulations. "Prevention of Significant Air Deterioration of Air Quality." Title 40—Protection of the Environment, Part 52. Washington, DC: U.S. Government Printing Office. 2005.

† Multiply $\mu\text{g}/\text{m}^3$ value by 2.7×10^{-8} to convert units to oz/yd^3

‡ Not to be exceeded on more than 1 day/year on the average over 3 years.

§ Not to be exceeded more than once per year.

areas and have the most stringent set of allowable increments. Most other areas in the United States are categorized as Class II areas and have the less stringent set of allowable increments. One goal identified in the Clean Air Act is to address visibility impairment from haze at the Prevention of Significant Deterioration Class I areas in the country. Regional haze is visibility impairment caused by cumulative air pollutant emissions from numerous sources over a wide geographic area (EPA, 1999). Key contributors to regional haze are sulfur dioxide, nitrogen oxides, and particulate matter. One source of particulate matter is soil dust or fugitive dust. The EPA in 40 CFR Part 51 requires states to address regional haze in their implementation plans.

The Wyoming West Uranium Milling Region air quality description focuses on two topics: NAAQS attainment status and PSD classifications in the region.

NAAQS compliance attainment status is typically determined at the county level. Each NAAQS pollutant is designated into one of the following categories: attainment, nonattainment, or maintenance. Areas are designated as attainment for a particular pollutant if atmospheric concentrations meet NAAQS. If atmospheric concentrations of a pollutant do not meet NAAQS, that area is designated as nonattainment for that pollutant. The maintenance category describes areas formerly designated as nonattainment, but that now meet NAAQS




Figure 3.2-15. Air Quality Attainment Status for Wyoming and Surrounding Areas (EPA, 2007)

requirements. Figure 3.2-15 identifies counties in Wyoming and surrounding areas that are partially or entirely designated as nonattainment or maintenance for NAAQS at the time this Draft GEIS was prepared (EPA, 2007b). All of the area within the Wyoming West Uranium Milling Region is classified as attainment. In fact, Wyoming only has one area that is not in attainment. The City of Sheridan in Sheridan County is designated as nonattainment for PM₁₀. Portions of several Colorado counties along the southern Wyoming border are classified as not in attainment. However, the southern boundary of the Wyoming West Uranium Milling Region is north of the Wyoming/Colorado border.

Table 3.2-10 identifies the Prevention of Significant Deterioration Class I areas in Wyoming. These areas are shown in Figure 3.2-16. There are no Class I areas in the Wyoming West Uranium Milling Region (40 CFR Part 81).

Table 3.2-10. U.S. Environmental Protection Agency Class I Prevention of Significant Deterioration Areas in Wyoming*

Bridger Wilderness
Fitzpatrick Wilderness
Grand Teton National Park
North Absaroka Wilderness
Teton Wilderness
Washakie Wilderness
Yellowstone National Park

*Modified from Code of Federal Regulations. "Prevention of Significant Air Deterioration of Air Quality." Title 40—Protection of the Environment, Part 81. Washington, DC: U.S. Government Printing Office. 2005.

EPA also encourages states to work with tribes and federal agencies in regional partnerships to address the regional haze issue. Wyoming is a member of the Western Regional Air Partnership. Also, specific provisions in 40 CFR Part 51 allow nine western states, including Wyoming, to implement the recommendations of the Grand Canyon Visibility Transport Commission within the regional haze program.


3.2.7 Noise

Noise is technically defined as unwanted sound. Noise is a potential occupational hazard because prolonged exposure to noise may cause long-term hearing loss. In the United States, noise levels are regulated at the federal level by the Occupational Health and Safety Administration and the Mining Safety and Health Administration (Bauer and Kohler, 2000). To provide a sense of magnitude, noise levels associated with common activities are presented in Figure 3.2-17.

Existing ambient noise levels can be used to establish baseline conditions and determine potential site-specific disturbances associated with ISL milling activities. The Wyoming West Uranium Milling Region is predominantly rural and undeveloped. Rural areas tend to be quiet, open sagebrush-grass and forested areas where natural phenomena such as wind, rain,

What are sound and noise?

When an object vibrates, some of the energy causes air molecules to vibrate. Nearby people or animals translate these vibrations into sound using the eardrum and brain. Noise is simply unwanted sound. Sound waves are characterized by frequency and measured in hertz (Hz); sound pressure is expressed as decibels (dB). Noises that are perceptible to human hearing range vary from 31 to 20,000 Hz. Audible sounds (those that can be heard) range from about 60 dB at a frequency of 31 Hz to less than about 1 dB between 900 and 8,000 Hz. Noise levels for perceptible frequencies are typically reported in A-weighted decibels to account for the way people respond to noise; this type of measurement assumes a human receptor to a particular noise-producing activity.



WYOMING

30 15 0 30 Miles
 30 15 0 30 Kilometers

Prevention of Significant Deterioration
 Class I Area
 Wyoming East Milling Region
 Wyoming West Milling Region

South Dakota - Nebraska Milling Region
 Interstate Highway
 Water bodies (Lakes, Bays, ...)
 State Boundary
 Counties

Figure 3.2-16. Prevention of Significant Deterioration Class I Areas in the Wyoming East Uranium Milling Region and Surrounding Areas (40 CFR Part 81)

insects, birds, and other wildlife account for most natural background sounds. Baseline noise levels for typical undeveloped desert or arid environments range from day-night sound levels of 22 dB on calm days to 38 dB on windy days (Brattstrom and Bondello, 1983; DOE, 2007).

Larger communities in the region include Riverton and Lander, with populations of between 5,000 and 10,000. Fort Washakie (population about 1,500), the location of the headquarters for the Wind River Indian Reservation is within the region. In addition, Rawlins (population about 8,500) is just east of the southeast corner of the region on Interstate 80 (see Section 3.2.10). In these more urbanized areas, ambient noise levels would be expected to be influenced by noise generating activities such as street noise, traffic, emergency vehicles, and construction equipment. Noise levels in these types of suburban residential/urban areas range from 45 to about 78 dB, with lower noise levels at night (Washington State Department of Transportation, 2006).

As described in Section 2.8, several highways cross the region, including U.S. Highways 20, 26, and 287, as well as Interstate 80. A summary of noise effects on wildlife populations (Federal Highway Administration, 2004) includes reference to measured average traffic noise levels at 15 m [50 ft] of 54–62 dBA for passenger cars and 58–70 dBA for heavy trucks (Federal Highway Administration, 2004) along Interstate 80. Baseline ambient noise levels would be similar or less for the United States and state highways in the region, as they are mostly undivided highways and tend to carry less traffic (particularly heavy trucks) than a major interstate highway like Interstate 80. For example, a 2005 traffic analysis at Interstate 80 milepost 208.65 just west of Rawlins indicates an average traffic count of about 12,400 vehicles per day. Of this, almost 50 percent was heavy truck traffic (Wyoming Department of Transportation, 2005). In comparison, for U.S. Highway 26 milepost 125.75 northwest of Riverton, the 2005 traffic count was about 3,700 vehicles with almost 90 percent passenger truck and car traffic (Wyoming Department of Transportation, 2005).

The two principal uranium districts in the Wyoming West Uranium Milling Region (the Great Divide Basin in the southeast part of the region and the Wind River Basin in the northeast part of the region) are located more than about 30 to 80 km [20 to 50 mi] from the larger communities, in rural undeveloped areas where the ambient noise levels would be expected to be low. There are a number of smaller communities along highways and roads through the uranium districts, including Jeffrey City and Bairoil near U.S. Highway 287 in the Great Divide Basin and Ervay and Sand Draw in the Wind River Basin, where noise levels would be expected to be slightly higher as a result of human activities. Areas of special sensitivity may be located on the Wind River Indian Reservation in the northwest corner of the region, but the reservation boundary is more than 16 km [10 mi] from

How is sound measured?

The human ear responds to a wide range of sound pressures. The range of sounds people normally experience extends from low to high pressures by a factor of 1 million. Sound is commonly measured using decibels (dB). Another common sound measurement is the A-weighted sound level (dBA). The A-weighting measures different sound frequencies and the variation of the human ear's response over the frequency range. Higher frequencies receive less A-weighting than lower ones. Noise levels are often reported as the equivalent sound level (DOE, 2007). The equivalent sound level is expressed as an A-weighted sound level over a specified period of time—usually 1 or 24 hours. The equivalent sound level is an equivalent steady sound level that, if it continued during a specified time period, would contain the same total energy as the actual time-varying sound over the monitored or modeled time period. Noise levels are also expressed as day-night sound levels: the average of the day and nighttime A-weighted sound level with a built-in penalty of 10 dBA at night when noise levels are likely lower. The day-night sound level is particularly useful for evaluating community-level noise effects. If noise is regulated, municipalities often have local ordinances specifying upper limits on evening noise levels, with specific hours for residential and commercial zones.

Description of the Affected Environment

the closest potential uranium ISL facility near Sand Draw, and more than 50 km [30 mi] from the center of the two uranium districts.

COMMON SOUNDS	DECIBELS*	EFFECT
Jet Operation	140	Painfully Loud
	130	
Jet Takeoff Thunder Rock Concert	120	Maximum Vocal Effort
Pile Drivers	110	
Garbage Truck	100	
Heavy Truck (50 ft)	90	Very Annoying Hearing Damage at 8 hr
Alarm Clock Hair Dryer	80	Annoying
Freeway Traffic Man's Voice (3 ft)	70	Telephone Use Difficult
Air Conditioning Unit (20 ft)	60	Intrusive
Light Auto Traffic (100 ft)	50	Quiet
Living Room Quiet Office	40	
Library Soft Whisper (15 ft)	30	Very Quiet
Broadcasting Studio	20	
	10	Just Audible

*To the ear, each 10 dB increase seems twice as loud. 70 dB is the point at which noise begins to harm hearing.

Figure 3.2-17. Comparison of Noise Levels Associated With Common Activities (After EPA, 1981)

3.2.8 Historical and Cultural Resources

The following summarizes the historical and cultural resources background and legislation and authorities regarding historical and cultural resources for the Uranium GEIS regions in the states of Nebraska, New Mexico, South Dakota, and Wyoming. The information is provided on a state-by-state basis rather than by the regions of interest as the historical and cultural resource information and agencies are organized at the state level.

3.2.8.1 Cultural Resources Overview

The Wyoming State Historic Preservation Office (SHPO) administers and is responsible for oversight and compliance with the National Register of Historic Places (NRHP), compliance and review for Section 106 of the National Historic Preservation Act (NHPA), and Traditional Cultural Properties review, enforcement of the Native American Graves Protection and Repatriation Act (NAGPRA) and compliance with other federal and state historic preservation laws, regulations, and statutes. The Wyoming SHPO and BLM have also entered into a Programmatic Agreement that describes the manner in which the Wyoming SHPO and the Wyoming BLM would interact and cooperate under the BLM national Programmatic Agreement. State level agreements between Wyoming and the National Resource Conservation Service (NRCS) and the USFS are in draft form. Wyoming SHPO's webpage with links to all of their resources can be found at: <http://wyoshpo.state.wy.us/>. The State of Wyoming also has a law pertaining to archaeological sites and human remains, entitled Archaeological Sites (Wyoming Statute Ann. §36-1-114, et seq).

A brief discussion of cultural and historical resource management processes is included in Appendix D.

The following provides a brief overview of prehistoric and historical cultures recognized in the central and northern plains region which includes the Wyoming West Uranium Milling Region. Figure 3.2-18 illustrates the division of the plains into regional subdivisions. The dating of cultural periods for the prehistoric period is provided in years before present (BP). Most prehistoric archaeological sites are concentrated along major river systems and their tributaries, but can also be found along many drainage basins in the eastern and central portions of the state.

Paleoindian Big Game Hunters (12,000 to 6,500 BP). The earliest well-defined cultural tradition in the northern and central plains region is the Paleoindian. Early humans entered the plains shortly after deglaciation allowed movement onto the northern and central plains sometime after 14,000 BP. A variety of cultures, each defined by the presence of distinctive, lanceolate projectile points, are recognized during the Paleoindian period: Clovis, Goshen, Folsom, Hell Gap-Agate Basin, Alberta, Cody Complex, and the late Paleoindian-Early Archaic Foothills/Mountain Complex. Most post-Clovis Paleoindian sites on the northern and upper central plains are known from bison kill sites. The Clovis culture (12,000 to 10,000 BP) is recognized by a distinctive projectile point style and a subsistence mode heavily reliant on hunting large, now-extinct mammals, notably mammoth, which became extinct at the end of the Clovis period, and ancient bison. The poorly defined Goshen Complex is found at the Carter/Kerr-McGee site in northeastern Wyoming and the Jim Pitts site in the Black Hills at the Wyoming-South Dakota border. Goshen is technologically similar to Clovis and may be contemporary with Clovis and perhaps Folsom. The Folsom culture (ca. 10,000 to 8,500 BP) is




Figure 3.2-18. General Location of Native American Plains Tribes

also known for a distinctive fluted, projectile point style and has been found at the Carter/Kerr-McGee site associated with bison and red ochre deposits. Folsom subsistence is also characterized by reliance on large game (the ancient bison). Folsom sites consist of campsites and kill sites. The latter tend to be located near cliffs and around water, such as ponds and springs.

The Hell Gap-Agate Basin Complex, Alberta Complex, and Cody Complex are widely distributed in the northern and central portions of the southern plains region at the Agate Basin, Hell Gap, and Carter/Kerr-McGee archaeological sites in eastern Wyoming. These late Paleoindian cultural complexes are, in their earliest forms, a continuation of preceding Paleoindian hunting traditions. The distinctive projectile point forms which define these cultural complexes in central and eastern Wyoming and western South Dakota are, in comparison to earlier Clovis, Goshen, and Folsom, much more restricted in geographic distribution. Toward the end of the Paleoindian period, however there is a transition in subsistence modes following the extinction of the ancient bison and the transition to hunting the modern form of bison ultimately leading to the transition to Archaic broad-spectrum foraging. Post molds and stone circles suggesting the presence of ephemeral shelters are sometimes found, primarily toward the end of the period.

The late Paleoindian Foothills/Mountain Complex is characterized by a reliance on medium-sized game animals rather than big game hunting. Sites are found in upland, mountainous regions leading some to suggest that Paleoindian groups may have split into lowland big game hunters and upland/mountain small and medium game hunters (Frison, 1991). The upland/mountain sites show increased use of small seed-bearing plants as indicated by the presence of groundstone implements, and suggests the presence of an early archaic lifestyle. Habitation sites of this complex are found in rockshelters and caves such as Mummy Cave in the Absaroka Mountains of northwestern Wyoming.

Archaic Foragers (6,500 to 2,500 BP). The Plains Archaic period represents the continuation of change in subsistence and settlement linked to an increasingly arid environment that occurs in the latter portion of the preceding late Paleoindian cultures. At the end of the Paleoindian period there is also a change in projectile point styles from lanceolate to somewhat smaller corner- and side-notched projectile points suggesting that the atlatl (spearthrower) was in use. Distinctive Archaic cultures, from early to late, include Mummy Cave, Oxbow, McKean, and Pelican Lake complexes and are found throughout the northern plains. Large bison kill sites, characteristic of the preceding Paleoindian period are virtually absent. Hunting and gathering wild plant foods is the primary mode of subsistence. Dietary breadth, indicated by increasing diversity and numbers of subsistence items, is believed to expand significantly with more medium and small mammals being hunted and the introduction of seed-bearing plants dietary staples indicated by the introduction of stone seed-grinding implements. The Early Archaic Medicine House site in the southeastern Wyoming contained evidence of structures, hearths, storage pits, and milling basins. At the McKean site in the Black Hills of Wyoming, a shallow pithouse was found. Through time, settlement is increasingly tethered to highly productive resource areas and sites tend to become larger and increasingly complex indicating the presence of somewhat more sedentary lifestyles relative to earlier periods. Settlement is focused on river valleys and elevated areas. Artifact styles, principally projectile points, become increasingly diversified suggesting increasing regionalization and cultural differentiation. In southeastern Wyoming, Pelican Lake projectile points are sometimes found in association with stone circles, firepits, and pithouses.

Late Prehistoric/Plains Woodland (2,500 to 300 BP). Early in the period, the preceding late Archaic broad-spectrum foraging subsistence and settlement patterns continue with little

change. In the Northern Plains, the Besant and Avonlea Complexes continued the Archaic lifestyles virtually unchanged until contact with European and American cultures. A significant technological change from atlatl to bow and arrow occurs during the Late Prehistoric period. Subsistence focused on scheduled small and medium game hunting, gathering plant foods, and bison hunting according to a seasonal round. In central and northeastern Wyoming, a basic hunting and gathering lifestyle differing little from the preceding Late Archaic period predominates. Although eastern Wyoming is considered peripheral to the eastern Woodland tradition, Woodland pottery is sometimes found in association with Besant points in the northern plains. The Butler-Risser site south of Casper, Wyoming, contained both Besant points and pottery. Food procurement and site location during this period appears to be focused primarily on elevated landforms near larger riverine systems and tributaries with increasing utilization of upland resources later in time. The Late Prehistoric/Plains Woodland of Wyoming is also characterized by the appearance of ceramics late in the period (Besant and Avonlea Complexes), introduced from the Eastern Woodland cultural area. The late Avonlea Complex and later Old Woman Complex sites in northern Wyoming contain artifact types that suggest a high degree of specialization in hunting large, upland game animals, primarily bison.

In the eastern portions of Wyoming the Upper Republican phase (ca. 1000–300 BP) is characterized by the presence of seasonal or permanent sedentary villages. These sites are usually on ridges and bluffs and have evidence of domesticated plants (corn, beans, squash, and sunflowers). Although horticulture was an important part of the subsistence base, wild plants and game animals formed a substantial part of the diet. Storage pits for food and other items are located within the structures and grinding tools are common. Pottery was diverse with globular jars and decorated exterior rims are common. The later Dismal River Aspect (ca. 500–300 BP) in southeastern Wyoming is focused primarily on hunting and gathering with only limited evidence of horticultural pursuits and a distinctive form of pottery.

In the 1500s to early 1700s AD, large migrations by Indian tribes occurred. The ancestors of modern the Apache, Arapaho, Comanches, Apache-Kiowas, and Kiowas migrated southward through western Wyoming in the 1500s and 1600s.

Post-Contact Tribes (300 to 100 BP). The post-contact period on the northern plains is that period after initial contact with Europeans and Americans. Although Euro-American trade goods may have appeared as early as the mid-1600s, the earliest documented contact in the northern and central plains is by Spanish and French explorers in the early 1700s AD. The horse appears to have been introduced at about the same time. The lifeways of the late Avonlea and post-Avonlea/Old Woman nomadic bison-hunting cultural complexes in central and northeastern Wyoming and the Upper Republican and Dismal River horticulturalists of eastern and southeastern Wyoming appear to have continued well into the mid to late 1700s AD. At the time of European exploration, the Dakota and Nakota moved into eastern Wyoming from what is now Minnesota. The Shoshone were present in southeastern Wyoming in the 1600s and 1700s. About this time the Crow moved into northeastern and north-central Wyoming and the Apache-Kiowas moved out of the Black Hills into southeastern Wyoming. The Apache-Kiowa migration through the Black Hills was followed by that of the Cheyenne who moved through western South Dakota and then into central Wyoming where they were joined by the Arapaho who settled in southern Wyoming (Reher, 1977). By the mid-1800s, much of the eastern and central portions of the state was occupied by nomadic Siouan-speaking tribes, primarily the Hunkpapa, Minneconjou, Brule, and Oglala.

Europeans and Americans (300 to 100 BP). The earliest European presence in Wyoming was by French explorers of the de la Vérendrye family in 1743. In 1803, the United States

completed the purchase of the Louisiana Territory from France. Early expeditions and trappers provide descriptions of varying quality for some of the early historical tribes in the region. In the later 1700s and early 1800s more intensive contact and settlement occurred first through missionaries and the fur trade period in the 1810s through the 1840s. In 1807 Manuel Lisa of St. Louis established a trading post on the Bighorn River. Others, including Jedediah Smith, fur trading companies quickly spread along the major river systems of Wyoming. Each year the fur traders and trappers would establish a rendezvous site where they would gather. Rendezvous sites are known throughout much of central and western Wyoming. By the late 1830s, the fur trade in Wyoming was in decline. By the mid-1800s, missionary, settler, and military contacts led to increasing conflict with the Siouan tribes of Wyoming. The slowly increasing number of settlers passing through traditional tribal use areas on well-established trails in the mid-1800s led to increasing conflict over time. The establishment of military forts on tribal lands to protect the settlers was yet another irritant to tribes.

Treaties, notably the Fort Laramie Treaty of 1851 were signed with the intent of removing tribes from along the emigrant trails and to allow for the building of trails and forts to protect settlers moving west on the Texas, Oregon, California, Mormon, Bozeman, and Bridger Trails in central and eastern Wyoming. Continued conflict resulted in the creation of the Great Sioux Reservation bounded by the Missouri River on the east, the Big Horn Mountains on the west, and the 46th and 43rd parallels to the north and south, respectively. Continued conflict with the U.S. military over the failure of the government to abide by treaty obligations led to several punitive expeditions to return tribes to reservations. In 1874, General George Armstrong Custer led an expedition to the Black Hills of Wyoming and South Dakota where the presence of gold, previously only rumored, was confirmed. The intense interest by Americans to go to the Black Hills to mine for gold led to numerous treaty violations; the Black Hills region was, by treaty, part of the Sioux reservation. The continued conflict over the Black Hills, along with reduction of the buffalo herds, led to the final military conquest of the Great Sioux Nation and their confinement to small reservations. In November 1875, President Grant ordered the Indians of the Powder River and Big Horn country in eastern and central Wyoming to return to their tribal agencies. The Sioux refused and were forced militarily onto their reservations. The Black Hills gold rush facilitated the subsequent settlement of much of Wyoming and the development of towns and cattle ranching.

Ranching, a livelihood well suited to the grassland plains of Wyoming, was practiced by settlers by the early 1870s. Most of the early ranching occurred in well-watered areas along existing trail systems to facilitate moving cattle to market. The arrival of the railroads in 1868 (first the Union Pacific in southern Wyoming, then branch lines in other parts of Wyoming) led to increased settlement and opened Wyoming to a flood of new settlers. In the 1880s, farmers began homesteading much of the open range leading to conflict with ranchers over fencing. They settled mostly around well-watered regions, with many of the new farmers pursuing newly developed dry-land farming techniques. These homestead farmers began a period of extensive agriculture throughout the state that lasted from the 1880s to the 1930s. The Great Depression and the droughts that occurred at the same time led to the abandonment of many farms and the outmigration of a significant portion of Wyoming's population. Many of the individual homesteads were bought out in the 1930s and 1940s to create larger farms using mechanized equipment.

3.2.8.2 Historic Properties Listed in the National and State Registers

Table 3.2-11 includes a summary of sites in the Wyoming West Uranium Milling Regions that are listed on the Wyoming state and/or National Register of Historic Places. Most of the sites

Table 3.2-11. National Register Listed Properties in Counties Included in the Wyoming West Uranium Milling Region

County	Resource Name	City	Date Listed YYYY/MM/DD
Carbon	Duck Lake Station Site	Wamsutter	1978-12-06
Fremont	BMU Bridge Over Wind River	Ethete	1985-02-22
Fremont	Decker, Dean, Site (48FR916; 48SW541)	Honeycomb Buttes	1986-03-12
Fremont	Delfelder Schoolhouse	Riverton	1978-03-29
Fremont	ELY Wind River Diversion Dam Bridge	Morton	1985-02-22
Fremont	Fort Washakie Historic District	Fort Washakie	1969-04-16
Fremont	Green Mountain Arrow Site (48FR96)	Stratton Rim	1986-03-12
Fremont	Jackson Park Town Site Addition Brick Row	Lander	2003-02-27
Fremont	King, C.H., Company, and First National Bank of Shoshoni	Shoshoni	1994-09-08
Fremont	Lander Downtown Historic District	Lander	1987-05-05
Fremont	Quien Sabe Ranch	Shoshoni	1991-04-18
Fremont	Riverton Railroad Depot	Riverton	1978-05-22
Fremont	Shoshone-Episcopal Mission	Fort Washakie	1973-04-11
Fremont	South Pass	South Pass City	1966-10-15
Fremont	South Pass City	South Pass City	1970-02-26
Fremont	St. Michael's Mission	Ethete	1971-06-21
Fremont	Union Pass	Unknown	1969-04-16
Fremont	U.S. Post Office and Courthouse--Lander Main	Lander	1987-05-19
Fremont	Wind River Agency Blockhouse	Ft. Washakie	2000-12-23
Natrona	Archeological Site No. 48NA83	Arminto	1994-05-13
Natrona	Big Horn Hotel	Arminto	1978-12-18
Natrona	Bishop House	Casper	2001-03-12
Natrona	Bridger Immigrant Road--Waltman Crossing	Casper	1975-01-17
Natrona	Casper Army Air Base	Casper	2001-08-03
Natrona	Casper Buffalo Trap	Casper	1974-06-25
Natrona	Casper Federal Building	Casper	1998-12-21
Natrona	Casper Fire Department Station No. 1	Casper	1993-11-04
Natrona	Casper Motor Company--Natrona Motor Company	Casper	1994-02-23
Natrona	Chicago and Northwestern Railroad Depot	Powder River	1988-01-07
Natrona	Church of Saint Anthony	Casper	1997-01-30
Natrona	Consolidated Royalty Building	Casper	1993-11-04
Natrona	DUX Bessemer Bend Bridge	Bessemer Bend	1985-02-22
Natrona	Elks Lodge No. 1353	Casper	1997-01-30
Natrona	Fort Caspar	Casper	1971-08-12
Natrona	Fort Caspar (Boundary Increase)	Casper	1976-07-19
Natrona	Independence Rock	Casper	1966-10-15
Natrona	Martin's Cove	Casper	1977-03-08
Natrona	Masonic Temple	Casper	2005-08-24
Natrona	Midwest Oil Company Hotel	Casper	1983-11-17
Natrona	Natrona County High School	Casper	1994-01-07
Natrona	North Casper Clubhouse	Casper	1994-02-18
Natrona	Ohio Oil Company Building	Casper	2001-07-25

Table 3.2-11. National Register Listed Properties in Counties Included in the Wyoming West Uranium Milling Region (continued)

County	Resource Name	City	Date Listed YYYY/MM/DD
Natrona	Pathfinder Dam	Casper	1971-08-12
Natrona	Rialto Theater	Casper	1993-02-11
Natrona	Roosevelt School	Casper	1997-01-30
Natrona	South Wolcott Street Historic District	Casper	1988-11-23
Natrona	Split Rock, Twin Peaks	Muddy Gap	1976-12-22
Natrona	Stone Ranch Stage Station	Casper	1982-11-01
Natrona	Townsend Hotel	Casper	1983-11-25
Natrona	Tribune Building	Casper	1994-02-18
Sweetwater	Eldon—Wall Terrace Site (48SW4320)	Westvaco	1985-12-13

are located in Fremont County, at least 32 km [20 mi] west of the two uranium districts in the Gas Hills and near Crooks Gap.

3.2.8.3 Tribal Consultation

There are several Native American Tribes located within or immediately adjacent to the state of Wyoming that have interests in the state (Figure 3.2-19). These include the

- Arapaho Tribe of the Wind River Reservation
- Shoshone Tribe of the Wind River Reservation
- Cheyenne River Sioux
- Flandreau Santee Sioux
- Lower Brulé Sioux
- Oglala Sioux
- Rosebud Sioux
- Sisseton-Whapeton Oyate
- Standing Rock Sioux
- Yankton Sioux
- Crow Tribe of Montana

The Siouan tribes are located throughout South and North Dakota, and the Crow are located in Montana but have interests in Wyoming. Other Siouan-speaking tribes as well as other tribes in North Dakota, Wyoming, Montana, and Nebraska may have traditional land use claims in the Wyoming West Uranium Milling Region.

The U.S. government and the State of Wyoming recognize the sovereignty of certain Native American tribes. These tribal governments have legal authority for their respective reservations. Executive Order 13175 requires executive branch federal agencies to undertake consultation and coordination with Indian tribal governments on a government-to-government basis. NRC, as an independent federal agency, has agreed to voluntarily comply with Executive Order 13175.

In addition, the NHPA provides these tribal groups with the opportunity to manage cultural resources within their own lands under the legal authority of a Tribal Historic Preservation Officer (THPO). To date, no tribes in Wyoming have applied for status as a THPO as provided

Description of the Affected Environment

by the NHPA. Some tribes have historic and cultural preservation offices that are not recognized as THPOs, but they should be consulted where they exist. NRC, in meeting its responsibilities under the NHPA, contacts tribal cultural resources personnel as part of the consultation process, along with consulting with the Wyoming SHPO.




Figure 3.2-19. Regional Distribution of Native American Tribes in Wyoming, South Dakota, and Nebraska

3.2.8.4 Places of Cultural Significance

Traditional cultural properties are places of special heritage value to contemporary communities because of their association with cultural practices and beliefs that are rooted in the histories of those communities and are important in maintaining the cultural identity of the communities (Parker and King 1998; also see King, 2003). Religious places are often associated with prominent topographic features like mountains, peaks, mesas, springs and lakes. In addition shrines may be present across the landscape to denote specific culturally significant locations and vision quest sites where an individual can place offerings.

Information on traditional land-use and the location of culturally significant places is often protected information within the community (e.g., see King, 2003). Therefore, the information presented on religious places is limited to those that are identified in the published literature and

are therefore restricted to a few highly recognized places on the landscape within southwestern South Dakota.

There are no known culturally significant places in the NRHP or state register located in the Wyoming West Uranium Milling Region. However, the Lakota Sioux or other Sioux bands (Cheyenne River Sioux, Lower Brule Sioux, Oglala Sioux, Rosebud Sioux) along with the Crow Tribe, the Arapaho, the Kiowa and Wind River Shoshone who once occupied portions of the Wyoming West Uranium Milling Region consider the Black Hills in Wyoming and South Dakota, Devil's Tower in northeastern Wyoming, and Bear Butte in southwestern South Dakota to be culturally significant; these were once used for personal rituals, the Sun Dance and are the source of origin legends.

Areas of central and eastern Wyoming once used by these tribes may contain additional, undocumented culturally significant sites and traditional cultural properties. Mountains, peaks, buttes, prominences, and other elements of the natural and cultural environment are often considered important elements of a traditional culturally significant landscape.

Traditional cultural properties are ones that refer to beliefs, customs, and practices of a living community that have been passed down over the generations. Native American traditional cultural properties are often not found on the state or national registers of historic properties or described in the extant literature or in SHPO files. There are, however, a range of cultural properties types of religious or traditional use that might be identified during the tribal consultation process. These might include:

- Sites of ritual and ceremonial activities and related features
- Shrines
- Marked and unmarked burial grounds
- Traditional use areas
- Plant and mineral gathering areas
- Traditional hunting areas
- Caves and rock shelters
- Springs
- Trails
- Prehistoric archaeological sites

The U.S. Bureau of Indian Affairs web site contains a list, current as of May 2007, of tribal leaders and contact information <<http://www.doi.gov/bia/Tribal%20Leaders-June%202007-2.pdf>>. These tribal groups should be contacted for consultations associated with ISL milling activities in their respective states (see Table 3.2-12). Additional tribal contact information may be obtained from the respective SHPO in Nebraska, Montana, South Dakota, and Wyoming.

Table 3.2-12. List of Tribal Contacts for Tribes With Interests in Nebraska, Montana, South Dakota, and Wyoming

Nebraska
Santee Sioux Nation, 108 Spirit Lake Ave. West, P(402) 857-2772 F(402) 857-2307, Roger Trudell, Chairman, Niobrara, NE 68760-7219
Ponca Tribe of Nebraska, P.O. Box 288, P(402) 857-3391 F(402) 857-3736, Larry Wright, Jr., Chairman, Niobrara, NE 68760
Omaha Tribal Council, P.O. Box 368, P(402) 837-5391 F(402) 837-5308, Mitchell Parker, Chairperson, Macy, NE 68039
Iowa Tribe of Kansas & Nebraska, 3345 Thrasher Rd., P(785) 595-3258 F(785) 595-6610, Leon Campbell, Chairman, White Cloud, KS 66094
Sac and Fox Nation of Missouri, 305 N. Main Street, P(785) 742-7471 F(785) 742-3785, Fredia Perkins, Chairperson, Reserve, KS 66434
Ponca Tribe of Nebraska, P.O. Box 288, P(402) 857-3391 F(402) 857-3736, Larry Wright, Jr., Chairman, Niobrara, NE 68760
Montana
Blackfeet Tribal Business Council, P.O. Box 850, P(406) 338-7276 F(406) 338-7530, Earl Old Person, Chairman, Browning, MT 59417 <btbc@3rivers.net>
Chippewa Cree Business Committee, RR 1, P.O. Box 544, P(406) 395-4282 F(406) 395-4497, John "Chance" Houle, Chairman, Box Elder, MT 59521
Crow Tribal Council, P.O. Box 169, P(406) 638-3715 F(406) 638-3773, Carl Venne, Chairman, Crow Agency, MT 59022
Fort Belknap Community Council, RR 1, Box 66, P(406) 353-2205 F(406) 353-4541, Julia Doney, President, Harlem, MT 59526
Fort Peck Tribal Executive Board, P.O. Box 1027, P(406) 768-5155 F(406) 768-5478, John Morales, Chairman, Poplar, MT 59255
Northern Cheyenne Tribal Council, P.O. Box 128, P(406) 477-6284 F(406) 477-6210, Eugene Littlecoyote, President, Lame Deer, MT 59043
Confederated Salish & Kootenai Tribes, Tribal Council, Box 278, P(406) 675-2700 F(406) 675-2806, James Steele, Jr., Chairman, Pablo, MT 59855 <csktadm@ronan.net>
South Dakota
Cheyenne River Sioux Tribe, P.O. Box 590, P(605) 964-4155 F(605) 964-4151, Joseph Brings Plenty, Chairman, Eagle Butte, SD 57625
Crow Creek Sioux Tribal Council, P.O. Box 50, P(605) 245-2221 F(605) 245-2470, Lester Thompson, Chairman, Fort Thompson, SD 57339
Flandreau Santee Sioux Executive Committee, P.O. Box 283, P(605) 997-3891 F(605) 997-3878, Joshua Weston, President, Flandreau, SD 57028 <president@fsst.org>
Lower Brule Sioux Tribal Council, 187 Oyate Circle, P(605) 473-5561 F(605) 473-5606, Michael Jandreau, Chairman, Lower Brule, SD 57548
Oglala Sioux Tribal Council, P.O. Box 2070, P(605) 867-6074 F(605) 867-6076, John Yellow Bird Steele, President, Pine Ridge, SD 57770
Rosebud Sioux Tribal Council, P.O. Box 430, P(605) 747-2381 F(605) 747-2905, Rodney Bordeaux, President, Rosebud, SD 57570 <www.rosebudsiouxtribe.org>
Sisseton-Wahpeton Oyate of the Lake Traverse Reservation, P.O. Box 509, P(605) 698-3911 F(605) 698-7907, Michael Selvage, Sr., Chairman, Agency Village, SD 57262 <http://swcc.cc.sd.us/>

Table 3.2-12. List of Tribal Contacts for Tribes With Interests in Nebraska, Montana, South Dakota, and Wyoming (continued)***South Dakota (continued)***

Standing Rock Sioux Tribal Council, P.O. Box D, P(701) 854-8500 F(701) 854-7299, Ron His Horse Is Thunder, Chairman, Fort Yates, ND 58538

Yankton Sioux Tribal Business & Claims Committee, P.O. Box 248, P(605) 384-3641 F(605) 384-5687, Robert Cournoyer, Chairman, Marty, SD 57361-0248 <bobbycournoyer@yahoo.com>
<www.yanktonsiouxtribe.org/index.html>

Wyoming

Arapaho Business Committee, P.O. Box 396, P(307) 332-6120 F(307) 332-7543, Richard B. Brannon, Chairman, Fort Washakie, WY 82514

Shoshone Business Committee, P.O. Box 217, P (307) 332-3532 F(307) 332-3055, Ivan D. Posey, Chairman, Fort Washakie, WY 82514

3.2.9 Visual/Scenic Resources

Assigning values to visual and scenic resources is subjective, but basic design elements such as form, line, color, and texture can be used to describe and evaluate landscapes.

Modifications that repeat the landscape's basic elements tend to match the surroundings well.

Modifications that do not match basic landscape features can look out of place and jar the viewer.

Potential visual impacts can be evaluated based on likely features that may result from anticipated activities (drilling masts, well heads, header houses, satellite ion exchange facilities, and centralized milling facilities) from the perspective of both design (space, height, color) and time (permanent versus temporary structures).

Federal land management agencies such as the BLM and the U.S. Forest Service (USFS) have established guidelines to inventory and manage visual resources. Because there are a variety of visual values, different levels of management are necessary. These activities are typically part of a visual resource management (VRM) system.

The BLM guidelines for VRM are identified in BLM Manual 8400 (BLM, 2007a). The VRM system identifies and inventories existing scenic values (BLM, 2007a–c) and establishes management objectives for those values. These area-specific objectives provide the standards for planning, designing, and evaluating the

**Objectives for Visual Resource Classes
(After BLM, 2007a,b)**

Class I: To preserve the existing character of the landscape. This class provides for natural ecological changes; however, it does not preclude very limited management activity. The level of change to the characteristic landscape should be very low and must not attract attention.

Class II: To retain the existing character of the landscape. The level of change to the characteristic landscape should be low. Management activities may be seen, but should not attract the attention of the casual observer. Any changes must repeat the basic elements of form, line, color, and texture found in the predominant natural features of the characteristic landscape.

Class III: To partially retain the existing character of the landscape. The level of change to the characteristic landscape should be moderate. Management activities may attract attention but should not dominate the view of the casual observer. Changes should repeat the basic elements found in the predominant natural features of the characteristic landscape.

Class IV: To provide for management activities that require major modifications of the existing character of the landscape. The level of change to the characteristic landscape can be high. These management activities may dominate the view and be the major focus of viewer attention. However, every attempt should be made to minimize the impact of these activities through careful location, minimal disturbance, and repeating the basic elements.

Description of the Affected Environment

potential visual resource impacts resulting from future management projects. The VRM system also provides for mitigation measures that can reduce potentially adverse visual impacts.

In practice, the VRM system as described by BLM consists of two stages:

- Inventory—Visual Resource Inventory (BLM, 2007b)
- Analysis—Visual Resource Contrast Rating (BLM, 2007c)

Landscape inventories are determined by taking scenic quality, visual sensitivity, and distance from the existing travel routes and dividing these factors into as many as four classes. The final VRM class determinations are typically established in the resource management plans developed by BLM field offices. The USFS system for VRM is slightly different from that used by the BLM, with five classifications based on visual quality and scenic integrity objectives (USFS, 1974, 1995).

Based on the BLM Visual Resource Handbook, the uranium districts in the Wyoming West Uranium Milling Region are located in the Wyoming Basin physiographic province (BLM, 2007a). Although BLM does not manage all of the land in the Wyoming West Uranium Milling Region, the BLM resource management plans prepared by the regional field offices establish VRM classifications for all of the region, including private land or land managed by other agencies. The regional management plans that cover the Wyoming West Uranium Milling Region include the Casper (BLM, 2007d; Bennett, 2003), Lander (BLM, 1987), Rock Springs (BLM, 2007e), and Rawlins (BLM, 2008) field offices (see the BLM Wyoming website at <http://www.blm.gov/wy/st/en.html>). The VRM classifications assigned within these resource plans are presented in Figure 3.2-20. The Lander resource management plan is in the process of being revised; as a result, the current VRM classification for the northern part of the Wyoming West Uranium Milling Region is not available at this time (BLM, 2007f). Public concerns expressed to BLM include visual and scenic resources relating to the quality of recreational experiences on public lands and protecting landscapes along sensitive resources such as the National Historic Trails (BLM, 2007d).

Visual Quality and Scenic Integrity Objectives of the USFS (From USFS, 1974, 1995)

The USFS established visual quality objectives as part of a visual management system in its 1974 forest landscape management handbook. These objectives described the different degrees of alteration associated with a proposed management strategy that the USFS would find acceptable in terms of visual contrast with the surrounding natural landscape. The visual quality objectives have been updated and replaced by scenic integrity objectives as part of the USFS scenery management system (USFS, 1995). There has been some overlap in their application, and both systems have been used by the USFS to define visual resources.

Preservation: This visual quality objective represents essentially unaltered landscape with only minute if any deviations. This is equivalent to an area with very high scenic integrity.

Retention: This visual quality objective represents landscape that appears to be intact to the casual viewer. Alterations may be present, but are consistent with the form, line, color, and texture of the landscape. It is equivalent to a classification of high scenic integrity.

Partial Retention: This visual quality objective represents landscape that appears slightly altered. New form, line, color, or texture may be introduced as long as they remain visually subordinate. This objective is equivalent to a classification of moderate scenic integrity.

Modification: This visual quality objective represents landscape that appears moderately altered. Changes may be introduced that visually dominate the characteristic landscape, but must reflect naturally established form, line, color, and texture to be compatible with natural surroundings. This objective is equivalent to a classification of low scenic integrity.




Figure 3.2-20 BLM Visual Resource Classifications for the Wyoming West Uranium Milling Region (BLM, 2008, 2007d,e)

Description of the Affected Environment

The bulk of the Wyoming West Uranium Milling Region is categorized by BLM as VRM Class III (along highways) and Class IV (open grassland, oil and natural gas, urban areas) (Figure 3.2-20). The BLM resource management plans do not identify any VRM Class I (most sensitive) resources that fall entirely within the Wyoming West Uranium Milling Region. Located in the northwestern corner of Carbon County, however, the Ferris Mountains Wilderness Study Area is identified as Class I (BLM, 2008) and borders the eastern boundary of the region, about 72 km [45 mi] north of Rawlins. The closest potential uranium ISL facility, however, is located about 24 km [15 mi] from the closest boundary of the Ferris Mountains Wilderness Study Area. VRM Class II areas are generally identified in ranges such as the Granite Mountains, and the Rock Springs field office identifies Red Lake, Alkali Basin, Alkali Draw, South Pinnacles, and Honeycomb Buttes Wilderness Study Areas in the southwestern corner of the region as Class II (Figure 3.2-20). These Class II areas, however, are more than 32 km [20 mi] from the closest point in either of the two uranium districts located within the Wyoming West Uranium Milling Region. In addition, scenic areas along the Sweetwater and Powder Rivers provide unique viewsheds (USFS, 2005). One potential facility may be located near Jeffrey City, within a few kilometers [miles] of the Sweetwater. All of the other potential facilities are located 24 km [15 mi] or more from these two rivers. As described in Section 3.2.6.2, there are no areas identified by EPA as Class 1 prevention of significant deterioration areas in the Wyoming West Uranium Milling Region (see Figure 3.2-16). In addition, the state of Wyoming Environmental Quality Council also has developed two designations for scenic resources, Unique and Irreplaceable and Rare or Uncommon. These designations are limited to a small number of locations (seven), and none are located within the two uranium districts in the Wyoming West Uranium Milling Region (Girardin, 2006).

The Wind River Indian Reservation occupies the northwestern corner of the region, including the Boysen and Pilot Reservoirs managed by the U.S. Bureau of Reclamation. These areas fall within the area covered by the BLM Lander field office, and VRM classifications are not available. These regions are more than 16 km [10 mi] northwest from the closest potential ISL facility at Sand Draw, however, and more than 50 km [30 mi] from the center of the two uranium districts at Gas Hills and Crooks Gap.

3.2.10 Socioeconomics

For the purpose of this Draft GEIS, the socioeconomic description for the Wyoming West Region includes communities within the region of influence for a potential ISL facility. Communities that have the highest potential for socioeconomic impacts are considered the affected environment. These potentially affected communities are defined by (1) proximity to an ISL facility {generally within 48 km [30 mi]}, (2) economic profile, such as potential for income growth or destabilization, (3) employment structure, such as potential for job placement or displacement, and (4) community profile, such as potential for growth or destabilization to local emergency services, schools, or public housing. The affected environment are listed in Table 3.2-13.

1

Table 3.2-13. Summary of the Affected Environment Within the Wyoming West Uranium Milling Region		
Counties Within Wyoming West	Towns Within Wyoming West	Native American Communities Within Wyoming West
Carbon	Arapahoe	Wind River Indian Reservation
Fremont	Ethete	
Natrona		
Sweetwater	Ft. Washakie	
	Lander	
	Riverton	
	St. Stephens	

2

3 The following sub-sections, describe areas most likely to have implications to socioeconomics.
 4 In some sub-sections, Core-Based Statistical Areas and Metropolitan Areas are also discussed.
 5 A Core-Based Statistical Area, according to the U.S. Census Bureau, is a collective term for
 6 areas ranging from a population of 10,000 to 50,000. A Metropolitan Area is greater than
 7 50,000 and a town is considered less than 10,000 in population (U.S. Census Bureau, 2008).
 8 A number of small towns with populations less than 1,000 exist in the affected environment but
 9 are not called out by name in Table 3.2-13 or in data presented in this section. Town such as
 10 Moneta, Jeffrey City, Bairoil, Lamont, Wamsutter and others are represented collectively by the
 11 applicable county level socioeconomic information provided in this section.

12

13 3.2.10.1 Demographics

14

15 For the Draft GEIS, demographics are based on 2000 Census data on population and racial
 16 characteristics of the affected environment (Table 3.2-14) and Figure 3.2-21 illustrates the
 17 population of communities within the Wyoming West Uranium Milling Region. Most 2006 data
 18 compiled by the U.S. Census Bureau is not yet available for the region.

19

20 The most populated county in the Wyoming West Uranium Milling Region is Natrona County
 21 and the most sparsely populated county is Carbon County. Riverton has the largest population
 22 in the region and, and the smallest populated town is Ethete (Wind River Indian Reservation).
 23 The county with the largest percentage of non-minorities is Natrona County with a white
 24 population of 94.2 percent, and Lander has a white population of 90.8 percent. The largest
 25 minority-based county is Fremont County with a white population of 76.5 percent. The largest
 26 minority-based town is Ethete, with a white population of only 4.9 percent.

27

28 Although not listed in Table 3.2-14, the total population counts based on 2000 U.S. Census
 29 Bureau of the Wind River Indian Reservation was 23,250. The Wind River Indian Reservation is
 30 shared by the Eastern Shoshone and Northern Arapahoe tribes and is located in Fremont and
 31 Hot Springs Counties, Wyoming. Riverton is the largest town on the reservation (U.S. Census
 32 Bureau, 2008).

33

34

**Table 3.2-14. 2000 U.S. Bureau of Census Population and Race Categories of the
Wyoming West Uranium Milling Region***

Affected Environment	Total Population	White	African American	Native American	Some Other Race	Two or More Races	Asian	Hispanic Origin†	Native Hawaiian and Other Pacific Islander
Wyoming	493,782	454,670	3,722	11,133	12,301	8,883	2,771	31,669	302
<i>Percent of total</i>		92.1%	0.8%	2.3%	2.5%	1.8%	0.6%	6.4%	0.1%
Carbon County	15,639	14,092	105	9	808	321	105	2,163	9
<i>Percent of total</i>		90.1%	0.7%	0.1%	5.2%	2.1%	0.7%	13.8%	0.1%
Fremont County	35,804	27,388	44	7,047	417	793	106	1,566	9
<i>Percent of total</i>		76.5%	0.1%	19.7%	1.2%	2.2%	0.3%	4.4%	0.0%
Natrona County	66,533	62,644	505	686	1,275	1,121	277	3,257	25
<i>Percent of total</i>		94.2%	0.8%	1.0%	1.9%	1.7%	0.4%	4.9%	0.0%
Sweetwater County	37,613	34,461	275	380	1,349	892	240	3,545	16
<i>Percent of total</i>		91.6%	0.7%	1.0%	3.6%	2.4%	0.6%	9.4%	0.0%
Lander	6,867	6,236	10	411	48	140	22	239	0
<i>Percent of total</i>		90.8%	0.1%	6.0%	0.7%	2.0%	0.3%	3.5%	0.0%
Arapahoe (Wind River Indian Reservation)	1,766	318	2	1,423	9	13	0	91	1
<i>Percent of total</i>		18.0%	0.1%	80.6%	0.5%	0.7%	0.0%	5.2%	0.1%
Ethete (Wind River Indian Reservation)	1,455	72	0	1,371	1	10	1	30	0
<i>Percent of total</i>		4.9%	0.0%	94.2%	0.1%	0.7%	0.1%	2.1%	0.0%

**Table 3.2-14. 2000 U.S. Bureau of Census Population and Race Categories of the
Wyoming West* Uranium Milling Region (continueud)**

Affected Environment	Total Population	White	African American	Native American	Some Other Race	Two or More Races	Asian	Hispanic Origin†	Native Hawaiian and Other Pacific Islander
Fort Washakie (Wind River Indian Reservation)	1,477	87	1	1,368	10	11	0	48	0
<i>Percent of total</i>		5.9%	0.1%	92.6%	0.7%	0.7%	0.0%	3.2%	0.0%
Riverton (Wind River Indian Reservation)	9,310	8,082	16	752	173	240	44	660	3
<i>Percent of total</i>		86.8%	0.2%	8.1%	1.9%	2.6%	0.5%	7.1%	0.0%
St. Stephens (Wind River Indian Reservation)	NA	NA	NA	NA	NA	NA	NA	NA	NA
<i>Percent of total</i>		NA	NA	NA	NA	NA	NA	NA	NA

*U.S. Census Bureau. "American FactFinder." <http://factfinder.census.gov/home/saff/main.html?_lang=en> (18 October 2007 and 25 February 2008).
†Hispanic origin can be any race and is calculated as a separate component of the total population (i.e., if added to the other races would total more than 100 percent).
§NA—not available.




Figure 3.2-21. Wyoming West Uranium Milling Region With Population

3.2.10.2 Income

Income information from the 2000 Census including labor force, income, and poverty levels for the affected environment, is based on data collected at the state and county levels. Data collected at the state level also includes information on towns, Core-Based Statistical Areas, or Metropolitan Areas and was done to take into consideration an outside workforce. An outside workforce may be a workforce willing to commute long distances (greater than 30 miles) for income opportunities or may be a workforce necessary to fulfill specialized positions (if local workforce is unavailable or does not have the appropriate skill set). In Wyoming, the workforce frequently commutes long distances to work. For example, in the Wyoming West Uranium Milling Region, all of the affected counties experienced net inflows of workers during the 4th Quarter of 2005. Net inflows ranged from 370 for Carbon County to 10,600 for Natrona County, predominantly for jobs related to the energy industry (Wyoming Workforce Development Council, 2007). Data collected at the county level is generally the same as the affected environment presented in Table 3.2-13, and also includes information on Native American communities. State level information for the surrounding region is provided in Table 3.2-15 for comparison and county data is listed in Table 3.2-16.

For the surrounding region, the state with the largest labor force population is Montana. The population with the largest labor force is Billings, Montana 320 km [200 mi] to the nearest potential ISL facility. The population in the surrounding region with the highest per capita income is Cheyenne, Wyoming 225 km [140 mi] from the nearest potential ISL facility and the lowest per capita income population is Laramie, Wyoming 160 km [100 mi] to the nearest potential ISL facility. The population with the highest percentage of individuals and families below poverty levels is Billings, Montana.

Based on review of Table 3.2-16, the county in the Wyoming West Uranium Milling Region with the largest labor force population is Natrona County and the smallest labor force population is in Carbon County. The town with the largest labor force population in the region is Riverton (Wind River Indian Reservation) and the smallest labor force population is in Ethete (Wind River Indian Reservation). Sweetwater County has the highest per capita income and the smallest per capita income is in Fremont County. Per capita income ranges from Lander (\$18,389) and the town of Ethete (\$7,129). The county with the highest percentage of individuals and families below poverty levels is Fremont County. The town with the highest percentage of individuals and families below poverty levels is Fort Washakie (Wind River Indian Reservation).

3.2.10.3 Housing

Housing information from the 2000 Census is provided in Table 3.2-17. Housing information for the Wind River Indian Reservation was only available for the town of Riverton (U.S. Census Bureau, 2008).

The availability of housing within the immediate vicinity of the potential ISL facilities in the Wyoming West Uranium Milling Region is limited. The majority of housing is available in larger populated areas such as the towns of Riverton (20 miles to nearest ISL facility) and Casper (60 miles to nearest ISL facility). Temporary housing such as apartments, lodging, and trailer camps within the immediate vicinity of the proposed ISL facilities is not as limited. The majority of apartments are available in larger populated areas such as the towns of Lander, Riverton, and Rawlins with a total of 18 apartment complexes (MapQuest, 2008). There are also

Table 3.2-15. U.S. Bureau of Census State Income Information for the Region Surrounding the Wyoming West Uranium Milling Region*

Affected Environment	2000 Labor Force Population (16 years and over)	Median Household Income in 1999	Median Family Income in 1999	Per Capita Income in 1999	Families Below Poverty Level in 2000	Individuals Below Poverty Level in 2000
Montana	458,306	\$33,024	\$40,487	\$17,151	25,004	128,355
Wyoming	257,808	\$37,892	\$45,685	\$19,134	10,585	54,777
Billings, Montana	47,584	\$35,147	\$45,032	\$19,207	2,130	10,402
<i>Percent of total†</i>	67.7%	NA	NA	NA	9.2%	12.0%
Cheyenne, Wyoming	27,647	\$38,856	\$46,771	\$19,809	891	4,541
<i>Percent of total†</i>	66.7%	NA	NA	NA	6.3%	8.8%
Lander, Wyoming	3,337	\$32,397	\$41,958	\$18,389	178	859
<i>Percent of total†</i>	62.5%	NA	NA	NA	9.95%	13.2%
Laramie, Wyoming	15,504	\$27,319	\$43,395	\$16,036	633	5,618
<i>Percent of total†</i>	67.2%	NA	NA	NA	11.1%	22.6%

*U.S. Census Bureau. "American FactFinder." <http://factfinder.census.gov/home/saff/main.html?_lang=en> (18 October 2007, 25 February 2008, and 15 April 2008).

†Percent of total based on a population of 16 years and over.

Table 3.2-16. U.S. Bureau of Census County and Native American Income Information for the Wyoming West Uranium Milling Region*

Affected Environment	2000 Labor Force Population (16 years and over)	Median Household Income in 1999	Median Family Income in 1999	Per Capita Income in 1999	Families Below Poverty Level in 2000	Individuals Below Poverty Level in 2000
Carbon County, Wyoming	7,744	\$36,060	\$41,991	\$18,375	411	1,879
<i>Percent of total†</i>	62.5%	NA	NA	NA	9.8%	12.9%
Fremont County, Wyoming	17,637	\$32,503	\$37,983	\$16,519	1,267	6,155
<i>Percent of total†</i>	64.9%	NA	NA	NA	13.3%	17.6%
Natrona County, Wyoming	35,081	\$36,619	\$45,575	\$18,913	1,548	7,695
<i>Percent of total†</i>	68.3%	NA	NA	NA	8.7%	11.8%
Sweetwater County, Wyoming	20,022	\$46,537	\$54,173	\$19,575	548	2,871
<i>Percent of total†</i>	70.6%	NA‡	NA	NA	5.4%	7.8%
Arapahoe (Wind River Indian Reservation)	636	\$22,679	\$24,659	\$8,943	134	784
<i>Percent of total†</i>	58.1%	NA	NA	NA	35.5%	45.0%
Ethete (Wind River Indian Reservation)	517	\$24,130	\$24,762	\$7,129	95	453
<i>Percent of total†</i>	60.5%	NA	NA	NA	33.9%	34.4%

Table 3.2-16. U.S. Bureau of Census County and Native American Income Information for the Wyoming West Uranium Milling Region* (continued)

Affected Environment	2000 Labor Force Population (16 years and over)	Median Household Income in 1999	Median Family Income in 1999	Per Capita Income in 1999	Families Below Poverty Level in 2000	Individuals Below Poverty Level in 2000
Fort Washakie (Wind River Indian Reservation)	567	\$18,906	\$20,658	\$7,700	151	636
<i>Percent of total†</i>	57.6%	NA	NA	NA	42.9%	42.7%
St. Stephens (Wind River Indian Reservation)	na	na	na	na	na	na
<i>Percent of total†</i>	na	NA	NA	NA	na	na
Riverton (Wind River Indian Reservation)	4,694	\$31,531	\$37,079	\$16,720	267	1,400
<i>Percent of total†</i>	64.5%	NA	NA	NA	11.0%	15.7%
* U.S. Census Bureau. "American FactFinder." < http://factfinder.census.gov/home/saff/main.html?_lang=en > (18 October 2007 and 25 February 2008). †Percent of total based on a population of 16 years and over. ‡NA—Not applicable. §na—not available.						

Table 3.2-17. U.S. Bureau of Census Housing Information for Wyoming*

Affected Environment	Single Family Owner-Occupied Homes	Median Value in Dollars	Median Monthly Costs With a Mortgage	Median Monthly Costs Without a Mortgage	Occupied Housing Units	Renter-Occupied Units
Wyoming	95,591	\$96,600	\$825	\$229	193,608	55,793
Carbon County	7,744	\$76,500	\$685	\$196	6,129	1,708
Fremont County	6,281	\$89,300	\$714	\$217	13,545	3,496
Natrona County	15,250	\$84,600	\$746	\$218	26,819	7,993
Sweetwater County	7,283	\$104,200	\$953	\$231	14,105	3,488
Lander	1,479	\$97,300	\$701	\$226	2,777	833
Riverton (Wind River Indian Reservation)	2,146	\$83,200	\$683	\$203	3,792	1,221

Source: U.S. Census Bureau. "American FactFinder." 2000.

<http://factfinder.census.gov/home/saff/main.html?_lang=en> (18 October 2007 and 25 February 2008).

5 hotels/motels along major highways or towns near potential ISL facilities in the two uranium districts in the Wyoming West Uranium Milling Regions. In addition to apartments and lodging, there are trailer camps situated near potential ISL facilities (along major roads or near towns) in this region (MapQuest, 2008)

3.2.10.4 Employment Structure

Employment structure from the 2000 Census including employment rate and type is based on data collected at the state and county level. Data collected at the state level also includes information on towns, Core-Based Statistical Areas, or Metropolitan Areas and was done to take into consideration an outside workforce. An outside workforce includes workers willing to commute long distances {more than 48 km [30 mi]} for employment opportunities or external labor necessary to fulfill specialized positions (if the local workforce is unavailable or does not have the necessary skill sets). Data collected at the county level is the same as the affected environment presented in Table 3.2-13, and also includes information on Native American communities.

Based on review of state level information, Wyoming has a low unemployment rate (3.5 percent).

Unemployment at the county level ranges from 3.3 percent (Carbon County) to 5.7 percent (Fremont County). The town with the highest percentage of employment is Lander and the town with the highest unemployment rate is Arapaho on the Wind River Indian Reservation.

Description of the Affected Environment

3.2.10.4.1 State Data

3.2.10.4.1.1 Montana

The State of Montana has an employment rate of 60.8 percent and unemployment rate of 4.1 percent. The largest sector of employment is management, professional, and related occupations at 33.1 percent. The largest type of industry is educational, health, and social services. The largest class of worker is private wage and salary workers (U.S. Census Bureau, 2008).

Billings

Billings has an employment rate of 64.8 percent and unemployment rate of 2.8 percent. The largest sector of employment is sales and office occupations at 31.9 percent. The largest type of industry is educational, health, and social services. The largest class of worker is private wage and salary workers (U.S. Census Bureau, 2008).

3.2.10.4.1.2 Wyoming

The State of Wyoming has an employment rate of 63.1 percent and unemployment rate of 3.5 percent. The largest sector of employment is sales and office occupations. The largest type of industry is educational, health, and social services. The largest class of worker is private wage and salary workers (U.S. Census Bureau, 2008).

Cheyenne

Cheyenne has an employment rate of 59.2 percent and unemployment less than the state at 3.3 percent. The largest sector of employment is management, professional, and related occupations at 33.0 percent. The largest type of industry is educational, health, and social services. The largest class of worker is private wage and salary workers (U.S. Census Bureau, 2008).

Lander

Lander has an employment rate of 59.4 percent and an unemployment rate lower than that of the state at 2.8 percent. The largest sector of employment is management, professional, and related occupations at 39.3 percent. The largest type of industry is educational, health, and social services at 37.9 percent. The largest class of worker is private wage and salary workers at 62.6 percent (U.S. Census Bureau, 2008).

Laramie

Laramie has an employment rate of 63.4 percent and unemployment less than the state at 3.7 percent. The largest sector of employment is management, professional, and related occupations at 40.5 percent. The largest type of industry is educational, health, and social services. The largest class of worker is private wage and salary workers (U.S. Census Bureau, 2008).

3.2.10.4.2 County Data

Carbon County, Wyoming

Carbon County has an employment rate of 59.2 percent and an unemployment rate lower than that of the state at 3.3 percent. The largest sector of employment is management, professional, and related occupations at 23.4 percent followed by sales and office occupations at 21.9 percent. The largest type of industry is educational, health, and social services at 17.1 percent. The largest class of worker is private wage and salary workers at 65.6 percent (U.S. Census Bureau, 2008).

Fremont County, Wyoming

Fremont County has an employment rate of 59.0 percent and an unemployment rate relatively high at 5.7 percent when compared to the state average. The largest sector of employment is management, professional, and related occupations at 33.9 percent followed by sales and office occupations at 22.5 percent. The largest type of industry is educational, health, and social services at 28.5 percent. The largest class of worker is private wage and salary workers at 64.1 percent (U.S. Census Bureau, 2008).

Natrona County, Wyoming

Natrona County has an employment rate of 64.6 percent and an unemployment rate similar to that of the state at 3.5 percent. The largest sector of employment is sales and office occupations at 29.9 percent followed by management, professional, and related occupations at 28.5 percent. The largest type of industry is educational, health, and social services at 21.2 percent. The largest class of worker is private wage and salary workers at 76.2 percent (U.S. Census Bureau, 2008).

Sweetwater County, Wyoming

Sweetwater County has an employment rate of 66.4 percent and an unemployment rate slightly higher than that of the state at 4.0 percent. The largest sector of employment is sales and office occupations at 23.4 percent followed by management, professional, and related occupations at 23.3 percent. The largest type of industry is educational, health, and social services at 18.2 percent. The largest class of worker is private wage and salary workers at 76.5 percent (U.S. Census Bureau, 2008).

Native American Communities

Information on labor force and poverty levels for the Wind River Indian Reservation is based on 2003 Bureau of Indian Affairs data and is provided in Table 3.2-18. The Northern Arapaho Tribe reports unemployment rates much higher than the statewide levels (U.S. Department of the Interior, 2003).

Description of the Affected Environment

Table 3.2-18. Employment Structure of the Wind River Indian Reservation Within the Affected Area*

Affected Environment	2003 Labor Force Population	Unemployed as Percent of Labor Force	Employed Below Poverty Guidelines	
Arapaho Tribe of the Wind River Indian Reservation	1,386	72%	106	8%

* U.S. Department of the Interior. "Affairs American Indian Population and Labor Force Report 2003." <<http://www.doi.gov/bia/labor.html>>. Washington, DC: U.S. Department of the Interior, Bureau of Indian Affairs, Office of Tribal Affairs. 2003.

3.2.10.5 Local Finance

Local finance such as revenue and tax information for the affected environment is provided below and in Table 3.2-19.

Table 3.2-19. 2007 Sales and Use Tax Distribution of the Affected Counties Within Wyoming West Uranium Milling Region*

Affected Counties	Use Tax		Sales Tax		Lodging Option Tax
	General	Specific	General	Specific	
Carbon County	\$8,546.95	\$64,236.31	\$465,469.37	\$47,391.45	\$40,974.56
Fremont County	\$0.0	\$116,086.27	\$0.0	\$580,209.10	\$40,792.32
Natrona County	\$132,453.29	\$0.0	\$1,572,768.04	\$0.0	\$98,624.31
Sweetwater County	\$124,140.09	\$250,559.08	\$1,459,877.63	\$1,327,426.97	\$73,276.64

* Wyoming Department of Revenue. "Sales and Tax Distribution Report by County 2007." <<http://revenue.state.wy.us/PortalVBVS/DesktopDefault.aspx?tabindex=3&tabid=10>> (18 October 2007 and 25 February 2008).

Wyoming

The State of Wyoming does not have an income tax nor does it assess tax on retirement income received from another state. Wyoming has a 4 percent state sales tax, 2 percent to 5 percent county lodging tax, and 5 percent use tax. Counties have the option of collecting an additional 1 percent tax for general revenue and 2 percent tax for specific purposes. Wyoming also imposes "ad valorem taxes" on mineral extraction properties. Taxes levied for uranium production were 4.0 percent in 2007 and totaled \$17 million dollars (Wyoming Department of Revenue, 2007). The majority of tax revenue came from Converse County with a small amount (\$7,159) from Sweetwater County (Wyoming Department of Revenue, 2007). Sales and use tax distribution information for the affected counties is presented in Table 3.2-19.

Native American Communities

The Wind River Indian Reservation's largest sources of revenue come from the Northern Arapaho and Eastern Shoshone Tribal Governments; the Bureau of Indian Affairs; the Ethete, Fort Washakie, and Arapahoe School Districts; the Indian Health Service; and Native American household income (University of Wyoming, 1997).

3.2.10.6 Education

Based on review of the affected environment, the county with the largest number of schools is Natrona County and the county with the smallest number of schools is Carbon County. The town with the largest number of schools is Lander and the towns with the smallest number of schools (Ethete, Arapaho) are located on the Wind River Indian Reservation.

Lander

Lander has one school district, Fremont County School District No. 1, with a total 2007 enrollment of approximately 1,930 students. There are 5 elementary schools, 4 middle schools, 3 high schools, 7 public schools, and 1 private school. The majority of schools provide bus services (Greatschools.com, 2008).

Carbon County

Carbon County has two school districts, Carbon County School District #1 and #2, with a combined total 2007 enrollment of approximately 2,650 students. There are a total of 9 elementary schools, 2 middle school, 2 high school, and 2 private schools. The majority of schools within each school district provide bus services (Carbon County School District No.1 and No. 2, 2008a,b).

Fremont County

Fremont County has over eight school districts, with a combined total 2007 enrollment of approximately 7,125 students. There are more than 25 public and private elementary, middle, and high schools. The majority of school districts provide bus services (Schoolbug.org 2007).

Natrona County

Natrona County has one school district: Natrona County School District No. 1, with a total enrollment of approximately 11,500 students in 2007. There are more than 30 public and private elementary and secondary schools. The majority of schools provide bus services (Natrona County School District No. 1, 2007).

Sweetwater County

Sweetwater County has 2 school districts with a total of 10 elementary schools, 3 intermediate/middle schools, 4 high schools, and 4 private or parochial schools. There are a total of about 7,175 students. The majority of schools within each district provide bus services (Sweetwater County School District No.1, 2007; Sweetwater County School District No. 2, 2005).

Native American Communities

The Wind River Indian Reservation has several school districts in the towns of Arapaho, Ethete, Fort Washakie, and Saint Stephens. There are a total of approximately 1,060 students. Schools are the Arapaho School, Wyoming Indian School, Fort Washakie School, and Saint Stephens Indian School. All four schools accommodate elementary through 12th grades. Information is not available if bus services are provided by any of these schools (Easternshoshone.net, 2008).

3.2.10.7 Health and Social Services

Health Care

The majority of the health care facilities that provide service in the vicinity of the Wyoming West Uranium Milling Region are located within the larger population centers. The closest health care facilities within the vicinity of the potential ISL facilities are located in Riverton, Lander, Casper, Cheyenne, Laramie, and Thermopolis with a total of 14 facilities (MapQuest, 2008). These consist of hospitals, clinics, emergency centers, and medical services. Hospitals located within the vicinity of the potential ISL facilities include Lander (1), Riverton (1), Rock Springs (1), Rawlind (1), Caspter (2), Laramie (1), and Thermopolis (1).

Local Emergency

Local police in the Wyoming West Uranium Milling Region is under the jurisdiction of each county. There are 16 police, sheriff, or marshals offices within the region: Carbon County (6), Fremont County (3), Natrona County (4), and Sweetwater County (3) (USACops, 2008a).

Fire departments within the Wyoming West Uranium Milling Region are comprised at the County, town, Core-Based Statistical Areas, or city level. There are 7 fire departments within the milling region: Lander (1), Natrona County (1), Dubois (1), Rawlins (2), Fort Washakie (1), and Riverton (1) (50States, 2008a).

3.2.11 Public and Occupational Health

3.2.11.1 Background Radiological Conditions

For a U.S. resident, the average total effective dose equivalent from natural background radiation sources is approximately 3 mSv/yr [300 mrem/yr] but varies by location and elevation (National Council of Radiation Protection and Measurements 1987). In addition, the average American receives 0.6 mSv/yr [60 mrem/yr] from man-made sources including medical diagnostic tests and consumer products (National Council of Radiation Protection and Measurements, 1987). Therefore the total from natural background and man-made sources for the average U.S. resident is 3.6 mSv/yr [360 mrem/yr]. For a breakdown of the sources of this radiation, see Figure 3.2-22.

How is Radiation Measured?

Radiation dose is measured in units of either sievert or rem and often referred to in either milliSv/mSv or millirem/mrem where 1,000 mSv=1 Sv and 1,000 mrem=1 rem. The conversion for sieverts to rem is Sv=100 rem. These units are used in radiation protection to measure the amount of damage to human tissue from a dose of ionizing radiation. Total effective dose equivalent, or TEDE, refers to the sum of the deep-dose equivalent (for external exposures) and the committed effective dose equivalent (for internal exposures). See Table 3.2-20 for public radiation doses from common activities.

Background dose varies by location primarily because of elevation changes and variations in the dose from radon. As elevation increases so does the dose from cosmic radiation and hence the total dose. Radon is a radioactive gas produced from the decay of U-238, which is naturally found in soil. The amount of radon in the soil/bedrock depends on the type the porosity and moisture content. Areas which have types of soils/bedrock like granite have higher radon levels than those with other types of soils/bedrock (EPA, 2006). For the Wyoming West Uranium Milling Region, the average background radiation dose for the state of Wyoming is used, which is 3.16 mSv/yr [316 mrem/yr] (EPA, 2006). This value includes natural and manmade sources. This dose is slightly lower than the U.S. average primarily because the radon dose is lower {U.S. average of 2 mSv/yr [200 mrem/yr] versus Wyoming average of 1.33 mSv/yr [133 mrem/yr]}. Because of the higher elevation, the dose from cosmic radiation is slightly higher than the U.S. average: 0.515 mSv/yr [51.5 mrem/yr] versus 0.27 mSv/yr [27 mrem/yr]. The remaining contributions from terrestrial, internal, and man-made radiation combined are the same as the U.S. average of 1.318 mSv/yr [131.8 mrem/yr].

Table 3.2-20. Public Radiation Doses*

Activity or Event	Dose
Flying from NY to LA	2.5 mrem/trip
Chest x-ray	10 mrem/exam
Full mouth dental x-ray	9 mrem/exam
U.S. average background	360 mrem/yr
* Voss, J.T. "Los Alamos Radiation Monitoring Notebook." LA-UR-00-2584. Los Alamos, New Mexico: Los Alamos National Laboratory. 2000.	

Outdoor radon concentrations are generally a small fraction of the average indoor concentrations. Outdoor radon concentrations can also be influenced by prior mining of any mineral (e.g., uranium, copper) in the area. To develop an open-pit or underground mine, soil and rock need to be excavated to reach the ore. This excavated rock, or overburden, can naturally contain higher levels of uranium and thorium than was present on the surface. Additionally, low grade ore may be left in the area around the mine, especially in the case of abandoned mines. Also, ore processed to extract elements other than uranium and thorium (such as copper, titanium, ruthenium, and other rare earth elements) could result in concentrating the natural uranium or thorium that was in the ore. The process of removing the rock or processing these ores could also change the physical and chemical characteristics controlling radon release, thus allowing additional radon to be released. The overburden and any ore left around the mine could elevate the local outdoor radon concentrations above the levels seen in other parts of the region. In close proximity to the mines, the level of terrestrial radiation could be elevated by the presence of mine waste. The overburden, low grade ore, and tailings from ore processed for other than uranium or thorium is called "technologically enhanced naturally occurring radioactive material" (TENORM). TENORM is not regulated by NRC. Radiation from these sources is considered part of background for compliance with NRC regulations.

3.2.11.2 Public Health and Safety

NRC has the statutory responsibility, under the Atomic Energy Act of 1954, as amended, to protect the public health and safety and the environment. NRC's regulations in 10 CFR Part 20 specify annual dose limits to members of the public of 1 mSv [100 mrem] total effective dose equivalent and 0.02 mSv/hr [2 mrem/hr] from any external sources.

3.2.11.3 Occupational Health and Safety

Occupational health and safety risks to workers include exposure to radioactive materials. Radiation safety practices for workers at uranium ISL facilities should be such that the dose to the workers is kept as low as is reasonably achievable. Radiation exposure limits are specified in 10 CFR Part 20. Occupational dose is determined by the more limiting of (1) 0.05 Sv [5 rem] total effective dose equivalent or (2) sum of the deep-dose equivalent and the committed dose equivalent to any individual organ or tissue other than the lens of the eye being equal to 0.5 Sv [50 rem]. The lens of the eye is limited to a dose equivalent of 0.15 Sv [15 rem] and the skin (of the whole body or any extremity) is limited to a shallow dose equivalent of 0.5 Sv [50 rem]. The monitoring requirements for occupational dose are covered in greater detail in Section 2.9 and Chapter 8.




Figure 3.2-22 Average Annual Background Radiation in the United States {Units of mrem [1 mSv=100 mrem]} (NRC, 2006)

3.2.12 References

- 50States. "Wyoming." <www.50states.com.> (15 April 2008).
- AATA International Inc. (2005). Environmental and social due diligence report Great Divide Basin ISL Uranium Project. Lost Soldier and Lost Creek Claim Areas, Wyoming. Fort Collins, Colorado.
- Anderson, D.C. "Uranium Deposits of the Gas Hills." Contributions to Geology, Wyoming Uranium Issue. Laramie, Wyoming: University of Wyoming. Vol. 8, No. 2.1. pp. 93–104. 1969.

Atomic Safety and Licensing Board. "Atomic Safety and Licensing Board Transcript in the Matter of Hydro Resources Inc." Docket No. 40-8968-ML, ASLBP No. 95-706-01-ML Washington, DC: Atomic Safety and Licensing Board. August 21, 2006.

Bailey, R.V. "Uranium Deposits in the Great Divide Basin-Crooks Gap Area, Fremont and Sweetwater Counties, Wyoming." Contributions to Geology, Wyoming Uranium Issue. Laramie, Wyoming: University of Wyoming. Vol. 8, No. 2.1. pp. 105-120. 1969.

Bauer, E.R. and J.L. Kohler. "Cross-Sectional Survey of Noise Exposure in the Mining Industry." G. Bockosh, M. Karmis, J. Langton, M.K. McCarter, and B. Rowe, eds. Proceedings of the 31st Annual Institute of Mining Health, Safety and Research, Roanoke, Virginia, August 27-30, 2000. Roanoke, Virginia: Institute of Mining Health, Safety, and Research. 2000.

Bennett, E. "The Visual Resource Inventory for the Casper Field Office." Casper, Wyoming: BLM, Casper Field Office. March 2003.

BLM. "Proposed Resource Management Plan and Final Environmental Impact Statement for Public Lands Administered by the Bureau of Land Management Rawlins Field Office." Rawlins, Wyoming: BLM, Rawlins Field Office. 2008. <<http://www.blm.gov/rmp/wy/rawlins/>> (3 March 2008).

BLM. "Visual Resource Management." Manual 8400. Washington, DC: BLM. 2007a. <<http://www.blm.gov/nstc/VRM/8400.html#Anchor-.06-23240>> (17 October 2007).

BLM. "Visual Resource Inventory." Manual H-8410-1. Washington, DC: BLM. 2007b. <<http://www.blm.gov/nstc/VRM/8410.html>> (17 October 2007).

BLM. "Visual Resource Contrast Rating." Manual 8431. Washington, DC: BLM. 2007c. <<http://www.blm.gov/nstc/VRM/8431.html>> (17 October 2007).

BLM. "Proposed Resource Management Plan and Final Environmental Impact Statement for the Casper Field Office Planning Area." BLM/WY/PL-07/017+1610. Casper, Wyoming: BLM, Casper Field Office. June 2007d. <<http://www.blm.gov/rmp/casper/PRMP-FEIS.htm>> (17 October 2007).

BLM. "Green River Resource Management Plan." Rock Springs, Wyoming: BLM. Rock Springs Field Office. 2007e. <<http://www.blm.gov/rmp/WY/application/index.cfm?rmpid=87>> (17 October 2007).

BLM. "Notice of Intent to Prepare a Resource Management Plan Revision and Associated Environmental Impact Statement, for the Lander Field Office, Wyoming." Lander, Wyoming: BLM, Lander Field Office. 2007f. *Federal Register* 72. pp. 6741-6742.

BLM. "Lander Resource Management Plan." Lander, Wyoming: BLM, Lander Field Office. June 1987. <<http://www.blm.gov/rmp/WY/Lander/rmp.pdf>> (17 October 2007).

Boberg, W.W. Some Speculations on the Development of Central Wyoming as a Uranium Province. Wyoming Geological Association 32nd Annual Field Conference Guidebook. Casper, Wyoming: Wyoming Geological Association. pp. 161-180. 1981.

Description of the Affected Environment

- 1 Brattstrom, B.H. and M.C. Bondello. "Effects of Off-Road Vehicle Noise on Desert Vertebrates."
2 *Environmental Effects of Off-Road Vehicles, Impacts and Management in Arid Regions*.
3 R.N. Webb and H.G. Wilshire, eds. New York City, New York: Springer-Verlag Publishing.
4 1983.
- 5
6 Bureau of Indian Affairs. "American Indian Population and Labor Force Report."
7 Washington, DC: U.S. Department of the Interior, Bureau of Indian Affairs. 2003.
- 8
9 Carbon County School District No.1. "Carbon County School District 1." 2008
10 <www.crb2.k11.wy.us/#> (27 February 2008).
- 11
12 Carbon County School District No. 2. "Carbon County School District 2." 2008.
13 <www.crb2.k12.wy.us/#>. (27 February 2008).
- 14
15 Center for Plant Conservation. "National Protection Plant Profile, Recovering Americas
16 Vanishing Flora." 2008 <[http://www.centerforplantconservation.org/
17 ASP/CPC_ViewProfile.asp?CPCNum=3241](http://www.centerforplantconservation.org/ASP/CPC_ViewProfile.asp?CPCNum=3241)> (15 February 2008).
- 18
19 Chapman, S.S., S.A. Bryce, J.M. Omernik, D.G. Despain, J. ZumBerge, and M. Conrad.
20 "Ecoregions of Wyoming." U.S. Geological Survey Map. Scale 1:1,400,000. 2004.
- 21
22 Chenoweth, W.L. "A Summary of Uranium Production in Wyoming." Mineral Resources of
23 Wyoming, Wyoming Geological Association, 42nd Annual Field Conference Guidebook.
24 Casper, Wyoming: Wyoming Geological Association. pp. 169–179. 1991.
- 25
26 Collentine, M., Libre, R., and K.R. Feathers (1981) . Occurrence and Characteristics of Ground
27 Water in the Great Divide and Washakie Basins, Wyoming. Wyoming Water Resources
28 Research Institute report to U.S. Environmental Protection Agency, vol. VI-A.
- 29
30 Colorado Division of Wildlife. "National Diversity Information Source." 2008.
31 <<http://ndis.nrel.colostate.edu/wildlifespx.asp?SpCode=010595>> (12 February, 2008).
- 32
33 Curtis J. and Grimes K. (2004). Wyoming Climate Atlas
34 <<http://www.wrds.uwyo.edu/wrds/wsc/climateatlas/>> (29 April 2008).
- 35
36 Davis, J.F. "Uranium Deposits of the Powder River Basin." Contributions to Geology, Wyoming
37 Uranium Issue. Laramie, Wyoming: University of Wyoming. Vol. 8, No. 2.1. pp. 131–142.
38 1969.
- 39
40 DOE. DOE/EA–1535, "Uranium Leasing Program Final Programmatic Environmental
41 Assessment." Washington, DC: DOE, Office of Legacy Management. 2007.
42 <http://www.eh.doe.gov/nepa/ea/EA1535/ulm_ea2007.pdf> (12 October 2007).
- 43
44 Driscoll, F.G. (1986). Groundwater and Wells. Second edition. Johnson Filtration Systems Inc.,
45 St. Paul, Minnesota, pp. 1089.
- 46
47 Easternshoshone.net. "Eastern Shoshone Tribe Local Area Schools and Colleges of
48 Wyoming." <<http://www.easternshoshone.net/EasternShoshoneLocalSchools2.htm>>
49 (02 April 2008).
- 50

- 1 EPA. "National Assessment Database." 2008. <<http://www.epa.gov/waters/305b/index.html>>
2 (28 February 2008).
- 3
- 4 EPA. "Counties Designate Nonattainment or Maintenance for Clean Air Act's National Ambient
5 Air Quality Standards (NAAQS)." 2007a. <[http://www.epa.gov/oar/](http://www.epa.gov/oar/oaqps/greenbk/mapnmpoll.html)
6 [oaqps/greenbk/mapnmpoll.html](http://www.epa.gov/oar/oaqps/greenbk/mapnmpoll.html)> (29 September 2007).
- 7
- 8 EPA. "Prevention of Significant Deterioration (PSD) Permit Program Status: May 2007."
9 2007b. <<http://www.epa.gov/nsr/where.html>> (26 September 2007).
- 10
- 11 EPA. "Assessment of Variations in Radiation Exposure in the United States (Revision 1)."
12 Contract Number EP-D-05-02. Washington, DC: EPA. 2006.
- 13
- 14 EPA. "Regional Haze Regulations, Final Rule." Title 40—Protection of Environment,
15 Chapter 51, Requirements for Preparation, Adoption, and Submittal of Implementation Plans.
16 40 CFR Part 51. *Federal Register*. Vol. 64, No. 126. pp. 35714–35774. July 1, 1999.
- 17
- 18 EPA. "Noise and Its Measurement." OPA 22/1. Washington, DC: EPA. January, 1981.
- 19
- 20 Federal Highway Administration. "Synthesis of Noise Effects on Wildlife Populations."
21 FHWA-HEP-06-016. Washington, DC: Federal Highway Administration, Department of
22 Transportation. 2004.
- 23
- 24 Frison, G.C. *Prehistoric Hunters of the High Plains*. 2nd Edition. San Diego, California:
25 Academic Press. 1991.
- 26
- 27 Girardin, J. "A List of Areas Designated Unique and Irreplaceable or Designated Rare or
28 Uncommon by the Council." Letter From J. Girardin to T. Lorenzon. Cheyenne, Wyoming:
29 Council on Environmental Quality. November 29, 2006.
- 30
- 31 Greatschools.com. "Lander, Wyoming." 2008. <www.greatschools.com> (27 February 2008).
- 32
- 33 Harris, R.E. and J.K. King. "Geological Classification and Origin of Radioactive Mineralization in
34 Wyoming." A.W. Snoke, J.R. Steidtmann, and S.M. Roberts, eds. *Geology of Wyoming*:
35 Geological Survey of Wyoming Memoir No. 5. pp. 898–916. 1993.
- 36
- 37 Harshman, E.N. "Uranium Deposits of Wyoming and South Dakota." *Ore Deposits in the*
38 *United States 1933–1967*. New York City, New York: American Institute of Mining,
39 Metallurgical, and Petroleum Engineers. pp. 815–831. 1968.
- 40
- 41 Houston, R.S. "Aspects of the Geologic History of Wyoming Related to the Formation of
42 Uranium Deposits." *Contributions to Geology, Wyoming Uranium Issue*. Laramie, Wyoming:
43 University of Wyoming. Vol. 8, No. 21. pp. 67–79. 1969.
- 44
- 45 King, T. *Places That Count: Traditional Cultural Properties in Cultural Resources Management*.
46 Walnut Creek, California: Altamira Press. 2003.
- 47
- 48 Lageson, D. and D. Spearing. *Roadside Geology of Wyoming*. Missoula, Montana: Mountain
49 Press Publishing Company. 1988.
- 50

Description of the Affected Environment

- 1 Langden, R.E. "Geology and Geochemistry of the Highland Uranium Deposit." Wyoming
2 Geological Association Earth Science Bulletin. pp. 41–48. 1973.
3
- 4 Lost Creek ISR, LLC. Lost Creek Project, South-Central Wyoming. Environmental Report.
5 Docket No. 40-9068.
6
- 7 Love, J.D. "Preliminary Report on Uranium Deposits in the Pumpkin Buttes Area, Powder River
8 Basin, Wyoming." U.S. Geological Survey Circular 176. 1952.
9
- 10 MapQuest. "Wyoming." <www.mapquest.com> (15 April 2008).
11
- 12 Montana Natural Heritage Program. "Animal Field Guide." 2008.
13 <http://fwp.mt.gov/fieldguide/detail_ABNRB02020.aspx> (12 February 2008).
14
- 15 Munn, L.C. and C.S. Arneson. "Soils of Wyoming—A Digital Statewide Map at 1:500,000-
16 Scale." B-1069. Laramie, Wyoming: University of Wyoming Agricultural Experiment Station,
17 College of Agriculture. 1998.
18
- 19 National Climatic Data Center. "NCDC U.S. Storm Events Database." 2007.
20 <<http://www4.ncdc.noaa.gov/cgi-win/wwcgl.dll?wwEvent~Storms>> (14 April 2008).
21
- 22 National Climatic Data Center. "Climates of the States, Climatology of the United States No. 60
23 (New Mexico, Nebraska, South Dakota, and Wyoming)." Asheville, North Carolina: National
24 Oceanic and Atmospheric Administration. 2005. <[http://cdo.ncdc.noaa.gov/cgi-](http://cdo.ncdc.noaa.gov/cgi-bin/climatenormals/climatenormals.pl?directive=prod_select2&prodtype=CLIM60&subrnum=>)
25 [bin/climatenormals/climatenormals.pl?directive=prod_select2&prodtype=CLIM60&subrnum=>](http://cdo.ncdc.noaa.gov/cgi-bin/climatenormals/climatenormals.pl?directive=prod_select2&prodtype=CLIM60&subrnum=>)
26 (30 January 2005).
27
- 28 National Climatic Data Center. "Climatography of the United States No. 20: Monthly Station
29 Climate Summaries, 1971–2000." Asheville, North Carolina: National Oceanic and
30 Atmospheric Administration. 2004.
31
- 32 National Council on Radiation Protection and Measurements. "Report No. 094—Exposure of
33 the Population in the United States and Canada From Natural Background Radiation."
34 Bethesda, Maryland: National Council on Radiation Protection & Measurements. 1987.
35
- 36 National Park Service. "Cultural Resource Management Guidelines." NPS-28. Washington,
37 DC: National Park Service. 1991.
38
- 39 National Weather Service. "NOAA Technical Report NWS 33: Evaporation Atlas for the
40 Contiguous 48 United States." Washington, DC: National Oceanic and Atmospheric
41 Administration. 1982.
42
- 43 Natrona County School District No. 1. "Natrona County School District." 2007.
44 <www.natronaschools.org> (15 October 2007).
45
- 46 NRC. NUREG/BR-0322, "Radiation Protection and the NRC." Washington, DC: NRC.
47 February 2006.
48
- 49 NRC. "Environmental Assessment for the Operation of the Gas Hills Project Satellite In Situ
50 Leach Uranium Recovery Facility." Docket No. 40-8857. 2004.
51

- 1 NRC. Regulatory Guide 8.31, "Information Relevant to Ensuring That Occupational Radiation
2 Exposures at Uranium Recovery Facilities Will Be as Low as Is Reasonably Achievable."
3 Washington, DC: NRC. May 2002.
4
- 5 NRC. NUREG-1508, "Final Environmental Impact Statement to Construct and Operate the
6 Crown Point Uranium Solution Mining Project, Crown Point, New Mexico." Washington, DC:
7 NRC. February 1997.
8
- 9 NRC. "Ferret Exploration Company of Nebraska, Inc. Crow Butte Project Semiannual ALARA
10 Report." License No. SUA-1441, Docket No. 40-8829. Washington, DC: NRC. 1989.
11
- 12 Parker, P. and T. King. "Guidelines for Evaluating and Documenting Traditional Cultural
13 Properties." National Register Bulletin 38. Washington, DC: National Park Service. 1998.
14
- 15 Platte River Endangered Partnership. "Platte River Endangered Partnership." 2008.
16 <<http://www.platteriver.org/>> (15 February 2008).
17
- 18 Rackley, R.I. "Environment of Wyoming Tertiary Uranium Deposits." American Association
19 Petroleum Geologists Bulletin. Vol. 56, No. 4. 1972.
20
- 21 Reher, C.A. "Chapter 6: Ethnology and Ethnohistory." Archaeology of the Eastern Powder
22 River Basin Wyoming. G.M. Zeimens and D. Walker, eds. Laramie, Wyoming: University of
23 Wyoming, Office of the Wyoming State Archaeologist, Department of Anthropology. 1977.
24
- 25 Schoolbug.org. "Fremont County, Wyoming Public Schools." 2007. <www.schoolbug.org>
26 (15 October 2007).
27
- 28 Stephens, J.G. "Geology and Uranium Deposits at Crooks Gap, Fremont County, Wyoming."
29 U.S. Geological Survey Bulletin 1147F. pp. F1-F82. 1964.
30
- 31 Sweetwater County School District No. 1. 2007. "Sweetwater County School District One."
32 <www.sweetwater1.org> (15 October 2007).
33
- 34 Sweetwater County School District No. 2. "Sweetwater County School District Two." 2005.
35 <www.sw2.k12.wy.us> (15 October 2007).
36
- 37 Texas Parks and Wildlife Department. "Hunting and Wildlife." Austin, Texas: Texas Parks and
38 Wildlife Department. 2007. <www.tpwd.state.us/huntwild> (15 October 2007).
39
- 40 Trentham, R.C. and I.P. Orajaka. "Leaching of Uranium From Felsic Volcanic Rocks:
41 Experimental Studies." *Uranium*. Vol. 3. pp. 55-67. 1986.
42
- 43 University of Wyoming. "Wyoming Natural Diversity Database." Laramie, Wyoming: University
44 of Wyoming. 2008. <<http://uwadmnweb.uwyo.edu/wyndd/>> (15 February 2008).
45
- 46 University of Wyoming. "Economic Impact of the Wind River Reservation on Fremont County."
47 Laramie, Wyoming: University of Wyoming. November 1997.
48
- 49 Ur-Energy USA, Inc. "Application for USNRC Source Material License Lost Creek Project,
50 South-Central Wyoming, Environmental Report." Littleton, Colorado: Ur-Energy USA, Inc.
51 ML073190539. October 2007.

Description of the Affected Environment

1 USACE. "Interim Regional Supplement to the Corps of Engineers Wetland Delineation Manual:
2 Arid West Region." 2006. Washington, DC: USACE. 2006.

3
4 USACE. "Find Notice of Issues and Modification of Nationwide Permits." Vol. 65, No. 47. pp.
5 12,897–12,899. 2000.

6
7 USACE. "Corps of Engineers Wetlands Delineation Manual." Technical Report Y-87-1.
8 Washington, DC: USACE. 1987

9 USACops. "Wyoming." <www.usacops.com> (15 April 2008).

10
11 U.S. Census Bureau. "American FactFinder 2000 Census Data." 2008.
12 <<http://factfinder.census.gov>> (25 February 2008).

13
14 USDA, Forest Service. "Landscape Aesthetics: A Handbook for Scenery Management."
15 Agriculture Handbook No. 701. Washington DC: USFS. 1995.

16
17 USFS. "Bighorn National Forest, Final Environmental Impact Statement for the Revised Land
18 and Resource Management Plan." Golden, Colorado: USFS, Rocky Mountain Region.
19 November 2005. <[http://www.fs.fed.us/r2/bighorn/projects/planrevision/](http://www.fs.fed.us/r2/bighorn/projects/planrevision/documents/final/index.shtml)
20 [documents/final/index.shtml](http://www.fs.fed.us/r2/bighorn/projects/planrevision/documents/final/index.shtml)> (17 October 2007).

21
22 U.S. Forest Service. "National Forest Landscape Management. Volume 2, Chapter 1. The
23 Visual Management System." Agriculture Handbook No. 462. Washington, DC: USFS. 1974.

24
25 U.S. Fish and Wildlife Service. "Mountain Prairie Region." 2008. <[http://www.fws.gov/](http://www.fws.gov/mountain%2Dprairie/)
26 [mountain%2Dprairie/](http://www.fws.gov/mountain%2Dprairie/)> (15 February, 2008).

27
28 U.S. Fish and Wildlife Service. "National Wetland Inventory Mapper." 2007.
29 <<http://www.fws.gov/nwi/>> (29 February 2008).

30
31 U.S. Geological Survey. "Water Watch." 2008. <<http://water.usgs.gov/waterwatch>>
32 (28 February 2008).

33
34 U.S. Geological Survey. "A Tapestry of Time and Terrain." Denver, Colorado: U.S. Geological
35 Survey. 2004. <<http://tapestry.usgs.gov/Default.html>> (25 February 2008).

36
37 Washington State Department of Transportation. "WSDOT's Guidance for Addressing Noise
38 Impacts in Biological Assessments—Noise Impacts." Seattle, Washington: Washington State
39 Department of Transportation. 2006. <[http://www.wsdot.wa.gov/TA/](http://www.wsdot.wa.gov/TA/Operations/Environmental/NoiseChapter011906.pdf)
40 [Operations/Environmental/NoiseChapter011906.pdf](http://www.wsdot.wa.gov/TA/Operations/Environmental/NoiseChapter011906.pdf)> (12 October 2007).

41
42 Whitehead, R.L. "Groundwater Atlas of the United States, Montana, North Dakota, South
43 Dakota, Wyoming." U.S. Geological Survey Report HA 730-I. Denver, Colorado:
44 U.S. Geological Survey. 1996. <http://capp.water.usgs.gov/gwa/ch_i/index.html>.

45
46 Wind River Country Wyoming. "Wind River Country.com." <www.windrivercountry.com>
47 (15 October 2007).

48
49 Wind River Visitor's Council. "Welcome to the Wind River Indian Reservation."
50 <<http://www.wind-river.org/info/wrindianreservation.php>> (02 April 2008).

- World Wildlife Fund. "Wildfinder—Mapping the World's Species: Ecoregion NA1313 (Wyoming Basin Shrub Steppe)." Washington, DC: World Wildlife Fund. 2007a. <<http://www.worldwildlife.org/wildfinder/searchByPlace.cfm?ecoregion=Na1313>> (15 October 2007).
- World Wildlife Fund. "Wyoming Basin Shrub Steppe." Washington, DC: World Wildlife Fund. 2007b. <http://www.worldwildlife.org/wildworld/profiles/terrestrial/na/na1313_full.html> (13 September 2007).
- World Wildlife Fund. "Colorado Rockies Forests." Washington, DC: World Wildlife Fund. 2007c. <http://www.worldwildlife.org/wildworld/profiles/terrestrial/na/na0511_full.html> (10 October 2007).
- World Wildlife Fund. "Northern Short Grasslands." Washington, DC: World Wildlife Fund. 2007d. <http://www.worldwildlife.org/wildworld/profiles/terrestrial/na/na0811_full.html> (13 September 2007).
- World Wildlife Fund. "South Central Rockies Forests." Washington, DC: World Wildlife Fund. 2007e. <http://www.worldwildlife.org/wildworld/profiles/terrestrial/na/na0528_full.html> (10 October 2007).
- WDEQ. "Chapter 2, Ambient Standards." 2006. <<http://deq.state.wy.us/aqd/standards.asp>> (23 October 2007).
- WDEQ. "Wyoming Surface Water Classification List." Cheyenne, Wyoming: WDEQ, Water Quality Division, Surface Water Standards. 2001.
- Wyoming Department of Revenue. "Sales and Tax Distribution Report by County." Cheyenne, Wyoming: Wyoming Department of Revenue. 2008. <<http://revenue.state.wy.us>> (25 February 2008).
- Wyoming Department of Transportation. "WYDOT Traffic Analysis." Cheyenne, Wyoming: Wyoming Department of Transportation. 2005. <<http://www.dot.state.wy.us/Default.jsp?sCode=hwyta>> (25 February 2008).
- Wyoming Game and Fish Department. "Comprehensive Wildlife Conservation Strategy." Cheyenne, Wyoming: Wyoming Game and Fish. 2008. <<http://gf.state.wy.us/wildlife/CompConvStrategy/Species>> (19 February 2008).
- Wyoming Game and Fish Department. "Official State List of Birds, Mammals, Amphibians, and Reptiles in Wyoming." Cheyenne, Wyoming: Wyoming Game and Fish. 2007a. <<http://gf.state.wy.us/wildlife/nongame/SpeciesList/index.asp>> (15 October 2007).
- Wyoming Geographic Information Science Center. "Land Ownership and Management for Wyoming." Laramie, Wyoming: University of Wyoming, Wyoming Geographic Information Service Center. 2008. <www.sdvc.uwyo.edu/clearinghouse/management.html> (15 February 2008).
- Wyoming Game and Fish Department. "Terrestrial Habitat/Aquatic Habitat/Habitat Management." Cheyenne, Wyoming: Wyoming Game and Fish. 2007b. <<http://gf.state.wy.us/habitat/aquatic/index.asp>> (13 September 2007).

Description of the Affected Environment

- 1 Wyoming Geographic Information Science Center. "Wyoming Gap Analysis." Laramie,
2 Wyoming: University of Wyoming, Wyoming Geographic Information Science Center. 2007a.
3 <<http://www.wygisc.uwyo.edu/wbn/gap.html>> (25 February 2008).
4
- 5 Wyoming Geographic Information Science Center. "Wyoming Gap Analysis—Download."
6 Laramie, Wyoming: University of Wyoming, Wyoming Geographic Information Science Center,
7 Wyoming Bioinformation Node. 2007b. <<http://www.sdvc.uwyo.edu/wbn/data.html>>
8 (25 February 2008).
9
- 10 Wyoming State Geological Survey, Industrial Minerals and Uranium Section. "Uranium Page."
11 <<http://www.wsgs.uwyo.edu/minerals/uranium.aspx>> July 15, 2005.
12
- 13 Wyoming Workforce Development Council. "Wyoming Workers Commuting Patters Study."
14 Cheyenne, Wyoming: Wyoming Workforce Development Council. 2007.
15
- 16 Zielinski, R.A. "Volcanic Rocks as Sources of Uranium." *Uranium Deposits in Volcanic Rocks*.
17 Vienna, Austria: International Atomic Energy Agency. pp. 83–95. 1984.
18
- 19 Zielinski, R.A. "Tuffaceous Sediments as Source Rocks for Uranium—A Case Study of the
20 White River Formation, Wyoming." *Journal of Geochemical Exploration*. Vol. 18. pp. 285–306.
21 1983.

3.3 Wyoming East Uranium Milling Region

3.3.1 Land Use

As shown on Figure 3.3-1, the Wyoming East Uranium Milling Region encompasses parts of eight counties (Albany, Campbell, Carbon, Converse, Johnson, Natrona, Platte, and Weston), although it predominantly lies within Converse and Campbell counties. This region straddles portions of the Wyoming Basin to the east and the upper part of the Missouri Plateau to the north (U.S. Geological Survey, 2004). In this region, past, current, and potential uranium milling operations are generally found in the four-corner area of Campbell, Converse, Natrona, and Johnson counties, (known as the Pumpkin Buttes District) and in the northern-central part of Converse County (known as the Monument Hill District). The Shirley Basin Uranium District located south of Casper is the past site of a conventional uranium milling facility (Figures 3.3-1 and 3.3-2). The geology and soils of these three uranium districts are detailed in Section 3.3.3.

While 53.3 percent of the land in Wyoming is federal and state public land, land ownership in this region is predominantly private (68 percent) (Table 3.3-1). Within the Wyoming East Uranium Milling Region there are portions of two large tracts of federal land that are managed by the U. S. Forest Service (USFS):

- The Thunder Basin National Grassland, which straddles Campbell, Converse, and Weston Counties in the Powder River Basin between the Big Horn Mountains to the west and the South Dakota Black Hills to the east, represents 15 percent of the region.
- The Medicine Bow National Forest, which occupies the southern part of Converse County and extends farther south into Albany County represents almost 6 percent of the region.

Although federal grasslands and forests occupy an important portion of the region (approximately 21 percent), most rangeland is privately owned (68 percent) and is primarily used for grazing cattle and sheep. Campbell County, for example has more private land ownership than any other county in Wyoming. Other federal lands managed by BLM, the U.S. Bureau of Reclamation, and the Department of Defense (Table 3.3-1) comprise scattered tracts mixed with state and private lands and represent only approximately 10 percent of the land in the Wyoming East Uranium Milling Region (Figure 3.3-1).

The open rangelands of this region consist of gently rolling hills covered by sagebrush and short grass prairies capable of supporting year-round cattle and sheep grazing. Compared to the productivity of the open rangeland, farmland is marginal. It consists of dry or locally irrigated grain, hay, and pasture crops for livestock grazing or for preparing livestock feed. Agriculture is limited in the region due to low precipitation and because other water resources are insufficient for irrigation.

In addition to providing forage for livestock and grazing, the Thunder Basin National Grassland provides a variety of recreational activities, such as sightseeing, hiking, camping, hunting, and fishing (USFS, 2008). The historic Bozeman, Oregon, and Bridger Trail Corridors (see Figure 3.1-2), extending north and north-northeast through Natrona and Johnson counties along the western edge of the Wyoming East Uranium Milling Region, also offer a variety of




Figure 3.3-2. Map Showing Outline of the Wyoming East Region and Locations of the Pumpkin Buttes and Monument Hill Districts in the Powder River Basin and the Shirley Basin Uranium District in the Shirley Basin

Table 3.3-1. Land Ownership and General Use in the Wyoming East Uranium Milling Region

Land Ownership and General Use	Area (mi ²)	Area (km ²)	Percent
Private Lands	5,503	14,252	68.3
U.S. Forest Service, National Grassland	1,238	3,207	15.4
U.S. Bureau of Land Management, Public Domain Land	797	2,064	9.9
U.S. Forest Service, National Forest	466	1,208	5.8
Bureau of Reclamation	36	92	0.4
U.S. Department of Defense (Navy)	14	35	0.2
Totals	8,054	20,859	100

recreational activities, including sightseeing, museums, historic sites and small state parks (Fort Phil Kearny/Bozeman Trail Association, 2008).

Oil and gas production facilities, coal mines and coal bed methane (CBM) facilities have been, and continue to be, developed throughout the federal and private rangeland of the Powder River basin. These coal, CBM, and oil and gas facilities are more prevalent and concentrated in the central and northern part of the Powder River basin in Campbell and Johnson counties. Given the abundance and density of CBM facilities in these counties, current and future permitted areas of ISL facilities of the Pumpkin Buttes District would be likely near or intermixed with such CBM sites. In the southern part of the Powder River basin in the Monument Hill District, there are only a few scattered CBM sites (U.S. Geological Survey, 2001). Future ISL facilities in the Monument Hill District therefore would not interfere with land use for CBM facilities.

3.3.2 Transportation

Past experience at NRC-licensed ISL facilities indicate these facilities rely on roads for transportation of goods and personnel (Section 2.8). As shown on Figure 3.3-3, the Wyoming East Uranium Milling Region is accessible from the west by Interstate 25, U.S. Highway 20 and State Route 220. From the north, the region is accessible via Gillette by State Route 59 or State Route 50. Travel from the east reaches the Wyoming East Uranium Milling Region using State Route 450 in the northern portion of the region and U.S. Highway 18 or U.S. Highway 26 further to the south. Southern access is from U.S. Highway 26 in the southeastern corner near Glendo and State Route 487 from the southwestern corner of the region. Rail lines traverse the southern part of the region following the path of Interstate 25. A rail spur forks north of Orin and generally follows State Route 59 north in the direction of Gillette.

Areas of interest in uranium milling in the region are shown in Figure 3.3-3. For discussion purposes, these areas are located in four main sub-regions when considering site access by local roads. Areas of milling interest that are located in the northwestern part of the region between Edgerton and Wright are accessed from Gillette to the north or from Casper to the south. A cluster of northernmost sites are accessed by local roads leading east to State Route 50 and then south to State Route 387 and either north to Gillette or south to Casper and Interstate 25. Along State Route 387, north of Edgerton, is another sub-region of Uranium milling interest. The midsection of the Wyoming East Uranium Milling Region, north of Douglas, Orpha, and Rolling Hills, is the third sub-region of concentrated milling interest. Local roads




Figure 3.3-3. Wyoming East Uranium Milling Region General Map With Current and Future Uranium Milling Site Transportation Corridor

including Ross Road provide access to this sub-region from the south using State Routes 93 and 95 that connect to Interstate 25. A rail spur runs north and dead ends into this area from the main line that follows Interstate 25. Further to the west in the direction of Casper, State Route 256 from Interstate 25 provides access for another milling site. The fourth sub-region of interest is in the southwestern corner of the Wyoming East Uranium Milling Region. This is the location of the Shirley Basin conventional milling site which is accessed using State Route 487 and 251 from Casper (and Interstate 25) to the north, or from the south on State Routes 487 and U.S. Highway 30 from Laramie.

Table 3.3-2 provides available traffic count data for roads that support areas of past or future milling interest in the Wyoming East Uranium Milling Region. Counts are variable with the minimum all vehicle count at 340 vehicles per day on State Route 93 at Orpha and the maximum on Interstate 25 Casper to State Route 95 at 10,220 vehicles per day. Most all vehicle counts in the Wyoming East Uranium Milling Region are above 900 vehicles per day.

Yellowcake product shipments are expected to travel from the milling facility to a uranium hexafluoride production (conversion) facility in Metropolis, Illinois (the only facility currently licensed by NRC in the United States for this purpose). Major interstate transportation routes are expected to be used for these shipments, which are required to follow NRC packaging and transportation regulations in 10 CFR Part 71 and U.S. Department of Transportation hazardous material transportation regulations at 49 CFR Parts 171–189. Table 3.3-3 describes representative routes and distances for shipments of Yellowcake from locations of Uranium milling interest in the Wyoming East Uranium Milling Region. Representative routes are considered owing to the number of routing options available that could be used by a future ISL facility.

3.3.3 Geology and Soils

As noted in Section 3.2.3, Wyoming contains the largest known reserves of uranium in the United States and has been the nation's leading producer of uranium ore since 1995 (Wyoming State Geological Survey, 2005). Sandstone-hosted uranium deposits account for the vast majority of the ore produced in Wyoming (Chenoweth, 1991). In the Wyoming East Uranium Milling Region, uranium mineralization is found in fluvial sandstones in two major areas: the Powder River Basin and the Shirley Basin (Figure 3.3-2). Uranium mineralization in sandstones in these two districts is in a geologic setting favorable for recovery by ISL milling. Since 1991, all uranium produced from sandstones in the Wyoming East Uranium Milling Region has been by the ISL method (Wyoming State Geological Survey, 2005).

The Powder River Basin encompasses an area of about 31,000 km² (12,000 mi²) in Converse and Campbell Counties. Uranium was first discovered in the Powder River Basin in 1951 near Pumpkin Buttes in the central part of the basin (Davis, 1969). Other uranium deposits were found along a 97-kilometer [60-mile] northwest-southeast trend in the southwest part of the Powder River Basin, and production began in 1953. Prior to 1968, total production from the Powder River Basin was slightly over 455,000 metric tons [500,000 tons] of U₃O₈ (Davis, 1969). The most important uranium deposits are in the Monument Hill district, which produced over 90% of the ore from the basin prior to 1968.

The Shirley Basin uranium area is mainly in the northeastern part of Carbon County (Figure 3.3-4). Uranium was discovered in the Shirley Basin in 1955 (Melin, 1969). Production began in 1960 from underground and open-pit mines. Milling by ISL began in 1964. Prior to

1

Table 3.3-2. Average Annual Daily Traffic Counts for Roads in the Wyoming East Uranium Milling Region*					
Road Segment	Distance (mi)	Trucks		All Vehicles	
		2005	2006	2005	2006
State Route 59 at Reno Junction (north of intersection with State Route 387)	—	690	750	3,630	3,930
State Route 387 at Pine Tree Junction (between State Routes 50 and 59)	20	210–410	220–410	970–3,130	970–3,130
State Route 387 at Edgerton North	—	380	440	2,110	2,140
Interstate 25 at Casper North (between Casper and State Route 259)	20	570–690	610–690	2,460–3,760	2,560–3,800
State Route 487 at Shirley Basin North (at intersection with State Route 251)	—	70	80	710	700
State Route 256 North Of Interstate 25	—	140	140	2,270	2,290
U.S. Highway 20/26 at Casper East (between Evansville and Parkerton)	0.5	200	230	2,900	2,900
Interstate 25 Casper to State Route 95	21	570–1,030	610–1,030	2,610–10,220	2,710–10,220
State Route 95 at Rolling Hills	—	50	50	1,800	1,810
State Route 93 at Orpha	—	50	50	340	340
State Route 59 Douglas to Bill	35	380–450	410–440	1,940–3,690	1,940–3,690
*Wyoming Department of Transportation. "Wyoming Department of Transportation Vehicle Miles." Data for Calendar Year 2005 and 2006 Provided on Request. District 2 Office, Casper, Wyoming: Wyoming Department of Transportation. April 18, 2008. 1 mi = 1.61 km					

2
3
4

Table 3.3-3. Representative Transportation Routes for Yellowcake Shipments From the Wyoming East Uranium Milling Region			
Origin	Destination	Major Links	Distance* (mi)
West of Savageton, Wyoming	Metropolis, Illinois	Local access road east to State Route 50 State Route 50 south to Route 387 State Route 387 south to Edgerton, Wyoming State Route 259 south to Interstate 25 Interstate 25 south to Casper, Wyoming Interstate 25 south to Denver, Colorado Interstate 70 east to St. Louis, Missouri Interstate 64 east to Interstate 57	1,420

1

Table 3.3-3. Representative Transportation Routes for Yellowcake Shipments From the Wyoming East Uranium Milling Region (continued)

Origin	Destination	Major Links	Distance (mi)
		Interstate 57 south to Interstate 24 Interstate 24 south to U.S. Highway 45 U.S. Highway 45 west to Metropolis, Illinois	
Northwest of Douglas, Wyoming	Metropolis, Illinois	Ross Road south to State Route 93 State Route 93 south to Interstate 25 Interstate 25 south to Denver, Colorado Denver, Colorado to Metropolis, Illinois (as above)	1,300
Shirley Basin Area, Wyoming	Metropolis, Illinois	Local access roads west to State Route 487 State Route 487 north to State Route 251 State Route 251 north to Casper, Wyoming Interstate 25 south to Denver, Colorado Denver, Colorado to Metropolis, Illinois (as above)	1,370
*American Map Corporation. "Road Atlas of the United States, Canada, and Mexico." Long Island City, New York: American Map Corporation. p. 144. 2006. 1 mi = 1.61 km			

1970, approximately 1,500 metric tons [1,600 tons] of U_3O_8 was produced from mines in the Shirley Basin (Chenoweth, 1991). The dominant source of sediment in the Powder River Basin and the Shirley Basin was Precambrian (greater than 453 million year old) granitic rock of the Sweetwater Arch and northern Laramie Range (Rackley, 1972; Harris and King, 1993). The Sweetwater Arch is also referred to as the Granite Mountains (Bailey, 1969; Anderson, 1969; Lageson and Spearing, 1988). The Sweetwater Arch and northern Laramie Range are mountain ranges composed of uraniferous granitic rock. Uplift of the Sweetwater Arch and Laramie Range began to affect sedimentation in the adjacent Powder River Basin and Shirley Basin in Late Cretaceous time (65 to 99 million years ago). Rapidly subsiding portions of these basins received thick clastic wedges (i.e., wedges made of fragments of other rocks) of predominantly arkosic sediments (i.e., sediments containing a significant fraction of feldspar), while larger, more slowly subsiding portions of the basins received a greater proportion of paludal (marsh) and lacustrine (lake) sediments.

Sediment in the west Shirley Basin was deposited on an alluvial fan, but in the east Shirley Basin and in the Powder River Basin sedimentation was channel and flood-plain deposits of a meandering stream (Rackley, 1972). Beginning in the middle Eocene (41 to 49 million years ago) and increasing in the Oligocene (23.8 to 33.7 million years ago), regional volcanic activity contributed a significant amount of tuffaceous materials (i.e., materials made from volcanic rock and mineral fragments in a volcanic ash matrix) to local sediments. Deposition within the basins probably continued through the Miocene (5.3 to 23.8 million years ago), but post-Miocene erosion has completely removed Oligocene and Miocene units.

A generalized stratigraphic section of Tertiary (1.8 to 65 million-year old) formations in the Wyoming East Uranium Milling Region is shown in Figure 3.3-5. Stratigraphic descriptions presented here are limited to formations that may be involved in potential milling operations or formations that may have environmental significance, such as important aquifers and confining units above and below potential milling zones.

1




Figure 3.3-4. Index and Structure Map of East-Central Wyoming Showing Relation of the Sweetwater Arch and Laramie Range to the Powder River Basin and the Shirley Basin. The Distribution of Arkosic, Carbonaceous Sediments and Mineralized Areas in the Powder River and Shirley Basins Are also Shown (Modified From Rackley, 1972).

1

Central Wyoming			
System	Series		Formation
Tertiary	Pliocene		Moonstone Formation
	Miocene		Split Rock Formation Arikaree Formation
	Oligocene		White River Formation
	Eocene	Upper	Wagon Bed Formation
		Middle	
		Lower	Wind River Formation Wasatch Formation
	Paleocene		Fort Union Formation
Cretaceous	Upper		Lance Formation

Figure 3.3-5. Stratigraphic Section of Tertiary Age Formations in the Powder River Basin and Shirley Basin of Central Wyoming. Major Sandstone-Type Uranium Deposits Are Hosted in the Wasatch Formation in the Powder River Basin and the Wind River Formation in the Shirley Basin (Modified From Harshman, 1968).

2

3 Formations hosting major sandstone-type uranium deposits in the Wyoming East Uranium
 4 Milling Region are the Wasatch Formation in the Powder River Basin and the Wind River
 5 Formation in the Shirley Basin. Both the Wasatch and Wind River are lower Eocene (49 to
 6 54.8 million years old) in age (Houston, 1969), and consist of interbedded, arkosic sandstone,
 7 conglomerate, siltstone, mudstone, and carbonaceous shale, all compacted but poorly
 8 cemented (Harshman, 1968). In the Powder River Basin, recoverable ore that can be exploited
 9 by ISL milling is located in parts of the Wasatch Formation extending from depths of 120 to
 10 300 m [400 to 1,000 ft] below the surface (Davis, 1969). Uranium deposits in the Shirley Basin
 11 lie at depths of 30 to 150 m [100 to 500 ft], almost entirely in the lower 90 m [300 ft] of the Wind
 12 River Formation (Melin, 1969; Bailey, 1969).

13

14 The Wagon Bed Formation conformably overlies the Wasatch and Wind River formations. The
 15 Wagon Bed comprises a series of interbedded arkosic sandstones and silicified claystones.
 16 Regionally, the Wagon Bed Formation may not be present in the central parts of the basins,
 17 having been removed by erosion. The White River Formation unconformably overlies the
 18 Wagon Bed Formation or the Wasatch and Wind River formations where the Wagon Bed has
 19 been removed by erosion. The White River consists of tuffaceous siltstone, claystone, and
 20 conglomerate with subordinate amounts of tuff. The White River overlaps older Tertiary
 21 formations and wedges out against pre-Tertiary rocks on the flanks of the basins. The White
 22 River Formation is overlain by the Split Rock Formation in the Shirley Basin and the Arikaree
 23 Formation in the Powder River Basin. The Split Rock and Arikaree consist of tuffaceous
 24 siltstone and sandstone beds that sometimes cap prominent ridges (Harshman, 1968).

1 The Fort Union Formation underlies the Wasatch and Wind River formations and, to a limited
2 extent, is also a host to sandstone-type uranium deposits (Davis, 1969; Langden, 1973). The
3 Fort Union is a fluvial deposit consisting of alternating and discontinuous mudstones, siltstones,
4 carbonaceous shales, and coarser arkosic sandstone. The Fort Union is unconformably
5 underlain by sediments of the Lance Formation, which is in turn underlain by a thick sequence
6 of older sandstones, mudstones, and shales.

7
8 The uranium deposits in the Wyoming East Uranium Milling Region are stratabound and
9 genetically related to geochemical interfaces, or roll-fronts (see Section 3.1.2). The roll-front ore
10 deposits in the Powder River Basin are usually multiple "C"-shaped rolls distorted by variations
11 in gross lithology (Davis, 1969). The principal ore minerals are uraninite, coffinite,
12 metatyuyamunite, and carnotite. Gangue minerals (i.e., low-value minerals intermixed with ore
13 minerals) are calcite, gypsum, pyrite, iron oxide, and barite (Mrak, 1968). Although most of the
14 uranium in the Shirley Basin is in roll-front deposits, important amounts also occur in tabular
15 bodies near the rolls. Tabular sandstone-hosted uranium deposits are found as blanket-like,
16 roughly parallel ore bodies along sandstone trends. The uranium mineralization in both the roll-
17 front and tabular deposits consists of disseminations and impregnations of uraninite, calcite,
18 pyrite, and marcasite in arkosic sandstones.

19
20 The source of uranium in sandstone-type uranium deposits in central Wyoming is a topic of
21 conjecture. Four theories on the source of uranium in these occurrences have been suggested:
22 (1) leached uranium from overlying ash-fall tuffs, (2) leached uranium from igneous and
23 metamorphic rocks in the highlands surrounding the basins, (3) leached uranium from the host
24 sandstones themselves, and (4) hydrothermal uranium from a magma source at depth (Harris
25 and King, 1993). Combinations of these theories have been proposed as well (Boberg, 1981).
26 The most popular theories are the tuff leach (1) and the highland leach (2). The tuff leach
27 theory is supported by extensive geochemical studies on uranium removal from tuff (Zielinski,
28 1983, 1984; Trentham and Orajaka, 1986). Further, it was the tuff leach theory that led to the
29 discovery of most of the large uranium deposits in Wyoming (Love, 1952). On the other hand,
30 many sandstone-hosted uranium deposits in Wyoming are found adjacent to crystalline rocks,
31 especially the uraniferous granites of the northern Laramie and Granite mountains (Harris and
32 King, 1993). Oxidized uranium leached from these crystalline terrains could have been
33 transported to the sites of present mineralization.

34
35 Soils within the Wyoming East Uranium Milling Region are diverse and can vary substantially in
36 terms of characteristics over relatively short distances. The distribution and occurrence of soils
37 in east-central Wyoming can vary both on a regional basis (mountains, foothills, basins) and
38 locally with changes in slope, geology, vegetation, climate, and time. In the Powder River Basin
39 and Shirley Basin, old, tilted sedimentary rocks occur in bands along the margins of the basins,
40 whereas younger sediments showing varying degrees of incision by erosion are found in the
41 basin centers.

42
43 The topographic position and texture of typical soils in the Powder River Basin and Shirley
44 Basin areas of east-central Wyoming was obtained from the Soils Map of Wyoming (Munn and
45 Arneson, 1998). This map was designed primarily for statewide study of ground water
46 vulnerability to contamination and would not be expected to be used for site-specific soil
47 interpretations at proposed ISL milling facilities. For site-specific evaluations, detailed soils
48 information would be expected to be obtained from published county soil surveys or the Natural
49 Resources Conservation Service (NRCS).

In the Powder River and Shirley basins, shallow loamy-skeletal (stony soils) with little or no subsoil development occupy ridge crests along the margins of the basins. These soils contain hard clasts (i.e., rock fragments) and tend to be much coarser than soils on the adjacent lower slopes. Loamy-skeletal soils with little subsoil development are also found in the foothills along the margins of the basin and along eroded drainageways. Fine to fine-loamy soils with moderate- to well-developed soil horizons are found on gently sloping to moderately steep slopes associated with alluvial fans and alluvial terraces. These soils are generally light-colored and depleted in moisture. Moderately-deep soils with well-developed soil horizons occur on low relief surfaces, such as stream terraces and floodplains, across broad expanses of the basins. Fine-loamy over sandy and coarse loamy soils occurs on stream terraces. Soils found on floodplains include fine loamy and fine sand loams. Dark-colored, base-rich soils formed under grass are generally associated with floodplains along streams with permanent high water.

3.3.4 Water Resources

3.3.4.1 Surface Waters

The Wyoming East Uranium Milling Region (Figure 3.3-6) includes portions of Albany, Campbell, Carbon, Converse, Johnson, Natrona, Platte, and Weston counties in east-central Wyoming. The watersheds within the Wyoming East Uranium Milling Region are listed in Table 3.3-4 along with range of designated uses of surface water bodies assigned by the State of Wyoming (WDEQ, 2001). Because surface water uses are designated for specific water bodies, such as stream segments and lakes, within a watershed and the specific locations of future uranium milling activities are not known at this time, the range of designated uses is provided rather than a listing of designated uses for each water body within a watershed. Not all water bodies within a watershed may have all of the designated uses listed in Table 3.3-4. For information regarding specific water bodies, the reader is referred to the Wyoming Department of Environmental Quality Surface Water Standards webpage deq.state.wy.us/wqd/watershed/surfacestandards.

The historical uranium milling districts included in the Wyoming East Uranium Milling Region are the Shirley Basin within the Little Medicine Bow watershed in the southwest and uranium deposits in the area known as the Powder River Basin that actually includes watersheds in addition to those contributing to the Powder River. Watersheds containing historical or potential uranium milling sites are: Middle North Platte-Casper, Lightning Creek, Dry Fork Cheyenne River, Antelope Creek, Salt Creek, and Upper Power River.

The Shirley Basin uranium district is located within the Little Medicine Bow River watershed (Figure 3.3-6) in Carbon and Albany counties. In addition to the Little Medicine Bow River, other significant surface water features associated with the Shirley Basin are Sand Creek and Muddy Creek. Several small reservoirs are located on these streams. Several unnamed springs are also shown on the topographic maps covering the Shirley Basin. The Little Medicine Bow River and most of its tributaries are generally Class 2AB waters with some classified as 2C and 3B (Table 3.3-4). The difference between Class 2AB and Class 2C waters is that Class 2C waters do not have drinking water supply or game fish as designated uses. Class 3B also excludes non-game fish and fish consumption as designated uses. Although the Little Medicine Bow River flows directly through an area of historic uranium mining and milling, it is not listed as an impacted or threatened water body (WDEQ, 2006). The average flow of the Little Medicine Bow River at Boles Spring, Wyoming is 0.3m³/s [11 ft³/s] (U.S. Geological Survey, 2008).

3.3-13




Figure 3.3-6. Watersheds Within the Wyoming East Uranium Milling Region

**Table 3.3-4. Primary Watersheds in the Wyoming West Uranium Milling Region
Range of Designated Uses of Water Bodies Within Each Watershed**

Watershed	Range of State Classification of Designated Uses *
Little Medicine Bow River and Tributaries	Generally 2AB with some tributaries 2B and 3C
Glendo Reservoir and Tributaries	2AB and 3B
Middle North Platte River	2AB with some tributaries 3B
Salt Creek	2C
Lightning Creek	3B
Dry Fork Cheyenne River	3B
Antelope Creek	3B
Upper Cheyenne River	3B
Upper Powder River	2ABww with some tributaries 3B
Upper Belle Fourche River and Tributaries	2ABww and 3B
<p>*Class 1 waters have designated uses including: Drinking Water, Game Fish, Non-Game Fish, Fish Consumption, Other Aquatic Life, Recreation, Wildlife Agriculture, Industry, Scenic Value.</p> <p>Class 2AB waters have designated uses including: Drinking Water, Game Fish, Non-Game Fish, Fish Consumption, Other Aquatic Life, Recreation, Wildlife Agriculture, Industry, Scenic Value.</p> <p>Class 2A waters have designated uses including: Drinking Water, Other Aquatic Life, Recreation, Wildlife Agriculture, Industry, Scenic Value.</p> <p>Class 2B waters exclude drinking water from the Class 2AB uses. Class 2C waters exclude drinking water and game fish from the Class 2AB uses.</p> <p>Class 3A, 3B and 3C waters have designated uses including: Other Aquatic Life, Recreation, Wildlife Agriculture, Industry, Scenic Value.</p> <p>Class 4A, 4B and 4C waters have designated uses include: Recreation, Wildlife Agriculture, Industry, Scenic Value.</p> <p>Class 2ABww and 2Bww are warm water fisheries.</p>	

The Powder River Basin contains the most extensive uranium deposits in Wyoming, covering a large portion of east-central Wyoming in Converse, Campbell and Johnson counties. Principal watersheds within the Powder River Basin uranium district are (from south to north, Glendo Reservoir (on the North Platte River), Middle North Platte-Casper, Lightning Creek, Dry Fork of the Cheyenne River, Antelope Creek, Salt Creek, Upper Cheyenne River, Upper Belle Fourche and Upper Powder River. The Lightning Creek, Antelope Creek, Dry Fork of the Cheyenne River and Upper Cheyenne River watersheds contain ephemeral and intermittent streams that flow to the Cheyenne River east of the uranium districts in the Powder River Basin. Other surface water features in these watersheds include stock ponds. The ephemeral and intermittent water bodies are generally Class 3B. These watersheds include areas of oil and natural gas as well as coal bed methane development.

The Middle North Platte-Casper watershed is drained by the North Platte River which is feed by numerous small tributaries. The North Platte River and most of its tributaries are classed as 2AB (Table 3.3-4). Portions of the North Platte River and some tributaries are impacted by elevated selenium concentrations (WDEQ, 2006). The flow of the North Platte River is not measured in this watershed.

The Salt Creek watershed is located north of Casper, Wyoming in Natrona County upstream from the Upper Powder River watershed. Salt Creek is a Class 2C water body (Table 3.3-4).

The water quality of Salt Creek is impaired due to elevated chloride and threatened by oil and

grease attributed to oil and natural gas production in the watershed. Flow in Salt Creek is not measured.

The Upper Belle Fourche River watershed is located in the northeastern portion of the Wyoming East Uranium Milling Region in Campbell County (Figure 3.3-6). The Upper Belle Fourche River in Wyoming is classed as 2ABww where "ww" indicates "warm water fishery" (Table 3.3-4). Water quality in some portions of the Upper Belle Fourche River is listed as impaired due to fecal coliform from livestock grazing east of the Wyoming East Uranium Milling Region (WDEQ, 2006). Average flow in the Upper Belle Fourche River at Moorcroft, Wyoming (just east of the Wyoming East Uranium Milling Region) is 0.4 m³/min [15 cubic ft/min] (U.S. Geological Survey, 2008).

The Upper Powder River watershed is located downstream of the Salt Creek watershed in Johnson and Campbell counties. The Upper Powder River is classified as 2ABww with its smaller tributaries classed as 3B (Table 3.3-4). The Upper Powder River is listed as impacted by high chloride (WDEQ, 2006). Average flow in the Upper Powder River at Sussex, Wyoming is 5.6 m³/min [199 cubic ft/s] (U.S. Geological Survey, 2008).

3.3.4.2 Wetlands and Waters of the United States

The majority of waterways in this region are comprised of ephemeral and intermittent streams. Some perennial slow moving rivers are also present in the region. Regulatory guidance and jurisdictional determination are the same as those found in Section 3.2.4.2 for Wyoming West Uranium Milling Region.

Freshwater emergent marshes are found in depressions, as fringes around lakes, and sloughs along slow-moving streams. These wetlands maybe temporarily to permanently inundated and are typically dominated by floating-leaved plants in deeper areas (e.g., *Lemna*, *Potamogeton*, *Brasenia*, *Nuphar*) and sedges (*Carex*, *Cyperus*, *Rhynchospora*), bulrushes (*Scirpus*, *Schoenoplectus*), spikerushes (*Eleocharis*), cattails (*Typha*), rushes, (*Juncus*), and grasses (e.g., *Phalaris*, *Spartina*) in seasonal wetlands (USACE, 2006).

Floodplain and riparian systems occur along rivers and streams across Wyoming East Uranium Milling Region. Common woody species in riparian and floodplain wetlands in the region include plains cottonwood (*Populus deltoides* ssp. *monilifera*), narrowleaf cottonwood (*P. angustifolia*), various willows, green ash (*Fraxinus pennsylvanica*), cedar elm, eastern swampprivet (*Forestiera acuminata*), and the introduced saltcedar (*Tamarix ramosissima*) (USACE, 2006).

Waters of the United States and special aquatic sites that include wetlands would need to be identified and the impact delineated upon individual site selection. Based on impacts and consultation with each area, appropriate permits would be obtained from the local USACE district. Section 401 state water quality certification is required for work in Waters of the United States. Within this region, the state of Wyoming regulates isolated wetlands and waters. Cumulative total project impacts greater than 1 acre would require a general permit for wetland mitigation by the WDEQ.

3.3.4.3 Groundwater

Groundwater resources in the Wyoming East Uranium Milling Region are part of regional aquifer systems that extend well beyond the areas of uranium milling interest in this part of Wyoming. Uranium bearing aquifers exist within these regional aquifer systems in the Wyoming East Uranium Milling Region. This section provides a general overview of the regional aquifer systems to provide context for a more focused discussion of the uranium bearing aquifers in the Wyoming East Uranium Milling Region, including hydrologic characteristics, level of confinement, groundwater quality, water uses, and important surrounding aquifers.

3.3.4.3.1 Regional Aquifer Systems

The location of the Wyoming East Uranium Milling Region is shown in Figures 3.3-1 and 3.3-2. The Northern Great Plains aquifer system is the major regional aquifer system in the Wyoming East Uranium Milling Region. The Northern Great Plains aquifer system extends over one-third of Wyoming (Whitehead, 1996).

The Northern Great Plains aquifer system includes confined Tertiary- and Cretaceous-aged sandstone aquifers and Paleozoic carbonate aquifers. The regional groundwater flow direction in this confined aquifer system is generally from southwest to northeast. The aquifer system is overlain by Quaternary-aged unconsolidated glacial and alluvial deposits that host shallow groundwater flow system. The Northern Great Plains aquifer system is underlain by crystalline rocks with low water yields. Recharge to the aquifer is by precipitation, water seeps from streambeds, and local irrigation. Discharge from the aquifer system is mainly by upward leakage of water into the shallower aquifers.

Whitehead (1996) grouped the Northern Great Plains aquifer system into five major aquifers. These aquifers, from shallowest to deepest, are the Lower Tertiary, Upper Cretaceous, Lower Cretaceous, Upper Paleozoic, and Lower Paleozoic aquifers. The Lower Tertiary aquifers consist of sandstone beds within the Wasatch Formation and the Fort Union Formation. Both formations consist of alternating beds of sandstone, siltstone, and claystone, but most water is stored in and flows through the more permeable sandstone beds. In the Powder River Basin, the Fort Union Formation and the Wasatch Formation are as thick as 1,095 m [3,600 ft] and 305 m [1,000 ft], respectively. In the Lower Tertiary aquifers, the regional groundwater flow direction is northward and northeastward from recharge areas in northeastern Wyoming.

The Upper Cretaceous aquifers consist of sandstone beds interbedded with siltstone and claystone in the Lance and the Hell Creek Formations and the Fox Hills Sandstone, which are 105 to 1,035 m [350 to 3,400 ft] and 90 to 135 m [300 to 450 ft thick]. The Fox Hills Sandstone is one of the most continuous water-yielding formations in the Northern Great Plains aquifer system. Groundwater in the Upper Cretaceous aquifers moves from aquifer recharge areas at higher altitudes toward discharge areas along major rivers. The general groundwater flow direction is northward in the Powder River Basin. In Wyoming, the potentiometric surface of the lower Tertiary aquifers is locally 122 m [400 ft] higher than that of the underlying upper Cretaceous aquifers. Hence, groundwater moves locally vertically downward from the lower Tertiary aquifers into the upper Cretaceous aquifers through the confining layer separating these two aquifers.

The Lower Cretaceous aquifers are separated from the overlying Upper Cretaceous aquifers by several thick confining units. The Pierre Shale, the Lewis Shale and the Steele Shale are the

regionally thickest and most extensive confining units. Water across the Pierre Shale can leak into the underlying Lower Cretaceous aquifers where the Pierre Shale is fractured.

The Lower Cretaceous aquifers are the most widespread aquifers in the Northern Great Plains aquifer system and contain several sandstones. The principal water-yielding units are the Muddy Sandstone and the Inyan Kara Group in the Powder River Basin. The Lower Cretaceous aquifers contain little freshwater. The water becomes saline in the deep parts of the Powder River Basin. Locally, the Sundance, Swift, Rierdon, and Piper Formations yield small to moderate quantities of water.

The Paleozoic aquifers cover a larger area, but they are deeply buried in most places and contain little freshwater. They are divided into Upper Paleozoic aquifers and Lower Paleozoic aquifers. In much of the Powder River Basin, the Upper and Lower Paleozoic aquifers are hydraulically connected and locally are called the Madison aquifer system.

The Upper Paleozoic aquifers are confined everywhere except in recharge areas. They consist primarily of the Madison Limestone, the Tensleep Sandstone in the western parts of the Powder River Basin and sandstone beds of the Minnelusa Formation in the eastern part of the Powder River Basin. The Pennsylvanian sandstones yield less water than the Madison Limestone and contain freshwater locally at the outcrop areas. Pennsylvanian rocks are not usually considered to be a principal aquifer. In the Upper Paleozoic aquifers, the regional groundwater flow direction is northeastward from recharge areas where the aquifers crop out adjacent to structural uplifts near the southern and western limits of the aquifer system.

Lower Paleozoic aquifers consist of sandstone and carbonate rocks. The principal geologic units that compose the lower Paleozoic aquifers are the Flathead Sandstone, sandstone beds of the Winnipeg Formation, limestones of the Red River and the Stonewall Formations, and the Bighorn and the Whitehead Dolomites. The groundwater flow direction is generally northeastward. Lower Paleozoic aquifers contain freshwater only in a small area in north-central Wyoming. These aquifers contain slightly saline to moderately saline water throughout the southern half of their extent.

The Madison Limestone exhibits karst features (features formed by the dissolution of a layer or layers of soluble bedrock, usually carbonate rock such as limestone or dolomite) at the outcrop areas in north-central Wyoming (Wyoming East region). Several large springs formed from some of the solution conduits in the Madison Limestone, including the Thermopolis hot springs system in central Wyoming with a discharge rate of about 11,355 L/min [3,000 gal/min] of geothermal water.

Recharge to the aquifers in most of the area is likely small, due to low annual precipitation and high evaporation. The mean annual precipitation in the Wyoming East Uranium Milling Region is typically in the range of 28-38 cm/year [11-15 in/year], but at high elevations, it locally exceeds 50 cm/year [20 in/year] based on precipitation data from 1971 to 2000. The evaporation rate was estimated to be 105.9 ± 7.1 cm/year [41.7 ± 2.8 in/year] using the Kohler-Nordenson- Fox equation with data from the station in Lander, Wyoming (Curtis and Grimes, 2004).

3.3.4.3.2 Aquifer Systems In The Vicinity Of Uranium Milling Sites

The hydrogeological system in areas of uranium milling interest in the Wyoming East Uranium Milling Region consists of a thick sequence of primarily sandstone aquifers and shale aquitards.

Description of the Affected Environment

Uranium-bearing sandstone aquifers in the Fort Union Formation at the active Uranium milling sites are also important for water supplies in the milling region.

Areas of uranium milling interest at the Reynolds and Smith Ranch area are underlain, from shallowest to deepest, by the alluvium, the Wasatch Formation, the Fort Union Formation, the Lance Formation, and the Fox Hills Formation. The alluvium has a thickness of 0 – 9 m [0 – 30 ft] and has small yields in stream valleys. The Wasatch Formation and the Fort Union Formation contain important sandstone aquifers for water supplies. Groundwater production from the Lance and the Fox Hills Formations are largely unknown at the ISL facilities in the Reynolds and Smith Ranch areas in Converse County (PRI, 2004).

As discussed in Section 3.3.4.3.1, this aquifer system is separated from the underlying aquifers including, from shallowest to deepest where they are continuous, the Muddy Sandstone, the Inyan Kara Group, and the Paleozoic aquifers by shale layers. The Paleozoic aquifers are deeply buried in most places and contain little freshwater (Whitehead, 1996).

3.3.4.3.3 Uranium-Bearing Aquifers

Uranium mineralization at locations of milling interest is typically hosted by the Paleocene-age confined sandstone aquifers in the Wyoming East Uranium Milling Region.

Confined sandstone beds in the Fort Union Formation are the uranium bearing aquifers in the Wyoming East Uranium Milling Region. At the Smith Ranch and Reynolds Ranch ISL sites the Pumpkin Buttes district in Converse County, the Fort Union Formation contains multiple confined sandstone aquifers in the eastern and northeastern parts of the permit area, but it is unconfined in the southwestern and western parts. Among the confined sandstone aquifers, the U- and S-Sandstones are the primary uranium mineralization zone and they are referred to as the U/S sand. O-Sandstone aquifers also contain economic uranium mineralization in the Fort Union Formation (NRC, 2006).

For ISL operations to begin, portions of the uranium-bearing sandstone aquifers in the Fort Union Formation in the Wyoming East Uranium Milling Region would need to be exempted by the UIC program administered by WDEQ (Section 1.7.2.1).

Hydrogeological characteristics: In the Wyoming East Uranium Milling Region, the production aquifer system typically consists of confined sandstone aquifers. Aquifer properties (e.g., transmissivity, thickness, storage coefficient) vary spatially in the region.

At the Smith Ranch and Reynolds Ranch areas, the mean effective transmissivity of the U/S sandstone aquifer and O-sandstone aquifer is 6,700 L/day/m [540 gal/day/ft {8.2 m²/day}] and 7,900 L/day/m [640 gal/day/ft {9.7 m²/day}], respectively. The storage coefficient for the U/S sandstone aquifer and O-sandstone aquifer ranges between 1.5×10^{-5} and 1.7×10^{-5} and 6.3×10^{-5} to 7.8×10^{-5} , respectively, indicating the confined nature of the production aquifer (typical storage coefficients for confined aquifers range from 10^{-5} - 10^{-3} (Driscoll, 1986; p.68)). The average groundwater velocities through the U/S-sandstone aquifer and O-sandstone aquifer were reported to be 2.4 m/yr [8 ft/yr] and 0.17 m/yr [0.56 ft/yr] (NRC, 2006). The approximate thickness of the of the Fort Union Formation is 910 – 1,100 m [3000 -3600 ft] in the Powder River Basin (PRI, 2004; Whitehead, 1996). Groundwater production from the Fort Union Formation is generally good with water yields as high as 2,080 L/min [550 gal/min] (PRI, 2004; NRC, 2006).

Level of confinement: The production aquifer is typically confined in the Wyoming East Uranium Milling Region. The thickness of the confinement varies spatially.

At the Smith Ranch and Reynolds Ranch ISL sites, the U/S sandstone is confined above by a 6–20 m [20–70 ft] thick shale aquitard (V Shale). It is confined below by a 45 m [150 m] thick

shale aquitard (R Shale) (NRC, 2006). Aquifer tests revealed that the confining shale members would be effective aquitards to the vertical movement of leaching solution (PRI, 2005).

As discussed in Section 3.3.4.3.1, the aquifer sequence that includes, from the shallowest to deepest, the Wasatch Formation, the Fort Union Formation, the Lance Formation, and the Fox Hills Formation are confined below by regionally extensive and thick low permeability layers that include the Pierre Shale, the Lewis Shale and the Steele Shale. The vertical hydraulic conductivity of the Pierre Shale is reported to be $1.5 \times 10^{-8} - 1.5 \times 10^{-4}$ m/day [$5 \times 10^{-8} - 5 \times 10^{-4}$ ft/day] outside the Wyoming East Uranium Milling Region (Kansas Geological Survey, 1991). The Pierre Shale is fractured in some parts of the region and may leak water to the underlying lower Cretaceous aquifers (Whitehead, 1996). Hence, where the Pierre Shale is fractured, the aquifer sequence may not be effectively confined below.

Groundwater quality: In some parts of the Wyoming East Uranium Milling Region, the total dissolved solids (TDS) levels in the uranium-bearing aquifers exceed the EPA's drinking water standards. The uranium and radium-226 concentrations in the uranium-bearing aquifers typically exceed their respective EPA Maximum Contaminant Levels.

At the Smith Ranch and Reynolds Ranch ISL area, the water quality is usually good in the U/S-sandstone and O-sandstone aquifers and meets the EPA's drinking water standards except for radium-226. Radium-226 naturally exists in the U/S sandstone and O-sandstone aquifers at a level of 296 pCi/L and 86 pCi/L, respectively, which exceeds the EPA's primary drinking water standard of 5 pCi/L. Both aquifers have TDS ranging from 234–952 mg/L [234–952 ppm] {the limit of dissolved solids recommended by the EPA for drinking water is 500 mg/L [500 ppm]} (NRC, 2006).

Current groundwater uses: In the vicinity of the Smith Ranch and Reynolds Ranch ISL area permit area, groundwater is largely pumped for livestock watering, and to a lesser extent, for domestic water supply (NRC, 2006).

3.3.4.3.4 Other Important Surrounding Aquifers for Water Supply

At the regional scale, the Wasatch Formation and the Fort Union Formation are important aquifers for water supplies. The Fox Hills Sandstone is one of the most continuous water-yielding formations in the Northern Great Plains aquifer system. Except at outcrop areas, the Paleozoic aquifers are not usually used for water production, because they are either deeply buried or contain saline water (Whitehead, 1996).

At the ISL facilities in the Reynolds and Smith Ranches, The Wasatch Formation and the Fort Union Formation contain important sandstone aquifers for water supplies. The thickness of the Wasatch Formation ranges from 0–150 m [0–500 ft] and yields as high as 530 L/min. Water yields from the Lance Formation and the Fox Hills Formations are largely unknown at the Reynolds and Smith Ranch areas. The thickness of the Lance Formation is about 915 m

[3,000 ft] and its water yield is estimated to not exceed 75 L/min [20 gal/min]. The thickness of the underlying Fox Hills Formation is about 150–210 m [500–700 ft] and its water yield is estimated to be not exceeding 380 L/min [100 gal/min] (PRI, 2004 and the references therein).

3.3.5 Ecology

3.3.5.1 Wyoming East Uranium Milling Flora

According to the EPA, the identified ecoregions in the Wyoming East Uranium Milling Region primarily consist of Wyoming Basin, Northern Great Plains, Southern Rockies, and the Western High plains ecoregions (Figure 3.3-7). Uranium milling districts in this region are generally found in the Rolling Sagebrush Steppe and the Powder River Basin of the Wyoming Basin. Habitat types and species found in these areas are based on the Wyoming Gap Analysis project (Wyoming Geographic Information Science Center, 1007) as described in Section 3.2.5.

The Rolling Sagebrush Steppe and the Salt Desert Shrub Basins ecoregions of the Wyoming Basin have been described in the Wyoming West Uranium Milling Region (Section 3.2.5). An excellent description of the Wyoming East Uranium Milling Region Fauna is provided by Chapman, et al. (2004) and is summarized below.

The Southern Rockies are characterized by rugged, steep mountains, intermontane depressions and open meadows, and high-elevation plateaus. Ponderosa pines are found at lower elevations with pinyon-juniper below that, grasslands are located in the lowest areas. Lodgepole pine is more common in the Middle Rockies region; white pine, grand fir, and cedar, prevalent in the Northern Rockies region, are absent from the Alpine zone. A greater portion of the Middle Rockies is used for summer grazing of livestock (Chapman, et al., 2004).

The Subalpine Forests ecoregion of the Southern Rockies is a forested area found on the steep forested slopes of the Medicine Bow and Sierra Madre mountains with a greater extent on the north slopes. The dense forests are dominated by lodgepole pine (*Pinus contorta*), Englemann spruce (*Picea engelmannii*) and subalpine fir (*Abies lasiocarpa*); some areas are locally dominated by aspen. Whortleberry dominates the forest understory. Subalpine meadows also occur in some areas (Chapman, et al., 2004).

The Mid-Elevation Forests and Shrublands ecoregion of the Southern Rockies is found in the 2,300 to 2,750 m [7,500 to 9,000 ft] elevation range within the Laramie, Medicine Bow, and Sierra Madre mountains. Vegetation located in the region from the southwest to northeast are comprised of aspen (*Populus tremula*), Douglas fir (*Pseudotsuga menziesii*), lodgepole pine, limber pine (*Pinus flexilis*), and ponderosa pine (*Pinus ponderosa*). Due to the increased availability of moisture Ponderosa pine grows mainly on the eastern slopes of the Laramie Mountains, as it does on the eastern Bighorn Mountains. The understory is composed of grasses and shrubs. Perennial streams are diverted for irrigation in lower elevations and are often dry in their lower reaches in the summer (Chapman, et al., 2004).

The Foothill Shrublands ecoregion of the Southern Rockies is a transitional between the higher elevation forests of the Laramie, Medicine Bow, and Sierra Madre mountains to the more arid grassland and sagebrush regions in the Wyoming Basin and the High Plains. On the east side of the Laramie Mountains, this ecoregion is a continuation of high plains prairie grasslands of blue grama, prairie junegrass, and western wheatgrass interspersed with mountain big sagebrush and mountain mahogany shrubland. Pockets of aspen, limber pine, and Douglas fir

1




Figure 3.3-7. Ecoregions of the Wyoming East Uranium Milling Region

2

Description of the Affected Environment

are often found on north-facing slopes. Riparian vegetation along the water courses originating in higher mountains include willow species and narrowleaf cottonwood, with boxelder (*Acer negundo*) and wild plum in the north. Land use is mainly livestock grazing and some irrigated hayland adjacent to perennial streams (Chapman, et al., 2004).

The High Plains ecoregion consists of rolling plains and tablelands formed by uplift and the erosion of the Rocky Mountains. Due to the rainshadow of the Rocky Mountains drought resistant shortgrass and mixed-grass prairie dominate the plains vegetation. Seasonal precipitation in this region generally falls during the growing season. This region occupies the southeastern corner of Wyoming where the Southern Rockies, Wyoming Basin, and the Northwestern Great Plains ecoregions meet. The boundaries of these regions fade into one another and some characteristics of each region can be found near the borders, making the boundary of the High Plains in Wyoming a transitional area.

The Moderate Relief Rangeland ecoregion of the High Plains consists of mixed-prairie vegetation dominated by grass species such as blue gramma, western winter wheatgrass, junegrass, Sandberg blue stem needle-and-thread, prairie junegrass, and winter fat. Other species found in the prairie include rabbitbrush, fringed sage, scattered yucca, and other various forbs. Patches of mountain mahogany and skunkbush sumac grow on bluffs and hilltops. The plains surface steadily increases in elevation as it rises to a subtle boundary transition with the Laramie Mountains (Chapman, et al., 2004).

The Pine Bluffs and Hills ecoregion of the High Plains is composed of escarpments, bluffs, and badlands. Ponderosa pine woodland and open grasslands alternate along the rocky outcrops. Common species found in this region include little blue stem, common juniper, and bearberry (*Arctostaphylos uva-ursi*). Areas of limber pine and sliver sagebrush may also be present (Chapman, et al., 2004).

The Sandy and Silty Tablelands ecoregion of the High Plains is characterized by tablelands with areas of moderate relief. This region consists of mixed-grass prairies dominated by blue gramma, western wheatgrass, june grass, needle-and-thread grass, rabbit brush, fringe sage, and various forbs. Since the 1880s Ecoregion 25g has been mainly used for livestock grazing (Chapman, et al., 2004).

The Northwestern Great Plains encompass the Missouri Plateau section of the Great Plains. This area includes semiarid rolling plains of shale and sandstone derived soils punctuated by occasional buttes and badlands. For the most part, it has not been influenced by continental glaciation. Cattle grazing and agriculture with spring wheat and alfalfa farming are common land uses. Agriculture is affected by erratic precipitation and limited opportunities for irrigation. In Wyoming, mining for coal and coal-bed methane production is prevalent, with a large increase in the number of coal-bed methane wells drilled in recent years. Native grasslands and some woodlands persist, especially in areas of steep or broken topography (Chapman, et al., 2004).

The Pine Scoria Hills ecoregion is composed of rugged broken land and stony rough hills covered by open ponderosa pine-Rocky Mountain juniper forest or ponderosa pine savannas. Coal, sandstone, and shale bedrock underlie the region. Savannas and extensive open grassland are found in areas with less available moisture. Species found in this region include little blue stem (*Schizachyrium scoparium*), bluebunch wheatgrass (*Pseudoroegneria spicata*), Idaho fescue (*Festuca idahoensis*), western wheatgrass, blue grama, and Sandberg bluegrass. Skunkbush sumac (*Rhus trilobata*) and western snowberry (*Symphoricarpos occidentalis*) are

common shrubs. Land use includes woodland grazing and areas of historical small-scale coal mining (Chapman, et al., 2004).

The Casper Arch ecoregion of the Northwestern Great Plains is a transitional region between the Northern Great Plains and the Wyoming Basin. Soils are weathered from sodic Cody shale; they are generally well drained to slowly permeable, and are moderately to very shallow. Shrubland dominated by sagebrush steppe, which may include, Wyoming big sagebrush, Gardner saltbush (*Atriplex gardneri*), Indian ricegrass (*Oryzopsis hymenoides*), birdfoot sagebrush (*Artemisia pedatifida*), western wheatgrass, bluebunch wheatgrass, needle-and-thread grass, blue grama, Sandberg bluegrass, junegrass, rabbitbrush, fringed sage, and other grasses, forbs, and shrubs (Chapman, et al., 2004).

The Powder River Basin ecoregion of the Northwestern Great Plains covers rolling prairie and dissected river breaks surrounding the Powder, Cheyenne, and upper North Platte rivers. The Powder River Basin has less precipitation and less available water than the neighboring regions. Vegetation within this region is composed of mixed-grass prairie dominated by blue grama, western wheatgrass, junegrass, Sandberg bluegrass, needle-and-thread grass, rabbitbrush, fringed sage, and other forbs, shrubs and grasses (Chapman, et al., 2004).

Wyoming East Uranium Milling Region Fauna

The animal species that may occur in the Wyoming Basin and the Middle/Southern Rockies have been discussed previously in the Wyoming West Uranium Milling Region (see Section 3.2.5.1)

The Northwest Great Plains/Northern short grasslands region of Wyoming is home to approximately 337 different species. Many of these species are found in the adjacent Wyoming Basin Shrub Steppe (World Wildlife Fund, 2007d,e). Many of the animals in this region are associated with prairie potholes. Birds include the Ferruginous hawk (*Buteo regalis*), Swainson's hawk (*Buteo swainsoni*), golden eagle, sharp tailed grouse (*Tympanuchus phasinellus*), sage grouse (*Centrocercus urophasianus*), the greater prairie chicken (*Tympanuchus cupido*), numerous migratory birds such as ducks and song birds, and one of the largest breed populations of the endangered piping plover (*Charadrius melodus*). Blacktail and whitetail deer, pronghorns, bighorn sheep, American bison (*Bison bison*), bobcat (*Lynx rufus*), and cougars (*Felis concolor*) are typical large animals. This region is also known for its abundance of white-tailed prairie dog towns, which the black-footed ferret uses as a habitat (World Wildlife Fund, 2007a–e).

The Western High Plains/Western Short Grasslands is home to approximately 431 different species. Many of these species can be found in the adjacent Northwest Great Plains region to the north. Rodents are the most numerous type of mammals of this region. These include Desert and Eastern cotton tail rabbits, gophers (*Thomomys sp.*), shrews (*Sorex sp.*), voles (*Microtus sp.*), kangaroo rats (*Dipodomys sp.*), black tailed prairie dogs, and numerous rats and mouse species. Larger mammals include the pronghorns, elk, big horn sheep, coyote, beaver (*Castor canadensis*), porcupine, bobcats, and foxes. The largest diversity of animals of the region is birds. Birds include the Ferruginous hawk, Swainson's hawk, golden eagle, sharp tailed grouse, prairie chickens, wrens, kingbirds, vireos sparrows, flycatchers, and ducks. This region contains numerous reptile and amphibians. Amphibian species include the northern cricket frog, leopard frog, bull frog, Rio Grande frog, narrowmouth toad, great plains toad, green toad, tiger salamander, and Woodhouse's toad. Western rattle snake ring-necked snake, king

snakes, hog-nose snake, and garter snake can be found in the region. Numerous lizards and turtles are also found within the region (World Wildlife Fund, 2007 a–e).

According to the Wyoming Game and Fish Department, crucial wintering habitats are found within this region for large game mammals and nesting leks for the sage grouse. Figures 3.3-8 to 3.3-14 show the crucial winters and yearlong ranges for large mammal found in this region. Most of the crucial areas are located either in the Thunder Basin National Grassland in the northeast portion of the region, the Medicine Bow National Forest in the Laramie Mountains, or along the North Platte River and its tributaries that traverse west-east across the lower half of the region. Within this region, the area of milling interest nearest to Casper is situated in close proximity to a crucial wintering area for antelopes. Numerous Sage Grouse leks are clustered near the Pumpkin Buttes Uranium District northwestern part of the study region. In addition, a large concentration of leks is found in the southwestern corner of the study region in the vicinity of the Shirley Basin Uranium District.

3.3.5.2 Aquatic

Within the Wyoming East Uranium Milling Region, watersheds identified as aquatic habitat areas include the Lower Salt Creek Basin, the middle North Platte River Corridor, the La Bonte Creek and Horseshoe Creek watersheds, and the North Platte River, Bolton Creek, and Bates Creek watersheds. Additional information on watersheds in the region is provided in Section 3.3.4.1. The three uranium districts within the Wyoming West Uranium Milling Region are located in the following regional watersheds: Salt Creek, Middle North Platte-Casper, Lightning Creek, Dry Fork Cheyenne River, Antelope Creek, and Upper Powder River.

The Lower Salt Creek basin located in the northeastern portion of the Wyoming West Uranium Milling Region (near the Pumpkin Buttes Uranium District) is a relatively dry basin with little vegetation. This basin includes and intermittent streams with few perennial streams. Many of the stream channels are degraded or actively degrading. Small reservoirs in the basin are dewatered for live stock and have diminished water storage capacity from sedimentation due to erosion. Native species like the Fathead minnow, flathead chub, longnose dace, plains minnow, sand shiner, and white sucker are found in this watershed (Wyoming Game and Fish Department, 2007a,b).

The La Bonte Creek and Horseshoe Creek watersheds are located in the southeastern portion of the Wyoming West Uranium Milling Region. These watersheds are subject to short periods of high water flow which contribute to the scouring of stream channels leaving wide channels which decrease during low flow periods during the summer, winter and fall seasons thus limiting habitat. Native species found in the watersheds include the brassy minnow, fathead minnow, long dace, sand shiner, longnose sucker, stonecat and plains killifish (*Fundulus kansae*). Sport fish that can be found in the systems include rainbow and Brown Trout (Wyoming Game and Fish Department, 2007a,b).

The middle North Platte River Corridor (near the Monument Hill Uranium District) is discussed for the Wyoming West Uranium Milling Region (Section 3.2.5.2).




Figure 3.3-8. Antelope Wintering Area for the Wyoming East Uranium Milling Region




Figure 3.3-9. Big Horn Wintering Area for the Wyoming East Uranium Milling Region





Figure 3.3-10. Elk Wintering Area for the Wyoming East Uranium Milling Region



SOURCE: Parturition & Crucial Wintering Area Data Provided by The Wyoming Game & Fish Department - 2005

Figure 3.3-11. Moose Wintering Area for the Wyoming East Uranium Milling Region




Figure 3.3-12. Mule Deer Wintering Area for the Wyoming East Uranium Milling Region




Figure 3.3-13. Sage-Grouse Leks Nesting Areas for the Wyoming East Uranium Milling Region




Figure 3.3-14. White Tailed Deer Wintering Area for the Wyoming East Uranium Milling Region

Description of the Affected Environment

The North Platte River, Bolton Creek, and Bates Creek watersheds are located in the southwestern portion of the Wyoming East Uranium Milling Region (in the vicinity of the Shirley Basin Uranium District). Soil erosion and sediment loading to these waterways have diminished the potential for fish to naturally reproduce. Sedimentation is further increased by erosive soils, intense grazing, road density, and poorly engineered stream crossings. Native fish within these watersheds include the big mouth shiner, brassy minnow, common shiner, creek chub, fathead minnow, longnose dace, sand shiner, stoneroller, longnose sucker, white sucker, and the plains killifish. Sports fish in the watershed include rainbow trout, cutthroat trout, brook trout, and green sunfish (Wyoming Game and Fish Department, 2007a,b).

3.3.5.3 Threatened and Endangered Species

A number of federally listed threatened and endangered species which are known to exist within habitats found within the region have been discussed previously for the Wyoming West uranium Milling Region in Section 3.2.5.3.

- Black Footed Ferret—discussed in Section 3.2.5.3
- Blowout Penstemon—discussed in Section 3.2.5.3
- Bony Tail—discussed in Section 3.2.5.3
- Canada Lynx—Section 3.2.5.3
- Colorado Butterfly Plant (*Gaura neomexicana* ssp. *Coloradensis*) —The Colorado butterfly plant typically occurs on subirrigated, stream deposited soils on level floodplains and drainage bottoms. Subpopulations are often found in low depressions or along bends in wide, active, meandering stream channels just a short distance upslope of the active channel. The plant occurs on soils derived from conglomerates, sandstones and tufaceous mudstones and siltstones of the Tertiary White River, Arikaree, and Ogallala Formations. Average annual precipitation within its range is 33–41 cm [13–16 in] primarily in the form of rainfall. The Colorado butterfly plant requires early- to mid-succession riparian habitat experiencing periodic disturbance. It commonly occurs in communities including redtop and Kentucky bluegrass on wetter sites, or wild licorice, Flodmans's thistle, curlytop gumweed, and smooth scouring rush on drier sites (U.S. Fish and Wildlife Service, 2008).
- Colorado Pikeminnow—discussed in Section 3.2.5.3.
- Humpback Chub—discussed in Section 3.2.5.3.
- Interior Least Tern—discussed in Section 3.2.5.3.
- Pallid Sturgeon—discussed in Section 3.2.5.3.
- Piping Plover—discussed in Section 3.2.5.3.
- Preble's Meadow Jumping Mouse—discussed in Section 3.2.5.3.
- Razor Sucker—discussed in Section 3.2.5.3.

- Ute Ladies's Tresses—discussed in Section 3.2.5.3.
- Western Prairie Fringed Orchid—discussed in Section 3.2.5.3.
- Whooping Crane—discussed in Section 3.2.5.3.
- Wyoming Toad (*Bufo baxteri*)—Wyoming Toad—This toad is a glacial known only from Albany County, Wyoming. It formerly inhabited flood plains, ponds, and small seepage lakes in the shortgrass communities of the Laramie Basin. The diet of this species includes ants, beetles, and a variety of other arthropods. Adults emerge from hibernation in May or June, after daytime maximum temperatures reach 70 degrees F (U.S. Fish and Wildlife Service, 2008).
- Yellow Billed Cuckoo—(candidate) discussed in Wyoming West Uranium Milling Region

State species of concern special status Wyoming Native Species Status matrix 1 (populations are greatly restricted or declining—extirpation appears possible); and 2 (populations are declining or restricted in numbers and or distribution—extirpation is not imminent); Wyoming state species of concern, which may be found in the Wyoming East Uranium Milling Region include the following:

- Kendall Warm Spring Dace (*Rhinichthys osculus thermalis*) Native Species Status 1—It resides solely in a warm spring tributary to the Green River within the Bridger-Teton National Forest. Kendall Warm Springs dace are found well distributed throughout all but the upper portion of the 300-m [984-ft] long spring creek. Kendall Warm Springs has a near constant temperature of 29 °C [85 °F]. Habitat consists of moderate to fast riffles, several man-made pools less than 1 m [3 ft] deep and shallower boggy areas. Adults are seen in the main current and pools while juveniles are seen in vegetated lateral habitats (Wyoming Game and Fish Department, 2008).
- Bluehead Sucker (*Catostomus discobolus*) Native Species Status 1—Bluehead suckers are usually found in the main current of streams, although its streamlined body form and narrow caudal peduncle indicate adaptation to living in the strong currents of larger rivers. Bluehead suckers prefer turbid to muddy streams often with high alkalinity and are rarely found in clear water (Wyoming Game and Fish Department, 2008).
- Black-footed Ferret Native Species Status 1—The black-footed ferret is found almost exclusively in prairie dog colonies in basin-prairie shrublands, sagebrush-grasslands, and grasslands. It is dependent on prairiedogs for food and all essential aspects of its habitat, especially prairie dog burrows where it spends most of its life underground (Wyoming Game and Fish Department, 2008).
- Bonneville Cutthroat (*Oncorhynchus clarki utah*) Native Species Status 2—Cutthroat trout prefer gravel-bottomed creeks and small rivers as well as lakes. The Bonneville cutthroat trout is well known for its ability to survive in harsh and often degraded (by man) habitats. In Wyoming, the Bonneville cutthroat is found in the Smith Fork and Thomas Fork drainages of the Bear River system. It is also native to some drainages in Idaho, Utah and Nevada with the bulk of its historic range within Utah (Wyoming Game and Fish Department, 2008).

Description of the Affected Environment

- Western Silvery Minnow (*Hybognathus argyritus*) Native Species Status 2—This minnow prefers large to medium sized rivers with sluggish flow and silted bottoms. They are typically found in shallow backwaters and slow pools with sand or gravel substrates. They are more abundant in clear water and show intolerance for turbidity and pollution. Western silvery minnows occur in the Belle Fourche, Little Powder, and Little Missouri rivers. They are believed to persist in the Powder River but recent surveys did not find them. They are believed extirpated from the Big Horn River. Often, it is associated with the more common plains minnow (Wyoming Game and Fish Department, 2008).
- Swift Fox (*Vulpes velox*), Native Species Status 4—The Swift Fox historically inhabited Montana and the Dakotas through the Great Plains states to northwestern Texas and eastern New Mexico. In Wyoming, it occurs primarily east of the continental divide, and is considered common in Wyoming. Habitat consists of shortgrass and mixed grass prairies, although it often uses highway and railroad right of ways, agricultural areas, and sagebrush-grasslands. Closely associated with prairie dog colonies, the Swift Fox uses underground dens year round. It selects habitat with low growing vegetation, relatively flat terrain, friable soils, and high den availability. Although expected to be stable, Wyoming classifies it as Native Species Status 4 because habitat is vulnerable though there is no ongoing significant loss of habitat (Wyoming Game and Fish Department, 2008).
- Plains Topminnow (*Fundulus sciadicus*) Native Species Status 2—The plains topminnow is considered to be of special concern in Minnesota, Missouri, Kansas, Nebraska, and Colorado. In Wyoming plains topminnows are considered rare and their distribution appears to be declining. The plains topminnow occupies habitats that are impacted by natural and anthropogenic dewatering. Introductions of western mosquito fish have been implicated in current restricted distribution of plains topminnow in Nebraska (Wyoming Game and Fish Department, 2008).
- Great Basin Gopher Snake—discussed in Section 3.2.5.3.
- Canada Lynx—discussed in Section 3.2.5.3.
- Pale Milk Snake Native Species Status 2—discussed in Section 3.2.5.3.
- Smooth Green Snake—discussed in Section 3.2.5.3.
- Yellow-billed Cuckoo—discussed in Section 3.2.5.3.
- Greater Sage Grouse—discussed in Section 3.2.5.3.
- Bald Eagle—discussed in Section 3.2.5.3.
- Trumpeter Swan—discussed in Section 3.2.5.3.
- Fringed Myotis—discussed in Section 3.2.5.3.
- Long-legged Myotis—discussed in Section 3.2.5.3.

- Pallid Bat—discussed in Section 3.2.5.3.
- Spotted Bat—discussed in Section 3.2.5.3.

3.3.6 Meteorology, Climatology, and Air Quality

3.3.6.1 Meteorology and Climatology

Wyoming's elevation results in relatively cool temperatures. Much of the temperature variations within the state can be attributed to elevation with average values dropping 1 to 2 °C [1.8 to 3.6 °F] per 300 m [1,000 ft] (National Climatic Data Center, 2005). Summer nights are normally cool although daytime temperatures may be quite high. The fall, winter, and spring can experience rapid changes with frequent variations from cold to mild periods. Freezes in early fall and late spring are typical and result in long winters and a short growing season. In the mountains and high valleys, freezes can occur any time in the summer. During winter warm spells, nighttime temperatures can remain above freezing. Valleys protected from the wind by mountain ranges can provide ideal pockets for cold air to settle and temperatures in the valley can be considerably lower than on nearby mountainsides. Table 3.3-5 identifies two climate stations located in the Wyoming East Uranium Milling Region. Climate data for these stations are found in the National Climatic Data Center's Climatology of the United States No. 20 Monthly Station Climate Summaries for 1971–2000 (National Climatic Data Center, 2004). This summary contains climate data for 4,273 stations throughout the United States and some territories. Table 3.3-6 contains temperature data for two stations in the Wyoming East Uranium Milling Region.

Precipitation within Wyoming varies with spring and early summer being the wettest time for much of the state. Mountain ranges are generally oriented in a north-south direction. This is perpendicular to the prevailing westerlies. Therefore, these mountains often act as moisture barriers. Air currents from the Pacific Ocean rise and drop much of their moisture along the western slopes of the mountains. Summer showers are frequent but typically result in rainfall amounts of a few hundredths of an inch. Usually several times a year in the state, local thunderstorms will result in 2.5 to 5 cm [1 to 2 in] of rain in a 24-hour period. On rare occasions,

Table 3.3-5. Information on Two Climate Stations in the Wyoming East Uranium Milling Region*

Station (Map Number)	County	State	Longitude	Latitude
Glenrock 5 ESE (044)	Converse	Wyoming	105°47W	42°50N
Midwest (062)	Natrona	Wyoming	106°17W	43°25N

*National Climatic Data Center. "Climatology of the United States No. 20: Monthly Station Climate Summaries, 1971–2000." Asheville, North Carolina: National Oceanic and Atmospheric Administration. 2004.

1

Table 3.3-6. Climate Data for Stations in the Wyoming East Uranium Milling Region*			
		Glenrock 5 ESE	Midwest
Temperature (°C)†	Mean—Annual	8.8	7.5
	Low—Monthly Mean	-3.1	-5.7
	High—Monthly Mean	22.4	21.5
Precipitation (cm)‡	Mean—Annual	31.0	35.0
	Low—Monthly Mean	0.90	1.4
		Glenrock 5 ESE	Midwest
	High—Monthly Mean	6.1	6.5
Snowfall (cm)	Mean—Annual	58.4	135
	Low—Monthly Mean	0	0
	High—Monthly Mean	13.5	22.6
*National Climatic Data Center. "Climatology of the United States No. 20: Monthly Station Climate Summaries, 1971–2000." Asheville, North Carolina: National Oceanic and Atmospheric Administration. 2004. †To convert Celsius (°C) to Fahrenheit (°F), multiply by 1.8 and add 32. ‡To convert centimeters (cm) to inches (in), multiply by 0.3937.			

2

3

4 rainfall in a 24-hour period can reach 7.5 to 12.5 cm [3 to 5 in] (National Climatic Data Center,
 5 2005). Heavy rains can create flash flooding in headwater streams and this flooding intensifies
 6 if these storms coincide with snow pack melting. Table 3.3-6 contains precipitation data for two
 7 stations in the Wyoming East Uranium Milling Region. The wettest month for both stations
 8 identified in Table 3.3-6 is May which, based on the snow depth data, coincides with snow pack
 9 melting (National Climatic Data Center, 2004). One of the stations is in Converse County and
 10 the other is in Natrona County. Data from the National Climatic Data Center's Storm Events
 11 Database from 1950 to 2007 indicates that the vast majority of thunderstorms in Converse and
 12 Natrona Counties occur between June and August with the most occurring in June (National
 13 Climatic Data Center, 2007).

14

15 Hailstorms are the most destructive storm event for Wyoming. Most hailstorms pass over open
 16 rangeland with minimal impact. When a hailstorm passes over a city or farmland, the property
 17 and crop damage can be severe. Most of the severe hailstorms occur in the southeast corner of
 18 the state.

19

20 Low elevations typically experience light to moderate snowfall from November to May. Snowfall
 21 within Wyoming varies by location with the mountain ranges typically receiving the most.
 22 Significant storms of 25 to 40 cm [10 to 16 in] of snow fall are infrequent outside of the
 23 mountains. Wind often coincides or follows snowstorms and can form snow drifts several
 24 meters deep. Snow can accumulate to considerable depths in the high mountains. Blizzards
 25 that last more than 2 days are uncommon. Table 3.3-6 contains snowfall data for two stations in
 26 the Wyoming East Uranium Milling Region.

27

28 Wyoming is windy and ranks first in the United States with an annual average speed of 6 m/s
 29 [12.9 mph]. During winter Wyoming frequently experiences periods where wind speed reaches
 30 13 to 18 m/s [30 to 40 mph] with gusts to 22 to 27 m/s [50 or 60 mph] (National Climatic Data
 31 Center, 2005). Prevailing wind direction varies by location but usually ranges between
 32 west-southwest through west to northwest. Since the wind is normally strong and constant from
 33 those directions, trees often lean to the east or southeast.

34

The pan evaporation rates for the Wyoming East Uranium Milling Region range from about 102 to 127 cm [40 to 50 in] (National Weather Service, 1982). Pan evaporation is a technique that measures the evaporation from a metal pan typically 121 cm [48 in] in diameter and 25 cm [10 in] tall. Pan evaporation rates can be used to estimate the evaporation rates of other bodies of water such as lakes or ponds. Pan evaporation rate data is typically available only from May to October. Freezing conditions often prevent collection of quality data during the other part of the year

3.3.6.2 Air Quality

The air quality general description for the Wyoming East Uranium Milling Region is similar to the description in Section 3.2.6 for the Wyoming West Uranium Milling Region.

As described in Section 1.7.2.2, the permitting process is the mechanism used to address air quality. If warranted, permits may set facility air pollutant emission levels, require mitigation measures, or require additional air quality analyses. Except for Indian Country, New Source Review permits in Wyoming are regulated under the EPA-approved State Implementation Plan. For Indian Country in Wyoming, the New Source Review permits are regulated under 40 CFR 52.21 (EPA, 2007a).

State Implementation Plans and permit conditions are based in part on federal regulations developed by the EPA. The NAAQS are federal standards that define acceptable ambient air concentrations for six common nonradiological air pollutants: nitrogen oxides, ozone, sulfur oxides, carbon monoxide, lead, and particulates. In June 2005, EPA revoked the 1-hour ozone standard nationwide in all locations except certain Early Action Compact Areas. None of the 1-hour ozone Early Action Compact Areas is in Wyoming. States may develop standards that are stricter or supplement the NAAQS. Wyoming has a more restrictive annual average standard for sulfur dioxide at $60 \mu\text{g}/\text{m}^3$ [1.6×10^{-6} oz/yd³] and a supplemental $50 \mu\text{g}/\text{m}^3$ [1.3×10^{-6} oz/yd³] PM₁₀ standard with an annual averaging time (Wyoming Department of Environmental Quality, 2006).

Prevention of Significant Deterioration requirements identify maximum allowable increases in concentrations for particulate matter, sulfur dioxide, and nitrogen dioxide for areas designated as attainment. Different increment levels are identified for different classes of areas and Class I areas have the most stringent requirements.

The Wyoming East Uranium Milling Region air quality description focuses on two topics: NAAQS attainment status and PSD classifications in the region.

All of the area within the Wyoming East Uranium Milling Region is classified as attainment for NAAQS. Figure 3.3-15 identifies counties in Wyoming and surrounding areas that are partially or entirely designated as nonattainment or maintenance for NAAQS at the time this GEIS was prepared (EPA, 2007b). All of the area within the Wyoming East Uranium Milling Region is classified as attainment. In fact, Wyoming only has one area that is not in attainment. The City of Sheridan in Sheridan County is designated as nonattainment for PM₁₀. Portions of several Colorado counties along the southern Wyoming border are classified as not in attainment. However, the southern boundary of the Wyoming East Uranium Milling Region is north of the Wyoming/Colorado border.




Figure 3.3-15. Air Quality Attainment Status for Wyoming and Surrounding Areas (EPA, 2007)

Table 3.3-7 identifies the Prevention of Significant Deterioration Class I areas in Wyoming. These areas are shown in Figure 3.3-16. There are no Class I areas in the Wyoming East Uranium Milling Region (40 CFR Part 81).

Table 3.3-7. U.S. Environmental Protection Agency Class I Prevention of Significant Deterioration Areas in Wyoming*

Bridger Wilderness
Fitzpatrick Wilderness
Grand Teton National Park
North Absaroka Wilderness
Teton Wilderness
Washakie Wilderness
Yellowstone National Park

*Modified from Code of Federal Regulations. "Prevention of Significant Air Deterioration of Air Quality." Title 40, Protection of the Environment, Part 81. Washington, DC: U.S. Government Printing Office. 2005.

3.3.7 Noise

The existing ambient noise levels in the undeveloped rural and more urban areas of the Wyoming East Uranium Milling Region would be 22 to 38 dB, similar to those described in Section 3.2.7 for the Wyoming West Uranium Milling Region. The largest community is Casper, the second largest city in Wyoming with a population near 50,000. Smaller communities include Glenrock and Douglas, with populations between 2,000 and about 6,000 (see Section 3.3.10). Ambient noise levels in these communities would be expected to be similar to other urban areas (up to 78 dB) (Washington State Department of Transportation, 2006).

As described in Section 3.3.2, major highways in the region include Interstate 25 and U.S. Highways 20, 26, 18, and 87. Sections of these highways are multi-lane, limited access freeways, and traffic is highest to the east (about 7,200 vehicles per day) and north (about 5,300 vehicles per day) of Casper on Interstate 25 (Wyoming Department of Transportation, 2005). Passenger cars make up about 75 percent of the traffic count on Interstate 25, indicating that ambient noise levels would likely be less than those measured at up to 70 dBA along Interstate 80 where traffic count and heavy truck traffic is higher (Federal Highway Administration, 2004; see also Section 3.2.7).

The current ISL uranium facilities (Smith Ranch-Highland, and Reynolds Ranch) and those that are anticipated for the Wyoming East Uranium Milling Region are located at least 16 km [10 mi] from the larger communities in the region. For the three uranium districts in the Wyoming East Uranium Milling Region, most of the ambient noise levels would therefore be anticipated to be similar to rural, undeveloped areas. As in the Wyoming West Uranium Milling Region, a number of small communities are located along the highways and roads that run through the region. For example, Linch, Savageton, and Sussex are located in the Pumpkin Buttes uranium district in the northwest corner of the region. In the central uranium district, the closest small communities include Orpha and Bill, and Shirley Basin is located in the uranium district in the southeast corner of the region. Noise levels in these areas would be anticipated to be higher than the undeveloped areas (22 to 38 dB), but less than the larger urban areas like Casper and Douglas.




Figure 3.3-16. Prevention of Significant Deterioration Class I Areas in the Wyoming East Uranium Milling Region and Surrounding Areas (40 CFR Part 81)

3.3.8 Historical and Cultural Resources

3.3.8.1 Cultural Resources Overview

A general overview of historical and cultural resources in Wyoming is provided in Section 3.2.8.1. As described in Section 3.2.8.1, the Wyoming SHPO administers and is responsible for oversight and compliance with the NRHP, compliance and review for Section 106 of the NHPA, and Traditional Cultural Properties review, enforcement of NAGPRA, and compliance with other federal and state historic preservation laws, regulations, and statutes.

3.3.8.2 Historic Properties Listed in the National and State Registers

Table 3.3-8 includes a summary of sites in the Wyoming East Uranium Milling Region that is listed on the Wyoming state and/or NRHP. Many of the sites are located in Casper, Glenrock, and Douglas, at least 16 km [10 mi] from potential and existing uranium ISL facilities. Several sites near Sussex in Johnson County are located near the uranium district in the northwest corner of the Wyoming East Uranium Milling Region.

Table 3.3-8. National Register Listed Properties in Counties Included in the Wyoming East Uranium Milling Region

County	Resource Name	City	Date Listed YYYY/MM/DD
Campbell	Basin Oil Field Tipi Rings (48CA1667)	Piney	1985-12-13
Campbell	Bishop Road Site (48CA1612)	Piney	1985-12-13
Campbell	Nine Mile Segment, Bozeman Trail (48CA264)	Pine Tree Junction	1989-07-23
Converse	Antelope Creek Crossing (48CO171 and 48CO165)	City Unavailable	1989-07-23
Converse	Braehead Ranch	Douglas	1995-09-07
Converse	Christ Episcopal Church and Rectory	Douglas	1980-11-17
Converse	College Inn Bar	Douglas	1979-07-10
Converse	Commerce Block	Glenrock	2005-01-21
Converse	Douglas City Hall	Douglas	1994-03-17
Converse	Fort Fetterman	Orpha	1969-04-16
Converse	Fremont, Elkhorn & Missouri Valley Railroad Passenger Depot	Douglas	1994-08-03
Converse	Glenrock Buffalo Jump	Glenrock	1969-04-16
Converse	Holdup Hollow Segment, Bozeman Trail (48CO165)	City Unavailable	1989-07-23
Converse	Hotel Higgins	Glenrock	1983-11-25
Converse	Jenne Block	Douglas	1998-01-06
Converse	La Prele Work Center	Douglas	1994-04-11
Converse	Morton Mansion	Douglas	2001-01-11
Converse	North Douglas Historic District	Douglas	2002-11-25
Converse	Officer's Club, Douglas Prisoner of War	Douglas	2001-09-08
Converse	Ross Flat Segment, Bozeman Trail (48CO165)	City Unavailable	1989-07-23
Converse	Sage Creek Station (48CO104)	Glenrock	1989-07-23
Converse	Stinking Water Gulch Segment, Bozeman Trail (48CO165)	City Unavailable	1989-07-23
Converse	U.S. Post Office—Douglas Main	Douglas	1987-05-19
Johnson	AJX Bridge over South Fork and Powder River	Kaycee	1985-02-22
Johnson	Cantonment Reno	Sussex	1977-07-29
Johnson	Dull Knife Battlefield	Barnum	1979-08-15

Table 3.3-8. National Register Listed Properties in Counties Included in the Wyoming East Uranium Milling Region (continued)

County	Resource Name	City	Date Listed YYYY/MM/DD
Johnson	EDZ Irigary Bridge	Sussex	1985-02-22
Johnson	Fort Reno	Sussex	1970-04-28
Johnson	Lake Desmet Segment, Bozeman Trail	City Unavailable	1989-07-23
Johnson	Powder River Station—Powder River Crossing (48JO134 and 48JO801)	Sussex	1989-07-23
Johnson	Sussex Post Office and Store	Kaycee	1998-11-12
Natrona	Archeological Site No. 48NA83	Arminto	1994-05-13
Natrona	Big Horn Hotel	Arminto	1978-12-18
Natrona	Bishop House	Casper	2001-03-12
Natrona	Bridger Immigrant Road—Waltman Crossing	Casper	1975-01-17
Natrona	Casper Army Air Base	Casper	2001-08-03
Natrona	Casper Buffalo Trap	Casper	1974-06-25
Natrona	Casper Federal Building	Casper	1998-12-21
Natrona	Casper Fire Department Station No. 1	Casper	1993-11-04
Natrona	Casper Motor Company—Natrona Motor Company	Casper	1994-02-23
Natrona	Church of Saint Anthony	Casper	1997-01-30
Natrona	Consolidated Royalty Building	Casper	1993-11-04
Natrona	DUX Bessemer Bend Bridge	Bessemer Bend	1985-02-22
Natrona	Elks Lodge No. 1353	Casper	1997-01-30
County	County	County	County
Natrona	Fort Casper	Casper	1971-08-12
Natrona	Fort Casper (Boundary Increase)	Casper	1976-07-19
Natrona	Independence Rock	Casper	1966-10-15
Natrona	Martin's Cove	Casper	1977-03-08
Natrona	Masonic Temple	Casper	2005-08-24
Natrona	Midwest Oil Company Hotel	Casper	1983-11-17
Natrona	Natrona County High School	Casper	1994-01-07
Natrona	North Casper Clubhouse	Casper	1994-02-18
Natrona	Ohio Oil Company Building	Casper	2001-07-25
Natrona	Pathfinder Dam	Casper	1971-08-12
Natrona	Rialto Theater	Casper	1993-02-11
Natrona	Roosevelt School	Casper	1997-01-30
Natrona	South Wolcott Street Historic District	Casper	1988-11-23
Natrona	Split Rock, Twin Peaks	Muddy Gap	1976-12-22
Natrona	Stone Ranch Stage Station	Casper	1982-11-01
Natrona	Teapot Rock	Midwest	1974-12-30
Natrona	Townsend Hotel	Casper	1983-11-25
Natrona	Tribune Building	Casper	1994-02-18

3.3.8.3 Tribal Consultation

Section 3.2.8.3 includes a discussion on Native American Tribes located within or immediately adjacent to the state of Wyoming that have interests in the state, including

- Arapaho Tribe of the Wind River Reservation
- Shoshone Tribe of the Wind River Reservation
- Cheyenne River Sioux

- Flandreau Santee Sioux
- Lower Brulé Sioux
- Oglala Sioux
- Rosebud Sioux
- Sisseton-Whapeton Oyate
- Standing Rock Sioux
- Yankton Sioux
- Crow Tribe of Montana

The Siouan tribes are located throughout South and North Dakota and the Crow are located in Montana but have interests in Wyoming. Other Siouan-speaking tribes as well as other tribes in North Dakota, Wyoming, Montana and Nebraska may have traditional land use claims in the Wyoming East Uranium Milling Region.

3.3.8.4 Places of Cultural Significance

Section 3.2.8.4 includes a more detailed discussion of culturally significant places and traditional cultural properties in Central and Eastern Wyoming. As described in Section 3.2.8, there are no known culturally significant places listed in the Wyoming East Uranium Milling Region.

3.3.9 Visual/Scenic Resources

Based on the BLM Visual Resource Handbook (BLM, 2007a–c), the uranium districts in the Wyoming East Uranium Milling Region are located at the junction of the Northern and Southern Rocky Mountain, Wyoming Basin, and Great Basin physiographic provinces (Bennett, 2003). The BLM resource management plans covering this region include the Casper (BLM, 2007d), Buffalo (BLM, 2001), Rawlins (BLM, 2008), and Newcastle (BLM, 2000) field offices (see the BLM Wyoming website at <http://www.blm.gov/wy/st/en.html>). The VRM classifications assigned within these resource plans are presented in Figure 3.3-17.

The bulk of the Wyoming East Uranium Milling Region is categorized as VRM Class III (along highways) and Class IV (open grassland, oil and natural gas, urban areas). The landscape has been extensively modified in urban areas and in several areas of oil, natural gas, and coal production, such as Natrona and Converse Counties near Casper and Douglas (Bennett, 2003; BLM, 2007d) and Johnson and Campbell Counties near Gillette (BLM, 2001). As a result, these areas are predominantly classified as VRM Class IV or as Class V/Rehabilitation. The BLM resource management plans do not identify any VRM Class I resources that fall within the Wyoming East Uranium Milling Region. VRM Class II areas are generally identified south of Interstate 25 in the region, ranging from the Laramie Mountains in the southwestern portion of the region and the North Platte River and its tributaries across the southern part of the region (BLM, 2007d, 1992). Additional areas of potentially sensitive visual resources include the Bozeman, Oregon, and Bridger historic trails that cross the southern part of the region, traveling east to west roughly parallel to the North Platte River (Bennett, 2003; BLM, 2007d, 1992) on the north side of the Laramie Mountains. All of the current and potential ISL facilities identified in the three uranium districts in the Wyoming East Uranium Milling Region are located within Class III through Class V/Rehabilitation VRM areas (Figure 3.3-17). There are no prevention of significant deterioration Class I regions or Wyoming Unique/Irreplaceable or Rare/Uncommon designated areas within the Wyoming East Uranium Milling Region (Girardin, 2006).




Figure 3.3-17. BLM Visual Resource Classifications for the Wyoming East Uranium Milling Region (BLM, 2008, 2007d, 2001, 2000)

3.3.10 Socioeconomics

For the purpose of this GEIS, the socioeconomic description for the Wyoming East Region includes communities within the region of influence for potential ISL facilities in the three uranium districts in the region. These include communities that have the highest potential for socioeconomic impacts and are considered the affected environment. Communities that have the highest potential for socioeconomic impacts are defined in the GEIS by (1) proximity to an ISL facility {generally within 48 km [30 mi]}, (2) economic profile, such as potential for income growth or destabilization, (3) employment structure, such as potential for job placement or displacement and (4) community profile, such as potential for growth or de-stabilization to local emergency services, schools, or public housing. The affected environment within the Wyoming East Uranium Milling Region consists of counties and Core-Based Statistical Areas. A Core-Based Statistical Areas, according to the U.S. Census Bureau, is a collective term for both metro and micro areas ranging from a population of 10,000 to 50,000 (U.S. Census Bureau, 2008). The major political divisions of the affected environment are listed in Table 3.3-9. The following sub-sections describe areas most likely to have implications to socioeconomics and are listed below. In some sub-sections Metropolitan Areas are also discussed. A Metropolitan Area is greater than 50,000 and a town is considered less than 10,000 in population (U.S. Census Bureau, 2008). Smaller communities such as Bill and Lynch are considered as part of the county demographics.

Table 3.3-9. Summary of Affected Environment Within the Wyoming East Uranium Milling Region

Counties Within Wyoming East	CBSAs Within Wyoming East
Albany	Casper
Campbell	
Carbon	
Converse	
Johnson	
Natrona	
Niobrara	
Platte	
Weston	

3.3.10.1 Demographics

Demographics are based on 2000 Census data population and racial characteristics of the affected environment (Table 3.3-10). (Figure 3.3-18 illustrates the populations of communities within the Wyoming East Uranium Milling Region). Most 2006 data compiled by the U.S. Census Bureau is not yet available for the geographic area of interest.

Table 3.3-10. 2000 U.S. Bureau of Census Population and Race Categories of the Wyoming East Uranium Milling Region*

Affected Environment	Total Population	White	African American	Native American	Some Other Race	Two or More Races	Asian	Hispanic Origin†	Native Hawaiian and Other Pacific Islander
Wyoming	493,782	454,670	3,722	11,133	12,301	8,883	2,771	31,669	302
Percent of total		92.1%	0.8%	2.3%	2.5%	1.8%	0.6%	6.4%	0.1%
Albany County	32,014	29,235	354	18	847	710	545	2,397	18
Percent of total		91.3%	1.1%	0.1%	2.6%	2.2%	1.7%	7.5%	0.1%
Campbell County	33,698	32,369	51	313	378	450	108	1,191	29
Percent of total		96.1%	0.2%	0.9%	1.1%	1.3%	0.3%	3.5%	0.1%
Carbon County	15,639	14,092	105	9	808	321	105	2,163	9
Percent of total		90.1%	0.7%	0.1%	5.2%	2.1%	0.7%	13.8%	0.1%
Converse County	12,052	11,416	18	110	296	177	32	660	3
Percent of total		94.7%	0.1%	0.9%	2.5%	1.5%	0.3%	5.5%	0.0%
Johnson County	7,075	6,865	6	45	39	112	8	148	0
Percent of total		97.0%	0.1%	0.6%	0.6%	1.6%	0.1%	2.1%	0.0%
Natrona County	66,533	62,644	505	686	1,275	1,121	277	3,257	25
Percent of total		94.2%	0.8%	1.0%	1.9%	1.7%	0.4%	4.9%	0.0%
Niobrara County	2,407	2,360	3	12	12	17	3	36	0
Percent of total		98.0%	0.1%	0.5%	0.5%	0.7%	0.1%	1.5%	0.0%
Platte County	8,807	8,471	14	44	149	112	15	465	2
Percent of total		96.2%	0.2%	0.5%	1.7%	1.3%	0.2%	5.3%	0.0%
Weston County	6,644	6,374	8	84	62	102	13	137	1
Percent of total		95.9%	0.1%	1.3%	0.9%	1.5%	0.2%	2.1%	0.0%
Casper	49,644	46,680	428	495	1,011	775	245	2,656	10
Percent of total		94.0%	0.9%	1.0%	2.0%	1.6%	0.5%	5.4%	0.0%

*U.S. Census Bureau. "American FactFinder." <http://factfinder.census.gov/home/saff/main.html?_lang=en> (18 October 2007 and 25 February 2008).

†Hispanic origin can be any race and is calculated as a separate component of the total population (i.e., if added to the other races would total more than 100 percent).




Figure 3.3-18. Wyoming East Uranium Milling Region With Population

The most populated county in the Wyoming East Uranium Milling Region is Natrona County and the most sparsely populated county is Niobrara County. The county with the largest percentage of non-minorities is Niobrara County with a white population of 98.0 percent. The largest minority based county is Carbon County with a white population of 90.1 percent or a minority-based population of 9.9 percent. The Core-Based Statistical Areas of Casper is demographically similar to the counties within the Wyoming East Uranium Milling Region.

3.3.10.2 Income

Income information from the 2000 Census including labor force, income, and poverty levels for the affected environment is based on data collected from state and county levels. Data collected at the state level also includes information on towns, Core-Based Statistical Areas, or Metropolitan Areas and was done to take into consideration an outside workforce. An outside workforce may be a workforce willing to commute long distances {greater than 48 km [30 mi]} for income opportunities or may be a workforce necessary to fulfill specialized positions (if local workforce is unavailable or un-specialized). In Wyoming, the workforce frequently commutes long distances to work. For example, in the Wyoming East Uranium Milling Region, most of the affected counties experienced net inflows of workers during the 4th Quarter of 2005. Net inflows ranged from about 160 for Johnson County to about 7,500 for Campbell County. These inflows were predominately for jobs related to the energy industry in the Powder River Basin (Wyoming Workforce Development Council, 2007). Converse (-1,063) and Platte (-228) Counties experienced net outflows during the same period. Data collected at the county level is generally the same as the affected environment presented in Table 3.3-9. State level information for the surrounding region is provided in Table 3.3-11 and county data is listed in Table 3.3-12.

For the surrounding region, the state with both the largest labor force population and families and individuals living below poverty level is Colorado. The largest labor force population is Billings, Montana {128 km [80 mi] from the nearest potential ISL facility in the region} and the smallest labor force population is Laramie, Wyoming { 96 km [60 mi] from the nearest potential ISL facility}. The population with the highest per capita income is Fort Collins, Colorado {240 km [150 mi] from the nearest potential ISL facility) and the lowest per capita income population is Laramie, Wyoming. The population with the highest percentage of individuals and families below poverty levels is Laramie, Wyoming (Table 3.3-11).

The county with the largest labor force is Natrona County and the smallest labor force is located in Niobrara County. The county with the highest per capita income is Campbell County and the smallest per capita income at the county level is Niobrara County. The county with the highest percentage of individuals and families living below the poverty level is Albany County (Table 3.3-12).

1

Table 3.3-11. U.S. Bureau of Census State Income Information for Wyoming East Uranium Milling Region*

Affected Environment	2000 Labor Force Population (16 years and over)	Median Household Income in 1999	Median Family Income in 1999	Per Capita Income in 1999	Families Below Poverty Level in 2000	Individuals Below Poverty Level in 2000
Colorado	2,331,898	\$47,203	\$55,883	\$24,049	67,614	388,952
South Dakota	394,945	\$35,282	\$42,237	\$17,562	18,172	95,900
Wyoming	257,808	\$37,892	\$45,685	\$19,134	10,585	54,777
Casper	26,343	\$36,567	\$46,267	\$19,409	1,122	5,546
<i>Percent of total†</i>	68.4%	NA‡	NA‡	NA‡	8.5%	11.4%
Cheyenne, Wyoming	27,647	\$38,856	\$46,771	\$19,809	891	4,541
<i>Percent of total†</i>	66.7%	NA‡	NA‡	NA‡	6.3%	8.8%
Ft. Collins, Colorado	69,424	\$44,459	\$59,332	\$22,133	1,417	15,835
<i>Percent of total†</i>	72.4%	NA‡	NA‡	NA‡	5.5%	14.0%
Laramie, Wyoming	15,504	\$27,319	\$43,395	\$16,036	633	5,618
<i>Percent of total†</i>	67.2%	NA‡	NA‡	NA‡	11.1%	22.6%
Rapid City, South Dakota	31,948	\$35,978	\$44,818	\$19,445	1,441	7,328
<i>Percent of total†</i>	68.8%	NA‡	NA‡	NA‡	9.4%	12.7%
* U.S. Census Bureau. "American FactFinder." < http://factfinder.census.gov/home/saff/main.html?_lang=en > (18 October 2007, 25 February 2008, and 15 April 2008). †Percent of total based on a population of 16 years and over. ‡NA—Not applicable.						

2

Table 3.3-12. U.S. Bureau of Census County Income Information for Wyoming East Uranium Milling Region*

Affected Environment	2000 Labor Force Population (16 years and over)	Median Household Income in 1999	Median Family Income in 1999	Per Capita Income in 1999	Families Below Poverty Level in 2000	Individuals Below Poverty Level in 2000
Albany County, Wyoming	18,182	\$28,790	\$44,334	\$16,706	763	6,228
Percent of total†	67.7%	NA‡	NA‡	NA‡	10.8%	21.0%
Campbell County, Wyoming	18,805	\$49,536	\$53,92	\$20,063	507	2,544
Percent of total†	76.6%	NA‡	NA‡	NA‡	5.6%	7.6%
Carbon County, Wyoming	7,744	\$36,060	\$41,991	\$18,375	411	1,879
Percent of total†	62.5%	NA‡	NA‡	NA‡	9.8%	12.9%
Converse County, Wyoming	6,244	\$39,603	\$45,905	\$18,744	319	1,379
Percent of total†	68.6%	NA‡	NA‡	NA‡	9.2%	11.6%
Johnson County, Wyoming	3,472	\$34,012	\$42,299	\$19,030	147	712
Percent of total†	61.7%	NA‡	NA‡	NA‡	7.2%	10.1%
Natrona County, Wyoming	35,081	\$36,619	\$45,575	\$18,913	1,548	7,695
Percent of total†	68.3%	NA‡	NA‡	NA‡	8.7%	11.8%
Niobrara County, Wyoming	1,193	\$29,701	\$33,714	\$15,757	74	309
Percent of total†	61.5%	NA‡	NA‡	NA‡	10.7%	13.4%
Platte County, Wyoming	4,540	\$33,866	\$41,449	\$17,530	216	1,021
Percent of total†	66.1%	NA‡	NA‡	NA‡	8.5%	11.7%
Weston County	3,183	\$32,348	\$40,472	\$17,366	119	628
Percent of total†	60.0%	NA‡	NA‡	NA‡	6.3%	9.9%

* U.S. Census Bureau. "American FactFinder." <http://factfinder.census.gov/home/saff/main.html?_lang=en> (18 October 2007 and 25 February 2008).
†Percent of total based on a population of 16 years and over.
‡NA—Not applicable.

3.3.10.3 Housing

Housing information based on 2000 Census data is provided in Table 3.3-13.

The availability of housing within the immediate vicinity of potential ISL facilities in the Wyoming East Uranium Milling Region is limited. The majority of housing is available in larger populated areas such as the towns of Casper {48 km [30 mil] to the nearest potential ISL facility} and Riverton {193 km [120 mil] to the nearest potential ISL facility}. Temporary housing such as apartments, lodging, and trailer camps within the immediate vicinity of the proposed ISL facilities is not as limited. There are 17 apartment complexes available in larger populated areas such as the Core-Based Statistical Areas or towns of Casper, Douglas, Lusk, and Orpha (MapQuest, 2008). There are also 15 hotels/motels along major highways or towns near the uranium districts located within the Wyoming East Uranium Milling Regions. In addition to apartments and lodging, there are more than 25 trailer camps situated along major roads or near towns (MapQuest, 2008).

Table 3.3-13. U.S. Bureau of Census Housing Information for the Wyoming East Uranium Milling Region*

Affected Environment	Single Family Owner-Occupied Homes	Median Value in Dollars	Median Monthly Costs With a Mortgage	Median Monthly Costs Without a Mortgage	Occupied Housing Units	Renter-Occupied Units
Wyoming	95,591	\$96,600	\$825	\$229	193,608	55,793
Albany County	4,987	\$118,600	\$916	\$225	13,269	6,345
Campbell County	5,344	\$102,900	\$879	\$247	12,207	3,174
Carbon County	7,744	\$76,500	\$685	\$196	6,129	1,708
Converse County	2,290	\$84,900	\$714	\$206	4,694	1,142
Johnson County	1,414	\$115,500	\$849	\$227	2,959	677
Natrona County	15,250	\$84,600	\$746	\$218	26,819	7,993
Niobrara County	480	\$60,300	\$562	\$200	1,011	222
Platte County	1,659	\$84,100	\$698	\$205	3,625	800
Weston County	1,174	\$66,700	\$664	\$199	2,624	549
Casper	12,642	\$84,500	\$744	\$220	20,437	6,645
Source: U.S. Census Bureau. "American FactFinder." < http://factfinder.census.gov/home/saff/main.html?_lang=en > (18 October 2007 and 25 February 2008).						

3.3.10.4 Employment Structure

Employment structure from the 2000 Census, including employment rate and type is based on data collected at the state and county levels. Data collected from the state level also includes

Description of the Affected Environment

information on towns, Core-Based Statistical Areas, or metropolitan areas and was done to take into consideration an outside workforce. An outside workforce may include workers willing to commute long distances {greater than 48 km [30 mil]} for employment opportunities or external labor necessary to fulfill specialized positions (if local workforce is unavailable or unspecialized). Data collected at the county level is generally the same as the affected environment presented in Table 3.3-9.

Based on review of regional state level information, Colorado has the highest percentage of employment.

At the county level, the county in the Wyoming East Uranium Milling Region with the highest percentage of employment is Campbell County and the county with the highest unemployment rate is Albany County.

3.3.10.4.1 State Data

3.3.10.4.1.1 Colorado

The State of Colorado has an employment rate of 66.3 percent and unemployment rate of 3.0 percent. The largest sector of employment is management, professional, and related occupations at 37.4 percent. The largest type of industry is educational, health, and social services. The largest class of worker is private wage and salary workers (U.S. Census Bureau, 2008).

Ft. Collins

Ft. Collins has an employment rate of 68.5 percent and unemployment higher than the state at 3.8 percent. The largest sector of employment is management, professional, and related occupations at 42.9 percent. The largest type of industry is educational, health, and social services. The largest class of worker is private wage and salary workers (U.S. Census Bureau, 2008).

3.3.10.4.1.2 South Dakota

The State of South Dakota has an employment rate of 64.9 percent and unemployment rate of 3.0 percent. The largest sector of employment is management, professional, and related occupations at 32.6 percent. The largest type of industry is educational, health, and social services. The largest class of worker is private wage and salary workers (U.S. Census Bureau, 2008).

Rapid City

Laramie has an employment rate of 63.7 percent and unemployment higher than the state at 3.2 percent. The largest sector of employment is management, professional, and related occupations at 32.8 percent. The largest type of industry is educational, health, and social services. The largest class of worker is private wage and salary workers (U.S. Census Bureau, 2008).

3.3.10.4.1.3 Wyoming

The State of Wyoming has an employment rate of 63.1 percent and unemployment rate of 3.5 percent. The largest sector of employment is sales and office occupations. The largest type of industry is educational, health, and social services. The largest class of worker is private wage and salary workers (U.S. Census Bureau, 2008).

Casper

Casper has an employment rate of 64.9 percent and an unemployment rate lower than that of the state at 3.4 percent. The largest sector of employment is sales and office occupations at 30.6 percent followed by management, professional, and related occupations at 29.7 percent. The largest type of industry is educational, health, and social services at 22.1 percent. The largest class of worker is private wage and salary workers at 76.6 percent (U.S. Census Bureau, 2008).

Cheyenne

Cheyenne has an employment rate of 59.2 percent and unemployment less than the state at 3.3 percent. The largest sector of employment is management, professional, and related occupations at 33.0 percent. The largest type of industry is educational, health, and social services. The largest class of worker is private wage and salary workers (U.S. Census Bureau, 2008).

Laramie

Laramie has an employment rate of 63.4 percent and unemployment less than the state at 3.7 percent. The largest sector of employment is management, professional, and related occupations at 40.5 percent. The largest type of industry is educational, health, and social services. The largest class of worker is private wage and salary workers (U.S. Census Bureau, 2008).

3.3.10.4.2 County Data

Albany County, Wyoming

Albany County has an employment rate of 63.9 percent and an unemployment rate higher than that of the state at 3.7 percent. The largest sector of employment is management, professional, and related occupations at 40.4 percent. The largest type of industry is educational, health, and social services at 37.1 percent. The largest class of worker is private wage and salary workers at 61.9 percent (U.S. Census Bureau, 2008).

Campbell County, Wyoming

Campbell County has an employment rate of 73.2 percent and an unemployment rate lower than that of the state at 3.4 percent. The largest sector of employment is management, professional, and related occupations at 23.9 percent followed by construction, extraction, and maintenance occupations at 23.7 percent. The largest type of industry is agriculture, forestry, fishing and hunting, and mining at 23.3 percent followed by educational, health, and social

Description of the Affected Environment

services at 16.7 percent. The largest class of worker is private wage and salary workers at 78.4 percent (U.S. Census Bureau, 2008).

Carbon County, Wyoming

Carbon County has an employment rate of 59.2 percent and an unemployment rate lower than that of the state at 3.3 percent. The largest sector of employment is management, professional, and related occupations at 23.4 percent followed by sales and office occupations at 21.9 percent. The largest type of industry is educational, health, and social services at 17.1 percent. The largest class of worker is private wage and salary workers at 65.6 percent (U.S. Census Bureau, 2008).

Converse County, Wyoming

Converse County has an employment rate of 65.4 percent and an unemployment rate lower than that of the state at 3.2 percent. The largest sector of employment is management, professional, and related occupations at 23.2 percent followed by sales and office occupations at 21.4 percent. The largest type of industry is agriculture, forestry, fishing and hunting, and mining at 20.1 percent followed by educational, health, and social services at 18.5 percent. The largest class of worker is private wage and salary workers at 71.1 percent (U.S. Census Bureau, 2008).

Johnson County, Wyoming

Johnson County has an employment rate of 57.6 percent and an unemployment rate slightly higher than that of the state at 3.7 percent. The largest sector of employment is management, professional, and related occupations at 37.5 percent followed by sales and office occupations at 20.3 percent. The largest type of industry is educational, health, and social services at 20.5 percent followed by agriculture, forestry, fishing and hunting, and mining at 19.5 percent. The largest class of worker is private wage and salary workers at 61.1 percent (U.S. Census Bureau, 2008).

Natrona County, Wyoming

Natrona County has an employment rate of 64.6 percent and an unemployment rate similar to that of the state at 3.5 percent. The largest sector of employment is sales and office occupations at 29.9 percent followed by management, professional, and related occupations at 28.5 percent. The largest type of industry is educational, health, and social services at 21.2 percent. The largest class of worker is private wage and salary workers at 76.2 percent (U.S. Census Bureau, 2008).

Niobrara County, Wyoming

Niobrara County has an employment rate of 59.4 percent and an unemployment rate lower than that of the state at 2.1 percent. The largest sector of employment is management, professional, and related occupations at 34.4 percent. The largest type of industry is agriculture, forestry, fishing and hunting, and mining at 24.7 percent. The largest class of worker is private wage and salary workers at 62.6 percent (U.S. Census Bureau, 2008).

Platte County, Wyoming

Platte County has an employment rate of 63.1 percent and an unemployment rate lower than that of the state at 2.9 percent. The largest sector of employment is management, professional, and related occupations at 30.3 percent. The largest type of industry is educational, health, and social services at 21.4 percent. The largest class of worker is private wage and salary workers at 64.4 percent (U.S. Census Bureau, 2008).

Weston County, Wyoming

Weston County has an employment rate of 56.6 percent and an unemployment rate lower than that of the state at 3.3 percent. The largest sector of employment is management, professional, and related occupations at 24.3 percent. The largest type of industry is agriculture, forestry, fishing and hunting, and mining at 22.4 percent. The largest class of worker is private wage and salary workers at 68.9 percent (U.S. Census Bureau, 2008).

3.3.10.5 Local Finance

Local finance such as revenue and tax information for the affected environment is provided below and in Table 3.3-14.

Local finance such as revenue and tax distribution information for the affected counties is presented in Table 3.3-14.

Wyoming

The State of Wyoming does not have an income tax nor does it assess tax on retirement income received from another state. Wyoming has a 4 percent state sales tax, 2 percent to 5 percent county lodging tax, and 5 percent use tax. Counties have the option of collecting an additional 1 percent tax for general revenue and 2 percent tax for specific purposes. Wyoming also imposes "ad valorem taxes" on mineral extraction properties. Taxes levied for uranium production was 4.0 percent in 2007 and totaled \$17 million dollars. The majority of tax revenue came from Converse County with a small amount (\$7,159) from Sweetwater County (Wyoming Department of Revenue, 2007). Sales and use tax distribution information for the affected counties is presented in Table 3.3-14.

Table 3.3-14. 2007 Sales and Use Tax Distribution of the Affected Counties Within the Wyoming East Uranium Milling Region*

Affected Counties	Use Tax		Sales Tax		Lodging Option Tax
	General	Specific	General	Specific	
Albany County	\$35,223.87	\$35,223.87	\$427,731.38	\$427,731.38	\$75,599.10
Campbell County	\$387,522.93	\$97,111.27	\$2,334,282.49	\$583,201.87	\$0.0
Carbon County	\$8,546.95	\$64,236.31	\$465,469.37	\$47,391.45	\$40,974.56

Table 3.3-14. 2007 Sales and Use Tax Distribution of the Affected Counties Within the Wyoming East Uranium Milling Region* (continued)

Affected Counties	Use Tax		Sales Tax		Lodging Option Tax
	General	Specific	General	Specific	
Converse County	\$46,192.16	\$0.0	\$236,705.84	\$0.0	\$18,649.94
Johnson County	\$23,318.00	\$0.0	\$246,961.51	\$0.0	\$28,700.89
Natrona County	\$132,453.29	\$0.0	\$1,572,768.04	\$0.0	\$98,624.31
Niobrara County	\$6,119.06	\$34,411.65	\$6,119.06	\$34,411.65	\$5,137.77
Platte County	\$26,652.78	\$0.0	\$103,473.55	\$0.0	\$703.15
Weston County	\$28,152.44	\$0.0	\$60,466.76	\$0.0	\$6,682.25
* Wyoming Department of Revenue. "Sales and Tax Distribution Report by County 2007." < http://revenue.state.wy.us/Portal/VBVS/DesktopDefault.aspx?tabindex=3&tabid=10 > (18 October 2007 and 25 February 2008).					

Casper

Sources of revenue for Casper, the largest city in the Wyoming East Uranium Milling Region, include sales, use, lodging, and property taxes as well as mill levies. The sales and use tax rate is 5 percent and lodging is 3 percent. The largest distribution of property tax is school district tax at a rate of 32.5 percent (Casper Chamber of Commerce, 2007).

Campbell County

Campbell County has 1 school district with 24 schools consisting of 15 elementary schools, 2 junior high schools, 1 junior/senior high school, 1 high school, 1 alternative school, and 1 aquatic center. There are a total of approximately 7,441 students. The majority of schools provide bus services (Campbell County School District No. 1, 2007).

Carbon County

Carbon County has two school districts, Carbon County School District #1 and #2, with a combined total of approximately 2,647 students. There are a total of 9 elementary schools, 2 middle school, 2 high school, and 2 private schools. The majority of schools within each school district provide bus services (Carbon County School District No.1 and No. 2, 2008a,b).

Converse County

Converse County has two school districts, Converse County School Districts No. 1 and No. 2, with a total of approximately 2,455 students. There are a total of 9 elementary schools, 4 middle/intermediate schools, and 2 high schools. The majority of schools within each school district provide bus services (Schoolbug.org, 2007b).

Johnson County

Johnson County has one school district with two elementary schools, one middle school, two high schools, and one learning center. There are a total of approximately 1,257 students. The majority of schools provide bus services (Johnson County School District No. 1, 2007).

Natrona County

Natrona County has one school district, Natrona County School District No. 1, with a total of approximately 11,500 students. There are more than 30 public and private elementary and secondary schools. The majority of schools provide bus services (Natrona County School District No. 1, 2007).

Niobrara County

Niobrara County has one school district, Niobrara County School District No. 1, with a total of approximately 422 students. There are 1 elementary and middle schools, 1 high school, and 1 private school. Information as to whether these schools provide bus services is not available (Niobrara County School District No. 1, 2008).

Platte County

Platte County has the Platte County School District No. 1, with a total of approximately 1,571 students. There are 2 elementary schools, 1 middle school, 1 high school, and 2 private or parochial schools. Information as to whether these schools provide bus services is not available (Platte County School District No.1, 2008).

Weston County

Weston County has one school district, Weston County School District No. 1, with a total of approximately 1,134 students. There are 2 elementary schools, 1 middle school, and 1 high school. Information as to whether these schools provide bus services is not available (Weston County School District No. 1, 2008).

3.3.10.6 Education

Information on education for the affected communities within the region of influence is presented next.

Based on review of the affected environment, the county with the largest number of schools is Natrona County and the county with the smallest number of schools is Niobrara County. The Core-Based Statistical Area of Casper was average to the county level when compared to the aforementioned schools.

Casper

Casper has one school district, Natrona County School District No. 1, with a total of approximately 11,500 students. There are more than 25 public and private elementary, middle, and high schools. The majority of schools provide bus services (Schoolbug.org, 2007a).

Description of the Affected Environment

Albany County

Albany County has one school district, Albany County School District No. 1, with a total of approximately 3,790 students. There are 13 elementary schools, 6 middle schools, and 3 high schools. The majority of schools provide bus services (Greatschools.com, 2008).

Campbell County

Campbell County has 1 school district with 24 schools consisting of 15 elementary schools, 2 junior high schools, 1 junior/senior high school, 1 high school, 1 alternative school, and 1 aquatic center. There are a total of approximately 7,441 students. The majority of schools provide bus services (Campbell County School District No. 1, 2007).

Carbon County

Carbon County has two school districts, Carbon County School District #1 and #2, with a combined total of approximately 2,647 students. There are a total of 9 elementary schools, 2 middle school, 2 high school, and 2 private schools. The majority of schools within each school district provide bus services (Carbon County School District No.1 and No. 2, 2008a,b).

Converse County

Converse County has two school districts, Converse County School Districts No. 1 and No. 2, with a total of approximately 2,455 students. There are a total of 9 elementary schools, 4 middle/intermediate schools, and 2 high schools. The majority of schools within each school district provide bus services (Schoolbug.org, 2007b).

Johnson County

Johnson County has one school district with two elementary schools, one middle school, two high schools, and one learning center. There are a total of approximately 1,257 students. The majority of schools provide bus services (Johnson County School District No. 1, 2007).

Natrona County

Natrona County has one school district, Natrona County School District No. 1, with a total of approximately 11,500 students. There are more than 30 public and private elementary and secondary schools. The majority of schools provide bus services (Natrona County School District No. 1, 2007).

Niobrara County

Niobrara County has one school district, Niobrara County School District No. 1, with a total of approximately 422 students. There are 1 elementary and middle schools, 1 high school, and 1 private school. Information as to whether these schools provide bus services is not available (Niobrara County School District No. 1, 2008).

Platte County

Platte County has the Platte County School District No. 1, with a total of approximately 1,571 students. There are 2 elementary schools, 1 middle school, 1 high school, and 2 private or parochial schools. Information as to whether these schools provide bus services is not available (Platte County School District No.1, 2008).

Weston County

Weston County has one school district, Weston County School District No. 1, with a total of approximately 1,134 students. There are 2 elementary schools, 1 middle school, and 1 high school. Information as to whether these schools provide bus services is not available (Weston County School District No. 1, 2008).

3.3.10.7 Health and Social Services

Health Care

The majority of the health care facilities that provide service in the vicinity of the Wyoming East Uranium Milling Region are located within populated areas of the affected environment. The closest health care facilities within the vicinity of the ISL facilities are located in Riverton, Lander, Casper, Douglas, Wheatland, Cheyenne, and Laramie and have a total of 15 facilities (MapQuest, 2008). These consist of hospitals, clinics, emergency centers, and medical services. The following hospitals are located proximate to the Wyoming East Milling Region: Riverton (1), Cheyenne (1), Laramie (1), and Wheatland (1).

Local Emergency

Local police within the Wyoming East Uranium Milling Region is under the jurisdiction of each county. There are 28 police, sheriff, or marshals offices within the region: Albany County (2), Campbell County (2), Carbon County (6), Converse County (3), Johnson County (3), Natrona County (4), Niobrara County (2), Platte County (3), and Weston County (3) (USACops, 2008b).

Fire departments within the Wyoming East Uranium Milling Region comprised at the county, town, Core-Based Statistical Areas, or city level. There are 7 fire departments within the milling region: Campbell County (1), Casper (1), Douglas (2), Lusk (1), Natrona County (1), and Wheatland (1) (50states, 2008b).

3.3.11 Public and Occupational Health

3.3.11.1 Background Radiological Conditions

For a U.S. resident, the average total effective dose equivalent from natural background radiation sources is approximately 3 mSv/yr [300 mrem/yr] but varies by location and elevation (National Council of Radiation Protection and Measurements, 1987). In addition, the average American receives 0.6 mSv/yr [60 mrem/yr] from man-made sources including medical diagnostic tests and consumer products (National Council of Radiation Protection and Measurements, 1987). Therefore the total from natural background and man-made sources for the average U.S. resident is 3.6 mSv/yr [360 mrem/yr]. For a breakdown of the sources of this radiation, see Figure 3.2-22.

Description of the Affected Environment

Background dose varies by location primarily because of elevation changes and variations in the dose from radon. As elevation increases so does the dose from cosmic radiation and hence the total dose. Radon is a radioactive gas produced from the decay of ^{238}U , which is naturally found in soil. The amount of radon in the soil/bedrock depends on the type the porosity and moisture content. Areas which have types of soils/bedrock like granite and limestone have higher radon levels than those with other types of soils/bedrock (EPA, 2006).

For the Wyoming East region, the average background radiation dose for the state of Wyoming is used which is 3.16 mSv/yr [316 mrem/yr] (EPA, 2006). This value includes natural and manmade sources. This dose is slightly lower than the U.S. average primarily because the radon dose is lower (U.S. average of 2 mSv/yr [200 mrem/yr] versus Wyoming average of 1.33 mSv/yr [133 mrem/yr]). The cosmic dose is slightly higher than the U.S. average: 0.515 mSv/yr [51.5 mrem/yr] versus 0.27 mSv/yr [27 mrem/yr]. The remaining contributions from terrestrial, internal, and manmade radiation combined are the same as the U.S. average of 1.318 mSv/yr [131.8 mrem/yr].

3.3.11.2 Public Health and Safety

Public health and safety standards are the same regardless of a facility's location. See Section 3.2.11.2 for further discussion of these standards.

3.3.11.3 Occupational Health and Safety

Occupational health and safety standards are the same regardless of facility's location. See Section 3.2.11.3 for further discussion of these standards.

3.3.12 References

50States. "Wyoming." <www.50states.com> (15 April 2008).

Anderson, D.C. "Uranium Deposits of the Gas Hills." *Contributions to Geology, Wyoming Uranium Issue*. Laramie, Wyoming: University of Wyoming. Vol. 8, No. 2.1. pp. 93–104. 1969.

Bailey, R.V. "Uranium Deposits in the Great Divide Basin-Crooks Gap Area, Fremont and Sweetwater Counties, Wyoming." *Contributions to Geology, Wyoming Uranium Issue*. Laramie, Wyoming: University of Wyoming. Vol. 8, No. 2.1. pp. 105–120. 1969.

Bennett, E. "The Visual Resource Inventory for the Casper Field Office." Casper, Wyoming: BLM, Casper Field Office. March 2003.

BLM. "Proposed Resource Management Plan and Final Environmental Impact Statement for Public Lands Administered by the Bureau of Land Management Rawlins Field Office." Rawlins, Wyoming: BLM, Rawlins Field Office. January 2008. <<http://www.blm.gov/rmp/wy/rawlins/>> (03 March 2008).

BLM. "Visual Resource Management." Manual 8400. Washington, DC: BLM. 2007a. <<http://www.blm.gov/nstc/VRM/8400.html#Anchor-.06-23240>> (17 October 2007).

- BLM. "Visual Resource Inventory." Manual H-8410-1. Washington, DC: BLM. 2007b. <<http://www.blm.gov/nstc/VRM/8410.html>> (17 October 2007).
- BLM. "Visual Resource Contrast Rating." Manual 8431. Washington, DC: BLM. 2007c. <<http://www.blm.gov/nstc/VRM/8431.html>> (17 October 2007).
- BLM. "Proposed Resource Management Plan and Final Environmental Impact Statement for the Casper Field Office Planning Area." BLM/WY/PL-07/017+1610. Casper, Wyoming: BLM, Casper Field Office. June 2007d. <<http://www.blm.gov/rmp/casper/PRMP-FEIS.htm>> (17 October 2007).
- BLM. "Buffalo Resource Management Plan." Buffalo, Wyoming: BLM, Buffalo Field Office. April 2001. <<http://www.blm.gov/rmp/WY/application/index.cfm?rmpid=101>> (17 October 2007).
- BLM. "Newcastle Resource Management Plan." Newcastle, Wyoming: BLM, Newcastle Field Office. 2000. <http://www.blm.gov/rmp/WY/application/rmp_browse.cfm?hlevel=1&rmpid=89&idref=28910> (17 October 2007).
- BLM. "Nebraska Record of Decision and Approved Resource Management Plan." Casper, Wyoming: BLM, Casper District Office. 1992. <<http://www.blm.gov/rmp/WY/Nebraska/rmp.pdf>> (17 October 2007).
- Boberg, W.W. "Some Speculations on the Development of Central Wyoming as a Uranium Province." Wyoming Geological Association 32nd Annual Field Conference Guidebook. Casper, Wyoming: Wyoming Geological Association. pp. 161-180. 1981.
- Campbell County School District No. 1. "Campbell County School District No. 1, Educational Services Center." Gillette, Wyoming. 2007. <www.esc.ccsd.k12.wy.us> (15 October 2007).
- Carbon County School District No.1. "Carbon County School District One." Rawlins, Wyoming: Carbon County School District. 2008. <www.crb1.k12.wy.us/#> (27 February 2008).
- Carbon County School District No. 2. "Carbon County School District #2, Carbon County, Wyoming." Saratoga, Wyoming: Carbon County School District #2. 2008. <www.crb2.k12.wy.us> (27 February 2008).
- Casper Chamber of Commerce. *Community Profile*. Casper, Wyoming: Casper Chamber of Commerce. 2006.
- Chapman, S.S., S.A. Bryce, J.M. Omernik, D.G. Despain, J. ZumBerge, and M. Conrad. "Ecoregions of Wyoming." U.S. Geological Survey Map. Scale 1:1,400,000. 2004.
- Chenoweth, W.L. "A Summary of Uranium Production in Wyoming." Mineral Resources of Wyoming, Wyoming Geological Association, 42nd Annual Field Conference Guidebook. Casper, Wyoming: Wyoming Geological Association. pp. 169-179. 1991.
- Curtis J. and Grimes K. (2004). Wyoming Climate Atlas, <<http://www.wrds.uwyo.edu/wrds/wsc/climateatlas/>> (29 April 2008).

Description of the Affected Environment

- 1 Davis, J.F. "Uranium Deposits of the Powder River Basin." Contributions to Geology, Wyoming
2 Uranium Issue. Laramie, Wyoming: University of Wyoming. Vol. 8, No. 2.1. pp. 131–142.
3 1969.
4
- 5 Driscoll, F.G. (1986). Groundwater and Wells. Second edition. Johnson Filtration Systems Inc.,
6 St. Paul, Minnesota. pp. 1,089.
7
- 8 EPA. "National Assessment Database." 2008. <<http://www.epa.gov/waters/305b/index.html>>
9 (28 February 2008).
10
- 11 EPA. "Counties Designate Nonattainment or Maintenance for Clean Air Act's National Ambient
12 Air Quality Standards (NAAQS)." 2007a. <[http://www.epa.gov/oar/oaqps/greenbk/](http://www.epa.gov/oar/oaqps/greenbk/mapnmpoll.html)
13 [mapnmpoll.html](http://www.epa.gov/oar/oaqps/greenbk/mapnmpoll.html)> (29 September 2007).
14
- 15 EPA. "Prevention of Significant Deterioration (PSD) Permit Program Status: May 2007."
16 2007b. <<http://www.epa.gov/nsr/where.html>> (26 September 2007).
17
- 18 EPA. "Assessment of Variations in Radiation Exposure in the United States (Revision 1)."
19 Contract Number EP–D–05–02. Washington, DC: EPA. 2006.
20
- 21 Federal Highway Administration. "Synthesis of Noise Effects on Wildlife Populations."
22 FHWA–HEP–06–016. Washington, DC: Federal Highway Administration, Department of
23 Transportation. 2004.
- 24 Fort Phil Kearny/Bozeman Trail Association. "The Bozeman Trail Corridor." Sheridan,
25 Wyoming: The Fort Phil Kearny/Bozeman Trail Association. 2008.
26 <<http://www.bozemantrail.org/btcorridor-map.html>> (27 February 2008).
27
- 28 Girardin, J. "A List of Areas Designated Unique and Irreplaceable or Designated Rare or
29 Uncommon by the Council." Letter From J. Girardin to T. Lorenzon. Cheyenne, Wyoming:
30 Council on Environmental Quality. November 29, 2006.
31
- 32 Greatschools.com. "Albany County." 2008. <www.greatschools.com> (27 February 2008)
33
- 34 Harris, R.E. and J.K. King. "Geological Classification and Origin of Radioactive Mineralization in
35 Wyoming." A.W. Snoke, J.R. Steidtmann, and S.M Roberts, eds. Geology of Wyoming:
36 Geological Survey of Wyoming Memoir No. 5. pp. 898–916. 1983.
37
- 38 Harshman, E.N. "Uranium Deposits of Wyoming and South Dakota." *Ore Deposits in the*
39 *United States 1933–1967*. New York City, New York: American Institute of Mining,
40 Metallurgical, and Petroleum Engineers. pp. 815–831. 1968.
41
- 42 Houston, R.S. "Aspects of the Geologic History of Wyoming Related to the Formation of
43 Uranium Deposits." Contributions to Geology, Wyoming Uranium Issue. Laramie, Wyoming:
44 University of Wyoming. Vol. 8, No. 2.1. pp. 67–79. 1969.
45
- 46 Johnson County School District No. 1. 2007. <www.natronaschools.org> (15 October 2007).
47
- 48 Kansas Geological Survey (1991). Open-File Report. 91-1
49 <<http://www.kgs.ku.edu/Dakota/vol3/fy91/rep06.htm>> (29 April 2008).
50

- 1 Lageson, D. and D. Spearing. *Roadside Geology of Wyoming*. Missoula, Montana: Mountain
2 Press Publishing Company. 1988.
3
- 4 Langden, R.E. "Geology and Geochemistry of the Highland Uranium Deposit." Wyoming
5 Geological Association Earth Science Bulletin. pp. 41–48. 1973. ^a
6
- 7 Love, J.D. "Preliminary Report on Uranium Deposits in the Pumpkin Buttes Area, Powder River
8 Basin, Wyoming." U.S. Geological Survey Circular 176. 1952.
9
- 10 MapQuest. "Wyoming." <www.mapquest.com> (15 April 2008).
11
- 12 Melin, R.E. "Uranium Deposits in Shirley Basin, Wyoming." Laramie, Wyoming: University of
13 Wyoming. Vol. 8, No. 2.1. pp. 143–149. 1969.
14
- 15 Mrak, V.A. "Uranium Deposits in the Eocene Sandstones of the Powder River Basin,
16 Wyoming." *Ore Deposits of the United States 1933–1967*. New York City, New York:
17 American Institute of Mining Engineers. pp. 838–848. 1968.
18
- 19 Munn, L.C. and C.S. Arneson. "Soils of Wyoming—A Digital Statewide Map at 1:500,000-
20 Scale." B-1069. Laramie, Wyoming: University of Wyoming Agricultural Experiment Station,
21 College of Agriculture. 1998.
22
- 23
- 24 National Climatic Data Center. "NCDC U.S. Storm Events Database." 2007.
25 <<http://www4.ncdc.noaa.gov/cgi-win/wwcgi.dll?wwEvent~Storms>> (14 April 2008).
26
- 27 National Climatic Data Center. "Climates of the States, Climatology of the United States No. 60
28 (New Mexico, Nebraska, South Dakota, and Wyoming)." Asheville, North Carolina: National
29 Oceanic and Atmospheric Administration. 2005. <[http://cdo.ncdc.noaa.gov/cgi-](http://cdo.ncdc.noaa.gov/cgi-bin/climatenormals/climatenormals.pl?directive=prod_select2&prodtype=CLIM60&subnum=>)
30 [bin/climatenormals/climatenormals.pl?directive=prod_select2&prodtype=CLIM60&subnum=>](http://cdo.ncdc.noaa.gov/cgi-bin/climatenormals/climatenormals.pl?directive=prod_select2&prodtype=CLIM60&subnum=>)
31 (30 January 2005).
32
- 33 National Climatic Data Center. "Climatography of the United States No. 20: Monthly Station
34 Climate Summaries, 1971–2000." Asheville, North Carolina: National Oceanic and
35 Atmospheric Administration. 2004.
36
- 37 National Weather Service. "NOAA Technical Report NWS 33: Evaporation Atlas for the
38 Contiguous 48 United States." Washington, DC: National Oceanic and Atmospheric
39 Administration. 1982.
40
- 41 Natrona County School District No. 1. 2007. <www.natronaschools.org> (15 October 2007).
42
- 43 Niobrara County School District No. 1. 2008. <www.lusk.k12.wy.us/> (27 February 2008).
44
- 45 NRC. "Environmental Assessment For the Addition of the Reynolds Ranch Mining Area to
46 Power Resources, Inc's Smith Ranch/Highland Uranium Project Converse County." Wyoming.
47 Source Material License No. SUA-1548. Docket No. 40-8964. 2006.
48

Description of the Affected Environment

1 NRC. NUREG-1508, "Final Environmental Impact Statement to Construct and Operate the
2 Crown Point Uranium Solution Mining Project, Crown Point, New Mexico." Washington, DC:
3 NRC. February 1997.

4
5 Platte County School District No. 1. 2008. <<http://platte.schoolfusion.us/>> (4 March 2008).

6
7 PRI. "License Amendment Request-Addition of Reynolds Ranch Amendment Area."
8 ML050390076. 2005.

9
10 PRI. Reynolds Ranch Amendment. Permit Mine No. 1548-Smith Ranch-Highland Uranium
11 Project. Volume III. 2004.

12
13 Rackley, R.I. "Environment of Wyoming Tertiary Uranium Deposits." *American Association
14 Petroleum Geologists Bulletin*. Vol. 56, No. 4. 1972.

15
16 Schoolbug.org. "City of Casper." 2007a. <www.schoolbug.org> (15 October 2007).

17
18 Schoolbug.org. "Converse County, Wyoming." 2007b. <www.schoolbug.org>
19 (15 October 2007).

20
21 Trentham, R.C. and I.P. Orajaka. "Leaching of Uranium From Felsic Volcanic Rocks:
22 Experimental Studies." *Uranium*. Vol. 3. pp. 55-67. 1986.

23
24 USACE. "Interim Regional Supplement to the Corps of Engineers Wetland Delineation Manual:
25 Great Plains Region." Vicksburg, Mississippi: USACE, U.S. Army Engineer Research and
26 Development Center. August 2006.

27
28 USACops. "Wyoming." <www.usacops.com> (15 April 2008)

29
30 U.S. Census Bureau. "American FactFinder 2000 Census Data." 2008.
31 <<http://factfinder.census.gov>> (25 February 2008).

32
33 U.S. Census Bureau. "Metropolitan and Micropolitan Statistical Areas." 2007.
34 <www.census.gov> (15 October 2007).

35
36 U.S. Fish and Wildlife Service. "Mountain Prairie Region." 2008a. <[http://www.fws.gov/
37 mountain%2Dprairie/](http://www.fws.gov/mountain%2Dprairie/)> (15 February 2008).

38
39 U.S. Fish and Wildlife Service. "Medicine Bow-Routt National Forests, Thunder Basin National
40 Grassland." Laramie, Wyoming: USFS. 2008b. <<http://www.fs.fed.us/r2/mbr/index.shtml>>
41 (27 February 2008).

42
43 U.S. Geological Survey. "Water Watch." 2008. <<http://water.usgs.gov/waterwatch>>
44 (28 February 2008).

45
46 U.S. Geological Survey. "A Tapestry of Time and Terrain." Denver, Colorado: U.S. Geological
47 Survey. 2004. <<http://tapestry.usgs.gov/Default.html>> (25 February 2008).

48
49 U.S. Geological Survey. "A Field Conference on Impacts of Coalbed Methane Development in
50 The Powder River Basin, Wyoming." U.S. Geological Survey Open-File Report 01-126.

- 1 Reston, Virginia: U.S. Geological Survey. 2001. <<http://pubs.usgs.gov/of/2001/ofr-01-126/figures.html>> (25 February 2008).
- 2
3
- 4 Washington State Department of Transportation. "WSDOT's Guidance for Addressing Noise
5 Impacts in Biological Assessments—Noise Impacts." Seattle, Washington: Washington State
6 Department of Transportation. 2006. <[http://www.wsdot.wa.gov/TA/Operations/
7 Environmental/NoiseChapter011906.pdf](http://www.wsdot.wa.gov/TA/Operations/Environmental/NoiseChapter011906.pdf)> (12 October 2007).
- 8
- 9 Weston County School District No. 1. 2008. <www.weston1.k12.wy.us/> (27 February 2008).
- 10
- 11 Whitehead, R.L. "Groundwater Atlas of the United States, Montana, North Dakota, South
12 Dakota, Wyoming." U.S. Geological Survey Report HA 730–I. Denver, Colorado:
13 U.S. Geological Survey. 1996. <http://capp.water.usgs.gov/gwa/ch_i/index.html>
14
- 15 Wyoming Department of Environmental Quality. "Chapter 2, Ambient Standards." 2006.
16 <<http://deq.state.wy.us/aqd/standards.asp>> (23 October 2007).
- 17
- 18 Wyoming Department of Revenue. 2008. <<http://revenue.state.wy.us>> (25 February 2008).
- 19
- 20 Wyoming Department of Revenue. "Sales and Tax Distribution Report by County." Cheyenne,
21 Wyoming: Wyoming Department of Revenue. 2007. <<http://revenue.state.wy.us>>
22 (25 February 2008).
- 23
- 24 Wyoming Department of Revenue. "Sales and Tax Distribution Report by County." Cheyenne,
25 Wyoming: Wyoming Department of Revenue. 2007. <<http://revenue.state.wy.us>> (18 October
26 2007).
- 27
- 28 World Wildlife Fund. "Colorado Rockies Forests." Washington, DC: World Wildlife Fund.
29 2007a. <http://www.worldwildlife.org/wildworld/profiles/terrestrial/na/na0511_full.html>
30 (10 October 2007).
- 31
- 32 World Wildlife Fund. "Northern Short Grasslands." Washington, DC: World Wildlife Fund.
33 2007b. <http://www.worldwildlife.org/wildworld/profiles/terrestrial/na/na0811_full.html>
34 (13 September 2007).
- 35
- 36 World Wildlife Fund. "South Central Rockies Forests." Washington, DC: World Wildlife Fund.
37 2007c. <http://www.worldwildlife.org/wildworld/profiles/terrestrial/na/na0528_full.html>
38 (10 October 2007).
- 39
- 40 World Wildlife Fund. "Wildfinder—Mapping the World's Species: Ecoregion NA1313 (Wyoming
41 Basin Shrub Steppe)." Washington, DC: World Wildlife Fund. 2007d.
42 <<http://www.worldwildlife.org/wildfinder/searchByPlace.cfm?ecoregion=Na1313>>
43 (15 October 2007).
- 44
- 45 World Wildlife Fund. "Wyoming Basin Shrub Steppe." Washington, DC: World Wildlife Fund.
46 2007e. <http://www.worldwildlife.org/wildworld/profiles/terrestrial/na/na1313_full.html>
47 (13 September 2007).
- 48
- 49 Wyoming Department of Transportation. "WYDOT Traffic Analysis." Cheyenne, Wyoming:
50 Wyoming Department of Transportation. 2005. <<http://www.dot.state.wy.us/>>

Description of the Affected Environment

1 Default.jsp?sCode=hwyta> (25 February 2008).

2
3 Wyoming Game and Fish Department. "Comprehensive Wildlife Conservation Strategy."
4 Cheyenne, Wyoming: Wyoming Game and Fish. 2008.

5 <<http://gf.state.wy.us/wildlife/CompConvStrategy/Species>> (19 February 2008).

6
7 Wyoming Game and Fish Department. "Official State List of Birds, Mammals, Amphibians, and
8 Reptiles in Wyoming." Cheyenne, Wyoming: Wyoming Game and Fish. 2007a.

9 <<http://gf.state.wy.us/wildlife/nongame/SpeciesList/index.asp>> (15 October 2007).

10
11 Wyoming Game and Fish Department. "Terrestrial Habitat/Aquatic Habitat/Habitat
12 Management." Cheyenne, Wyoming: Wyoming Game and Fish. 2007b.

13 <<http://gf.state.wy.us/habitat/aquatic/index.asp>> (13 September 2007).

14
15 Wyoming Geographic Information Science Center. "Wyoming Gap Analysis." Laramie,
16 Wyoming: University of Wyoming, Wyoming Geographic Information Science Center. 2007.

17 <<http://www.wygisc.uwyo.edu/wbn/gap.html>> (25 February 2008).

18
19 Wyoming State Geological Survey, Industrial Minerals and Uranium Section. "Uranium Page."

20 <<http://www.wsgs.uwyo.edu/minerals/uranium.aspx>> (15 July 2005).

21
22 Wyoming Workforce Development Council. "Wyoming workers Commuting Patterns Study."
23 Cheyenne, Wyoming: Wyoming Workforce Development Council. 2007.

24 Zielinski, R.A. "Volcanic Rocks as Sources of Uranium." *Uranium Deposits in Volcanic Rocks*.
25 Vienna, Austria: International Atomic Energy Agency. pp. 83–95. 1984.

26
27 Zielinski, R.A. "Tuffaceous Sediments as Source Rocks for Uranium—A Case Study of the
28 White River Formation, Wyoming." *Journal of Geochemical Exploration*. Vol. 18. pp. 285–306.
29 1983.

3.4 Nebraska-South Dakota-Wyoming Uranium Milling Region

3.4.1 Land Use

The Nebraska-South Dakota-Wyoming Uranium Milling Region defined in this GEIS, is represented by a south-southeast–north-northwest swath of land encompassing parts of Sioux and Dawes counties in Nebraska, Fall River, Custer, Pennington and Lawrence counties in South Dakota, and Niobrara, Weston and Crook counties in Wyoming (Figure 3.4-1).

This region lies within portions of the Missouri Plateau, the Black Hills and the High Plains sections of the Great Plains province (U.S. Geological Survey, 2004). The locations of past, current and potential uranium milling operations are found in the Crow Butte Uranium District located in Dawes County, Nebraska; in the Southern Black Hills Uranium District in Fall River County, South Dakota and Niobrara County, Wyoming; and in the Northern Black Hills Uranium District in Crook County, Wyoming (Figure 3.4-2). Details on the geology and soils of these three districts are provided in Section 3.4.3.

The general land ownership and use statistics for the Nebraska-South Dakota-Wyoming Uranium Milling Region shown below were calculated using the Geographic Information System used to construct the map shown in Figure 3.4-1. Private lands (59 percent) and National Forest and National Grassland (38 percent combined) account for 97 percent of this region (Table 3.4-1).

In the areas of interest in Dawes and Sioux Counties in Nebraska, the predominant land cover consists of a mix of western short grass prairie and western wheat grass prairie, followed by agricultural fields and ponderosa pine forests and woodlands (Henebry, et al., 2005). A large portion of Dawes and Sioux Counties is occupied by the Oglala National Grassland to the north and west and by the Nebraska National Forest in the center, which are both administered by the USFS (Figure 3.4-1). These federal lands offer general recreational activities, including camping, fishing and hunting (USFS, 2008b). Chadron, a 394-ha [972-acre] state park in the heart of the Nebraska National Forest and Fort Robinson, a 8,900-ha [22,000-acre] state park of Pine Ridge scenery west of Crawford, also offer general recreational activities to the public. (Nebraska Game and Parks Commission, 2008). Similar to nearby Niobrara County in Wyoming to the west and Fall River County in South Dakota to the north, the dominant land use in these two northwestern Nebraska counties is cattle grazing on both public and private rangeland and associated livestock feed production. Cultivated lands mixed with the rangeland are used primarily to produce winter wheat and hay, which is both grazed and harvested.

Approximately half of Fall River County in the southwest corner of South Dakota is occupied by the Buffalo Gap National Grassland to the south and by the Black Hills National Forest to the north, which are both managed by the USFS. Higher elevation areas to the north into the Black Hills National Forest create favorable growing conditions for ponderosa pine. The lower elevation areas surrounding the Black Hills to the south are primarily used as rangeland for livestock grazing and as agricultural land. Hay and winter wheat farming are the principal agricultural uses in dry land areas, and alfalfa, corn, and vegetables are typically grown in wetter valley areas and on irrigated land (South Dakota State University, 2001). A large part of Shannon County, South Dakota, which abuts Fall River County to the East, is occupied entirely by the Pine Ridge Indian Reservation (Figure 3.4-1).



- Legend**
- | | | | |
|---|--|--|---------------------------|
| ▲ | Ur Milling Site (NRC) | | Forest Service |
| ● | City | | Department of Defense |
| | South Dakota - Nebraska Milling Region | | Bureau of Land Management |
| | Interstate Highway | | National Park Service |
| | US Highway | | Bureau of Indian Affairs |
| | State Highway | | Bureau of Reclamation |
| | Railroad | | |
| | Water bodies (Lakes, Bays, ...) | | |
| | Rivers and Streams | | |
| | State Boundary | | |
| | Counties | | |

3.4-2




Figure 3.4-2. Map Showing the Nebraska-South Dakota-Wyoming Uranium Milling Region and Uranium Milling Sites in the Black Hills Uranium Districts in South Dakota and Wyoming and in the Crow Butte Uranium District in Nebraska

Table 3.4-1. Land Ownership and General Use in the Nebraska-South Dakota-Wyoming Uranium Milling Region

Land Ownership and General Use	Area (mi ²)	Area (km ²)	Percent
State and Private Lands	5,379	13,932	58.6
U.S. Forest Service (USFS), National Forest	1,979	5,125	21.5
USFS, National Grassland	1,553	4,022	16.9
U.S. Bureau of Land Management, Public Domain Land	185	480	2
National Park Service, National Park	41	107	0.5
Bureau of Reclamation	16	42	0.2
USFS, Wilderness	22	56	0.2
USFS, National Recreation Area	4	11	0.05
National Park Service, National Monument	4	11	0.05
Totals	9,185	23,788	100

More than half of Custer, Pennington and Lawrence counties in South Dakota is also occupied by the Black Hills National Forest (Figure 3.4-1). In these counties the majority of the land cover consists of ponderosa pine forest associated with short to tall grass lands and agricultural fields (South Dakota State University, 2001).

Historically, the Black Hills have been prospected and mined for many minerals, metals, and materials. Recreational activities provided in the Buffalo Gap National Grassland and in the Black Hills National Forest are similar to those described for USFS lands in Nebraska and in the Wyoming East Uranium Milling Region (USFS, 2008a,b).

In the eastern and northeastern Wyoming Counties of Niobrara and Crook, land ownership is predominantly private as it is in the Wyoming East Uranium Milling Region. BLM administered lands, which are scattered and mixed with state and private lands, represent less than 10 percent of the land. In Weston County, located between Niobrara and Crook counties, land ownership is dominated by the USFS Thunder Basin National Grassland. In its eastern half, a large portion of Crook County is occupied by the Black Hills National Forest. To the west of the forest on Route 24, Devils Tower National Monument, administered by the National Park Service, provides additional recreational activities in Crook County (Figure 3.4-1).

The characteristics of open rangeland in these three eastern Wyoming counties are similar to those of the Wyoming East Uranium Milling Region described in Section 3.3.1. Cattle and sheep grazing represent the primary land use on private and federal lands. Recreational activities available on federal lands are also similar to those described above for parts of Nebraska, South Dakota and the Wyoming East Uranium Milling Region (Section 3.3.1).

3.4.2 Transportation

Past experience at NRC licensed ISL facilities indicate these facilities rely on roads for transportation of goods and personnel (Section 2.8). As shown on Figure 3.4-3, the Nebraska-South Dakota-Wyoming Uranium Milling Region is accessible by a variety of highways. In the northern part of the region, Interstate 90 connects Gillette, Wyoming and Rapid City, South Dakota. U.S. Highway 212 enters the region from Montana to the north intersecting U.S. Highway 85 and then crossing Interstate 90 to the south and traversing the region



Description of the Affected Environment

southbound to intersect U.S. Highway 20. U.S. Highway 20 traverses the south portion of the region and connects with Interstate 25 to the west. A rail line services the central portion of the South Dakota/Nebraska region along U.S. Highway 16 from the west to the intersection with U.S. Highway 85 at Newcastle and then south to Crawford at the southern boundary of the region.

Areas of past, present, or future uranium milling interest in the region are shown in Figure 3.4-3. These areas are located in three subregions when considering site access by local roads. The area of milling interest in the northeastern part of the region (north of Aladdin, Wyoming) is accessible by local access roads to U.S. Highway 212 southeast to U.S. Highway 85 south which intersects Interstate 90. Traveling west from Aladdin, State Route 24 connects to U.S. Highway 14 and Interstate 90 continuing west to Gillette. Milling sites further to the southwest of the region (near Burdock, South Dakota) are served by local access roads and U.S. Highway 18 west to connect with U.S. Highway 85 southbound that exits the region from the southwest. At Lusk, Wyoming U.S. Highway 20 west provides access to Interstate 25. Areas of milling interest near the southern border of the region (near Crawford, Nebraska) are served by local access roads to U.S. Highway 20 which exits the region to the west to intersect Interstate 25.

Table 3.4-2 provides available traffic count data for roads that support areas of past or future milling interest in the Nebraska-South Dakota-Wyoming Uranium Milling Region. Counts are variable with the minimum all vehicle count at 333 vehicles per day on U.S. Highway 16 West of Custer (westbound) and the maximum on Interstate 90 East of Spearfish (between Spearfish and Whitewood) at 9,491 vehicles per day. Most of the vehicle counts in the Nebraska-South Dakota-Wyoming Uranium Milling Region are above 400 vehicles per day.

Yellowcake product shipments are expected to travel from the milling facility to a uranium hexafluoride production (conversion) facility in Metropolis, Illinois (the only facility currently licensed by NRC in the U.S. for this purpose). Major interstate transportation routes are expected to be used for these shipments, which are required to follow NRC packaging and transportation regulations in 10 CFR Part 71 and U.S. Department of Transportation hazardous material transportation regulations at 49 CFR Parts 171–189. Table 3.4-3 describes representative routes and distances for shipments of Yellowcake from locations of Uranium milling interest in the South Dakota/Nebraska Uranium Milling Region. Representative routes are considered owing to the number of routing options available that could be used by a future ISL facility.

3.4.3 Geology and Soils

Sandstone-hosted uranium ore deposits have been identified in western South Dakota, northeastern Wyoming, and in northwestern Nebraska (Figure 3.4-2). In the Nebraska-South Dakota-Wyoming Uranium Milling Region, uranium mineralization is found in fluvial sandstones in two major areas: the Black Hills of western South Dakota and northeastern Wyoming and the Crawford Basin of northwestern Nebraska. Uranium mineralization in the sandstone-hosted uranium deposits in these two areas is in a geologic setting amenable to recovery by ISL milling.

3.4.3.1 The Black Hills (Western South Dakota-Northeastern Wyoming)

The Black Hills are an asymmetrical domal uplift elongated in a northwest direction (Figure 3.4-4). Economically significant uranium discoveries in the Black Hills are contained within strata of the Inyan Kara Group (Chenoweth, 1988). Prior to 1968, the Black Hills produced approximately 1,800 metric tons [2,000 tons] of U_3O_8 (Hart, 1968). The bulk of this

Table 3.4-2. Average Annual Daily Traffic Counts for Roads in the Nebraska-South Dakota-Wyoming Uranium Milling Region*

Road Segment	County, State	All Vehicles
State Route 24 at Devils Tower Junction (intersection with U.S. Highway 14)	Crook, Wyoming	982–1,236
State Route 14 at Devils Tower Junction (west intersection with State Route 24)	Crook, Wyoming	610–675
Interstate 90 at County Border East (near Beulah, Wyoming)	Crook, Wyoming	4,048–5,272
U.S. Highway 85 North of Belle Fourche (southbound in direction of U.S. Highway 212)	Butte, South Dakota	468–905†
Interstate 90 East of Spearfish (between Spearfish and Whitewood)	Lawrence, South Dakota	5,201–9,491†
U.S. Highway 16 West of Custer (westbound)	Custer, South Dakota	333–1,231†
U.S. Highway 385 North of Hot Springs (near north county line)	Fall River, South Dakota	425–1,243†
U.S. Highway 18 at Mule Creek Junction (intersection with U.S. Highway 85)	Niobrara, Wyoming	817–1,192
U.S. Highway 85 at Mule Creek Junction (south of intersection with U.S. Highway 18)	Niobrara, Wyoming	1,327–2,037
U.S. Highway 20 at Van Tassell (at east county line)	Niobrara, Wyoming	415–552
U.S. Highway 20 at Manville South (intersection with State Route 270)	Niobrara, Wyoming	1,418–1,891

*Wyoming Department of Transportation. "Wyoming Department of Transportation Traffic Analysis." 2005. <<http://dot.state.wy.us/Default.jsp?sCode=hwyta>> (27 December 2005). South Dakota Department of Transportation. "Automatic Traffic Recorder Data." 2008. <<http://gis.sd.gov/dot%5Fctsys/>> (January 2008).
†Data for South Dakota are monthly averages of daily counts; Wyoming data are the arithmetic mean of average annual daily counts for each day of the week.

Table 3.4-3. Representative Transportation Routes for Yellowcake Shipments From the Nebraska-South Dakota-Wyoming Uranium Milling Region*

Origin	Destination	Major Links	Distance (mi)
North of Aladdin, Wyoming	Metropolis, Illinois	Local access road northeast to U.S. Highway 212 U.S. Highway 212 southeast to U.S. Highway 85 U.S. Highway 85 south to Interstate 90 Interstate 90 east to Sioux Falls, South Dakota Interstate 29 south to Kansas City, Missouri Interstate 70 east to St. Louis, Missouri Interstate 64 east to Interstate 57 Interstate 57 south to Interstate 24 Interstate 24 south to U.S. Highway 45 U.S. Highway 45 west to Metropolis, Illinois	1,230

Table 3.4-3. Representative Transportation Routes for Yellowcake Shipments From the Nebraska-South Dakota-Wyoming Uranium Milling Region* (continued)

Origin	Destination	Major Links	Distance (mi)
Edgemont, South Dakota	Metropolis, Illinois	Local access road south to U.S. Highway 18 U.S. Highway 18 west to U.S. Highway 85 U.S. Highway 85 south to U.S. Highway 20 U.S. Highway 20 west to Interstate 25 Interstate 25 south to Denver, Colorado Interstate 70 east to St. Louis, Missouri Interstate 64 east to Interstate 57 Interstate 57 south to Interstate 24 Interstate 24 south to U.S. Highway 45 U.S. Highway 45 west to Metropolis, Illinois	1,410
Crawford, Wyoming	Metropolis, Illinois	Local access roads north to U.S. Highway 20 U.S. Highway 20 west to Interstate 25 Interstate 25 south to Denver, Colorado Denver, Colorado, to Metropolis, Illinois (as above)	1,360

*American Map Corporation. "Road Atlas of the United States, Canada, and Mexico." Long Island City, New York: American Map Corporation. p. 144. 2006.

production came from the Hulett Creek and Carlile districts of the northern Black Hills and the Edgemont district of the southern Black Hills (Figure 3.4-4).

Stratigraphic units present in the Black Hills area are shown in Figure 3.4-5. Jurassic (144 to 206 million year old) and Cretaceous (65 to 144 million year old) rocks crop out low on the flanks of the Black Hills and form the eroded surface upon which younger rocks were deposited (Harshman, 1968). Sedimentary rocks of Tertiary (1.8 to 65 million year old) age are virtually absent from the Black Hills. However, remnants of Miocene (5.3 to 23.8 million year old) and/or Paleocene (54.8 to 65 million year old) age rocks on the flanks of the Black Hills indicate that at one time rocks of middle and late Tertiary age may have extended across the area and at least partially buried the Black Hills uplift. The Tertiary rocks are tuffaceous (i.e., they contain materials made from volcanic rock and mineral fragments in a volcanic ash matrix) and clastic (i.e., they contain fragments or grains of older rocks) and are of fluvial (river), lacustrine (lake), and paludal (marsh) origin.

The Inyan Kara Group is Lower Cretaceous (99 to 144 million years old) in age and consists of subequal amounts of complexly interbedded sandstone and claystone (Renfro, 1969). The Inyan Kara is bounded below by continental Jurassic sediments of the Morrison Formation and is overlain by marine sediments of the Lower Cretaceous Skull Creek Shale. Resistant Inyan Kara sediments form the outermost ring of hogback ridges that crop out in a roughly oval pattern around the flanks of the Black Hills. Major uranium deposits occur from 2 to 8 km [1 to 5 mi] downdip from the main Inyan Kara escarpment at depths ranging from 30 to 180 m [100 to 600 ft].

The Inyan Kara Group is formally subdivided into the Lakota Formation and the Fall River Formation, which are generally accepted to be respectively continental and marginal marine in origin (Robinson, et al., 1964). The source of sediment for the Lakota and Fall River is considered to include all pre-Cretaceous sediments that were exposed to the south and east of the Black Hills (Renfro, 1969).




Figure 3.4-4. Outcrop Map of the Inyan Kara Group in the Black Hills of Western South Dakota and Northeastern Wyoming Showing the Locations of Principal Uranium Mining Districts (From Hart, 1968)

Black Hills Area			
System	Series	Formation	
Tertiary	Pliocene	Ogallala Formation	
	Miocene	Arikaree Formation	
	Oligocene	White River Formation	
	Eocene	(Absent)	
	Paleocene	Fort Union Formation	
Cretaceous	Upper	Hell Creek Formation	
		Fox Hills Sandstone	
		Pierre Shale	
		Niobrara Formation	
		Carlile Shale, Greenhorn Formation, and Belle Fourche Shale	
	Lower	Mowry Shale	
		Newcastle Sandstone and Skull Creek Shale	
		Fall River and Lakota Formations	Inyan Kara Group
Jurassic	Morrison Formation		
	Sundance Formation		
	Gypsum Spring Formation		

Figure 3.4-5. Principal Stratigraphic Units in the Black Hills Area of Western South Dakota and Northeastern Wyoming (Modified From Harshman, 1968)

1 The Lakota is a sequence of coastal-plain deposits of fine-grained, poorly sorted sandstone and
2 mudstone; channel-fill deposits of cross-bedded sandstone; natural levee and overbank
3 deposits of lenticular (i.e., deposits with a lens-shaped cross section), fine-grained,
4 carbonaceous sandstone and siltstone; and floodplain deposits of bedded siltstone, mudstone,
5 and claystone (Maxwell, 1974). The Lakota Formation is from 15 to 90 m [50 to 300 ft] thick and
6 thickens regionally from northwest to southeast (Chenoweth, 1988).

7
8 The oldest Lakota strata are thin, discontinuous dark gray to olive black, humic sandstone and
9 claystone containing sparse sub-bituminous coal seams (Renfro, 1969). These strata appear to
10 conform with the underlying Morrison Formation. The lowermost Lakota grades upward to a
11 sequence of dark gray, medium- to coarse-grained, cherty and quartzose sandstone containing
12 abundant disseminated carbon and pore-filling, massive pyrite. The uppermost Lakota consists
13 of lenticular greenish gray to dark gray, fine- to medium-grained, quartzose sandstone and
14 vari-colored claystone.

15
16 Dondanville (1963) divided the Fall River Formation into deltaic and marine facies. The deltaic
17 facies forms approximately 50 percent of the formation and consists of channel sandstone,
18 interchannel sandstone and mudstone, and blanket sandstones formed during erosion of
19 abandoned deltas. The marine and marginal-marine rocks consist of offshore and lagoonal
20 mudstone and shale, and bar and spit sandstone. The Fall River is from 30 to 45 m [100 to
21 150 ft] thick and thickens regionally from southeast to northwest at the expense of the
22 underlying Lakota Formation.

23
24 Renfro (1969) describes the Fall River as a light to dark gray, fine- to medium-grained quartzose
25 sandstone containing traces of glauconite and abundant disseminated carbon, pyrite, and
26 detrital chert. Thin beds of claystone and siltstone are common. The Fall River is in
27 conformable contact and regionally intertongues with the overlying Skull Creek Shale.

28
29 Uranium deposits in the Inyan Kara Group are typified by roll-front accumulations (see
30 Section 3.1.1). Geometric complexity of individual roll-fronts is governed by the stratigraphic
31 complexity of the Inyan Kara host sediments. Most roll-fronts are within tabular sandstones of
32 the Fall River Formation or widespread cherty sandstone facies of the Lakota Formation and
33 have simple C-shaped cross sections that extend laterally for tens of miles (Figure 3.4-6).
34 Roll-front deposits in the more complex sandstone and claystone facies of the upper Lakota
35 Formation are very erratic and generally contain relatively weak mineralization. Mineralization
36 in the roll limbs seldom extends more than 90 to 120 m [300 or 400 ft] up-plunge from the roll
37 fronts. Although roll fronts in the Inyan Kara are common, ore grade mineralization is restricted
38 vertically and laterally. Ore most often occurs in terminal lobes of the roll-front trends.

39
40 Within Inyan Kara ore bodies, uranium minerals coat sand grains, fill interstices between grains,
41 and are finely disseminated in organic matter (Renfro, 1969). In oxidized deposits, the uranium
42 vanadates, carnotite, tyuyamunite, and meta-tyuyamunite are the principal ore minerals.
43 Uraninite and coffinite are the main minerals in unoxidized ore. Pyrite, marcasite, and calcite
44 are present as gangue minerals (i.e., low-value minerals intermixed with ore minerals). Tongues
45 of hematite-stained pinkish-red sandstone are present at most of the deposits. This alteration is
46 due to the oxidation of pyrite in the sandstone by migrating groundwater.

47
48 The source of uranium in the Inyan Kara deposits is unknown, but two main theories have been
49 proposed. Renfro (1969) proposed that the uranium and other metals indigenous to the Lakota
50 and Fall River sediments were mobilized by oxidizing groundwater and transported downdip,
51




Figure 3.4-6. Schematic Cross Section Through a Typical Inyan Kara Roll-Front Deposit Showing Differences in U_3O_8 Concentration Between “Fresh” (i.e., Unoxidized) and “Altered” Ground (Modified From Renfro, 1969)

where they were precipitated along an oxidation-reduction boundary. Hart (1968) proposed that uranium was leached by groundwater from tuffaceous beds of the White River Group that were unconformably deposited across the eroded Black Hills uplift. Migrating groundwater carried the uranium into the permeable host rocks where it traveled downdip into reducing environments. Later groundwater movements remobilized and redeposited some of the ore bodies.

The surface of the Black Hills range is still largely mantled by sedimentary rocks that form an outer ring of hogback ridges that crop out in a roughly oval pattern around the flanks of the range. Soils in low lying areas adjacent to the Black Hills of western South Dakota and northeastern Wyoming consist of the weathering products of these sedimentary rocks. The topographic position and texture of typical soils in the Black Hills were obtained from the Soils Map of Wyoming (Munn and Arneson, 1998). This map was designed primarily for a statewide study of groundwater's vulnerability to contamination and would not be expected to be used for site-specific soil interpretations at proposed ISL milling facilities. For site-specific evaluations, detailed soils information would be expected to be obtained from published county soil surveys or NRCS.

Soils within the Black Hills area of western South Dakota and northeastern Wyoming are mostly fine textured (fine or fine-loamy soils). Shallow fine and fine-loamy soils with little or no subsoil development are found on ridges and steep slopes on the flanks of Black Hills. On gently sloping to moderately steep slopes adjacent to ridges, moderately deep fine and fine-loamy soils with moderate- to well-developed soil horizons are found. These soils are generally light-colored and depleted in moisture. On low gradient surfaces, such as terraces and floodplains, deep fine and fine-loamy soils with well developed subsoil horizons are found. Dark-colored, base-rich soils formed under grass are generally associated with floodplains along streams with permanent high water tables.

3.4.3.2 The Crawford Basin (Northwestern Nebraska)

Uranium deposits in northwestern Nebraska are located in Dawes and Sioux Counties in what has been named the Crawford Basin (Figure 3.4-2) (DeGraw, 1969). In 1979, an area west of the city of Crawford in Sioux County and an area north of Crawford in Dawes County were identified as having considerable weak uranium mineralization associated with vague

oxidation-reduction boundaries (Collings and Knode, 1984). In 1981 and 1982, the Crow Butte mineralized trend was discovered southeast of Crawford in Dawes County. The Crow Butte mineralized trend is about 10 km [6 mi] long and up to 900 m [3,000 ft] wide with ore reserves calculated to be over 13,600 metric tons [15,000 tons] of U_3O_8 having an average grade exceeding 0.25 percent U_3O_8 (Collings and Knode, 1984). Uranium mineralization in the Crow Butte area occurs exclusively within the Chadron Sandstone.

The Crawford Basin is a triangular, asymmetrical basin bounded by the Black Hills Uplift on the northwest, the Chadron Arch to the west, and the Cochran Arch to the south (Figure 3.4-7). As

Black Hills Uplift




Figure 3.4-7. Bedrock Geology and Major Structural Features of the Crawford Basin (Modified From Gjelsteen and Collings, 1988)

a result of the Black Hills Uplift, formations underlying the uranium milling areas in the Crawford Basin dip gently to the south. The single most prominent structural feature within the Crawford

Description of the Affected Environment

1 Basin is the White River Fault. It is located north of Crawford and strikes northeast to southwest
2 with the upthrown side to the south. The total vertical displacement is 60 to 120 m [200 to
3 400 ft].

4
5 A generalized stratigraphic section of sedimentary strata in the Crow Butte mining area of
6 northwestern Nebraska is shown in Figure 3.4-8. Stratigraphic descriptions presented here are
7 limited to formations that may be involved in potential milling operations or formations that may
8 have environmental significance, such as important aquifers or confining units above and below
9 potential milling zones.

10
11 The Upper Cretaceous (65 to 99 million year old) Pierre Shale is a widespread, compositionally
12 uniform, dark gray to black marine shale, which outcrops extensively in Dawes County north of
13 the Crow Butte mining area (Collings and Knode, 1984). In Dawes County, the Pierre shale is
14 365 to 460 m [1,200 to 1,500 ft] thick and is essentially impermeable. Due to aerial exposure
15 and subsequent erosion, the top of the present-day Pierre contact marks a major unconformity
16 and exhibits a paleotopography with considerable relief (DeGraw, 1969). As a result of the
17 extended exposure to atmospheric weathering, an ancient soil horizon, or paleosol, from 0 to
18 10 m [0 to 33 ft] thick was formed on the surface of the Pierre Shale.

19
20 The Oligocene (23.8 to 33.7 million year old) White River Group lies unconformably on top of
21 the Pierre Shale. The White River Group consists of the Chadron and Brule Formations. The
22 Chadron comprises three distinct units: the Basal Chadron Sandstone Member, Middle Chadron
23 Member, and Upper Chadron Member.

24
25 Uranium mineralization in the Crow Butte mineralized trend occurs exclusively within the
26 Basal Chadron Sandstone. The Basal Chadron Sandstone Member consists of coarse-grained
27 arkosic sandstone (i.e., sandstone containing a significant fraction of feldspar) with frequent
28 interbedded thin clay beds. Occasionally, the lower portion of the Basal Member is a very
29 coarse, poorly sorted conglomerate. The Basal Sandstone is the depositional product of a
30 large, braided stream system and ranges from 0 to 105 m [0 to 350 ft] thick.

31
32 The Middle Chadron Member overlies the Basal Sandstone Member. The lower part of the
33 Middle Member is impermeable brick-red clay with occasional interbedded gray-green clay. The
34 brick-red clay grades upward to a light green-gray sandy claystone. The upper part of the
35 Middle Member is light gray bentonitic clay. The Middle Member ranges from 12 to 30 m [40 to
36 100 ft] thick. The Upper Chadron Member consists of massive claystones and siltstones,
37 generally considered to be fluvial in origin (Vondra, 1958). The Upper Chadron Member
38 averages 30 m [100 ft] thick throughout the Crow Butte mining area.

39
40 The Brule Formation lies conformably on top of the Chadron Formation and consists almost
41 entirely of siltstones with minor sand channels. The Brule is subdivided into two members: the
42 Orella and the Whitney. The Orella lies directly on the Chadron and is composed of buff to
43 brown siltstones. The Whitney comprises massive buff to brown siltstones and contains several
44 volcanic ash horizons.

45
46 Uranium deposits in the Basal Chadron Sandstone are associated with oxidation-reduction
47 boundaries or roll-fronts (see Section 3.1.1) adjacent to the White River Fault (Figure 3.4-9).
48 Within the Crow Butte uranium ore trend, the Basal Chadron is about 12 m [40 ft] thick (Collings

1

Northwestern Nebraska			
Age	Group	Formation	Member
Miocene	Arikaree	Monroe Creek	
		Gering	
Oligocene	White River	Brule	Whitney
			Orella
		Chadron	Upper
			Middle
			Basal
Eocene ?		Paleosol	
Cretaceous		Pierre Shale	

Figure 3.4-8. Generalized Stratigraphic Units in the Crow Butte Area of Northwestern Nebraska (Modified From Collings and Knode, 1984)

1




Figure 3.4-9. Location of Oxidation-Reduction Fronts Detected During Exploration Drilling Within the Chadron Sandstone in Northwestern Nebraska. Arrow Shows Direction of Groundwater Flow at the Time of Mineralization as Indicated by Roll-Front Geometry (Modified From Gjølsteen and Collings, 1988).

and Knode, 1984). Depth to mineralization varies from 85 to 250 m [275 to 820 ft]. Uranium is present in the matrix and as a coating on grains as coffinite and uraninite and occurs locally in concentrations as high as 3.0% (Gjølsteen and Collings, 1988). The volcanoclastic sediments contained in and overlying the Chadron sandstone are considered to be the most likely source of the uranium of the roll-front deposits in the Crawford Basin because of their abundance, close proximity, and susceptibility to dissolution (Gjølsteen and Collings, 1988).

The distribution and occurrence of soils in Nebraska-South Dakota-Wyoming Uranium Milling Region varies regionally with respect to landform development (e.g., ridges, floodplains, hills) and locally with changes in slope, geology, vegetation, climate, and time. The general characteristics of soils associated with landforms in Dawes County was obtained from the U.S. Department of Agriculture (NRCS, 2007). For site-specific evaluations at proposed ISL

1 milling facilities, more detailed soils information can be obtained from published county soil
2 surveys or the NRCS.

3
4 In Dawes County, silt loam and silty clay loam soils having little to moderate horizon
5 development are found on ridges. These shallow to moderately shallow soils occur on steep
6 slopes where erosion activity is greatest. Soils on hillslopes vary from soils having little or
7 moderate horizon development to soils that have well-developed horizons (deep soils). Silty
8 clay and silty clay loam soils having little to moderate horizon development are found on the
9 steeper parts of hillslopes where erosional activity is greatest. Silty clay loam and loamy very
10 fine sand soils having well-developed horizons are found on gently sloping parts of hillslopes.
11 On plains, which are nearly level or gently sloping, silt loam soils with well-developed clay
12 horizons are found. Soils found on stream terraces and flood plains are generally very deep,
13 with soil textures that are highly variable, depending on the local geology. Silty clay, silty clay
14 loam, silt loam and loam soils are found on stream terraces. Clay, loamy very fine sand, and
15 sandy loam soils are found on flood plains.

16 17 **3.4.4 Water Resources**

18 19 **3.4.4.1 Surface Waters**


20
21 The Nebraska-South Dakota-Wyoming Uranium Milling Region includes portions of
22 northwestern Nebraska, eastern Wyoming, and southwest South Dakota. Watersheds in the
23 Nebraska-South Dakota-Wyoming Uranium Milling Region are shown in Figure 3.4-10. The
24 watersheds within the Nebraska-South Dakota-Wyoming Uranium Milling Region are listed in
25 Table 3.4-4 along with the generic designated uses of surface water bodies in these
26 watersheds. The designated uses of water bodies in these watersheds differ slightly from state
27 to state. Thus, the designated uses for water bodies in watersheds that cross state boundaries
28 may be different. To simplify the discussion of the water quality characteristics of water bodies
29 in each watershed, the designated uses in Table 3.4-4 have been grouped into the following
30 generic categories: fisheries, fish and wildlife propagation, recreation, drinking water supply,
31 agriculture, industrial and aesthetic. Water bodies with the generic use as a fishery may support
32 either warmwater or coldwater species. More detailed descriptions of the designated uses in
33 each state can be found in the following references

- 34
- 35 • Wyoming – WDEQ (2001; 2006)
- 36 • Nebraska – Nebraska Department of Environmental Quality (2008)
- 37 • South Dakota – South Dakota Department of Environmental and Natural Resources
- 38 (2008)
- 39

40 Surface water features in specific areas of uranium mineralization within the Nebraska-South
41 Dakota-Wyoming Uranium Milling Region are discussed next.

42 43 **Nebraska**

44
45 The area of known uranium mineralization in Nebraska is located in Dawes County within the
46 Upper White River watershed (Figure 3.4-10) The average annual flow of the White River at the
47 Nebraska-South Dakota state line, near the northern limit of known uranium deposits is
48 approximately 1.7 m³/s [60 ft³/s] (U.S. Geological Survey, 2008a). The state designated uses
49 for the White River above Chadron, Nebraska are: drinking water supply, aquatic life (cold
50 water), agriculture, and aesthetics (Nebraska Department of Environmental Quality, 2008).



SOUTH DAKOTA - NEBRASKA REGION



Figure 3.4-10. Watersheds Within the Nebraska-South Dakota-Wyoming Uranium Milling Region

The immediate area of uranium mineralization is drained by White Clay Creek, Squaw Creek, and English Creek with headwaters in the Nebraska National Forest along Pine Ridge. Small surface impoundments are present along these creeks used for stock watering. The state designated uses for these perennial creeks are: aquatic life (cold water), fish consumption, agriculture, and aesthetics (Nebraska Department of Environmental Quality, 2008). These streams are not identified as having impaired water quality.

Table 3.4-4. Primary Watersheds in the Nebraska-South Dakota-Wyoming Uranium District and Range of Generic Designated Uses of Water Bodies Within Each Watershed

Watershed	Generic State Designated Uses of Water Bodies in the Watershed	
Upper White River	Nebraska	Fisheries Fish and Wildlife Propagation Drinking Water Recreation Agriculture Aesthetics
Hat Creek	Nebraska	Fisheries Fish and Wildlife Propagation Drinking Water Recreation Agriculture Aesthetics
	South Dakota	Fisheries Fish and Wildlife Propagation Drinking Water Recreation Agriculture Aesthetics
Angostura Reservoir	South Dakota	Fisheries Fish and Wildlife Propagation Drinking Water Recreation Agriculture Aesthetics
Cheyenne River Above Angostura Reservoir	South Dakota	Fisheries Fish and Wildlife Propagation Recreation Agriculture Aesthetics
	Wyoming	Fisheries Fish and Wildlife Propagation Drinking Water Recreation Agriculture Industrial Aesthetics

1

Table 3.4-4. Primary Watersheds in the Nebraska-South Dakota-Wyoming Uranium District and Range of Generic Designated Uses of Water Bodies Within Each Watershed (continued)

Watershed	Generic State Designated Uses of Water Bodies in the Watershed	
Beaver Creek	South Dakota,	Fisheries Fish and Wildlife Propagation Recreation Agriculture Aesthetics
	Wyoming	Fisheries Fish and Wildlife Propagation Drinking Water Recreation Agriculture Industrial Aesthetics
Upper Belle Fourche River and Tributaries	Wyoming	Fisheries Fish and Wildlife Propagation Drinking Water Recreation Agriculture Industrial Aesthetics
Lower Belle Fourche River and Tributaries	South Dakota	Fisheries Fish and Wildlife Propagation Recreation Agriculture Aesthetics
	Wyoming	Fisheries Fish and Wildlife Propagation Drinking Water Recreation Agriculture Industrial Aesthetics
Redwater River and Tributaries	South Dakota	Fisheries Fish and Wildlife Propagation Recreation Agriculture Aesthetics
	Wyoming	Fisheries Fish and Wildlife Propagation Drinking Water Recreation Agriculture Industrial Aesthetics

2

3

4

The Nebraska-South Dakota-Wyoming Uranium Milling Region also includes a portion of Sioux County and the Hat Creek watershed. Hat Creek is tributary to the Cheyenne River above

Angostura Reservoir in South Dakota. The average flow of Hat Creek at the gauging station near Edgement, South Dakota is $0.14 \text{ m}^3/\text{s}$ [$5.1 \text{ ft}^3/\text{s}$] (U.S. Geological Survey, 2008a). The only impaired water body reported in the Hat Creek watershed is Meng Lake which has high conductivity and impaired pH (Nebraska Department of Environmental Quality, 2008).

South Dakota and Wyoming

The uranium deposits in the Nebraska-South Dakota-Wyoming Uranium Milling Region of South Dakota and Wyoming occur around the western and northern flanks of the Black Hills. The principal uranium deposits are in Fall River County, South Dakota within the Angostura Reservoir watershed and in Niobrara, Weston and Crook counties in Wyoming (Hart, 1968) within the Angostura Reservoir and Lower Belle Fourche River watersheds. Although Custer, Pennington, and Lawrence counties in South Dakota are included within the Nebraska-South Dakota-Wyoming Uranium Milling Region, uranium deposits are not known to exist in these counties. The primary watersheds in South Dakota and Wyoming that may contain uranium deposits within the Nebraska-South Dakota-Wyoming Uranium Milling Region are listed in Table 3.4-4 along with their generic state designated uses and any known impairments to these uses. Although the Nebraska-South Dakota-Wyoming Uranium Milling Region shown in Figure 3.4-10 includes small portions of additional watersheds on its periphery, these secondary watersheds are not in areas of anticipated uranium milling activities.

The uranium deposits in South Dakota occur within the watersheds of the Cheyenne River upstream of Angostura Reservoir, Beaver Creek, Redwater River, and Lower Belle Fourche River (Figure 3.4-10). Within South Dakota, the Cheyenne River has generic designated uses of fisheries, fish and wildlife propagation, recreation, irrigation, and aesthetics. According to South Dakota Department of Environment and Natural Resources (2008), the Cheyenne River above Angostura Reservoir is impaired due to high salinity from natural salts. The average flow of the Cheyenne River at Edgemont, South Dakota is $1.6 \text{ m}^3/\text{s}$ [$58 \text{ ft}^3/\text{s}$] (U.S. Geological Survey, 2008a). The upland portions of the uranium district are primarily drained by ephemeral and intermittent streams with the exception of the lower reach of Red Canyon Creek which is perennial and fed by springs on the flanks of the Black Hills.

The Beaver Creek watershed includes portions of Custer and Pennington counties in South Dakota and Weston County in Wyoming. The generic designated uses of Beaver Creek and its tributaries are listed in Table 3.4-4. Portions of Beaver Creek and its tributaries within South Dakota are impaired due to elevated temperature, salinity, and turbidity (South Dakota Department of Environment and Natural Resources, 2008). The average flow of Beaver Creek at Mallo Camp, Wyoming is $0.048 \text{ m}^3/\text{s}$ [$1.7 \text{ ft}^3/\text{s}$].

The Upper Belle Fourche watershed is located in Wyoming northwest of the Beaver Creek watershed in Weston and Crook counties. The generic designated uses of the Upper Belle Fourche River and its tributaries are listed in Table 3.4-4. A number of perennial streams flowing from the flanks of the Black Hills, such as Inyan Kara Creek, are also present in this watershed. These streams are fed by springs on the flanks of the Black Hills. Streams in portions of the Upper Belle Fourche watershed are impacted by elevated fecal coliform from unidentified sources (WDEQ, 2006).

The Lower Belle Fourche watershed extends from northeastern Crook County in Wyoming (downstream of the Upper Belle Fourche watershed) into Butte, Meade, and Lawrence counties in South Dakota. The designated uses of the Lower Belle Fourche watershed and some of its tributaries are impacted by elevated temperature, salinity, turbidity, and fecal coliform (South

Description of the Affected Environment

Dakota Department of Environment and Natural Resources, 2008). The elevated salinity, turbidity, and fecal coliform are from agricultural livestock grazing activities. Some of the tributaries to the Belle Fourche River drain historical mining districts and are impacted by metals and acidity due to mine drainage. The average flow of the Belle Fourche River at the Wyoming-South Dakota state line is 1.4 m³/s [49 ft³/s] (U.S. Geological Survey, 2008a).

The Redwater River watershed straddles the Wyoming-South Dakota state line between the upper and lower Belle Fourche watersheds (Figure 3.4-10). The generic designated uses of the Redwater River and its tributaries are listed in Table 3.4-4. The average flow of the Redwater River at the gaging station above Belle Fourche, South Dakota is 4.2 m³/s [148 ft³/s] (U.S. Geological Survey, 2008a). Water bodies in this watershed are not listed as impaired.

3.4.4.2 Wetlands and Waters of the United States

Wetland areas found in this region are consistent with those found in the Wyoming East Uranium Milling Region (Section 3.3.4.2). Waters of the United States and special aquatic sites that include wetlands would be expected to be identified and the impact delineated upon individual site selection. Based on impacts and consultation with each area, appropriate permits would be obtained from the local USACE district. Section 401 state water quality certification is required for work in Waters of the United States. Within Wyoming, the state of Wyoming regulates isolated wetlands and waters. Cumulative total project impacts greater than 0.4 ha [1 acre] require a general permit for wetland mitigation by WDEQ. Within Nebraska, waters of the state are under the authority of the Nebraska Department of Environmental Quality. Isolated wetlands are included in Title 117, Nebraska Surface Water Quality Standards. No permitting mechanism is in place to authorize projects in isolated waters; however, state water quality standards apply.

3.4.4.3 Groundwater

Groundwater resources in the Nebraska-South Dakota-Wyoming Uranium Milling Region are part of regional aquifer systems that extend well beyond the areas of uranium milling interest in this part of Nebraska, South Dakota, and Wyoming. Uranium bearing aquifers exist within these regional aquifer systems in the Nebraska-South Dakota-Wyoming Uranium Milling Region. This section provides a general overview of the regional aquifer systems to provide context for a more focused discussion of the uranium bearing aquifers in the Nebraska-South Dakota-Wyoming Uranium Milling Region, including hydrologic characteristics, level of confinement, groundwater quality, water uses, and important surrounding aquifers.

3.4.4.3.1 Regional Aquifer Systems

Major regional aquifers in the Nebraska-South Dakota-Wyoming Uranium Milling Region include the Northern Great Plains aquifer system (Whitehead, 1996) and the High Plains aquifer system (Miller and Appel, 1997).

Northern Great Plain Aquifer System (underlying South Dakota). The Northern Great Plains aquifer system underlies most of South Dakota section of the Nebraska-South Dakota-Wyoming Uranium Milling Region (Whitehead, 1996). The Upper Cretaceous aquifers (important for uranium mineralization and water supplies) and the Paleozoic aquifers (important only for water supplies) of the Northern Great Plains aquifer system are the most extensive aquifers in the South Dakota section of the Nebraska-South Dakota-Wyoming Uranium Milling Region.

Groundwater in the upper Cretaceous aquifers (including minor aquifers in the region) contains less than 3,000 mg/L [3,000 ppm] dissolved solids except for small areas in South Dakota where concentrations are as large as 10,000 mg/L [10,000 ppm]. Water with dissolved-solids concentrations of less than 1,000 mg/L [1,000 ppm] is near the Black Hills Uplift (in west South Dakota) and in smaller areas near the boundaries of the aquifers. Groundwater from the upper Cretaceous aquifers provides domestic and livestock-watering supplies as well as several small communities in northwestern South Dakota.

The lower Cretaceous aquifers are composed of several sandstones. The principal water-yielding units are the Newcastle Sandstone (equivalent to the Dakota Sandstone) and the Inyan Kara Group in the Williston Basin. The Newcastle Sandstone is only a few tens of feet thick where it crops out on the flanks of the Black Hills Uplift, but its subsurface equivalent, the Dakota Sandstone, is more than 122 m [400 ft] thick in southeastern South Dakota. In many places, the Newcastle Sandstone is separated from the underlying Inyan Kara Group through the Skull Creek Shale. The Inyan Kara Group merges eastward into the lower part of the Dakota Sandstone in South Dakota.

The Lower Cretaceous aquifers are confined except at outcrop areas that encircle structural uplifts, such as the Black Hills Uplift and the Bighorn Mountains. In South Dakota, the lower Cretaceous aquifers are overlain by poorly permeable till and glacial-lake deposits, and the aquifers behave like a confined to semiconfined aquifer. The regional groundwater flow direction is northeastward from aquifer recharge areas at high altitudes to discharge areas. Although the groundwater in the lower Cretaceous aquifers is slightly saline in most of South Dakota, the aquifers are the principal source of water for livestock watering and domestic use. The water is very saline or a brine in the deep parts of the Williston Basin.

The upper Paleozoic aquifers consist primarily of the Madison Limestone, which is called the Madison Group in the Williston Basin. The Tensleep Sandstone in the western parts of the Powder River Basin and sandstone beds of the Minnelusa Formation in the Williston Basin and the eastern part of the Powder River Basin are treated as separated aquifers at the regional scale. The Pennsylvanian sandstones are not usually considered to be a principal aquifer. The Madison Limestone exhibits karst features in outcrop areas of the Madison in western South Dakota where large springs originate from solution conduits. In the upper Paleozoic aquifers, the regional groundwater flow direction is northeastward from recharge areas near structural uplifts close to the southern and western limits of the aquifer system. Withdrawal of the oil and gas from the hydrocarbon reservoir have resulted in water leaking downward from the upper Paleozoic aquifers through confining units into deeper permeable zones. Groundwater in the upper Paleozoic aquifers is fresh only in small zones near recharge areas, including the area of freshwater encircling the Black Hills Uplift in western South Dakota. The water becomes slightly saline to saline away from the recharge areas into the Williston Basin. Due to the upward leakage of the mineralized water from the upper Paleozoic aquifers into upper Cretaceous aquifers in central South Dakota, the groundwater becomes saline in shallower aquifers.

Lower Paleozoic aquifers are deeply buried for the most part. They consist of sandstone and carbonate rocks. There are great uncertainties in water yield characteristics of these aquifers at the regional scale. The regional groundwater flow direction is northeastward. Lower Paleozoic aquifers contain fresh water only in a small area near the Black Hills Uplift, but contains slightly saline to moderately saline groundwater throughout the southern one-half of their extent. In a large area in central South Dakota, some of the slightly saline water in the Lower Paleozoic aquifers leaks upward into shallower aquifers.

High Plains Aquifer System (underlying Nebraska). The High Plains aquifer underlies the southernmost part of Nebraska-South Dakota-Wyoming Uranium Milling Region. The High Plains aquifer is the principal source of groundwater for the High Plains region. The High Plains aquifer is unconfined for the most part. The water table is usually less than 61 m [200 ft] below the land surface in western Nebraska. However, the water table is between 61 and 91 m [200 and 300 ft] below the land surface in parts of western Nebraska. The regional groundwater flow direction is from west to east at an average velocity of 0.3 m/day [1 ft/day]. The saturated thickness of the High Plains aquifer ranged from 0 to approximately 305 m [0 to 1,000 ft] in 1980 with an average saturated thickness of 104 m [340 ft]. The average specific yield for entire aquifer is 15 percent. Recharge to the aquifer includes precipitation infiltrating through dune sands in western Nebraska, infiltration locally from streams and canals, by a small quantity of water moving upward from the underlying bedrock. The rates of recharge are highly variable and range from about 0.3 to 20 percent of the average annual precipitation. Discharge from the aquifer includes water losses to springs, seeps, and streams, evapotranspiration, minor water losses to bedrocks, and withdrawals mostly for irrigation.

The High Plains aquifer consists of all or parts of several geologic units of Quaternary and Tertiary age. Clay to gravel size unconsolidated deposits of Quaternary age overlie the Ogallala Formation. These unconsolidated deposits are considered to be part of the High Plains aquifer, if they are saturated as in southeastern Nebraska. The High Plains aquifer is locally confined above by thick loess that consists mostly of silt and clay sized materials. Highly porous dune sands of Quaternary age, where they are saturated, are also considered to be part of the aquifer (e.g., in west-central Nebraska) and recharges the High Plains aquifers.

The Ogallala Formation is underlain by the Arikaree Group. The Arikaree Group, which is composed of massive sandstone, overlies the Brule Formation. The maximum thickness of the Arikaree Group is about 305 m [1,000 ft] in western Nebraska. The Oligocene-aged Brule Formation of Oligocene, which is the upper unit of the White River Group, underlies much of western Nebraska. It is predominantly composed of massive siltstone and sandstone and is considered to be an aquifer only where it is fractured or it contains solution openings.

In large parts of Nebraska, the High Plains aquifer is underlain by upper Cretaceous rocks that primarily consist of shale, chalk, limestone, and sandstone. Only the chalk, where it is fractured or contains solution openings, yields enough water for irrigation. The Chadron Formation, part of the White River Group, directly underlies the High Plains aquifer in most of western Nebraska. It is predominantly composed of clay and silt units with minimal permeability.

In parts of western Nebraska, the High Plains aquifer is underlain by Jurassic- and Triassic-age rocks that primarily consist of shale and sandstone. The Jurassic and Triassic age rocks generally have low permeability, but some sandstone beds are locally permeable enough to yield water. In other areas, the High Plains aquifer is underlain by Tertiary and Permian rocks that predominantly consist of red shale, siltstone, sandstone, gypsum, anhydrite, and dolomite and locally include limestone and halite (rock salt) as beds or disseminated grains.

During 1990, about 17 million L/day [4.6 million gal/day] groundwater was pumped from the High Plains aquifer, mostly (97 percent) for agricultural purposes. The potential water yield from wells in most of Nebraska is typically greater than 4.1 million L/day [1.1 million gal/day], although the water yield varies with the geologic formation tapped. For example, water yields from the Brule Formation are typically less than 1.6 million L/day [430,000 million gal/day]. Water yields from the Arikaree Group are not usually large, but locally in Western Nebraska are as large as 1.9 million L/day [500,000 million gal/day]. The water yields from the Brule

1 Formation and the Arikaree Group are relatively larger where these rocks have secondary
2 fractures. Water yields from the Ogallala Formation are 5.5 million L/day [1.4 million gal/day] in
3 many parts of Nebraska.

4
5 In most of Nebraska, dissolved-solids concentrations in the High Plains aquifer are less than
6 500 mg/L [500 ppm], but locally exceed 1,000 mg/L [1,000 ppm] {the limit of dissolved solids
7 recommended by the EPA for drinking water is 500 mg/L [500 ppm]}. Sodium concentrations in
8 the High Plains aquifer are less than 25 mg/L [25 ppm] in most of Nebraska. However,
9 excessive fluoride concentrations are a widespread problem in the High Plains aquifer. High
10 fluoride concentrations in the range of {2–8 mg/L [2–8 ppm]} are reported for the High Plains
11 aquifer where the aquifer contains volcanic ash deposits or it is underlain by rocks of
12 Cretaceous age.

13
14 The unconfined nature of the High Plains aquifer system along with the shallow water table
15 makes the aquifer vulnerable to contamination by fertilizers and organic pesticides. Elevated
16 concentrations of sodium, alkalinity, nitrate, and triazine (a herbicide) have been found in the
17 aquifer in Nebraska. For example, during 1984–1985, nearly 33 percent of well samples in
18 Nebraska showed measurable concentrations {greater than 0.04 µg/L [0.04 ppb]} of the
19 herbicide atrazine (Whitehead, 1996).

20 21 3.4.4.3.2 Aquifer Systems in the Vicinity of Uranium Milling Sites

22
23 An underlying hydrogeological system in past and current areas of uranium milling interest in
24 the Nebraska section of the Nebraska-South Dakota-Wyoming Uranium Milling Region consists
25 of a thick sequence of primarily sandstone and also limestone aquifers typically separated by
26 shale aquitards. Uranium-bearing sandstone aquifers in the Inyan Kara Group at the potential
27 ISL sites are used for local irrigation water supplies.

28
29 Areas of uranium milling interest in the South Dakota section of the Nebraska-South Dakota-
30 Wyoming Uranium Milling Region are underlain by water-bearing layers including, from
31 shallowest to deepest, the alluvial aquifers, the Newcastle sandstone (equivalent to the Muddy
32 Sandstone), the sandstone aquifers in the Inyan Kara Group, the Morrision Formation, the
33 Sundance Formation, the Spearfish Formation, the Minnekahta Limestone, the Minnelusa
34 Formation, the Madison Formation, and the Deadwood Formation. Among these aquifers, the
35 Inyan Kara Group, the Minnekahta Limestone, the Minnelusa Formation, the Madison
36 Formation, and the Deadwood Formation contain important aquifers for water supplies. The
37 rest of the water-bearing units in the region are pumped for limited local water uses (Williamson
38 and Carter, 2001).

39
40 An underlying hydrogeological system in past and current areas of uranium milling interest in
41 the Nebraska section of the Nebraska-South Dakota-Wyoming Uranium Milling Region consists
42 of a thick sequence of primarily sandstone and also limestone aquifers typically separated by
43 shale aquitards.

44
45 At the Crow Butte ISL sites in Nebraska, only the Basal Chadron sandstone is considered to be
46 an aquifer (NRC, 1998). The Arikaree and Brule Formations are not considered to be important
47 aquifers for water supplies in this region (Miller and Appel, 1997; NRC, 1998).

3.4.4.3.3 Uranium-Bearing Aquifers

In the South Dakota section of the Nebraska-South Dakota-Wyoming Uranium Milling Region, the sandstone aquifers in the Inyan Kara Group are important aquifers for uranium mineralization (Driscoll et al., 2002). In this region, uranium may have been introduced into the Inyan Kara Group through upward leakage of uranium-rich water from the Minnelusa aquifer (Gott, et al., 1974). In the Nebraska section of the Nebraska-South Dakota-Wyoming Uranium Milling Region, the Basal Chadron sandstone aquifer (in the Chadron Formation) hosts uranium mineralization (NRC, 1998).

For ISL operations to begin, portions of the uranium-bearing sandstone aquifers in the Inyan Kara Group and the Basal Chadron Sandstone of aquifer the Nebraska-South Dakota-Wyoming Uranium Milling Region would need to be exempted by the appropriate EPA- or state-administered underground injection program (Section 1.7.2.1).

Hydrogeological characteristics: In the South Dakota section of the Nebraska-South Dakota-Wyoming Uranium Milling Region, the Inyan Kara sandstone aquifers are typically confined except at outcrop areas. Transmissivity of the Inyan Kara aquifer ranges from 0.08–560 m²/day [0.8 - 6,000 ft²/day]. For ISL operations to be practical, the hydraulic conductivity of the production aquifer must be large enough to allow reasonable water flow from injection to production wells. Hence, the portions of the Inyan Kara aquifer with low hydraulic conductivities may not be readily amenable to uranium recovery using ISL techniques. The storage coefficient is in the range of 2.5×10^{-5} – 1.0×10^{-4} (Driscoll et al., 2002) indicating the confined nature of the production aquifer (typical storage coefficients for confined aquifers range from 10^{-5} – 10^{-3} (Driscoll et al., 1986; p.68)).

In the Nebraska section of the Nebraska-South Dakota-Wyoming Uranium Milling region, the Basal Chadron sandstone aquifer is confined by a thick sequence of aquitards. Transmissivity of the Basal Chadron sandstone aquifer ranges from 30 to 45 m²/day [350 to 480 ft²/day] and the average aquifer storage coefficient is in the range of 1.3×10^{-5} – 8.4×10^{-4} (NRC, 1998), indicating the confined nature of the production aquifer (typical storage coefficients for confined aquifers range from 10^{-5} – 10^{-3} (Driscoll, 1986; p.68)).

Level of confinement: The production aquifer is typically confined in the Nebraska-South Dakota-Wyoming Uranium Milling. The thickness of the confinement varies spatially.

In South Dakota, the Inyan Kara Group is generally confined by several thick shale layers, except in the outcrop area around structural uplifts, such as the Black Hills. The Inyan Kara Group is confined above by the Skull Creek Shale with a thickness of 46-80 m [150-270 ft]. The Skull Creek Shale is confined above by the regionally continuous Pierre Shale unit with a thickness of 1,220 m [4,000 ft] in the Black Hills area. The Inyan Kara Group is hydraulically separated from the underlying Minnekahta limestone by low permeability units including, from shallowest to deepest, the Morrison Formation, the Sundance Formation, and the Spearfish Formation. The total thickness of these low permeability layer varies from 190 to 450 m [625 to 1,470 ft] at the Black Hills. Thus, except at the outcrop areas, the sandstone aquifers in the Inyan Kara Group are confined above and below by thick confining units in the Nebraska-South Dakota-Wyoming Uranium Milling. A vertical hydraulic conductivity of 0.4×10^{-6} m/day [1.3×10^{-6} ft/day] for the Skull Creek Shale and 1.5×10^{-8} – 1.5×10^{-4} m/day [5×10^{-8} – 5×10^{-4} ft/day] for the Pierre Shale is estimated in South Dakota (Kansas Geological Survey, 1991).

In Nebraska, the ore-bearing aquifer is confined below by the Pierre shale with an average thickness of 365 m [1,200 ft] and a vertical hydraulic conductivity of 3.4×10^{-11} to 3.6×10^{-12} m/s [11.2×10^{-11} to 11.8×10^{-12} ft/s]. The upper confinement unit is composed of a red clay bed up to 3–8 m [10–25 ft] thick with a vertical hydraulic conductivity of 3×10^{-8} to 2×10^{-7} m/day [1×10^{-7} to 7×10^{-7} ft/day]. The red clay bed is overlain by another thick confining layer (the Middle Chadron) with an average thickness of 95–100 m [315–325 ft]. The thickness of the upper confining unit is about 60–90 m [200–300 ft] in the permit area. Aquifer testing indicates that movement of lixiviant would be vertically contained by the confining units and horizontally captured in the production zone in the Crow Butte region (NRC, 1998).

Groundwater quality: Water from the Inyan Kara aquifer in South Dakota is locally fresh to slightly saline. However, generally high concentrations of dissolved solids, iron, sulfate, and manganese may hamper the use of water from Inyan Kara aquifer. Hard water from wells located on or near the outcrop may require special treatment. Suitability for irrigation may be affected by high specific conductance and sodium adsorption ratio (the ratio of the sodium (detrimental element) concentration to the combined concentration of calcium and magnesium (beneficial elements)). Almost 18 percent of samples collected from the Inyan Kara aquifer exceed the maximum concentration level for combined radium-226 and radium-228. About 4 percent of these samples exceed the maximum concentration level for uranium. The uranium and radium-226 concentrations ranged from 0.1 to 109 ppm and 7.4×10^{-3} –1.59 Bq/L [0.2–43 pCi/L] in the Inyan Kara aquifer, respectively. In the southern Black Hills, radium-226 and uranium concentrations may preclude use of untreated water from Inyan Kara aquifer for drinking (Williamson and Carter, 2001).

Based on baseline (pre-operational) water quality data, the Basal Chadron Sandstone is generally of good quality (with the total uranium less than 3.7×10^{-4} – 8.9×10^{-2} Bq/L [0.01–2.40 pCi/L] and the total conductivity in the range of 1,500–2,500 mhos). The State of Nebraska Department of Environmental Quality defines the Basal Chadron sandstone as an underground source of drinking water (NRC, 1998). However, in the vicinity of the mineralized zone, uranium and radium concentrations are elevated. Radium-226 levels range from 3.7×10^{-3} –22.9 Bq/L [0.1–619 pCi/L], which exceeds the 5 pCi/L EPA primary drinking water standard. As a result, water drawn from Chadron sandstone is not considered potable near the mineralization zone (NRC, 1998).

Current groundwater uses: Groundwater from Inyan Kara aquifer is typically pumped for local irrigation. Groundwater from the Basal Chadron Sandstone is pumped for agricultural and domestic uses.

3.4.4.3.4 Other Important Surrounding Aquifers for Water Supply

The major aquifers in the hydrologic setting of the Black Hill area all underlie the Inyan Kara Group. The major aquifers include, from shallowest to deepest, the Minnekahta Limestone, the Minnelusa Formation, the Madison Formation, and the Deadwood Formation. These aquifers are separated by relatively impermeable layers, but they are (including the Inyan Kara Group) collectively confined by the underlying Precambrian basement rocks and the overlying the Skull Creek and the Pierre Shales. These aquifers are used extensively for water supplies in the region (Williamson and Carter, 2001). The average saturated thicknesses of the the Minnekahta Limestone, the Minnelusa Formation, the Madison Formation, and the Deadwood Formation are 15 m [50 ft], 224 m [736 ft], 159 m [521 ft], and 152 m [500 ft], respectively. The aquifer transmissivity for the Minnelusa Formation, the Madison Formation, and the Deadwood Formation are estimated to be 2.8–28 m²/day [30–300 ft²/day], 9.2×10^{-4} –5,000 m²/day [0.01–

Description of the Affected Environment

54,000 ft²/day], and 23–93 m²/day [250–1,000 ft²/day], respectively. The storage coefficient for the Minnelusa Formation and the Madison Formation are estimated to be 6.6×10^{-5} – 2.0×10^{-4} and 1.12×10^{-6} –0.002 (Driscoll et al., 2002). At the Crow Butte ISL sites in Nebraska, only the Basal Chadron sandstone is considered to be an aquifer (NRC, 1998).

3.4.5 Ecology

3.4.5.1 Nebraska-South Dakota-Wyoming Uranium Milling Region Flora

According to the EPA, the identified ecoregions in the Nebraska-South Dakota-Wyoming Uranium Milling Region primarily consist of Middle Rockies, Northwestern Great Plains, Western High Plains, and the Nebraska Sand Hills ecoregions (Figure 3.4-11). Uranium districts are located in sub-ecoregions including the Black Hills Foothills, Sagebrush Steppe, the Pine Ridge Escarpment, and the Powder River Basin.

The Middle Rockies ecoregion is discussed in the Wyoming West region (section 3.2.5).




Figure 3.4-11. Ecoregions for the Nebraska-South Dakota-Wyoming Uranium Milling Region

1 The Black Hills Foothills ecoregion is composed of the Hogback Ridge and the Red Valley. The
2 Hogback Ridge forms a ring of foot hills surrounding the Black Hills. The Red Valley encircles
3 most of the Black Hills dome and acts as a buffer between the Hogback Ridge. Natural
4 vegetation within this region includes ponderosa pine woodlands and open savannas with an
5 understory of western wheat grass, needle-and-thread grass, little bluestem, blue grama, buffalo
6 grass (*Hierochloe odorata*), and leadplant. In addition, some burr oak is found in the north and
7 Rocky Mountain juniper occurs in the south (Chapman, et al., 2004).

8
9 The Black Hills Plateau ecoregion is a relatively flat, elevated expanse, with broad ridges and
10 entrenched canyons, covering the mid-elevation slopes of the Black Hills. The Black Hills, a
11 mountainous outlier in the Great Plains, have a highly diverse vegetative cover, with an overlap
12 of eastern, boreal, and Rocky Mountain species. The dominate tree spies found in the region is
13 the ponderosa pine, however, it blends with eastern boxelder, burr oak, boreal paper birch.
14 White spruce and sedges can be found in moist areas. The understory includes grasses like
15 little bluestem and timber oatgrass (*Danthonia intermedia*) and shrubs such as juniper,
16 snowberry, bearberry, and buffaloberry (*Shepherdia argentea*) (Chapman, et al., 2004).

17
18 The Black Hills Core Highlands ecoregion includes the higher portions of the limestone plateau
19 above 1,500 m [5,000 ft] and the granitic intrusions that form the major peaks to elevations
20 greater than 2,130 m [7,000 ft]. Due to the high elevation, temperature, and high rainfall boreal
21 species such as white spurce, quaking aspen, and paper bitch can be found on the northern
22 slopes and moist canyons. Ponderosa pine forests interspersed with high meadows are
23 predominant in the region. Understory species include sedges in moist areas, bearded
24 wheatgrass, oatgrass, brone grass, common juniper, snowberry, Oregon grass, bearberry, and
25 iris (Chapman, et al., 2004)

26
27 The Northwestern Great Plains is discussed in Section 3.3.5.1.

28
29 The Montana Central Grassland ecoregion is found mostly in Montana with only a small area
30 continuing into northern Wyoming. The dominate vegetation within this region is a mixed grass
31 prairie comprised of blue gramma, western wheatgrass, june grass, Sandberg bluegrass,
32 needle-and thread grass, rabbit bush, fringed sage, and grama-needlegrass-wheatgrass. The
33 shrub or woodland component found in other ecoregions (Sagebrush Steppe) is absent
34 (Chapman, et al., 2004).

35
36 The Sagebrush Steppe ecoregion is found in Montana and in the Dakotas with only a small area
37 extending into Wyoming. Vegetation types in this region consist of big sagebrush, Nuttall
38 saltbush (*Atriplex nuttallii*), and short grass prairie. The sparse sagebrush communities consist
39 of dusky gray sagebrush (*Artemisia arbuscula* ssp. *Arbuscula*), dwarf sage (*Artemisia*
40 *columbiensis*), and big sagebrush. Prairie vegetation that can be found include western
41 wheatgrass, green needlegrass, blue grama, Sandberg bluegrass, junegrass, rabbit brush,
42 fringe sage, and buffalograss. The shrub vegetation of this ecoregion is transitional between
43 the grasslands of the Montana Central Grassland and the woodland of the Pine Scoria Hills
44 (Bryce, 1996)

45
46 The Semiarid Pierre Shale Plains relatively treeless consisting of rolling hills and grasslands.
47 This is an arid region with rainfall between 38 to 43 cm [15 to 17 in] annually (Bryce, 1996). The
48 natural mixed-grass prairies of the region include shortgrass species, such as buffalograss,
49 western wheatgrass, bluebunch wheatgrass, needle-and-thread grass, blue gramma, and
50 sandberg bluegrass. This ecoregion the sagebrush component found in the neighboring
51 Sagebrush Steppe (Chapman, et al., 2004).

Description of the Affected Environment

The Powder River Basin and Pine Scoria Hills ecoregions are discussed in Section 3.3.5.1.

The White River Badlands in Nebraska border the northern edges of the Pine Ridge escarpment and are southern outliers of a more extensive area in South Dakota. The landscape is broken by grass-covered, perched "sod tables" that may be grazed or tilled typical native vegetation found in this region consists of silver sagebrush, western wheatgrass saltbush, and rabbitbrush (Chapman, et al., 2001).

Western High Plains

The Pine Ridge Escarpment forms the boundary between the Missouri Plateau to the north and the High Plains to the south. This escarpment consists of a Ponderosa pine woodland composed of Rocky Mountain juniper, western soapberry, skunkbush sumac, choke cherry (*Prunus virginiana*), and Arkansas rose (*Rosa arkansana*). The vegetation found in the mixed-grass prairies of the region consists of little bluestem, western wheatgrass, preaires and reed, needle-and-thread grass, blue grama, and threadleaf sedges in moist areas (Chapman, et al., 2001).

The Pine Bluffs and Hills ecoregion is discussed in Section 3.3.5.1.

The Sandy and Silty Tablelands ecoregion is discussed in Section 3.3.5.1.

The Flat to Rolling Cropland ecoregion has extensive drylands farming, irrigated crops, and rangelands throughout this region. Winter wheat, grain sorghum, corn, and alfalfa are the main cash crops, with smaller acreages in forage crops consisting of grain (Chapman, et al., 2001).

The Dense Clay Prairie differs from the surrounding ecoregions in its relative lack of vegetative cover. The grassland in this ecoregion is missing its short- and mid-level layers, however it does include tall grasses comprised mostly of western wheatgrass are found in this ecoregion. Little to no woodlands are found along waterways (Bryce 1996).

Nebraska Sand Hills Ecoregions

The Nebraska Sand Hills consist of one of the most distinct and homogeneous ecoregions in North America. One of the largest areas of grass stabilized sand dunes in the world, this region is generally devoid of cropland agriculture, and except for some riparian areas in the north and east, the region is treeless. Numerous lakes and wetlands dot the region and parts of the region are without streams (Chapman, et al., 2001).

The Sand Hills include grass stabilized sand dunes and open sand areas. Dune size, pattern, and alignment generally follow a west to east trending axis, with the larger dune hills in the west having local relief as great as about 120 m [400 ft]. Grasses found in the area consist of prairie sandreed (*Calamovilfa longifolia*), little blue stem, sand blue stem (*Andropogon hallii*), switchgrass (*Panicum virgatum*), sand love grass (*Eragrostis trichodes*), needle-and-thread grass, blue gramma (*Bouteloua gracilis*), and hairy gramma (*Bouteloua hirsuta*) (Chapman, et al., 2001).

The Alkaline Lakes Area is dominated by sand dunes and many scattered alkaline lakes. These lakes are located in what is commonly referred to as the "closed basin area." This area is generally devoid of streams. The high alkalinity around lake restricts wetland vegetation growth with the exception of alkaline tolerant species such as certain alkaline bulrush (*Schoenoplectus*

1 *maritimus*), alkali sacaton (*Sporobolus airoides*) and inland saltgrass (*Distichlis stricta*). Grass
2 species found in the region are similar to those found in the Sand Hills region consisting of
3 prairie sandreed, little blue stem, sand blue stem, switchgrass, sand love grass, needle-and-
4 thread grass, blue gramma, and hairy gramma (Chapman, et al., 2001).

6 **Nebraska-South Dakota-Wyoming Uranium Milling Region Fauna**

8 Animal species that may occur in the Middle/Southern Rockies which include the Black Hills,
9 the Northwest Great Plains/Northern short grasslands, and Western High Plains/Western
10 Short Grasslands have been discussed in the Wyoming East Uranium Milling Region
11 (Section 3.3.5.1). According to the WGFD crucial wintering habitats are found with this region
12 for large game animals and nesting leks for the sage grouse. Figures 3.4-12 to 3.4-18 depict
13 the crucial winters, yearlong areas ranges for large game found in this region. Within this region
14 the Northern Black Hills Uranium District located in the northeastern portion of the region is near
15 the crucial winter/year long area for white tail deer. Sage grouse Leks appear to be located on
16 the western side of the Nebraska-Suth Dakota-Wyoming Uranium Milling Region in the vicinity
17 of the Southern Black Hills Uranium District.

19 A comprehensive listing of habitat types and species that have been surveyed within
20 South Dakota are compiled as part of the South Dakota Gap Analysis Project (South Dakota
21 State University, 2007).

23 According to the Nebraska Game and Parks Commission, Nebraska has approximately 400 bird
24 species, 95 mammal species, and more than 60 reptile and amphibian species.

25 A comprehensive listing of habitat types and species that have been surveyed within Nebraska
26 are compiled as part of the Gap Analysis Project (University of Nebraska, 2007).

28 **3.4.5.2 Aquatic**

30 **Wyoming**

32 As previously discussed there are approximately 49 native fish species found in the watersheds
33 throughout the state of Wyoming. These species are identified in Table 3.2-5. Current
34 conditions of these watersheds found within the Nebraska-South Dakota-Wyoming Uranium
35 Milling Region have been evaluated, and fish species that would benefit from conservation
36 measures within the watersheds found within the Nebraska-South Dakota-Wyoming Uranium
37 Milling Region have been identified. These watersheds include the Little Missouri watershed
38 and the Cheyenne River Watershed.

40 The Little Missouri watershed is composed of numerous creeks such as Prairie and Cottonwood
41 creek and the north fork of the Little Missouri River. This watershed is located in the
42 northwestern portion of the Nebraska-South Dakota-Wyoming Uranium Milling Region in the
43 vicinity of the Northern Black Hills Uranium District. The game fish habitat in the watershed is
44 restricted to reservoirs and the stream flow in the Little Missouri River. Limiting conditions
45 include small stream size, periods of low flow, high turbidity and sedimentation. Game fish
46 species found in the watershed include brook trout, black bullhead, channel catfish, large mouth
47 bass, rainbow trout, small mouth bass, and stonecat. Nongame species include brassy
48 minnow, flathead chub, fathead minnow, goldeye, green sun fish, lake chub, longnose dace,
49 shorthead redhorse, sand sucker, western silvery minnow, and white sucker (Wyoming Game
50 and Fish Department, 2007).





Figure 3.4-12. Antelope Wintering Areas for the Nebraska-South Dakota-Wyoming Uranium Milling Region

1



SOURCE: Parturition & Crucial Wintering Area Data Provided by The Wyoming Game & Fish Department - 2005

Figure 3.4-13. Big Horn Wintering Areas for the Nebraska-South Dakota-Wyoming Uranium Milling Region

2





Figure 3.4-14. Elk Wintering Areas for the Nebraska-South Dakota-Wyoming Uranium Milling Region



SOURCE: Parturition & Crucial Wintering Area Data Provided by The Wyoming Game & Fish Department - 2005

Figure 3.4-15. Moose Wintering Areas for the Nebraska-South Dakota-Wyoming Uranium Milling Region

1




Figure 3.4-16. Mule Deer Wintering Areas for the Nebraska-South Dakota-Wyoming Uranium Milling Region

2

1




Figure 3.4-17. Sage Grouse/LEK Nesting Areas for the Nebraska-South Dakota-Wyoming Uranium Milling Region

2

1




Figure 3.4-18. White Tailed Deer Wintering Areas for the Nebraska-South Dakota-Wyoming Uranium Milling Region

2
3

The Cheyenne River Watershed is composed of the Lower Cheyenne River, Upper Cheyenne River, Bear Creek, Upper and Lower Antelope Creek, Little Thunder Creek, Black Thunder, and the Lodgepole Creek. This watershed is located in the central western portion of the Nebraska-South Dakota-Wyoming Uranium Milling Region in the vicinity of the Southern Black Hills Uranium District. The Cheyenne River is a free-flowing prairie stream until it reaches the Angostura reservoir in South Dakota. Most of the tributaries are intermittent with some perennial stream segments. Most game species are limited to small reservoirs and impoundments. Species found in the watershed include game fish such as the black bull head and channel catfish and nongame fish such as the carp, flathead minnow, green sunfish, longnose dace long nose sucker, plains killi fish, river carpsucker, sand shiner, and white sucker (Wyoming Game and Fish Department, 2007).

South Dakota

The major watersheds in South Dakota include the Red Water, Beaver, Middle Cheyenne-Spring, Rapid Creek, Angostura Reservoir watershed, which includes the Cheyenne River. The list of fishes present in the South Dakota is summarized in Table 3.4-5.

The South Dakota Division of Wildlife (2004) indicates that the Angostura Reservoir watershed has an area of approximately 23,570 km² [9,100 mi²]. Primary game fish in the watershed include walleye, channel catfish, smallmouth bass (*Micropterus dolomieu*), gizzard shad (*Dorosoma cepedianum*), largemouth bass, black crappie, and emerald shiner (*Notropis atherinoides*). (South Dakota Game ,Fish, and Parks, 2008)

The Cheyenne River originates in eastern Wyoming flowing on the south side of the Black Hills Uplift in the vicinity of the Southern Black Hills Uranium Districtg. The Cheyenne River Watershed Assessment study area is approximately 4,690 km² [1,811 mi²] in Pennington, Custer, and Fall River Counties in South Dakota. Approximately 45 fish species can be found in the Cheyenne River (South Dakota Game and Fish, 2008).

Nebraska

The White River-Hat Creek Basin is located in northwestern Nebraska above the Niobrara River basin north of the Crow Butte Uranium District. This basin originates in Nebraska and drains in northeast to the confluence with the Missouri River (White River) and the Cheyenne River (Hat Creek) in South Dakota. The basin encompasses approximately 5,450 km² [2,130 mi²]. Key aquatic species identified in the basin are the brown trout, rainbow trout, rainbow trout, and channel catfish (Nebraska Department of Environmental Quality, 2005a).

The Niobrara River Basin located in the vicinity of the Crow Butte Uranium District in northwestern and north-central Nebraska originates in eastern Wyoming. The watershed covers approximately 30,745 km² [11,870 mi²] and has approximately 4,054 km [2,519 mi] of streams. The basin also has watersheds that originate in South Dakota. Streamflow in the basin is a function of surface runoff and groundwater contributions. Major tributaries to the watershed include Ponca Creek, Verdigre Creek, Keya Paha River, Long Pine Creek, Plum Creek, Snake River, and Minnechaduza Creek (Nebraska Department of Environmental Quality, 2005b). Fish species found in the Niobrara watershed region are listed in Table 3.4-6.

Table 3.4-5. Fishes of the Angostura Reservoir, Cheyenne River Watershed*

Common Name	Scientific Name
American Eel	<i>Anguilla rostrata</i>
Banded Killifish	<i>Fundulus diaphanus</i>
Bighead Carp	<i>Aristichthys nobilis</i>
Bigmouth Buffalo	<i>Ictiobus cyprinellus</i>
Bigmouth Shiner	<i>Notropis dorsalis</i>
Black Buffalo	<i>Ictiobus niger</i>
Black Bullhead	<i>Ameiurus melas</i>
Black Crappie	<i>Pomoxis nigromaculatus</i>
Blackchin Shiner	<i>Notropis hederdon</i>
Blacknose Dace	<i>Rhinichthys atratulus</i>
Blacknose Shiner	<i>Notropis hedrolepis</i>
Blackside Darter	<i>Percina maculata</i>
Blackspot Shiner	<i>Notropis atrocaudalis</i>
Blue Catfish	<i>Ictalurus furcatus</i>
Blue Sucker	<i>Cycleptus elongatus</i>
Bluegill	<i>Lepomis macrochirus</i>
Bluegill/Green Sunfish Hybrid	<i>Lepomis macrochirus</i> x <i>L. cyanellus</i>
Bluntnose Minnow	<i>Pimephales notatus</i>
Bowfin	<i>Amia calva</i>
Brassy Minnow	<i>Hybognathus hankinsoni</i>
Brook Silverside	<i>Labidesthes sicculus</i>
Brook Stickleback	<i>Culaea inconstans</i>
Brook Trout	<i>Salvelinus fontinalis</i>
Brown Bullhead	<i>Ameiurus nebulosus</i>
Brown Trout	<i>Salmo trutta</i>
Bullhead Minnow	<i>Pimephales vigilax</i>
Burbot	<i>Lota lota</i>
Central Mudminnow	<i>Umbri limi</i>
Central Stoneroller	<i>Campostoma anomalum</i>
Channel Catfish	<i>Ictalurus punctatus</i>
Chinook Salmon	<i>Oncorhynchus tshawytscha</i>
Coho Salmon	<i>Oncorhynchus kisutch</i>
Common Carp	<i>Cyprinus carpio</i>
Common Shiner	<i>Luxilus cornutus</i>
Creek Chub	<i>Semotilus atromaculatus</i>
Cutthroat Trout	<i>Oncorhynchus clarki</i>
Emerald Shiner	<i>Notropis atherinoides Rafinesque</i>
European Rudd	<i>Scardinius erythrophthalmus</i>
Fathead Minnow	<i>Pimephales promelas</i>
Finescale Dace	<i>Phoxinus neogaeus Cope</i>
Flathead Catfish	<i>Pylodictis olivaris</i>
Flathead Chub	<i>Platygobio gracilis</i>
Freshwater Drum	<i>Aplodinotus grunniens Rafinesque</i>
Gizzard Shad	<i>Dorosoma cepedianum</i>
Golden Redhorse	<i>Moxostoma erythrurum</i>
Golden Shiner	<i>Notemigonus crysoleucas</i>

Table 3.4-5. Fishes of the Angostura Reservoir, Cheyenne River Watershed* (continued)

Common Name	Scientific Name
Goldeye	<i>Hiodon alosoides</i>
Grass Carp	<i>Ctenopharyngodon idella</i>
Greater Redhorse	<i>Moxostoma valenciennesi</i>
Green Sunfish	<i>Lepomis cyanellus</i>
Hornyhead Chub	<i>Nocomis biguttatus</i>
Iowa Darter	<i>Etheostoma exile</i>
Johnny Darter	<i>Etheostoma nigrum</i>
Kokanee Salmon	<i>Oncorhynchus nerka</i>
Lake Chub	<i>Couesius plumbeus</i>
Lake Herring	<i>Coregonus artedii</i>
Lake Sturgeon	<i>Acipenser flavescens Rafinwsque</i>
Lake Trout	<i>Salvelinus namaycush</i>
Lake Whitefish	<i>Coregonus clupeaformis</i>
Largemouth Bass	<i>Micropterus salmoides</i>
Logperch	<i>Percina caprodes</i>
Longnose Dace	<i>Rhinichthys cataractae</i>
Longnose Gar	<i>Lepisosteus osseus</i>
Longnose Sucker	<i>Catostomus catostomus</i>
Mississippi Silvery Minnow	<i>Hybognathus nuchalis</i>
Mooneye	<i>Hiodon tergisus Lesueur</i>
Mottled Sculpin	<i>Cottus bairdi</i>
Mountain Sucker	<i>Catostomus platyrhynchus</i>
Muskellunge	<i>Esox masquinongy</i>
Northern Hog Sucker	<i>Hypentelium nigricans</i>
Northern Pike	<i>Esox lucius</i>
Northern Redbelly Dace	<i>Phoxinus eos</i>
Orangespotted Sunfish	<i>Lepomis humilis</i>
Paddlefish	<i>Polyodon spathula</i>
Pallid Sturgeon	<i>Scaphirhynchus albus</i>
Pearl Dace	<i>Margariscus margarita Cope</i>
Plains Killifish	<i>Fundulus zebrinus</i>
Plains Minnow	<i>Hybognathus placitus</i>
Plains Topminnow	<i>Fundulus sciadicus</i>
Pugnose Shiner	<i>Notropis anogenus</i>
Pumpkinseed	<i>Lepomis gibbosus</i>
Quillback	<i>Carpionodes cyprinus</i>
Rainbow Smelt	<i>Osmerus mordax</i>
Rainbow Trout	<i>Oncorhynchus mykiss</i>
Red Shiner	<i>Cyprinella lutrensis</i>
Redear Sunfish	<i>Lepomis microlophus</i>
Ribbon Shiner	<i>Lythrurus Fumeus</i>
River Carpsucker	<i>Carpionodes carpio</i>
River Darter	<i>Percina shumardi</i>
River Shiner	<i>Notropis blennioides</i>
Rock Bass	<i>Ambloplites rupestris</i>
Rosyface Shiner	<i>Notropis rubellus</i>

Description of the Affected Environment

Sand Shiner	<i>Notropis stramineus</i>
Table 3.4-5. Fishes of the Angostura Reservoir, Cheyenne River Watershed* (continued)	
Common Name	Scientific Name
Sauger	<i>Stizostedion canadense</i>
Saugeye	<i>Stizostedion vitreum</i> x <i>S. canadense</i>
Shorthead Redhorse	<i>Moxostoma macrolepidotum</i>
Shortnose Gar	<i>Lepisosteus platostomus</i>
Shovelnose Sturgeon	<i>Scaphirhynchus platyrhynchus</i>
Sicklefin Chub	<i>Macrhybopsis meeki</i>
Silver Chub	<i>Macrhybopsis storeriana</i>
Silver Lamprey	<i>Ichthyomyzon unicuspis</i>
Silverband Shiner	<i>Notropis shumardi</i>
Skipjack Herring	<i>Alosa chrysochloris</i>
Slender Madtom	<i>Noturus exilis</i> Nelson
Slenderhead Darter	<i>Percina phoxocephala</i>
Smallmouth Bass	<i>Micropterus dolomieu</i>
Smallmouth Buffalo	<i>Ictiobus bubalus</i>
Spotfin Shiner	<i>Cyprinella spiloptera</i>
Spottail Shiner	<i>Notropis hudsonius</i>
Stonecat	<i>Noturus flavus</i>
Sturgeon Chub	<i>Macrhybopsis gelida</i>
Suckermouth Minnow	<i>Phenacobius mirabilis</i>
Tadpole Madtom	<i>Noturus gyrinus</i>
Threadfin Shad	<i>Dorosoma petenense</i>
Tiger Muskie	<i>Esox lucius</i> X <i>E. masquinongy</i>
Topeka Shiner	<i>Notropis topeka</i>
Trout-perch	<i>Percopsis omiscomaycus</i>
Walleye	<i>Stizostedion vitreum</i>
Western Silvery Minnow	<i>Hybognathus argyritis</i>
White Bass	<i>Morone chrysops</i>
White Crappie	<i>Pomoxis annularis</i>
White Perch	<i>Morone americana</i>
White Sucker	<i>Catostomus commersoni</i>
Wiper (hybrid)	<i>Morone saxatilis</i>
Yellow Bullhead	<i>Ameiurus natalis</i>
Yellow Perch	<i>Perca flavescens</i>
*South Dakota Department of Game, Fish, and Parks. "Fishing in South Dakota." Pierre, South Dakota: South Dakota Game, Fish, and Parks. 2008 < www.sdgamefish.com/Wildlife/fishing > (15 February 2008)..	

1

Table 3.4-6. Fishes of the Niobrara River Watershed*	
Common Name	Scientific Name
Black Crappie	<i>Pomoxis nigromaculatus</i>
Blacknose Shiner	<i>Notropis hedrolepis</i>
Blue Catfish	<i>Ictalurus furcatus</i>
Bluegill	<i>Lepomis macrochirus</i>
Brook Stickleback	<i>Culaea inconstans</i>
Brook Trout	<i>Salvelinus fontinalis</i>
Brown Trout	<i>Salmo trutta</i>
Channel Catfish	<i>Ictalurus punctatus</i>

Table 3.4-6. Fishes of the Niobrara River Watershed* (continued)

Common Name	Scientific Name
Common Shiner	<i>Luxilus cornutus</i>
Finescale Dace	<i>Phoxinus neogaeus</i> Cope
Flathead Catfish	<i>Pylodictis olivaris</i>
Golden Shiner	<i>Notemigonus crysoleucas</i>
Iowa Darter	<i>Etheostoma exile</i>
Johnny Darter	<i>Etheostoma nigrum</i>
Lake Chub	<i>Couesius plumbeus</i>
Lake Sturgeon	<i>Acipenser flavescens</i> Rafinwsque
Largemouth Bass	<i>Micropterus salmoides</i>
Muskellunge	<i>Esox masquinongy</i>
Northern Pike	<i>Esox lucius</i>
Northern Redbelly Dace	<i>Phoxinus eos</i>
Orange Throat Darter	<i>Etheostoma spectabile</i>
Paddlefish	<i>Polyodon spathula</i>
Pallid Sturgeon	<i>Scaphirhynchus albus</i>
Pearl Dace	<i>Margariscus margarita</i> Cope
Pumpkinseed	<i>Lepomis gibbosus</i>
Rainbow Trout	<i>Oncorhynchus mykiss</i>
Redear Sunfish	<i>Lepomis microlophus</i>
Rock Bass	<i>Ambloplites rupestris</i>
Sauger	<i>Stizostedion canadense</i>
Shovelnose Sturgeon	<i>Scaphirhynchus platyrhynchus</i>
Smallmouth Bass	<i>Micropterus dolomieu</i>
Spotted Bass	<i>Micropterus punctulatus</i>
Striped Bass	<i>Morone saxatilis</i>
Sturgeon Chub	<i>Macrhybopsis gelida</i>
Topeka Shiner	<i>Notropis topeka</i>
Walleye	<i>Stizostedion vitreum</i>
White Bass	<i>Morone chrysops</i>
White Crappie	<i>Pomoxis annularis</i>
Yellow Perch	<i>Perca flavescens</i>
*Nebraska Department of Environmental Quality. "Total Maximum Daily Loads for the Niobrara River Basin." Lincoln, Nebraska: Nebraska Department of environmental Quality. December 2005.	

3.4.5.3 Threatened and Endangered Species

Federally listed threatened and endangered species which are known to exist within habitats found within the region include the following:

- Black-Footed Ferret—discussed in Section 3.2.5.3
- Blowout Penstemon—discussed in Section 3.2.5.3
- Interior Least Tern—discussed in Section 3.2.5.3
- Piping Plover—discussed in Section 3.2.5.3
- Pallid Sturgeon—discussed in Section 3.2.5.3
- Ute Ladies' Tresses Orchid—discussed in Section 3.2.5.3
- Western Prairie Fringed Orchid—discussed in Section 3.2.5.3
- Whooping Crane—discussed in Section 3.2.5.3

State listed Threaten and Endangered species for South Dakota, Nebraska, and special status 1 and 2 species of concern for Wyoming that occur within the region include the following.

South Dakota

- American Dipper (*Cinclus mexicanus*), State Threatened—A unique bird of the cold, fast streams in the Black Hills. American Dippers feed on insects found on stream bottoms, swimming underwater to depths of up to 6 m [20 ft] and even walking on the stream bed. Often nests on the underside of bridges over mountain streams (South Dakota Birds and Birding, 2008).
- Osprey (*Pandion haliaetus*), State Threatened—Osprey habitat includes lakes, large rivers and coastal bays. It is adapted to its fish-eating diet with a reversible front toe and spiny nodules under its toes (spicules) to aid in grasping fish captured by plunge-diving feet first. Ospreys nest at the tops of large living or dead trees, on cliffs, on utility poles or on other tall manmade structures. Clutch size ranges from two to four eggs with hatching in about 30 days. Young fly at 44–59 days and are dependent on parents for 6–12 weeks. This species has a worldwide distribution. In North America, the osprey breeds from northern Saskatchewan, Labrador and Newfoundland in Canada, to the Great Lakes states and along the Pacific and Atlantic coasts. In South Dakota, it is a historical nester in the southeastern part of the state and an uncommon migrant. Many summer observations and the first modern (1991) successful osprey nest in the state raise hopes for the future of this species in South Dakota (U.S. Geological Survey, 2008b).
- Swift Fox State Threatened—discussed in Section 3.2.5.3
- Finescale Dace (*Phoxinus neogaeus*) State Threatened—The Finescale Dace ranges widely but populations existing in Wyoming and Nebraska are considered glacial relics. Commonly occurs in the Niobrara River and several sites in Crook County where they are native to the North Fork Cow Creek in the Cheyenne River drainage. Typically occur in cool, boggy lakes and sluggish acidic streams. They are commonly found in lakes and ponds and are often associated with beaver ponds. Considered to be widespread, abundant, and globally secure but are considered threatened in South Dakota and of special concern in North Dakota, Nebraska, and Wyoming. Distribution is believed to be stable at drainage or sub-drainage scale but declining on the site and stream scale (Wyoming Game and Fish Department, 2008).
- Longnose Sucker, State Threatened—The longnose sucker is found in cool, spring-fed creeks where it feeds on the bottom on algae, crustaceans, snails and insect larvae (caddisflies, mayflies, midges). It spawns in lakes or in shallow-flowing streams over gravel, where fry remain until 1–2 weeks old. Longnose suckers do not sexually mature until 4–9 years of age. The longnose is the most widespread sucker species in North America. It is found in Canada and Alaska; south from western Maryland, north to Minnesota, west and north through northern Colorado and through Washington. South Dakota populations are on the edge of its range and are found in the Belle Fourche River drainage north of the Black Hills (U.S. Geological Survey, 2008b).

- Bald Eagle, State Threatened—discussed in Section 3.2.5.3
- Piping Plover, State Threatened—The piping plover is present on breeding grounds from late March through August. It nests on sandbars and sand and gravel beaches with short, sparse vegetation along inland lakes, on natural and dredge islands in rivers, in gravel pits along rivers and on salt-encrusted bare areas of sand, gravel or pebbly mud on interior alkali ponds and lakes. Nests are shallow, scraped depressions, occasionally lined with small pebbles, shells or other material. A clutch of four eggs is usually laid in late May or early June, with hatching in 27–31 days. Both eggs and young are tended by both parents. Piping plovers feed along the water's edge on small insects, crustaceans and mollusks. In South Dakota, the piping plover is a common breeding associate of the endangered interior least tern. Three North American breeding populations of piping plovers are recognized and have the following distributions: the Atlantic Coast from Newfoundland to Virginia; the Great Lakes, excluding the rocky north shores of Lakes Superior and Huron; and the northern Great Plains. The greatest number of piping plovers breed in the northern Great Plains. This breeding population occurs in scattered alkaline wetlands of the northern Great Plains and on the Missouri River and its tributaries in the Dakotas and Nebraska. In South Dakota, nesting occurs primarily on the natural stretches of the Missouri River below the Gavins Point and Fort Randall Dams, although some nesting may occur on tributaries. Piping plovers have also been reported from Bitter and Waubay Lakes in Day County and Horseshoe Lake in Codington County in northeastern South Dakota. This species overwinters along the Atlantic coast from North Carolina to Florida, along the Gulf coast and in the Bahamas and West Indies (U.S. Geological Survey, 2008b).
- Northern River Otter State Threatened—The river otter is found in rivers, ponds, lakes and unpolluted waters in wooded areas. Key habitat components are riparian vegetation, temporary den and resting sites (cavities under tree roots, shrub patches, tall grass) and adequate food. It is active all year, mainly at night. Air trapped in the fur insulates the river otter while underwater, where it can stay for up to four minutes. Long, stiff whiskers to locate prey and good underwater vision aid in hunting success. The river otter is sexually mature at two years, breeding in early spring. The female has two–three pups (range one–six) in a secluded natal den site. Young leave the den at 2 months, are weaned by 3 months, but remain with the female until just prior to the birth of the mother's next litter. It occupies dens built by other animals, log jams and unused human structures. River otters primarily eat fish. Other aquatic foods include frogs, crayfish and turtles, making the river otter a good barometer of water quality. The river otter is distributed throughout North America north of Mexico, except for the extreme southwestern United States. In South Dakota, it has been reported from Hughes County along the Missouri River, with unverified reports from adjacent counties.

Nebraska

- Finescale Dace State Special Concern—discussed previously for South Dakota
- Swift Fox State Endangered—discussed in Section 3.3.5.3
- Ute Ladies' Tresses Orchid, State Endangered—discussed in Section 3.2.5.3
- Whooping Crane State Endangered—discussed in Section 3.3.5.3

Wyoming

- Finescale Dace, Native Species Status 1—discussed previously for South Dakota

Description of the Affected Environment

- Pearl Dace (*Margariscus margarita*) Native Species Status 1—the pearl dace occurs in cool bogs, ponds, lakes, creeks and clear streams. It spawns in the spring in clear water with a weak to moderate current over sand or gravel. This species feeds on invertebrates (insects and zooplankton) and algae (U.S. Geological Survey, 2008b).
- Western Silvery Minnow, Native Species Status 1—discussed in Section 3.2.5.3
- Canda Lynx, Native Species Status 1—discussed in Section 3.2.5.3
- Plains Topminnow Native Species Status 2— discussed in Section 3.2.5.3
- Goldeye (*Hiodon alosoides*), Native Species Status 2—In Wyoming, the goldeye can be found in the Powder, Little Powder and Little Missouri rivers and in Clear and Crazy Woman creeks. It prefers large rivers and their associated backwaters and marshes, or the shallow waters of large lakes and reservoirs. Young goldeye have never been found in Wyoming, it is thought that populations in the northeastern part of the state are maintained by the migration of adult fish seeking spawning grounds (Wyoming Game and Fish Department, 2008).
- Pale Milk Snake (*Lampropeltis triangulum multistrata*), Native Species Status 2—The pale milksnake prefers grasslands, sandhills and scarp woodlands below 1,830 m [6,000 ft] in elevation. It is distributed throughout the northern Great Plains. In Wyoming, it can be found in the eastern counties and the Big Horn Basin (Wyoming Game and Fish Department, 2008).
- Smooth Green Snake, Native Species Status 2— discussed in Section 3.2.5.3
- Yellow-Billed Cuckoo, Native Species Status 2— discussed in Section 3.2.5.3
- Greater Sage Grouse, Native Species Status 2— discussed in Section 3.2.5.3
- Bald Eagle, Native Species Status 2— discussed in Section 3.2.5.3
- Trumpeter Swan Native, Species Status 2— discussed in Section 3.2.5.3
- Fringed Myotis Native Species Status 2— discussed in Section 3.2.5.3
- Long-Eared Myotis, Native Species Status 2— discussed in Section 3.2.5.3
- Long-Legged Myotis Native Species Status 2—discussed in previous regions.
- Pallid Bat, Native Species Status 2— discussed in Section 3.2.5.3
- Spotted Bat, Native Species Status 2— discussed in Section 3.2.5.3
- Townsend's Big-Eared Bat, Native Species Status 2— discussed in Section 3.2.5.3

3.4.6 Meteorology, Climatology, and Air Quality

3.4.6.1 Meteorology and Climatology

The Nebraska-South Dakota-Wyoming Uranium Milling Region contains portions of three states: Wyoming, Nebraska, and South Dakota. This region is characterized by hot summers and cold winters and rapid temperature fluctuations are common. The Rocky Mountains have a great influence on the climate. As air crosses the Rockies from the west much moisture is lost on the windward sides of the mountains and becomes warmer as it descends on the eastern slopes. Table 3.4-7 identifies three climate stations located in the Nebraska-South Dakota-Wyoming Uranium Milling Region. Climate data for these stations are found in the National Climatic Data Center's Climatology of the United States No. 20 Monthly Station Climate Summaries for 1971–2000 (National Climatic Data Center, 2004). This summary contains climate data for 4,273 stations throughout the United States and some territories. Table 3.4-8 contains temperature data for three stations in the Western South Dakota/Nebraska Uranium Milling Region.

Most precipitation in the Nebraska-South Dakota-Wyoming Uranium Milling Region occurs in the spring and summer. Rainstorms, hailstorms, and lightning are most likely to occur in the summer. Heavy rain can accompany thunderstorms and may cause some flooding. This flooding intensifies if these storms coincide with snow pack melting. Table 3.4-8 contains precipitation data for three stations in the Nebraska-South Dakota-Wyoming Uranium

Table 3.4-7. Information on Three Climate Stations in the Nebraska-South Dakota-Wyoming Uranium Milling Region*

Station (Map Number)	County	State	Longitude	Latitude
Colony	Crook	Wyoming	104°11W	44°55N
Newcastle	Weston	Wyoming	104°13W	43°51N
Ardmore 2 N	Fall River	South Dakota	103°39W	43°03N

*National Climatic Data Center. "Climatology of the United States No. 20: Monthly Station Climate Summaries, 1971–2000." Asheville, North Carolina: National Oceanic and Atmospheric Administration. 2004.

Table 3.4-8. Climate Data for Stations in the Nebraska-South Dakota-Wyoming Uranium Milling Region*

		Colony	Newcastle	Ardmore 2 N
Temperature (°C)†	Mean—Annual	8.3	7.9	8.1
	Low—Monthly Mean	–5.3	–5.7	–6.0
	High—Monthly Mean	22.4	22.5	22.5
Precipitation (cm)‡	Mean—Annual	37.8	40.7	43.7
	Low—Monthly Mean	0.9	1.1	1.0
	High—Monthly Mean	6.8	6.5	7.3
Snowfall (cm)	Mean—Annual	93.2	95.5	105
	Low—Monthly Mean	0	0	0
	High—Monthly Mean	19.6	19.8	18.5

*National Climatic Data Center. "Climatology of the United States No. 20: Monthly Station Climate Summaries, 1971–2000." Asheville, North Carolina: National Oceanic and Atmospheric Administration. 2004.

†To convert Celsius (°C) to Fahrenheit (°F), multiply by 1.8 and add 32

‡To convert centimeters (cm) to inches (in), multiply by 0.3937

Description of the Affected Environment

Milling Region. The wettest month varies for the stations identified in Table 3.4-8. May is the wettest month for the Newcastle (Weston County, Wyoming) and Ardmore (Fall River County, South Dakota) stations and June is the wettest month for the Colony (Crook County, Wyoming) station. Based on the snow depth data, the wettest months coincide with melting snow pack (National Climatic Data Center, 2004). Data from National Climatic Data Center's Storm Events Database from 1950 to 2007 indicates that the vast majority of thunderstorms in Crook, Weston, and Fall River Counties occur between May and August with most occurring in July (National Climatic Data Center, 2007).

The mountains typically receive the most snow. Occasionally snow can accumulate to a considerable depth. During snow periods there is often wind that may cause a large proportion to collect in gullies and behind windbreaks. Peak snow fall generally occurs in February and early March. Table 3.4-8 contains snowfall data for three stations in the Nebraska-South Dakota- Wyoming Uranium Milling Region.

The pan evaporation rates for the Western South Dakota/Nebraska Uranium Milling Region range from about 102 - 127 cm [40 to 50 in] (National Weather Service, 1982). Pan evaporation is a technique that measures the evaporation from a metal pan typically 121 cm [48 in] in diameter and 25 cm [10 in] tall. Pan evaporation rates can be used to estimate the evaporation rates of other bodies of water such as lakes or ponds. Pan evaporation rate data are typically available only from May to October. Freezing conditions often prevent collection of quality data during the other part of the year.

3.4.6.2 Air Quality

The air quality general description for the Western South Dakota/Nebraska Uranium Milling Region would be similar to the description in Section 3.2.6 for the Wyoming West Uranium Milling Region. The Nebraska-South Dakota-Wyoming Uranium Milling Region information in Section 3.4.6.2 is limited to the modification, supplementation, or summarization of the Wyoming West Uranium Milling Region information presented in Section 3.2.6.

As described in Section 1.7.2.2, the permitting process is the mechanism used to address air quality. If warranted, permits may set facility air pollutant emission levels, require mitigation measures, or require additional air quality analyses. The Nebraska-South Dakota-Wyoming Uranium Milling Region covers portions of Wyoming, South Dakota, and Nebraska. Except for Indian Country, New Source Review permits in these three states are regulated under the EPA-approved State Implementation Plan except for the Prevention of Significant Deterioration permits in South Dakota, which are regulated by 40 CFR 52.21 (EPA, 2007a). For Indian Country in these three states, the New Source Review permits are regulated under 40 CFR 52.21 (EPA, 2007a).

State Implementation Plans and permit conditions are based in part on federal regulations developed by the EPA. The NAAQS are federal standards that define acceptable ambient air concentrations for six common nonradiological air pollutants: nitrogen oxides, ozone, sulfur oxides, carbon monoxide, lead, and particulates. In June 2005, EPA revoked the 1-hour ozone standard nationwide in all locations except certain Early Action Compact Areas. None of the 1-hour ozone Early Action Compact Areas are in Wyoming, South Dakota, or Nebraska. States may develop standards that are stricter or supplement the NAAQS. Wyoming has a more restrictive annual average standard for sulfur dioxide at $60 \mu\text{g}/\text{m}^3$ [1.6×10^{-6} oz/yd³] and a supplemental $50 \mu\text{g}/\text{m}^3$ [1.3×10^{-6} oz/yd³] PM₁₀ standard with an annual averaging time (Wyoming Department of Environmental Quality, 2006). Nebraska has a $50 \mu\text{g}/\text{m}^3$

[1.3 × 10⁻⁶ oz/yd³] PM₁₀ standard with an annual averaging time (Nebraska Department of Environmental Quality, 2002). South Dakota standards implement NAAQS straightforward (South Dakota Department of Environment and Natural Resources, 2007).

Prevention of Significant Deterioration requirements identify maximum allowable increases in concentrations for particulate matter, sulfur dioxide, and nitrogen dioxide for areas designated as attainment. Different increment levels are identified for different classes of areas and Class I areas have the most stringent requirements.

The Nebraska-South Dakota-Wyoming Uranium Milling Region Air Quality description focuses on two topics: NAAQS attainment status and PSD classifications in the region.

Figure 3.4-19 identifies the counties in and around the Western South Dakota/Nebraska Uranium Milling Region that are partially or entirely designated as nonattainment or maintenance for NAAQS at the time this GEIS was prepared (EPA, 2007b). All of the area within the Nebraska-South Dakota-Wyoming Uranium Milling Region is classified as attainment. Wyoming only has one area that is not in attainment. The City of Sheridan in Sheridan County is designated as nonattainment for PM₁₀. Nebraska only has one area not in attainment. A portion of the city of Omaha in Douglas County is designated as maintenance for lead but this is in eastern Nebraska, about 500 km [311 mi] from the Nebraska-South Dakota-Wyoming Uranium Milling Region. No areas in South Dakota are designated as nonattainment or maintenance. Two counties in southeast Montana are not in attainment. However, the two Montana counties that border the Nebraska-South Dakota-Wyoming Uranium Milling Region are in attainment.

Table 3.4-9 identifies the Prevention of Significant Deterioration Class I areas in Wyoming, South Dakota, Nebraska, and Montana. These areas are shown in Figure 3.4-20. The Nebraska-South Dakota-Wyoming Uranium Milling Region does contain a Class I area for the Wind Cave National Park in South Dakota (40 CFR Part 81).

3.4.7 Noise

The existing ambient noise levels for undeveloped rural and more urban areas in the Nebraska-South Dakota-Wyoming Uranium Milling Region would be similar to those described in Section 3.2.7 for the Wyoming West Uranium Milling Region. This is a large region spanning parts of three different states. The largest community within the region, with a population of about 12,500, is Spearfish, South Dakota in the northeastern portion. Smaller communities with populations from around 1,000 to 6,000 include Sundance and Newcastle, Wyoming, Hot Springs and Custer, South Dakota, and Crawford and Chadron in Dawes County, Nebraska (see Section 3.4.10). Ambient noise levels in these communities would likely be in the range of 45 to about 78 dB (Washington State Department of Transportation, 2006). In addition, the Pine Ridge Indian Reservation is just to the east of the South Dakota/Nebraska Uranium Milling Region.

A number of major highways cross the region, including Interstate 90 in the northern portion and a number of U.S. and state undivided highways. Ambient noise levels near these highways would be similar to or less than those measured at up to 70 dBA for Interstate 80, as the total traffic count and the percentages of heavy truck traffic are less (Wyoming Department of Transportation, 2005; Federal Highway Administration, 2004; see also Section 3.2.7 and 3.4.2).

3.4-50




Figure 3.4-19. Air Quality Attainment Status for Western South Dakota/Nebraska Uranium Milling Region and Surrounding Areas (EPA, 2007b)

Table 3.4-9. U.S. Environmental Protection Agency Class I Prevention of Significant Deterioration Areas in Wyoming, South Dakota, Nebraska, and Montana*

WYOMING Bridger Wilderness Fitzpatrick Wilderness Grand Teton National Park North Absaroka Wilderness Teton Wilderness Washakie Wilderness Yellowstone National Park	MONTANA Anaconda-Pintlar Wilderness Bob Marshall Wilderness Cabinet Mountains Wilderness Gates of the Mountain Wilderness Glacier National Park Medicine Lake Wilderness Mission Mountain Wilderness Red Rock Lakes Wilderness Scapegoat Wilderness Selway-Bitterroot Wilderness U.L. Bend Wilderness Yellowstone National Park
SOUTH DAKOTA Badlands Wilderness Wind Cave National Park	NEBRASKA None

*Modified from Code of Federal Regulations. "Prevention of Significant Air Deterioration of Air Quality."
 Title 40—Protection of the Environment, Part 81. Washington, DC: U.S. Government Printing Office. 2005.

A number of scenic byways through the Black Hills could be more sensitive to noise impacts, but these are located more than 16 km [10 mi] east of the areas of interest for ISL uranium recovery.

For the three uranium districts located in the Nebraska-South Dakota-Wyoming Uranium Milling Region, there are several National Park Service and U.S. Forest Service properties, state parks, and other properties (see Figure 3.4-1) that may be sensitive to noise impacts. Much of this area is protected from extensive development, and the ambient noise levels would be expected to be similar to undeveloped rural areas (up to 38 dB) (DOE, 2007).

Northernmost uranium district (Wyoming)

- Devil's Tower National Monument (Wyoming)
- Black Hills National Forest (Wyoming-South Dakota)


Central uranium district (Wyoming, South Dakota)

- Thunder Basin National Grassland (Wyoming)
- Buffalo Gap National Grassland (South Dakota)

Southern uranium district (Nebraska)

- Oglala National Grassland (Nebraska)
- Nebraska National Forest (Nebraska)
- Fort Robinson State Park (Nebraska)

1



SOUTH DAKOTA - NEBRASKA REGION




Figure 3.4-20. Prevention of Significant Deterioration Class I Areas in the Western South Dakota/Nebraska Uranium Milling Region and Surrounding Areas (40 CFR Part 81)

2

Small communities are located within and near each of the three uranium districts, including Aladdin, Wyoming in the northernmost district, Riverview, Wyoming and Burdock and Edgemont, South Dakota in the central district, and Crawford, Nebraska near the Crow Butte ISL facility in the southern district. In general, these small towns are located 8 km [5 mi] or more from the uranium projects.

3.4.8 Historical and Cultural Resources

Appendix D provides a general overview of historical and cultural resource impact assessment at the federal level. As noted in Section 3.2.8, specific cultural resources in Wyoming, South Dakota, Nebraska, and New Mexico are described at the state level by the responsible state agencies. For the purposes of describing cultural and historical resources for the Nebraska-South Dakota-Wyoming Uranium Milling Region, an overview of Wyoming cultural and historical resources is provided in Section 3.2.8. Cultural and historical resources in South Dakota and Nebraska are described separately in this section (Section 3.4.8).

The South Dakota SHPO is a division of the South Dakota State Historical Society. The director of the South Dakota State Historical Society serves as the state's Historic Preservation Officer. The South Dakota SHPO administers and is responsible for oversight and compliance with the NRHP, compliance and review for Section 106 of the NHPA, Preservation of Historic Property Procedures (South Dakota Codified Law 1-19-11.1), and Traditional Cultural Properties, NAGPRA and archaeological survey through its Archaeology Division as well as compliance with other federal and state historic preservation laws, regulations, and statutes. Their webpage can be found at: <<http://www.sdhistory.org>>. The State of South Dakota also has laws regarding human remains, entitled Cemeteries and Burials (SDCL 1-20-32, Chapter 34-27).

The Nebraska SHPO is a division of the Nebraska State Historical Society. The director of the Nebraska State Historical Society serves as the state's Historic Preservation Officer. The NSHPO administers and is responsible for oversight and compliance with the NRHP, the Nebraska Historic Buildings Survey, compliance and review for Section 106 of the NHPA and Traditional Cultural Properties, NAGPRA and archaeological survey through its Archaeology Division and compliance with other federal and state historic preservation laws, regulations, and statutes. Their webpage can be found at: <<http://www.nebraskahistory.org/histpres>>. The State of Nebraska also has laws regarding human remains, entitled Unmarked Human Burials Sites (Revised Statutes of Nebraska 1989 Supplement Article 12 [12-1201 to 12-1212]) and Human Skeletal Remains or Burial Goods, Prohibited Acts; Penalty (Article 28-1301).

3.4.8.1 Cultural Resources Overview

3.4.8.1.1 Cultural Resources of Western and Southwestern South Dakota

The following provides a brief overview of prehistoric and historical cultures recognized in the central and northern plains region which includes western South Dakota. The dating of cultural periods for the prehistoric period are provided in years before present (BP). Most prehistoric archaeological sites are concentrated along the James, Missouri, White, Cheyenne and Big Sioux river valleys, but can be found along many drainage basins in the state. Figures 3.2-18 and 3.2-19 illustrate the division of the plains into regional subdivisions.

Paleoindian Big Game Hunters (12,000 to 6,500 BP). The earliest well-defined cultural tradition in the central plains region is the Paleoindian. Early humans entered the plains shortly

Description of the Affected Environment

after deglaciation allowed movement onto the central plains sometime after 14,000 BP. A variety of cultures, each defined by the presence of distinctive projectile points, are recognized during the Paleoindian period: Clovis, Goshen, Folsom, Hell Gap-Agate Basin, Cody Complex and Plano. Most post-Clovis Paleoindian sites on the northern and upper central plains are known from bison kill sites. The Clovis culture (12,000 to 10,000 BP) is recognized by a distinctive projectile point style and a subsistence mode heavily reliant on hunting large, now-extinct mammals, notably mammoth and mastodon, which became extinct at the end of the Clovis period. The poorly defined Goshen Complex found at the Jim Pitts site in the Black Hills may be contemporary with Clovis and is technologically similar. The Folsom culture (ca. 10,000 to 8,500 BP) is also known for a distinctive projectile point style. Folsom subsistence is also characterized by reliance on large game, the ancient bison. Folsom sites consist of camp sites and kill sites. The latter tend to be located near cliffs and around water, such as ponds and springs. The Plano, Hell Gap-Agate Basin, and Cody Complex cultures (ca. 8,500 to 6,500 BP) are, in their earliest forms, a continuation of earlier Paleoindian hunting traditions. The distinctive projectile point forms which define these cultural complexes are, in comparison to earlier Clovis and Folsom, much more restricted in geographic distribution. Toward the middle and end of the period encompassing these cultures, however there is a transition in subsistence modes following with the extinction of the ancient bison form to the modern form of bison and ultimately, a transition to Archaic broad-spectrum foraging. Post molds and stone circles suggesting the presence of ephemeral shelters are sometimes found, primarily toward the end of the period.

Archaic Foragers (6,500 to 3,500 BP). The Plains Archaic period represents the continuation of change in subsistence and settlement linked to an increasingly arid environment that occurs in the latter portion of the preceding late Paleoindian cultures. Distinctive Archaic cultures, from early to late, include Mummy Cave, Oxbow, McKean, and Pelican Lake complexes. Kill sites, characteristic of the preceding Paleoindian period are virtually absent. Hunting and gathering wild plant foods is the primary mode of subsistence. Dietary breadth, indicated by increasing diversity and numbers of subsistence items, is believed to expand significantly with more medium and small mammals being hunted and the introduction of seed-bearing plants dietary staples indicated by the introduction of stone seed-grinding implements. Through time, settlement is increasingly tethered to highly productive resource areas and sites tend to become larger and increasingly complex indicating the presence of somewhat more sedentary lifestyles relative to earlier periods. Settlement is focused on river valleys and elevated areas. Artifact styles, principally projectile points, become increasingly diversified suggesting increasing regionalization and cultural differentiation.

Late Prehistoric/Plains Woodland (3,500 to 300 BP). Early in the period, the preceding late Archaic broad-spectrum foraging subsistence and settlement patterns continue with little change. In the Northern Plains, the Besant and Avonlea Complexes continued the Archaic lifestyles virtually unchanged until contact with European and American cultures. Subsistence focused on scheduled small and medium game hunting, gathering plant foods and bison hunting according to a seasonal round. In western South Dakota, a basic hunting and gathering lifestyle differing little from the preceding Late Archaic period predominates. At the very end of the period, some villages located along water courses in western South Dakota may have practiced horticulture, but its contribution to diet among such Northern Plains groups was limited. Food procurement and site location appears to be focused primarily on elevated landforms near larger riverine systems and tributaries with increasing utilization of upland resources later in time. The Late Prehistoric/Plains Woodland of South Dakota is also characterized by the appearance of ceramics late in the period (Avonlea Complex), perhaps introduced from the Eastern Woodland cultural area. The late Avonlea Complex and later Old Woman Complex

1 sites contain artifact types that suggest a high degree of specialization in hunting large, upland
2 game animals, primarily bison.

3
4 In the eastern portions of South Dakota along the Missouri River, seasonal or permanent
5 sedentary villages of various sizes occur. These villages were largely reliant on domesticated
6 plants (corn, beans, and squash). Although horticulture was an important part of the
7 subsistence base, wild plants and game animals formed a substantial part of the diet. Villages
8 were primarily located along major river systems and larger tributaries. Most sites consisted of
9 small clusters of rectangular wattle and daub lodges with a few larger village sites. Storage pits
10 for food and other times are located within the structures. Pottery was diverse with globular jars
11 and decorated exterior rims common.

12
13 In the 1500s to early 1700s A.D., large migrations occurred. The ancestors of the modern
14 Apache, Arapaho, Comanches, and Kiowas migrated southward through western South Dakota
15 in the 1500s and 1600s. The Crow also resided in western South Dakota for a time. The
16 central portion of the state was occupied by the Arika, Mandan, and Cheyenne while the Lakota,
17 Omahas, Poncas, Otos and Ioway occupied the eastern portion of the state.

18
19 **Post-contact Tribes (300 to 100 BP).** The post-contact period on the northern plains is that
20 period after initial contact with Europeans and Americans. Although Euro-American trade goods
21 may have appeared as early as the mid-1600s, the earliest documented contact in the northern
22 and central plains is by Spanish and French explorers in the early 1700s AD. The horse
23 appears to have been introduced at about the same time. The lifeways of the late Avonlea and
24 post-Avonlea/Old Woman nomadic bison-hunting cultural complexes appear to have continued
25 well into the mid to late 1700s AD. At the time of European exploration, Arikara and Mandan
26 farming villages were noted along the Missouri river in central South Dakota. In the 1700s, the
27 Cheyenne moved westward along with the Lakota and displaced the Mandan and Arikara. The
28 Dakota and Nakota moved into eastern South Dakota from Minnesota and displaced the
29 Poncas and the Omaha. By the mid-1800s, the entire state was occupied by nomadic
30 Siouan-speaking tribes, primarily the Santee, Yankton, and Teton.

31
32 **Europeans and Americans (300 to 100 BP).** The earliest European presence in South Dakota
33 was by French explorers of the de la Vérendrye family in 1743. In 1803, the United States
34 completed the purchase of the Louisiana Territory from France. A portion of South Dakota was
35 visited by the Lewis and Clark Expedition in 1804–1806. These early expeditions provide
36 descriptions of varying quality for some of the early historical tribes in the region. In the later
37 1700s and early 1800s more intensive contact and settlement occurred first through
38 missionaries and the fur trade period in the 1830s through the 1860s. The American Fur
39 Company and its fur trading posts located along the Big Sioux, James, Vermillion, Missouri,
40 Cheyenne, White, and Big Stone Lake formed the foundation for later settlements. By the
41 mid-1800s missionary, settler, and military contacts led to increasing conflict with the Siouan
42 tribes of South Dakota. The slowly increasing number of settlers passing through traditional
43 tribal use areas in the mid-1800s led to increasing conflict over time and the establishment of
44 military forts in tribal lands, yet another irritant to tribes.

45
46 Treaties, notably the Fort Laramie Treaty of 1851 were signed with the intent of removing tribes
47 from along the emigrant trails and to allow for the building of trails and forts to protect settlers
48 moving west. Continued conflict resulted in the creation of the Great Sioux Reservation
49 bounded by the Missouri River on the east, the Big Horn Mountains on the west, and the 46th
50 and 43rd parallels to the north and south, respectively. Continued conflict with the U.S. military

over the failure of the government to abide by treaty obligations led to several punitive expeditions to return tribes to reservations. In 1874, General George Armstrong Custer led an expedition to the Black Hills where the presence of gold, previously only rumored, was confirmed. The intense interest by Americans to go to the Black Hills to mine for gold led to numerous treaty violations; the Black Hills region was, by treaty, part of the Sioux reservation. The continued conflict over the Black Hills, along with reduction of the buffalo herds, led to the final military conquest of the Great Sioux Nation and their confinement to small reservations. The Black Hills gold rush led to the rapid settlement of much of South Dakota and the development of towns and cattle ranching.

Ranching, a livelihood well suited to the grassland plains of South Dakota, was practiced by settlers by the early 1870s. The arrival of the railroads (the Milwaukee) led to increased settlement and opened South Dakota to a flood of new settlers, most of them recent European immigrants intent on farming. These early settlers began a period of extensive agriculture throughout the state, mostly around well-watered regions, with many of the new farmers pursuing newly developed dry-land farming techniques. During the Great Depression and the droughts that occurred at the same time led to the abandonment of many farms and the out-migration of a significant portion of South Dakota's population.

3.4.8.1.2 Cultural Resources of Western Nebraska

The following provides a brief overview of prehistoric and historical cultures recognized in the central plains region which includes Nebraska. The dating of cultural periods for the prehistoric period are provided in years before present (BP). Figures 3.2-18 and 3.2-19 illustrate the division of the plains into regional subdivisions.

Paleoindian Big Game Hunters (12,000 to 8,000 BP). The earliest well-defined cultural tradition in the central plains region is the Paleoindian. Early humans entered the plains shortly after deglaciation allowed movement onto the central plains sometime after 14,000 BP. Three cultures are recognized during the Paleoindian period: Clovis, Folsom, and Plano. The Clovis culture (12,000 to 10,000 BP) is recognized by a distinctive projectile point style and a subsistence mode heavily reliant on big-game hunting, notably mammoth and mastodon, which became extinct at the end of the period. The Folsom culture (ca. 10,000 to 8,500 BP) is also known for a distinctive projectile point style. Folsom subsistence is also characterized by reliance on large game, the ancient bison. Folsom sites consist of camp sites and kill sites. The latter tend to be located near cliffs and around water, such as ponds and springs. The Plano culture (ca. 8,500 to 6,500 BP) is, in its earliest form, a continuation of earlier Paleoindian hunting traditions. Toward the end of the period, however there is a transition in subsistence modes with the extinction of the ancient bison to the modern form of bison and a transition to Archaic foragers. Plano sites containing circular rock alignments and post mold circles suggest the presence of structures.

Archaic Foragers (6,500 to 2,000 BP). The Plains Archaic period represents the continuation of change in subsistence and settlement linked to an increasingly arid environment that occurs in the latter portion of the preceding late Paleoindian Plano culture. Kill sites, characteristic of the preceding Paleoindian period are virtually absent. Although hunting and gathering is the only mode of subsistence, dietary breadth, indicated by increasing diversity and numbers of subsistence items, is believed to expand significantly with more medium and small mammals being hunted and the introduction of seed-bearing plants as staples. Through time, settlement is increasingly tethered to highly productive resource areas and sites tend to become larger and increasingly complex indicating the presence of more sedentary lifestyles relative to earlier

periods. Artifact styles, principally projectile points, become increasingly diversified suggesting increasing regionalization and cultural differentiation.

Plains Woodland (2,000 to 1,000 BP). The Plains Woodland period is characterized by largely sedentary lifestyles and a mixed subsistence economy consisting of wild game animals and plants and horticulture utilizing the domesticates, maize and beans. The defining settlement pattern of the Woodland Period consists of earth lodge villages, some of which may have been occupied only seasonally. There is variability in the size of Plains Woodland communities. The communities can be small with as few as two or three structures, to very large (two to three hectares) with numerous contemporary structures. The majority of the larger settlements tended to be located along larger drainages (e.g., Missouri, Republican, Arkansas, and Red rivers) with permanent water and located near abundant biotic and abiotic resources. The Plains Woodland is also characterized by the appearance of ceramics, perhaps introduced from the Eastern Woodland cultural area.

Plains Village (1,000 to 600 BP). The Plains Village period continues the trend toward increasing sedentism and increasing reliance on domesticated plants (corn, beans, and squash). Although horticulture was an important part of the subsistence base, wild plants and game animals formed a substantial part of the Plains Village diet. Villages were primarily located along major river systems and larger tributaries. Most sites, however, consisted of small clusters of rectangular wattle and daub lodge. Storage pits for food and other times are located within the structures. Pottery was diverse with globular jars and decorated exterior rims being common. Small, triangular side- and corner-notched projectile points are common. Early historical Plains Village groups include the Siouan-speaking Omaha, Ponca, Otoe-Missouria, Ioway, and Kansa along with the Caddoan-speaking groups including the Arikara and Pawnee. The Plains Village period is divided into several regional phases and include the St. Helena, Nebraska, Itskari and Smokey Hill phases.

Post-Contact Tribes (400 to 100 BP). The post-contact period on the central plains is that period after initial contact with Europeans and Americans. The earliest documented contact in the central plains is by Spanish and French explorers in the early 1700s AD. Tribes in present include the Caddoan farming villages of the Pawnee and Arikara in eastern Nebraska. Siouan-speaking tribes were the Omaha, Ponca, Otoe-Missouria, Ioway, and Kansa. Both Caddoan and Siouan-speaking groups lived in permanent earth lodge villages, were agriculturalists and hunted bison in western Nebraska. Western Nebraska was also home to "nomadic" tribes that resided in tepee villages and were dependent on bison hunting. These tribes include the Apache, Crow, Kiowa, Cheyenne, Teton, Comanche, and Arapahoe. The Lakota, Northern Cheyenne, and Arapaho resided in northwestern Nebraska, and the Oglala and Brule Sioux were concentrated around the Black Hills and the upper White and Niobrara rivers in northern Sioux County. By the mid 1800s, the Oglala and Brule had extended their range to include the Platte River region.

Europeans and Americans (300 to 100 BP). The earliest European presence in Nebraska was by French and Spanish explorers in the early AD 1700s and possibly earlier in the late 1600s. The Villasur expedition to explore the area was led by Pedro de Villasur out of the Spanish province of New Mexico in 1720 AD. Later explorers included Lewis and Clark and Zubulon Pike among others. These early expeditions provide descriptions of varying quality for some of the early historical tribes in the region. In the later 1700s and early 1800s more intensive contact and settlement occurred first through the fur trade in the 1830s and 1840s,

Description of the Affected Environment

and then through missionary and military contacts. By the mid-1800s, emigrant trails, notably the Oregon-California Trail, among others, traversed the Nebraska area. The large number of settlers moving along the emigrant trails passing through tribal use areas led to increasing conflict over time and the establishment of military forts in tribal lands, yet another irritant to tribes. Treaties, notably the Fort Laramie Treaty of 1851 were signed with the intent of removing tribes from along the emigrant trails and to allow for the building of trails and forts to protect settlers moving west. Continued conflict resulted in the creation of the Great Sioux Reservation bounded by the Missouri River on the east, the Big Horn Mountains on the west, and the 46th and 43rd parallels to the north and south, respectively. Fort Robinson in Dawes County was established in 1874 adjacent to the Red Cloud Agency near the White River. Fort Robinson served as a military outpost to contain the Sioux tribes on the Great Sioux Reservation, the Sioux Wars and the Cheyenne Outbreak. Fort Robinson continued in use through World War I and in World War II trained soldiers and served a prisoner of war camp. It ceased to be used as a military camp in 1948 and today is a Nebraska state park and historic site.

Ranching, a livelihood well suited to the grassland plains of western Nebraska, was practiced by early settlers by the early 1870s. The arrival of the railroads (Chicago and Northwestern and the Fremont, Elkhorn, and Missouri Valley) in 1885 opened northwestern Nebraska to a flood of settlers, most of them recent European immigrants. These early settlers began a period of extensive agriculture throughout western Nebraska, mostly around well-watered regions, but many of the settlers pursued newly developed dry-land farming techniques. The established ranching community relied on open range cattle grazing. Agricultural practices relied on fencing cattle out of fields. In response, ranchers would often fence off public lands to prevent settlement. This and other issues often led to conflict between farmers and ranchers and the eventual decline of ranching. In 1903, the North Platte irrigation project was authorized by Congress. The project included the construction of five reservoirs, six power plants and an irrigation canal system (the Interstate Canal).

3.4.8.2 Historic Properties Listed in the National and State Registers

3.4.8.2.1 Historic Properties in Western South Dakota

In addition to the sites listed in Table 3.4-10, the following sites in western South Dakota are listed on South Dakota state and/or the National Register of Historic Places. There are no listed sites in Butte, Fall River, or Pennington counties as of this writing.

Custer County

- Custer Campsite #1 RR
- Borglum Ranch & Studio Historic District RR

Lawrence County

- Thoen Stone & Site
- Frawley Ranch

Table 3.4-10. National Register Listed Properties in Counties Included in the Nebraska-South Dakota-Wyoming Uranium Milling Region

County	Resource Name	City	Date Listed YYYY/MM/DD
Wyoming			
Crook	DXN Bridge Over Missouri River	Hulett	1985-02-22
Crook	Entrance Road—Devils Tower National Monument	Devils Tower	2000-07-24
Crook	Entrance Station—Devils Tower National Monument	Devils Tower	2000-07-24
Crook	Inyan Kara Mountain	Sundance	1973-04-24
Crook	Old Headquarters Area Historic District	Devils Tower	2000-07-20
Crook	Ranch A	Beulah	1997-03-17
Crook	Sundance School	Sundance	1985-12-02
Crook	Sundance State Bank	Sundance	1984-03-23
Crook	Tower Ladder—Devils Tower National Monument	Devils Tower	2000-07-24
Crook	Vore Buffalo Jump	Sundance	1973-04-11
Crook	Wyoming Mercantile	Aladdin	1991-04-16
Niobrara	DSD Bridge Over Cheyenne River	Riverview	1985-02-22
Weston	Cambria Casino	Newcastle	1980-11-18
Weston	Jenney Stockade Site	Newcastle	1969-09-30
Weston	U.S. Post Office—Newcastle Main	Newcastle	1987-05-19
Weston	Weston County Courthouse	Newcastle	2001-09-01
Weston	Wyoming Army National Guard Cavalry Stable	Newcastle	1994-07-07
South Dakota			
Custer	Archeological Site No. 39CU1619	Custer	1999-06-03
Custer	Archeological Site No. 39CU70	Custer	1993-10-20
Custer	Archeological Site No. 39CU890	Hermosa	1993-08-06
Custer	Ayres, Lonnie and Francis, Ranch	Custer	1991-01-25
Custer	Badger Hole	Custer	1973-03-07
Custer	Bauer, Maria, Homestead Ranch	Custer	1992-06-09
Custer	Beaver Creek Bridge	Hot Springs	1984-08-08
Custer	Beaver Creek Rockshelter	Pringle	1993-10-25
Custer	Buffalo Gap Cheyenne River Bridge	Buffalo Gap	1988-02-08
Custer	Buffalo Gap Historic Commercial District	Buffalo Gap	1995-06-30
Custer	CCC Camp Custer Officers' Cabin	Custer	1992-06-09
Custer	Cold Springs Schoolhouse	Custer	1973-03-07
Custer	Custer County Courthouse	Custer	1972-11-27
Custer	Custer State Game Lodge	Custer	1983-03-30
Custer	Custer State Park Museum	Hermosa	1983-03-30
Custer	Fairburn Historic Commercial District	Fairburn	1995-06-30
Custer	First National Bank Building	Custer	1982-03-05
Custer	Fourmile School No. 21	Custer	1991-01-25
Custer	Garlock Building	Custer	2004-01-28
Custer	Grace Coolidge Memorial Log Building	Custer	2001-06-21
Custer	Historic Trail and Cave Entrance	Custer	1995-04-19
Custer	Lampert, Charles and Ollie, Ranch	Custer	1990-07-05
Custer	Mann, Irene and Walter, Ranch	Custer	1990-07-05
Custer	Norbeck, Peter, Summer House	Custer	1977-09-13
Custer	Pig Tail Bridge	Hot Springs	1995-04-07
Custer	Ranger Station	Custer	1995-04-05
Custer	Roetzel, Ferdinand and Elizabeth, Ranch	Custer	1991-01-25
Custer	Site No. 39 Cu 510	City Restricted	1982-05-20
Custer	Site No. 39 Cu 511	City Restricted	1982-05-20

Table 3.4-10. National Register Listed Properties in Counties Included in the Nebraska-South Dakota-Wyoming Uranium Milling Region (continued)

County	Resource Name	City	Date Listed YYYY/MM/DD
Custer	Site No. 39 Cu 512	City Restricted	1982-05-20
Custer	Site No. 39 Cu 513	City Restricted	1982-05-20
Custer	Site No. 39 Cu 514	City Restricted	1982-05-20
Custer	Site No. 39 Cu 515	City Restricted	1982-05-20
Custer	Site No. 39 Cu 516	City Restricted	1982-05-20
Custer	Site No. 39 Cu 91	City Restricted	1982-05-20
Custer	South Dakota Dept. of Transportation Bridge No. 17-289-107	Custer	1993-12-09
Custer	Stearns, William, Ranch	Custer	1990-07-05
Custer	Streeter, Norman B., Homestead	Buffalo Gap	1995-06-30
Custer	Towner, Francis Averill (T.A.) and Janet Leach, House	Custer	1990-06-21
Custer	Tubbs, Newton Seymour, House	Custer	1993-12-09
Custer	Ward, Elbert and Harriet, Ranch	Custer	1990-07-05
Custer	Way Park Museum	Custer	1973-03-07
Custer	Wind Cave National Park Administrative and Utility Area Historic District	Custer	1984-07-11
Custer	Young, Edna and Ernest, Ranch	Custer	1990-07-05
Fall River	Allen Bank Building and Cascade Springs Bath House-Sanitarium	Hot Springs	1984-02-23
Fall River	Archeological 39FA1638	Edgemont	2005-07-14
Fall River	Archeological Site 39FA1336	Edgemont	2005-07-14
Fall River	Archeological Site 39FA1937	Edgemont	2005-07-14
Fall River	Archeological Site No. 39FA1010	Hot Springs	1993-10-20
Fall River	Archeological Site No. 39FA1013	Hot Springs	1993-10-20
Fall River	Archeological Site No. 39FA1046	Edgemont	1993-10-20
Fall River	Archeological Site No. 39FA1049	Hot Springs	1993-08-06
Fall River	Archeological Site No. 39FA1093	Hot Springs	1993-10-20
Fall River	Archeological Site No. 39FA1152	Hot Springs	1993-10-20
Fall River	Archeological Site No. 39FA1154	Hot Springs	1993-10-20
Fall River	Archeological Site No. 39FA1155	Hot Springs	1993-10-20
Fall River	Archeological Site No. 39FA1190	Edgemont	1993-10-20
Fall River	Archeological Site No. 39FA1201	Edgemont	1993-08-06
Fall River	Archeological Site No. 39FA1204	Hot Springs	1993-10-20
Fall River	Archeological Site No. 39FA243	Edgemont	1993-10-20
Fall River	Archeological Site No. 39FA244	Edgemont	1993-10-20
Fall River	Archeological Site No. 39FA316	Edgemont	1993-10-20
Fall River	Archeological Site No. 39FA321	Edgemont	1993-10-20
Fall River	Archeological Site No. 39FA395	Edgemont	1993-10-20
Fall River	Archeological Site No. 39FA446	Edgemont	1993-10-20
Fall River	Archeological Site No. 39FA447	Edgemont	1993-10-20
Fall River	Archeological Site No. 39FA448	Edgemont	1993-10-20
Fall River	Archeological Site No. 39FA542	Edgemont	1993-10-25
Fall River	Archeological Site No. 39FA678	Edgemont	1993-08-06
Fall River	Archeological Site No. 39FA679	Edgemont	1993-10-20
Fall River	Archeological Site No. 39FA680	Edgemont	1993-10-20
Fall River	Archeological Site No. 39FA682	Edgemont	1993-10-20
Fall River	Archeological Site No. 39FA683	Edgemont	1993-10-20
Fall River	Archeological Site No. 39FA686	Edgemont	1993-10-20
Fall River	Archeological Site No. 39FA688	Edgemont	1993-10-20

Table 3.4-10. National Register Listed Properties in Counties Included in the Nebraska-South Dakota-Wyoming Uranium Milling Region (continued)

County	Resource Name	City	Date Listed YYYY/MM/DD
Fall River	Archeological Site No. 39FA690	Edgemont	1993-10-20
Fall River	Archeological Site No. 39FA691	Edgemont	1993-10-20
Fall River	Archeological Site No. 39FA767	Edgemont	1993-10-20
Fall River	Archeological Site No. 39FA788	Edgemont	1993-10-20
Fall River	Archeological Site No. 39FA806	Hot Springs	1993-08-06
Fall River	Archeological Site No. 39FA819	Edgemont	1993-10-20
Fall River	Archeological Site No. 39FA86	Edgemont	1993-08-06
Fall River	Archeological Site No. 39FA88	Edgemont	1993-10-20
Fall River	Archeological Site No. 39FA89	Edgemont	1993-08-06
Fall River	Archeological Site No. 39FA90	Hot Springs	1993-10-20
Fall River	Archeological Site No. 39FA99	Edgemont	1993-10-20
Fall River	Bartlett—Myers Building	Edgemont	2006-05-31
Fall River	Chilson Bridge	Edgemont	1993-12-09
Fall River	Flint Hill Aboriginal Quartzite Quarry	Edgemont	1978-07-14
Fall River	Hot Springs High School	Hot Springs	1980-05-07
Fall River	Hot Springs Historic District	Hot Springs	1974-06-25
Fall River	Jensen, Governor Leslie, House	Hot Spring	1987-09-25
Fall River	Log Cabin Tourist Camp	Hot Springs	2004-01-28
Fall River	Lord's Ranch Rockshelter	Edgemont	2005-07-14
Fall River	Petty House	Hot Springs	1999-02-12
Fall River	Site 39FA1303	Edgemont	2005-06-08
Fall River	Site 39FA1639	Edgemont	2005-06-09
Fall River	Site No. 39 FA 277	City Restricted	1982-05-20
Fall River	Site No. 39 FA 389	City Restricted	1982-05-20
Fall River	Site No. 39 FA 554	City Restricted	1982-05-20
Fall River	Site No. 39 FA 58	City Restricted	1982-05-20
Fall River	Site No. 39 FA 676	City Restricted	1982-05-20
Fall River	Site No. 39 FA 677	City Restricted	1982-05-20
Fall River	Site No. 39 FA 681	City Restricted	1982-05-20
Fall River	Site No. 39 FA 684	City Restricted	1982-05-20
Fall River	Site No. 39 FA 685	City Restricted	1982-05-20
Fall River	Site No. 39 FA 687	City Restricted	1982-05-20
Fall River	Site No. 39 FA 7	City Restricted	1982-05-20
Fall River	Site No. 39 FA 75	City Restricted	1982-05-20
Fall River	Site No. 39 FA 79	City Restricted	1982-05-20
Fall River	Site No. 39 FA 91	City Restricted	1982-05-20
Fall River	Site No. 39 FA 94	City Restricted	1982-05-20
Fall River	St. Martin's Catholic Church and Grotto	Oelrichs	2005-05-30
Fall River	Wesch, Phillip, House	Hot Springs	1984-02-23
Lawrence	Ainsworth, Oliver N., House	Spearfish	1990-10-25
Lawrence	Baker Bungalow	Spearfish	1996-10-24
Lawrence	Buskala, Henry Ranch	Dumont	1985-11-13
Lawrence	Cook, Fayette, House	Spearfish	1988-07-13
Lawrence	Corbin, James A., House	Spearfish	1990-10-25
Lawrence	Court, Henry, House	Spearfish	1990-10-25
Lawrence	Dakota Tin and Gold Mine	Spearfish	2005-06-08
Lawrence	Deadwood Historic District	Deadwood	1966-10-15
Lawrence	Dickey, Eleazer C. and Gwinnie, House	Spearfish	1989-07-13
Lawrence	Dickey, Walter, House	Spearfish	1988-05-16

Description of the Affected Environment

Table 3.4-10. National Register Listed Properties in Counties Included in the Nebraska-South Dakota-Wyoming Uranium Milling Region (continued)

County	Resource Name	City	Date Listed YYYY/MM/DD
Lawrence	Driskill, William D., House	Spearfish	1989-07-13
Lawrence	Episcopal Church of All Angels	Spearfish	1976-04-22
Lawrence	Evans, Robert H., House	Spearfish	1991-11-01
Lawrence	Frawley Historic Ranch	Spearfish	1974-12-31
Lawrence	Halloran-Matthews-Brady House	Spearfish	1976-12-12
Lawrence	Hewes, Arthur, House	Spearfish	1990-10-25
Lawrence	Hill, John, Ranch—Keltomaki	Brownsville	1985-11-13
Lawrence	Homestake Workers House	Spearfish	1991-11-01
Lawrence	Keets, Henry, House	Spearfish	1988-07-13
Lawrence	Knight, Webb S., House	Spearfish	1989-07-13
Lawrence	Kroll Meat Market and Slaughterhouse	Spearfish	1988-05-20
Lawrence	Lead Historic District	Lead	1974-12-31
Lawrence	Lown, William Ernest, House	Spearfish	1976-05-28
Lawrence	Mail Building, The	Spearfish	1988-05-16
Lawrence	McLaughlin Ranch Barn	Spearfish	2002-02-14
Lawrence	Mount Theodore Roosevelt Monument	Deadwood	2005-12-22
Lawrence	Old Finnish Lutheran Church	Lead	1985-11-13
Lawrence	Redwater Bridge, Old	Spearfish	1993-12-09
Lawrence	Riley, Almira, House	Spearfish	1989-07-13
Lawrence	Spearfish City Hall	Spearfish	1990-10-25
Lawrence	Spearfish Filling Station	Spearfish	1988-05-16
Lawrence	Spearfish Fisheries Center	Spearfish	1978-05-19
Lawrence	Spearfish Historic Commercial District	Spearfish	1975-06-05
Lawrence	Spearfish Post Office (Old)	Spearfish	1999-02-12
Lawrence	St. Lawrence O'Toole Catholic Church	Central City	2003-02-05
Lawrence	Tomahawk Lake Country Club	Deadwood	2005-10-26
Lawrence	Toomey House	Spearfish	1997-11-07
Lawrence	Uhlig, Otto L., House	Spearfish	1989-07-13
Lawrence	Walsh Barn	Spearfish	2003-05-30
Lawrence	Walton Ranch	Spearfish	2005-05-30
Lawrence	Whitney, Mary, House	Spearfish	1990-10-25
Lawrence	Wolzmuth, John, House	Spearfish	1988-07-13
Pennington	Archeological Site No. 39PN376	Spearfish	1989-07-13
Pennington	Burlington and Quincy High Line Hill City to Keystone Branch	Spearfish	1990-10-25
Pennington	Byron, Lewis, House	Spearfish	1988-05-16
Pennington	Calumet Hotel	Spearfish	1978-05-19
Pennington	Casper Supply Company of SD	Spearfish	1975-06-05
Pennington	Cassidy House	Spearfish	1999-02-12
Pennington	Church of the Immaculate Conception	Central City	2003-02-05
Pennington	Dean Motor Company	Deadwood	2005-10-26
Pennington	Dinosaur Park	Spearfish	1997-11-07
Pennington	Emmanuel Episcopal Church	Spearfish	1989-07-13
Pennington	Fairmont Creamery Company Building	Spearfish	2003-05-30
Pennington	Feigel House	Spearfish	2005-05-30
Pennington	First Congregational Church	Spearfish	1990-10-25
Pennington	Gambrill Storage Building	Spearfish	1988-07-13
Pennington	Harney Peak Hotel	Custer	1993-10-25
Pennington	Harney Peak Tin Mining Company Buildings	Hill City	2003-02-05

Table 3.4-10. National Register Listed Properties in Counties Included in the Nebraska-South Dakota-Wyoming Uranium Milling Region (continued)

County	County	County	County
Pennington	Otho Mining District	Hermosa	1999-12-17
Pennington	Pennington County Courthouse	Hill City	1977-04-11
Pennington	Quinn, Michael, House	Custer	1983-03-10
Pennington	Rapid City Carnegie Library	Hill City	1977-07-21
Pennington	Rapid City Garage	Keystone	1981-02-22
Pennington	Rapid City Historic Commercial District	Keystone	1982-06-17
Pennington	Rapid City Laundry	Hill City	1994-06-03
Pennington	Site No. 39 PN 108	City Restricted	1982-05-20
Pennington	Site No. 39 PN 438	City Restricted	1982-05-20
Pennington	Site No. 39 PN 439	City Restricted	1982-05-20
Pennington	Site No. 39 PN 57	City Restricted	1982-05-20
Pennington	Von Woehrmann Building	Hill City	1977-04-13
Nebraska			
Dawes	Army Theatre	Crawford	1988-07-07
Dawes	Bordeaux Trading Post	Chadron	1972-03-16
Dawes	Chadron Public Library	Chadron	1990-06-21
Dawes	Co-operative Block Building	Crawford	1985-09-12
Dawes	Crites Hall	Chadron	1983-09-08
Dawes	Dawes County Courthouse	Chadron	1990-07-05
Dawes	Fort Robinson and Red Cloud Agency	Crawford	1966-10-15
Dawes	Hotel Chadron	Chadron	2002-08-15
Dawes	Library	Chadron	1983-09-08
Dawes	Miller Hall	Chadron	1983-09-08
Dawes	Sparks Hall	Chadron	1983-09-08
Dawes	U.S. Post Office—Crawford	Crawford	1992-05-11
Dawes	Wohlers, Henry, Sr., Homestead	Crawford	2004-10-15
Dawes	Work, Edna, Hall	Chadron	1983-09-08
Sioux	Cook, Harold J., Homestead Cabin	Agate	1977-08-24
Sioux	Hudson-Meng Bison Kill Site	Crawford	1973-08-28
Sioux	Sioux County Courthouse	Harrison	1990-07-05

3.4.8.2.2 Historic Properties in Western Nebraska

In addition to the sites listed in Table 3.4-10, the following sites in western Nebraska are listed on Nebraska state and/or the National Register of Historic Places:

Dawes County

- James Bordeaux Trading Post [DW00-002] Listed 1972/03/16
- Henry Wohlers, Sr. Homestead [DW00-043] Listed 2004/10/15
- Chadron Commercial Historic District [DW03] Listed 2007/3/27
- Chadron State College Historic Buildings [DW03] Listed 1983/09/08
- Hotel Chadron [DW03-023] Listed 2002/08/15
- Dawes County Courthouse [DW03-081] Listed 1990/07/05
- Chadron Public Library [DW03-091] Listed 1990/06/21
- Crawford United States Post Office [DW04-007] Listed 1992/05/11
- Co-Operative Block Building [DW04-024] Listed 1985/09/12
- Fort Robinson and Red Cloud Agency [DW07] Listed 1966/10/15

These sites are located within about 5–8 km [3–5 mi] of the existing Crow Butte ISL Facility.

Sioux County

- Hudson-Meng Bison Kill Site [25-SX-115] Listed 1973/08/28
- Harold J. Cook Homestead (Bone Cabin Complex) [SX00-028] Listed 1977/08/24
- Sandford Dugout [SX00-032] Listed 2000/03/09
- Wind Springs Ranch Historic and Archeological District [SX00-033, 25-SX-77, 25-SX-600-655] Listed 2000/11/22
- Sioux County Courthouse [SX04-002] Listed 1990/07/05

3.4.8.3 Tribal Consultations

3.4.8.3.1 South Dakota Tribal Consultation

There are 10 Native American Tribes located within or immediately adjacent to the state of South Dakota. These are the Cheyenne River Sioux, Flandreau Santee Sioux, Lower Brulé Sioux, the Crow Tribe of Montana Oglala Sioux, Rosebud Sioux, Sisseton-Whapeton Oyate, Standing Rock Sioux, Yankton Sioux, and the Ponca Tribe of Nebraska. The Siouan tribes are located throughout South and North Dakota, whereas the Ponca are located in northeastern Nebraska, but have interests in South Dakota. These and other Siouan-speaking tribes in North Dakota, Wyoming, Montana and Nebraska may have traditional land use claims in western South Dakota.

The United States government and the State of South Dakota recognize the sovereignty of certain Native America tribes. These tribal governments have legal authority for their respective reservations. Executive Order 13175 requires federal agencies to undertake consultation and coordination with Indian tribal governments on a government-to-government basis. In addition, the National Historic Preservation Act provides these tribal groups with the opportunity to manage cultural resources within their own lands under the legal authority of a Tribal Historic Preservation Office (THPO).** The THPO therefore replaces the South Dakota SHPO as the agency responsible for the oversight of all federal and state historic preservation compliance laws. To date, no tribes in South Dakota have applied for Status as a THPO as provided by the NHPA. Projects proponents must, however, contact tribal cultural resources personnel as part of the consultation process along with the South Dakota SHPO. The SHPO ensures compliance with applicable federal laws on tribal lands and undertakes consultation with the tribes and the Bureau of Indian Affairs for undertakings that might occur on tribal reservation lands. Some tribes have historic and cultural preservation offices that are not recognized as THPOs, but must also be consulted where they exist.

3.4.8.3.2 Nebraska Tribal Consultation

There are six Native American Tribes located within the state of Nebraska. These are the Omaha, Ponca, Winnebago, Santee Sioux, the Iowa Tribe of Kansas and Nebraska, and the Sac and Fox Nation of Missouri, Kansas, and Nebraska. These tribes are located near the Missouri River in eastern Nebraska. There are no reservation lands in western Nebraska. However, the Oglala Sioux Tribe of the Pine Ridge Reservation are located at the Nebraska-South Dakota border adjacent to the Nebraska-South Dakota-Wyoming Uranium Region.

1 These and other Siouan-speaking tribes in South Dakota, Wyoming and Nebraska may have
2 traditional land use claims in western Nebraska.

3
4 The United States government and the State of Nebraska recognize the sovereignty of certain
5 Native America tribes. These tribal governments have legal authority for their respective
6 reservations. Executive Order 13175 requires executive branch federal agencies to undertake
7 consultation and coordination with Indian tribal governments on a government-to-government
8 basis. NRC, as an independent federal agency, has agreed to voluntarily comply with Executive
9 Order 13175.

10
11 In addition, the National Historic Preservation Act provides these tribal groups with the
12 opportunity to manage cultural resources within their own lands under the legal authority of a
13 THPO. The THPO therefore replaces the Nebraska SHPO as the agency responsible for the
14 oversight of all federal and state historic preservation compliance laws. To date, no tribes in
15 Nebraska have applied for status as a THPO as provided by the NHPA. Some tribes have
16 historic and cultural preservation offices that are not recognized as THPOs, but they should be
17 consulted where they exist. NRC, in meetings its responsibilities under the NHPA, contacts
18 tribal cultural resources personnel as part of the consultation process, along with consulting with
19 the Nebraska SHPO.

20 21 **3.4.8.4 Places of Cultural Significance**

22
23 As described in Section 3.2.8.4, Traditional Cultural Properties are places of special heritage
24 value to contemporary communities because of their association with cultural practices and
25 beliefs that are rooted in the histories of those communities and are important in maintaining the
26 cultural identity of the communities (Parker and King, 1998; King, 2003). Religious places are
27 often associated with prominent topographic features like mountains, peaks, mesas, springs
28 and lakes. In addition shrines may be present across the landscape to denote specific culturally
29 significant locations and vision quest sites where an individual can place offerings.

30
31 Information on traditional land use and the location of culturally significant places is often
32 protected information within the community (King, 2003). Therefore, the information presented
33 on religious places is limited to those that are identified in the published literature and are
34 therefore restricted to a few highly recognized places on the landscape within southwestern
35 South Dakota.

36
37 Traditional cultural properties are ones that refer to beliefs, customs, and practices of a living
38 community that have been passed down over the generations. Native American traditional
39 cultural properties are often not found on the state or national registers of historic properties or
40 described in the extant literature or in SHPO files. There are, however, a range of cultural
41 properties types of religious or traditional use that might be identified during the tribal
42 consultation process. These might include:

- 43
44 • Sites of ritual and ceremonial activities and related features
45 • Shrines
46 • Marked and unmarked burial grounds
47 • Traditional use areas
48 • Plant and mineral gathering areas
49 • Traditional hunting areas
50 • Caves and rock shelters

- Springs
- Trails
- Prehistoric archaeological sites

The U.S. Bureau of Indian Affairs web site contains a list, current as of May 2007, of tribal leaders and contact information <<http://www.doi.gov/bia/Tribal%20Leaders-June%202007-2.pdf>>. These tribal groups should be contacted for consultations associated with ISL milling activities in their respective states (see Table 3.2-12). Additional tribal contact information may be obtained from the respective State Historic Preservation Offices in Nebraska, Montana, South Dakota, and Wyoming.

3.4.8.4.1 Places of Cultural Significance in South Dakota

There are no known culturally significant places listed in Butte, Lawrence, Pennington, Custer, or Fall River counties. However, the Siouan tribes who once occupied portions of South Dakota (Cheyenne River Sioux, Flandreau Santee Sioux, Lower Brule Sioux, Oglala Sioux, Rosebud Sioux, Sisseton-Whapeton Oyate, Standing Rock Sioux, Yankton Sioux, and the Ponca Tribe of Nebraska) consider the Black Hills in Wyoming and South Dakota, Devil's Tower in northeastern Wyoming, and Bear Butte in southwestern South Dakota to be culturally significant.

Areas of western South Dakota, once used by these tribes may contain additional, undocumented culturally significant sites and traditional cultural properties. Mountains, peaks, buttes, prominences, and other elements of the natural and cultural environment are often considered important elements of a traditional culturally significant landscape.

3.4.8.4.2 Places of Cultural Significance in Nebraska

There are no known culturally significant places listed in Dawes and Sioux counties. However, the tribes who once occupied western Nebraska (Lakota, Northern Cheyenne, Arapaho, Oglala and Brule Sioux) along the upper White and Niobrara rivers and extending into the Black Hills of South Dakota all consider the Black Hills in Wyoming and South Dakota, Devil's Tower in northeastern Wyoming, and Bear Butte in southwestern South Dakota to be culturally significant.

Areas of western Nebraska once used by these tribes may contain additional, undocumented culturally significant sites and traditional cultural properties. Mountains, peaks, buttes, prominences, and other elements of the natural and cultural environment are often considered important elements of a traditional culturally significant landscape.

3.4.9 Visual/Scenic Resources

Based on the BLM Visual Resource Handbook, the Nebraska-South Dakota-Wyoming Uranium Milling Region (BLM, 2007a–c) is located within the Great Plains physiographic province, adjacent to the southern end of the Black Hills. The northwestern corner of Wyoming (see Figure 3.3-17) is located within the area managed by the Newcastle BLM field office (BLM, 2000). Most of the area is categorized as VRM Class III, but there are some Class II areas identified around Devils Tower National Monument and the Black Hills National Forest along the Wyoming-South Dakota border (see Figure 3.4-1). One potential uranium ISL facility has been identified for development in the northeast corner of Nebraska-South Dakota-Wyoming Uranium Milling Region, about 16 km [10 mi] northeast of the Black Hills National Forest, and about

45 km [28 mi] northeast of Devils Tower. There are no Wyoming Unique/Irreplaceable or Rare/Uncommon designated areas within the Nebraska-South Dakota-Wyoming Uranium Milling Region (Girardin, 2006).

Uranium resources in South Dakota are being evaluated near Fall River County in the southwestern corner of the state. Although it does not assign a VRM classification to the region, the Nebraska and South Dakota BLM field offices resource management plan classifies this region as having natural vegetation of wheatgrass, grama grass, sagebrush, and pine savanna (BLM, 1992, 1985). Similar areas are identified as Class III VRM areas in Wyoming. The USFS has also performed some visual resource classification in association with its forest and grasslands management plans in the region (see text box in Section 3.2.9). The revisions to Northern Great Plains Management Plans (USFS, 2001a) indicate that for the grasslands in Fall River County, almost 95 percent of the area is categorized with a scenic integrity objective of low to moderate (moderately to heavily altered). The Black Hills National Forest land and resource management plan and subsequent amendments (USFS, 1997, 2001b, 2005) identified management plans to maintain about 85 percent of the region for low to moderate scenic integrity objectives. About 15 percent is identified as high (13.6 percent) to very high (1.2 percent) scenic integrity objectives (USFS, 2005). In areas lacking human-caused disturbances, the landscape has attributes that potentially have a high level of scenic integrity (USFS, 2005). There is a prevention of significant deterioration Class 1 Areas identified for the Wind Cave National Park in South Dakota as described in Section 3.4.6.2 and shown in Figure 3.4-20, but this is at least 40 km [25 mi] east of the closest potential uranium ISL facility.

Similar to South Dakota, uranium resources in Dawes County in northwestern Nebraska are located in the Great Plains physiographic province. The Crow Butte ISL facility in Dawes County is located near the Pine Ridge Unit of the Nebraska National Forest. The revisions to Northern Great Plains Management Plans (USFS, 2001a) indicate that for the Oglala National Grassland and the Pine Ridge Unit of the Nebraska National Forest, about 87 percent of the landscape is classified as having low to moderate scenic integrity objective classification, with the remaining 13 percent roughly divided between high (7.3 percent) to very high (5.4 percent).

3.4.10 Socioeconomics

For the purpose of this GEIS, the socioeconomic description for the Nebraska-South Dakota-Wyoming Region includes communities within the region of influence for potential ISL facilities in the three uranium districts in the region. These include communities that have the highest potential for socioeconomic impacts and are considered the affected environment. Communities that have the highest potential for socioeconomic impacts are defined by (1) proximity to an ISL facility {generally within 48 km [30 mi]}, (2) economic profile, such as potential for income growth or de-stabilization, (3) employment structure, such as potential for job placement or displacement and (4) community profile, such as potential for growth or destabilization to local emergency services, schools, or public housing. The affected environment within the Nebraska-South Dakota-Wyoming Uranium Milling Region consists of counties and Native American communities. The affected environment is listed in Table 3.4-11. The following subsections describe areas most likely to have implications to socioeconomics and are listed below. A Core-Based Statistical Areas, according to the U.S. Census Bureau, is a collective term for both metro and micro areas ranging from a population of 10,000 to 50,000. A Metropolitan Area is greater than 50,000 and a town is considered less than 10,000 in population (U.S. Census Bureau, 2007). Smaller communities are considered as part of the county demographics.

Table 3.4-11. Summary of Affected Environment Within the Nebraska-South Dakota-Wyoming Uranium Milling Region

Counties Within Nebraska	Counties Within South Dakota	Counties Within Wyoming	Native American Communities Within South Dakota
Dawes	Butte	Campbell	Pine Ridge Indian Reservation
Sioux	Custer	Crook	
	Fall River	Niobrara	
	Shannon	Weston	

3.4.10.1 Demographics

Demographics for the year 2000 are based on population and racial characteristics of the affected environment and are provided in Tables 3.4-12 through 3.4-14. Figure 3.4-21 illustrates the populations of communities within the Nebraska-South Dakota/-Wyoming Uranium Milling Region. Most 2006 data compiled by the U.S. Census Bureau is not yet available for the geographic areas of interest.

Based on review of Tables 3.4-12 – 3.4-14, the most populated county is Campbell County, Wyoming and the most sparsely populated county is Sioux County, Nebraska. For communities located within 48 km [30 mi] of potential ISL facilities, the most populated town is Pine Ridge, South Dakota (Pine Ridge Indian Reservation) and the smallest populated town is Oglala, South Dakota (Pine Ridge Indian Reservation). The county with the largest percentage of non-minorities is Niobrara County, Wyoming with a white population of 98.0 percent. The town with the largest minority population is Pine Ridge, South Dakota with a white population of 3.7 percent. The largest minority based county is Shannon County, South Dakota with a white population of only 4.5 percent. The largest minority-based town is Oglala, South Dakota with a white population of only 0.7 percent.

Although not listed in Table 3.4-12, the total population counts based on 2000 Census data for the Pine Ridge Indian Reservation totaled 15,521 individuals (U.S. Census Bureau, 2008), with approximately 93 percent Native American. However, recent studies suggest that the population may be larger (Housing Assistance Council, 2002).

3.4.10.2 Income

Income information from the 200 Census including labor force, income, and poverty levels for the affected environment in the Nebraska-South Dakota-Wyoming Uranium Milling Region is based on data collected at the state and county levels.

Data collected at the state level also includes information on towns, Core-Based Statistical Areas, or Metropolitan Areas and was done to take into consideration an outside workforce. An outside workforce may be a workforce willing to commute long distances {greater than 48 km [30 mi]} for income opportunities or may be a workforce necessary to fulfill specialized positions (if local workforce is unavailable or unspecialized). Data collected from a county level is generally the same affected environment previously discussed in Table 3.4-11 and also includes information on Native American communities near the Nebraska-South Dakota-Wyoming Uranium Milling Region. State-level information is provided in Table 3.4-15 and county data are listed in Table 3.4-16.

Table 3.4-12. 2000 U.S. Bureau of Census Population and Race Categories of Nebraska*

Affected Environment	Total Population	White	African American	Native American	Some Other Race	Two or More Races	Asian	Hispanic Origin†	Native Hawaiian and Other Pacific Islander
Nebraska	1,711,263	1,533,261	68,541	14,896	47,845	23,953	21,931	94,425	836
Percent of total		89.6%	4.0%	0.9%	2.8%	1.4%	1.3%	5.5%	0.0%
Dawes County	9,060	8,457	73	261	93	143	28	220	5
Percent of total		93.3%	0.8%	2.9%	1.0%	1.6%	0.3%	2.4%	0.1%
Sioux County	1,475	1,440	0	2	17	13	3	34	0
Percent of total		97.6%	0.0%	0.1%	1.2%	0.9%	0.2%	2.3%	0.0%

*U.S. Census Bureau. "American FactFinder." 2000. <http://factfinder.census.gov/home/saff/main.html?_lang=en> (18 October 2007 and 26 February 2008).

†Hispanic origin can be any race and is calculated as a separate component of the total population (i.e., if added to the other races would total more than 100%).

Table 3.4-13. 2000 U.S. Bureau of Census Population and Race Categories of South Dakota*

Affected Environment	Total Population	White	African American	Native American	Some Other Race	Two or More Races	Asian	Hispanic Origin†	Native Hawaiian and Other Pacific Islander
South Dakota	754,854	669,404	4,685	62,283	3,677	10,156	4,378	10,903	261
Percent of total		88.7%	0.6%	8.3%	0.5%	1.3%	0.6%	1.4%	0.0%
Butte County	9,094	8,687	9	150	99	127	22	266	0
Percent of total		95.5%	0.1%	1.6%	1.1%	1.4%	0.2%	2.9%	0.0%
Custer County	7,275	6,851	20	227	26	137	13	110	1
Percent of total		94.2%	0.3%	3.1%	0.4%	1.9%	0.2%	1.5%	0.0%
Fall River County	7,453	6,746	24	451	22	189	17	130	4
Percent of total		90.5%	0.3%	6.1%	0.3%	2.5%	0.2%	1.7%	0.1%
Shannon County	12,466	562	10	11,743	28	114	3	177	6
Percent of total		4.5%	0.1%	94.2%	0.2%	0.9%	0.0%	1.4%	0.0%
Oglala (Pine Ridge Indian Reservation)	1,229	9	0	1,214	1	4	1	4	0
Percent of total		0.7%	0.0%	98.8%	0.1%	0.3%	0.1%	0.3%	0.0%

Table 3.4-13. 2000 U.S. Bureau of Census Population and Race Categories of South Dakota* (continued)


Affected Environment	Total Population	White	African American	Native American	Some Other Race	Two or More Races	Asian	Hispanic Origin†	Native Hawaiian and Other Pacific Islander
Pine Ridge (Pine Ridge Indian Reservation)	3,171	118	3	2,987	16	43	1	57	3
Percent of total		3.7%	0.1%	94.2%	0.5%	1.4%	0.0%	1.8%	0.1%
*U.S. Census Bureau. "American FactFinder." < http://factfinder.census.gov/home/saff/main.html?_lang=en > (18 October 2007, 26 February 2008, and 15 April 2008).									
†Hispanic origin can be any race and is calculated as a separate component of the total population (i.e., if added to the other races would total more than 100 %).									

Table 3.4-14. 2000 U.S. Bureau of Census Population and Race Categories of Northwestern Wyoming*

Affected Environment	Total Population	White	African American	Native American	Some Other Race	Two or More Races	Asian	Hispanic Origin†	Native Hawaiian and Other Pacific Islander
Wyoming	493,782	454,670	3,722	11,133	12,301	8,883	2,771	31,669	302
Percent of total		92.1%	0.8%	2.3%	2.5%	1.8%	0.6%	6.4%	0.1%
Campbell County	33,698	32,369	51	313	378	450	108	1,191	29
Percent of total		96.1%	0.2%	0.9%	1.1%	1.3%	0.3%	3.5%	0.1%
Crook County	5,887	5,761	3	60	15	44	4	54	0
Percent of total		97.9%	0.1%	1.0%	0.3%	0.7%	0.1%	0.9%	0.0%
Niobrara County	2,407	2,360	3	12	12	17	3	36	0
Percent of total		98.0%	0.1%	0.5%	0.5%	0.7%	0.1%	1.5%	0.0%
Weston County	6,644	6,374	8	84	62	102	13	137	1
Percent of total		95.9%	0.1%	1.3%	0.9%	1.5%	0.2%	2.1%	0.0%

*U.S. Census Bureau. "American FactFinder." <http://factfinder.census.gov/home/saff/main.html?_lang=en> (18 October 2007, 25 February 2008, and 25 April 2008).

†Hispanic origin can be any race and is calculated as a separate component of the total population (i.e., if added to the other races would total more than 100 percent).



SOUTH DAKOTA - NEBRASKA REGION




Figure 3.4-21. Nebraska-South Dakota-Wyoming Uranium Milling Region With Population

Table 3.4-15. U.S. Bureau of Census State Income Information for the Nebraska-South Dakota-Wyoming Uranium Milling Region*

Affected Environment	2000 Labor Force Population (16 Years and Over)	Median Household Income in 1999	Median Family Income in 1999	Per Capita Income in 1999	Families Below Poverty Level in 2000	Individuals Below Poverty Level in 2000
Nebraska	917,470	\$39,250	\$48,032	\$19,613	29,977	161,269
South Dakota	394,945	\$35,282	\$43,237	\$17,562	18,172	95,900
Wyoming	257,808	\$37,892	\$45,685	\$19,134	10,585	54,777
Alliance, Nebraska	4,531	\$39,408	\$47,766	\$18,584	255	979
<i>Percent of total†</i>	66.7%	NA	NA	NA	10.6%	11.2%
Chadron, Nebraska	3,228	\$27,400	\$44,420	\$16,312	127	1,025
<i>Percent of total†</i>	68.26%	NA‡	NA	NA	11.0%	21.4%
Gering, Nebraska	3,927	\$35,185	\$42,378	\$18,775	130	590
<i>Percent of total†</i>	64.1%	NA	NA	NA	5.9%	7.8%
Rapid City, South Dakota	31,948	\$35,978	\$44,818	\$19,445	1,441	7,328
<i>Percent of total†</i>	68.8%	NA	NA	NA	9.4%	12.7%
Scottsbluff, Nebraska	7,122	\$29,938	\$37,778	\$17,065	562	2,654
<i>Percent of total†</i>	62.5%	NA	NA	NA	14.5%	18.3%

Affected Environment	2000 Labor Force Population (16 Years and Over)	Median Household Income in 1999	Median Family Income in 1999	Per Capita Income in 1999	Families Below Poverty Level in 2000	Individuals Below Poverty Level in 2000
Spearfish, South Dakota	4,635	\$26,887	\$40,257	\$16,565	189	1,362
<i>Percent of total†</i>	65.1%	NA	NA	NA	9.8%	17.4%
Sturgis, South Dakota	3,199	\$30,253	\$38,698	\$16,763	187	756
<i>Percent of total†</i>	63.0%	NA	NA	NA	11.0%	12.0%
Casper, Wyoming	26,343	\$36,567	\$46,267	\$19,409	1,122	5,546
<i>Percent of total†</i>	68.4%	NA	NA	NA	8.5%	11.4%

U.S. Census Bureau. "American FactFinder." <http://factfinder.census.gov/home/saff/main.html?_lang=en> (18 October 2007, 26 February 2008, 15 April 2008, and 25 April 2008).
†Percent of total based on a population of 16 years and over.
‡NA = not applicable.

‡NA = not applicable.

Table 3.4-16. U.S. Bureau of Census County and Native American Income Information for the Nebraska-South Dakota-Wyoming Uranium Milling Region*

South Dakota*						
Affected Environment	2000 Labor Force Population (16 Years and Over)	Median Household Income in 1999	Median Family Income in 1999	Per Capita Income in 1999	Families Below Poverty Level in 2000	Individuals Below Poverty Level in 2000
Butte County	4,683	\$29,040	\$34,173	\$13,997	234	1,147
<i>Percent of total†</i>	68.3%	NA	NA	NA	9.4%	12.8%
Custer County	3,535	\$36,303	\$43,628	\$17,945	129	659
<i>Percent of total†</i>	59.6%	NA‡	NA	NA	6.2%	9.4%
Fall River County	3,408	\$29,631	\$37,827	\$17,048	153	951
<i>Percent of total†</i>	59.6%	NA	NA	NA	7.8%	13.6%
Shannon County	3,884	\$20,916	\$20,897	\$6,286	1,056	6,385
<i>Percent of total†</i>	52.4%	NA	NA	NA	45.1%	52.3%
Oglala (Pine Ridge Indian Reservation)	339	\$17,300	\$19,688	\$3,824	88	733
<i>Percent of total†</i>	49.9%	NA	NA	NA	45.1%	55.8%
Pine Ridge (Pine Ridge Indian Reservation)	1,149	\$21,089	\$20,170	\$6,067	320	2,057
<i>Percent of total†</i>	57.0%	NA	NA	NA	49.2%	61.0%

Table 3.4-16. U.S. Bureau of Census State Income Information for Nebraska-South Dakota-Wyoming* (continued)

Affected Environment	2000 Labor Force Population (16 Years and Over)	Median Household Income in 1999	Median Family Income in 1999	Per Capita Income in 1999	Families Below Poverty Level in 2000	Individuals Below Poverty Level in 2000
Dawes County	4,989	\$29,476	\$41,092	\$16,353	207	1,548
<i>Percent of total†</i>	66.8%	NA‡	NA	NA	9.8%	18.9%
Sioux County	749	\$29,851	\$31,406	\$15,999	48	227
<i>Percent of total†</i>	64.7%	NA	NA	NA	11.1%	15.4%
Wyoming*						
Campbell County	18,805	\$49,536	\$53,927	\$20,063	507	2,544
<i>Percent of total†</i>	76.6%	NA	NA	NA	5.6%	7.6%
Crook County	2,937	\$35,601	\$43,105	\$17,379	129	529
<i>Percent of total†</i>	64.4%	NA	NA	NA	7.8%	9.1%
Niobrara County	1,193	\$29,701	\$33,714	\$15,757	74	309
<i>Percent of total†</i>	61.5%	NA	NA	NA	10.7%	13.4%
Weston County	3,183	\$32,348	\$40,472	\$17,366	119	628
<i>Percent of total†</i>	60.0%	NA	NA	NA	6.3%	9.9%

U.S. Census Bureau. "American FactFinder." <http://factfinder.census.gov/home/saff/main.html?_lang=en> (18 October 2007, 26 February 2008, 15 April 2008, and 25 April 2008).
†Percent of total based on a population of 16 years and over.
‡NA = not applicable.

For the surrounding region, the state with the largest labor force population and families and individuals below poverty level is Nebraska (Table 3.4-15). The population with the largest labor force is Rapid City, South Dakota {48 km [30 mi] from the nearest potential ISL facility} and the smallest labor force population is Sturgis, South Dakota {32 km [20 mi] from the nearest potential ISL facility}. The population with the largest per capita income is Rapid City, South Dakota and the smallest per capita income population is Chadron, Nebraska {16 km [10 mi] from the nearest ISL facility}. The population with the highest percentage of individuals and families below poverty levels is Scottsbluff, Nebraska {32 km [20 mi] from the nearest ISL facility}.

Within the Nebraska-South Dakota-Wyoming Uranium Milling Region, the county with the largest labor force population is Campbell County, Wyoming and the county with the smallest labor force population is Sioux County, Nebraska (Table 3.4-16). The town with the largest labor force population is Pine Ridge, South Dakota (Pine Ridge Indian Reservation) and the town with the smallest labor force population is Oglala, South Dakota (Pine Ridge Indian Reservation). The county with the largest per capita income is Campbell County, Wyoming, and the lowest per capita income county is Shannon County, South Dakota. The county with the highest percentage of individuals and families below poverty levels is Shannon County, South Dakota, and the town with the highest percentage of individuals and families below poverty levels is Pine Ridge, South Dakota.

3.4.10.3 Housing

Housing information from the 2000 Census data for the affected environment is provided in Table 3.4-17 through 3.4-19.

The availability of housing within the immediate vicinity of the proposed ISL facilities is limited (Housing Assistance Council, 2002). The majority of housing is available in larger populated areas such as the Core-Based Statistical Areas and towns of Rapid City, South Dakota {48 km [30 mi] from the nearest ISL facility}, Spearfish, South Dakota {16 km [10 mi] to nearest potential ISL facility}, Sturgis, South Dakota {32 km [20 mi] from the nearest ISL facility}, Chadron, Nebraska {16 km [10 mi] to nearest ISL facility}, Alliance, Nebraska {16 km [10 mi] from the nearest ISL facility}, and Gillette, Wyoming {64 km [40 mi] from the nearest ISL facility}. There are approximately 10 housing units including manufactured housing (trailer homes) and residential property (neighborhoods) currently available in the region (mapquest, 2008c).

Temporary housing such as apartments, lodging, and trailer camps within the immediate vicinity of the proposed ISL facilities is not as limited. The majority of apartments are available in larger populated areas such as the Core-Based Statistical Areas and towns of Rapid City, Spearfish, and Sturgis in South Dakota; Chadron and Alliance in Nebraska; and Gillette in Wyoming, with about 25 apartment complexes currently available (MapQuest, 2008). There are also approximately 10 hotels/motels located along major highways or towns near the proposed ISL facilities. In addition to apartments and lodging, there are 20 trailer camps situated along major roads or near towns (MapQuest, 2008c).

Table 3.4-17. U.S. Bureau of Census Housing Information for the Nebraska Uranium Milling Region*

Affected Environment	Single Family Owner-Occupied Homes	Median Value in Dollars	Median Monthly Costs With a Mortgage	Median Monthly Costs Without a Mortgage	Occupied Housing Units	Renter-Occupied Units
Nebraska	370,495	\$88,000	\$895	\$283	666,184	207,216
Dawes County	1,553	\$55,200	\$684	\$262	3,512	1,211
Sioux County	140	\$42,600	\$600	\$257	605	106

*U.S. Census Bureau. "American FactFinder." <http://factfinder.census.gov/home/saff/main.html?_lang=en> (18 October 2007 and 26 February 2008).

Table 3.4-18. U.S. Bureau of Census Housing Information for South Dakota*

Affected Environment	Single Family Owner-Occupied Homes	Median Value in Dollars	Median Monthly Costs With a Mortgage	Median Monthly Costs Without a Mortgage	Occupied Housing Units	Renter-Occupied Units
South Dakota	137,531	\$79,600	\$828	\$279	290,245	87,887
Butte County	1,360	\$60,200	\$706	\$272	3,516	841
Custer County	1,073	\$89,100	\$884	\$292	2,970	1,073
Fall River County	1,286	\$54,300	\$687	\$271	3,127	901
Shannon County	631	\$25,900	\$515	\$192	2,785	1,323
Oglala (Pine Ridge Indian Reservation)	29	\$70,700	\$450	\$99	239	145
Pine Ridge (Pine Ridge Indian Reservation)	126	\$15,000	\$0	\$185	709	473

*U.S. Census Bureau. "American FactFinder." <http://factfinder.census.gov/home/saff/main.html?_lang=en> (18 October 2007, 26 February 2008, and 15 April 2008).

Table 3.4-19. U.S. Bureau of Census Housing Information for the Nebraska-South Dakota-Wyoming Uranium Milling Region*

Affected Environment	Single Family Owner-Occupied Homes	Median Value in Dollars	Median Monthly Costs With a Mortgage	Median Monthly Costs Without a Mortgage	Occupied Housing Units	Renter-Occupied Units
Wyoming	95,591	\$96,600	\$825	\$229	193,608	55,793
Campbell County	5,344	\$102,900	\$879	\$247	12,207	3,174
Crook County	836	\$85,4000	\$682	\$207	2,308	411
Niobrara County	480	\$60,300	\$562	\$200	1,011	222
Weston County	1,174	\$66,700	\$664	\$199	2,624	549

Source: U.S. Census Bureau. "American FactFinder." <http://factfinder.census.gov/home/saff/main.html?_lang=en> (18 October 2007, 25 February 2008, and 25 April 2008).

3.4.10.4 Employment Structure

The regional employment structure from the 2000 Census data, including employment rate and type is collected at the state and county levels. Data collected at the state level also include information on towns, Core-Based Statistical Areas, or Metropolitan Areas and was done to take into consideration an outside workforce. An outside workforce may be a workforce willing to commute long distances {greater than 48 km [30 mi]} for employment opportunities or may be a workforce necessary to fulfill specialized positions (if local workforce is unavailable or unspecialized). Data collected from a county level is the same affected environment previously discussed in Table 3.4-11 and also includes information on Native American communities.

For the region surrounding the Nebraska-South Dakota-Wyoming Uranium Milling Region, the state with the highest percentage of employment is Nebraska. The population with the highest percentage of employment is the town of Chadron, Nebraska and the population with the highest unemployment rate is Spearfish, South Dakota.

Within the Nebraska-South Dakota-Wyoming Uranium Milling Region, the county with the highest percentage of employment is Campbell County, Wyoming and the county with the highest unemployment rate is Shannon County, Nebraska. The towns with the highest unemployment rate are located on the Pine Ridge Indian Reservation (Table 3.4-20).

3.4.10.4.1 State Data

3.4.10.4.1.1 Nebraska

The State of Nebraska has an employment rate of 66.7 percent and unemployment rate of 2.5 percent. The largest sector of employment is management, professional, and related occupations at 33.0 percent. The largest type of industry is educational, health, and social services at 20.7 percent. The largest class of worker is private wage and salary workers at 77.1 percent (U.S. Census Bureau, 2007).

Gering

Gering has an employment rate of 61.6 percent and unemployment rate the same as that of the state at 2.5 percent. The largest sector of employment is management, professional, and related occupations at 34.0 percent. The largest type of industry is educational, health, and social services. The largest class of worker is private wage and salary workers (U.S. Census Bureau, 2008).

Scottsbluff

Scottsbluff has an employment rate of 57.6 percent and unemployment rate much higher than that of the state at 4.6 percent. The largest sector of employment is management, professional, and related occupations at 29.6 percent. The largest type of industry is educational, health, and social services. The largest class of worker is private wage and salary workers (U.S. Census Bureau, 2008).

Alliance

Alliance has an employment rate of 63.1 percent and unemployment rate higher than that of the state at 3.6 percent. The largest sector of employment is production, transportation, and

Description of the Affected Environment

material moving occupations at 25.9 percent. The largest type of industry is transportation and warehousing, and utilities. The largest class of worker is private wage and salary workers (U.S. Census Bureau, 2008).

**Table 3.4-20. Employment Structure of the Pine Ridge Indian Reservation
Within the Affected Area***

Affected Environment	2003 Labor Force Population	Unemployed as Percent of Labor Force	Employed Below Poverty Guidelines	
Oglala Sioux Tribe of Pine Ridge	27,778	87%	716	21%

* U.S. Department of the Interior. "Affairs American Indian Population and Labor Force Report 2003." <<http://www.doi.gov/bia/labor.html>>. Washington, DC: U.S. Department of the Interior, Bureau of Indian Affairs, Office of Tribal Affairs. 2003.

Chadron

Chadron has an employment rate of 65.2 percent and unemployment rate lower than that of the state at 2.8 percent. The largest sector of employment is management, professional, and related occupations at 29.2 percent. The largest type of industry is educational, health, and social services. The largest class of worker is private wage and salary workers (U.S. Census Bureau, 2008).

3.4.10.4.1.2 South Dakota

The State of South Dakota has an employment rate of 64.9 percent and unemployment rate of 3.0 percent. The largest sector of employment is management, professional, and related occupations at 32.6 percent. The largest type of industry is educational, health, and social services at 22.0 percent. The largest class of worker is private wage and salary workers at 72.9 percent (U.S. Census Bureau, 2007).

Rapid City

Rapid City has an employment rate of 63.7 percent and unemployment rate higher than that of the state at 3.2 percent. The largest sector of employment is management, professional, and related occupations at 32.8 percent. The largest type of industry is educational, health, and social services. The largest class of worker is private wage and salary workers (U.S. Census Bureau, 2008).

Spearfish

Spearfish has an employment rate of 53.5 percent and unemployment rate much higher than that of the state at 11.5 percent. The largest sector of employment is management, professional, and related occupations at 33.5 percent. The largest type of industry is educational, health, and social services. The largest class of worker is private wage and salary workers (U.S. Census Bureau, 2008).

Sturgis

Sturgis has an employment rate of 59.5 percent and unemployment rate lower than that of the state at 2.8 percent. The largest sector of employment is sales and occupations at 27.6 percent. The largest type of industry is educational, health, and social services. The largest class of worker is private wage and salary workers (U.S. Census Bureau, 2008).

3.4.10.4.1.3 Wyoming

The State of Wyoming has an employment rate of 63.1 percent and unemployment rate of 3.5 percent. The largest sector of employment is sales and office occupations. The largest type of industry is educational, health, and social services. The largest class of worker is private wage and salary workers (U.S. Census Bureau, 2007).

Casper

Casper has an employment rate of 64.9 percent and an unemployment rate lower than that of the state at 3.4 percent. The largest sector of employment is sales and office occupations at 30.6 percent followed by management, professional, and related occupations at 29.7 percent. The largest type of industry is educational, health, and social services at 22.1 percent. The largest class of worker is private wage and salary workers at 76.6 percent (U.S. Census Bureau, 2007).

3.4.10.4.2 County Data

3.4.10.4.2.1 Nebraska

Dawes County

Dawes County has an employment rate of 63.8 percent and unemployment rate slightly higher than that of the state at 2.7 percent. The largest sector of employment is management, professional, and related occupations at 32.4 percent. The largest type of industry is educational, health, and social services at 28.9 percent. The largest class of worker is private wage and salary workers at 58.8 percent (U.S. Census Bureau, 2007).

Sioux County

Sioux County has an employment rate of 62.1 percent and unemployment rate slightly higher than that of the state at 2.7 percent. The largest sector of employment is management, professional, and related occupations at 50.3 percent. The largest type of industry is agriculture, forestry, fishing and hunting, and mining at 40.5 percent. The largest class of worker is private wage and salary workers at 52.8 percent (U.S. Census Bureau, 2008).

3.4.10.4.2.2 South Dakota

Butte County

Butte County has an employment rate of 64.3 percent and unemployment rate higher than that of the state at 3.9 percent. The largest sector of employment is management, professional, and related occupations at 27.0 percent. The largest type of industry is agriculture, forestry, fishing,

Description of the Affected Environment

and hunting, and mining at 19.4 percent. The largest class of worker is private wage and salary workers at 66.8 percent (U.S. Census Bureau, 2008).

Custer County

Custer County has an employment rate of 57.5 percent and unemployment rate lower than that of the state at 2.0 percent. The largest sector of employment is management, professional, and related occupations at 34.6 percent. The largest type of industry is educational, health, and social services at 20.6 percent. The largest class of worker is private wage and salary workers at 58.5 percent (U.S. Census Bureau, 2007).

Fall River County

Custer County has an employment rate of 52.9 percent and unemployment rate higher than that of the state at 3.9 percent. The largest sector of employment is management, professional, and related occupations at 34.7 percent. The largest type of industry is educational, health, and social services at 31.1 percent. The largest class of worker is private wage and salary workers at 58.2 percent (U.S. Census Bureau, 2007).

Shannon County

Shannon County has an employment rate of 35.1 percent and unemployment rate considerably higher than that of the state at 17.3 percent. The largest sector of employment is management, professional, and related occupations at 37.8 percent. The largest type of industry is educational, health and social services. The largest class of worker is government workers (U.S. Census Bureau, 2008).

3.4.10.4.2.3 Wyoming

Campbell County

Campbell County has an employment rate of 73.2 percent and an unemployment rate lower than that of the state at 3.4 percent. The largest sector of employment is management, professional, and related occupations at 23.9 percent followed by construction, extraction, and maintenance occupations at 23.7 percent. The largest type of industry is agriculture, forestry, fishing and hunting, and mining at 23.3 percent followed by educational, health, and social services at 16.7 percent. The largest class of worker is private wage and salary workers at 78.4 percent (U.S. Census Bureau, 2007).

Crook County

Crook County has an employment rate of 62.2 percent and an unemployment rate lower than that of the state at 2.1 percent. The largest sector of employment is management, professional, and related occupations at 29.9 percent. The largest type of industry is agriculture, forestry, fishing and hunting, and mining at 24.7 percent. The largest class of worker is private wage and salary workers at 59.5 percent (U.S. Census Bureau, 2007).

Niobrara County

Niobrara County has an employment rate of 59.4 percent and an unemployment rate lower than that of the state at 2.1 percent. The largest sector of employment is management, professional,

and related occupations at 34.4 percent. The largest type of industry is agriculture, forestry, fishing and hunting, and mining at 24.7 percent. The largest class of worker is private wage and salary workers at 62.6 percent (U.S. Census Bureau, 2008).

Weston County

Weston County has an employment rate of 56.6 percent and an unemployment rate lower than that of the state at 3.3 percent. The largest sector of employment is management, professional, and related occupations at 24.3 percent. The largest type of industry is agriculture, forestry, fishing and hunting, and mining at 22.4 percent. The largest class of worker is private wage and salary workers at 68.9 percent (U.S. Census Bureau, 2008).

3.4.10.4.3 Native American Communities

Information on labor force and poverty levels for the Pine Ridge Indian Reservation is based on 2003 Bureau of Indian Affairs data and is provided in Table 3.4-20. The Oglala Sioux Tribe reports unemployment rates of more than 80 percent, much higher than the statewide levels that range from 2.5 percent for Nebraska to 3.5 percent for Wyoming (U.S. Census Bureau, 2007; U.S. Department of the Interior, 2003).

3.4.10.5 Local Finance

Local finance information such as revenue and tax information for the affected environment is provided in the following sections.

3.4.10.5.1 Nebraska

Sources of revenue for the State of Nebraska come from income, sales, cigarette, motor, and lodging taxes. Personal income tax rates for Nebraska range from 2.56 percent to 6.84 percent. The sales and use tax rate is 5.5 percent. Information on "ad valorem taxes" or mineral taxes such as that from uranium extraction is not available (Nebraska Department of Revenue, 2007). Information on local finance for the affected communities within the region of influence is presented next.

Dawes County

Sources of revenue for Dawes County come from real estate and property taxes. The net property taxes levied in 2003 were \$1,634,113 with a state aid of \$634,793 (Nebraska Department of Revenue, 2007).

Sioux County

Sources of revenue for Sioux County come from real estate and property taxes (Nebraska Department of Revenue, 2007).

3.4.10.5.2 South Dakota

Sources of revenue for the State of South Dakota come from 36 different state taxes. These taxes are grouped into four main categories: sales, use, and contractors' excise taxes; motor fuel taxes; motor vehicle fees and taxes; and special taxes. Once collected, these tax revenues are distributed into the state's general fund, local units of government, and the state highway

Description of the Affected Environment

fund. In 2006, 72 percent came from sales, use, and contractors' excise taxes; 11 percent from motor fuel taxes; 9 percent from special taxes; and 8 percent from vehicle taxes. South Dakota also imposes an energy minerals tax on owners of energy minerals (such as uranium). In 2006, the tax rate base was 4.5 percent of the taxable value and approximately 50 percent was disbursed to local government (South Dakota Department of Revenue and Regulation, 2007). Information on local finance for the affected communities within the region of influence is presented next.

Butte County

The majority of revenue for Butte County comes from sales, use, and property taxes. In 2004, a total revenue of \$1,578,000 was collected from property taxes (City-Data.com, 2008).

Custer County

The majority of revenue for Custer County is from property taxes. In 2006, there were approximately 13,000 parcels of land in Custer County and \$9.3 million was collected in real estate taxes. Other sources of revenue come from motor vehicle fees (Custer County South Dakota, 2007).

Fall River County

In 2004, the majority of revenue for Fall River County was from property taxes (\$2,101,000) and motor vehicle fees (\$482,000) (City-Data.com, 2007).

Shannon County

The majority of revenue for Shannon County comes from retail sales at \$30,594 as of 2002 and federal grants at \$197,565 as of 2004 (US Census Bureau, 2008).

3.4.10.5.3 Wyoming

The State of Wyoming does not have an income tax nor does it assess tax on retirement income received from another state. Wyoming has a 4 percent state sales tax, 2 percent to 5 percent county lodging tax, and 5 percent use tax. Counties have the option of collecting an additional 1 percent tax for general revenue and 2 percent tax for specific purposes. Wyoming also imposes "ad valorem taxes" on mineral extraction properties. Sales and use tax distribution information for the affected counties is presented in Table 3.4-21.

3.4.10.5.4 Native American Communities

The Pine Ridge Indian Reservation is the poorest reservation in the United States. The majority of revenue for Pine Ridge comes from employment by the Oglala Sioux Tribe, Oglala Lakota College, Bureau of Indian Affairs, and the Indian Health Service. Some revenue also comes from agricultural production, gaming, hunting, and ranching (Housing Assistance Council, 2002)).

1

Table 3.4-21. 2007 Sales and Use Tax Distribution of the Affected Counties Within Wyoming (Through September 28, 2007)					
Affected Counties	Use Tax		Sales Tax		Lodging Option Tax
	General	Specific	General	Specific	
Campbell County	\$387,522.93	\$97,111.27	\$2,334,282.49	\$583,201.87	\$0.0
Crook County	\$23,375.38	\$83,017.39	\$23,325.92	\$82,636.59	\$10,096.20
Niobrara County	\$6,119.06	\$34,411.65	\$6,119.06	\$34,411.65	\$5,137.77
Weston County	\$28,152.44	\$0.0	\$60,466.76	\$0.0	\$6,682.25
* Wyoming Department of Revenue. "Sales and Tax Distribution Report by County 2007." < http://revenue.state.wy.us/PortalVBVS/DesktopDefault.aspx?tabindex=3&tabid=10 > (18 October 2007, 25 February 2008, and April 25, 2008).					

2

3 **3.4.10.6 Education**

4

5 Information on education for the affected communities is presented in the following paragraphs.

6

7 Based on review of the affected environment, the county with the largest number of schools is

8 Campbell County, WY and the county with the smallest number of schools is Niobrara, WY.

9 The towns with the smallest number of schools or smaller schools are located on the Pine Ridge
10 Indian Reservation.

11

12 **3.4.10.6.1 Nebraska**

13

14 Dawes County

15

16 Dawes County has a total of 17 schools including public schools, elementary schools, middle
17 schools, high schools, and 1 academy. There are a total of approximately 5,500 students. The
18 majority of schools provide bus services (Schoolbug.org, 2007a).

19

20 Sioux County

21

22 Sioux County has a total of 6 schools including 5 public schools and 1 high school, with a total
23 of approximately 565 students. Information as to whether these schools provide bus services is
24 not available (Publicschoolsreport.com, 2008).

25

26 **3.4.10.6.2 South Dakota**

27

28 Butte County

29

30 Butte County has 3 elementary schools, 2 middle schools, and 2 high schools. There are a total
31 of approximately 1,789 students. Information as to whether these schools provide bus services
32 is not available (Schoolbug.org, 2008).

33

34

Description of the Affected Environment

Custer County

Custer County has 5 elementary schools, 1 middle school, 1 high school, and 1 alternative school for a total of nine schools. There are a total of approximately 1,207 students. Information as to whether these schools provide bus services is not available (Schoolbug.org, 2007b).

Fall River County

Fall River County has 4 elementary schools, 2 middle schools, and 1 junior high school, and 3 high schools for a total of 10 schools. There are a total of approximately 1,200 students. Information as to whether these schools provide bus services is not available (Schoolbug.org, 2007c).

Shannon County

Shannon County has one school district, which consists of 4 elementary and junior high schools. There are approximately 991 students. Information as to whether these schools provide bus services is not available (Greatschools, 2008d).

Native American Communities

The Pine Ridge Indian Reservation has the Pine Ridge School and the Oglala elementary school (Housing Assistance Council, 2002; Pine Ridge School, 2008). Specific information pertaining to school population or bus services is not available.

3.4.10.6.3 Wyoming

Campbell County

Campbell County has 1 school district with 24 schools consisting of 15 elementary schools, 2 junior high schools, 1 junior/senior high school, 1 high school, 1 alternative school, and 1 aquatic center. There are a total of approximately 7,441 students. The majority of schools provide bus services (Campbell County School District No. 1, 2007).

Crook County

Crook County has 1 school district with 2 elementary schools, 2 secondary schools, and 1 high school, with a total of approximately 1,142 students. Information as to whether these schools provide bus services is not available (Crook County School District, 2008).

Niobrara County

Niobrara County has one school district, Niobrara County School District No. 1, with a total of approximately 422 students. There are 1 elementary and middle schools, 1 high school, and 1 private school. Information as to whether these schools provide bus services is not available (Niobrara County School District No. 1, 2008).

Weston County

Weston County has one school district, Weston County School District No. 1, with a total of approximately 1,134 students. There are 2 elementary schools, 1 middle school, and 1 high school. Information as to whether these schools provide bus services is not available (Weston County School District No. 1, 2008).

3.4.10.7 Health and Social Services

The majority of health care facilities are located within populated areas of the affected environment. The closest health care facilities within the vicinity of the potential ISL facilities are located in Spearfish, Edgemont, Rapid City and Sturgis, South Dakota; Alliance, Gordon, and Chadron, Nebraska; Gillette, Sundance, and Torrington, Wyoming, and have a total of at least 18 facilities (MapQuest, 2008b). These consist of hospitals, clinics, emergency centers, and medical services. The following hospitals are located proximate to the Nebraska-South Dakota-Wyoming Uranium Milling Region: Spearfish, South Dakota (1), Rapid City, South Dakota (2), Alliance, Nebraska (1), Gordon, Nebraska (1), Chadron, Nebraska (2), Gillette, Wyoming (2), and Torrington, Wyoming (1).

Local police within the Nebraska-South Dakota-Wyoming Uranium Milling Region are under the jurisdiction of each county. There are 20 police, sheriff, or marshals offices within the region: Butte County, South Dakota (2), Custer County, South Dakota (1), Fall River County, South Dakota (2), Shannon County, South Dakota (1), Dawes County, Nebraska (3), Sioux County, Nebraska (1), Campbell County, Wyoming (2), Crook County, Wyoming (3), Niobrara County, Wyoming (2), and Weston County, Wyoming (3) (usacops, 2008c).

Fire departments within the affected area are comprised at the County, town or CBSA level. There are 45 fire departments within the milling region: Rapid City, South Dakota (16), Sturgis, South Dakota (14), Spearfish, South Dakota (5), Alliance, Nebraska (1), Campbell County, Wyoming (2), Crook County, Wyoming (1), and Gillette, Wyoming (2) (50states, 2008).

3.4.11 Public and Occupational Health

3.4.11.1 Background Radiological Conditions

For a U.S. resident, the average total effective dose equivalent from natural background radiation sources is approximately 3 mSv/yr [300 mrem/yr] but varies by location and elevation (National Council of Radiation Protection and Measurements, 1987). In addition, the average American receives 0.6 mSv/yr [60 mrem/yr] from man-made sources including medical diagnostic tests and consumer products (National Council of Radiation Protection & Measurements 1987). Therefore the total from natural background and man-made sources for the average U.S. resident is 3.6 mSv/yr [360 mrem/yr]. For a breakdown of the sources of this radiation, see Figure 3.2-22.

Background dose varies by location primarily because of elevation changes and variations in the dose from radon. As elevation increases so does the dose from cosmic radiation and hence the total dose. Radon is a radioactive gas produced from the decay of ^{238}U , which is naturally found in soil. The amount of radon in the soil/bedrock depends on the type the porosity and moisture content. Areas which have types of soils/bedrock like granite and limestone have higher radon levels than those with other types of soils/bedrock (EPA, 2006).

Description of the Affected Environment

Radiological background for Wyoming is provided in Section 3.2.11.1. For the States of South Dakota and Nebraska the average background rate including natural and manmade sources is 6.0 mSv/yr [600 mrem/yr] and 3.5mSv/yr [350 mrem/yr], respectively (EPA, 2006). The average background rate for South Dakota is significantly higher than the U.S. average background rate of 3.6 mSv/yr [360 mSv/yr] and for Nebraska it is very similar.

For South Dakota, the radon dose is 4.4 mSv/yr [440 mrem/yr] compared to the U.S. average radon dose of 2.0 mSv/yr [200 mrem/yr]. For South Dakota, the indoor average radon rate is significantly higher than the U.S. average due to geological reasons as well as poor ventilation within homes (EPA, 2006). For the western region of South Dakota which of interest here, the radon levels are half as much when compared to the state average (South Dakota Department of Environmental and Natural Resources, 2008) and therefore, background dose is expected to be closer to the national average for this region.

3.4.11.2 Public Health and Safety

Public health and safety standards are the same regardless of a facility's location. Therefore, see Section 3.2.11.2 for further discussion of these standards.

3.4.11.3 Occupational Health and Safety

Occupational health and safety standards are the same regardless of facility's location. Therefore, see Section 3.2.11.3 for further discussion of these standards.

3.4.12 References

BLM. "Visual Resource Management." Manual 8400. Washington, DC: BLM. 2007a. <<http://www.blm.gov/nstc/VRM/8400.html#Anchor-.06-23240>> (17 October 2007).

BLM. "Visual Resource Inventory." Manual H-8410-1. Washington, DC: BLM. 2007b. <<http://www.blm.gov/nstc/VRM/8410.html>> (17 October 2007).

BLM. "Visual Resource Contrast Rating." Manual 8431. Washington, DC: BLM. 2007c. <<http://www.blm.gov/nstc/VRM/8431.html>> (17 October 2007).

BLM. "Newcastle Resource Management Plan." Newcastle, Wyoming: BLM, Newcastle Field Office. 2000. <http://www.blm.gov/rmp/WY/application/rmp_browse.cfm?hlevel=1&rmpid=89&idref=28910> (17 October 2007).

BLM. "Nebraska Record of Decision and Approved Resource Management Plan." Casper, Wyoming: BLM, Casper District Office. 1992. <<http://www.blm.gov/rmp/WY/Nebraska/rmp.pdf>> (17 October 2007).

BLM. "Final Resource Management Plan, South Dakota Resource Area, Miles City District." Miles City, Montana: BLM, South Dakota Resource Area, Miles City District. 1985. <http://www.blm.gov/mt/st/en/prog/planning/south_dakota.html> (17 October 2007).

Bryce, S.A., J.M. Omernik, D.A. Pater, M. Ulmer, J. Schaar, J. Freeouf, R. Johnson, P. Kuck, and S.H. Azevedo. "Ecoregions of North Dakota and South Dakota." U.S. Geological Survey. Scale 1:1,500,000. Reston, Virginia: U.S. Geological Survey. 1996.

- 1 Bureau of Indian Affairs. "American Indian Population and Labor Force Report." 2003.
2
- 3 Chapman, S.S., S.A. Bryce, J.M. Omernik, D.G. Despain, J. ZumBerge, and M. Conrad.
4 "Ecoregions of Wyoming." U.S. Geological Survey Map. Scale 1:1,400,000. 2004.
5
- 6 Chapman, S.S., J.M. Omernik, J.A. Freeouf, D.G. Huggins, J.R. McCauley, C.C. Freeman,
7 G. Steinauer, R.T. Angelo, and R.L. Schlepp. "Ecoregions of Nebraska and Kansas."
8 U.S. Geological Survey Map. Scale 1:1,950,000. Reston, Virginia: U.S. Geological Survey.
9 2001.
10
- 11 Chenoweth, W.L. "Geology and Production History of the Uranium Deposits in the Northern
12 Black Hills, Wyoming—South Dakota." Eastern Powder River Basin, Wyoming Geological
13 Association, 39th Annual Field Conference Guidebook. Casper, Wyoming: Wyoming
14 Geological Association. pp. 263–270. 1988.
15
- 16 City-Data.com. "Butte County, South Dakota." 2008. <www.city-data.com>
17 (28 February 2008).
18
- 19 City-Data.com. "Fall River County, South Dakota." 2007. <www.city-data.com>
20 (14 October 2007).
21
- 22 Collings, S.F. and R.H. Knode. "Geology and Discovery of the Crow Butte Uranium Deposit,
23 Dawes County, Nebraska." Practical Hydromet '83, 7th Annual Symposium on Uranium and
24 Precious Metals. Littleton, Colorado: American Institute of Mining, Metallurgical, and Petroleum
25 Engineering. 1984.
26
- 27 Crook County School District. "Crook County, Wyoming." 2008. <www.crooknet.k12.wy.us>
28 (25 April 2008).
29
- 30 Custer County South Dakota. "Custer County." 2007. <www.custercountysd.com/>
31 (15 October 2007).
32
- 33 DeGraw, H.M. "Subsurface Relations of the Cretaceous and Tertiary in Western Nebraska."
34 MS Thesis. University of Nebraska. Lincoln, Nebraska. 1969.
35
- 36 Dondanville, R.F. "The Fall River Formation, Northwestern Black Hills; Lithology and Geology
37 History." Joint Field Conference Guidebook. Northern Powder River Basin, Wyoming:
38 Wyoming Geological Association, Billings Geological Society. pp. 87–99. 1963.
39
- 40 Driscoll, F.G. "Groundwater and Wells." Second edition. St Paul, Minnesota: Johnson
41 Filtration Systems Inc. 1986.
42
- 43 Driscoll, D.G., J.M. Carter, J.E. Williamson, and L.D. Putnam. "Hydrology of the Black Hills
44 Area, South Dakota." U.S. Geological Survey Water Resources Investigation Report 02-4094.
45 2002.
46
- 47 EPA. "National Assessment Database." 2008. <<http://www.epa.gov/waters/305b/index.html>>
48 (28 February 2008).
49

Description of the Affected Environment

- 1 EPA. "Counties Designate Nonattainment or Maintenance for Clean Air Act's National Ambient
2 Air Quality Standards (NAAQS)." 2007. <[http://www.epa.gov/oar/oaqps/greenbk/
3 mapnmpoll.html](http://www.epa.gov/oar/oaqps/greenbk/mapnmpoll.html)> (29 September 2007).
4
- 5 EPA. "Prevention of Significant Deterioration (PSD) Permit Program Status: May 2007."
6 2007b. <<http://www.epa.gov/nsr/where.html>> (26 September 2007).
7
- 8 EPA. "Assessment of Variations in Radiation Exposure in the United States (Revision 1)."
9 Contract Number EP-D-05-02. Washington, DC: EPA. 2006.
10
- 11 Federal Highway Administration. "Synthesis of Noise Effects on Wildlife Populations."
12 FHWA-HEP-06-016. Washington, DC: Federal Highway Administration, Department of
13 Transportation. 2004.
14
- 15 Girardin, J. "A List of Areas Designated Unique and Irreplaceable or Designated Rare or
16 Uncommon by the Council." Letter (November 29) to T. Lorenzon. Cheyenne, Wyoming:
17 Council on Environmental Quality. 2006.
18
- 19 Gjelsteen, T.W. and S.P. Collings. "Relationship Between Groundwater Flow and Uranium
20 Mineralization in the Chadron Formation, Northwest Nebraska." Eastern Powder River Basin,
21 Wyoming Geological Association, 39th Annual Field Conference Guidebook. Casper, Wyoming:
22 Wyoming Geological Association. pp. 271-284. 1988.
23
- 24 Gott, G.B, D.E. Wolcott, and C.G. Bowles. "Stratigraphy of the Inyan Kara Group and
25 Localization of Uranium Deposits, Southern Black Hills, South Dakota and Wyoming."
26 U.S. Geological Survey Professional Paper 763. 1974.
27
- 28 Greatshools.com. "Shannon County, South Dakota." 2008. <<http://www.greatshools.com>.
29 (16 April 2008).
30
- 31 Harshman, E.N. "Uranium Deposits of Wyoming and South Dakota." *Ore Deposits in the*
32 *United States 1933-1967*. New York City, New York: American Institute of Mining,
33 Metallurgical, and Petroleum Engineers. pp. 815-831. 1968.
34
- 35 Hart, O.M. "Uranium in the Black Hills." *Ore Deposits of the United States, 1933-1967*.
36 J.D. Ridge, ed. New York City, New York: American Institute of Mining, Metallurgical, and
37 Petroleum Engineers. pp. 832-837. 1968.
38
- 39 Henebry, G.M., B.C. Putz, M.R. Vaitkus, and J.W. Merchant. "The Nebraska Gap Analysis
40 Project Final Report." Lincoln, Nebraska: University of Nebraska, School of Natural Resources.
41 2005.
42
- 43 Housing Assistance Council. "Taking Stock: Rural People, Poverty, and Housing at the Turn of
44 the 21st Century." Washington, DC: Housing Assistance Council. December 2002.
45
- 46 Kansas Geological Survey. Open-File Report. 91-1. 1991. <[http://www.kgs.ku.edu/Dakota/
47 vol3/fy91/rep06.htm](http://www.kgs.ku.edu/Dakota/vol3/fy91/rep06.htm)> (29 April 2008).
48
- 49 King, T. *Places That Count: Traditional Cultural Properties in Cultural Resources Management*.
50 Walnut Creek, California: Altamira Press. 2003.
51

1 MapQuest. "Nebraska, South Dakota, and Wyoming." 2008. <<http://mapquest.com>> (16 April
2 2008).

3
4 Maxwell, C.H. "Map and Stratigraphic Sections Showing Distribution of Some Channel
5 Sandstones in the Lakota Formation, Northwestern Black Hills, Wyoming. U.S. Geological
6 Survey Miscellaneous Field Studies Map MF-632. 1974.

7
8 Miller J.A. and C.L. Appel. "Groundwater Atlas of the United States, Kansas, Missouri, and
9 Nebraska." U.S. Geological Survey Report HA 730-D. Denver, Colorado: U.S. Geological
10 Survey. <http://capp.water.usgs.gov/gwa/ch_d/index.html> 1997.

11
12 Munn, L.C. and C.S. Arneson. "Soils of Wyoming—A Digital Statewide Map at 1:500,000-
13 Scale." B-1069. Laramie, Wyoming: University of Wyoming Agricultural Experiment Station,
14 College of Agriculture. 1998.

15
16 National Climatic Data Center. "NCDC U.S. Storm Events Database." 2007.
17 <<http://www4.ncdc.noaa.gov/cgi-win/wwcgi.dll?wwEvent~Storms>> (14 April 2008).

18
19 National Climatic Data Center. "Climates of the States, Climatology of the United States No. 60
20 (New Mexico, Nebraska, South Dakota, and Wyoming)." Asheville, North Carolina: National
21 Oceanic and Atmospheric Administration. 2005. <[http://cdo.ncdc.noaa.gov/cgi-
22 bin/climatenormals/climatenormals.pl?directive=prod_select2&prodtype=CLIM60&subnum=>](http://cdo.ncdc.noaa.gov/cgi-bin/climatenormals/climatenormals.pl?directive=prod_select2&prodtype=CLIM60&subnum=>)
23 (30 January 2005).

24
25 National Climatic Data Center. "Climatography of the United States No. 20: Monthly Station
26 Climate Summaries, 1971–2000." Asheville, North Carolina: National Oceanic and
27 Atmospheric Administration. 2004.

28
29 National Council on Radiation Protection and Measurements. "Report No. 094—Exposure of
30 the Population in the United States and Canada From Natural Background Radiation."
31 Bethesda, Maryland: National Council on Radiation Protection & Measurements. 1987.

32
33 National Weather Service. "NOAA Technical Report NWS 33: Evaporation Atlas for the
34 Contiguous 48 United States." Washington, DC: National Oceanic and Atmospheric
35 Administration. 1982.

36
37 Nebraska Department of Environmental Quality. "Total Maximum Daily Loads for the White
38 River Hat Creek Basin." Lincoln, Nebraska: Nebraska Department of Environmental Quality.
39 December 2005a.

40
41 Nebraska Department of Environmental Quality. "Total Maximum Daily Loads for the Niobrara
42 River Basin." Lincoln, Nebraska: Nebraska Department of Environmental Quality.
43 December 2005b.

44
45 Nebraska Department of Environmental Quality. "Ambient Air Quality Regulations." 2002.
46 <<http://www.deq.state.ne.us/>> (23 October 2007).

47
48 Nebraska Department of Revenue. 2007. <www.revenue.ne.gov/> (15 October 2007).
49

Description of the Affected Environment

- 1 Nebraska Game and Parks Commission. "Park Areas in the Panhandle Region." Lincoln,
2 Nebraska: Nebraska Game and Parks Commission. 2008.
3 <<http://www.ngpc.state.ne.us/parks/guides/parksearch/getregion.asp?District=1>> (29 February
4 2008).
5
6 NRC. "Environmental Assessment for Renewal of Source Material License No. SUA-1534:
7 Crow Butte Resources, Incorporated Crow Butte Uranium Project Dawes County, Nebraska.
8 Docket No. 40-8943. Washington, DC: NRC. 1998
9
10 NRC. NUREG-1508, "Final Environmental Impact Statement to Construct and Operate the
11 Crown Point Uranium Solution Mining Project, Crown Point, New Mexico." Washington, DC:
12 NRC. February 1997.
13
14 NRCS. "United States Department of Agriculture, National Resources Conservation Service.
15 Dawes County, Nebraska." <<http://soildatamart.nrcs.usda.gov/survey.aspx?County=NE045>>
16 October 2007.
17
18 Parker, P. and T. King. "Guidelines for Evaluating and Documenting Traditional Cultural
19 Properties." National Register Bulletin 38. Washington, DC: National Park Service. 1998.
20
21 Pine Ridge School. "Pine Ridge School, Pine Ridge, South Dakota." 2008. <www.prs.bia.edu>
22 (29 April 2008).
23
24 Publicschoolsreport.com. "Sioux County, Nebraska." 2008. <publicschoolsreport.com>
25 (27 February 2008).
26
27 Renfro, A.R. "Uranium Deposits in the Lower Cretaceous of the Black Hills." *Contributions to*
28 *Geology*. Laramie, Wyoming: University of Wyoming. Vol. 8, No. 2-1. pp. 87-92. 1969.
29
30
31 Robinson, C.S., W.J. Mapel, and M.H. Bergendahl. "Stratigraphy and Structure of the Northern
32 and Western Flank of the Black Hills Uplift, Wyoming, Montana, and South Dakota."
33 U.S. Geological Survey Professional Paper 404. 1964.
34
35 Schoolbug.org. "Butte County, South Dakota." 2008. <www.schoolbug.org> (27 February
36 2008).
37
38 Schoolbug.org. "Custer County, South Dakota." 2007a. <www.schoolbug.org>
39 (13 October 2007).
40
41 Schoolbug.org. "Fall River County, South Dakota." 2007b. <www.schoolbug.org>
42 (13 October 2007).
43
44 Schoolbug.org. "Dawes County, Nebraska." 2007c. <www.schoolbug.org> (13 October 2007).
45
46 South Dakota Birds and Birding. "American Dipper." 2008. <[http://sdakotabirds.com/](http://sdakotabirds.com/species/american_dipper_info.htm)
47 [species/american_dipper_info.htm](http://sdakotabirds.com/species/american_dipper_info.htm)> (12 February 2008).
48
49 South Dakota Department of Environmental and Natural Resources. "Radon."
50 <<http://www.state.sd.us/denr/DES/AirQuality/radon1.htm>> 2008. (15 April 2008).
51

1 South Dakota Department of Environment and Natural Resources. "Air Quality Program."
2 2007. <<http://www.state.sd.us/denr/DES/AirQuality/airprogr.htm>> (23 October 2007).

3
4 South Dakota Department of Revenue and Regulation. "South Dakota Department of Revenue
5 and Regulation." 2007. <www.state.sd.us/drr2/revenue.html> (18 October 2007).

6
7 South Dakota Division of Wildlife. "South Dakota Statewide Fisheries Survey." Angostura
8 Reservoir County: Fall River, No. 2102-/F21-R-37. 2004. <[http://www.sdgfp.info/](http://www.sdgfp.info/wildlife/Fishing/WesternLakes/Angostura04.pdf)
9 [wildlife/Fishing/WesternLakes/Angostura04.pdf](http://www.sdgfp.info/wildlife/Fishing/WesternLakes/Angostura04.pdf)> (04 April 2008).

10
11 South Dakota Game, Fish, and Parks. "Fishing in South Dakota." Pierre, South Dakota:
12 South Dakota Game, Fish, and Parks. 2008 <<http://www.sdgfp.info/Wildlife/fishing/>>
13 (15 February 2008).

14
15 South Dakota State University. "South Dakota GAP Analysis Project. "Brookings,
16 South Dakota: South Dakota State University, Department of Wildlife and Fisheries Sciences.
17 2007. <<http://wfs.sdstate.edu/sdgap/sdgap.htm>> (29 November 2007).

18
19 South Dakota State University. "South Dakota GAP Analysis Project. "Brookings, South
20 Dakota: South Dakota State University, Department of Wildlife and Fisheries Sciences. 2001.
21 <<http://wfs.sdstate.edu/sdgap/sdgap.htm>> (29 February 2008).

22
23 University of Nebraska. "Gap Analysis Project." Lincoln, Nebraska: University of Nebraska.
24 2007.

25
26 Usacops. South Dakota, Nebraska, and Wyoming. 2008. <<http://www.usacops.com>>.
27 (16 April 2008).

28
29 U.S. Census Bureau. "American FactFinder 2000 Census Data." 2008.
30 <<http://factfinder.census.gov>> (25 February 2008).

31 U.S. Census Bureau. "QuickFacts." <http://factfinder.census.gov>>. (16 April 2008).

32
33 USFS. "Medicine Bow-Routt National Forests, Thunder Basin National Grassland."
34 Laramie, Wyoming: USFS. 2008a. <<http://www.fs.fed.us/r2/mbr/index.shtml>>
35 (27 February 2008).

36
37 USFS. "Nebraska & Samuel R. McKelvie National Forests Buffalo Gap, Fort Pierre, and Oglala
38 National Grasslands." Chadron, Nebraska: USDA Forest Service Nebraska, Samuel R.
39 McKelvie National Forests Buffalo Gap, Fort Pierre, and Oglala National Grasslands. 2008b.

40
41 USFS. "Black Hills National Forest, Phase II Amendment: 1997 Land and Resource
42 Management Plan Final Environmental Impact Statement." Custer, South Dakota: USFS,
43 Black Hills National Forest. 2005. <[http://www.fs.fed.us/r2/blackhills/](http://www.fs.fed.us/r2/blackhills/projects/planning/amendments/phase_II/index.shtml)
44 [projects/planning/amendments/phase_II/index.shtml](http://www.fs.fed.us/r2/blackhills/projects/planning/amendments/phase_II/index.shtml)> (17 October 2007).

45
46 USFS. "Land and Resource Management Plan for the Nebraska National Forest and
47 Associated Units (Rocky Mountain Region)." Chadron, Nebraska: USFS, Nebraska National
48 Forest. 2001a. <http://www.fs.fed.us/ngp/plan/feis_plan_nebraska.htm> (17 October 2007).

Description of the Affected Environment

- 1 USFS. "Black Hills National Forest, Phase I Amendment: 1997 Land and Resource
2 Management Plan Environmental Assessment." Custer, South Dakota: USFS, Black Hills
3 National Forest. 2001b. <[http://www.fs.fed.us/r2/blackhills/projects/planning/
4 amendments/phase_i/index.shtml](http://www.fs.fed.us/r2/blackhills/projects/planning/amendments/phase_i/index.shtml)> (17 October 2007).
5
- 6 USFS. "Revised Land and Resource Management Plan for the Black Hills National Forest."
7 Custer, South Dakota: USFS, Black Hills National Forest. 1997. <[http://www.fs.fed.us/r2/
8 blackhills/projects/planning/97Revision/fp/index.shtml](http://www.fs.fed.us/r2/blackhills/projects/planning/97Revision/fp/index.shtml)> (17 October 2007).
9
- 10 U.S. Geological Survey. 2008a. "Water Watch." <<http://water.usgs.gov/waterwatch>>
11 (28 February 2008).
12
- 13 U.S. Geological Survey. "Fragile Legacy Endangered, Threatened and Rare Animals of
14 South Dakota." 2008b. <[http://www.npwrc.usgs.gov/resource/wildlife/
15 sdrare/index.htm#contents](http://www.npwrc.usgs.gov/resource/wildlife/sdrare/index.htm#contents)> (1 March 2008).
16
- 17 U.S. Geological Survey. "A Tapestry of Time and Terrain." Denver, Colorado: U.S. Geological
18 Survey. 2004. <<http://tapestry.usgs.gov/Default.html>> (25 February 2008).
19
- 20 Vondra, C.F. "The Stratigraphy of the Chadron Formation in Northwestern Nebraska." MS
21 Thesis. University of Nebraska. Laramie, Wyoming. 1958.
22
- 23 Washington State Department of Transportation. "WSDOT's Guidance for Addressing Noise
24 Impacts in Biological Assessments—Noise Impacts." Seattle, Washington: Washington State
25 Department of Transportation. 2006. <[http://www.wsdot.wa.gov/TA/Operations/
26 Environmental/NoiseChapter011906.pdf](http://www.wsdot.wa.gov/TA/Operations/Environmental/NoiseChapter011906.pdf)> (12 October 2007).
27
- 28 WDEQ. "Chapter 2, Ambient Standards." 2006. <<http://deq.state.wy.us/aqd/standards.asp>>
29 (23 October 2007).
30
- 31 Whitehead, R.L. "Groundwater Atlas of the United States, Montana, North Dakota, South
32 Dakota, Wyoming." U.S. Geological Survey Report HA 730-I. Denver, Colorado:
33 U.S. Geological Survey. <http://capp.water.usgs.gov/gwa/ch_i/index.html> 1996.
34
- 35 Williamson J.E. and J.M. Carter J.M. Water-Quality Characteristics in the Black Hills Area,
36 South Dakota. U.S. Geological Survey Water Resources Investigation Report 01-4194. 2001.
37
- 38 Wyoming Department of Transportation. "WYDOT Traffic Analysis." Cheyenne, Wyoming:
39 Wyoming Department of Transportation. 2005. <[http://www.dot.state.wy.us/
40 Default.jsp?sCode=hwyta](http://www.dot.state.wy.us/Default.jsp?sCode=hwyta)> (25 February 2008).
41
- 42 Wyoming Game and Fish Department. "Comprehensive Wildlife Conservation Strategy."
43 Cheyenne, Wyoming: Wyoming Game and Fish. 2008.
44 <<http://gf.state.wy.us/wildlife/CompConvStrategy/Species>> (19 February, 2008)
45
- 46 Wyoming Game and Fish Department. "Terrestrial Habitat/Aquatic Habitat/Habitat
47 Management." Cheyenne, Wyoming: Wyoming Game and Fish. 2007.
48 <<http://gf.state.wy.us/habitat/aquatic/index.asp>> (13 September 2007).

3.5 Northwestern New Mexico Uranium Milling Region

3.5.1 Land Use

The Northwestern New Mexico Uranium Milling Region defined in this GEIS lies within the Navajo section of the Colorado Plateau (U.S. Geological Survey, 2004). This region includes McKinley County and the northern part of Cibola County (Figure 3.5-1). Past, current and potential uranium milling operations are found in two areas: (1) the central western part of McKinley County, east of Gallup, New Mexico and (2) the southeastern part of McKinley County and the northern part of Cibola County, east and northeast of Grants, New Mexico. These two areas are parts of the Grants Uranium District (Figure 3.5-2). Details on the geology and soils of this district and its subdivisions are provided in Section 3.5.3.

Land distribution statistics in Table 3.5-1 were calculated using the Geographic Information System used to construct the map shown in Figure 3.5-1. The data show that 91 percent of the Northwestern New Mexico Uranium Milling Region is composed of private land (50 percent), Indian Reservation land (27 percent) and U.S. National Forest land (14 percent).

Indian Reservation land, administered by the Bureau of Indian Affairs, comprises Acoma Pueblo, Laguna, Navajo, Ramah Navajo, and Zuni Indian land. Navajo land forms the northwest corner of McKinley County and abuts the northwestern part of the Grants Uranium District. Portions of any potential new ISL facility in this area of this district could fall within Navajo allottees, who own the surface and mineral rights. BIA administers the leases needed for both the surface use and mineral rights on such land. In this area of McKinley County, the Crownpoint and Church Rock Chapters of the Navajo Nation are part of an area known as the checkerboard due to its mixed private tribal and government property rights. Certain properties are under the Navajo Tribal Trust while individual Navajo allotments are privately held, with some BIA oversight (NRC, 1997).

Land use issues in the area of the Navajo Nation are a sensitive issue and consideration should be paid to ongoing jurisdictional disputes over the checkerboard lands. In addition, contamination of water supplies within the Rio San Jose Basin as a result of uranium milling has further heightened the Navajo Nation's sensitivity to land uses that may affect their ability to use tribal lands for raising livestock.

BLM lands occupy only approximately 8 percent of the region and are mostly concentrated in the northeastern corner of McKinley County (Figure 3.5-1). Other federal lands managed by the DoD (Fort Wingate Military Reservation) and the National Park Service represent less than 1 percent of the region.

Although sparsely populated, this region has three fairly large population centers: Gallup, with more than 20,000 people, Grants with approximately 9,000 people, and Zuni Pueblo with about 6,400 people. Smaller communities are scattered along the Interstate 40 corridor (Figure 3.5-2). Generally, private, federal and Indian Reservations land in this region are rural, mainly undeveloped, sparsely populated and are mostly used for livestock grazing, and to a lesser extent, for timber and agricultural production. In McKinley County, for example, more than 85 percent of the land is used for agricultural purposes and 83 percent of that land is used for livestock grazing. Only 9 percent and 0.6 percent of the land is used for timber production and for dry and irrigated crop production, respectively. Coal and uranium milling activities use less than 1 percent of the land in McKinley County (NRC, 1997).




Figure 3.5-1. Northwestern New Mexico Uranium Milling Region General Map With Current and Future Uranium Milling Site Locations




Figure 3.5-2. Map Showing Outline of the Northwestern New Mexico Region and the Location of the Grants Uranium District Along the Southern Margin of the San Juan Basin

Table 3.5-1. Land Ownership and General Use in the Northwestern New Mexico Uranium Milling Region

Land Ownership and General Use	Area (mi²)	Area (km²)	Percent
State and Private Lands	3,682	9,537	50.1
Bureau of Indian Affairs, Indian Reservations	1,999	5,176	27.2
U.S. Forest Service, National Forest	1,028	2,662	14
U.S. Bureau of Land Management (BLM), Public Domain Land	579	1,501	7.9
U.S. Department of Defense (Army)	29	75	0.4
National Park Service, National Monument	25	64	0.3
National Park Service, National Historic Park	6	16	0.08
BLM, National Conservation Area	1	2	0.01
BLM, Wilderness	0.5	1	0.01
Totals	7,350	19,035	100

Recreational and cultural activities for the public are available in the Mt. Taylor Ranger District, part of the Cibola National Forest. This forest includes the Zuni Mountains to the west of Grants and the San Mateo Mountains and Mount Taylor, about 24 km [15 mi] to the east-northeast of Grants. Mount Taylor is designated by the Navajo Nation as one of six sacred mountains. In Navajo tradition, Mount Taylor has a special significance as it represents the southern boundary of the Navajo traditional homeland (USFS, 2006), and in February 2008, the New Mexico Cultural Properties Review Committee approved listing the Mount Taylor Traditional Cultural Property in the State Register of Cultural Properties (see Section 3.5.8.3).

El Malpais National Monument in Cibola County and the Chaco Culture National Historical Park, which has several sites in McKinley County and San Juan County further north, are the two main recreational and cultural areas managed by the National Park Service in the Northwestern New Mexico Uranium Milling Region.

3.5.2 Transportation

Past experience at NRC licensed ISL facilities indicate these facilities rely on roads for transportation of most goods and personnel (Section 2.8). As shown on Figure 3.5-3, the New Mexico Uranium Milling Region is accessed from the east and west by Interstate 40, from the north by U.S. Highway 491 (formerly U.S. Highway 666) and State Routes 371 and 509 from the north, and State Route 36 and 602 from the south. A rail line traverses the region east and west along the path of Interstate 40.

Areas of past, present, or future interest in uranium milling in the region are shown in Figure 3.5-3. These areas are located in three sub-regions when considering site access by local roads. Areas of milling interest from west to east include areas near Pinedale northeast of Gallup, the area near Crownpoint north of Thoreau, and the area northeast of Milan and Grants near Ambrosia Lake and San Mateo. All these areas have access to Interstate 40 to the south using local access roads to State Routes 566 near Pinedale, 371 near Crownpoint, and 509 and 605 near Ambrosia Lake and San Mateo.

Table 3.5-2 provides available traffic count data for roads that support areas of past, present, or future milling interest in the Northwestern New Mexico Uranium Milling Region. Counts are




Figure 3.5-3. Northwestern New Mexico Uranium Milling Region Transportation Corridor Locations

1

Table 3.5-2. Average Annual Daily Traffic Counts for Roads in the Northwestern New Mexico Uranium Milling Region*

Road Segment	County	All Vehicles	
		2005	2006
State Route 566 North at State Route 118	McKinley	4,605	4,637
State Route 371 at Interstate 40 (Thoreau)	McKinley	5,514	5,552
State Route 371 North at Navajo 9 to Mariano Lake	McKinley	3,842	3,868
State Route 605 North at County Line North of Milan	McKinley	2,522	2,488
State Route 605 North at State Route 509 to Ambrosia Lake	McKinley	1,595	1,562
State Route 509 North at State Route 605	McKinley	338	330
Interstate 40, Thoreau Interchange North	McKinley	11,676	11,709
State Route 605 North at State Route 122 in Milan	Cibola	1,232	1,196
Interstate 40, Grants-Milan Interchange	Cibola	10,186	9,993
*NMDOT. "Road Segments by Traffic (AADT) Info." Data for Cibola and McKinley Counties from the New Mexico State Highway and Transportation Department's Consolidated Highway Data Base, provided by request. Santa Fe, New Mexico: New Mexico Department of Transportation. April 2008.			

2

3 variable with the minimum all vehicle count at 330 vehicles per day on State Route 509 North at
 4 State Route 605 and the maximum on Interstate 40, Thoreau Interchange North at 11,709
 5 vehicles per day. Most all vehicle counts in the Northwestern New Mexico Uranium Milling
 6 Region are above 1500 vehicles per day.

7

8 Yellowcake product shipments are expected to travel from the milling facility to a uranium
 9 hexafluoride production (conversion) facility in Metropolis, Illinois (the only facility currently
 10 licensed by NRC in the U.S. for this purpose). Major interstate transportation routes are
 11 expected to be used for these shipments, which are required to follow NRC packaging and
 12 transportation regulations in 10 CFR Part 71 and U.S. Department of Transportation hazardous
 13 material transportation regulations at 49 CFR Parts 171–189. Table 3.5-3 describes
 14 representative routes and distances for shipments of Yellowcake from locations of Uranium
 15 milling interest in the Northwestern New Mexico Uranium Milling region. Representative routes
 16 are considered owing to the number of routing options available that could be used by a future
 17 ISL facility.

18

19 3.5.3 Geology and Soils

20

21 New Mexico ranks second in uranium reserves in the United States. In the Northwestern New
 22 Mexico Uranium Milling Region, uranium resources are located primarily in the Grants uranium
 23 district (see Figure 3.5-2). The Grants uranium district includes a belt of sandstone-type
 24 uranium deposits stretching 135 km [85 mi] along the south side of the San Juan Basin. The
 25 Grants district consists of eight subdistricts, which extend from east of Laguna to west of Gallup
 26 (Figure 3.5-4) (McLemore and Chenoweth, 1989). The sandstone-type uranium deposits in the
 27 Grants district are generally in a geologic setting favorable for exploitation by ISL milling. More
 28 than 150,000 metric tons [170,000 tons] of U_3O_8 have been produced from these deposits from

Table 3.5-3. Representative Transportation Routes for Yellowcake Shipments From the Northwestern New Mexico Uranium Milling Region*

Origin	Destination	Major Links	Distance (mi)
North of Pinedale, New Mexico	Metropolis, Illinois	Local access road to State Route 566 State Route 566 south to Interstate 40 Interstate 40 east to Memphis, Tennessee Interstate 55 north to Interstate 155 Interstate 155 north to Interstate 24 Interstate 24 north to Metropolis, Illinois	1,360
Crownpoint, New Mexico	Metropolis, Illinois	Local access road to State Route 371 State Route 371 south to Interstate 40 Interstate 40 east to Metropolis, Illinois (as above)	1,360
North of San Mateo, New Mexico	Metropolis, Illinois	Local access road to State Route 334 at San Mateo State Route 334 west to State Route 605 State Route 605 to Interstate 40 at Milan near Grants	1,300

*American Map Corporation. "Road Atlas of the United States, Canada, and Mexico." Long Island City, New York: American Map Corporation. p. 144. 2006.

1948 to 2002, accounting for 97 percent of the total production in New Mexico and more than 30 percent of the total production in the United States (McLemore and Chenoweth, 1989).

The San Juan Basin is a structural depression occupying a major portion of the southeastern Colorado Plateau physiographic province (Hunt, 1974). The plateau encompasses much of western Colorado, eastern Utah, northeastern Arizona, and northwestern New Mexico. The San Juan Basin is underlain by up to 3,000 m [10,000 ft] of sedimentary strata, which generally dip gently from the margins toward the center of the basin. The margins of the basin are characterized by relatively small elongate domes, uplifts, and synclinal depressions.

Uranium mineralization in Grants district occurs within Upper Jurassic (144 to 159 million year old) and Cretaceous (65 to 144 million year old) sandstones. Stratigraphic descriptions presented here are limited to formations that would be involved in potential milling operations or formations that may have environmental significance, such as important aquifers and confining units above and below potential milling zones. A generalized stratigraphic column of formations in the Grants uranium district is shown in Figure 3.5-5.

The Morrison Formation is composed of the Recapture, Westwater Canyon, and Brushy Basin Members and is the host formation for major uranium deposits in the Grants uranium district. Most of the deposits are within the main sandstone bodies of the Westwater Canyon Member. In addition, the Westwater Canyon is an important regional aquifer. Large uranium deposits are also found in a series of sandstone beds, known collectively as the Poison Canyon sandstones of economic usage, which occur near the base of the Brushy Basin Member in the Blackjack (Smith Lake), Poison Canyon, and Ambrosia Lake mining areas (Holen and Hatchell, 1986). Deposits also occur in sandstone lenses higher in the Brushy Basin in the Blackjack (Smith Lake) mining area. In the Laguna district a bed of sandstone overlying the Brushy Basin, the Jackpile Sandstone Member of the Morrison (Owen, 1984), contains the large Jackpile-Paguete, L-Bar and Saint Anthony deposits. Relationships of the deposits in the various Morrison units are shown in Figure 3.5-6.




Figure 3.5-4. Index Map of the Grants Uranium District, San Juan Basin, New Mexico, Showing Eight Subdistricts (Modified From McLemore, 2007)

Elsewhere in the San Juan Basin, significant but relatively small sandstone-type deposits also occur in the Dakota Sandstone in the Church Rock area and in the Burro Canyon Formation in the Carjilon area (Holen and Hatchell, 1986). The Todilto Limestone in the Grants district, which has accounted for about two percent of total production, is quite impermeable and is unlikely to be amenable to production by ISL. Beyond the San Juan Basin, significant but relatively small sandstone-type deposits occur in the Galisteo Formation in the Hagan Basin, and in the Crevasse Canyon and Baca Formations in the Riley-Pie Town areas.

The following regional descriptions of the stratigraphic units within the San Juan Basin are derived from reports by Green and Pierson (1977), Hilpert (1963, 1969), Chenoweth and Learned (1980), and Holen and Hatchell (1986).

The Recapture Member is the bottommost member of the Morrison Formation. It is as thick as 150 m [500 ft] northwest of Gallup but thins to 45 to 90 m [150 to 300 ft] in outcrops near Gallup and eastward. The Recapture is one of the most variable stratigraphic units in the area. It

1




Figure 3.5-5. Generalized Stratigraphic Section of Upper Jurassic and Cretaceous Formations in the Grants Uranium District (NRC, 1997)

2

3




Figure 3.5-6. Generalized Geologic Section Showing the Stratigraphic Relations of the Morrison Formation Between the Ambrosia Lake and Laguna Areas (From Hilpert, 1969)

occurs in the Gallup mining district as a sequence of interbedded siltstone, mudstone, and sandstone strata. Individual strata range from centimeters to meters in thickness. Sandstone beds are generally less than 5 m [15 ft] thick (Hilpert, 1969). The Recapture is believed to interfinger with the underlying Cow Springs Sandstone, and several authors have combined the two units as one. No significant uranium deposits occur in the Recapture Member.

The Westwater Canyon Member of the Morrison Formation consists of interbedded fluvial red, tan, and light gray arkosic sandstone (i.e., sandstone containing a significant fraction of feldspar), claystone, and mudstone. It is a major water-bearing member of the Morrison. The unit ranges from 53 to 85 m [175 to 275 ft] thick in outcrop from Gallup to the continental divide (Hilpert, 1969) and is known to be considerably thicker locally. In most places, the Westwater Canyon displays one or more mudstone units that range from thin partings to units up to 6 m [20 ft] thick. The mudstone units have limited lateral continuity, and only the thicker ones are extensive. The Westwater Canyon is host for the major uranium deposits in the region. The uranium occurs in coarse-grained, poorly sorted sandstone units and is closely associated with the carbonaceous material that coats the sand grains.

Three types of stratabound uranium deposits are present in the Westwater Canyon Member: primary (trend or tabular), roll-front (redistributed), and remnant-primary sandstone uranium deposits (Figure 3.5-7) (McLemore, 2007). Primary sandstone-hosted uranium deposits, also known as prefault, trend, blanket, and black-band ores, are found as blanket-like, roughly parallel ore bodies along sandstone trends. These deposits are characteristically less than 2.5 m [8 ft] thick, average more than 0.20 percent U_3O_8 , and have sharp ore-to-waste boundaries. The largest deposits in the Grants uranium district contain more than 13,600 metric tons [15,000 tons] of U_3O_8 .

During the Tertiary (1.8 to 65 million years ago), oxidizing groundwaters migrated through the Morrison Formation and remobilized some of the primary sandstone uranium deposits (Saucier, 1981). Uranium was reprecipitated ahead of the oxidizing waters forming roll-front sandstone uranium deposits (see Section 3.1.1). Roll-front uranium deposits are also known as post-fault, stack, secondary, and redistributed ores. A schematic diagram of the formation of a redistributed or roll-front uranium deposit is shown in Figure 3.1-5. They are discordant, asymmetrical, irregularly shaped, characteristically more than 2.5 m [8 ft] thick, have diffuse ore-to-waste contacts, and cut across sedimentary structures. The average deposit contains approximately 8,500 metric tons [9,400 tons] U_3O_8 with an average grade of 0.16 percent. Some redistributed uranium deposits are vertically stacked along faults (see Figure 3.5-7).

Remnant sandstone-hosted uranium deposits were preserved in sandstone after oxidizing waters that formed roll-front uranium deposits had passed. Some remnant sandstone-hosted uranium deposits were preserved because they were surrounded by or found in less permeable sandstone and could not be reached by oxidizing groundwaters. These deposits are similar to primary sandstone-hosted uranium deposits, but are difficult to locate because they occur sporadically within the oxidized sandstone. The average size is approximately 1,200 metric tons [1,400 tons] U_3O_8 at a grade of 0.20 percent.

There is no consensus on the origin of the Morrison Formation sandstone uranium deposits and the source of uranium is not well constrained (Sanford, 1992). Uranium could be derived from alteration of volcanic detritus and shales within the Morrison Formation (Thamm et al., 1981; Adams and Saucier, 1981) or from groundwater derived from a volcanic highland to the southwest. The majority of the proposed models for their formation suggest that deposition occurred at a groundwater interface between two fluids of different chemical compositions




Figure 3.5-7. Schematic Diagram of the Different Types of Uranium Deposits in the Morrison Formation, Grants Uranium District, New Mexico (Modified from Holen and Hatchell, 1986). See Text for Description.

and/or oxidation/reduction states. Bleaching of the Morrison sandstones and the geometry of tabular uranium bodies floating in sandstone beds supports the reaction of two chemically different waters, most likely a dilute meteoric water and saline brine from deeper in the basin (McLemore, 2007).

The Brushy Basin Member overlies the Westwater Canyon and ranges from 12 to 40 m [40 to 125 ft] thick in the Gallup region. It is mainly composed of light greenish gray and varicolored claystone, interbedded with sandstone lenses having similar lithology and appearance to sandstones found in the Westwater Canyon Member (Ristorcelli, 1980). The mudstones are largely derived from volcanic ash falls (Peterson, 1980) and contain considerable amounts of bentonite. The contact between the Brushy Basin and the Westwater Canyon is gradational and interfingering.

The Dakota Sandstone is the basal formation of the Cretaceous System and unconformably overlies the Morrison Formation. The Dakota is a gray-brown quartz sandstone with some interbedded conglomerate, shale, carbonaceous shale, and coal. The Dakota Sandstone is marine in origin and is considered to represent the earliest transgression of late Cretaceous seas. The Dakota crops out around the margins of the San Juan Basin and thickens towards

the center of the basin to about 60 m [200 ft]. The Mancos Shale overlies the Dakota Sandstone and is a thick, mostly uniform gray marine shale containing thin lenses of fine-grained sandstone.

Approximately 227 metric tons [250 tons] of U_3O_8 have been produced from roll-front uranium deposits in the Dakota Sandstone in the southern part of the San Juan Basin (Chenoweth, 1989). Uranium deposits in the Dakota Sandstone are typically tabular masses that range in size from thin pods a few meters (feet) long and wide to masses as much as 760 m [2,500 ft] long and 300 m [1,000 ft] wide. The larger deposits are only a few meters (feet) thick, but a few are as much as 8 m [25 ft] thick (Hilpert, 1969). Ore grades range from 0.12 to 0.30 percent and average 0.21 percent U_3O_8 . Uranium is found with carbonaceous plant material near or at the base of channel sandstones or in carbonaceous shale and lignite and is associated with fractures, joints, or faults and with underlying permeable sandstone of the Brushy Basin or Westwater Canyon Members. The largest deposits in the Dakota Sandstone are found in the Old Church Rock mine in the Church Rock subdistrict, where uranium is associated with a major northeast-trending fault. More than 81 metric tons [90 tons] of U_3O_8 have been produced from the Dakota Sandstone in the Old Church Rock mine (Chenoweth, 1989).

The San Juan Basin is part of the Colorado Plateau physiographic province, which is generally characterized by rough, broken terrain, including small steep mountainous areas, plateaus, cuestas, and mesas intermingled with steep canyon walls, escarpments, and valleys. Thick colluvium deposits are commonly found forming a mantle on steep slopes surrounding sandstone mesas and cuestas in the San Juan Basin. In contrast, Quaternary alluvium is found on the valley floors of the region. These deposits consist of fine sand, silt, and clay derived from the weathering of sandstone, siltstone, and mudstone exposed at the surface. Alluvial deposits generally are thin but are known to exceed a thickness of 10 m [30 ft] in larger valleys.

General soils information associated with landforms in the southern part of the San Juan Basin was obtained from the Soil Survey of McKinley County Area, New Mexico, McKinley County and Parts of Cibola and San Juan Counties (NRCS, 2001). For site-specific evaluations at proposed ISL milling facilities, more detailed soils information would be expected to be obtained from published county soil surveys or the U.S. Department of Agriculture NRCS.

In the southern part of the San Juan Basin, soils on hills and mountains vary greatly in horizon development, from soils with no development to soils that have well-developed clay horizons. Gravelly clay loams having little or no horizon development are usually found on steeper slopes where erosional activity is greatest. Clay loam soils that have well-developed horizons are generally found on gently sloping to moderately steep slopes, where erosion is slight to moderate. Gravelly to fine sand loam soils characterized by well-developed clay horizons are found on mesa summits and cuesta dip slopes, which are nearly level to gently sloping. Sandy to fine sandy loam soils with little or no horizon development are found on the escarpment of mesas and cuestas and on hogbacks, where erosional activity is great. Fine sandy loam soils are found on the summits of ridges and are mostly shallow, whereas sandy loam soils are found on the side slopes of ridges and are generally shallow but sometimes deeper. Soils on alluvial fans are generally very deep, and their soil textures are highly variable, depending on the local geology. Soils found on alluvial fans include clay loam and fine sandy loam. Soils on stream terraces are underlain by stratified sand, gravel, loamy, silty, or clayey sediments and, in some cases, buried paleosols. Typical soils that represent stream terraces are sandy clay loam and silt loam. Soils on floodplains and drainageways are generally very deep, with soil textures that are highly variable, depending on the local geology. Clay loam and fine sand loam soils are found in drainageways and fine sand and clay loam soils are found on floodplains.

3.5.4 Water Resources

3.5.4.1 Surface Waters

The Northwestern New Mexico Uranium Milling Region includes McKinley and the northern portion of Cibola County and a small portion western Bernalillo County. Watersheds in the Northwestern New Mexico Uranium Milling Region are Rio San Jose, Zuni, Chaco Canyon, Upper Puerco River,¹ Arroyo Chico, and a small portion of Rio Puerco (EPA, 2008) (Figure 3.5-8). The named uranium deposits shown in Figure 3.5-4 are listed with their corresponding watershed in Table 3.5-4. The unnamed uranium deposits northeast of Chaco Canyon are located in the Arroyo Chico and Rio Puerco watersheds. Historical and potential uranium milling sites are located in the Upper Puerco, Chaco, Arroyo Chico, and Rio San Jose watersheds. The Zuni River watershed does not contain any identified uranium deposits that are being considered for ISL uranium recovery. The Rio San Jose is the watershed only water watershed with perennial stream reaches within the area of potential uranium milling.

The Rio San Jose and associated tributaries drain the south-central portion of McKinley County and northeastern portion of Cibola County. The Rio San Jose flows into Rio Puerco east of the Northwestern New Mexico Uranium Milling Region. The state designated uses of Rio San Jose and its tributaries are listed in Table 3.5-5 along with known impairments to these uses. Impairments to water quality within the Rio San Jose watershed include elevated nutrients, metals (aluminum), turbidity, temperature and sediment. Flow of the Rio San Jose is not gauged within the region.

The Rio Puerco drains a small portion of the east-central part of the Northwestern New Mexico Uranium Milling Region (Figure 3.5-8). The Rio Puerco flows southeast to the Rio Grande southeast of the Northwestern New Mexico Uranium Milling Region. The mainstem of the Rio Puerco is east of the Northwestern New Mexico Uranium Milling Region and none of the tributaries of Rio Puerco are perennial within the Northwestern New Mexico Uranium Milling Region.

The other watersheds within the area of potential uranium recovery with Northwestern New Mexico Uranium Milling Region contain ephemeral streams that flow only after precipitation events. The only surface water features in these watershed are springs and stock ponds. Many springs are present within the Northwestern New Mexico Uranium Milling Region in McKinley and Cibola counties. These springs occur on the flanks of mountainous areas, such as the Chuska Mountains in the western portion of the region and the Mt. Taylor area in the southeastern portion of the region as well as in the intermontane areas. These springs are fed by both local and regional aquifer systems (see Section 3.5.4.3).

3.5.4.2 Wetlands and Waters of the United States

Wetlands and other shallow aquatic habitats occupy only about 1–5 percent of the land surface in this region (USACE, 2006).

Within this region no digital data are available. However, hardcopy National Wetland Inventory Maps can be obtained from the U.S. Fish and Wildlife Service. In general Waters of the U.S. in

¹ The Rio Puerco watershed is located in north-central New Mexico and drains into the Rio Grande. The Puerco River watershed is located in west-central New Mexico and drains into the Little Colorado River in Arizona.




Figure 3.5-8. Watersheds in the Northwestern New Mexico Uranium Milling Region

1

Table 3.5-4. Named Uranium Deposits in New Mexico and Corresponding Watersheds	
Uranium Deposit	Watershed
Barnabe Montano	Rio San Jose
Marquez	Rio San Jose
Laguna	Rio San Jose
Grants	Rio San Jose
Smith Lake	Rio San Jose
Nose Rock	Chaco Canyon
Chaco Canyon	Chaco Canyon
Church Rock	Puerco River
Crownpoint	Chaco Canyon

2
3

Table 3.5-5. Primary Watersheds in New Mexico, Designated Uses and Known Impairments			
Watershed	Tributary or Reach	State Designated Uses	Known Impairments
Rio San Jose	Bluewater Creek	Wildlife Habitat Irrigation Fish Culture Domestic Water Supply Cold Water Fishery Primary Contact Livestock Watering	Nutrients Aluminum Turbidity Temperature Sedimentation
	Bluewater Lake	Wildlife Habitat Irrigation Fish Culture Domestic Water Supply Cold Water Fishery Primary Contact Livestock Watering	None
	Rio Moquino	Wildlife Habitat Irrigation Fish Culture Domestic Water Supply Cold Water Fishery Primary Contact Livestock Watering	Temperature Sedimentation

1

Table 3.5-5. Primary Watersheds in New Mexico, Designated Uses and Known Impairments (continued)

Watershed	Tributary or Reach	State Designated Uses	Known Impairments
	Rio Paquate	Wildlife Habitat Irrigation Fish Culture Domestic Water Supply Cold Water Fishery Primary Contact Livestock Watering	Selenium Temperature Sedimentation
	Rio San Jose	Wildlife Habitat Livestock Watering	None
	Seboyeta Creek	Wildlife Habitat Irrigation Fish Culture Domestic Water Supply Cold Water Fishery Primary Contact Livestock Watering	None
Rio Puerco	No Perennial Reaches in New Mexico Region		
Upper Puerco River	No Perennial Reaches in New Mexico Region		
Arroyo Chico	No Perennial Reaches in New Mexico Region		
Chaco	No Perennial Reaches in New Mexico Region		
Zuni River	No Known Uranium Recovery Activities in Zuni Watershed		

2
3 this region consist of ephemeral stream/arroyos with few perennial rivers. Bands of wetlands
4 are concentrated along rivers and streams within this region. Seasonally emergent wetland
5 areas may be found within woody habitat at high elevations. Within this region springs and
6 seeps often support small marshes (ciénegas), oases, and other wetland types (USACE, 2006).
7 Desert playas are intermittent shallow lakes that develop in the flat, lower portions of arid basins
8 during the wet season. Most are unvegetated and may not contain water every year.

9
10 Waters of the United States and special aquatic sites that include wetlands would be expected
11 to be identified and the impact delineated upon individual site selection. Based on impacts and
12 consultation with each area, appropriate permit would be expected to be obtained from the local
13 USACE district. Within this region the state does not regulate wetlands; however, Section 401
14 state water quality certification is required for work in Waters of the United States.

15 3.5.4.3 Groundwater

16
17
18 Groundwater resources in the Northwestern New Mexico Uranium Milling Region are part of
19 regional aquifer systems that extend well beyond the areas of uranium milling interest in this
20 part of New Mexico. Uranium bearing aquifers exist within these regional aquifer systems in the
21 Northwestern New Mexico Uranium Milling Region. This section provides a general overview of
22 the regional aquifer systems to provide context for a more focused discussion of the

uranium-bearing aquifers in northwestern New Mexico, including hydrologic characteristics, level of confinement, groundwater quality, water uses, and important surrounding aquifers.

3.5.4.3.1 Regional Aquifer Systems

The Colorado Plateau aquifers underlie northwestern New Mexico and most parts of the Northwestern New Mexico Uranium Milling Region (Robson and Banta, 1995). The principal aquifers are present only in the San Juan Basin in northwest New Mexico. The geographical region in New Mexico underlain by the Colorado Plateaus aquifers is sparsely populated and the quality and quantity of the groundwater pumped from these aquifers are suitable for most agricultural or domestic uses. The aquifers are typically composed of permeable sedimentary rocks of Permian to Tertiary ages.

Robson and Banta (1995) grouped the Colorado Plateau aquifers into four principal aquifers, which are, from shallowest to deepest, the Uinta-Animas aquifer, the Mesaverde aquifer, the Dakota-Glen Canyon aquifer system, and the Coconino-De Chelly aquifer. These four principal aquifers are hydraulically separated by relatively impermeable confining layers. The Mancos shale confining unit that underlies the Mesaverde aquifer and the Chinle-Moenkopi confining unit that underlies the Dakota-Glen Canyon aquifer system are the thickest confining layers. Among these four aquifer systems, the Mesaverde aquifer system (for water supplies) and the Dakota-Glen Canyon aquifer system (for water supplies and uranium milling) are the most important aquifer systems in the Northwestern New Mexico Uranium Milling Region.

The Mesaverde Aquifer: The Mesaverde aquifer is a regionally important aquifer for water supplies. It consists of sandstone, coal, siltstone, and shale of the Mesaverde Group in the San Juan Basin. The formations of the Mesaverde Group extensively interbedded with the Mancos Shale and, to a lesser extent, with the Lewis Shale. The thickness of the Mancos Shale typically ranges from 305 to 1,830 m [1,000 to 6,000 ft], and in general it forms a thick barrier to vertical and lateral groundwater flow. The maximum thickness of the Mesaverde aquifer is about 1,370 m [4,500 ft] in the southern part of San Juan Basin. The recharge to aquifer is by precipitation and discharge from aquifer is to streams, springs, and seeps, by upward movement across confining layers and into overlying aquifers, and by withdrawals. In general water pumpage from the Mesaverde aquifer is small; therefore, water-level declines are usually localized. The altitude of the potentiometric surface ranges from 1,525 to 2,440 m [5,000 to 8,000 ft] in the San Juan Basin. In most parts of the basin, transmissivity of the Mesaverde aquifer is typically less than 4.65 m²/day [50 ft²/day]. However, where the aquifer is fractured, the local transmissivities could be 100 times higher.

The water quality in the Mesaverde aquifer is variable. The dissolved solids concentration ranges from about 1,000 to 4,000 mg/L [1,000 to 4,000 ppm] in parts of the San Juan Basins, which exceed EPA's Secondary Drinking Water Standard of 500 mg/L [500 ppm].

Dakota-Glen Canyon Aquifer System: Large depths to the water table or poor water quality make the aquifers of the Dakota-Glen Canyon aquifer system unsuitable for production in most parts of the New Mexico Uranium Milling Region. Where an aquifer is close to the land surface, however, it can be important source of water. The Dakota-Glen Canyon aquifer system is confined by Mancos confining unit above and by Chinle-Moenkopi confining unit below. The thickness of the Chinle-Moenkopi confining unit is typically 305 to 610 m [1,000 to 2,000 ft]. These confining units substantially limit the Dakota-Glen Canyon aquifer system's hydraulic connection with the overlying and underlying aquifers.

The Dakota-Glen Canyon aquifer system consists of four major aquifers: the Dakota aquifer (including the Dakota Sandstone and adjacent water-yielding rocks), the Morrison aquifer (including water-yielding rocks generally of the lower part of the Morrison Formation), the Entrada aquifer (including the Entrada Sandstone and the Preuss Sandstone), and the Glen Canyon aquifer (including the Glen Canyon Sandstone or Group and the Nugget Sandstone). The aquifer systems typically include confining units that separate these aquifers. At the regional scale, recharge areas, discharge areas, groundwater flow directions, and water quality are similar among these four aquifers.

The top of the Dakota aquifer is less than 610 m [2,000 ft] below the surface in the San Juan Basins. The transmissivity of the Dakota aquifer is poorly defined in the region. The Dakota aquifer is underlain by the Morrison Formation. In most parts of the basin, the relatively impermeable Morrison confining unit is present in the upper parts of the Morrison Formation. The middle and lower parts of the Morrison Formation forms the Morrison aquifer, but only the coarser-grained strata generally yields water. In the San Juan Basin, the Morrison aquifer includes two underlying water-yielding sandstone units, the Cow Springs and Junction Creek Sandstones. In most places, the Morrison aquifer is underlain by the relatively impermeable Curtis-Stump confining unit.

The Entrada aquifer underlies either the Curtis-Stump confining unit or the Morrison aquifer. The Entrada aquifer consists mainly of the Entrada Sandstone. In the western part of the Uinta Basin, the aquifer is composed of the Preuss Sandstone, which is an equivalent of the Entrada aquifer. In part of the basins, the Entrada aquifer directly overlies the Glen Canyon aquifer that consists of Wingate Sandstone, Kayente Formation, and the Navajo Sandstone. The Glen Canyon is the thickest and where fractured has relatively high transmissivities. The transmissivity of the Glen Canyon aquifer typically ranges from about 9.23- 92.9 m²/day [100 to 1,000 ft²/day]. Groundwater flow in the Glen Canyon aquifer is toward major discharge areas along the San Juan Rivers. The depth to the top of the Glen Canyon aquifer is typically less than 610 m [2,000 ft]. The dissolved-solids concentration in the Glen Canyon aquifer is less than 1,000 mg/L [1,000 ppm].

3.5.4.3.2 Aquifer Systems In The Vicinity Of Uranium Milling Sites

The underlying hydrogeological system in past and current areas of uranium milling interest in the Northwestern New Mexico Uranium Milling Region consists of a thick sequence of primarily sandstone aquifers and shale aquitards.

Areas of uranium milling interest at the Crownpoint, Unit 1, and Church Rock areas are underlain, from shallowest to deepest, by water-bearing layers in the Mesaverde Formation, the Dakota sandstone, the Morrison Formation (including the uranium-bearing Westwater Canyon aquifer), the Cow Springs Sandstone, and Entrada Sandstone. The Mesaverde Formation is regionally important for water supplies. The uranium-bearing Westwater Canyon aquifer at the active Uranium milling sites is also important for water supplies in the milling region. Little information is available for the Cow Springs sandstone aquifer, but the existing data suggests that Cow Springs aquifer underlying the Wastewater Canyon aquifer contain good quality water (HRI, 1996). Although the Dakota sandstone at the town of Crownpoint is qualified as a drinking water supply according to EPA's National Primary Drinking Water Regulations, it is locally (e.g., in McKinley County) unused as a water supply because of its poor water quality (NRC, 2007).

Description of the Affected Environment

3.5.4.3.3 Uranium-Bearing Aquifers

The most important uranium deposits in the northwestern New Mexico Region are hosted by the Westwater Canyon sandstone aquifer in the Morrison Formation (NRC, 1997; McLemore, 2007). The uranium-bearing sandstone aquifers in the Westwater Canyon aquifer and the Dakota sandstone near the town of Crownpoint must be exempted (Section 1.7.2) by EPA's UIC program (40 CFR § 144.3) before ISL operations begin.

Hydrogeological characteristics: The groundwater flow velocities in the Westwater Canyon aquifer at the Crownpoint site ranged from 3.9 m/yr [12.9 ft/yr] in the east to 2.4 m/yr [8 ft/yr] in the west side of the site. Transmissivity estimates for the Westwater Canyon aquifer range from 235 to 250 m²/day [2,550 to 2,700 gal/day/ft]. The storage coefficient values ranged from 4.50×10^{-5} to 1.39×10^{-4} (NRC, 1997).

At Unit 1, the aquifers are the same as those at the Crownpoint site. The calculated average groundwater velocity is 1.5 m/yr [5 ft/yr] in the Westwater Canyon aquifer. In the Westwater Canyon aquifer, transmissivity ranges from 84 to 133 m²/day [905 to 1,432 gal/day/ft] and the storage coefficient values range from 9.40×10^{-5} to 1.60×10^{-4} (NRC, 1997).

The aquifers located beneath the Church Rock site are similar to those beneath the Crownpoint and Unit 1 sites. The average groundwater flow velocity in the Westwater Canyon at Church Rock is 2.7 m/yr [8.7 ft/yr]. Transmissivity of the Westwater Canyon aquifer ranges from 86 to 123 m²/day [926 to 1,326 gal/day/ft] and the storage coefficient ranges from 8.90×10^{-5} to 4.13×10^{-4} (NRC, 1997).

The average storage coefficient of the Westwater Canyon aquifer is on the order of 10^{-5} – 10^{-4} at the Crownpoint, Unit 1, and Church Rock sites, indicating the confined nature of the production aquifer [typical storage coefficients for confined aquifers range from 10^{-5} – 10^{-3} (Driscoll, 1986)].

Level of confinement: At the Crownpoint site, the Westwater Canyon aquifer is confined below by the Recapture Shale and confined above by the Brushy Basin Shale. The upper aquitard is about 80 m [260 ft] thick and is continuous at the site. The lower confinement unit consists entirely of shale and is continuous at the site. Aquifer tests revealed no significant vertical flow across the Recapture Shale and Brushy Basin Shale aquitards. At Unit 1, both the upper (Brushy Basin Shale) and lower (Recapture Shale) aquitards that confine the Westwater Canyon aquifer are continuous beneath Unit 1. No significant vertical flow across the aquitards was detected. At the Church Rock site, the upper aquitard above the Westwater Canyon aquifer (Brushy Basin Shale) is 4–9 m [13–28 ft] thick. The thickness of the lower aquitard (Recapture Shale) was reported to be 55 m [180 ft] thick (NRC, 1997).

Groundwater quality: At the Crownpoint site, the artesian uranium-ore bearing Westwater Canyon sandstone aquifer is a valuable resource for high-quality groundwater, which fits the definition of underground sources of drinking water in the EPA National Primary Drinking Water Regulations (NRC, 1997). The TDS concentrations in groundwater range from 281 to 3,180 mg/L [281 to 3,180 ppm] and averages 773 mg/L [773 ppm]. The TDS levels in four town water wells ranged from 325 to 406 mg/L [325 to 406 ppm], which are lower than the EPA's Secondary Drinking Water Standard of 500 mg/L [500 mg/L]. Even though the town's water supply wells are completed in sandstones that contain uranium deposits, radionuclide

concentrations in the Crownpoint public water supply are low. The uranium and radium-226 concentrations at the Crownpoint ISL site's monitoring wells were in the range of less than 0.001 to 0.007 mg/L [0.001 to 0.007 ppm] and 0.3 to 0.6 pCi/L, respectively (EPA's drinking water standard for uranium is 0.03 mg/L (0.03 ppm) and for radium-226 is 5.0 pCi/L) (NRC, 1997).

At the Unit 1 site, groundwater in the Westwater Canyon aquifer in general meets New Mexico drinking water quality standards, except for radium-226 and uranium concentrations. The average radium-226 concentration at the Unit 1 ISL site's monitoring wells is 10.3 pCi/L, which exceeds the EPA drinking water standard for radium-226 (5.0 pCi/L). The average uranium concentration at the Unit 1 site is about 2.0 mg/L [2 ppm], which is higher than at the Crownpoint site. The average TDS of 285.0 mg/L [285 ppm] was lower than the EPA drinking water standard of 500 mg/L [500 ppm] (NRC, 1997).

At the Church Rock site, the groundwater quality is generally good in Westwater Canyon aquifer and meets the New Mexico drinking water quality standards, except for radium-226 concentration. However, the average radium-226 concentration at the monitoring wells was 10.2 pCi/L, exceeding the EPA drinking water standard of 5.0 pCi/L for radium. The average uranium concentration was 0.01 mg/L [0.01 ppm]. The average TDS of 369.75 mg/L [369.75 ppm] was lower than the EPA drinking water standard of 500 mg/L [500 ppm] (NRC, 1997).

Current groundwater uses: Groundwater in the northwestern New Mexico Region area is suitable for drinking. Groundwater has been used for domestic supplies, especially in the Crownpoint and Unit 1 areas. Most of the wells in and near the Church Rock site either owned by Hydro Resources, Inc. or are private wells (NRC, 1997).

3.5.4.3.4 Other Important Surrounding Aquifers for Water Supply

The Dakota Sandstone at the town of Crownpoint is qualified as a drinking water supply according to EPA's National Primary Drinking Water Regulations. Little information is available for the Cow Springs aquifer, but the existing data suggests that Cow Springs aquifer underlying the Westwater Canyon aquifer contains good quality water (HRI, 1996).

3.5.5 Ecology

3.5.5.1 Northwestern New Mexico Flora

According to EPA, the Northwestern New Mexico Uranium Milling Region contains two ecoregions, the Arizona/New Mexico Plateau and the Arizona/New Mexico Mountains (Figure 3.5-9). These regions and subregions are as follows. The Grants Uranium District in the region is located in the Semi Arid Tablelands, Conifer Woodlands, and Savannas ecoregions and near the San Juan/Chaco Tablelands and Mesas ecoregions.

The Arizona/New Mexico Plateau is a transitional region between shrublands and wooded higher relief tablelands of the Colorado Plateaus in the north, the lower less vegetated Mojave Basin and Range in the west, and forested mountain ecoregions that border the region on the




Figure 3.5-9. Ecoregions for the Northwestern New Mexico Uranium Milling Region

1 northeast and south. The topography in the region changes from a few meters [feet] on plains
2 and mesa tops to well over 305 m [1,000 ft] along tableland side slopes. This region extends
3 across northern Arizona, northwestern New Mexico, and into Colorado in the San Luis Valley
4 (Griffith, et al., 2006).

6 The San Juan/Chaco Tablelands and Mesas ecoregion of plateaus, valleys, and canyons
7 contains a mix of desert scrub, semi-desert shrub-steppe, and semi-desert grasslands. Native
8 vegetation found within the region include shadscale, fourwing saltbush, mat saltbush,
9 greasewood, mormon tea, Indian ricegrass, alkali sacaton, galleta (*Pleuraphis jamesii*), and blue
10 and black grammas are typical. Rocky Mountain (*Juniperus scopulorum*), one-seed (*Juniperus*
11 *monosperma*), and Utah Junipers (*Juniperus osteosperma*) can be found on higher mesas
12 (Griffith, et al., 2006).

14 The Semiarid Tablelands consists of mesas, plateaus, valleys, and canyons. This region
15 contains areas of high and low relief plains. Grass, shrubs, and woodland cover the tablelands.
16 The vegetation is not as sparse as that found in the San Juan/Chaco Table lands to the north or
17 the Albuquerque basin to the east. Scattered junipers occur on shallow, stony soils, and are
18 dense in some areas. Pinyon-juniper woodland is also common in some areas. Fourwing
19 saltbush, alkali sacaton, sand dropseed, and mixed gramma grasses are common species
20 found in this region (Griffith, et al., 2006).

22 The Lava Malpais can be found in the south central portion of the region. The lava substrate
23 has the ability in places to trap and retain moisture, allowing for a more mesophytic
24 vegetation, such as stunted Douglas fir and ponderosa pine, to occur in some areas. Other
25 species which are found in this region include grasses like blue grama and side oats with
26 shrubs of Apache Plume (*Fallugia paradoxa*) and New Mexico Olive (*Forestiera pubescens*)
27 (Griffith, et al., 2006).

29 The Near-Rockies Valleys and Mesas ecoregion is a region comprised of mostly pinyon-juniper
30 woodland, juniper savanna, and mesa and valley topography, with influences of higher elevation
31 vegetation in drainages from the adjacent Southern Rockies. Other natural species that can be
32 found in this region include one seed and Rock mountain junipers, indian ricegrass, big
33 sagebrush, sand dropseed, gallets, threeawns, blue gramma, and rabbitbrush (Griffith, et al.,
34 2006).

36 The Arizona/New Mexico Mountains region is distinguished from neighboring mountainous
37 ecoregions by lower elevations and associated vegetation indicative of drier, warmer
38 environments. Forests of spruce, fir, and Douglas fir, which are common in mountainous
39 regions are limited to the highest elevations in this region. Chaparral is common at lower
40 elevations in some areas, pinyon-juniper and oak woodlands are found at lower and middle
41 elevations. Higher elevations in the region are mostly covered with open to dense ponderosa
42 pine forests. These mountains are the northern extent of some Mexican plant and animal
43 species. Surrounded by deserts or grasslands, these mountains in New Mexico can be
44 considered biogeographical islands (Griffith, et al., 2006).

46 The Montane Conifer Forests are found west of the Rio Grande at elevations from about 2,130
47 to 2,900 m [7,000 to 9,500 ft]. Ponderosa pine and Gambel oak (*Quercus gambelii*) are
48 common, along with mountain mahogany and serviceberry (*Amelanchier alnifolia*). Some
49 Douglas fir, southwestern white pine (*Pinus strobiformis*), and white fir occur in a few areas
50 (Griffith, 2006). This region also includes mixed conifer/aspen stands. Seven different conifers
51 can be found growing in the same region, and there are a number of common cold-deciduous

Description of the Affected Environment

shrub and grass species, including a few maple (*Acer spp.*), blueberry (*Vaccinium*) species, gray alder (*Alnus incana*), kinnikinnick (*Arctostaphylos uva-ursi*), water birch (*Betula occidentalis*), redosier dogwood (*Cornussericea*), Arizona fescue (*Festuca arizonica*), fivepetal cliffbush (*Jamesia Americana*), creeping barberry (*Mahonia repens*), Oregon boxleaf (*Paxistima myrsinites*), Kuntze mallow ninebark (*Physocarpus malvaceus*), New Mexico locust (*Robinia neomexicana*), mountain snowberry, and Gambel oak (*Quercus gambelii*). Herbaceous species include fringed brome (*Bromus ciliatus*), Geyer's sedge (*Carex geyeri*), Ross' sedge (*Carex rossii*), dryspike sedge (*Carex siccata*), screwleaf muhly, bluebunch wheatgrass, sprucefir fleabane (*Erigeron eximius*), Virginia strawberry (*Fragaria virginiana*), smallflowered woodrush (*Luzula parviflora*), sweetcicely (*Osmorhiza berteroi*), bittercress ragwort (*Packera cardamine*), western meadow-rue (*Thalictrum occidentale*), and Fendler's meadow-rue (*Thalictrum fendleri*) (New Mexico Department of Game and Fish, 2006).

The Conifer Woodlands and Savannas ecoregion is an area of mostly pinyon-juniper woodlands consisting of one-seed, alligator, and Rocky Mountain Junipers with some ponderosa pine at higher elevations. It often intermingles with grasslands and shrublands consisting of blue gramma, junegrass, gallet, bottlebrush squirreltail. In addition, some areas may have Gambel oak. Utah juniper and big sagebrush can be found in the Chuska Mountains. At lower elevations yuccas and cactus can be found (Griffith, et al., 2006)

The Arizona/New Mexico Subalpine Forests occur west of the Rio Grande at the higher elevations, generally above about 2,900 m [9,500 ft]. The region includes parts of the Mogollon Mountains, Black Range, San Mateo Mountains, Magdalena Mountains, and Mount Taylor. Although there are some vegetational differences from mountain range to mountain range within the region, the major forest trees include Engelmann spruce, corkbark fir (*Abies lasiocarpa* var. *arizonica*), blue spruce, white fir, and aspen. Some Douglas fir occurs at lower elevations (Griffith, et al., 2006).

Northwestern New Mexico Fauna

According to the Biota Information System of New Mexico, more than 1,100 species of amphibians, reptiles, mammals, birds, invertebrates, and fish are found throughout the state. Bird fauna is diverse with more than 500 species. Mammal diversity is high compared to other southwestern states, with approximately 184 species. New Mexico has approximately 26 species of amphibians and over 100 species of reptiles.

Common mammals found within the Northwestern New Mexico Uranium Milling Region include numerous myotis bat species, black bear, bobcat, numerous rodents, coyotes, bighorn sheep, Gunnison's prairie dogs, skunks, and squirrels. In addition, critical elk winter habitat and calving areas are located in the area (Figure 3.5-10). Currently, most of the proposed or existing ISL facilities are located within designated critical elk winter habitat. Most of the habitat in this region is found within the southern half of McKinley County and most of Cibola County. Common bird species found in the region include bluebirds, buntings, doves, ducks, cormorants, hummingbirds, jays, flycatchers, kingbirds, mockingbird, sparrows, and ravens. Raptor species include hawks such as the ferruginous hawk, red-tailed hawk, sharp shinned hawk, and Swainson's hawk; noted owl species found in the counties are the barn owl, burrowing owl, elf owl, flammulated owl, great horned owl, pygmy owl, and Mexican owl. The climax raptor found in the region is the golden eagle (Biota Information System of New Mexico, 2007).




Figure 3.5-10. Elk Winter Habitat and Calving Areas for the Northwestern New Mexico Uranium Milling Region

Description of the Affected Environment

Individual county listings can be obtained through the Biota Information System of New Mexico. A comprehensive listing of habitat types and species (with their scientific names) have been surveyed within New Mexico are compiled as part of the Southwest Regional Gap Analysis Project (New Mexico State University, 2007).

3.5.5.2 Aquatic

According to the Biota Information system of New Mexico-M, there are approximately 161 different species of fish located within the state, with approximately 48 species found in the watersheds of the region (Table 3.5-6) (Biota Information System of New Mexico, 2007). The New Mexico Comprehensive Wildlife Conservation Strategy Plan indicates that the majority of the areas in which milling would occur lie within the Zuni, Rio Grande, and the lower portion of the San Juan watersheds (New Mexico Department of Game and Fish, 2006).

Table 3.5-6. Native Fish Species Found in New Mexico

Common Name	Scientific Name
Bass, Largemouth	<i>Micropterus salmoides salmoides</i> (NM)
Bass, Smallmouth	<i>Micropterus dolomieu</i>
Bass, Striped	<i>Morone saxatilis</i>
Bass, White	<i>Morone chrysops</i>
Bluegill	<i>Lepomis macrochirus</i>
Buffalo, Smallmouth	<i>Ictiobus bubalus</i>
Bullhead, Black	<i>Ameiurus melas</i>
Bullhead, Yellow	<i>Ameiurus natalis</i>
Carp, Common	<i>Cyprinus carpio</i>
Carp, Grass	<i>Ctenopharyngodon idella</i>
Carp sucker, River	<i>Carpionodes carpio carpio</i>
Catfish, Blue	<i>Ictalurus furcatus</i>
Catfish, Channel	<i>Ictalurus punctatus</i>
Catfish, Chihuahua	<i>Ictalurus sp</i> (NM)
Catfish, Flathead	<i>Pylodictis olivaris</i>
Chub, Flathead	<i>Platygobio gracilis</i>
Chub, Gila	<i>Gila intermedia</i>
Chub, Rio Grande	<i>Gila pandora</i>
Chub, Roundtail	<i>Gila robusta</i>
Crappie, Black	<i>Pomoxis nigromaculatus</i>
Crappie, White	<i>Pomoxis annularis</i>
Dace, Longfin	<i>Agosia chrysogaster</i>
Dace, Longnose	<i>Rhinichthys cataractae</i>
Dace, Speckled	<i>Rhinichthys osculus</i> (Gila pop.)
Dace, Speckled	<i>Rhinichthys osculus</i> (Non-Gila pop.)
Killifish, Rainwater	<i>Lucania parva</i>
Minnow, Fathead	<i>Pimephales promelas</i>
Minnow, Loach	<i>Tiaroga cobitis</i>
Minnow, Roundnose	<i>Dionda episcopa</i>

1

Table 3.5-6. Native Fish Species Found in New Mexico (continued)	
Common Name	Scientific Name
Minnow, Silvery, Rio Grande	<i>Hybognathus amarus</i>
Perch, Yellow	<i>Perca flavescens</i>
Shad, Gizzard	<i>Dorosoma cepedianum</i>
Shad, Threadfin	<i>Dorosoma petenense</i>
Shiner, Golden	<i>Notemigonus crysoleucas</i>
Shiner, Red	<i>Cyprinella lutrensis</i>
Shiner, Rio Grande	<i>Notropis jemezianus</i>
Spikedance	<i>Meda fulgida</i>
Stoneroller, Central	<i>Campostoma anomalum</i>
Sucker, Bluehead, Zuni	<i>Catostomus discobolus yarrowi</i> (NM)
Sucker, Desert	<i>Catostomus clarki</i>
Sucker, Rio Grande	<i>Catostomus plebeius</i>
Sucker, Sonora	<i>Catostomus insignis</i>
Sucker, White	<i>Catostomus commersoni</i>
Sunfish, Green	<i>Lepomis cyanellus</i>
Trout, Brown	<i>Salmo trutta</i>
Trout, Gila	<i>Oncorhynchus gilae</i>
Trout, Rainbow	<i>Oncorhynchus mykiss</i>
Western Mosquito Fish	<i>Gambusia affinis</i>

2

3 The Zuni watershed also encompasses the upper Puerco watershed. The Zuni watershed has
4 an impacted water system due to settlement changes, overgrazing, and logging. The loss of
5 vegetative cover led to increased erosion, gullyng, head cutting, wide discharge fluctuations,
6 and loss of water in the system (New Mexico Department of Game and Fish, 2006). Eight
7 nonnative fish have been found in the watershed, with the green sunfish (*Lepomis cyanellus*),
8 fathead minnow (*Pimephales promelas*), and the plains killifish (*Fundulus zebrinus*)
9 comparatively common and widespread. Several sport fish have been introduced to the system
10 such as northern pike (*Esox lucius*), rainbow trout (*Oncorhynchus mykiss*), and channel catfish
11 (*Ictalurus punctatus*). Crayfish (*orconectes virilis*) have also been introduced into the system
12 (New Mexico Department of Game and Fish, 2006).

13

14 Two fish, the Roundtail Chub (*Gila robusta*) and Zuni bluehead sucker (*Catostomus discobolus*
15 *yarrowi*) and one crustacean (*Hyaella Spp.*) have been identified as species of greatest
16 conservation need (New Mexico Department of Game and Fish, 2006).

17

18 The Rio Grande watershed originates in the San Juan Mountains of Southern Colorado and
19 flows south through the entire length of New Mexico. This watershed also encompasses the
20 Arroyo Chico, Rio San Jose and Rio Puerco watersheds as previously discussed. The aquatic
21 habitats in the Rio Grande consist of reservoirs, marshes, and perennial streams (New Mexico
22 Department of Game and Fish, 2006). Numerous species have been introduced into the
23 Rio Grande Watershed. Common carp (*Cyprinus carpio*) are widespread and nonnative
24 salmonids, including rainbow trout, cutthroat subspecies (*O. clarki*) brook trout (*Salvelinus*
25 *fontinalis*), and brown trout (*Salmo trutta*) live in mountain streams. Kokanee salmon
26 (*Oncorhynchus nerka*), rainbow trout, and brown trout are present in reservoirs. Warm/cool
27 water fish include largemouth bass (*Micropterus salmoides*), smallmouth bass (*M. dolomieu*),
28 walleye (*Sander vitreus*), northern pike, white bass (*Morone chrysops*), crappie (*Pomoxis spp.*),
29 and sunfishes (*Lepomis spp.*) (New Mexico Department of Game and Fish, 2006).

Eleven fish species have been designated as a species of greatest conservation need. The Mexican tetra (*Astyanax mexicanus*), speckled chub (*Macrhybopsis aestivalis*), Rio Grande shiner (*Notropis jemezianus*), blue sucker (*Cycleptus elongates*), and gray redhorse (*Moxostoma congestum*) have disappeared from key habitats in the Rio Grande watershed. The following fish are in conservation need: Rio Grande cutthroat trout, Rio Grande chub, Rio Grande sucker, smallmouth sucker, and blue catfish (New Mexico Department of Game and Fish, 2006).

Noted native fish species historically found within the watersheds associated with sites in the Grants Uranium District include blue catfish (*Ictalurus furcatus*), desert sucker (*Catostomus clarki*), Gila chub (*Gila intermedia*), Gila topminnow (*Poeciliopsis occidentalis*), Gila trout (*Oncorhynchus gilae*), loach minnow (*Rhinichthys cobitis*), Rio Grande sucker (*Catostomus plebeius*), Rio Grande silver minnow (*Hybognathus amarus*), Rio Grande shiner, Rio Grande cutthroat trout (*Oncorhynchus clarki virginianalis*), Rio Grande chub (*Gila Pandora*), roundtail chub, spikedace (*Meda fulgida*), smallmouth buffalo (*Ictiobus bubalus*), Sonora sucker (*Catostomus insignis*), and the Zuni Bluehead sucker (Biota Information System of New Mexico, 2007).

The San Juan watershed which contains many first and second order streams found in the Chaco watershed within the milling region. The San Juan River Basin is the second largest of the three sub-basins which comprise the Upper Colorado River Basin. The San Juan River Basin drains about 97,300 km² [38,000 mi²] of southwestern Colorado, northwestern New Mexico, northeastern Arizona, and southeastern Utah (U.S. Fish and Wildlife Service, 2006). At least eight native fish species cutthroat trout, roundtail chub, Colorado pikeminnow, speckled dace, flannelmouth sucker, bluehead sucker, razorback sucker, and mottled sculpin are located within the basin. Colorado pikeminnow, razorback sucker, and the bonytail chub are federally listed as endangered species, with New Mexico listing the roundtail chub as endangered. Noted non native fish found within the higher order streams in the watershed include red shiner, common carp, fathead minnow, plains killfish, whiter sucker, brown trout, rainbow trout, and channel catfish (New Mexico Department of Game and Fish, 2006).

3.5.5.3 Threatened and Endangered Species

Federally listed threatened and endangered and species which are known to exist within habitats found within the region include the following:

- Bald Eagle—(delisted monitored).
- Black-Footed Ferret— (extirpated).
- Mexican Spotted Owl (*Strix occidentalis lucida*)—(critical habitat designated)- Mexican spotted owls nest, roost, forage, and disperse in a diverse assemblage of biotic communities. Mixed-conifer forests are commonly used throughout most of the range which may include Douglas fir and/or white fir, with codominant species including southwestern white pine, limber pine, and ponderosa pine. The understory often contains the above coniferous species as well as broadleaved species, such as Gambel oak, maples, box elder, and/or New Mexico locust. In southern Arizona and Mexico, Madrean pine-oak forests are also commonly used. Spotted owls nest and roost primarily in closed-canopy forests or rocky canyons. They nest in these areas on cliff ledges, in stick nests built by other birds, on debris platforms in trees, and in tree

cavities. In southern Utah, Colorado, and some portions of northern New Mexico, most nests are in caves or on cliff ledges in rocky canyons. Forests used for roosting and nesting often contain mature or old-growth stands with complex structure, are typically uneven-aged, multistoried, and have high canopy closure. A wider variety of trees are used for roosting, but again Douglas-fir is the most commonly used species (U.S. Fish and Wildlife Service, 2008)

- Pecos Puzzle Sunflower (*Helianthus paradoxus*)—This species is found in areas that have permanently saturated soils, including desert wetlands (cienegas) that are associated with springs, but may include stream and lake margins. When found around lakes, these lakes are usually natural cienega habitats that have been impounded (Center for Plant Conservation, 2008).
- South Western Willow Fly Catcher (*Empidonax traillii extimus*)—The southwestern willow flycatcher breeds in patchy to dense riparian habitats along streams, reservoirs, or other wetlands. Common tree or shrub species include willow, seep willow, boxelder, stinging nettle, blackberry, cottonwood, arrowweed, tamarisk (salt cedar), and Russian olive. Habitat characteristics vary across the subspecies' range. However, occupied sites usually consist of dense vegetation in the patch interior, or dense patches interspersed with openings, creating a mosaic that is not uniformly dense. In almost all cases, slow-moving or still water, or saturated soil is present at or near breeding sites during non-drought years (U.S. Fish and Wildlife Service, 2008).
- Yellow Billed Cuckoo—previously described in Section 3.2.5.3.
- Zuni Blue Head Sucker (*Catostomus dicobolus yarrowi*) (candidate)—More recent surveys (early to mid 1990s) determined the distribution of Zuni bluehead sucker in New Mexico to be limited mainly to the Río Nutria drainage upstream of the mouth of the Nutria Box Canyon. This included the mouth of Río Nutria box canyon, upper Río Nutria, confluence of Tampico Draw and Río Nutria, Tampico Spring, and Agua Remora. Definitive habitat associations for Zuni bluehead sucker have not been determined. Zuni bluehead sucker are primarily found in shaded pools and pool-runs, about 0.3 to 0.5-m 1 to 1.5-ft] deep with water velocity less than 10 cm/s [4 in/s]. Zuni bluehead suckers were found over clean, hard substrate, from gravel and cobble to boulders and bedrock (New Mexico Department Game and Fish, 2004).
- Zuni Fleabane (*Erigeron rhizomatus*)—Zuni fleabane grows in selenium-rich red or gray detrital clay soils derived from the Chinle and Baca formations. Plants are found at elevations from 2,230-2,440 m [7,300–8,000 ft] in pinyon-juniper woodland. Zuni fleabane prefers slopes of up to 40 degrees, usually with a north-facing aspect. Although the overall vegetative cover is usually high, there are few other competing plants on the steep easily erodible slopes that are Zuni fleabane's primary habitat. Zuni fleabane is found only in areas of suitable soils. These soils occur most extensively in the Sawtooth Mountains and in the northwestern part of the Datil Mountains in Catron County, New Mexico. There are 29 known sites in this area, which range in size from a fraction of an acre to about 105 hectares [260 acres]. There are two sites on the northwest side of the Zuni Mountains in McKinley County, New Mexico, and one site in Apache County, Arizona (U.S. Fish and Wildlife Service, 2008).

Description of the Affected Environment

- Rio Grande Silvery Minnow (*Hybognathus amarus*)—Currently, the Rio Grande silvery minnow is believed to occur only in one reach of the Rio Grande in New Mexico, a 280-km (174-mi) stretch of river that runs from Cochiti Dam to the headwaters of Elephant Butte Reservoir. Its current habitat is limited to about 7 percent of its former range. The Rio Grande silvery minnow uses only a small portion of the available aquatic habitat. In general, the species most often uses silt substrates in areas of low or moderate water velocity (e.g., eddies formed by debris piles, pools, and backwaters). The Rio Grande silvery minnow is rarely found in habitats with high water velocities, such as main channel runs, which are often deep and swift. The species is most commonly found in depths of less than 20 cm [7.9 in] in the summer and 31–40 cm [12.2–15.75 in] in the winter (U.S. Fish and Wildlife Service, 2007).

State listed threatened and endangered species for the region include the following:

- American marten (*Martes americana*)—The American marten is broadly distributed. It extends from the spruce-fir forests of northern New Mexico to the northern limit of trees in arctic Alaska and Canada. American martens live in mature, dense conifer forests or mixed conifer-hardwood forests. They prefer woods with a mixture of conifers and deciduous trees including hemlock, white pine, yellow birch, maple, fir and spruce. Especially critical is presence of many large limbs and fallen trees in the understory, known as coarse woody debris. These forests provide prey, protection and den sites (New Mexico Department of Game and Fish, 2008).
- Arctic peregrine falcon (*Falco peregrinus tundrius*)—Peregrine falcons live mostly along mountain ranges, river valleys, and coastlines. Historically, they were most common in parts of the Appalachian Mountains and nearby valleys from New England south to Georgia, the upper Mississippi River Valley, and the Rocky Mountains. Peregrines also inhabited mountain ranges and islands along the Pacific Coast from Mexico north to Alaska and in the Arctic tundra (U.S. Fish and Wildlife Service, 2008).
- Bald Eagle (*Haliaeetus leucocephalus*)—In New Mexico, migrating bald eagles can be found near rivers and lakes, where occasional tall trees provide lookout perches and night roosts. Reservoirs with sizable populations of migrating bald eagles include Ute, Conchas, Ft. Sumner, Santa Rosa, Elephant Butte, Caballo, Cochiti, El Vado, Heron, and Navajo (New Mexico Department of Game and Fish, 2008).
- Baird's sparrow (*Ammodramus bairdii*)—Breeds in native mixed-grass and fescue prairie. Winters in grasslands; specific winter habitat requirements not well described. Baird's Sparrow does not inhabit prairie lands where fire suppression and changes in natural grazing patterns have allowed woody vegetation to grow excessively. Some hayfields or pastures may support Baird's Sparrow where native grasses occur in sufficient quantity, but generally cultivated land is far inferior habitat relative to true prairie. Winters from southeast Arizona, southern New Mexico, and south Texas to north-central Mexico (Cornell, 2008)
- Broadbilled humming bird (*Cynanthus latirostris*)—In the United States this hummingbird is found in riparian woodlands at low to moderate elevations. In Guadalupe Canyon these woodlands are characterized by cottonwoods, sycamores, white oaks, and hackberries. Nests found in Guadalupe Canyon have been in a variety of trees, shrubs, and even forests (New Mexico Department of Game and Fish, 2004).

- 1 • Brown Pelican (*Pelecanus occidentalis*) —Brown pelicans nest on small, isolated
2 coastal islands where they are safe from predators such as raccoons and coyotes. This
3 is a potential migrant though the region (Texas Parks and Wildlife Department, 2007)
4
- 5 • Common black hawk (*Buteogallus anthracinus*) —Obligate riparian nester, dependent
6 on mature, relatively undisturbed habitat supported by a permanent flowing stream.
7 Streams less than 30-cm 12-in] deep of low to moderate gradient with many riffles, runs,
8 pools, and scattered boulders or lapped with branches provide ideal hunting conditions
9 (Public Employees for Environmental Responsibility, 2008).
10
- 11 • Costa's hummingbird (*Calypte costae*) —Occurs mainly in Southern California, Arizona,
12 Baja California, and western Mexico, but also extends into Nevada, extreme
13 southeastern Utah, and southeastern New Mexico. Habitats occupied by Costa's
14 Hummingbirds include Sonoran desert scrub, the Mojave Desert, California chaparral,
15 California coastal scrub, and the Cape deciduous forest of Baja California (Audubon
16 Society, 2007).
17
- 18 • Gray vireo (*Vireo vicinior*) —Gray Vireo breeds in some of the hottest, driest areas of
19 the American Southwest, favoring dry thorn scrub, chaparral, and pinyon-juniper and
20 oak-juniper scrub, in arid mountains and high plains scrubland. This species forages in
21 thickets, taking most of its prey from leaves, twigs, and branches of small trees and
22 bushes. Its diet on the breeding grounds consists of a variety of arthropods, including
23 large grasshoppers, cicadas, and caterpillars. Winter diet differs based on locality--birds
24 found in western Texas are primarily insectivorous, while those wintering in southern
25 Arizona and adjacent northern Mexico feed mainly on fruit (Audubon Society, 2007).
26
- 27 • Interior Least tern—previously described Section 3.3.5.3.
28
- 29 • Jemez Mountains Salamander (*Plethodon neomexicanus*) —Native to north-central
30 New Mexico. This species has been found in various localities in the Jemez Mountains
31 in Sandoval, Los Alamos, and Rio Arriba counties. This salamander typically lives on
32 shady, wooded sites at elevations of about 2,300 to 2,900 m [7,500 to 9,500 ft]. In
33 these habitats, characterized by coniferous trees, salamanders spend much of their
34 time under and in fallen logs. Old, stabilized talus slopes, especially those with a good
35 covering of damp soil and plant debris, are important types of cover for this species
36 (New Mexico Department of Game and Fish, 2008).
37
- 38 • Meadow jumping mouse (*Zapus hudsonius*) —Jumping mice are nocturnal, and in
39 New Mexico this species occurs in moist habitats dominated by damp and rich
40 vegetation. The meadow jumping mouse inhabits areas with streams, moist soil, and
41 lush streamside vegetation consisting of grasses, sedges, and forbs. Such habitats are
42 in the Jemez Mountains, and the edges of permanent ditches and cattail stands in the
43 Rio Grande Valley (New Mexico Department of Game and Fish, 2008).
44
- 45 • Neo tropic cormorant (*Phalacrocorax brasilianus*) —This cormorant is found from
46 southern New Mexico to southern Louisiana. Southward through Central America and
47 the Caribbean to South America. Neotropic cormorants also may wander northward to
48 the Bernalillo area and westward to the Gila Valley. This bird is rare in southern Hidalgo
49 County, the area near Alamogordo, and in the lower Pecos Valley from Bitter Lake
50 National Wildlife Refuge southward (New Mexico Department of Game and Fish, 2008).

Description of the Affected Environment

- Peregrine falcon (*Falco peregrines*)—In New Mexico the breeding sites of peregrine falcons are on cliffs in wooded and forested habitats, with large "gulfs" of air nearby in which these predators can forage (New Mexico Department of Game and Fish, 2008).
- Rio Grande shiner (*Notropis jemezianus*)—The Rio Grande shiner is found in the Rio Grande drainage, from just above the mouth to Pecos River (north in Pecos River to Sumner Lake, New Mexico) and (formerly) Rio Grande, New Mexico (where now extirpated); absent from large sections of Rio Grande and Pecos River in western Texas; occurs in Rio San Juan, Rio Salado, and Rio Conchos, Mexico; common in lower Rio Grande, less common elsewhere. Can be found in runs and flowing pools of large open weedless rivers and large creeks with bottom of rubble, gravel, and sand, often overlain with silt (NatureServe, 2008).
- Spotted bat (*Euderma maculatum*) —The rarity of this bat and the diverse habitats in which it has been seen have caused confusion about its preferences. Some have been captured in pine forests at high elevations (8,000-9,000 ft); others came from a pinyon pinejuniper association; and still others from desert scrub areas. Spotted Bats are known only from about 20 locations in western and southern New Mexico (New Mexico Department of Game and Fish, 2008).
- South Western Willow flycatcher—previously described in this section as a federally listed species.
- Wrinkled marsh snail (*Stagnicola caperata*)—The wrinkled marsh snail occurs in such habitats as vegetated ditches, marshes, streams, and poinds, typically that are seasonally dry. Such a site is occupied by the New Mexico population in the Jemez Mountains, where the habitat is a shallow pond at 2,600 m elevation. The species also occurs in areas of perennial water, including the former population at Bitter Lake National Wildlife Refuge (USACE, 2007).
- Zuni Bluehead sucker—previously described in this section as a federally listed species.

3.5.6 Meteorology, Climatology, and Air Quality

3.5.6.1 Meteorology and Climatology

Temperature in New Mexico is influenced more by elevation than latitude. Mean annual temperatures range from 17 °C [64 °F] in the southeast to less than 4 °C [40 °F] in the high mountains and northern valleys (National Climatic Data Center, 2005). New Mexico typically experiences variations between daytime and nighttime temperatures. Table 3.5-7 identifies two climate stations located in the Northwestern New Mexico Uranium Milling Region. Climate data for these stations are found in the National Climatic Data Center's Climatology of the United States No. 20 Monthly Station Climate Summaries for 1971–2000 (National Climatic Data Center, 2004). This summary contains climate data for 4,273 stations throughout the United States and some territories. Table 3.5-8 contains temperature data for two stations in the Northwestern New Mexico Uranium Milling Region.

The precipitation and snow that New Mexico receives comes from both the Pacific Ocean to the west and the Gulf of Mexico to the southeast. Average annual precipitation ranges from 25 cm [10 in] to more than 50 cm [20 in] at higher elevations (National Climatic Data Center, 2005). In

Table 3.5-7. Information on Two Climate Stations in the Northwestern New Mexico Uranium Milling Region*

Station (Map Number)	County	State	Longitude	Latitude
Grants Milan AP	Cibola	New Mexico	107°54W	35°10N
McGaffey 5 SE	McKinley	New Mexico	108°27W	35°20N

*National Climatic Data Center. "Climatography of the United States No. 20: Monthly Station Climate Summaries, 1971–2000." Asheville, North Carolina: National Oceanic and Atmospheric Administration. 2004.

Table 3.5-8. Climate Data for Stations in the Northwestern New Mexico Uranium Milling Region*

		Grants Milan AP	McGaffey 5 SE
Temperature (°C) †	Mean—Annual	10.4	5.9
	Low—Monthly Mean	−0.6	−4.5
	High—Monthly Mean	22.1	17.2
Precipitation (cm) ‡	Mean—Annual	27.6	51.6
	Low—Monthly Mean	1.1	1.7
	High—Monthly Mean	5.3	7.0
Snowfall (cm)	Mean—Annual	23.9	136
	Low—Monthly Mean	0	0
	High—Monthly Mean	7.4	26.9

*National Climatic Data Center. "Climatography of the United States No. 20: Monthly Station Climate Summaries, 1971–2000." Asheville, North Carolina: National Oceanic and Atmospheric Administration. 2004.
†To convert Celsius (°C) to Fahrenheit (°F), multiply by 1.8 and add 32.
‡To convert centimeters (cm) to inches (in), multiply by 0.3937.

summer, the source of precipitation is usually brief, but often intense thunderstorms. For most of the state, 30 to 40 percent of the year's annual moisture falls in July and August. Typically, New Mexico does not experience widespread floods. Heavy thunderstorms can cause local flash floods. Heavy rains or rain in conjunction with snowmelt can cause large rivers to flood. Table 3.5-8 contains precipitation data for two stations in the Western New Mexico Uranium Milling Region. The wettest month for both stations identified in Table 3.5-8 is August and, based on the snow depth data, snow pack melting usually occurs earlier in the summer (National Climatic Data Center, 2004). One of the stations is in Cibola County and the other is in McKinley County. Data from National Climatic Data Center's Storm Events Database from 1950 to 2007 indicates that the majority of thunderstorms in Cibola and McKinley Counties occur somewhat evenly between May and September (National Climatic Data Center, 2007).

In winter, the precipitation usually falls as snow in the mountains; however the precipitation in the valleys can be either rain or snow. Table 3.5-9 contains snowfall data for two stations in the Northwestern New Mexico Uranium Milling Region.

As an example, Figure 3.5-11 shows a wind rose for Gallup, New Mexico for 1991. Winds are predominantly from the west southwest and southwest. Wind speeds are depicted in knots where 1 knot is approximately equal to 0.51 m/s [1.7 ft/s]. Wind roses such as these should be

Table 3.5-9. U.S. Environmental Protection Agency Class I Prevention of Significant Deterioration Areas in New Mexico and Arizona*

New Mexico	Arizona
Bandelier Wilderness Bosque del Apache Wilderness Carlsbad Caverns National Park Gila Wilderness Pecos Wilderness Salt Creek Wilderness San Pedro Parks Wilderness Wheeler Peak Wilderness White Mountain Wilderness	Chiricahua National Monument Wilderness Chiricahua Wilderness Galiuro Wilderness Grand Canyon National Park Mazatzal Wilderness Mount Baldy Wilderness Petrified Forest National Park Pine Mountain Wilderness Saguaro Wilderness Sierra Ancha Wilderness Superstition Wilderness Sycamore Canyon Wilderness
*Modified from Code of Federal Regulations. "Prevention of Significant Air Deterioration of Air Quality." Title 40—Protection of the Environment, Part 81. Washington, DC: U.S. Government Printing Office. 2005.	

obtained for the actual location of the facility for preferably a period of time of 1 year or longer. This data can be used for dispersion estimates.

The pan evaporation rates for the Northwest New Mexico Uranium Milling Region range from about 114 to 152 cm [45 to 60 in] (National Weather Service, 1982). Pan evaporation is a technique that measures the evaporation from a metal pan typically 121 cm [48 in] in diameter and 25 cm [10 in] tall. Pan evaporation rates can be used to estimate the evaporation rates of other bodies of water such as lakes or ponds. Pan evaporation rate data is typically available only from May to October. Freezing conditions often prevent collection of quality data during the other part of the year.

3.5.6.2 Air Quality

The general air quality general description for the Northwestern New Mexico Uranium Milling Region would be similar to the description in Section 3.2.6. for the Wyoming West Uranium Milling Region.

As described in Section 1.7.2.2, the permitting process is the mechanism used to address air quality. If warranted, permits may set facility air pollutant emission levels, require mitigation measures, or require additional air quality analyses. Except for Indian Country, New Source Review permits in New Mexico are regulated under the EPA-approved State Implementation Plan. For Indian Country in New Mexico, the New Source Review permits are regulated under 40 CFR 52.21 (EPA, 2007a).

State Implementation Plans and permit conditions are based in part on federal regulations developed by the EPA. The NAAQS are federal standards that define acceptable ambient air concentrations for six common nonradiological air pollutants: nitrogen oxides, ozone, sulfur oxides, carbon monoxide, lead, and particulates. In June 2005, EPA revoked the 1-hour ozone

1




Figure 3.5-11. Windrose for Gallup, New Mexico, Airport for 1991 (New Mexico Environmental Department, 2007)

2
3 standard nationwide in all locations except certain Early Action Compact Areas. None of the 1-
4 hour ozone Early Action Compact Areas are in New Mexico. States may develop standards that
5 are stricter or supplement the NAAQS. New Mexico has a more restrictive standard for carbon
6 monoxide throughout the state and for sulfur dioxide in a small area around the city of Hurley.
7 This area around Hurley is not within the Northwest New Mexico Uranium Milling Region. New
8 Mexico also has a nitrogen dioxide standard with a 24-hour averaging time (New Mexico
9 Environment Department, 2002).

10

Prevention of Significant Deterioration requirements identify maximum allowable increases in concentrations for particulate matter, sulfur dioxide, and nitrogen dioxide for areas designated as attainment. Different increment levels are identified for different classes of areas and Class I areas have the most stringent requirements.

The Northwestern New Mexico uranium milling region air quality description focuses on two topics: NAAQS attainment status and PSD classifications in the region.

Figure 3.5-12 identifies the counties in and around the Northwestern New Mexico Uranium Milling Region that are partially or entirely designated as nonattainment or maintenance for NAAQS at the time this GEIS was prepared (EPA, 2007b). The Northwestern New Mexico Uranium Milling Region covers portions of New Mexico and borders Arizona. All of the area within this milling region is classified as attainment. Portions of two counties in New Mexico are not in attainment: Bernalillo County (central New Mexico) and Dona Ana County (south central New Mexico). The city of Albuquerque in Bernalillo County is designated as maintenance for carbon monoxide. The northwest part of Bernalillo County is only several kilometers from the Northwestern New Mexico Uranium Milling Region border, however, the Albuquerque is about 50 km [31 mi] from this border. The city of Anthony in Doña Ana County is designated as nonattainment for PM₁₀. The Sunland Park area of Doña Ana County was designated as nonattainment for the 1-hour ozone standard until the EPA revoked the standard in 2005. Several counties in southern Arizona, including one that borders New Mexico, are not in attainment. However, the one Arizona county (Apache County) that borders the Northwestern New Mexico Uranium Milling Region is in attainment.

Table 3.5-9 identifies the Prevention of Significant Deterioration Class I areas in New Mexico and Arizona. The Class I areas in and around the Northwestern New Mexico Uranium Milling Region are shown in Figure 3.5-13. There are no Class I areas in the Northwestern New Mexico Uranium Milling Region (Code of Federal Regulation, 2005).

3.5.7 Noise

The existing ambient noise levels for undeveloped rural in the Northwestern New Mexico Uranium Milling Region would be similar to those described in Section 3.2.7 for the Wyoming West Uranium Milling Region (up to 38 dB). The largest communities in the region include Gallup with a population of more than 20,000, Grants with a population of about 9,000, and Zuni Pueblo (about 6,400) (see Section 3.5.10). Urban noise levels in these communities and the smaller surrounding population centers would be similar to those (up to about 78 dB) for other urban areas (Washington State Department of Transportation, 2006).

As described in Section 3.5.2, two major highways cross the Northwestern New Mexico Uranium Milling Region, Interstate 40 runs east west, and U.S. Highway 491 runs north from Gallup. There are also several state undivided highways, but the area is only sparsely served by paved roads. Traffic counts for Interstate-40 are higher than those reported for I-80 in Wyoming, with annual average daily traffic reported at about 16,500 just east of the New Mexico/Arizona line (New Mexico Department of Transportation, 2007). Traffic counts for U.S. Highway 491 are less, with annual average daily traffic of about 9,700 north of Gallup (New Mexico Department of Transportation, 2007). This suggests that ambient noise levels near these highways might be higher than the levels measured for I-80 (Wyoming Department of Transportation, 2005; Federal Highway Administration, 2004; see also Section 3.2.7).




Figure 3.5-12. Air Quality Attainment Status for the Northwest New Mexico Uranium Milling Region and Surrounding Areas (EPA, 2007a)

The potential uranium projects in the region are more than 8 km [5 mi] from Interstate 40 and ambient noise levels would not be affected by highway noise. In some cases, such as at Crownpoint, the proposed facility would be located close to a small community, and the ambient noise levels would be expected to be slightly higher. Areas of special sensitivity to potential noise impacts could include areas of special significance to the Native American culture in the region (see Section 3.5.8).

3.5.8 Historical and Cultural Resources

The New Mexico State Historic Preservation Office (SHPO) is responsible for the oversight of federal and state historic preservation compliance laws, regulations and statutes. The Cultural

1




Figure 3.5-13. Prevention of Significant Deterioration Class I Areas in the Northwestern New Mexico Uranium Milling Region and Surrounding Areas (40 CFR Part 81)

Properties Act (Sections 16-6 through 18-6-23, New Mexico Statutes Annotated 1978) was enacted in 1969 and amended several times in the ensuing years. It established the State

Historic Preservation Division and Cultural Properties Review Committee which issues permits for survey and excavation on state lands, and for the excavation of burials. Burial excavation permits are specifically required by the Unmarked Burial Statute (18-6-11.2, 1989) and the Marked Burial Statute (30-12-12, 1989) for human remains found on state or private lands; whereas, the NAGPRA applies to federal lands. The Reburial Grounds Act (18-6-14, 2006) provides for the designation of reburial areas for unclaimed human remains. The Cultural Properties Act also requires that state agencies provide the New Mexico SHPO with the opportunity to participate in planning activities that would affect properties on the State Register of Cultural Properties or the National Register of Historic Places. The Prehistoric and Historic

Sites Preservation Act of 1969 (Sections 18-8-1 through 18-8-8, NMSA 1978) prohibits the use of state funds that would adversely affect sites on the State or National Registers, unless the state agency demonstrates that there is no feasible or prudent alternative. The Cultural Properties Protection Act (Sections 18-6A-1 through 18-6A-6, New Mexico Statutes Annotated 1978) enacted in 1993, encourages state agencies to consult with the New Mexico SHPO in order to develop programs that will identify cultural properties and ensure that they will not be inadvertently damaged or destroyed. Lastly, Executive Order No. 2005-003 recognizes the sovereignty of Native American tribes in the state of New Mexico and provides that state agencies should conduct tribal consultation on the protection of culturally significant places and the repatriation of human remains and cultural items. Information on the New Mexico SHPO can be found at the following link: <<http://www.nmhistoricpreservation.org>>.

The United States government and the State of New Mexico recognize the sovereignty of certain Native American tribes. These tribal governments have legal authority for their respective reservations. Executive Order 13175 requires executive branch federal agencies to undertake consultation and coordination with Indian tribal governments on a government-to-government basis. NRC, as an independent federal agency, has agreed to voluntarily comply with Executive Order 13175.

In addition, the National Historic Preservation Act provides these tribal groups with the opportunity to manage cultural resources within their own lands under the legal authority of a Tribal Historic Preservation Office (THPO). The THPO therefore replaces the New Mexico SHPO as the agency responsible for the oversight of all federal and state historic preservation compliance laws. Both the Navajo Nation and Zuni Pueblo have a recognized Tribal Historic Preservation Office (THPO) program. Other tribes have historic and cultural preservation offices that are not recognized as THPOs, but they should be consulted where they exist (see appended New Mexico tribal consultation list for Cibola and McKinley Counties).

The Navajo Nation has passed the Natural Resources Protection Act of 2005, which is designed to "ensure that no further damage to the culture, society, and economy of the Navajo Nation occurs because of uranium mining within the Navajo Nation ..." An insight into the affects of uranium exploration on traditional Navajo life is provided in the recent publication entitled *The Navajo People and Uranium Mining* (Udall, et al. 2007). The Navajo Nation Code also states that "the six culturally significant mountains...Tsoodzil...must be respected, honored and protected for they, as leaders, are the foundation of the Navajo Nation (Navajo Nation, 2005, pp. 22-23)." *Tsoodzil* (Turquoise Mountain) is the Navajo word for Mount Taylor some 24 km [15 mi] north of Grants, New Mexico and, in Navajo tradition, marks the southern boundary of the Navajo Dinétah or traditional homeland.

3.5.8.1 New Mexico Historic and Cultural Resources

McKinley and Cibola counties are rich in cultural resources. In fact, the first highway salvage archaeological excavations in the nation were conducted along old Route 66 in this vicinity during the 1950s. Archaeological compliance work continues through the 21st century in respect to a variety of economic activities, including highway construction, energy development, tourism at the national monuments and the realignment of military installations. Cultural resource overviews and Class II surveys of the region have therefore been provided by several federal agencies; however, they date to the 1980s when most of the energy related development was initiated. The San Juan Basin Regional Uranium Study was certainly one of the most important of these studies (Broster and Harrill, 1982; Dulaney and Dosh 1981; Plog and Wait 1979; Powers, et al., 1983; Tainter and Gillio, 1980).

Description of the Affected Environment

Interstate 40 passes through Albuquerque, Grants and Gallup, acting as a primary east-west link across the region. New Mexico State Road 491 heads north from Gallup to Shiprock and the Four-Corners area. Lastly, Grants is connected to Chaco Canyon National Monument by way of State Road 371. A variety of archaeological projects have therefore been conducted in respect highway-related compliance work (e.g., Damp, et al. 2000; Gilpin, 2007).

McKinley and Cibola counties have been a major focus of energy development activities, including coal, uranium and natural gas pipeline projects. The McKinley Coal Mine and the Laguna uranium mine represent two examples of extensive surface mining operations (Allen and Nelson, 1982; Kelley, 1982). In addition, the ENRON and El Paso pipeline projects have cross cut the region to supply the west with natural gas from sources in northwest New Mexico (Winter, 1994).


Three national monuments are located within the Norwestern New Mexico Uranium Milling Region, Chaco Canyon, El Morro, and El Malpais. Although Chaco Canyon is situated to the north of Grants, New Mexico in San Juan County, several outlying components of Chaco National Monument are present in Cibola and McKinley Counties including the Red Mesa Valley group east of Gallup, the Cebolleta Mesa Group, Puerco of the West Group and portions of the South Chaco Slope Group (Marshall, et al., 1979; Powers, et al., 1983). El Morro and El Malpais National Monuments are also located near Grants (Powers and Orcutt, 2005a; Murphy, et al., 2003).

Fort Wingate is a closed military installation that has been extensively surveyed for cultural resources. The former Army munitions depot is located south of I-40 between Gallup and Grants. These lands contain numerous archaeological sites and have ancestral ties to both Zuni Pueblo and the Navajo Nation (Schutt and Chapman, 1997; Perlman, 1997).

A total of 21,625 archaeological sites have been recorded in McKinley and Cibola counties as of this writing. A single Class II sample survey identified an average density of 6 sites/km² [15 sites/mi²] for the southern San Juan Basin (Dulaney and Dosh, 1981); however, site densities as high as 12 sites/km² [30 sites/mi²] were identified on Cebolleta Mesa (Broster and Harrill, 1982). Table 3.5-10 provides a summary of sites recorded by time period for McKinley and Cibola Counties and Figure 3.5-14 illustrates the distribution of these sites across the counties. However, this distribution only includes those areas that have been systematically surveyed for cultural resources. Together these resources represent over 10,000 years of human land-use in the region. The following is a brief review of the Native American occupation of the area.

Table 3.5-10. Number of Recorded Sites by Time Period and County

Period	County	
	McKinley	Cibola
Paleoindian	18	34
Archaic	426	359
Ancestral Pueblo	8,211	2,742
Historic Pueblo	575	290
Navajo	4,476	378
Other Historic	518	1,057
Undetermined	2,822	2,331
Total*	15,040	6585
*Note: Because many sites include multiple temporal components, the total number of sites presented above does not reflect the total number of components (occupations) that might exist at each site.		



Documented Archaeological Sites in McKinley and Cibola Counties, February 2008





Figure 3.5-14. Distribution of Recorded Archaeological Sites in McKinley and Cibola Counties, New Mexico

Paleoindian (ca. 10,000 to 6000 B.C.)

The Paleoindian occupation of the region is primarily represented by the presence of isolated projectile points with a few campsites (Figure 3.5-15). Clovis (10,000–9,000 B.C.), Folsom (9,000–8,000) and Late Paleoindian (8,000–6,000 B.C.) points have been identified at various locations across the landscape. The Clovis inhabitants presumably hunted a range of large animal species including mammoth; whereas, Folsom hunters focused on migratory bison herds and Late Paleoindian hunters on bison, with other animal and plant species (Amick, 1994; Broster and Harrill, 1982; Judge, 2004; Stanford, 2005).

Archaic (ca. 6,000 B.C to A.D. 400)

The Archaic occupation of the region is characterized by the presence of numerous temporary campsites (Figure 3.5-16). Early Archaic (6,000–4,000 B.C.) and Middle Archaic (4,000–2000 B.C.) sites appear to be less common than those occupied during the Late Archaic (2000 B.C.–A.D. 400); however, this may be a product of differential preservation and the exposure of subsurface deposits, rather than differences in the degree to which these groups occupied the area. Early and Middle Archaic groups gathered a variety of plant species, while hunting medium to small-size game. In contrast, domesticated maize first appears in New Mexico by 2100 B.C., probably as a supplement to gathered plant foods, with the first evidence of simple irrigation perhaps as early as 1000 B.C. (Damp, et al., 2002; Huber and Van West, 2005; Simmons, 1986; Vierra, 2008).

Ancestral Puebloan (ca. A.D. 400 to 1540)

For many years, archaeologists referred to the prehistoric culture that arose in the San Juan Basin after the Archaic period as the “Anasazi,” a word borrowed from the Navajo that means “old people” or “enemy ancestors” (Kantner, 2004); although this term continues to be widely used among archaeologists and the public alike, many contemporary Pueblo people find the use of Anasazi to be offensive. Although controversy about this issue continues (Kantner, 2004 and Riggs, 2005), archaeologists and government agencies increasingly use the term “Ancestral Puebloan” in place of Anasazi, a practice that is followed here.

The Ancestral Puebloan period appears to have emerged directly from the preceding Archaic period, and begins with the initial appearance of pottery and the bow and arrow, more elaborate pit structure architecture, and the more intensive use of maize agriculture. Although a number of chronological sequences for this period have been proposed for the region, the two major sequences currently in use are the Cebolleta Mesa and Pecos Chronologies (Kidder, 1927), (Table 3.5-11, Figure 3.5-17).

Basketmaker II (ca. 500 B.C. to A.D. 400)

Basketmaker II (or Late Archaic) represents a continuation of the previous hunting and gathering lifestyle. However, important changes in subsistence and social organization were occurring with a growing dependence on the cultivation of maize. Recent excavations in the region have documented habitation sites with houses, storage pits and refuse areas. High water table farming adjacent to playa settings appears to have been an important niche for early maize cultivation, with numerous storage features having been discovered in these contexts. In addition, the earliest evidence of water diversion through irrigation channels is also represented. Lastly, important changes in technology were also occurring including the use of ceramic containers, and the bow and arrow (Damp, et al. 2002; Kearns, et al., 1998; Vierra, 1994; 2008).

1





Figure 3.5-15. Paleoindian Sites

2

1



Density of Archaic-Period Sites in McKinley and Cibola Counties, February 2008





Figure 3.5-16. Distribution of Archaic-Period Sites

2

1

Table 3.5-11. Cebolleta Mesa and Pecos Chronologies		
Cebolleta Mesa Sequence	Dates B.C./A.D.	Pecos Classification
—	Ca. 500 BC–AD 500	Basketmaker II
Lobo Period	?–700 AD	Basketmaker III
White Mound Phase	700–800	Basketmaker III/Pueblo I
Kiatuthlana Phase	800–870	Pueblo I
Red Mesa Phase	850–950	Early Pueblo II
Cebolleta Phase	950–1100	Pueblo II
Pilares Phase	1100–1200	Pueblo III
Kowina Phase	1200–1400	Pueblo III to IV
Cubero Phase	1400–1540	Late Pueblo IV
Acoma Phase	1540–present	Pueblo V/Historic Pueblo

2

3

Basketmaker III (ca. A.D. 400 to 700)

4

In comparison to the preceding Late Archaic period, Basketmaker III material culture is characterized by the introduction of the bow and arrow and fired ceramic vessels.

7

Basketmaker III sites in the San Juan region also featured larger and more elaborate pit habitation structures, larger villages, and evidence for increased trade and greater reliance on agriculture, including both corn and beans (Reed, 2000b). Although Basketmaker III sites have been identified throughout McKinley and Cibola counties, these sites typically date to the later portion of this time period and transition gradually into Pueblo I occupations, with few major cultural differences between them (Tainter and Gillio, 1980). In general, Basketmaker III sites are fairly rare in most of the McKinley/Cibola region compared to other areas to the north and west (Cordell, 1979; Orcutt, et al., 2005; Powers and Orcutt, 2005b; Schutt and Chapman, 1997; Tainter and Gillio, 1980). In McKinley County, however, many sites that become important during the later Pueblo II period were initially occupied at this time (Powers, et al., 1983).

17

18

Pueblo I (ca. A.D. 700 to 900)

19

20


The Pueblo I period is distinguished from the Basketmaker III period by the first appearance of painted black-on-white pottery. Although a shift away from living in subterranean pit structures and into above-ground rooms is also typically part of the Basketmaker III/Pueblo I transition (Reed, 2000a), pithouses remained the dominant structure type in much of McKinley and Cibola counties until fairly late in the Pueblo I period, with small surface rooms primarily used for storage (Schutt and Chapman, 1997; Tainter and Gillio, 1980). Small above-ground pueblos constructed from masonry or jacal (wattle-and-daub) began to be used for habitation in some areas by the end of the Pueblo I period (Schutt and Chapman, 1997). Kivas—subterranean structures with a specialized ceremonial function—also made their first appearances during this period (Schutt and Chapman, 1997). Although Pueblo I-period sites are not particularly common in McKinley and Cibola counties, they are more numerous than Basketmaker III sites, and represent the first substantial Ancestral Puebloan occupations in many areas (Schachner and Kilby, 2005; Schutt and Chapman, 1997; Tainter and Gillio, 1980).

31

32

33

34



Density of Ancestral Pueblo Sites in McKinley and Cibola Counties, February 2008




Figure 3.5-17. Distribution of Ancestral Puebloan Sites

Pueblo II (ca. A.D. 900 to 1100)

The Pueblo II period represents a considerable change in Ancestral Puebloan culture throughout the Four Corners region, including the present study area (Powers, et al. 1983, Schutt and Chapman 1997, Tainter and Gillio 1980). Blocks of contiguous, above-ground masonry rooms become the primary focus of occupation, with below-ground structures increasingly shifting to a predominantly ceremonial function (Powers and Orcutt, 2005b; Schutt and Chapman, 1997). Sites are often much larger than in the preceding Pueblo I period, and populations increase steeply throughout McKinley and Cibola counties: in many areas, populations during Pueblo II reach a peak that is not exceeded during the prehistoric period (Tainter and Gillio, 1980).

This period also marks the development of the Chacoan regional system, an event with major repercussions for the entire Four Corners region (Kantner and Mahoney, 2000; Noble, 2004; Powers, et al., 1983). Beginning around A.D. 850, Ancestral Puebloan peoples living in Chaco Canyon, located just north of McKinley County (Judge, 2004; Powers, et al., 1983; Windes, 2004) began constructing a series of elaborate, carefully planned multistory masonry structures today known as "great houses" (Windes, 2004). Although rooted in the Puebloan architecture of previous periods, the great houses were larger than contemporary structures anywhere else in the Puebloan world (Mills, 2002b). By the mid-13th century, when major construction ceased, at least 18 great houses had been constructed in and around the canyon, the largest reaching 4 or more stories and incorporating hundreds of rooms and an elaborate, decorative core-and-veneer masonry style (Judge, 2004; Mahoney and Kantner, 2000; Mills, 2002b).

Nor was great house construction limited to Chaco Canyon. Starting at about A.D. 950, great houses began to be built beyond the canyon at numerous locations throughout the San Juan Basin. More than 200 great houses with Chacoan-style architecture and features have been identified to date across an area stretching from eastern Arizona and southern Colorado to the edges of the Jemez Mountains and the foothills of Mount Taylor. Outlier sites in McKinley and Cibola counties include Casamero, Kin Nizhoni, and Village of the Great Kivas (Mahoney and Kantner, 2000; Marshall, et al., 1979). Southern and eastern areas near Acoma and Laguna are less clearly part of the Chaco system, exhibiting clear differences from sites in the San Juan Basin, (Tainter and Gillio, 1980), but outliers may exist in these areas as well (Powers and Orcutt, 2005b). Outlying great houses are typically located among much smaller and less elaborate masonry pueblos and are often accompanied by distinctive structures including extremely large "great kivas" and Chacoan roads. These roads are intentionally constructed trails that typically measure 8 to 12 m [26 to 39 ft] in width and incorporate raised beds, borders, gates, stairways, and other features (Mahoney and Kantner, 2000; Mills, 2002b; Powers and Orcutt, 2005b). Their function is not well-understood, but recent studies suggest they may link ceremonially and ritually important features of the Chacoan landscape (Kantner, 1997; Van Dyke, 2004).

The function and meaning of Chacoan great houses are not well-understood, but most evidence suggests they were not simply residential structures. Excavated great houses in Chaco Canyon typically contain few rooms with cooking hearths and very little household trash, leading some archaeologists to suggest that even the largest structures never housed more than 100 permanent residents (Mills, 2002b). Most archaeologists now believe these structures served some sort of public function, perhaps as part of a ceremonial system centered around Chaco itself. However it functioned, Chaco's far-reaching influence served to funnel trade goods into the canyon. Recent studies of ceramic and lithic artifacts, wooden roof beams, and

even foodstuffs like corn from great houses in the canyon suggest that many of these goods were brought in from far-flung areas such as the Chuska Mountains in eastern Arizona, the Mesa Verde area in southern Colorado, and the Mount Taylor region (Cordell, 2004; Mills, 2002b; Toll, 2004).

Pueblo III (ca. A.D. 1100 to 1300)

Great house construction within Chaco Canyon itself ceased by about A. D. 1130, and most of the canyon's occupants appear to have moved elsewhere by the late twelfth century (Judge, 2004; Mills, 2002b). Many factors probably contributed to the demise of Chaco, but a series of major droughts that afflicted the region throughout much of the 12th century may have had a particularly influential role (Mills, 2002b). Beyond Chaco Canyon, however, many great house communities remained occupied throughout the 1100s, retaining many aspects of their Chacoan origins but incorporating new and distinctly different features as well (Mills, 2002b). Perhaps spurred by drought, populations declined throughout much of McKinley and Cibola counties (Kintigh, 1996; Roney, 1996; Tainter and Gillio, 1980). New settlements founded during this period were frequently larger and more compact than the great house communities of the preceding period as populations aggregated in areas more conducive to conserving and managing water (Kintigh, 1996). Populations in some areas appear to have recovered and stabilized somewhat by the early thirteenth century (Powers and Orcutt, 2005a; Roney, 1996). The process of abandonment and aggregation began to accelerate again by the late 1200s, however, as renewed drought increasingly pushed Pueblo populations into relatively well-watered areas along the Zuni River to the west and the Rio San Jose to the east (Kintigh, 1996; Roney, 1996; Tainter and Gillio, 1980).

Pueblo IV (ca. A.D. 1300 to 1540)

The settlement reorganization that began during the Pueblo III period continued during Pueblo IV. By A.D. 1400, most of the Four Corners region was abandoned, with remnant populations concentrated in the Zuni and Rio San Jose areas and at the Hopi mesas in Arizona (Huntley and Kintigh, 2004; Kintigh, 1996; Roney, 1996). The number of sites present in these areas continued to drop as populations aggregated in large villages, but the compactly laid-out pueblos that remained were often extremely large, with several including more than 1,000 rooms (Huntley and Kintigh, 2004). By the late Pueblo IV period, the vast majority of Puebloan people in west-central New Mexico were at least part-time residents of one of these large pueblos: the smaller habitation sites that characterized earlier periods were virtually absent in many areas (Huntley and Kintigh, 2004; Roney, 1996). These newly aggregated large villages shared many similarities across the region settlements typically consisted of blocks of contiguous rooms arranged around plaza areas used for domestic activities and public rituals. At larger sites, these roomblocks were often two or more stories tall. Sites were also frequently located in highly defensive locations, especially early in the period (Huntley and Kintigh, 2004; Roney, 1996; Tainter and Gillio, 1980).

Historic Pueblo (post A.D. 1540)

By the mid-16th century, Puebloan groups occupied no more than ten villages in west-central New Mexico: six to nine Zuni-speaking pueblos arrayed along the lower Zuni River and its tributaries south of modern Gallup (Huntley and Kintigh, 2004) and the single Keres-speaking village of Acoma, located on a mesa top in eastern Cibola county along the Rio San Jose (Adams and Duff, 2004) (Figure 3.5-18). The first contact between these villages and the Spanish came in 1539, when a small expedition led by Franciscan friar Marcos de Niza and the




Figure 3.5-18. Distribution of Historic Pueblo Sites

Description of the Affected Environment


former slave Esteban entered the Zuni region, only to return abruptly to Mexico when Esteban was killed. (Ferguson and Hart, 1985; Spicer, 1962). The much larger expedition of Francisco Vasquez de Coronado fought a battle with the Zuni in July 1540 outside the village of Hawikuh and stopped briefly at Acoma on its way to the Rio Grande valley (Ferguson and Hart, 1985; Flint and Flint, 2005). More sustained contact with the Spanish empire came in 1598, when both the Zuni and Acoma areas were formally subjugated by the expedition of Juan de Oñate (Spicer, 1962).

Franciscan missions were established at both Zuni and Acoma in 1629, but the distance between Zuni and the center of Spanish power along the Rio Grande allowed the Zuni to retain a degree of cultural and religious independence (Ferguson and Hart, 1985; Spicer, 1962). Franciscan missions at Acoma and the Zuni villages of Hawikuh and Halona:wa operated until the Pueblo Revolt of 1680, when the Spanish were driven from New Mexico for a dozen years, but missionization in the Zuni region continued only sporadically after the Spanish reconquest in the late 1600s. At both Acoma and Zuni, however, European infectious diseases and the economic demands of the colonizers decimated Puebloan populations: at Zuni, the six or more villages inhabited at contact dwindled to three by 1680, and only one village, the present pueblo of Zuni, was reoccupied after the reconquest (Mills, 2002a). To the east, Acoma remained the only village along the Rio San Jose until 1697, when the pueblo of Laguna was established by a group of Acoma dissidents and refugees from other villages after the Spanish reconquest (Ellis, 1979).

More benign aspects of colonialism included new economic opportunities afforded by the food crops and domesticated animals brought by the Spanish. Sheepherding, in particular, began at both Zuni and Acoma as early as the mid-17th century, and by the mid-eighteenth century the Zunis grazed more than 15,000 sheep across an area extending as far as 112 km [70 mi] from the central pueblo itself (Ferguson and Hart, 1985; Schutt and Chapman, 1997). Small, temporary campsites associated with sheepherding and agriculture are among the most common historic period Puebloan archaeological sites from the 1600s into the 20th century (Ferguson, 1996; Schutt and Chapman, 1997).

Navajo (ca. 1700 to present)

With the exception of the areas just discussed, much of the northern Southwest, including northwestern New Mexico was abandoned by Ancestral Puebloan groups during the 14th century, followed by the expansion of Athabaskan hunter-gatherers into these vacated areas, perhaps as early as the late 15th century (Dean, et al. 1994; Towner, 1996). The Athabaskan-speaking groups are believed to have been the ancestors of today's Navajo and Apachean groups in the Southwest. The ancestral Navajo groups subsequently adopted maize cultivation and later moved south into the southern San Juan Basin by the 1700s (Figure 3.5-19). The 18th century Navajo migration southward was due to several factors including conflict with the Comanches and Utes, and drought and disease outbreaks. Records of Navajo baptisms at the Cebolleta Mission occur after 1749, with Navajo raids on local settlers and Laguna Pueblo Indians being reported in the late 1700s (Brugge, 1968; Correll, 1976; Reeve, 1959). This conflict continued through the 1800s, although the Navajos in the Mount Taylor (Tsodzil) area were also involved in trade relations with both local Spanish and Pueblo Indians. Nonetheless, in 1864 all the Navajos residing in the region were forcibly moved to Fort Sumner in eastern New Mexico. By 1868 the Navajos were allowed to return to their lands

1
2

Density of Navajo Sites in McKinley and Cibola Counties, February 2008





Figure 3.5-19. Distribution of Navajo Archaeological Sites

1 within a newly designated reservation. The arrival of the railroad during the 1880s provided
2 them with a market for wool blankets and jewelry. However, this was a mixed blessing, with
3 pressures on the Navajo households to produce market items, versus subsistence self-
4 sufficiency. Ultimately, Navajos expanded into more marginal areas which could not sustain the
5 growing economic markets, with the long-term result being the partitioning of landholdings into
6 smaller family owned tracts, the overgrazing of these tracts and a shift towards wage earning
7 jobs (Kelley, 1986).

8 9 **3.5.8.2 Historic Properties Listed In The National And State Registers**

10
11 Table 3.5-12 includes a summary of sites in the Northwestern New Mexico Uranium Milling
12 Region that are listed on the New Mexico state and/or National Register of Historic Places.
13 Most of the sites are located in McKinley County, and the locations of many of the
14 archaeological sites are not identified to reduce the likelihood of vandalism. Historic sites are
15 located in the communities of Grants, Gallup, and Crownpoint, all of which are close to potential
16 uranium ISL milling locations.

17 18 **3.5.8.3 New Mexico Tribal Consultation**

19
20 There are 22 Native American Pueblos and Tribes located within the state of New Mexico. Most
21 of these groups are situated along the Rio Grande valley corridor from Albuquerque to Taos,
22 with several additional groups being represented in the northwest and southern parts of the
23 state. Five tribes have reservation lands within McKinley and Cibola counties, consisting of
24 Acoma Pueblo, Laguna Pueblo, Zuni Pueblo, the Navajo Nation and the Ramah Navajo Tribe.
25 These counties lie in the northwestern section of the state, along the southern periphery of the
26 San Juan Basin. The region is characterized by mesas and open grasslands which are
27 bounded by the Chuska Mountains, Zuni Mountains and Mount Taylor rising to heights of over
28 2,950 m [9,700 ft]. The Continental Divide bisects the area with drainages flowing towards the
29 north, west and east. Silko provides an insight into the Pueblo perspective of this environment
30 when she states that "there is no high mesa edge or mountain peak where one can stand and
31 not immediately be part of all that surrounds. Human identity is linked with all the elements of
32 Creation (Silko, 1990, pp. 884–885)."

33
34 Traditional Cultural Properties are places of special heritage value to contemporary
35 communities because of their association with cultural practices and beliefs that are rooted in
36 the histories of those communities and are important in maintaining the cultural identity of the
37 communities (Parker and King, 1998; King, 2003). Religious places are often associated with
38 prominent topographic features like mountains, peaks, mesas, springs and lakes (Silko, 1990).
39 In addition, shrines are present across the landscape to denote specific culturally significant
40 locations where an individual can place offerings (Ellis, 1974; Perlman, 1997; Rands, 1974a,b).
41 Ancestral villages also represent culturally significant places where the ancestors of these
42 contemporary communities once resided in the distant past, and are sometimes linked to
43 Pueblo migration stories (Ellis, 1974). In addition, specific resource collecting areas may have
44 significance for maintaining traditional lifeways (Ferguson and Hart, 1985; Perlman, 1997;
45 Rands 1974a,b). Lastly, pilgrimage trails with trail markers provide a link to all these areas
46 across the broad ethnic landscape (Ferguson and Hart, 1985; Fox, 1994; Parsons, 1918;
47 Sedgwick, 1926).

Table 3.5-12. National Register Listed Properties in Counties Included in the Northwestern New Mexico Uranium Milling Region

County	Resource Name	City	Date Listed YYYY-MM-DD
Cibola	Bowlin's Old Crater Trading Post	Bluewater	2006-03-21
Cibola	Candelaria Pueblo	Grants	1983-03-10
Cibola	Route 66 Rural Historic District: Laguna to McCarty's	Cubero	1994-01-13
Cibola	Route 66, State Maintained from McCarty's to Grants	Grants	1997-11-19
Cibola	Route 66, State maintained from Milan to Continental Divide	Continental Divide	1997-11-19
McKinley	Andrews Archeological District	Prewitt	1979-05-17
McKinley	Archeological Site # LA 15278 (Reservoir Site; CM 100)	Pueblo Pintado	1985-08-02
McKinley	Archeological Site # LA 45,780	Pueblo Pintado	1985-08-02
McKinley	Archeological Site # LA 45,781	Pueblo Pintado	1985-08-02
McKinley	Archeological Site # LA 45,782	Pueblo Pintado	1985-08-02
McKinley	Archeological Site # LA 45,784	Pueblo Pintado	1985-08-02
McKinley	Archeological Site # LA 45,785	Pueblo Pintado	1985-08-02
McKinley	Archeological Site # LA 45,786	Pueblo Pintado	1985-08-02
McKinley	Archeological Site # LA 45,789	Pueblo Pintado	1985-08-02
McKinley	Archeological Site # LA 50,000	Pueblo Pintado	1985-08-02
McKinley	Archeological Site # LA 50,001	Pueblo Pintado	1985-08-02
McKinley	Archeological Site # LA 50,013 (CM101)	Pueblo Pintado	1985-08-02
McKinley	Archeological Site # LA 50,014 (CM 102)	Pueblo Pintado	1985-08-02
McKinley	Archeological Site # LA 50,015 (CM 102A)	Pueblo Pintado	1985-08-02
McKinley	Archeological Site # LA 50,016 (CM 103)	Pueblo Pintado	1985-08-02
McKinley	Archeological Site # LA 50,017 (CM 104)	Pueblo Pintado	1985-08-02
McKinley	Archeological Site # LA 50,018	Pueblo Pintado	1985-08-02
McKinley	Archeological Site # LA 50,019 (CM 105)	Pueblo Pintado	1985-08-02
McKinley	Archeological Site # LA 50,020 (CM 106)	Pueblo Pintado	1985-08-02
McKinley	Archeological Site # LA 50,021	Pueblo Pintado	1985-08-02
McKinley	Archeological Site # LA 50,022 (CM 107)	Pueblo Pintado	1985-08-02
McKinley	Archeological Site # LA 50,023 (CM 118)	Pueblo Pintado	1985-08-02
McKinley	Archeological Site # LA 50,024 (CM 108)	Pueblo Pintado	1985-08-02
McKinley	Archeological Site # LA 50,025 (CM 109)	Pueblo Pintado	1985-08-02
McKinley	Archeological Site # LA 50,026 (CM 108)	Pueblo Pintado	1985-08-02
McKinley	Archeological Site # LA 50,027 (CM 111)	Pueblo Pintado	1985-08-02
McKinley	Archeological Site # LA 50,028 (CM 112)	Pueblo Pintado	1985-08-02
McKinley	Archeological Site # LA 50,030 (CM 114)	Pueblo Pintado	1985-08-02
McKinley	Archeological Site # LA 50,031 (CM 115)	Pueblo Pintado	1985-08-02
McKinley	Archeological Site # LA 50,033 (CM 117)	Pueblo Pintado	1985-08-02
McKinley	Archeological Site # LA 50,034	Pueblo Pintado	1985-08-02
McKinley	Archeological Site # LA 50,036	Pueblo Pintado	1985-08-02
McKinley	Archeological Site # LA 50,037	Pueblo Pintado	1985-08-02
McKinley	Archeological Site # LA 50,038	Pueblo Pintado	1985-08-02
McKinley	Archeological Site # LA 50,044	Pueblo Pintado	1985-08-02
McKinley	Archeological Site # LA 50,071 (CM 148)	Pueblo Pintado	1985-08-02
McKinley	Archeological Site # LA 50,072 (CM 94)	Pueblo Pintado	1985-08-02
McKinley	Archeological Site # LA 50,074 (CM 181)	Pueblo Pintado	1985-08-02
McKinley	Archeological Site # LA 50,077	Pueblo Pintado	1985-08-02
McKinley	Archeological Site # LA 50,080	Pueblo Pintado	1985-08-02
McKinley	Archeological Site No. LA 50,035	Pueblo Pintado	1985-10-09

Table 3.5-12. National Register Listed Properties in Counties Included in the Northwestern New Mexico Uranium Milling Region (continued)

County	Resource Name	City	Date Listed YYYY-MM-DD
McKinley	Ashcroft—Merrill Historic District	Ramah	1990-07-27
McKinley	Bee Burrow Archeological District	Seven Lakes	1984-12-10
McKinley	Casa de Estrella Archeological Site	Crownpoint	1980-10-10
McKinley	Chaco Culture National Historical Park	Thoreau	1966-10-15
McKinley	Chief Theater	Gallup	1988-05-16
McKinley	Cotton, C.N., Warehouse	Gallup	1988-01-14
McKinley	Cousins Bros. Trading Post	Chi Chil Tah	2006-03-22
McKinley	Dalton Pass Archeological Site	Crownpoint	1980-10-10
McKinley	Drake Hotel	Gallup	1988-01-14
McKinley	El Morro Theater	Gallup	1988-05-16
McKinley	El Rancho Hotel	Gallup	1988-01-14
McKinley	Fort Wingate Archeological Site	Fort Wingate	1980-10-10
McKinley	Fort Wingate Historic District	Fort Wingate	1978-05-26
McKinley	Grand Hotel	Gallup	1988-05-25
McKinley	Greenlee Archeological Site	Crownpoint	1980-10-10
McKinley	Halona Pueblo	Gallup	1975-02-10
McKinley	Harvey Hotel	Gallup	1988-05-25
McKinley	Haystack Archeological District	Crownpoint	1980-10-10
McKinley	Herman's, Roy T., Garage and Service Station	Thoreau	1993-11-22
McKinley	Lebanon Lodge No. 22	Gallup	1989-02-14
McKinley	Log Cabin Motel	Gallup	1993-11-22
McKinley	Manuelito Complex	Manuelito	1966-10-15
McKinley	McKinley County Courthouse	Gallup	1989-02-15
McKinley	Palace Hotel	Gallup	1988-05-16
McKinley	Peggy's Pueblo	Zuni	1994-08-16
McKinley	Redwood Lodge	Gallup	1998-02-13
McKinley	Rex Hotel	Gallup	1988-01-14
McKinley	Route 66, State maintained from Iyanbito to Rehobeth	Rehobeth	1997-11-19
McKinley	Southwestern Range and Sheep Breeding Laboratory Historic District	Fort Wingate	2003-05-30
McKinley	State Maintained Route 66—Manuelito to the Arizona Border	Mentmore	1993-11-22
McKinley	Upper Kin Klizhin Archeological Site	Crownpoint	1980-10-10
McKinley	US Post Office	Gallup	1988-05-25
McKinley	Vogt, Evon Zartman, Ranch House	Ramah	1993-02-04
McKinley	White Cafe	Gallup	1988-01-14

Of course the area of McKinley and Cibola counties only composes a small portion of the lands considered to be affiliated with traditional land-use activities. For example, the Navajo Nation bounds their traditional lands by the four culturally significant mountains: Hesperus Peak, Blanca Peak, Mount Taylor and the San Francisco Peaks which are located in Colorado, New Mexico and Arizona, respectively (Linford, 2000). Zuni Pueblo recognizes a shrine that is situated more than 240 km [150 mi] away at Bandelier National Monument near Los Alamos, New Mexico (Ferguson and Hart, 1985). On the other hand, Mount Taylor is significant to nearby Acoma and Laguna Pueblos for its role in their traditional origin myth where the Gambler held captive the Rainclouds until released by Sun Youth and Old Grandmother Spider (Sterling, 1942; Silko, 1990).

Information on traditional land-use and the location of culturally significant places is often protected information within the community (e.g., see King, 2003). Therefore, the information presented on religious places is limited to those that are identified in the published literature and are therefore restricted to a few highly recognized places on the landscape within McKinley and Cibola counties. Various documents pertaining to the Indian land claims also provide background information on local history and traditional land-use (Hawley Ellis, 1974; Minge, 1974; Rands, 1974a,b; Jenkins, 1974).

Linford's (2000) statement on the relation between mythology and place names is relevant to all traditional communities when he states that "a location's religious significance is more obscure, usually ascribed through its association with, or mention in, one or more of the stories that are the foundation of Navajo ceremonies" (ibid:17; also see Kelley and Francis, 1994; Holt 1981; Ortiz, 1992; Silko, 1990). The list of religious places provided in Table 3.5-13 is most often associated with traditional stories that recount the community's heritage through oral traditions. Ellis (1974) and Rand (1974a,b) do, however, provide a list of shrines that are associated with Laguna and Acoma Pueblos, and Ferguson and Hart (1985) of religious sites associated with Zuni Pueblo.

On February 22, 2008, the New Mexico Cultural Properties Review Committee accepted an emergency listing of the Mount Taylor Traditional Cultural Property to the State Register of Cultural Properties. The nomination was submitted by Acoma Pueblo, Hopi Tribe, Laguna Pueblo, the Navajo Nation and Zuni Pueblo. The boundaries of the Traditional Cultural Property have been tentatively set to include the summit and surrounding mesas above 2,440 m [8,000 ft], with the boundary dropping down to 2,224 m [7,300 ft] in the area of Horace Mesa. This application was specifically initiated to protect culturally sensitive sites that may be impacted by proposed uranium mining activities. The nominating group has 1 year to complete the final nomination to the state register; however, during this time the Traditional Cultural Property is given the full status of being listed.

The New Mexico Historic Preservation web site suggests that the following Pueblo and Tribal Groups should be contacted for consultation associated with activities in McKinley and Cibola counties: Acoma Pueblo, Hopi Tribe, Isleta Pueblo, Laguna Pueblo, Mescalero Apache Tribe, Navajo Nation, Sandia Pueblo, White Mountain Apache Tribe and Zuni Pueblo. This list was generated from the Pueblo and American land claims, Historic Preservation Division (HPD) ethnographic study, the National Park Service's Native American Consultation database and groups which directly contacted HPD requesting to be notified of potential activities in these areas. The Pueblo and Tribal contact information provided in Table 3.5-14 was obtained from the State of New Mexico, Indian Affairs Department web site: <<http://www.iad.state.nm.us/pueblogovandtribaloff.html>>.

3.5.8.4 Traditional Cultural Landscapes

Although archaeology and cultural resources management have historically focused on archaeological sites and artifact finds, past and present human interactions with their natural surroundings extend beyond the material traces of past human behavior. As a result, archaeologists and resource managers alike are increasingly focusing on the concept of traditional *cultural landscapes* as a broader, more accurate perspective on the way humans conceive of and use their environments. A cultural landscape is not the same as a natural

1

Table 3.5-13. Known Culturally Significant Places in McKinley and Cibola Counties		
Place	Affiliated Tribe	Reference
Bandera Crater	Zuni	Ferguson and Hart (p. 127)*
Cerro del Oro	Laguna	Parson,† Rands (p. 68)‡
Chuska Mountains (various locations)	Navajo	Linford (p. 194)§
Correo Snake Pit	Acoma and Laguna	Hawley Ellis (p. 92), Parsons,† Rands (p. 8)¶
Dowa Yalanne	Zuni	Ferguson and Hart (p. 124)*
El Malpais	Navajo	Linford (p. 204)§
El Morro	Zuni	Ferguson and Hart (p. 127)*
Hosta Butte	Navajo	Linford (p. 218)§
Ice Caves	Zuni	Ferguson and Hart (p. 125)*
Mount Taylor Shrines	Acoma Laguna Zuni	Parsons (p. 185);# Rands(p. 97),¶ Hawley-Ellis (p. 92), Ferguson and Hart (p. 126)*
Mount Taylor: Kaweshtima Tsiipiya T'se pina Tsoodzil Dewankwi Kyabachu Yalanne	Acoma Hopi Laguna Navajo Zuni	Application for Register. New Mexico State Register of Cultural Properties, February 22, 2008. New Mexico State Historic Preservation Office.
Pueblo Pintado	Navajo	Linford (p. 247)§
Red Lake	Navajo	Linford (p 250)§
Springs	Acoma Laguna Zuni	Rands (p. 97)¶, White (pp. 45–47),** Hawley-Ellis (p. 92), Ferguson and Hart (pp. 125–132)*
Zuni Salt Lake	Laguna Zuni Navajo	Rands (p. 68),‡ Ferguson and Hart (p. 126),* Linford (p. 284)§
Zuni Mountains (various locations)	Zuni	Ferguson and Hart (pp. 125, 132)*
<p>*Ferguson, T.J. and E. Hart. <i>A Zuni Atlas</i>. Norman, Oklahoma: University of Oklahoma Press. 1985. †Parsons, E.C. "War God Shrines of Laguna and Zuni." <i>American Anthropologist</i>. Vol. 20. pp. 381–405. 1918. ‡Rands, R. <i>Laguna Land Utilization: Pueblo Indians IV</i>. New York City, New York: Garland Publishing. 1974. §Linford, L. <i>Navajo Places: History, Legend and Landscape</i>. Salt Lake City, Utah: University of Utah Press. 2000. Hawley Ellis, F. <i>Archaeologic and Ethnologic Data: Acoma-Laguna Land Claims</i>. New York City, New York: Garland Publishing, Inc. 1974. ¶Rands, R. <i>Acoma Land Utilization: Pueblo Indians III</i>. New York City, New York: Garland Publishing. 1974. #Parsons, E.C. "Notes on Acoma and Laguna." <i>American Anthropologist</i>. pp. 162–186. 1918. **White, L.A. <i>The Acoma Indians</i>. Forty-Seventh Annual Report of the Bureau of American Ethnology to the Secretary of the Smithsonian Institution. Washington, DC: Smithsonian Institution. 1932.</p>		

2

3 "environment:" rather, it is produced by a cultural group's interaction with their environment. In
 4 simple terms, a cultural landscape is what results as members of a particular human group
 5 "project culture onto nature" (Crumley and Marquardt, 1990) by interacting with, modifying, and
 6 conceptualizing their natural surroundings over time (Ansuetz, et al., 2001).
 7 The notion of a cultural landscape includes the physical evidence of a group's interactions with
 8 the natural world, but is not limited to quantifiable material resources or patterns. A landscape
 9 perspective also incorporates the significance of particular places or landmarks for a group's

1

Table 3.5-14. 2008 Pueblo and Tribal Government Contacts for McKinley and Cibola Counties, New Mexico		
Affiliated Tribe	Contact	Address
Acoma Pueblo	Governor Chandler Sanchez	Pueblo of Acoma P.O. Box 309 Acoma, NM 87034 (505) 552-6604/6605
Acoma Pueblo	Teresa Pasqual, Director	Pueblo of Acoma Historic Preservation Office PO Box 309 Acoma, NM 87034 (505) 552-5170
Hopi Tribe	Chairman Benjamin Nuvamsa	Hopi Tribe P.O. Box 123 Kykotsmovi, AZ 86039 (928) 734-3000
Hopi Tribe	Leigh Kuwanwisiwma	Hopi Cultural Preservation Office The Hopi Tribe P.O. Box 123 Kykotsmovi, AZ 86039 (928) 734-6636 P (928) 734-3613 EX611 Lee (928) 734-3629 Fax
Jemez Pueblo	Governor Paul Chinana	Jemez Pueblo P.O. Box 100 Jemez Pueblo, NM 87024 (505) 834-7359
Jicarilla Apache Nation	President Levi Pesata	Jicarilla Apache Nation P.O. Box 507 Dulce, NM 507 (505) 759-3242
Isleta Pueblo	Governor Robert Benavides	Pueblo of Isleta P.O. Box 1270 Isleta Pueblo, NM 87022 (505) 869-3111/6333
Laguna Pueblo	Governor John Antonio, Sr.	Pueblo of Laguna P.O. Box 194 Laguna Pueblo, NM 87026 (505) 552-6654/6655/6598
Mescalero Apache Tribe	President Carleton Naiche- Palmer	Mescalero Apache Tribe P.O. Box 227 Mescalero, NM 88340 (505) 464-4494
Navajo Nation	President Joe Shirley, Jr.	Navajo Nation P.O. Box 9000 Window Rock, AZ 86515 (928) 871-6352/6357

1

Table 3.5-14. 2008 Pueblo and Tribal Government Contacts for McKinley and Cibola Counties (continued)		
Affiliated Tribe	Affiliated Tribe	Affiliated Tribe
Navajo Nation	Alan Downer	Tribal Preservation Officer Navajo Nation Historic Preservation Department P.O. Box 4950 Window Rock, AZ 86515 (928) 871-6437
Sandia Pueblo	Governor Robert Montoya	Pueblo of Sandia 481 Sandia Loop Bernalillo, NM 87004 (505) 867-3317
White Mountain Apache	Mr. Ramon Riley	White Mountain Apache Tribe P.O. Box 507 Fort Apache, AZ 85926
Zuni Pueblo	Governor Norman Cooney	Pueblo of Zuni P.O. Box 339 Zuni, NM 87327 (505) 782-7022
Zuni Pueblo	Jonathan Damp	Office of Heritage and Historic Preservation Pueblo of Zuni PO Box 339 Zuni, New Mexico 87327-0339 (928) 782-4814 P (928) 782-2393 F

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

histories, traditional stories, or religious beliefs (Ansuetz, 2007, Ansuetz, et al. 2001, Basso, 1996). Particular locations may serve as reminders of traditional beliefs or ways of life, or be venerated as supernatural beings in their own right. To quote a recent summary, a landscape perspective encompasses a "community's intimate relationships with the land and its resources in every aspect of its material life, including economy, society, polity, and recreation" (Ansuetz, 2007).

Understanding the importance of traditional cultural landscapes, then, means being aware of many overlapping dynamics of a culture's relationships with its environment. A landscape perspective must also take into account the overlapping, diverse cultural landscapes of many different cultures. In west-central New Mexico, for instance, a survey of cultural landscapes would include the distinct, extensive territories formerly used by the Zunis for economic activities ranging from farming and herding to gathering medicinal plants or collecting raw materials for stone tools (Ferguson and Hart, 1985). It would also recognize the culturally significant springs, caves and shrines dotting the world as conceived by the Keres people of Laguna and Acoma, or the culturally significant peaks at the four cardinal directions delineating this world's boundaries (Snead and Preucel 1999; White, 1932). Similar culturally significant landmarks recognized by the Navajo form part of yet another traditional landscape perspective, as described above. Finally, the roads and ruins of the ancient inhabitants of Chaco Canyon figure in the traditional histories of Zuni, Acoma, and Navajo alike, but also serve as clues to illuminate the traditional landscapes of the Chacoans themselves. Like their modern descendents, the ancient Chacoans seem to have placed importance on astronomical alignments, the cardinal directions, and prominent peaks, mesas and other landmarks (Van Dyke, 2004).

In summary, then, the distribution of archaeological sites, artifacts, and other physical markers of human activity are only one dimension of the processes in which past human groups used and conceptualized their surroundings. The traditional cultural landscapes of west-central New Mexico's indigenous groups include a wide variety of landmarks, traditional use areas, and other important features, many of which retain importance for contemporary groups. These traditional landscapes are increasingly recognized by agencies and archaeologists alike and play an expanding role in historic preservation and cultural resource management decision making.

3.5.9 Visual/Scenic Resources

Based on the BLM Visual Resource Handbook (BLM, 2007a–c), the Grants uranium district in the Northwestern New Mexico Uranium Milling Region is located in the Colorado Plateau physiographic province (BLM, 2007a). The Farmington and Albuquerque field offices of the BLM have classified most of the region as VRM Class III and IV (BLM, 2003, 2000). There are no VRM Class I VRM areas, and most of the Class II regions are located just north of Interstate 40. As described in NRC (1997), the primary viewers in the San Juan Basin and Grants Uranium Districts are likely to be Native American residents living on and near a proposed ISL facility (see Section 3.5.8). For this reason, their aesthetic sense at the landscape scale is important. In general, Native American thought is “integrative and comprehensive. It does not separate intellectual, moral, emotional, aesthetic, economic, and other activities, motivations, and functions” (Norwood and Monk, 1987). For both the Navajo and Zuni, moral good tends to be equated with aesthetic good: that which promotes or represents human survival and human happiness tends to be experienced as “beautiful.” The landscape is beautiful by definition because the Holy People designed it to be a beautiful, harmonious, happy, and healthy place (Norwood and Monk, 1987). Native Americans have not created an abstract category for unspecified vistas; the emphasis is on specific mountains, specific trees, and specific colors of the soil (Norwood and Monk 1987). References to the visual quality of a given area may be more meaningful when linked to an identifiable place and not to more generalized landscapes.

Natural and scenic attractions within the Grants uranium district in the Northwestern New Mexico Uranium Milling Region are minimal. Regionally, the Chaco Culture National Historic Park, El Malpais National Monument (BLM, 2000), El Morro National Monument, and the Red Rock State Park, among other features, attract tourists for scenic, historic, and cultural features (see Section 3.5.1). Near Gallup and south of Interstate 40, the USFS categorizes the visual quality objectives within the Cibola National Forest as predominantly (about 75 percent) in the Modification and Maximum Modification class (USFS, 1985), with some areas such as the Mt. Taylor district in the San Mateo Mountains having high scenic integrity (USFS, 2007). In addition, in February 2008, the New Mexico Cultural Properties Review Committee approved listing the Mount Taylor Traditional Cultural Property in the State Register of Cultural Properties (see Section 3.5.8.3). With the exception of major highways such as Interstate 40 and U.S. Highway 491, area roads are used mostly for local travel. The urban areas such as Gallup, Crownpoint, and Grants tend to dominate visual resources near these cities and towns (NRC, 1997).

The resource management plan for the Farmington field office of the BLM provides a VRM classification for the public lands in the Northwestern New Mexico Uranium Milling Region (BLM, 2003) (Figure 3.5-20). The visual context is also an important component of the cultural resource values of the Chacoan Outliers, Native American Use and Sacred Areas of Critical Environmental Concern, and additional traditional cultural properties (BLM, 2003). The approximately 2 million ha [5 million acres] of regional public lands and subsurface mineral resources BLM administers in the Farmington field office have a relatively small amount (about




Figure 3.5-20. BLM Visual Resource Classifications for the Northwestern New Mexico Uranium Milling Region (BLM, 2003, 2000)

13 percent) of VRM Classes I and II viewsheds associated with wilderness areas, wilderness study areas, specially designated areas, and special management areas. As categorized by BLM, the visual landscape in northwestern New Mexico is dominated by VRM Class IV (55 percent) and Class III (32 percent). The natural state has been considerably modified by human activities and structures associated with oil and gas development, including gas wells, pipelines, and the accompanying access roads. There are no Class I areas within the Northwestern New Mexico Uranium Milling Region. Areas categorized as Class II include locations where scenic vistas (from major highways), riverfronts, and high places are important because of associated sightseeing and recreational value (BLM, 2003).

Specific VRM Class II locations identified by BLM within and near the region include the Cabezon Peak, Cañon Jarido, Elk Springs, Ignacio Chavez, Jones Canyon, and La Lena special management areas and the Empedrado wilderness study areas (BLM 2003) at the eastern edge of the Northwestern New Mexico Uranium Milling Region. The USFS also identifies Corral Canyon and the western edge of the San Pedro Mountains in the La Jara area of the Santa Fe National Forest just to the east of the Northwestern New Mexico Uranium Milling Region as areas where recreation and timber are to be managed to preserve visual resource value (USFS, 2007). These Class II resource areas are adjacent to the Grants uranium district, but the closest potential uranium ISL facility to these resource areas is about 16 km [10 mi]. There are some Class II viewsheds associated with the Chaco Culture National Historic Park just to the north that extend into the region about 50 km [30 mi] north of the nearest potential uranium recovery facility (Figure 3.5-20). BLM National Conservation Areas, adjacent to the El Malpais National Monument and about 3 km [2 mi] south of Grants, are also identified as Class II. Two potential facilities are located near San Mateo Mesa about 16 km [10 mi] northwest of Mt. Taylor. In addition, two of the proposed facilities are located within about 3-8 km [2-5 mi] of the borders of the Navajo Nation (Figure 3.5-20). Current indications from industry are that these would be developed as conventional milling operations (NRC, 2008).

3.5.10 Socioeconomics

For the purpose of this GEIS, the socioeconomic description for the Northwestern New Mexico Uranium Milling Region includes communities within the region of influence for potential ISL facilities in the Grants Uranium District. These include communities that have the highest potential for socioeconomic impacts and are considered the affected environment. Communities that have the highest potential for socioeconomic impacts are defined by (1) proximity to an ISL facility (generally within about 48 km [30 mi]), (2) economic profile, such as potential for income growth or de-stabilization, (3) employment structure, such as potential for job placement or displacement and (4) community profile, such as potential for growth or destabilization to local emergency services, schools, or public housing. The affected environment consists of counties, towns, Core-Based Statistical Areas, and Native American communities (reservation land) (Table 3.5-15). A Core-Based Statistical Areas, according to the U.S. Census Bureau, is a collective term for both metro and micro areas ranging from a population of 10,000 to 50,000 (U.S. Census Bureau, 2007). The following sub-sections describe areas most likely to have implications with regard to socioeconomics. In some sub-sections Metropolitan Areas are also discussed. A Metropolitan Area is greater than 50,000 and a town is considered less than 10,000 in population (U.S. Census Bureau, 2007).

1

**Table 3.5-15. Summary of Affected Environment Within the
Northwestern New Mexico Uranium Milling Region**

Table 3.5-15. Summary of Affected Environment Within the Northwestern New Mexico Uranium Milling Region			
Counties Within New Mexico	Towns Within New Mexico	CBSAs Within New Mexico	Native American Communities Within New Mexico
Cibola	Grants	Gallup	Acoma Indian Reservation
McKinley			Tohajiilee Indian Reservation
Sandoval			Laguna Indian Reservation
			Navajo Nation Indian Reservation
			Ramah Navajo Indian Reservation
		Zuni Indian Reservation	

2

3

3.5.10.1 Demographics

5

Demographics are based on 2000 Census data on population and racial characteristics of the affected environment (Table 3.5-16). Figure 3.5-21 illustrates the populations of communities within the Northwestern New Mexico Uranium Milling Region. Most 2006 data compiled by the U.S. Census Bureau is not yet available for the geographic area of interest.

10

Based on review of Table 3.5-16, the most populated county is Sandoval County and the most sparsely populated county is Cibola County. The largest populated town/Core-Based Statistical Areas in the Northwestern New Mexico Uranium Milling Region is Gallup. The county with the largest percentage of non-minorities is Sandoval County with a white population of 65.1 percent. The town/Core-Based Statistical Areas with the largest percentage of non-minorities is Grants with a white population of 56.2 percent. The largest minority-based county is McKinley County with a white population of only 16.4 percent. The largest minority-based town is Gallup with a white population of 40.1 percent.

19

Although not listed in Table 3.5-16, total population counts based on 2000 U.S. Census Bureau data (U.S. Census Bureau, 2008) for the Native American communities (reservation land) that would be affected are

23

- Acoma Indian Reservation: 2,802
- Tohajiilee Indian Reservation: 1,649
- Laguna Indian Reservation: not available
- Navajo Nation Indian Reservation: 173,987*
- Ramah Navajo Indian Reservation: 2,167
- Zuni Indian Reservation: 7,758

30

*Includes Arizona, Utah, and New Mexico (131,166 were reported as living in Arizona).


31

**Table 3.5-16. 2000 U.S. Bureau of Census Population and Race Categories of the
Northwestern New Mexico Uranium Milling Region***

Affected Environment	Total Population	White	African American	Native American	Some Other Race	Two or More Races	Asian	Hispanic Origin†	Native Hawaiian and Other Pacific Islander
New Mexico	1,819,046	1,214,253	34,343	173,483	309,882	66,327	19,255	765,386	1,503
<i>Percent of total</i>		66.8%	1.9%	9.5%	3.6%	3.6%	1.1%	42.1%	0.1%
Cibola County	25,595	10,138	246	10,319	3,952	828	98	8,555	14
<i>Percent of total</i>		39.6%	1.0%	40.3%	15.4%	3.2%	0.4%	33.4%	40.3%
McKinley County	74,798	12,257	296	55,892	4,095	1,882	344	9,276	32
<i>Percent of total</i>		16.4%	0.4%	74.7%	5.5%	2.5%	0.5%	12.4%	0.0%
Sandoval County	89,908	58,512	1,535	14,634	11,118	3,117	894	26,437	98
<i>Percent of total</i>		65.1%	1.7%	16.3%	12.4%	3.5%	1.0%	29.4%	0.1%
Gallup	20,274	8,106	219	7,404	2,985	1,187	289	6,699	19
<i>Percent of total</i>		40.1%	1.1%	36.6%	14.8%	5.9%	1.4%	33.1%	0.1%
Grants	8,806	4,947	143	1,054	2,184	386	81	4,611	11
<i>Percent of total</i>		56.2%	1.6%	12.0%	24.8%	4.4%	0.9%	52.4%	0.1%

* U.S. Census Bureau. "American FactFinder." <http://factfinder.census.gov/home/saff/main.html?_lang=en> (18 October 2007 and 25 February 2008).

†Hispanic origin can be any race and is calculated as a separate component of the total population (i.e., if added to the other races would total more than 100 percent).



NEW MEXICO REGION



Figure 3.5-21. Northwestern New Mexico Uranium Milling Region With Population

3.5.10.2 Income

Income information from 2000 Census data including labor force, income, and poverty levels for the affected environment is collected at the state and county levels. Data collected from a state level also includes information on towns, Core-Based Statistical Areas, or Metropolitan Areas and was done to take into consideration an outside workforce. An outside workforce may be a workforce willing to commute long distances {greater than 48 km [30 mi]} for income opportunities or may be a workforce necessary to fulfill specialized positions (if local workforce is unavailable or un-specialized). Data collected from a county level is generally the same affected environment discussed previously in Table 3.5-15 and also includes information on Native American communities in the Northwestern New Mexico Uranium Milling Region. State level information is provided in Table 3.5-17 and county data is listed in Table 3.5-18.

For the region surrounding the Northwestern New Mexico Uranium Milling Region, the state with the largest labor force population is Arizona. The community with the largest labor force is Albuquerque, New Mexico {144 km [90 mi] from the nearest potential ISL facility} and the smallest community labor force is Grants, New Mexico {8 km [5 mi] from the nearest potential ISL facility}. The community with the highest per capita income is Santa Fe, New Mexico {96 km [60 mi] from the nearest potential ISL facility} and the lowest per capita income population is Silver City, New Mexico {161 km [100 mi] from the nearest potential ISL facility}. Outside of tribal lands, the community with the highest percentage of individuals and families below poverty levels is Grants, New Mexico.

The county with the largest labor force population in the Northwestern New Mexico Uranium Milling Region is Sandoval County and the county with the smallest labor force population is Cibola County. The county with the highest per capita income is Sandoval County and the lowest per capita income county is McKinley County. The county with the highest percentage of individuals and families below the poverty level is McKinley County (Table 3.5-18).

3.5.10.3 Housing

Housing information from the 2000 Census data is provided in Table 3.5-19.

The availability of housing within the immediate vicinity of the proposed ISL facilities is somewhat limited. The majority of housing is available in larger populated areas such as Gallup {24 km [15 mi] to the nearest potential ISL facility}, Grants {8 km [5 mi] to nearest potential ISL facility}, Albuquerque {144 km [90 mi] to the nearest potential ISL facility}, and Rio Rancho {161 km [100 mi] to the nearest potential ISL facility}. There are approximately 20 housing units, including manufactured housing parks or residential neighborhoods in this region (MapQuest, 2008d).

Temporary housing such as apartments, lodging, and trailer camps within the immediate vicinity of the Grants Uranium District ISL facilities is not as limited. The majority of apartments are available in larger populated areas such as the Gallup, Grants, Belen, Los Lunas, and Albuquerque with approximately 75 apartment complexes (MapQuest, 2008). There are 19 hotels/motels along major highways or towns near the ISL facilities. In addition to apartments and lodging, there are three trailer camps also located near potential ISL facilities (along major roads or near towns) (MapQuest, 2008).

Table 3.5-17. U.S. Bureau of Census State Income Information for the Northwestern New Mexico Uranium Milling Region*

Affected Environment	2000 Labor Force Population (16 years and over)	Median Household Income In 1999	Median Family Income In 1999	Per Capita Income In 1999	Families Below Poverty Level In 2000	Individuals Below Poverty Level In 2000
Arizona	2,387,139	\$40,558	\$46,723	\$20,275	128,318	698,669
New Mexico	834,632	\$34,133	\$39,425	\$17,261	68,178	328,933
Albuquerque, New Mexico	232,320	\$38,272	\$46,979	\$20,884	11,285	59,641
<i>Percent of total</i>	66.2%	NA	NA	NA	10.0%	13.5%
Farmington, New Mexico	18,204	\$37,663	\$42,605	\$18,167	1,328	5,910
<i>Percent of total</i>	65.0%	NA	NA	NA	12.9%	16.0%
Flagstaff, Arizona	30,822	\$37,146	\$48,427	\$18,637	1,255	8,751
<i>Percent of total</i>	73.7%	NA	NA	NA	10.6%	17.4%
Gallup, New Mexico	8,941	\$34,868	\$39,197	\$15,789	804	4,079
<i>Percent of total</i>	61.9%	NA	NA	NA	16.6%	20.8%
Grants, New Mexico	3,801	\$30,652	\$33,464	\$14,053	446	1,810
<i>Percent of total</i>	58.3%	NA	NA	NA	19.4%	21.9%
Rio Rancho, New Mexico	25,964	\$47,169	\$52,233	\$20,322	521	2,619
<i>Percent of total</i>	67.9%	NA	NA	NA	3.7%	5.1%

Table 3.5-17. U.S. Bureau of Census State Income Information for the Northwestern New Mexico Uranium Milling Region* (continued)

Affected Environment	2000 Labor Force Population (16 years and over)	Median Household Income In 1999	Median Family Income In 1999	Per Capita Income In 1999	Families Below Poverty Level In 2000	Individuals Below Poverty Level In 2000
Santa Fe, New Mexico	34,033	\$40,392	\$49,705	\$25,454	1,425	7,439
<i>Percent of total</i>	66.8%	NA	NA	NA	9.5%	12.3%
Silver City, New Mexico	4,249	\$25,881	\$31,374	\$13,813	483	2,237
<i>Percent of total</i>	52.5%	NA	NA	NA	17.7%	21.9%

*Source: U.S. Census Bureau. "American FactFinder." <http://factfinder.census.gov/home/saff/main.html?_lang=en> (18 October 2007, 25 February 2008, and 15 April 2008).
†Percent of total based on a population of 16 years and over.
‡NA—not applicable.

Table 3.5-18. U.S. Bureau of Census County Income Information for the Northwestern New Mexico Uranium Milling Region*

Affected Environment	2000 Labor Force Population (16 years and over)	Median Household Income In 1999	Median Family Income In 1999	Per Capita Income In 1999	Families Below Poverty Level In 2000	Individuals Below Poverty Level In 2000
Cibola County, New Mexico	9,848	\$27,774	\$30,714	\$11,731	1,365	6,054
<i>Percent of total</i>	53.0%	NA	NA	NA	21.5%	24.8%
McKinley County, New Mexico	26,498	\$25,005	\$26,806	\$9,872	5,303	26,664
<i>Percent of total</i>	53.4%	NA	NA	NA	31.9%	36.1%
Sandoval County, New Mexico	41,599	\$44,949	\$48,984	\$19,174	2,130	10,847
<i>Percent of total</i>	63.0%	NA	NA	NA	9.0%	12.1%

*Source: U.S. Census Bureau. "American FactFinder." <http://factfinder.census.gov/home/saff/main.html?_lang=en> (18 October 2007 and 25 February 2008).

†Percent of total based on a population of 16 years and over.

‡NA—not applicable.

Affected Environment	Single Family Owner-Occupied Homes	Median Value in Dollars	Median Monthly Costs With a Mortgage	Median Monthly Costs Without a Mortgage	Occupied Housing Units	Renter-Occupied Units
New Mexico	339,888	\$108,100	\$929	\$228	677,971	200,908
Cibola County	3,742	\$62,600	\$654	\$179	8,327	1,873
McKinley County	10,235	\$57,000	\$841	\$140	21,476	5,840
Sandoval County	21,873	\$115,400	\$979	\$233	31,411	5,097
Gallup	2,922	\$97,000	\$933	\$4,245	6,807	2,682
Grants	1,634	\$64,700	\$697	\$210	3,160	1,024

* U.S. Census Bureau. "American FactFinder." <http://factfinder.census.gov/home/saff/main.html?_lang=en> (18 October 2007 and 25 February 2008).

* U.S. Census Bureau. "American FactFinder." <http://factfinder.census.gov/home/saff/main.html?_lang=en> (18 October 2007 and 25 February 2008).

3.5.10.4 Employment Structure

Employment structure from the 2000 Census data including employment rate and type, is based on data collected at the state and county levels. Data collected at the state level also includes information on towns, Core-Based Statistical Areas, or Metropolitan Areas and was done to take into consideration an outside workforce. An outside workforce may be a workforce willing to commute long distances {greater than [48 km [30 mi]]} for employment opportunities or may be a workforce necessary to fulfill specialized positions (if local workforce is unavailable or unspecialized). Data collected from a county level is generally the same affected environment previously discussed in Table 3.5-15 and also includes information on Native American communities.

Based on review of state information, the state in the vicinity of the Northwestern New Mexico Uranium Milling Region with the highest percentage of employment is Arizona.

At the the county with the highest percentage of employment is Sandoval County and the county with the highest unemployment rate is McKinley County. Native American communities (Navajo Nation, Zuni, and Laguna Reservations) report unemployment rates of 60 percent or more, much greater than the state unemployment levels of 3.4 percent (Arizona) to 4.4 percent (New Mexico) Table 3.5-20).

3.5.10.4.1 State Data

3.5.10.4.1.1 Arizona

The State of Arizona has an employment rate of 57.2 percent and unemployment rate of 3.4 percent. The largest sector of employment is management, professional, and related occupations. The largest type of industry is educational, health, and social services. The largest class of worker is private wage and salary workers (U.S. Census Bureau, 2008).

Flagstaff

Flagstaff has an employment rate of 69.8 percent and an unemployment rate slightly higher than that of the state at 3.9 percent. The largest sector of employment is management, professional, and related occupations at 30.2 percent. The largest type of industry is educational, health, and social services. The largest class of worker is private wage and salary workers (U.S. Census Bureau, 2008).

3.5.10.4.1.2 New Mexico

The State of New Mexico has an employment rate of 55.7 percent and unemployment rate of 4.4 percent. The largest sector of employment is management, professional, and related occupations. The largest type of industry is educational, health, and social services. The largest class of worker is private wage and salary workers (U.S. Census Bureau, 2007).

Albuquerque

Albuquerque has an employment rate of 61.8 percent and an unemployment rate lower than that of the state at 3.8 percent. The largest sector of employment is management, professional, and related occupations at 38.5 percent. The largest type of industry is educational, health, and

social services. The largest class of worker is private wage and salary workers (U.S. Census Bureau, 2008).

Gallup

Gallup has an employment rate of 57.1 percent and an unemployment rate slightly higher than that of the state at 4.8 percent. The largest sector of employment is management, professional, and related occupations at 38.9 percent. The largest type of industry is educational, health, and social services at 31.5 percent. The largest class of worker is private wage and salary workers at 65.2 percent (U.S. Census Bureau, 2007).

Grants

Grants has an employment rate of 51.9 percent and an unemployment rate higher than that of the state at 6.2 percent. The largest sector of employment is management, professional, and related occupations at 30.0 percent. The largest type of industry is educational, health, and social services at 23.6 percent. The largest class of worker is private wage and salary workers at 61.3 percent (U.S. Census Bureau, 2008).

Farmington

Farmington has an employment rate of 60.4 percent and an unemployment rate slightly higher than that of the state at 4.5 percent. The largest sector of employment is management, professional, and related occupations at 30.2 percent. The largest type of industry is educational, health, and social services. The largest class of worker is private wage and salary workers (U.S. Census Bureau, 2008).

Rio Rancho

Rio Rancho has an employment rate of 64.3 percent and an unemployment rate slightly higher than that of the state at 3.2 percent. The largest sector of employment is management, professional, and related occupations at 34.5 percent. The largest type of industry is educational, health, and social services. The largest class of worker is private wage and salary workers (U.S. Census Bureau, 2008).

Santa Fe

Santa Fe has an employment rate of 63.7 percent and an unemployment rate much lower than that of the state at 3.0 percent. The largest sector of employment is management, professional, and related occupations at 43.0 percent. The largest type of industry is educational, health, and social services. The largest class of worker is private wage and salary workers (U.S. Census Bureau, 2008).

3.5.10.4.2 County Data

Cibola County, New Mexico

Cibola County has an employment rate of 46.8 percent and an unemployment rate relatively higher than that of the state at 6.1 percent. The largest sector of employment is management, professional, and related occupations at 29.6 percent. The largest type of industry is

Description of the Affected Environment

educational, health, and social services at 27.4 percent. The largest class of worker is private wage and salary workers at 58.4 percent (U.S. Census Bureau, 2007).

McKinley County, New Mexico

McKinley County has an employment rate of 44.2 percent and an unemployment rate relatively higher than that of the state at 9.2 percent. The largest sector of employment is management, professional, and related occupations at 32.4 percent. The largest type of industry is educational, health, and social services at 32.4 percent. The largest class of worker is private wage and salary workers at 55.9 percent (U.S. Census Bureau, 2007).

Sandoval County, New Mexico

Sandoval County has an employment rate of 58.8 percent and an unemployment rate lower than that of the state at 3.9 percent. The largest sector of employment is management, professional, and related occupations at 36.0 percent. The largest type of industry is educational, health, and social services at 17.4 percent. The largest class of worker is private wage and salary workers at 73.6 percent (U.S. Census Bureau, 2007).

Native American Communities

Information on labor force and poverty levels for the affected Native American communities within Northwestern New Mexico is based on 2003 Bureau of Indian Affairs data and is provided below in Table 3.5-20 (U.S. Department of the Interior, 2003).

3.5.10.5 Local Finance

Local finance such as revenue and tax information for the affected environment is provided below and in Tables 3.5-21 to 3.5-23.

Table 3.5-20. Employment Structure of Native American Communities Within the Affected Environment of the Northwestern New Mexico Uranium Milling Region*

Affected Areas	2003 Labor Force Population	Unemployed as Percent of Labor Force	Employed Below Poverty Guidelines	
Acoma Indian Reservation	NR†	NR	NR	NR
Canoncito Indian Reservation	NA‡	NA	NA	NA
Laguna Indian Reservation	828	81%	NR	NR
Navajo Nation Indian Reservation (Eastern Navajo Agency)	2,664	74%	62	2%
Ramah Navajo Indian Reservation	NR	NR	NR	NR
Zuni Indian Reservation	1,591	64%	110	7%

* U.S. Department of the Interior. "Affairs American Indian Population and Labor Force Report 2003." <<http://www.doi.gov/bia/labor.html>>. Washington, DC: U.S. Department of the Interior, Bureau of Indian Affairs, Office of Tribal Affairs. 2003.
†NR—Not reported by tribes.
‡NA—not available.

Table 3.5-21. Net Taxable Values for Affected Counties Within New Mexico for 2006*

Affected Counties	Residential	Nonresidential	Total
Cibola County	\$88,563,082	\$145,457,203	\$234,020,285
McKinley County	\$219,073,850	\$410,061,159	\$629,311,981
Sandoval County	\$1,631,727,293	\$449,148,142	\$6,755,265

*Source: New Mexico Taxation and Revenue Department. "2006 Property Tax Facts."
<<http://www.tax.state.nm.us/pubs/taxresstat.htm>>. Santa Fe, New Mexico: New Mexico Taxation and Revenue Department (18 October 2007 and 25 February 2008).

Table 3.5-22. Percent Change in Tax Values From 2005 to 2006 for the Affected Counties Within New Mexico*

Affected Counties	Residential	Nonresidential	Total
Cibola County	3.0 percent	3.6 percent	3.4 percent
McKinley County	4.1 percent	4.0 percent	4.0 percent
Sandoval County	18.8 percent	8.7 percent	16.5 percent

*New Mexico Taxation and Revenue Department. "2006 Property Tax Facts."
<<http://www.tax.state.nm.us/pubs/taxresstat.htm>>. Santa Fe, New Mexico: New Mexico Taxation and Revenue Department (18 October 2007 and 25 February 2008).

Table 3.5-23. Percent Distribution of New Mexico Property Tax Obligations Within Affected Counties for 2006*

Affected Counties	State	County	Municipal	School District	Other
Cibola County	4.4 percent	34.4 percent	9.8 percent	34.4 percent	17 percent
McKinley County	3.9 percent	32.3 percent	10.9 percent	31.6 percent	21.1 percent
Sandoval County	4.8 percent	26.6 percent	19.7 percent	39.7 percent	9.1 percent

* New Mexico Taxation and Revenue Department. "2006 Property Tax Facts." <<http://www.tax.state.nm.us/pubs/taxresstat.htm>>. Santa Fe, New Mexico: New Mexico Taxation and Revenue Department (18 October 2007 and 25 February 2008).

New Mexico

Sources of revenue for the State of New Mexico come from income, mineral extraction, and property taxes. Personal income tax rates for New Mexico range from 1.7 percent to 5.3 percent. New Mexico does not have a sales tax and instead has a 5 percent gross receipts tax. Combined gross receipts tax rates throughout the state range from 5.125 to 7.8125 percent. Net taxable values for affected counties in New Mexico are presented in Table 3.5-21 (New Mexico Taxation and Revenue Department, 2008).

Description of the Affected Environment

Percentages and sources of revenue for 2006 were counties at 32.3 percent, municipalities at 14.3 percent, school districts at 30.0 percent, conservancy districts at 0.1 percent, state debt service at 4.8 percent, health facilities at 8.8 percent, and higher education at 9.7 percent. Total tax values for the affected counties within New Mexico are listed below. Percent change in net taxable values from 2005 to 2006 for the affected counties is provided in Table 3.5-22 (New Mexico Taxation and Revenue Department, 2008).

New Mexico imposes "ad valorem production" and "ad valorem production equipment" taxes in lieu of property taxes on mineral extraction properties. Taxes are levied monthly on all owners and are imposed on products below the wellhead, such as oil and gas. Equipment is also levied against the operator of the property. In 2000, ad valorem production and production equipment taxes totaled approximately \$43.4 million in taxes. Of this total, 83 percent came from the oil and gas production tax. How revenues are distributed in a particular county is determined by property tax rates imposed at the county

Percent distribution of New Mexico property tax obligations for 2006 within the affected counties is listed in Table 3.5-23. Information on local finance for the Core-Based Statistical Areas of Gallup and town of Grants is presented below.

Gallup

Sources of revenue for Gallup consist of gross receipts taxes, compensating taxes, corporate income taxes, franchise taxes, property taxes, severance taxes, and workers' compensation taxes. The largest tax revenues are gross receipts at a rate of 7.6 percent and property tax ranging from 4.7 percent to 7.4 percent. Revenue from gross receipts totaled \$115,031,909 as of 2004 (City of Gallup Economic Development Center, 2007).

Grants

Sources of revenue for Grants consist of gross receipts taxes and property taxes (New Mexico Economic Development, 2008).

Native American Communities

The Acoma Indian Reservation's largest sources of revenue come from the Sky City Casino and big game hunting. Specific financial information including tax revenue is not available (Acoma New Mexico, 2007).

The Tohajiilee Indian Reservation receives revenue from local retail and gaming. Specific financial information including tax revenue is not available (Division of Economic Development of the Navajo Nation, 2006).

The Laguna Indian Reservation receives revenue from local retail and gaming. Specific financial information including tax revenue is not available (New Mexico Tourism Department, 2008).

The largest source of revenue for the Navajo Nation Indian Reservation comes from internal and external revenue. Internal revenue is referred to as General Fund revenues and consists of mining and taxes. Mining is the largest source of internal revenue. Taxes are the second largest sources of internal revenue and in 2005 accounted for \$75.0 million (Division of Economic Development of the Navajo Nation, 2006). Taxes include business gross receipts.

This tax could be levied on uranium production within the Navajo Reservation if production is determined to occur on the reservation (NRC, 1997). External sources of revenue consist of Federal, State, Private and other funds, and are mostly in the form of grants (Division of Economic Development of the Navajo Nation, 2006).

The Ramah Navajo Indian Reservation is one of 110 chapters that make up the larger Navajo Nation. The Ramah Navajo take no assistance from the Navajo Nation. The majority of revenue comes from federal funding because this group does not have a single, sustainable economic development program that generates significant income (Ramah Navajo Chapter, 2003).

The majority of revenue for the Zuni Indian Reservation comes from federal grants, such as the Community Services Block Grant. Other sources of income include local taxes such as sales tax from gross receipts (Pueblo of Zuni, 2008).

3.5.10.6 Education

Based on review of the affected environment, the county with the largest number of schools is McKinley County and the county with the smallest number of schools is Cibola County. The town/Core-Based Statistical Areas with the largest number of schools is Gallup and the town/Core-Based Statistical Areas with the smallest number of schools is Grants. The Native American community with the largest number of schools is the Navajo Nation and the Native American community with the smallest number of schools is the Tohajiilee Indian Reservation.

Grants

Grants has 2 elementary schools, 1 middle school, 1 high school, 3 private academies, and 1 public school, with a total of approximately 2,414 students (Localschooldirectory.com, 2008).

Gallup

Gallup has 33 public schools and 2 parochial schools, with a total of approximately 8,013 students. (City of Gallup Economic Development Center, 2007).

Cibola County

Public education in Cibola County is operated by Grants/Cibola County Schools, which is based in Grants, New Mexico. There are 7 elementary schools, 1 middle school, 1 middle-high school, and 1 high school, with a total of approximately 3,698 students. The majority of schools provide bus services (Grants-Cibola County Schools, 2007)).

McKinley County

Public education in McKinley County education system is operated by the Gallup-McKinley County school district, which serves students from Gallup and surrounding areas of McKinley County. There are 36 public and private elementary, middle, and high schools within the county, with a total of approximately 13,840 students. The majority of schools provide bus services (Greatschools, 2007c).

Description of the Affected Environment

Sandoval County

Sandoval County has a total of 11 elementary schools, 6 middle schools, and 5 high schools, with a total of approximately 8,580 students. The majority of schools provide bus services (Publicschoolreview.com, 2008).

Native American Communities

The Acoma Indian Reservation has the Sky City Community School located at Acoma Pueblo. The total number of students is approximately 275. Information as to whether this school provide bus services is not available (Public Schools Report, 2007).

The Tohajiilee Indian Reservation has one school that is located within the Tohajiilee Indian Reservation. Specific information pertaining to school population or bus services is not available (Tohajiilee Chapter, 2008).

The Laguna Indian Reservation has 1 elementary school, 1 middle school, 1 high school, and 1 academy. Specific information pertaining to school population or bus services is not available (Lat-Long.com, 2008).

The Navajo Nation Indian Reservation has over 150 public, private and Bureau of Indian Affairs schools serving students from kindergarten through high school. There are over 10,000 students. Information as to whether these schools provide bus services is not available (Division of Economic Development of the Navajo Nation, 2008)).

The Ramah Navajo Indian Reservation school system is operated by the Ramah Navajo School Board and the Ramah Navajo Chapter. It has an Indian-controlled contract school located in Pine Hill, New Mexico. It accommodates almost 600 students from elementary through 12th grade. Information as to whether this school provides bus services is not available (Ramah Navajo Chapter, 2003).

The Zuni Indian Reservation has 2 elementary schools, 1 middle school, and 2 high schools, with a total of approximately 2,000 students. Information as to whether these schools provide bus services is not available (Zuni Pueblo Public School District, 2008).

3.5.10.7 Health and Social Services

Health Care Facilities

The majority of health care facilities are located within populated areas of the affected environment. The closest health care facilities within the vicinity of the ISL facilities are located in Gallup, Zuni, Rio Rancho, and Albuquerque and total approximately 50 facilities (MapQuest, 2008). These consist of hospitals, clinics, emergency centers, and medical services. There are 13 hospitals located within or proximate of this region: Gallup (1), Zuni (1), Rio Rancho (1), and Albuquerque (greater than 10).

Local Emergency

Local police within the affected environment is within the jurisdiction of each county. There are 12 police, sheriff, or marshal's offices within the region: Cibola County (3), McKinley County (3), and Sandoval County (6) (usacops, 2008).

Fire departments within the affected area are comprised at the town, CBSA, or city level. There are 24 fire departments within the milling region: Grants (4), Gallup (13), and Albuquerque (7) (50states, 2008d).

3.5.11 Public and Occupational Health

3.5.11.1 Background Radiological Conditions

For a U.S. resident, the average total effective dose equivalent from natural background radiation sources is approximately 3 mSv/yr [300 mrem/yr] but varies by location and elevation (National Council of Radiation Protection and Measurements, 1987). In addition, the average American receives 0.6 mSv/yr [60 mrem/yr] from man-made sources including medical diagnostic tests and consumer products (National Council of Radiation Protection and Measurements 1987). Therefore the total from natural background and man-made sources for the average U.S. resident is 3.6 mSv/yr [360 mrem/yr]. For a breakdown of the sources of this radiation, see Figure 3.2-22.

Background dose varies by location primarily because of elevation changes and variations in the dose from radon. As elevation increases so does the dose from cosmic radiation and hence the total dose. Radon is a radioactive gas produced from the decay of ^{238}U , which is naturally found in soil. The amount of radon in the soil/bedrock depends on the type the porosity and moisture content. Areas which have types of soils/bedrock like granite and limestone have higher radon levels than those with other types of soils/bedrock (EPA, 2006).

The total effective dose equivalent is the total dose from external sources and internal material released from licensed operations. Doses from sources in the general environment (such as terrestrial radiation, cosmic radiation, and naturally occurring radon) are not included in the dose calculation for compliance with 10 CFR Part 20, even if these sources are from technologically enhanced naturally occurring radioactive material (TENORM), such as pre-existing radioactive residues from prior mining (Atomic Safety and Licensing Board, 2006).

For the Northwestern New Mexico Uranium Milling Region, the average background rate including natural and manmade sources for the state of New Mexico is used which is 3.15 mSv/yr [315 mrem/yr] (EPA, 2006). This average background rate in New Mexico is lower than the U.S. average rate of 3.6 mSv/yr [360 mrem/yr] primarily because average annual radon dose is less for New Mexico {1.32 mSv/yr [132 mrem/yr] versus the national average of 2 mSv/yr [200 mrem/yr]}. The background contribution from cosmic radiation is slightly higher for New Mexico versus the U.S. average {0.47 mSv/yr [47 mrem/yr] versus the national average of 0.27 mSv/yr [27 mrem/yr]}. The remaining contributors to background dose (terrestrial radiation, internal radiation, and manmade) are similar for New Mexico {1.36 mSv [136 mrem/yr]} and the U.S. average {1.33 mSv/yr [133 mrem/yr]}. The combination of these differences results in a decrease from the national average of about 0.45 mSv [45 mrem/yr].

3.5.11.2 Public Health and Safety

Public health and safety standards are the same regardless of a facility's location. Therefore, see Section 3.2.11.2 for further discussion of these public health and safety standards.

3.5.11.3 Occupational Health and Safety

Occupational health and safety standards are the same regardless of facility's location. Therefore, see Section 3.2.11.3 for further discussion of these occupational health and safety standards.

3.5.12 References

50states.com. "New Mexico." 2008. <<http://50states.com>> (15 April 2008).

Acoma, New Mexico. "Sky City." 2007. <www.skycity.com> (13 October 2007).

Adams, E.C. and A.I. Duff, eds. *The Protohistoric Pueblo World, A.D. 1275–1600*. Tucson, Arizona: University of Arizona Press. 2004.

Adams, S.S. and A.E. Saucier. "Geology and Recognition Criteria for Uraniferous Humate Deposits, Grants Uranium Region, New Mexico—Final Report." Open-File Report GJBX. Vol. 2, No. 81. Washington, DC: DOE. 1981

Allen, C. and B. Nelson. *Anasazi and Navajo Land Use in the McKinley Mine Area Near Gallup, New Mexico: Archaeology (Vol. 1)*. Albuquerque, New Mexico: University of New Mexico, Office of Contract Archaeology. 1982.

Amick, D. "Folsom Diet Breadth and Land Use in the American Southwest." Ph.D. dissertation. University of New Mexico, Department of Anthropology. Albuquerque, New Mexico. 1994.

Anschuetz, K.F. "Introducing a Landscape Approach for Evaluating Traditional Senses of Time and Place." *More Than a Scenic Mountain Landscape: Valles Caldera National Preserve Land Use History*. K.F. Anschuetz and T. Merlan, eds. Fort Collins, Colorado: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. pp. 249–262. 2007.

Anschuetz, K.L., R.H. Wilshusen, and C. Schick. "An Archaeology of Landscape: Perspectives and Directions." *Journal of Archaeological Research*. Vol. 9, No. 2. pp. 157–211. 2001.

Audubon Society. "2007 Watchlist." 2007. <<http://www.audubon2.org/watchlist>> (12 February 2008).

Basso, K.H. *Wisdom Sits in Places: Landscape and Language Among the Western Apache*. Albuquerque, New Mexico: University of New Mexico. 1996.

Biota Information System of New Mexico. "Species Booklets and Reports." Santa Fe, New Mexico: New Mexico Department of Game and Fish. 2007. <<http://www.bison-m.org>> (5 October 2007).

BLM. "Visual Resource Management." Manual 8400. Washington, DC: BLM. 2007a. <<http://www.blm.gov/nstc/VRM/8400.html#Anchor-.06-23240>> (17 October 2007).

- 1 BLM. "Visual Resource Inventory." Manual H-8410-1. Washington, DC: BLM. 2007b.
2 <<http://www.blm.gov/nstc/VRM/8410.html>> (17 October 2007).
3
- 4 BLM. "Visual Resource Contrast Rating." Manual 8431. Washington, DC: BLM. 2007c.
5 <<http://www.blm.gov/nstc/VRM/8431.html>> (17 October 2007).
6
- 7 BLM. "Farmington Proposed Resource Management Plan and Final Environmental Impact
8 Statement." BLM-NM-PL-03-014-1610. Farmington, New Mexico: BLM, Farmington Field
9 Office. 2003. <http://www.nm.blm.gov/ffo/ffo_p_rmp_feis/ffo_p_rmp_index.html>
10 (17 October 2007).
11
- 12 BLM. "Proposed El Malpais Plan and Final Environmental Impact Statement." Albuquerque,
13 New Mexico: BLM, Albuquerque Field Office. 2000.
14
- 15 Broster, J. and B. Harrill. *A Cultural Resource Management Plan for Timber Sale and Forest*
16 *Development Area on the Pueblo of Acoma (Vol. 1 and 2)*. Albuquerque, New Mexico: Bureau
17 of Indian Affairs. 1982.
18
- 19 Brugge, D. "Navajos in the Catholic Church Records of New Mexico 1694-1875." Research
20 Report No. 1. Window Rock, New Mexico: Parks and Recreation Department, Navajo Nation.
21 1968.
22
- 23 Bureau of Indian Affairs. "American Indian Population and Labor Force Report." Washington,
24 DC: Bureau of Indian Affairs. 2003.
25
- 26 Center for Plant Conservation. "National Protection Plant Profile, Recovering Americas
27 Vanishing Flora." 2008 <[http://www.centerforplantconservation.org/](http://www.centerforplantconservation.org/ASP/CPC_ViewProfile.asp?CPCNum=3241)
28 [ASP/CPC_ViewProfile.asp?CPCNum=3241](http://www.centerforplantconservation.org/ASP/CPC_ViewProfile.asp?CPCNum=3241)> (15 February 2008).
29
- 30 Chenoweth, W.L. "Geology and Production History of Uranium Deposits in the Dakota
31 Sandstone, McKinley County, New Mexico." *New Mexico Geology*. Vol. 11. pp. 21-29. 1989.
32
- 33 Chenoweth, W.L. and E.A. Learned. "Stratigraphic Section, Church Rock Area, McKinley
34 County, New Mexico." *Geology and Mineral Technology of the Grants Uranium Region* 1979.
35 C.A. Rautman, ed. New Mexico Bureau of Mines and Mineral Resources Memoir 38.
36 Socorro, New Mexico: New Mexico Bureau of Mines and Mineral Resources. 1980.
37
- 38 City of Gallup Economic Development Center. "City of Gallup Departments." 2007.
39 <www.ci.gallup.nm.us/dept.htm> (18 October 2007).
40
- 41 Cordell, L.S. "Chaco's Corn: Where Was It Grown?" *Search of Chaco: New Approaches to an*
42 *Archaeological Enigma*. D.G. Noble, ed. Santa Fe, New Mexico: School of American
43 Research Press. 2004.
44
- 45 Cordell, L.S. "Cultural Resources Overview: Middle Rio Grande Valley, New Mexico."
46 Albuquerque, New Mexico: U.S. Department of Agriculture, Forest Service, Southwestern
47 Region, and Santa Fe, New Mexico: Bureau of Land Management, New Mexico State Office.
48 1979.
49

Description of the Affected Environment

- Correll, J. "Through the White Man's Eyes: A Contribution to Navajo History. A Chronological Record of the Navajo People From Earliest Times to the Treaty of June 1, 1868." Publication No. 1. Window Rock, New Mexico: Navajo Heritage Center. 1976.
- Crumley, C. and W.H. Marquardt. "Landscape: A Unifying Concept in Regional Analysis." *Interpreting Space: GIS and Archaeology*. K.M.S. Allen, S.W. Green, and E.B.W. Zubrow, eds. London, England: Taylor and Francis. pp. 49–54. 1990.
- Damp, J., S. Hall, and S. Smith. "Early Irrigation on the Colorado Plateau Near Zuni Pueblo, New Mexico." *American Antiquity*. Vol. 67, No. 4. pp. 665–676. 2002.
- Dean, J.S., W.H. Doelle, and J.D. Orcutt. "Adaptive Stress: Environment and Demography." *Themes in Southwest Prehistory*. G.J. Gumerman, ed. Santa Fe, New Mexico: School of American Research Press. pp. 53–86. 1994.
- Division of Economic Development of the Navajo Nation. "Educational Facilities on the Navajo Nation." Window Rock, Arizona: Division of Economic Development of the Navajo Nation. 2008. <www.navajobusiness.com/infrastructure/Education%20Facilities.htm>. 23 February 2008).
- Division of Economic Development of the Navajo Nation. "2005–2006 Comprehensive Economic Development Strategy of Navajo Nation." Window Rock, New Mexico: Division of Economic Development of the Navajo Nation. 2006.
- Driscoll, F.G. "Groundwater and Wells." Second edition. St. Paul, Minnesota: Johnson Filtration Systems Inc. 1986.
- Dulaney, A. and S. Dosh. "A Class II Cultural Resources Inventory of the Southern Portion of the Chaco Planning Unit, McKinley and Sandoval Counties, New Mexico." Albuquerque, New Mexico: Bureau of Land Management. 1981.
- Ellis, F.H. "Laguna Pueblo." *Handbook of North American Indians: Southwest*. A. Ortiz, ed. Vol. 9. Washington, DC: Smithsonian Institution. 1979.
- Ellis, F.H. *Archaeologic and Ethnologic Data: Acoma-Laguna Land Claims: Pueblo Indians II*. New York City, New York: Garland Publishing. 1974a.
- Ellis, F.H. *Anthropology of Laguna Pueblo Land Claims: Pueblo Indians III*. New York City, New York: Garland Publishing. 1974b.
- EPA. "National Assessment Database." 2008. <<http://www.epa.gov/waters/305b/index.html>> (28 February 2008).
- EPA. "Counties Designate Nonattainment or Maintenance for Clean Air Act's National Ambient Air Quality Standards (NAAQS)." 2007a. <<http://www.epa.gov/oar/oaqps/greenbk/mapnmpoll.html>> (29 September 2007).
- EPA. "Prevention of Significant Deterioration (PSD) Permit Program Status: May 2007." 2007b. <<http://www.epa.gov/nsr/where.html>> (26 September 2007).
- EPA. "Assessment of Variations in Radiation Exposure in the United States (Revision 1)." Contract Number EP–D–05–02. Washington, DC: EPA. 2006.

- 1
2 Federal Highway Administration. "Synthesis of Noise Effects on Wildlife Populations."
3 FHWA-HEP-06-016. Washington, DC: Federal Highway Administration, Department of
4 Transportation. 2004.
5
6 Ferguson, T.J. *Historic Zuni Architecture and Society: An Archaeological Application of Space*
7 *Syntax: Anthropological Papers of the University of Arizona 60*. Tucson, Arizona: University of
8 Arizona Press. 1996.
9
10 Ferguson, T.J. and E. Hart. *A Zuni Atlas*. Norman, Oklahoma: University of Oklahoma Press.
11 1985.
12
13 Flint, R. and S.C. Flint. *Documents of the Coronado Expedition, 1539-1542*. Dallas, Texas:
14 Southern Methodist University Press. 2005.
15
16 Fox, S. "Sacred Pedestrians: The Many Faces of Southwest Pilgrimage." *Journal of the*
17 *Southwest*. Vol. 36. pp. 33-53. 1994.
18
19 Gilpin, D. "Social Transformations and Community Organization in the Southwest San Juan
20 Basin, New Mexico: Archaeological Investigations Along Navajo Route 9, Twin Lakes to
21 Standing Rock." Flagstaff, Arizona: SWCA Environmental Consultants. 2007.
22
23 Grants-Cibola County Schools. "Grants-Cibola County Schools." Grants, New Mexico:
24 Grants-Cibola County School District. 2007. <www.gccs.cc> 15 October 2007).
25
26 Greatschools. "McKinley County." 2007c. <www.greatschools.net> (15 October 2007).
27
28 Green, W.G. and C.T. Pierson. "A Summary of the Stratigraphy and Depositional Environments
29 of Jurassic and Related Rocks in the San Juan Basin." San Juan Basin III, New Mexico
30 Geological Society Guidebook, 28th Field Conference. J.E. Fassat, ed. pp. 147-152. 1977.
31
32 Griffith, G.E., J.M. Omerik, M.M. McGraw, G.Z. Jacobi, C.M. Canavan, T.S. Schrader,
33 D. Mercer, R. Hill, and B.C. Moran. "Ecoregions of New Mexico." U.S. Geological Survey Map.
34 Scale 1:1,400,000. Reston, Virginia: U.S. Geological Survey. 2006.
35
36 Hilpert, L.S. "Uranium Resources of Northwestern New Mexico." U.S. Geological Survey
37 Professional Paper 603. 1969.
38
39 Hilpert, L.S. "Regional and Local Stratigraphy of Uranium-Bearing Rocks." *Geology and*
40 *Technology of the Grants Uranium Region*. V.C. Kelley, ed. New Mexico Bureau of Mines and
41 Mineral Resources Memoir 15. Socorro, New Mexico: New Mexico Bureau of Mines and
42 Mineral Resources. pp. 6-18. 1963.
43
44 Holen, H.K. and W.O. Hatchell. "Geological Characterization of New Mexico Uranium Deposits
45 for Extraction by In-Situ Leach Recovery." New Mexico Bureau of Mines and Mineral
46 Resources Open-File Report 251. Socorro, New Mexico: New Mexico Bureau of Mines and
47 Mineral Resources. 1986.
48
49 Holt, B. "Navajo Sacred Areas: Guide for Management." *Contract Abstracts and CRM*
50 *Archaeology*. Vol. 2, No. 2. pp. 45-53. Albuquerque, New Mexico: Atechiston, Inc. 1981.
51

Description of the Affected Environment

- 1 Huber, E.K. and C.R. Van West. "Synthetic Studies. Archaeological Data Recovery at the New
2 Mexico Transportation Corridor and First Five-Year Permit Area, Fence Lake Coal Mine Project,
3 Catron County, New Mexico." Vol. 3. Technical Series 84. Tucson, Arizona: Statistical
4 Research. 2005.
- 5
6 Hunt, C.B. "Natural Regions of the United States and Canada." San Francisco, California:
7 W.H. Freeman and Company. 1974.
- 8
9 Huntley, D.L. and K.W. Kintigh. "Archaeological Patterning and Organizational Scale of Late
10 Prehistoric Settlement Clusters in the Zuni Region of New Mexico." *The Protohistoric Pueblo*
11 *World, A.D. 1275-1600*. E.C. Adams and A.I. Duff, eds. Tucson, Arizona: University of
12 Arizona. 2004.
- 13
14 Jenkins, M.E. *History of Laguna Pueblo Land Claims. Pueblo Indians IV*. New York City, New
15 York: Garland Press. 1974.
- 16
17 Judge, J.W. "Chaco's Golden Century." *Search of Chaco: New Approaches to an*
18 *Archaeological Enigma*. D.G. Noble, ed. Santa Fe, New Mexico: School of American
19 Research Press. pp. 1-6. 2004.
- 20
21 Kantner, J. *Ancient Puebloan Southwest*. Cambridge, United Kingdom: Cambridge University
22 Press. 2004.
- 23
24 Kantner, J. "Ancient Roads, Modern Mapping." *Expedition*. Vol. 39, No. 3. 1997.
- 25
26 Kantner, J. and N.M. Mahoney, eds. *Great House Communities Across the Chacoan*
27 *Landscape: Anthropological Papers of the University of Arizona No. 64*. Tucson, Arizona:
28 University of Arizona Press. 2000.
- 29
30 Kearns, T., C. Kugler, and P. Simmon. "The Dog Leg Site (LA 6448): A Basketmaker II Cache
31 Local in the Southern Chuska Valley, New Mexico." *Pipeline Archaeology 1990-1993: The*
32 *El Paso Natural Gas North System Expansion Project, New Mexico and Arizona (Vol. II)*.
33 T. Kearns and J. McVickar, eds. Farmington, New Mexico: Western Cultural Resource
34 Management. pp. 301-476. 1998.
- 35
36 Kelley, K. *Navajo Land Use: An Ethnoarchaeological Study*. New York City, New York:
37 Academic Press. 1986.
- 38
39 Kelley, K. "Anasazi and Navajo Land Use in the McKinley Mine Area Near Gallup, New Mexico
40 (Ethnohistory, Vol. 2)." Albuquerque, New Mexico: University of New Mexico, Office of
41 Contract Archeology. 1982.
- 42
43 Kelley, K. and H. Francis. *Navajo Sacred Places*. Bloomington, Indiana: University of Indiana
44 Press. 1994.
- 45
46 Kidder, A.V. "Southwestern Archaeological Conference." *Science*. Vol. 66. pp. 489-491.
47 1927.
- 48
49 King, T. *Places That Count: Traditional Cultural Properties in Cultural Resources Management*.
50 Walnut Creek, California: Altamira Press. 2003.
- 51

- 1 Kintigh, K.W. "The Cibola Region in the Post-Chacoan Era." *The Prehistoric Pueblo World, A.D. 1150–1350*. M.A. Adler, ed. Tucson, Arizona: University of Arizona Press. pp. 131–144.
2 1996.
3
4
- 5 Lat-Long.com. Laguna Indian Reservation. 2008. <www.lat-long.com> (28 February 2008).
6
7 Linford, L. *Navajo Places: History, Legend and Landscape*. Salt Lake City, Utah: University of
8 Utah Press. 2000.
9
- 10 Localschooldirectory.com. "Grants Schools in New Mexico". 2008.
11 <www.localschooldirectory.com> (28 February 2008).
12
- 13 Mahoney, N.M. and J. Kantner. "Chacoan Archaeology and Great House Communities." *Great*
14 *House Communities Across the Chacoan Landscape, Anthropological Papers of the University*
15 *of Arizona No. 64*. J. Kantner and N.M. Mahoney, eds. Tucson, Arizona: University of Arizona
16 Press. pp. 1–15. 2000.
17
- 18 Mapquest. New Mexico. <<http://www.mapquest.com>> (15 April 2008).
19
- 20 Marshall, M., J. Stein, R. Loose, and J. Novotny. *Anasazi Communities of the San Juan Basin*.
21 Public Service Company of New Mexico. Santa Fe, New Mexico: New Mexico Historic
22 Preservation Division. 1979.
23
- 24 McLemore, V.T. "Uranium Resources in New Mexico." Proceedings 2007 Annual Meeting and
25 Exhibit Society for Mining, Metallurgy, and Exploration, Denver, Colorado, February 25–28,
26 2007. Littleton, Colorado: Society for Mining, Metallurgy, and Exploration. 2007.
27
- 28 McLemore, V.T. and W.L. Chenoweth. "Uranium Resources in New Mexico." New Mexico
29 Bureau of Mines and Mineral Resources Map 18. Socorro, New Mexico: New Mexico Bureau
30 of Mines and Mineral Resources. 1989.
31
- 32 Mills, B.J. "Acts of Resistance: Zuni Ceramics, Social Identity, and the Pueblo Revolt." *Archaeologies of the Pueblo Revolt*. R.W. Preucel, ed. Albuquerque, New Mexico: University
33 of New Mexico Press. pp. 85–98. 2002a.
34
35
- 36 Mills, B.J. "Recent Research on Chaco: Changing Views on Economy, Ritual and Society." *Journal of Archaeological Research*. Vol. 10, No. 1. pp. 65–117. 2002b.
37
38
- 39 Minge, W. *Defense of the Pueblo of Acoma Land Claims: Pueblo Indians III*. New York City,
40 New York: Garland Publishing. 1974.
41
- 42 Murphy, D., G. Huey, and W. Clay. "El Morro National Monument." Globe, Arizona:
43 Southwestern Parks and Monuments. 2003.
44
45
- 46 National Climatic Data Center. "NCDC U.S. Storm Events Database." 2007.
47 <<http://www4.ncdc.noaa.gov/cgi-win/wwcgi.dll?wwEvent~Storms>> (14 April 2008).
48
- 49 National Climatic Data Center. "Climates of the States, Climatology of the United States No. 60
50 (New Mexico, Nebraska, South Dakota, and Wyoming)." Asheville, North Carolina: National
51 Oceanic and Atmospheric Administration. 2005. <<http://cdo.ncdc.noaa.gov/cgi->

Description of the Affected Environment

bin/climatenormals/climatenormals.pl?directive=prod_select2&prodtype=CLIM60&subnum=>
(30 January 2005).

National Climatic Data Center. "Climatology of the United States No. 20: Monthly Station Climate Summaries, 1971-2000." Asheville, North Carolina: National Oceanic and Atmospheric Administration. 2004.

National Council on Radiation Protection and Measurements. "Report No. 094—Exposure of the Population in the United States and Canada From Natural Background Radiation." Bethesda, Maryland: National Council on Radiation Protection & Measurements. 1987.
Navajo Nation. "Navajo Code Annotated. Titles 1–5." Window Rock, Arizona: Navajo Nation. 2005.

National Weather Service. "NOAA Technical Report NWS 33: Evaporation Atlas for the Contiguous 48 United States." Washington, DC: National Oceanic and Atmospheric Administration. 1982.

NatureServe. "NatureServe Explorer: An Online Encyclopedia of Life (Web Application)." Version 7.0. 2008. Arlington, Virginia: NatureServe. <<http://www.natureserve.org/explorer>> (14 April 2008).

Navajo Nation. <www.navajolorg/govt.htm> 25 February 2008).

New Mexico Department of Game and Fish. "Wildlife Notes." Albuquerque, New Mexico: New Mexico Department of Game and Fish. 2008. <http://www.wildlife.state.nm.us/education/wildlife_notes/WildlifeNotes.htm> (15 February 2008).

New Mexico Department of Game and Fish. "Comprehensive Wildlife Conservation Strategy for New Mexico." Santa Fe, New Mexico: New Mexico Department of Game and Fish. 2006.

New Mexico Department of Game and Fish. "Zuni Bluehead Sucker Recovery Plan." 2004. <http://www.wildlife.state.nm.us/conservation/threatened_endangered_species/documents/ZuniBlueheadSuckerRecoveryPlan.pdf> (14 February 2008).

New Mexico Department of Transportation. "New Mexico 2006 Traffic Survey." Albuquerque, New Mexico: New Mexico Department of Transportation. 2007. <<http://nmshtd.state.nm.us/main.asp?secid=15370>> (26 February 2008).

New Mexico Economic Development. "Grants." 2008. <www1.edd.statenn.us/datacenter/moredata/index.html> (27 February 2008).

New Mexico Environmental Department, Air Quality Bureau. "Dispersion Modeling—Meteorological Data." 2007. <http://www.nmenv.state.nm.us/aqb/modeling/metfiles/ISCST3/SCRAM_GALLUP_91.pdf> (13 September 2007).

New Mexico Environment Department. "Air Quality Regulations." 2002. <<http://www.nmenv.state.nm.us/aqb/>> (23 October 2007).

New Mexico State University. "Southwest Regional Gap Analysis Project." Las Cruces, New Mexico: New Mexico State University. 2007. <<http://fws-nmcfwru.nmsu.edu/swregap/>> (29 November 2007).

- 1
2 New Mexico Taxation and Revenue Department. "2006 Property Tax Facts." Albuquerque,
3 New Mexico: New Mexico Taxation and Revenue Department. 2008.
4
5 New Mexico Taxation and Revenue Department. "New Mexico Oil and Gas Ad Valorem
6 Production and Production Equipment Taxes—Description and Summary of Statistical
7 Reports." Albuquerque, New Mexico: New Mexico Taxation and Revenue Department.
8 Unknown.
9
10 New Mexico Tourism Department. "New Mexico: The Land of Enchantment Tourism
11 Department." 2008. <www.newmexico.org/place/loc/cities/page/DB-place/place/1200.html>
12 (25 February 2008).
13
14 Noble, D.G., ed. *Search of Chaco: New Approaches to an Archaeological Enigma*.
15 Santa Fe, New Mexico: School of American Research Press. 2004.
16
17 Norwood, V. and J. Monk, eds. *The Desert Is No Lady: Southwestern Landscapes in Women's*
18 *Writing and Art*. New Haven, Connecticut: Yale University Press. 1987.
19
20 NRC. "Expected New Uranium Recovery Facility Applications/Restarts/Expansions: Updated
21 1/24/2008." 2008. <[http://www.nrc.gov/info-finder/materials/uranium/2008-ur-projects-list-](http://www.nrc.gov/info-finder/materials/uranium/2008-ur-projects-list-public-012408.pdf)
22 [public-012408.pdf](http://www.nrc.gov/info-finder/materials/uranium/2008-ur-projects-list-public-012408.pdf)> (08 February 2008).
23
24 NRC. NUREG-1508, "Final Environmental Impact Statement To Construct and Operate the
25 Crownpoint Uranium Solution Mining Project, Crown Point, New Mexico." Washington, DC:
26 NRC. February 1997.
27
28 NRCS, United States Department of Agriculture. "Soil Survey of McKinley County Area, New
29 Mexico, McKinley County and Parts of Cibola and San Juan Counties. Albuquerque, New
30 Mexico: New Mexico NRCS State Office. 2001.
31
32 Orcutt, J.D., J.L. McVickar, and J.D. Kilby. "Environmental and Cultural Background." *The*
33 *El Malpais Archeological Survey, Phase I*. R.P. Powers and J.D. Orcutt, eds. Intermountain
34 Cultural Resources Management Professional Paper Vol. 70. Santa Fe, New Mexico: National
35 Park Service. pp. 31-44. 2005.
36 Ortiz, J. *Woven Stone*. Tucson, Arizona: University of Arizona Press. 1992.
37
38 Owen, D.E. "The Jackpile Sandstone Member of the Morrison Formation in West-Central New
39 Mexico—A Formal Definition." *New Mexico Geology*. Vol. 6, No. 3. pp. 45-52. 1984.
40
41 Parker, P. and T. King. "Guidelines for Evaluating and Documenting Traditional Cultural
42 Properties." National Register Bulletin 38. Washington, DC: National Park Service. 1998.
43
44 Parsons, E.C. "Notes on Acoma and Laguna." *American Anthropologist*. pp. 162-186. 1918.
45
46 Perlman, S. "Fort Wingate Depot Activity Ethnographic Study." Albuquerque, New Mexico:
47 University of New Mexico, Office of Contract Archaeology. 1997.
48
49 Peterson, R.J. "Geology of Pre-Dakota Uranium Geochemical Cell, Section 13, T.16. N, R.17.
50 W., Church Rock Area, McKinley County." *Geology and Mineral Technology of the Grants*

Description of the Affected Environment

- 1 Uranium District. New Mexico Bureau of Mines and Mineral Resources, Memoir 38. Socorro,
2 New Mexico: New Mexico Bureau of Mines and Mineral Resources. pp. 131–134. 1980.
3
- 4 Plog, F. and W. Wait. "The San Juan Basin Tomorrow: Planning for the Conservation of
5 Cultural Resources in the San Juan Basin." Santa Fe, New Mexico: National Park Service.
6 1979.
7
- 8 Powers, R. and J. Orcutt. "The El Malpais Archaeological Survey." Professional Report No. 70.
9 Santa Fe, New Mexico: Intermountain Cultural Resources Management. 2005a.
10
- 11 Powers, R. and J. Orcutt, eds. "Site Typology, Architecture, Population and Settlement
12 Patterns." *The El Malpais Archeological Survey, Phase I*. Professional Paper. Vol. 80.
13 Santa Fe, New Mexico: Intermountain Cultural Resources Management. pp. 71–97. 2005b.
14
- 15 Powers, R., B. Gillespie, and S. Lekson. "The Outlier Survey: A Regional View of Settlement in
16 the San Juan Basin." Reports of the Chaco Center No. 3. Albuquerque, New Mexico: National
17 Park Service. 1983.
18
- 19 Public Employees for Environmental Responsibility. "Common Black Hawk." 2008.
20 <<http://www.txpeer.org/limpia/CommonBlackHawk.html>> (14 February 2008).
21
- 22 Public Schools Report, Public School Information and Data—Acoma Indian Reservation." 2007.
23 <<http://schools.publicschoolsreport.com>>. (15 October 2007).
24
- 25 Publicschoolreview.com. "Sandoval County." 2008 <www.publicschoolreview.com>
26 (28 February 2008).
27
- 28 Pueblo Indians. "Pueblo Indians." 2008. <[www.everyculture.com/North-America/pueblo-](http://www.everyculture.com/North-America/pueblo-Indians.html)
29 [Indians.html](http://www.everyculture.com/North-America/pueblo-Indians.html)> (26 February 2008).
30
- 31 Pueblo of Zuni. "Pueblo Indians." 2008 <www.ashiwi.org/> (25 February 2008).
32
- 33 Ramah Navajo Chapter. "A History of the Ramah Navajo Community." Ramah, New Mexico:
34 Tohajiilee Navajo Chapter. 2008. <www.tohajiilee.ndes.org> (28 February 2008).
35
- 36 Rands, R. *Acoma Land Utilization: Pueblo Indians III*. New York City, New York: Garland
37 Publishing. 1974a.
38
- 39 Rands, R. *Laguna Land Utilization: Pueblo Indians IV*. New York City, New York: Garland
40 Publishing. 1974b.
41
- 42 Reed, P.F., ed. *Foundations of Anasazi Culture: The Basketmaker-Pueblo Transition*.
43 Salt Lake City, Utah: University of Utah Press. 2000a.
44
- 45 Reed, P.F., ed. "Fundamental Issues in Basketmaker Archaeology." *Foundations of Anasazi*
46 *Culture: The Basketmaker-Pueblo Transition*. Salt Lake City, Utah: University of Utah Press.
47 pp. 3–16. 2000b.
48
- 49 Reeve, F. "The Navajo-Spanish Peace, 1720s–1770s." *New Mexico Historical Review*.
50 Vol. 34. pp. 9–30. 1959.
51

- 1 Riggs, C.R. "Late Ancestral Pueblo or Mogollon Pueblo? An Architectural Perspective on
2 Identity." *Kiva*. Vol. 70, No. 4. pp. 323–348. 2005.
- 3
4 Ristorcelli, S.J. "Geology of Eastern Smith Lake Ore Trend, Grants Mineral Belt." *Geology and
5 Mineral Technology of the Grants Uranium Region*. New Mexico Bureau of Mines and Mineral
6 Resources Memoir 38. Socorro, New Mexico: New Mexico Bureau of Mines and Mineral
7 Resources. pp. 145–152. 1980.
- 8
9 Robson, S.G. and E.R. Banta. "Ground Water Atlas of the United States: Arizona, Colorado,
10 New Mexico, Utah, HA730–C." 1995. <http://capp.water.usgs.gov/gwa/ch_c/C-test6.html>
- 11
12 Roney, J.R. "The Pueblo III Period in the Eastern San Juan Basin and Acoma-Laguna Areas."
13 *The Prehistoric Pueblo World, A.D. 1150–1350*. M.A. Adler, ed. Tucson, Arizona: University of
14 Arizona Press. pp. 145–169. 1996.
- 15
16 Sanford, R.F. "A New Model for Tabular-Type Uranium Deposits." *Economic Geology*. Vol. 87.
17 pp. 2,041–2,055. 1992.
- 18
19 Saucier, A.E. "Tertiary Oxidation in Westwater Canyon Member of the Morrison Formation."
20 *Geology and Mineral Technology of the Grants Uranium Region 1979*. C.A. Rautman, compiler.
21 New Mexico Bureau of Mines and Mineral Resources Memoir 38. Socorro, New Mexico:
22 New Mexico Bureau of Mines and Mineral Resources. pp. 116–121. 1981.
- 23
24 Schachner, G. and J.D. Kilby. "Archaic and Puebloan Chronology." *The El Malpais
25 Archeological Survey, Phase I*. R.P. Powers and J.D. Orcutt, eds. Intermountain Cultural
26 Resources Management Professional Paper No. 70. Sante Fe, New Mexico: National Park
27 Service. 2005.
- 28
29 Schutt, J.A. and R.C. Chapman. "Cycles of Closure: A Cultural Resources Inventory of Fort
30 Wingate Depot Activity, New Mexico." Albuquerque, New Mexico: University of New Mexico,
31 Office of Contract Archaeology. 1997.
- 32
33 Sedgwick, W. *Acoma the Sky City: A Study in Pueblo-Indian History and Civilization*.
34 Cambridge, United Kingdom: Harvard University Press. 1926.
- 35
36 Silko, L. "Landscape, History, and Pueblo Imagination." *The Norton Book of Nature Writing*. R.
37 Finch and J. Elder, eds. New York City, New York: W.W. Norton and Company. pp. 882–893.
38 1990.
- 39
40 Simmons, A. "New Evidence for the Early Use of Cultigens in the American Southwest."
41 *American Antiquity*. Vol. 51. pp. 73–89. 1986.
- 42
43 Snead, J. and R. Preucel. "The Ideology of Settlement: Ancestral Keres Landscapes in the
44 Northern Rio Grande." *Archaeologies of Landscape: Contemporary Perspectives*.
45 W. Ashmore and A.B. Knapp, eds. Oxford, United Kingdom: Blackwell. pp. 169–200. 1999.
- 46
47 Spicer, E.H. *Cycles of Conquest: The Impact of Spain, Mexico, and the United States on the
48 Indians of the Southwest, 1533–1960*. Tucson, Arizona: University of Arizona Press. 1962.
- 49
50 Stanford, D. "Paleoindian Archaeology and Late Pleistocene Environments in the Plains and
51 Southwestern United States." *Ice Age Peoples of North America (Second Edition)*.

Description of the Affected Environment

- 1 R. Bonnicksen and K. Turnmire, eds. College Station, Texas: Texas A&M University, Center
2 for the Study of the First Americans. pp. 281–339. 2005.
- 3
- 4 Sterling, M.W. "Origin Myth of Acoma." Bulletin 135. Washington, DC: Bureau of American
5 Ethnology. 1942.
- 6
- 7 Tainter, J. and D. Gillio. "Cultural Resources Overview: Mt. Taylor Area, New Mexico."
8 Albuquerque, New Mexico: USFS. 1980.
- 9
- 10 Texas Parks and Wildlife Department. "Wildlife Facts Sheets." 2007.
11 <<http://www.tpwd.state.tx.us/huntwild/wild/species/>> (15 February 2008).
- 12
- 13 Thamm, J.K., A.A. Kovschak, and S.S. Adams. "Geology and Recognition Criteria for
14 Sandstone Uranium Deposits of the Salt Wash Type, Colorado Plateau Province—Final
15 Report." Report GJBX. Vol. 6, No. 81. Washington, DC: DOE. 1981.
- 16
- 17 Tohajiilee Chapter. <www.tohajiilee.ndes.org> (28 February 2008).
- 18
- 19 Toll, H.W. "Artifacts in Chaco: Where They Came From and What They Mean." *In Search of*
20 *Chaco: New Approaches to an Archaeological Enigma*. D.G. Noble, ed. Santa Fe,
21 New Mexico: School of American Research Press. pp. 32–40. 2004.
- 22
- 23 Towner, R.H., ed. *The Archaeology of Navaho Origins*. Salt Lake City, Utah: University of
24 Utah Press. 1996.
- 25
- 26 Udall, S., D. Brugge, T. Benally, and E. Yazzie-Lewis. *The Navajo People and Uranium Mining*.
27 Albuquerque, New Mexico: University of New Mexico Press. 2007.
- 28
- 29 USACE. "Final Environmental Assessment and Finding of No Significant Impact: The Los Utes
30 Acequia Pipeline Project Sandoval County, New Mexico." Albuquerque, New Mexico: USACE.
31 2007.
- 32
- 33 USACE. "Interim Regional Supplement to the Corps of Engineers Wetland Delineation Manual:
34 Arid West Region." Washington, DC: USACE. 2006.
- 35
- 36 U.S. Census Bureau. "American FactFinder 2000 Census Data." 2008.
37 <<http://factfinder.census.gov>> (25 February 2008).
- 38
- 39 Usacops. New Mexico. <<http://www.usacops.com>> (15 April 2008).
- 40
- 41 U.S. Census Bureau. American FactFinder 2000 Census Data. <<http://factfinder.census.gov>>
42 25 February 2008).
- 43 U.S. Department of the Interior, Bureau of Indian Affairs. "2003 Indian Population and Labor
44 Force Report." 2003. <<http://www.doi.gov/bia/laborforce/2003LaborForceReportFinalAll.pdf>>
45 (24 October 2007).
- 46
- 47 U.S. Fish and Wildlife Service. "Ecological Services." 2008. <[http://www.fws.gov/](http://www.fws.gov/southwest/es/)
48 [southwest/es/](http://www.fws.gov/southwest/es/)> (12 February 2008).
- 49 U.S. Fish and Wildlife Service. "Rio Grande Silvery Minnow Draft Recovery Plan." 2007.
50 <http://ecos.fws.gov/docs/recovery_plan/070118a.pdf> (12 February 2008).
- 51

- 1 U.S. Fish and Wildlife Service. "San Juan River Basin Recovery Implementation Program."
2 2006. <[http://www.fws.gov/southwest/sjrip/pdf/DOC_SJRIP_Final_](http://www.fws.gov/southwest/sjrip/pdf/DOC_SJRIP_Final_Program_Document_9_1_2006.pdf)
3 [Program_Document_9_1_2006.pdf](http://www.fws.gov/southwest/sjrip/pdf/DOC_SJRIP_Final_Program_Document_9_1_2006.pdf)> (14 February 2008).
4
- 5 USFS. "Environmental Assessment for Continental Divide National Scenic Trail Location—
6 Cibola National Forest, Mt. Taylor Ranger District, Cibola and McKinley Counties."
7 Albuquerque, New Mexico: USFS, Southwestern Region. March 2007.
8 <[http://www.fs.fed.us/r3/cibola/](http://www.fs.fed.us/r3/cibola/projects/nepa_reports/continental_divide_trail_0307.pdf)
9 [projects/nepa_reports/continental_divide_trail_0307.pdf](http://www.fs.fed.us/r3/cibola/projects/nepa_reports/continental_divide_trail_0307.pdf)> (17 October 2007).
10
- 11 USFS. "Cibola National Forest General Vicinity Map." Albuquerque, New Mexico: USFS,
12 Southwestern Region. 2006. <<http://www.fs.fed.us/r3/cibola/maps/vicinitysmap.shtml>>
13 (29 February 2008).
14
- 15 USFS. "Cibola National Forest Land and Resource Management Plan." Albuquerque,
16 New Mexico: USFS, Southwestern Region. <[http://www.fs.fed.us/r3/cibola/](http://www.fs.fed.us/r3/cibola/projects/forest_plan/forest_plan.shtml)
17 [projects/forest_plan/forest_plan.shtml](http://www.fs.fed.us/r3/cibola/projects/forest_plan/forest_plan.shtml)> 1985.
18
- 19 U.S. Geological Survey. "A Tapestry of Time and Terrain." Denver, Colorado: U.S. Geological
20 Survey. 2004. <<http://tapestry.usgs.gov/Default.html>> (25 February 2008).
21
- 22 Van Dyke, R.M. "Chaco's Sacred Geography." *In Search of Chaco: New Approaches to an*
23 *Archaeological Enigma*. D.G. Noble, ed. Santa Fe, New Mexico: School of American
24 Research Press. pp. 78–85. 2004.
25
- 26 Vierra, B. "Early Agriculture on the Southwestern Periphery of the Colorado Plateau: A Case in
27 Diversity of Tactics." *Archaeology Without Borders: Contact, Commerce and Change in the*
28 *U.S. Southwest and Northwestern Mexico*. M. McBrinn and L. Webster, eds. Boulder,
29 Colorado: University of Colorado Press. pp. 71–88. 2008.
30
- 31 Vierra, B. "Aceramic and Archaic Research Questions." *Across the Colorado Plateau:*
32 *Anthropological Studies for the Transwestern Pipeline Expansion Project: Excavation and*
33 *Interpretation of Aceramic and Archaic Sites (Vol. 14)*. T. Burchett, B. Vierra, and K. Brown,
34 eds. Albuquerque, New Mexico: University of New Mexico, Office of Contract Archaeology.
35 pp. 375–378. 1994.
36
- 37 Washington State Department of Transportation. "WSDOT's Guidance for Addressing Noise
38 Impacts in Biological Assessments—Noise Impacts." Seattle, Washington: Washington State
39 Department of Transportation. 2006. <[http://www.wsdot.wa.gov/TA/Operations/](http://www.wsdot.wa.gov/TA/Operations/Environmental/NoiseChapter011906.pdf)
40 [Environmental/NoiseChapter011906.pdf](http://www.wsdot.wa.gov/TA/Operations/Environmental/NoiseChapter011906.pdf)> (12 October 2007).
41
- 42 White, L.A. *The Acoma Indians*. Forty-Seventh Annual Report of the Bureau of American
43 Ethnology to the Secretary of the Smithsonian Institution. Washington, DC: Smithsonian
44 Institution. 1932.
45
- 46 Windes, T.C. "The Rise of Early Chacoan Great Houses." *In Search of Chaco: New*
47 *Approaches to an Archaeological Enigma*. D.G. Noble, ed. Santa Fe, New Mexico: School of
48 American Research Press. pp. 14–21. 2004.
49

Description of the Affected Environment

1 Winter, J. "Across the Colorado Plateau: Anthropological Studies for the Transwestern Pipeline
2 Expansion Project: Synthesis and Conclusions (Vol. 20)." Albuquerque, New Mexico:
3 University of New Mexico, Office of Contract Archaeology. 1994.

4
5 Wyoming Department of Transportation. "WYDOT Traffic Analysis." Cheyenne, Wyoming:
6 Wyoming Department of Transportation. 2005. <[http://www.dot.state.wy.us/
7 Default.jsp?sCode=hwyta](http://www.dot.state.wy.us/Default.jsp?sCode=hwyta)> (25 February 2008).

8
9 Zuni Pueblo Public School District. "Zuni Pueblo Official Website of the Zuni People's Public
10 School District." 2008. <www.zpsd.org/zuni/index.htm> (28 February 2008).
11