Joseph Mangano, Executive Director Radiation and Public Health Project 716 Simpson Avenue Ocean City NJ 08226 609-399-4343 odiejoe@aol.com

ROEWEL

M

4/85/08 73FR 82448

Chief, Rulemaking Directives and Editing Branch Division of Administrative Services Office of Administration U.S. Nuclear Regulatory Commission Washington DC 20055

By Email Vogtle LR EIS@nrc.gov

July 16, 2008

Dear Sir/Madam:

On behalf of my colleagues at the Radiation and Public Health Project (RPHP), please accept the following comments on the Environmental Impact Statement for the proposed new nuclear reactors at the Alvin Vogtle plant.

My comments are focused on the health risks of the proposed new reactors that were either not addressed or minimized in the draft EIS statement. Moreover, our statement is unique, as they are based on official data on radioactive contamination and health status.

Based on the following, we believe that new nuclear reactors would pose a serious health risk for local residents around Vogtle:

1. Releases of airborne radioactivity vary, and have greatly exceeded minimal levels. Releases during 2001-2004 from Vogtle 1 are about 10 times greater than from Vogtle 2. Releases in 1992 were over 1000 times greater than in 1987 or 1988 (Tables 1 and 2).

2. From 1987-1990 (as Vogtle began operating) to 1991-2003 (during full operation), average radioactivity levels in drinking water, river water, and sediment downriver or at the Vogtle plant rose (also see Table 3):

Beta in Raw Drinking Water Beta in Finished Drinking Water Beryllium-7 in Sediment Cesium-137 in Sediment Tritium in River Water + 37.1% + 17.8% + 39.5% + 37.4% + 44.6%

50NSI Beview Complete Nemplate = AJM-013

E-REDS=ADM-03 Coll = J.P. Leous (JPL1) S. Hermandez (SHQ)

3. From 1987-1990 to 1991-2003, the cancer death rate for children age 0-24 in Burke County rose 55.5%, vs. a 14.1% decline nationally. The Burke cancer death rate age 25-54 rose 55.1% vs. a 2.9% national decline (Table 4).

4. From 1985-1987 to 1988-1990, as the Vogtle reactors began operating, infant deaths in Burke County rose from 16 to 28, a 70% rise, compared to a 7% U.S. decline (Table 5).

5. From 1979-87 to 1988-2003, the Burke County infant death rate rose 19%, compared to a 30% decline nationally. Thus, the infant death rise in the late 1980s was not a fluke, but the start of a statistically significant change. Increases were significant for all races, blacks, and whites (Table 6).

6. From 1987-1990 to 1991-2003, the cancer death rate in Burke County GA rose 25.1% vs. a 4.2% national decline. Increases were statistically significant for both whites (+17.5%) and blacks (+30.7%), see Table 7.

These data suggest that Vogtle emissions have increased radioactivity levels in the local environment. They also suggest that radioactivity has caused unexplained increases in local mortality rates.

Findings should be considered in the context of Burke County as a high-poverty and high-minority area (Table 8). Populations with inadequate health care coverage have reduced access to needed care; thus, adding a carcinogen to bodies of local residents (through breathing and the food chain) may result in an increased health risk. In addition, the addition of new pollutants (reactors) in a county with a disproportionate percent of African-Americans amounts to environmental injustice.

The fact that the EIS ignores the above data means that it is an incomplete document. Thus, we strongly recommend that the NRC not approve licenses for new reactors at Vogtle, until the above information has been thoroughly examined and the true risks of nuclear power at Vogtle has been understood and related to the public.

Sincerely,

Joseph J. Mangano MPH MBA Executive Director Radiation and Public Health Project Table 1Annual Airborne Emissions from Vogtle Nuclear Plant, 1987-1993

| <u>Year</u> | Microcuries | |
|-------------|-------------|--|
| 1987 | - 20 | |
| 1988 | 18 | |
| 1989 | 1250 | |
| 1990 | 85 | |
| 1991 | 2080 | |
| 1992 | 5870 | |
| 1993 | 521 | |

Source: Tichler J, Doty K, Lucadamo K. Radioactive Materials Released from Nuclear Power Plants. NUREG/CR-2907. Upton NY: Brookhaven National Laboratory, prepared for the U.S. Nuclear Regulatory Commission, Annual Report 1993. Represents Iodine-131 and particulates, all airborne releases of particulates with a half life of at least 8 days. (Annual reports that listed all U.S. reactors ceased in 1993).

3

Table 2

Gaseous Emissions, Fission and Activation Products From Vogtle Nuclear Plant, 2001-2004, in Curies

| Year | Vogtle 1 | Vogtle 2 |
|-------|----------|----------|
| 2001 | 12.13 | 0.42 |
| 2002 | 23.89 | 2.36 |
| 2003 | 1.68 | 0.64 |
| 2004 | 0.64 | 1.31 |
| TOTAL | 38.34 | 4.73 |

Source: U.S. Nuclear Regulatory Commission, www.reirs.com/effluent

Table 3

Trends in Environmental Radioactivity Levels Near Vogtle, 1987-1990 to 1991-2003

| | Annual Avg. | Annual Avg. | |
|---|-------------|------------------|-------------|
| Type of Radioactivity | 1987-1990 | <u>1991-2003</u> | <u>% Ch</u> |
| Beta in Raw Drinking Water | | | |
| - Indicator (downriver) ¹ | 2.583 | 3.540 | + 37.1% |
| - Control (upriver) ² | 3.535 | 3.202 | - 9.4% |
| Beta in Finished Drinking Water | | | · |
| - Indicator (downriver) ¹ | 2.205 | 2.597 | + 17.8% |
| - Control (upriver) ² | 2.113 | 2:230 | + 5.6% |
| Beryllium-7 in Sediment | | | |
| - Indicator (at Vogtle) ³ | 930.5 | 1297.8 | + 39.5% |
| - Control (off site) ⁴ | 578.3 | 1229.8 | +112.7% |
| Cobalt-60 in Sediment | | · · · · · | |
| - Indicator (at Vogtle) ³ | 51:33 | 138.3 | +169.5% |
| Cesium-137 in Sediment | , | ·. | |
| - Indicator (at $Vogtle$) ³ | 192.3 | 264.2 | + 37.4% |
| - Control (off site) ⁴ | 137.8 | 112.5 | - 18.3% |
| Tritium in River Water, avg. 6 sites | 744.9 | 1077.3 | + 44.6% |

¹Beaufort/Jasper County Water Treatment Plant, Beaufort SC, 112 mi downriver, plus Cherokee Hill Water Treatment Plant, Port Wentworth SC, 122 mi. downriver. ²Augusta Water Treatment Plant, Augusta GA, 56 mi. upriver. ³Savannah River, 0.8 mi. ENE of Vogtle plant. ⁴Savannah River, 2.5 mi. N of Vogtle plant. Beta and tritium in picocuries per liter, others in picocuries per kilogram dry. Source: Vogtle Electric Generating Plant Annual Radiological Environmental Operating Report for 2005, www.nrc.gov.

Table 4

Change in Cancer Death Rate, Burke County vs. U.S. 1987-1990 to 1991-2003

Age 0-24

| <u>County</u> Burke GA United States | <u>Cancer Deaths</u> <u>'87-90</u> <u>'91-03</u> 1 5 | <u>Population 0-24</u> <u>'87-90</u> <u>'91-03</u> 36207 116431 | Deaths/ '87-90 2.76 4.33 | <u>100000</u> <u>'91-03</u> 4.29 3.72 | <u>% Ch</u> +55.5% - 14.1% |
|--|--|---|-----------------------------------|--|----------------------------------|
| Age 25-54 | | | | | |
| · | Cancer Deaths | Population 25-54 | Deaths/ | 100000 | |
| County / | '87-90 '91-03 | '87-90 '91-03 | '87-90 | '91-03 | % Ch |
| Burke GA | 15 84 | 30919 111666 | 48.5 | 75.2 | +55.1% p<.01 |
| United States | , | | 59.0 | 57.3 | - 2.9% |

Source: National Center for Health Statistics, <u>http://wonder.cdc.gov</u>, underlying cause of death. Uses codes for all cancers; ICD-9 codes 140.0-239.9 (until 1998), and ICD-10 codes C00-D48.9 (after 1998).

Table 5

Change in Infant Death Rate (under 1 year) Burke County vs. U.S., 1985-1987 to 1988-1990

| Rate/100000 (Deaths) | | | | | | |
|----------------------|----------------|----------------|------------|-------|--|--|
| County | 1985-87 | <u>1988-90</u> | % Ch. Rate | | | |
| Burke GA | 13.71 (16) | 23.31 (28) | +70.1% | p<.06 | | |
| U.S. | 10.36 (117329) | 9.66 (116916) | - 6.8% | | | |

Source: National Center for Health Statistics (http://wonder.cdc.gov, underlying cause of death).

Table 6

Change in Infant Death Rate (under 1 year), by Race Burke County vs. U.S., 1979-1987 to 1988-2003

| | United States | Burke County | |
|-----------|-----------------------|-----------------------|--------|
| Race | Rate/100,000 (Deaths) | Rate/100,000 (Deaths) | |
| All | | · · · | |
| 1979-1987 | 1132.9 (374433) | 1164.4 (42) | |
| 1988-2003 | 799.0 (507571) | 1382.2 (86) | |
| % Change | - 29.5% | +18.7% | p<.002 |
| White | | | |
| 1979-1987 | 973.7 (260243) | 455.9 (6) | |
| 1988-2003 | 661.1 (330750) | 744.6 (17) | |
| % Change | - 32.1% | +63.3% | p<.03 |
| Black | | | |
| 1979-1987 | 2017.9 (104439) | 1576.9 (36) | |
| 1988-2003 | 1549.9 (159045) | 1755.3 (69) | |
| % Change | - 23.2% | +11.3% | p<.04 |

Source: National Center for Health Statistics (http://wonder.cdc.gov, underlying cause of death).

Table 7

Change in Death Rate, All Ages Combined, by Race Cancer and All Other Causes of Death, 1987-1990 to 1991-2003 Burke County vs. U.S.

| | <u>Burke I</u> | <u>Deaths</u> | Burke I | Rate/1000 | US Rate | e/100000 | <u>% Ch I</u> | Rate |
|----------------------|----------------|----------------|----------------|---------------|---------------|----------------|---------------|-------------|
| Area | <u> *87-90</u> | <u> •91-03</u> | <u> *87-90</u> | <u>•91-03</u> | <u>*87-90</u> | <u> '91-03</u> | Burke | <u>U.S.</u> |
| Cancer -All | 135 | 570 | 185.0 | 231.5 | 216.6 | 207.6 | +25.1 | - 4.2 |
| Cancer - White | 73 | 310 | 190.3 | 223.5 | 212.4 | 204.5 | +17.5 | - 3.7 |
| Cancer – Black | 62 | 260 | 185.0 | 241.7 | 277.3 | 261.6 | +30.7 | - 5.7 |
| | | | | <u>_</u> | | | | |
| Other Causes – All | 701 | 2317 | 971.6 | 929.1 | 741.5 | 675.8 | - 4.4 | - 8.9 |
| Other Causes – White | 319 | 1130 | 894.3 | 863.0 | 717.3 | 655.7 | - 3.5 | - 8.6 |
| Other Causes – Black | 382 | 1186 | 1083.3 | 1042.1 | 988.9 | 894.2 | - 3.8 | - 9.6 |

Source: National Center for Health Statistics, <u>http://wonder.cdc.gov</u>, underlying cause of death. Uses codes for all cancers; ICD-9 codes 140.0-239.9 (until 1998), and ICD-10 codes C00-D48.9 (after 1998). Burke County cancer rate increases are significant at p < .00001 (all), p < .02 (white), and p < .0002 (black). Rates adjusted to 2000 U.S. standard population.

Table 8

Demographic Characteristics, Burke County vs. U.S.

| <u>Category</u> | <u>Burke</u> | <u>U.S.</u> |
|-------------------------|--------------|-------------|
| 2005 est. population | 23299 | 296M |
| 2000 population | 22243 | 281M |
| 2003 % below poverty | 21.7 | 12.5 |
| 2004 % black | 51.1 | 12.8 |
| 2004 % Hispanic | 1.6 | 14.1 |
| 2004 % Asian | 0.3 | 4.2 |
| 2000 % High School grad | 64.9 | 80.4 |
| 2000 % College grad | 9.5 | 24.4 |

Note: Percent high school and college graduates are for adults over age 25. Source: U.S Bureau of the Census, <u>www.census.gov</u>, your gateway to the 2000 census, state and county quick facts.

6