

### Subsection 2.5.3 Table of Contents

<u>Section</u>	<u>Title</u>	<u>Page</u>
2.5.3	Surface Faulting .....	2.5.3-1
2.5.3.1	Geological, Seismological, and Geophysical Investigations .....	2.5.3-2
2.5.3.2	Geological Evidence, or Absence of Evidence, for Surface Deformation .....	2.5.3-4
2.5.3.3	Correlation of Earthquakes with Capable Tectonic Sources .....	2.5.3-7
2.5.3.4	Ages of Most Recent Deformation .....	2.5.3-8
2.5.3.5	Relationship of Tectonic Structures in the Site Area to Regional Tectonic Sources .....	2.5.3-13
2.5.3.6	Characterization of Capable Tectonic Sources .....	2.5.3-13
2.5.3.7	Designation of Zones of Quaternary Deformation in the Site Region .....	2.5.3-13
2.5.3.8	Potential for Tectonic or Non-Tectonic Deformation at the Site .....	2.5.3-14
2.5.3.9	References .....	2.5.3-15

**VCS COL 2.0-28-A**

**2.5.3 Surface Faulting**

This section evaluates the potential for tectonic surface deformation and non-tectonic surface deformation at the VCS site. Information contained in this section is developed in accordance with RG 1.208 and is intended to demonstrate compliance with 10 CFR 100.23, *Geologic and Seismic Siting Criteria*. Specifically, this subsection addresses the following issues:

- Potential surface deformation associated with active tectonism, including capable seismic sources (faults) and volcanism
- Potential surface deformation associated with growth faults, which do not extend into the crystalline basement beneath the Coastal Plains and, as described in Appendix C of RG 1.208, are not considered to be potential tectonic sources of strong vibratory ground motion
- Potential surface deformation associated with non-tectonic processes such as collapse structures (karst collapse), subsurface salt migration (salt domes), and man-induced deformation (e.g., mining collapse, subsidence due to fluid withdrawal)

In summary, there are no capable faults within the 200-mile VCS site region (see [Figures 2.5.1-222](#) and [2.5.1-225](#)), and there is negligible potential for tectonic fault rupture within 25 miles of the site. The VCS site lies within the regionally extensive Tertiary age Vicksburg growth fault zone along the Texas Gulf Coast ([Figure 2.5.1-202a](#)). Detailed studies of the site area described in [Subsection 2.5.1.2.4.2](#) provide evidence for the absence of active growth faults whose surface projections lie within the footprint of the VCS plant. These studies further document evidence for localized Quaternary surface deformation above two growth faults, referred to as fault D (or fault GM-D) and fault E (or fault GM-E), within the site area (see [Figure 2.5.1-240](#)). The closest approach of the zone of Quaternary surface deformation associated with fault D to Seismic Category 1 structures is 787 feet (240 meters), and the closest approach of the zone of Quaternary surface deformation to remaining standard plant design structures is 731 feet (223 meters) ([Figure 2.5.1-243](#)) (see [Figure 2.5.1.2.4.2.3.2](#)). The closest approach of the zone of Quaternary surface deformation associated with fault E to Seismic Category 1 structures is approximately 2.6 miles. Detailed studies, including analysis of seismic reflection and LiDAR data, characterize the location, width, and long-term average rate of activity of the more proximal zone of surface deformation associated with fault D, and further indicate that any

potential future surface deformation due to growth fault activity will be confined to this zone and thus will not affect Units 1 and 2 Seismic Category 1 structures. There is no potential for volcanic activity within the site area or vicinity (see [Subsection 2.5.1.2.5](#)). There is no potential for other non-tectonic surface deformation within the site area due to processes such as karst collapse, salt migration, or glacial rebound. Based on available information on settlement due to withdrawal of petroleum and/or groundwater in the site vicinity, there is little risk of surface deformation from this source.

The following sections present the data, observations, and references to support these conclusions.

#### 2.5.3.1 **Geological, Seismological, and Geophysical Investigations**

Available information regarding the potential for permanent surface deformation at the VCS site is documented in several primary sources:

- Geologic mapping published by the U.S. Geological Survey (USGS), the state of Texas, and other researchers (e.g., [References 2.5.3-201](#), [2.5.3-202](#), [2.5.3-203](#), [2.5.3-204](#), and [2.5.3-205](#))
- Articles published by various researchers in referenced journals and field trip guidebooks (e.g., [References 2.5.3-202](#), [2.5.3-206](#), [2.5.3-207](#), [2.5.3-208](#), and [2.5.3-209](#))
- Seismicity data compiled and analyzed in published research (e.g., [References 2.5.3-210](#), [2.5.3-211](#), and [2.5.3-212](#))

In addition to reviewing this existing information, the following investigations were performed to assess the potential for tectonic and non-tectonic deformation within the 5-mile VCS site area

- Interpretation of aerial photography and remote sensing imagery
- Analysis and interpretation of seismic reflection data
- Detailed analysis of high resolution, LiDAR-derived topographic data
- Aerial and field reconnaissance
- Review of pre- and post-EPRI ([Reference 2.5.3-223](#)) seismicity (see [Subsection 2.5.2](#))
- Discussions with current researchers in the area

#### 2.5.3.1.1 Previous Site Investigations

The VCS site is a greenfield site with no existing nuclear facilities or other facilities for which a safety analysis or surface faulting hazard evaluation was performed. The VCS site lies within the Gulf Coastal Plains physiographic province, which has been the subject of regional geologic investigations. Previous geologic studies relevant to the VCS site, and in particular to the potential for permanent ground deformation, were reviewed as part of this COLA investigation and are summarized in [Subsection 2.5.1](#).

#### 2.5.3.1.2 Regional and Local Geological Studies

The USGS completed a compilation of all active or potentially active Quaternary features (e.g., faults, liquefaction features, young folds, and uplifts) in the central and eastern United States, including the Gulf Coastal Plains region (e.g., [References 2.5.3-213](#), [2.5.3-214](#), [2.5.3-215](#), [2.5.3-216](#), [2.5.3-217](#), and [2.5.3-218](#)). These compilations do not identify any Quaternary tectonic faults, features, or other evidence of tectonic activity within 25 miles (40 km) of the VCS site (see [Subsection 2.5.1.1.4.3.5](#)).

As described in [Subsection 2.5.1.1.4.3.5.3](#), evidence for Quaternary activity in the form of non-tectonic surface deformation has been documented on some growth faults in the Texas Coastal Plains. As noted by Wheeler ([Reference 2.5.3-219](#), Synopsis):

“The gulf-margin normal faults in Texas are assigned as Class B structures because [of] their low seismicity and because they may be decoupled from underlying crust, making it unclear if they can generate significant seismic ruptures that could cause damaging ground motion.”

The definition of a Class B structure, per USGS criteria ([Reference 2.5.3-215](#)), is:

“Class B: Geologic evidence demonstrates the existence of Quaternary deformation, but either (1) the fault might not extend deeply enough to be a potential source of significant earthquakes, or (2) the currently available geologic evidence is too strong to confidently assign the feature to Class C but not strong enough to assign it to Class A.”

In contrast, a Class A structure exhibits “geologic evidence (that) demonstrates the existence of a Quaternary fault of tectonic origin,

whether the fault is exposed by mapping or inferred from liquefaction or other deformational features” (Reference 2.5.3-215).

The assessment of the USGS (Reference 2.5.3-219) is consistent with studies published since the 1986 EPRI (Reference 2.5.3-223) study (see description in Subsection 2.5.1.2.4.2) that growth faults are shallow upper crustal features confined to the Coastal Plains section and do not extend into the crystalline basement. As described in Subsection 2.5.1.2.4.2, there is a consensus in the informed technical community that the sediments involved in growth faulting do not have the requisite elastic strength to store sufficient strain energy for release in moderate to large earthquakes. The USGS assessment (Reference 2.5.3-219) is consistent with the guidance provided in Section C.2.4 of Appendix C, NRC RG 1.208, which characterizes growth faults as non-tectonic and hence non-capable structures:

“Non-tectonic structures are those found in karst terrain and those resulting from growth faulting. . . Large, naturally occurring growth faults such as those found in the coastal plain of Texas and Louisiana can pose a surface displacement hazard, even though offset most likely occurs at a much less rapid rate than that of tectonic faults. They are not regarded as having the capacity to generate damaging vibratory ground motion, can often be identified and avoided in siting, and their displacements can be monitored.”

Geologic mapping and analyses of subsurface data relevant to identifying and characterizing growth fault activity in the site vicinity is presented in Subsection 2.5.1.2.4.2.

### **2.5.3.2 Geological Evidence, or Absence of Evidence, for Surface Deformation**

#### **2.5.3.2.1 Bedrock Faults**

As shown on Figure 2.5.1-204, no bedrock faults have been mapped within the site area. The entire site vicinity is located in the Texas Gulf Coastal Plains, and the site area specifically is directly underlain by approximately 41,000 feet (equivalent to 7.8 miles or 12.5 kilometers) of Mesozoic and Cenozoic strata deposited within the Gulf of Mexico Basin (Figure 2.5.1-226) (see Subsection 2.5.1.1.4.1.3). Analysis of time-migrated seismic reflection profiles (Figures 2.5.1-245 through 2.5.1-247) for this COLA documents that all growth faults in the site area that displace Coastal Plains strata in the upper 4000 to 8000 feet

(1200 to 2400 meters) of the sedimentary section terminate downward against or above a sub-horizontal detachment horizon within the Coastal Plains section ([Subsection 2.5.1.2.4.2.3.1](#)). This observation, along with other stratigraphic and structural relationships that document primary activity of the faults during deposition of the Oligocene Vicksburg Formation, is consistent with the interpretation that the faults are part of the Vicksburg zone of non-tectonic and non-capable growth faults.

#### 2.5.3.2.2 Growth Faults

##### 2.5.3.2.2.1 Previous Studies

As described in [Subsection 2.5.1.2.4.2.3](#), previously published studies and proprietary data documented the presence of five growth faults within the VCS site area. From southeast to northwest, the faults are designated: GM-E, GM-D, GM-K, GM-L, and GM-N ([Figure 2.5.1-240](#)). These faults are part of the well-documented Vicksburg growth fault zone ([Subsection 2.5.1.2.4.2](#)), which formed and was primarily active in Oligocene time. The growth faults are confined to the Mesozoic and Cenozoic Gulf Coastal Plains stratigraphic section, and do not extend into the underlying crystalline basement. Existing proprietary subsurface mapping ([Reference 2.5.3-220](#)) has documented that growth faults GM-E, GM-D, GM-K, GM-L, and GM-N deform marker horizons in the Oligocene-Miocene Frio Formation stratigraphic units overlying the Vicksburg Formation at depths as shallow as about 4000 to 6000 feet (1219 to 1829 meters) ([Subsection 2.5.1.2.4.2.1.2](#)). The youngest stratigraphic horizons documented by Geomap ([Reference 2.5.3-220](#)) are within the Oligocene-Miocene Frio Formation, and thus the Geomap ([Reference 2.5.3-220](#)) dataset provides no evidence to assess activity or non-activity of these growth faults during deposition of sediments younger than the Frio Formation.

##### 2.5.3.2.2.2 Current Investigations

[Subsection 2.5.1.2.4.2.3](#) presents compiled mapping and subsurface data that document the location and geometry of growth faults within the site area. In addition to this compilation effort, an array of four seismic reflection profiles were licensed by Exelon to: (1) document the presence or absence of other growth faults not included in the compilation mapping; (2) elucidate the subsurface geometry of the faults; and (3) assess the stratigraphic and structural relations that document activity or non-activity of the faults. As described in [Subsection 2.5.1.2.4.2.1.5](#),

these reflection profiles were processed and interpreted using modern, standard-of-practice techniques. Also, new analyses of air photos and LiDAR data, as well as aerial and field reconnaissance, were performed for this COLA project to assess whether anomalous topographic features commonly associated with active growth faults exist within the site area and are correlated with growth faults identified in the subsurface. The analyses are described in detail in [Subsection 2.5.1.2.4.2.3](#) and are summarized in the following subsections. Observations and interpretations developed from these analyses that are relevant to assessing activity or non-activity of growth faults in the site area are presented in [Subsection 2.5.3.4.2.1](#).

#### 2.5.3.2.2.2.1 Seismic Reflection Data

Seismic reflection data was used to investigate the subsurface geology of the site area, with the specific goal of identifying and mapping growth faults, documenting their geometry, and analyzing structural and stratigraphic relationships that constrain the timing of their last activity. As described in [Subsection 2.5.1.2.4.2.1.5](#), a total of four 2-D seismic reflection profiles were selected to provide coverage of the site area. The data was licensed from a private broker, processed, and analyzed using industry standard software and techniques.

Based on inspection and interpretation of the time-migrated seismic reflection profiles, the site area is underlain by a series of southeast-dipping listric growth faults, most of which sole into a main detachment horizon at a depth of approximately 3.9 to 4.5 seconds two-way travel time. If it is assumed that the average acoustic velocity for the imaged Gulf Coastal Plains strata is approximately 6000 feet (1829 meters) per second ([Reference 2.5.3-221](#)), then the depth to the main detachment is approximately 11,700 to 13,500 feet (3400 to 4100 meters), confirming that the growth-fault system is confined to the upper part of the Coastal Plains stratigraphic section. Through comparison with proprietary Geomap ([Reference 2.5.3-220](#)) data, several of the faults interpreted in the seismic data were correlated with previously recognized growth faults GM-E, GM-D, GM-K, GM-L, and GM-N ([Figures 2.5.1-245, 2.5.1-246, 2.5.1-247](#)). Several other growth faults not identified in the Geomap ([Reference 2.5.3-220](#)) compilation also were recognized and mapped on the seismic lines (e.g., SR-01 and SR-03). In addition to growth faults, four distinct and laterally correlative stratigraphic marker horizons were identified and mapped in the

subsurface among all four seismic lines. From oldest to youngest, the marker horizons were designated Horizon 1 (interpreted to be middle Oligocene in age), Horizon 2 (interpreted to be Oligocene to Miocene in age), Horizon 3 (interpreted to be early Pliocene in age), and Horizon 4 (interpreted to be Pliocene to Quaternary in age). The stratigraphic markers were used to assess and document the timing of last activity on the growth faults.

#### 2.5.3.2.2.2.2 LiDAR Derived Topography

As described in [Subsection 2.5.1.2.4.2.1.4](#), LiDAR technology provides extremely detailed imaging of topography. LiDAR data of the Texas coastal regions, including the majority of the VCS site vicinity and all of the site area, were acquired in 2007 by the Texas Natural Resources Information System, in conjunction with the Federal Emergency Management Agency. The LiDAR data for the majority of the site vicinity was analyzed in this site characterization investigation and used to develop high-resolution, shaded relief images of the site vicinity and site area (e.g., [Figures 2.5.1-237](#), [2.5.1-238](#), [2.5.1-239](#), [2.5.1-240](#), [2.5.1-241](#), [2.5.1-242](#), [2.5.1-243](#), and [2.5.1-249](#)). The images were analyzed specifically for lineaments and other anomalous topographic features, in particular those spatially associated with the surface projections of growth faults (see [Subsection 2.5.1.2.4.2.2](#)).

#### 2.5.3.2.2.2.3 Stereo Aerial Photography

Stereo-paired aerial photographs (1:20,000 scale), obtained from the United States Department of Agriculture Farm Service Agency, were used to identify topographic, tonal, and vegetation lineaments and assess their spatial relationship, if any, to the surface projections of growth faults within the site area. Features observed in the photos are described in [Subsection 2.5.1.2.4.2.3](#) and include lineaments associated with topographic features and linear tonal and vegetation changes ([Figure 2.5.1-237](#)). In several cases, the photo lineaments are coincident with topographic lineaments identified within the LiDAR data ([Figure 2.5.1-242](#)).

### 2.5.3.3 Correlation of Earthquakes with Capable Tectonic Sources

As part of this COL application, a comprehensive seismicity catalog was compiled for the greater site region (see [Subsection 2.5.2.1](#)). The seismicity catalog contains no earthquakes with  $m_b \geq 3.0$  within 25 miles (40 km) of the VCS site ([Figures 2.5.1-211](#) and [2.5.1-225](#)), and thus there

is no spatial correlation of earthquake epicenters with known faults, postulated faults, tectonic features, or geomorphic features within the site vicinity. In addition, no reported historical earthquake epicenters have been associated with buried bedrock faults, growth faults, or other geologic features within the site vicinity ([Figures 2.5.1-211](#) and [2.5.1-225](#)) ([References 2.5.3-210](#), [2.5.3-211](#), and [2.5.3-212](#)).

#### 2.5.3.4 **Ages of Most Recent Deformation**

##### 2.5.3.4.1 **Bedrock Faults**

As summarized in [Subsection 2.5.1.1.4.1](#), the most recent bedrock deformation within the Texas Gulf Coastal Plains occurred during the Mesozoic Era and is related to rifting that led to development of the Gulf of Mexico basin. As the Gulf of Mexico opened, the basement crust of the Coastal Plains subsided and became part of a passive continental margin. The passive margin crust was progressively buried by Mesozoic and Cenozoic marine deposits of the Gulf of Mexico, and, based on an interpretation of regional gravity data, presently is at a depth of approximately 41,000 feet (equivalent to 7.8 miles or 12.5 kilometers) beneath the VCS site ([Figure 2.5.1-226](#)). The basement underlying the site area is part of this extended continental crust. However, as described in [Subsection 2.5.1.1.4.1.3](#), the locations of basement features or faults involved in this extension and rifting are unknown due to the thick sedimentary deposits overlying the deep basement rocks. The current state of knowledge indicates that there is no evidence for capable bedrock faults within the site area, and there is no evidence of fault activity within the basement rocks of the site area since the Mesozoic, if at all.

##### 2.5.3.4.2 **Growth Faults**

###### 2.5.3.4.2.1 Evidence for Activity and Non-Activity of Growth Faults in the Site Area

As described in detail in [Subsection 2.5.1.2.4.2.3.1.4](#), fault GM-D is the only geologic structure directly imaged by seismic reflection data in the site area that exhibits stratigraphic evidence for late Cenozoic to Quaternary activity. Other subsurface faults (e.g., GM-K, GM-L, GM-M, SR-01, SR-03, SR-04, SR-05, and SR-07) ([Figures 2.5.1-240](#), [2.5.1-245](#) to [2.5.1-247](#)) are overlain by an undeformed Horizon 3 marker, which provides positive evidence for no activity of these faults since early Pliocene time (from about 5 to 4 million years before present).

Based on interpretation of depth-migrated seismic reflection data, fault GM-D displaces the Pliocene-Quaternary stratigraphic marker Horizon 4 approximately 70 feet down to the southeast. Reflectors at extremely shallow depth above the Horizon 4 marker also appear to be deformed, but primarily by distributed southeast-down tilting or folding within an upward-widening triangular zone, rather than by discrete fault offset. It is important to note that the lateral continuity and imaging of reflectors above Horizon 4 is relatively poor, probably due to a combination of near-shore and relatively energetic late Cenozoic depositional processes ([Reference 2.5.3-202](#)) that did not favor the formation of well-defined, laterally continuous horizontal strata, and the fact the seismic data was originally acquired to provide optimum imaging at much greater depths where oil and gas are located. The southeast-down tilting of reflectors above Horizon 4 is interpreted to represent monoclinial fault-propagation folding of the shallow crust in the hanging wall of fault GM-D. As imaged in the seismic line that approaches closest to the plant footprint, the monoclinial fold associated with fault GM-D is interpreted to be about 1600 feet (490 meters) wide in the shallow subsurface. The fold is directly correlated with a LiDAR lineament associated with a distinct southeast-down break in topography, which is also about 1600 feet (490 meters) wide at the ground surface where the seismic line crosses the lineament. Based on these relationships, the LiDAR lineament is interpreted to be a monoclinial fold in the upper surface of the Beaumont Formation related to Quaternary activity of fault GM-D. The closest approach of the zone of monoclinial folding associated with fault GM-D to Seismic Category 1 structures is 787 feet (240 meters), and the closest approach of the zone to remaining standard plant design structures is 731 feet (223 meters) ([Figure 2.5.1-243](#)) (see [Subsection 2.5.1.2.4.2.3.2](#)).

Fault GM-E is not adequately imaged by reflection data to assess activity or non-activity since deposition of the Horizon 3 and Horizon 4 seismic markers. In lieu of direct seismic imaging, activity or non-activity of fault GM-E was assessed through geomorphic analysis. Subtle topography imaged by LiDAR data indicates that a distinctly linear, southeast-facing slope break associated with the updip projection of fault GM-E is present in the upper surface of the middle to late Pleistocene Beaumont Formation. The lineament can be traced to the southwest where it is present in relatively younger deposits and landforms of the San Antonio river valley ([Figure 2.5.1-239](#); see [Subsection 2.5.1.2.4.2.3](#)). These

relationships provide strong evidence that the lineament and associated southeast-facing slope break were not formed by fluvial processes, and that the lineament is clearly post-Beaumont in age where it crosses the San Antonio River valley. Based on these observations, the lineament is interpreted to be associated with local Quaternary activity of growth fault GM-E.

The following sections provide more detailed characterizations of surface deformation associated with faults GM-D and GM-E, and estimates of long-term average activity rates.

#### 2.5.3.4.2.1.1 Fault GM-D

As described in [Subsection 2.5.1.2.4.2.3.1.4](#), fault GM-D is observed to cause down-to-the-southeast displacement of the Horizon 4 marker on all of the seismic profiles ([Figures 2.5.1-245](#), [2.5.1-246](#), [2.5.1-247](#), and [2.5.1-248](#)) ([Table 2.5.1-204](#)), except 2D Reflection Profile "GSI" Time Mitigated ([Figure 2.5.1-246](#)), which crosses the fault at a point where there is no LiDAR or photo lineament indicative of surface deformation. From these relations, it is interpreted that Quaternary activity of fault GM-D is limited to short reaches of the structure that have linear topographic anomalies discernable in LiDAR imagery, and not along the entire fault.

As described in [Subsections 2.5.3.4.2.1](#) and [2.5.1.2.4.2.3.2](#), the pattern of reflectors in the seismic data suggests that post-Horizon 4 activity of fault GM-D has produced distributed down-to-the-southeast monoclinical tilting or folding of strata within an upward-widening triangular zone in the hanging wall of the fault. Detailed topographic profiles extracted from the LiDAR data across the lineament within the site reveal that the slope break is characterized by a generally uniform increase in surface gradient across a horizontal distance that ranges from approximately 300 to 1600 feet (91 to 488 meters) ([Figures 2.5.1-250a](#) through [2.5.1-250c](#)). The broad tilting expressed in the topography is consistent with the interpretation that shallow deformation above Horizon 4 is characterized by monoclinical folding of strata in the hanging wall of fault GM-D. The zone of Quaternary deformation, therefore, is limited to the monoclinical fold in the hanging wall of the fault.

As described in [Subsection 2.5.1.2.4.2.3.2](#), total southeast-down relief on the upper surface of the Beaumont Formation across the monoclinical fold and associated LiDAR lineament at the site is approximately 4 feet (1.2

meters), with minor variability in relief along strike. The slope break is modified in places by post-Beaumont erosion, which locally has reduced the gradient and obscured the monoclinical profile. Even in locations along the LiDAR lineament where the slope break is subdued, however, the southeast-down structural relief on the upper surface of the Beaumont Formation due to activity on the fault can be readily determined from analysis of topographic profiles. Through analysis and interpretation of a series of closely spaced topographic profiles across the topographic lineament associated with fault GM-D, the extent of monoclinical surface deformation above the fault was delineated and mapped, and is shown in [Figure 2.5.1-243](#) (see [Subsection 2.5.1.2.4.2.3.2](#)). The closest approach of this zone of monoclinical folding to Seismic Category 1 structures is 787 feet (240 meters), and the closest approach of the zone to remaining standard plant design structures is 731 feet (223 meters) ([Figure 2.5.1-243](#)) (see [Subsection 2.5.1.2.4.2.3.2](#)).

#### 2.5.3.4.2.1.2 Fault GM-E

Growth fault GM-E is over 2.6 miles (4.2 kilometers) from the center-points of VCS Units 1 and 2, and as such any activity on the fault will not affect the site. Despite this fact, fault GM-E is still a structure of interest because it is the only fault, besides fault GM-D, that exhibits evidence for Quaternary activity within the site area. As described in [Subsection 2.5.1.2.4.2.3.1.3](#), fault GM-E is not imaged in the seismic reflection data because the seismic profiles do not extend far enough south to cross the fault and provide sufficient imaging resolution of the fault at depth. Despite the lack of direct reflection imaging, the distinct topographic lineament apparent in the LiDAR data and its spatial correlation with the surface projection of fault GM-E strongly suggest that fault GM-E has been active in the Quaternary and formed a monoclinical, southeast-facing slope break in the upper surface of the Beaumont Formation, similar to that associated with fault GM-D.

As described in [Subsection 2.5.1.2.4.2.4](#), fault GM-E crosses a variety of features including the deposits of the Beaumont Formation, younger Pleistocene stream terrace and floodplain deposits of the San Antonio River, and man-made features (i.e., FM 445, U.S. Highway 77, and SR 239) ([Figures 2.5.1-204](#) and [2.5.1-239](#)). Field reconnaissance of the fault across these features was unable to provide any refinements on the timing of activity other than that movement has occurred since deposition of the Beaumont Formation and younger sediments of the San Antonio

river valley. Topographic profiles of the fault along FM 445 derived from the LiDAR data reveal that the slope break associated with the fault has the same characteristics as the non-eroded profiles of fault GM-D (e.g., Profiles 4 and 8 in [Figures 2.5.1-250a](#) and [2.5.1-250b](#), respectively): a distinct inflection of the ground surface at the location of the lineament with the southeast side down. For fault GM-E, this step has an approximately 4.9 feet (1.5 meters) offset over 980 feet (300 meters), or equivalently a narrow, steepened region with a slope of approximately 0.29°.

#### 2.5.3.4.2.1.3 Long-Term Average Rates of Activity

Long-term average rates of surface deformation associated with growth fault GM-D were estimated from the age of the deformed upper surface of the Beaumont Formation and the total surface offset. Based on analysis of topographic profiles, the separation of the upper surface of the Beaumont Formation across fault GM-D ranges from approximately 1.5 feet to 4.5 feet (about 0.5 meter to 1.5 meters). As described in [Subsection 2.5.1.2.3](#), the precise age of the Beaumont Formation is uncertain. Current estimates of the age of the Beaumont vary between 100,000 and 350,000 years ([References 2.5.3-201](#), [2.5.3-202](#), [2.5.3-203](#), [2.5.3-205](#), and [2.5.3-222](#)). From the extremes in the range of offsets and ages, the corresponding range in long-term average separation rates across fault GM-D is approximately  $5.1 \times 10^{-5}$  inches per year to  $5.4 \times 10^{-4}$  inches per year (see [Subsection 2.5.1.2.4.2.3.2](#)). If it is assumed that fault GM-D slips continuously and uniformly at these rates, then the maximum down-to-the-southeast displacement of the land surface across the zone of monoclinical tilting in 100 years will be about 1/18th of an inch (see [Subsection 2.5.1.2.4.2.3.2](#)).

Using a similar approach, with topographic relief on the surface of the Beaumont Formation determined from analysis of LiDAR profiles, the range in long-term deformation rates for fault GM-E is  $1.7 \times 10^{-4}$  inches per year to  $5.9 \times 10^{-4}$  inches per year. This vertical relief and implied range of deformation rates are similar to those observed for fault GM-D. The similarities between the two faults could either be coincidental or may suggest that the mechanisms, rates, and characteristics of growth fault activity within the site area are fairly uniform.

#### **2.5.3.5 Relationship of Tectonic Structures in the Site Area to Regional Tectonic Sources**

The only geologic structures in the site area are growth faults associated with the Vicksburg fault zone, which has been mapped for a minimum of 500 miles along trend in the Gulf Coastal Plains (see description in [Subsection 2.5.1.2.4.2.3.2](#)). As described in [Subsection 2.5.3.2.2.1](#), the site area growth faults are confined to approximately the upper 13,000 feet (4000 meters) of the Gulf Coastal Plains stratigraphic section and do not extend into the crystalline basement, which is estimated to lie at a depth of about 41,000 feet (7.8 miles or 12.5 kilometers) beneath the site area, based on interpretation of regional gravity data. As noted in the introduction to [Section 2.5.3](#), Section C.2.4 of Appendix C, RG 1.208 states that growth faults are “non-tectonic” and are “not regarded as having the capacity to generate damaging vibratory ground motion.” Given that growth faults are the only geologic structures present in the site area, it is concluded that there are no tectonic structures in the site area. Furthermore, there is no correlation of geologic structures in the site area to regional, capable tectonic sources.

#### **2.5.3.6 Characterization of Capable Tectonic Sources**

Based on data presented in [Subsection 2.5.1](#) and previous descriptions in [Subsection 2.5.3.4](#), there are no capable tectonic sources within the VCS site area.

#### **2.5.3.7 Designation of Zones of Quaternary Deformation in the Site Region**

There are no zones of Quaternary deformation associated with tectonic faults requiring detailed investigation within the site area.

Interpretation of aerial photography and LiDAR data, coupled with aerial and field reconnaissance, has documented Quaternary surface deformation associated with growth faults GM-D and GM-E in the site area (see [Subsection 2.5.1.2.4.2.3](#)). Detailed analysis of seismic reflection and LiDAR data indicate that displacement on faults GM-D and GM-E can be traced upwards to very shallow depths in the stratigraphic section. Displacement on these structures at depth projects updip to distinct lineaments in the LiDAR data that coincide with zones of mappable southeast-side-down monoclinical tilting of the upper depositional surface of the Quaternary Beaumont Formation. The closest approach of a zone of monoclinical tilting to Seismic Category 1 structures

is 787 feet (240 meters), and the closest approach to remaining standard plant design structures is 731 feet (223 meters) ([Figure 2.5.1-243](#)) (see [Subsection 2.5.1.2.4.2.3.2](#)). The extent of monoclinical surface deformation is discernable from analysis of LiDAR data, and does not require detail investigations of the type used to characterize potential surface rupture on capable tectonic faults.

### **2.5.3.8 Potential for Tectonic or Non-Tectonic Deformation at the Site**

#### **2.5.3.8.1 Potential for Tectonic Deformation**

The potential for tectonic deformation at the site is negligible. There are no capable tectonic faults within the site vicinity.

Based on a review of geologic literature, there is no documented intrusive or extrusive volcanic activity of Tertiary age within the site region, and thus there is no potential for volcanic activity within the site vicinity (see [Subsection 2.5.1.2.5](#)). The youngest mapped volcanic and intrusive rocks in the site region are Mesozoic in age, and they primarily crop out to the north and northwest of the site at distances of 60 miles or greater ([Figures 2.5.1-212, 2.5.1-202a, and 2.5.1-202b](#)).

#### **2.5.3.8.2 Potential for Non-Tectonic Deformation**

##### **2.5.3.8.2.1 Growth Faults**

The potential for non-tectonic deformation within the VCS site from movement on growth faults is confined to a narrow zone associated with fault GM-D, as shown in [Figure 2.5.1-249](#) and [Figure 2.5.1-243](#) (see [Subsection 2.5.1.2.4.2.3](#)). Any potential future activity on fault D will not extend beneath the footprint of Seismic Category 1 structures at the VCS site.

##### **2.5.3.8.2.2 Other Potential Sources of Non-Tectonic Deformation**

There is no evidence of non-tectonic deformation at the site in the form of glacially induced faulting, collapse structures, salt migration, mining, or subsidence due to petroleum or groundwater extraction:

- All documented faulting within the site vicinity is caused by growth faults, the activity of which is likely related to sediment compaction, dewatering, flow of salt in the subsurface, or fluid withdrawal. There are no documented examples of glacially induced faulting in the site region.

- There are no deposits of limestone or other carbonate rocks at shallow depths within the site area that pose potential karst collapse and surface subsidence hazards.
- No piercement-type salt domes are located within the site vicinity (see [Subsection 2.5.1.1.4.3.4.1](#)).
- There are no mining activities within the site area that may produce man-induced surface collapse.
- There is no potential for subsidence due to withdrawal of petroleum and groundwater, as described in [Subsection 2.5.1.2.6.4](#). Further descriptions supporting this statement is found in [Subsections 2.4.12](#) and [2.5.4](#).

#### 2.5.3.9 References

- 2.5.3-201 Barnes, V.E., *Geologic Atlas of Texas Beeville-Bay City Sheet*, Bureau of Economic Geology, University of Texas, 1987.
- 2.5.3-202 Blum, M. D., and Aslan, A., "Signatures of Climate vs. Sea-Level Change within Incised Valley-Fill Successions: Quaternary Examples from the Texas Gulf Coast," *Sedimentary Geology*, v. 190, p. 177-211, 2006.
- 2.5.3-203 Dubar, J.R., Ewing, T., Lundelius, E.L., Otvos, E.G., and Winker, C.D., "Quaternary Geology of the Gulf of Mexico Coastal Plain," *Geology of North America*, v. K-2, *Quaternary Nonglacial Geology: Conterminous U.S.*, Geological Society of America, p. 583-610, 1991.
- 2.5.3-204 King, P.B., and Beikman, H.M., *Geologic Map of the United States (Exclusive of Alaska and Hawaii)*, U.S. Geological Survey, 1974.
- 2.5.3-205 Winker, C.D., *Late Pleistocene Fluvial-Deltaic Deposition: Texas Coastal Plain and Shelf* [MA thesis], University of Texas, 1979.
- 2.5.3-206 Collins, E., Hobday, D., and Kreitler, C., *Quaternary Faulting in East Texas*, Geological Circular 80-1, Bureau of Economic Geology, University of Texas, 1980.
- 2.5.3-207 Dodge, M.M., and Posey, J.S., *Structural Cross Sections, Tertiary Formations, Texas Gulf Coast*, 32 plates, Bureau of Economic Geology, University of Texas, 1981.

- 2.5.3-208 Galloway, W.E., Liu, X., Travis-Neuberger, D., and Xue, L., *Reference High-Resolution Correlation Cross Sections, Paleogene Section, Texas Coastal Plain*, Bureau of Economic Geology, University of Texas, 1994.
- 2.5.3-209 Kreitler, C., *Lineations and Faults in the Texas Coastal Zone*, Report of Investigations No. 85, p. 32., Bureau of Economic Geology, University of Texas, 1976.
- 2.5.3-210 Davis, D.M., Pennington, W., and Carlson, S., "Historical Seismicity of The State of Texas: A Summary," *Gulf Coast Association of Geological Societies Transactions*, v. 35, p. 39-44, 1985.
- 2.5.3-211 Davis, S.D., Pennington, W.D., and Carlson, S.M., *A Compendium of Earthquake Activity in Texas*, Geological Circular 89-3, Bureau of Economic Geology, University of Texas, 1989.
- 2.5.3-212 Frohlich, C., and Davis, S.D., *Texas Earthquakes*, 275 p., University of Texas Press, 2002.
- 2.5.3-213 Crone, A.J., and Wheeler, R.L., *Data for Quaternary Faults, Liquefaction Features, and Possible Tectonic Features in the Central and Eastern United States, East of the Rocky Mountain Front*, Open-File Report 00-260, U.S. Geological Survey, 2000.
- 2.5.3-214 U.S. Geological Survey and Texas Bureau of Economic Geology, *Quaternary Fault and Fold Database for the United States*, 2006, Available at: <http://earthquake.usgs.gov/regional/qfaults/>, accessed on November 26, 2007.
- 2.5.3-215 Wheeler, R.L., *Known or Suggested Quaternary Tectonic Faulting, Central and Eastern United States—New and Updated Assessments for 2005*, Open-File Report 2005-1336, U.S. Geological Survey, 2005.
- 2.5.3-216 Wheeler, R.L., "Quaternary Tectonic Faulting in the Eastern United States," *Engineering Geology*, v. 82, p. 165-186, 2006.
- 2.5.3-217 Wheeler, R.L., "Paleoseismic Targets, Seismic Hazard, and Urban Areas in the Central and Eastern United States," *Bulletin of the Seismological Society of America*, v. 98, p. 1572-1580, 2008.

- 2.5.3-218 Wheeler, R.L., and Crone, A.J., "Known and Suggested Quaternary Faulting in the Midcontinent United States," *Engineering Geology*, v. 62, p. 51-78, 2001.
- 2.5.3-219 Wheeler, R.L., "Fault Number 924, Gulf-Margin Normal Faults, Texas," *Quaternary Fault and Fold Database of the United States*, 1999, Available at <http://earthquakes.usgs.gov/regional/qfaults>, USGS, accessed January 11, 2007.
- 2.5.3-220 Geomap, *Upper Texas Gulf Coast Mapping Service Maps 327, 328, 331, 260, 262, and 263*, 2007. Licensed from Geomap Company to William Lettis and Associates, Inc.
- 2.5.3-221 Bally, A.W., "Seismic Expression of Structural Styles," *AAPG Studies in Geology*, v. 1-3, American Association of Petroleum Geologists, 1983.
- 2.5.3-222 Blum, M., and Price, D.M., *Quaternary Alluvial Plain Construction in Response to Glacio-Eustatic and Climatic Controls, Texas Gulf Coastal Plain, Relative Role of Eustacy, Climate, And Tectonism in Continental Rocks*, Special Publication 59, p. 31-48, Society for Sedimentary Geology, 1998.
- 2.5.3-223 Electric Power Research Institute, *Seismic Hazard Methodology for the Central and Eastern United States*, Report NP-4726, 1986.