MACROINVERTEBRATE ASSESSMENT OF MAYO CREEK NEAR THE VC SUMMER NUCLEAR STATION OPERATED BY SOUTH CAROLINA ELECTRIC AND GAS COMPANY, FAIRFIELD COUNTY, SOUTH CAROLINA

JULY 2008

Submitted To:

SOUTH CAROLINA ELECTRIC AND GAS COMPANY
Fairfield County, South Carolina

CARNAGEY BIOLOGICAL SERVICES, LLC

MACROINVERTEBRATE ASSESSMENT OF MAYO CREEK NEAR THE VC SUMMER NUCLEAR STATION OPERATED BY SOUTH CAROLINA ELECTRIC AND GAS COMPANY, FAIRFIELD COUNTY, SOUTH CAROLINA

JULY 2008

Submitted To:

SOUTH CAROLINA ELECTRIC AND GAS COMPANY
Fairfield County, South Carolina

Submitted by:

CARNAGEY BIOLOGICAL SERVICES, LLC
636 Westwood Drive
Lexington, South Carolina
803-233-6952

SCDHEC Laboratory Certification No. 32572

TABLE OF CONTENTS

		Page
	LIST OF TABLES	ii
	LIST OF FIGURES	iii
I	SUMMARY	1
II	INTRODUCTION	2
III	DESCRIPTION OF STUDY AREA	2
IV	METHODS	4
	A. Field Sampling	4
	B. Water Chemistry	4
	C. Sample Processing	4
	D. Data Analysis	4
V	RESULTS	7
	A. Physicochemical Analysis	7
	B. Macroinvertebrate Community Analysis	7
VI	DISCUSSION	9
VII	REFERENCES	10

LIST OF TABLES

Table		Page
1	Procedures used in the calculation of selected metrics used in this report.	6
2	Physicochemical data collected in conjunction with the macroinvertebrate assessment of Mayo Creek near the effluent discharge of the VC Summer Nuclear Station, Fairfield County, South Carolina, 18 July 2008.	7
3	Macroinvertebrates, their North Carolina biotic index tolerance values (TV), functional feeding groups (FG), and abundance collected from Mayo Creek near the effluent discharge of the VC Summer Nuclear Station, Fairfield County, South Carolina, 18 July 2008.	11
4	Rapid bioassessment metrics calculated for the three sampling stations on Mayo Creek near the effluent discharge of the VC Summer Nuclear Station, Fairfield County, South Carolina, 18 July 2008.	14
5	Dominant taxa (>5% of the collection) for the three sampling stations on Mayo Creek near the effluent discharge of the VC Summer Nuclear Station, Fairfield County, South Carolina, 18 July 2008.	15
6	Habitat assessment scores determined in conjunction with the macroinvertebrate assessment for the three sampling stations on Mayo Creek near the effluent discharge of the VC Summer Nuclear Station, Fairfield County, South Carolina, 18 July 2008.	16

LIST OF FIGURES

Figure		Page
1	Sampling locations for benthic macroinvertebrates collected from Mayo Creek near the VC Summer Nuclear Station, operated by SOUTH CAROLINA ELECTRIC AND GAS COMPANY, Fairfield County, South Carolina.	3

I. SUMMARY

On July 18, 2008, CARNAGEY BIOLOGICAL SERVICES, LLC (SCDHEC Laboratory Certification Number 32572) conducted a benthic macroinvertebrate community assessment on Mayo Creek, near the VC Summer Nuclear Station, operated by SOUTH CAROLINA ELECTRIC AND GAS COMPANY. The objective of this assessment was to determine the condition of the stream's macroinvertebrate community.

Results of the benthic macroinvertebrate assessment indicated the stream's macroinvertebrate community was stressed at all three stations,. This may be due to the drought conditions at the time of sampling. Station 2 had a SCDHEC rating of "good-fair", while Stations 2 and 3 had a rating of "fair". The NCBI rating for Stations 1 and 3 was "good-fair", while Station 2 had a value of "good". All three stations shared similar taxa richness, number of specimens, EPT indices, and EPT abundance.

The water chemistry data measured in conjunction with the macroinvertebrate assessment similar temperature, pH dissolved oxygen, and conductivity at all three stations. All parameters monitored were within water quality standards for Class FW waters of the State of South Carolina (SCDHEC, 1998).

II. INTRODUCTION

On July 18, 2008, a benthic macroinvertebrate community assessment was conducted on Mayo Creek near the VC Summer Nuclear Station, Fairfield County, South Carolina.

III. DESCRIPTION OF STUDY AREA

Collections of aquatic macroinvertebrates were made from three sampling locations in Mayo Creek near the VC Summer Nuclear Station (Figure 1).

Station 1 was located approximately 1.5 kilometers upstream of Parr Road below the confluence of a small unnamed tributary. The river at this point was approximately 0.3 to 0.4 meters wide, and less than 0.1 to 0.1 meters deep. The substrate consisted mainly of sand with some gravel, cobble, and boulders, and the canopy provided approximately 70% cover.

Station 2 was located approximately 170 meters upstream of Parr Road. The river at this point was 0.1 to 0.2 meters wide and less than 0.1 to 0.1 meters deep. The substrate consisted mainly of sand, with some cobble and exposed cobble and bedrock. This station had approximately 75% canopy cover.

Station 3 was established approximately 50 meters downstream of Parr Road. The river at this point was 0.2 to 0.3 meters wide and less than 0.1 to 0.1 meters deep. The substrate consisted mainly of sand, with some gravel, cobble, and boulders. This station had approximately 85% canopy cover.

Figure 1. Sampling locations for macroinvertebrates collected from Mayo Creek near the effluent discharge of the VC Summer Nuclear Station, operated by SOUTH CAROLINA ELECTRIC AND GAS COMPANY, Fairfield County, South Carolina.



IV. METHODS

A. Field Sampling

Qualitative collections of aquatic macroinvertebrates were made with a D-frame aquatic dip net, a U.S. Standard No. 30 sieve, and hand picking organisms from substrates with forceps. The multiple habitat approach, where specimens from all available habitats (stream margins, leaf packs, aquatic vegetation, water-soaked logs, and sand deposits) are pooled to form one aggregate sample was utilized as the sampling procedure. Samples were preserved in the field with 70% ethanol. Each sample represented 1.5 hours of sampling effort. Sampling procedures and habitat types were kept similar at each station to enable species and numerical population comparisons between stations. Habitat scores were determined using the *Habitat Assessment Field Data Sheet for Low Gradient Streams* (Barbour *et al.*, 1999).

B. Water Chemistry

Water chemistry parameters measured at each station in conjunction with the macroinvertebrate sampling included temperature, pH, conductivity, and dissolved oxygen.

C. Sample Processing

Upon return to the laboratory, macroinvertebrates were sorted from debris with the aid of a stereomicroscope. The macroinvertebrates were enumerated and identified to the lowest positive taxonomic level with the aide of appropriate microscopic techniques and taxonomic keys. All specimens will be maintained in CARNAGEY BIOLOGICAL SERVICES, LLC voucher collection for five years or placed into the permanent reference collection.

D. Data Analysis

Comparisons of the macroinvertebrate communities were based on the known tolerance levels and life history strategies of the organisms encountered and on changes in taxonomic composition between sampling stations. Changes in taxonomic composition were determined using metrics outlined in Rapid Bioassessment Protocol III of the US EPA's *Rapid Bioassessment Protocols for Use in Streams and Rivers* (Plafkin *et al.* 1989) and SCDHEC's *Standard Operating and Quality Control Procedures for Macroinvertebrate Sampling* (SCDHEC, 1999). These metrics included the following:

1) Taxa richness - The number of different taxa found at a particular location is an indication of diversity. Reductions in community diversity have been positively associated with various forms of environmental pollution, including nutrient loading, toxic substances, and sedimentation (Barbour *et al.*, 1996; Fore *et al.*, 1996; Rosenberg and Resh, 1993; Shackleford, 1988).

2) EPT Index - EPT Index is the number of taxa from the insect orders Ephemeroptera, Plecoptera and Trichoptera found at a station. These three insect orders are considered to be intolerant of adverse changes in water quality, especially temperature and dissolved oxygen, and therefore, a reduction in these taxa is indicative of reduced water quality (Barbour *et al.*, 1996; Lenat, 1988).

3) Chironomidae taxa and abundance - The Chironomidae are a taxonomically and ecologically diverse group with many taxa which are tolerant of various forms of pollution. The chironomids are often the dominant group encountered at impacted or stressed sites (Rosenberg and Resh, 1993).

4) Ratio of EPT and Chironomidae abundance - The relative abundance of these four indicator groups is a measure of community balance. When compared to a reference site, good biotic conditions are reflected in a fairly even distribution among these four groups (Plafkin *et al.*, 1989). The value of this ratio is reduced by impact due to the general reduction of the more sensitive EPT taxa and an increase in the more tolerant chironomid taxa.

5) Ratio of scraper/scraper and filtering collectors - When compared to a reference site, shifts in the dominance of a particular feeding type may indicate a community responding to an over-abundance of a particular food source or toxicants bound to a particular food source (Rosenberg and Resh, 1993).

6) Shredder/total number of specimens collected - When compared to a reference site, reductions in the relative abundance of shredders can indicate changes in the quality or quantity of riparian zone vegetation or the presence of toxic substances bound to organic carbon contained in the leaf and woody material which comprises their food source (Plafkin *et al.*, 1989).

7) Percent contribution of dominant taxon - This measures the redundancy and evenness of the community structure. It assumes a highly redundant community reflects an impaired community because as the more sensitive taxa are eliminated, there is often a significant increase in the remaining tolerant forms (Barbour *et al.*, 1996; Shackleford, 1988).

8) Dominant taxa in common - When compared to a reference site, major shifts in the composition and abundance of dominant taxa can indicate environmental stress (Barbour *et al.*, 1996; Shackleford, 1988).

9) Community loss index (Table 1) - This index measures the loss of taxa between a reference or control station and a study site. It is an index of dissimilarity, with values

increasing as the degree of dissimilarity from the reference station increases (Courtemanch and Davies, 1987; Plafkin *et al.*, 1989).

10) Jaccard coefficient of community similarity (Table 1) - This coefficient represents the degree of similarity in taxonomic composition between two stations in terms of taxon presence or absence. Values range from 0 to 1.0, increasing as the degree of similarity increases (Jaccard, 1912; Plafkin *et al.*, 1989).

11) Sörensen coefficient (Table 1) - This coefficient represents the degree of similarity in taxonomic composition between two stations in terms of taxon presence or absence. Values range from 0 to 1.0, increasing as the degree of similarity increases (Breitenmoser-Würsten and Satori, 1995).

12) North Carolina biotic index (Table 1) - This index utilizes a pollution tolerance value developed over a wide range of conditions and pollution types to assess the amount of impact (North Carolina Department of Environment, Health and Natural Resources, 1997). The values range from 0-10, increasing as water quality decreases. Taxa are designated as Rare (1-2 specimens), Common (3-9 specimens), or Abundant (≥10 specimens) and assigned a 1, 3, or 10 abundance code, respectively, for calculation of the NCBI.

13) SCDHEC bioclassification – Bioclassification is determined by averaging scores for the NCBI and EPT index at each station, then rating sites as "Excellent, Good, Good-Fair, Fair, or Poor" (SCDHEC, 1999).

Table 1. Procedures used in the calculation of selected metrics used in this report.

Metric	**Procedure**
Community Loss Index	CL = d-a/e Where: a = number of taxa common to both samples. d = total number of taxa present in sample A. e = total number of taxa present in sample B.
Jaccard Coefficient of Similarity	JCS = a/a+b+c Where: a = number of taxa common to both samples. b = number of taxa present in sample B but not A. c = number of taxa present in sample A but not B.
Sörensen Coefficient	$C_S = 2a/(d+e)$ Where: a = number of taxa common to both samples. d = the number of taxa present in sample A. e = the number of taxa present in sample B.
North Carolina Biotic Index	NCBI = $TV_i N_i / N$ Where: TV_i = the tolerance for the ith taxon. N_i = the abundance code of the ith taxon. N = sum of abundance codes for all taxa in the sample.

V. RESULTS

A. Physicochemical Analysis

The water chemistry data measured in conjunction with the macroinvertebrate assessment are presented in Table 2. The data reflected similar temperature, pH dissolved oxygen, and conductivity at all three stations.

Table 2. Physicochemical data collected in conjunction with the macroinvertebrate assessment of Mayo Creek near the effluent discharge of the VC Summer Nuclear Station, Fairfield County, South Carolina, 18 July 2008.

	Station		
Parameter	**1**	**2**	**3**
Water Temperature (°C)	22.60	21.34	20.91
pH (SU)	7.17	6.98	7.04
Conductivity (μmhos/cm)	122	123	126
Dissolved Oxygen (mg/l)	6.48	7.20	6.75

B. Macroinvertebrate Community Analysis

A total of 429 specimens representing 48 taxa were collected from Mayo Creek during this assessment. The taxa list, number of specimens, and relative abundance for each taxon are presented in Table 3. Bioassessment metrics for each sampling station are presented in Table 4. Table 5 lists the dominant taxa for each sampling station. Habitat assessment scores are presented in Table 6 for each station.

The sampling effort at Station 1, the upstream control, yielded 151 specimens representing 26 taxa (Table 3). An EPT index of 8 was calculated for this station (Table 4). The Chironomidae were represented by 5 taxa and contributed 16% of the total specimens collected. The NCBI value of 6.17 resulted in a water quality rating of "good-fair" for this station. The SC Bioclassification score of 2.3 indicated a "fair" rating for Station 1. The dominant functional feeding group was the collector-gatherers, which contributed 27% of the collection. The dominant taxon was *Caenis* sp., which contributed 24% of the collection (Table 5).

Station 2 yielded 149 specimens representing 33 taxa (Table 3). An EPT index of 9 was calculated for this station (Table 4). The Chironomidae were represented by 4 taxa and contributed 3% of the total specimens collected. The NCBI value of 5.73 results in a water quality rating of "good" for this station. The SC Bioclassification score of 2.8 indicated a

"good-fair" rating for Station 2. The dominant functional feeding group was the collector-gatherers, which contributed 30% of the collection. The community loss index value of 0.24 and the Sörensen coefficient value of 0.61 indicate this station is similar to Station 1. The dominant taxon was *Caenis* sp., which contributed 22% of the specimens collected (Table 5).

Station 3 yielded 129 specimens representing 26 taxa. An EPT index of 9 was calculated for this station. The Chironomidae were represented by 2 taxa and contributed a total of 2% of the specimens collected. The NCBI value of 5.93 results in a water quality rating of "good-fair" for this station. The SC Bioclassification score of 2.3 indicated a "fair" rating for Station 3. The dominant functional feeding group was the collector-gatherers, which contributed 50% of the collection. The community loss index value of 0.42 and the Sörensen coefficient value of 0.58 indicate this station is similar to Station 1. The community loss index value of 0.62 and the Sörensen coefficient value of 0.58 indicate this station is slightly dissimilar to Station 2. The dominant taxon was *Caenis* sp., which contributed 42% of the specimens collected (Table 5).

VI. DISCUSSION

Results of the benthic macroinvertebrate assessment conducted on Mayo Creek, July 18, 2008, indicated the river's macroinvertebrate community was stressed at all three stations,. This may be due to the drought conditions at the time of sampling. This drought has persisted for the past year. Station 2 had a SCDHEC ratings of "good-fair", while Stations 2 and 3 had a rating of "fair". The NCBI rating for Stations 1 and 3 was "good-fair", while Station 2 had a value of "good". All three stations shared similar taxa richness, number of specimens, EPT indices, and EPT abundance.

The water chemistry data measured in conjunction with the macroinvertebrate assessment similar temperature, pH dissolved oxygen, and conductivity at all three stations. All parameters monitored were within water quality standards for Class FW waters of the State of South Carolina (SCDHEC, 1998).

VII. REFERENCES

Barbour, M.T.; J. Gerritsen; G.E. Griffith; R. Frydenborg; E. McCarron; J.S. White; and M.L. Bastian. 1996. A framework for biological criteria for Florida streams using benthic macroinvertebrates. J. N. Am. Benthol. Soc. 15:185-211.

Barbour, M. T., J. Gerritsen, B. D. Snyder, and J. B. Stribling. 1999. Rapid bioassessment protocols for use in streams and wadeable rivers: periphyton, benthic macroinvertebrates, and fish, Second Edition. EPA 841-B-99-002. U.S. Environmental Protection Agency, Office of Water, Washington, D.C.

Breitenmoser-Würsten, C. and M. Satori. 1995. Distribution, diversity, life cycle and growth of a mayfly community in a prealpine stream (Insecta, Ephemeroptera). Hydrobiol. 308: 85-101.

Courtemanch, D.L. and S.P. Davies. 1987. A coefficient of community loss to assess detrimental change in aquatic communities. Water Research 21:217-222.

Fore, L.S.; J.R. Karr; and R.W. Wisseman. 1996 Assessing invertebrate responses to human activities: evaluation of alternative approaches. J. N. Am. Benthol. Soc. 15:212-232.

Jaccard, P. 1912. The distribution of flora in the alpine zone. New Phytologist 11: 37-50.

Lenat, D.R. 1988. Water quality assessment of streams using a qualitative collection method for benthic macroinvertebrates. J. N. Am. Benthol. Soc. 7: 222-233.

North Carolina Department of Environment, Health and Natural Resources. 1997. Standard operating procedures: biological monitoring. State of North Carolina. Division of Water Quality, North Carolina Department of Environment, Health and Natural Resources, Raleigh, NC, 65 pp.

Plafkin, J.L., M.T. Barbour, K.D. Porter, S.K. Gross and, R.M. Hughes. 1989. Rapid bioassessment protocols for use in streams and rivers. US EPA Assessment and Watershed Protection Division, Washington, D.C. EPA/444/4-89/001.

Rosenberg, D.M. and V.H. Resh (eds.) 1993. Freshwater biomonitoring and benthic macroinvertebrates. Chapman and Hall, New York, New York. 488pp.

Shackleford, B. 1988. Rapid bioassessment of lotic macroinvertebrate communities: Biocriteria development. Biomonitoring Section, Arkansas Dept. Poll. Contl. And Ecol. Little Rock, Ark. 45pp.

South Carolina Department of Health and Environmental Control. 1998. Water classifications and standards (Regulation 61-68), Classified waters (Regulation 61-69) State of South Carolina. Office of Environmental Quality Control, SC DHEC, Columbia, SC.

South Carolina Department of Health and Environmental Control. 1999. Standard Operating and Quality Control Procedures for Macroinvertebrate Sampling. Technical Report No. 004-98. Bureau of Water, Division of Water Monitoring, Assessment, and Protection, Aquatic Biology Section. SC DHEC, Columbia, SC. 18+ pp.

Table 3. Macroinvertebrates, their North Carolina biotic index tolerance values (TV), functional feeding groups (FG), and abundance collected from Mayo Creek near the effluent discharge of the VC Summer Nuclear Station, Fairfield County, South Carolina, 18 July 2008.

				No. of Individuals			Relative Abundance		
Seq	**Taxon**	**TV**	**FG**	**Sta. 1**	**Sta. 2**	**Sta. 3**	**Sta. 1**	**Sta. 2**	**Sta. 3**
Annelida									
Oligochaeta									
Lumbriculida									
Branchiobdellidae									
1	Branchiobdellidae Genus species		P		7			0.05	
Lumbriculidae									
2	Lumbriculidae Genus species	7.03	SC	1			0.01		
Tubificida									
Tubificidae									
3	Tubifex tubifex	10.00	SC		3			0.02	
Arthropoda									
Arachnoidea									
Acariformes									
Arrenuridae									
4	Arrenurus sp.	5.53	P	1		1	0.01		0.01
Hydrachnidae									
5	Hydrachna sp.	5.53	P	3	4	7	0.02	0.03	0.05
Crustacea									
Cyclopoida									
Cyclopidae									
6	Eucyclops agilis		OM		1			0.01	
Decapoda									
Cambaridae									
7	Cambaridae Genus species		OM	1	2		0.01	0.01	
8	Cambarus sp.	7.62	OM		1			0.01	
Hexapoda									
Coleoptera									
Dytiscidae									
9	Neoporus sp.		P			1			0.01
Elmidae									
10	Dubiraphia quadrinotata	5.93	CG	1	1		0.01	0.01	
11	Dubiraphia sp.	5.93	CG	1			0.01		
12	Oulimnius latiusculus	1.78	CG			2			0.02
13	Stenelmis sp.	5.10	SC		4			0.03	

* CG = collector-gatherer, CF = collector-filterer, OM = omnivore, P = predator, SH = shredder, SC = scraper

Table 3. Continued.

				No. of Individuals			Relative Abundance		
Seq	**Taxon**	**TV**	**FG**	**Sta. 1**	**Sta. 2**	**Sta. 3**	**Sta. 1**	**Sta. 2**	**Sta. 3**
Gyrinidae									
14	Dineutus sp.	5.54	P		1			0.01	
Diptera									
Ceratopogonidae									
15	Bezzia/Palpomyia sp.	6.86	P	4	1		0.03	0.01	
Chironomidae									
16	Ablabesmyia mallochi	7.19	P	7	1	2	0.05	0.01	0.02
17	Cryptotendipes sp.	6.19	CG	3			0.02		
18	Microtendipes pedellus gr.	5.53	CF	12			0.08		
19	Paratendipes sp.	5.11	CG		2			0.01	
20	Phaenopsectra obediens gr.	6.50	SC		1			0.01	
21	Procladius sp.	9.10	P			1			0.01
22	Xestochironomus sp.		P	1			0.01		
23	Zavrelimyia sp.	9.11	P	1	1		0.01	0.01	
Dixidae									
24	Dixella indiana		CG		5	5		0.03	0.04
Simuliidae									
25	Simulium sp.	4.00	CF		6	2		0.04	0.02
Tipulidae									
26	Antocha sp.	4.25	CG		1			0.01	
Ephemeroptera									
Baetidae									
27	Baetis sp.	4.71	CG		3	1		0.02	0.01
28	Centroptilum sp.	6.60	CG			1			0.01
Caenidae									
29	Caenis sp.	7.41	CG	36	33	54	0.24	0.22	0.42
Ephemerellidae									
30	Ephemerella sp.	2.04	CG			1			0.01
Heptageniidae									
31	Maccaffertium modestum	5.50	SC	15	14	11	0.10	0.09	0.09
32	Stenacron interpunctatum	6.87	SC	19	2	8	0.13	0.01	0.06
Isonychiidae									
33	Isonychia sp.	3.45	CF	2	12	4	0.01	0.08	0.03
Megaloptera									
Corydalidae									
34	Corydalus cornutus	5.16	P	1	1	1	0.01	0.01	0.01
35	Nigronia serricornis	4.95	P	1	3	1	0.01	0.02	0.01

* CG = collector-gatherer, CF = collector-filterer, OM = omnivore, P = predator, SH = shredder, SC = scraper

Table 3. Continued.

				No. of Individuals			Relative Abundance		
Seq	**Taxon**	**TV**	**FG**	**Sta. 1**	**Sta. 2**	**Sta. 3**	**Sta. 1**	**Sta. 2**	**Sta. 3**
Sialidae									
36	Sialis aequalis	7.17	P	7	1	1	0.05	0.01	0.01
Odonata									
Aeshnidae									
37	Boyeria vinosa	5.89	P	1	1	3	0.01	0.01	0.02
Calopterygidae									
38	Calopteryx sp.	7.78	P	7		3	0.05		0.02
Coenagrionidae									
39	Argia bipunctulata	8.17	P		1			0.01	
Gomphidae									
40	Gomphus sp.	5.80	P	1	1	4	0.01	0.01	0.03
41	Progomphus sp.		P			1			0.01
Libellulidae									
42	Macromia sp.	6.16	P		2	1		0.01	0.01
Plecoptera									
Perlidae									
43	Perlinella drymo		P	1			0.01		
Trichoptera									
Hydropsychidae									
44	Cheumatopsyche sp.	6.22	CF	15	18	6	0.10	0.12	0.05
45	Hydropsyche betteni	7.78	CF		4			0.03	
Leptoceridae									
46	Triaenodes ignitus	4.58	SH	6	10	6	0.04	0.07	0.05
Philopotamidae									
47	Chimarra sp.	2.76	CF	3	1		0.02	0.01	
Mollusca									
Bivalvia									
Unionoida									
Corbiculidae									
48	Corbicula fluminea	6.12	CF			1			0.01

* CG = collector-gatherer, CF = collector-filterer, OM = omnivore, P = predator, SH = shredder, SC = scraper

Table 4. Rapid bioassessment metrics calculated for the three sampling stations on Mayo Creek near the effluent discharge of the VC Summer Nuclear Station, Fairfield County, South Carolina, 18 July 2008.

	Station		
Metric	**1**	**2**	**3**
Taxa Richness	26	33	26
Number of Specimens	151	149	129
EPT Index	8	9	9
EPT Abundance	97	97	92
Chironomidae Taxa	5	4	2
Chironomidae Abundance	24	5	3
EPT/Chironomidae Abundance	4.04	19.40	30.67
North Carolina Biotic Index	6.17	5.73	5.93
SCDHEC Bioclassification	2.3	2.8	2.3
Percent Collector-Filterers	21.19	27.52	10.08
Percent Collector-Gatherers	27.15	30.20	49.61
Percent Omnivores	0.66	2.68	0.00
Percent Predators	23.84	16.78	20.93
Percent Scrapers	23.18	16.11	14.73
Percent Shredders	3.97	6.71	4.65
Scraper/Collector-Filterers	1.09	0.59	1.46
Shredders/Total	0.04	0.07	0.05
Percent Dominant Taxon	23.84	22.15	41.86
Number Of Dominant Taxa	5	5	4
Dominants In Common		3	3
			2
Community Loss Index		0.24	0.42
			0.62
Jaccard Coefficient of Similarity			0.44
			0.40
Sörensen Coefficient		0.61	0.58
			0.58

Table 5. Dominant taxa (>5% of the collection) for the three sampling stations on Mayo Creek near the effluent discharge of the VC Summer Nuclear Station, Fairfield County, South Carolina, 18 July 2008.

Station 1			**Station 2**			**Station 3**		
Taxon	**No.**	**Rel. Abd.**	**Taxon**	**No.**	**Rel. Abd.**	**Taxon**	**No.**	**Rel. Abd.**
Caenis sp.	36	23.84	Caenis sp.	33	22.15	Caenis sp.	54	41.86
Stenacron interpunctatum	19	12.58	Cheumatopsyche sp.	18	12.08	Maccaffertium modestum	11	8.53
Maccaffertium modestum	15	9.93	Maccaffertium modestum	14	9.40	Stenacron interpunctatum	8	6.20
Cheumatopsyche sp.	15	9.93	Isonychia sp.	12	8.05	Hydrachna sp.	7	5.43
Microtendipes pedellus gr.	12	7.95	Triaenodes ignitus	10	6.71			

Table 6. Habitat assessment scores determined in conjunction with the macroinvertebrate assessment for the three sampling stations on Mayo Creek near the effluent discharge of the VC Summer Nuclear Station, Fairfield County, South Carolina, 18 July 2008.

Habitat Parameter	***Sta. 1***	***Sta. 2***	***Sta. 3***
1. Epifaunal Substrate/Available Cover	12	12	13
2. Pool Substrate Characterization	11	11	11
3. Pool Variability	7	9	7
4. Sediment Deposition	17	11	16
5. Channel Flow Status	12	13	13
6. Channel Alteration	20	20	20
7. Channel Sinuosity	13	16	15
8. Bank Stability (Left Bank (LB*))	8	8	9
Bank Stability (Right Bank (RB*))	8	9	9
9. Vegetative Protection (LB*)	8	8	7
Vegetative Protection (RB*)	8	8	7
10. Riparian Vegetative Zone (LB*)	10	10	10
Riparian Vegetative Zone (RB*)	10	10	10
Total Score	**144**	**145**	**147**

* Left or right bank is determined when facing downstream.