



**Figure 2.4-88—{The Differences Between the Potentiometric Surfaces of the Magothy Aquifer, September 1975 and September 2003, in Southern Maryland}**




FSAR Section 2.4

**Figure 2.4-89—{The Differences Between the Potentiometric Surfaces of the Upper Patapsco Aquifer, September 1990 and September 2003, in Southern Maryland}**




FSAR Section 2.4

**Figure 2.4-90—{The Differences Between the Potentiometric Surfaces of the Lower Patapsco Aquifer, September 1990 and September 2003, in Southern Maryland}**



FSAR Section 2.4

Figure 2.4-91—{Calvert County Ground-Water-Level Monitoring Network, Location of Selected Water Level Monitoring Wells}



FSAR Section 2.4

Figure 2.4-92—{Well Hydrograph for Monitoring Well CA Fd 51 Screened in the Piney Point - Nanjemoy Aquifer at Calvert Cliffs State Park}




Figure 2.4-93—{Well Hydrograph for Monitoring Well CA Ed 42 Screened in the Aquia Aquifer at CCNPP}




Figure 2.4-94—(Well Hydrograph for Monitoring Well CA Dc 35 Screened in the Magothy Aquifer at Scientists Cliffs)




Figure 2.4-95—{Well Hydrograph for Monitoring Well CA Db 96 Screened in the Upper Patapsco Aquifer at Prince Frederick}






Figure 2.4-96—{Well Hydrograph for Monitoring Well CA Fd 85 Screened in the Lower Patapsco Aquifer at Chesapeake Ranch Estates}





Figure 2.4-97—{Modeled Post-Construction Depth to Surficial Aquifer Water Table Around Power Block 3}



**Note: Units are Feet below grade level**