

-----  
Part VI: Uncertainty Analysis  
=====

ORESAD Uncertainty Analysis Results

Probabilistic Input .....	2
Total Dose .....	3
Total Risk .....	4
Dose vs Pathway: Ground External .....	5
Dose vs Pathway: Inhalation (w/o Radon) .....	6
Dose vs Pathway: Radon (Water Ind.) .....	7
Dose vs Pathway: Plant (Water Ind.) .....	8
Dose vs Pathway: Meat (Water Ind.) .....	9
Dose vs Pathway: Milk (Water Ind.) .....	10
Dose vs Pathway: Soil Ingestion .....	11
Dose vs Pathway: Water Ingestion .....	12
Dose vs Pathway: Fish Ingestion .....	13
Dose vs Pathway: Radon (Water Dep.) .....	14
Dose vs Pathway: Plant (Water Dep.) .....	15
Dose vs Pathway: Meat (Water Dep.) .....	16
Dose vs Pathway: Milk (Water Dep.) .....	17
Cumulative Probability Summary.....	18
Summary of dose at graphical times, reptition 1.....	19
Summary of dose at graphical times, reptition 2.....	20
Summary of dose at graphical times, reptition 3.....	21
Peak of the mean dose at graphical times.....	22
Correlation and Regression coefficients (if any).....	23

Probabilistic Input

0Number of Sample Runs: 300

Number	Name	Distribution	Parameters							
1	DCACTC (1)	LOGNORMAL-N	7.78	2.76						
2	DCACTU1 (1)	LOGNORMAL-N	7.78	2.76						
3	DCACTS (1)	LOGNORMAL-N	7.78	2.76						
4	DCACTC (2)	LOGNORMAL-N	8.17	1.7						
5	DCACTU1 (2)	LOGNORMAL-N	8.17	1.7						
6	DCACTS (2)	LOGNORMAL-N	8.17	1.7						
7	BRTF (82,1)	LOGNORMAL-N	-5.52	.916291						
8	BRTF (82,2)	LOGNORMAL-N	-7.13	.693147						
9	BRTF (82,3)	LOGNORMAL-N	-8.11	.91629						
10	BBIO (82,1)	LOGNORMAL-N	5.7	1.1						
11	BRTF (88,1)	LOGNORMAL-N	-3.22	.916291						
12	BRTF (88,2)	LOGNORMAL-N	-6.91	.693147						
13	BRTF (88,3)	LOGNORMAL-N	-6.91	.47						
14	BBIO (88,1)	LOGNORMAL-N	3.9	1.1						
15	UW	UNIFORM	250	2500						
16	MLINH	CONTINUOUS LINEAR	8	0	0	.000008	.0151	.000016	.1365	.00003
.8119	.00004	.9495	.00006	.9937	.000076	.9983	.0001	1		
17	SHF3	UNIFORM	.15	.95						
18	DM	TRIANGULAR	0	.15	.6					
19	DROOT	UNIFORM	.3	4						
20	YV (1)	TRUNCATED LOGNORMAL-N	.56	.48	.001	.999				
21	WLAM	TRIANGULAR	5.1	18	84					
22	RWET (2)	TRIANGULAR	.06	.67	.95					

Nuclide (j)	Peak Time	Peak Dose	DOSE (j, t), mrem/yr								
			t=	0.00E+00	1.00E+00	3.00E+00	1.00E+01	3.00E+01	1.00E+02	3.00E+02	1.00E+03
-----											
Pb-210											
Min	0.00E+00	7.34E-02	7.34E-02	7.08E-02	6.60E-02	2.06E-02	2.41E-04	7.76E-12	0.00E+00	0.00E+00	
Max	0.00E+00	3.45E+00	3.45E+00	3.33E+00	3.11E+00	2.43E+00	1.20E+00	9.23E-02	1.89E-05	6.74E-15	
Avg	0.00E+00	3.89E-01	3.89E-01	3.74E-01	3.46E-01	2.64E-01	1.25E-01	9.14E-03	9.53E-08	7.34E-17	
Std	0.00E+00	4.35E-01	4.35E-01	4.18E-01	3.86E-01	2.95E-01	1.40E-01	1.07E-02	1.22E-06	6.20E-16	
Ra-226											
Min	0.00E+00	2.35E+00	2.35E+00	2.34E+00	1.99E+00	1.10E+00	2.10E-01	2.39E-03	0.00E+00	0.00E+00	
Max	4.16E+01	1.01E+01	9.56E+00	9.62E+00	9.72E+00	9.97E+00	1.01E+01	7.64E+00	1.93E-03	2.19E-04	
Avg	8.34E+00	2.92E+00	2.86E+00	2.87E+00	2.87E+00	2.87E+00	2.81E+00	2.20E+00	6.73E-06	8.27E-07	
Std	1.01E+01	7.76E-01	7.39E-01	7.42E-01	7.46E-01	7.62E-01	7.72E-01	5.86E-01	1.11E-04	1.27E-05	
-ALL											
Min	0.00E+00	2.43E+00	2.43E+00	2.43E+00	2.10E+00	1.19E+00	2.54E-01	5.84E-03	0.00E+00	0.00E+00	
Max	0.00E+00	1.26E+01	1.26E+01	1.25E+01	1.24E+01	1.21E+01	1.11E+01	7.72E+00	1.95E-03	2.19E-04	
Avg	0.00E+00	3.25E+00	3.25E+00	3.24E+00	3.22E+00	3.14E+00	2.93E+00	2.21E+00	6.83E-06	8.27E-07	
Std	0.00E+00	9.90E-01	9.90E-01	9.86E-01	9.77E-01	9.47E-01	8.64E-01	5.93E-01	1.12E-04	1.27E-05	
=====											

-ALL is total dose summed for all nuclides.

0 Probabilistic Risk Summary

0Nuclide (j)	t=	0.00E+00	1.00E+00	3.00E+00	1.00E+01	3.00E+01	1.00E+02	3.00E+02	1.00E+03
-----									
Pb-210									
Min		1.07E-06	1.03E-06	9.63E-07	3.26E-07	3.87E-09	1.25E-16	0.00E+00	0.00E+00
Max		4.99E-05	4.82E-05	4.49E-05	3.51E-05	1.73E-05	1.34E-06	2.29E-10	8.89E-20
Avg		5.64E-06	5.42E-06	5.01E-06	3.83E-06	1.82E-06	1.33E-07	1.15E-12	9.19E-22
Std		6.30E-06	6.06E-06	5.59E-06	4.27E-06	2.03E-06	1.54E-07	1.48E-11	7.84E-21
Ra-226									
Min		5.31E-05	5.22E-05	4.40E-05	2.42E-05	4.52E-06	3.81E-08	0.00E+00	0.00E+00
Max		1.37E-04	1.38E-04	1.40E-04	1.44E-04	1.46E-04	1.14E-04	2.31E-08	2.62E-09
Avg		5.90E-05	5.91E-05	5.91E-05	5.88E-05	5.71E-05	4.52E-05	8.07E-11	9.91E-12
Std		8.59E-06	8.69E-06	8.79E-06	9.23E-06	9.90E-06	8.40E-06	1.33E-09	1.52E-10
-ALL									
Min		5.43E-05	5.40E-05	4.57E-05	2.56E-05	5.17E-06	8.85E-08	0.00E+00	0.00E+00
Max		1.80E-04	1.80E-04	1.79E-04	1.74E-04	1.61E-04	1.15E-04	2.33E-08	2.62E-09
Avg		6.47E-05	6.45E-05	6.41E-05	6.27E-05	5.89E-05	4.53E-05	8.18E-11	9.91E-12
Std		1.24E-05	1.24E-05	1.23E-05	1.20E-05	1.12E-05	8.48E-06	1.35E-09	1.52E-10
=====		=====	=====	=====	=====	=====	=====	=====	=====

-ALL is total risk summed for all nuclides.

0Nuclide (j)	t=	0.00E+00	1.00E+00	3.00E+00	1.00E+01	3.00E+01	1.00E+02	3.00E+02	1.00E+03
-----									
Pb-210									
Min		1.26E-03	9.92E-04	6.12E-04	1.13E-04	9.06E-07	4.08E-14	0.00E+00	0.00E+00
Max		1.40E-03	1.36E-03	1.27E-03	1.02E-03	5.45E-04	5.91E-05	0.00E+00	0.00E+00
Avg		1.40E-03	1.35E-03	1.25E-03	9.82E-04	5.02E-04	5.01E-05	0.00E+00	0.00E+00
Std		1.41E-05	3.85E-05	7.38E-05	1.22E-04	1.02E-04	1.57E-05	0.00E+00	0.00E+00
Ra-226									
Min		2.22E+00	2.03E+00	1.71E+00	9.35E-01	1.66E-01	4.00E-04	0.00E+00	0.00E+00
Max		2.31E+00	2.31E+00	2.30E+00	2.27E+00	2.18E+00	1.80E+00	0.00E+00	0.00E+00
Avg		2.31E+00	2.31E+00	2.29E+00	2.25E+00	2.14E+00	1.70E+00	0.00E+00	0.00E+00
Std		5.83E-03	1.69E-02	3.66E-02	8.59E-02	1.48E-01	2.03E-01	0.00E+00	0.00E+00
-ALL									
Min		2.22E+00	2.04E+00	1.71E+00	9.36E-01	1.66E-01	4.59E-04	0.00E+00	0.00E+00
Max		2.31E+00	2.31E+00	2.30E+00	2.27E+00	2.19E+00	1.80E+00	0.00E+00	0.00E+00
Avg		2.31E+00	2.31E+00	2.30E+00	2.25E+00	2.14E+00	1.70E+00	0.00E+00	0.00E+00
Std		5.83E-03	1.69E-02	3.66E-02	8.59E-02	1.48E-01	2.03E-01	0.00E+00	0.00E+00
=====		=====	=====	=====	=====	=====	=====	=====	=====

-ALL is total pathway dose summed for all nuclides.

0Nuclide (j)	t=	0.00E+00	1.00E+00	3.00E+00	1.00E+01	3.00E+01	1.00E+02	3.00E+02	1.00E+03
-----									
Pb-210									
Min		5.01E-06	4.84E-06	4.51E-06	2.68E-06	2.96E-08	9.49E-16	0.00E+00	0.00E+00
Max		7.48E-04	7.23E-04	6.74E-04	5.26E-04	2.59E-04	2.00E-05	0.00E+00	0.00E+00
Avg		9.78E-05	9.41E-05	8.71E-05	6.70E-05	3.24E-05	2.53E-06	0.00E+00	0.00E+00
Std		7.01E-05	6.75E-05	6.29E-05	4.96E-05	2.54E-05	2.32E-06	0.00E+00	0.00E+00
Ra-226									
Min		1.96E-06	2.11E-06	2.38E-06	3.16E-06	4.35E-06	1.11E-06	0.00E+00	0.00E+00
Max		2.93E-04	3.15E-04	3.55E-04	4.72E-04	6.51E-04	5.82E-04	0.00E+00	0.00E+00
Avg		3.84E-05	4.12E-05	4.65E-05	6.17E-05	8.52E-05	7.95E-05	0.00E+00	0.00E+00
Std		2.75E-05	2.96E-05	3.34E-05	4.47E-05	6.35E-05	6.70E-05	0.00E+00	0.00E+00
-ALL									
Min		6.97E-06	6.94E-06	6.88E-06	6.68E-06	6.08E-06	3.04E-06	0.00E+00	0.00E+00
Max		1.04E-03	1.04E-03	1.03E-03	9.98E-04	9.10E-04	6.02E-04	0.00E+00	0.00E+00
Avg		1.36E-04	1.35E-04	1.34E-04	1.29E-04	1.18E-04	8.21E-05	0.00E+00	0.00E+00
Std		9.76E-05	9.71E-05	9.63E-05	9.40E-05	8.84E-05	6.92E-05	0.00E+00	0.00E+00
=====		=====	=====	=====	=====	=====	=====	=====	=====

-ALL is total pathway dose summed for all nuclides.

0Nuclide (j)	t=	0.00E+00	1.00E+00	3.00E+00	1.00E+01	3.00E+01	1.00E+02	3.00E+02	1.00E+03
-----									
Pb-210									
Min		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Max		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Avg		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Std		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Ra-226									
Min		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Max		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Avg		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Std		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
-ALL									
Min		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Max		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Avg		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Std		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
=====		=====	=====	=====	=====	=====	=====	=====	=====

-ALL is total pathway dose summed for all nuclides.

0Nuclide (j)	t=	0.00E+00	1.00E+00	3.00E+00	1.00E+01	3.00E+01	1.00E+02	3.00E+02	1.00E+03
-----									
Pb-210									
Min		9.40E-03	9.08E-03	8.46E-03	6.61E-03	1.86E-04	5.98E-12	0.00E+00	0.00E+00
Max		3.38E+00	3.26E+00	3.04E+00	2.38E+00	1.17E+00	9.02E-02	0.00E+00	0.00E+00
Avg		2.95E-01	2.83E-01	2.62E-01	2.00E-01	9.38E-02	6.66E-03	0.00E+00	0.00E+00
Std		4.31E-01	4.14E-01	3.82E-01	2.91E-01	1.38E-01	1.03E-02	0.00E+00	0.00E+00
Ra-226									
Min		1.41E-02	1.57E-02	1.87E-02	2.73E-02	2.80E-02	6.42E-04	0.00E+00	0.00E+00
Max		7.21E+00	7.27E+00	7.37E+00	7.63E+00	7.78E+00	5.71E+00	0.00E+00	0.00E+00
Avg		5.27E-01	5.33E-01	5.44E-01	5.74E-01	5.97E-01	4.32E-01	0.00E+00	0.00E+00
Std		7.37E-01	7.39E-01	7.42E-01	7.50E-01	7.45E-01	5.29E-01	0.00E+00	0.00E+00
-ALL									
Min		6.89E-02	6.85E-02	6.77E-02	6.51E-02	4.16E-02	1.69E-03	0.00E+00	0.00E+00
Max		1.01E+01	1.01E+01	9.98E+00	9.67E+00	8.78E+00	5.78E+00	0.00E+00	0.00E+00
Avg		8.21E-01	8.16E-01	8.06E-01	7.74E-01	6.91E-01	4.39E-01	0.00E+00	0.00E+00
Std		9.85E-01	9.80E-01	9.70E-01	9.34E-01	8.39E-01	5.36E-01	0.00E+00	0.00E+00
=====		=====	=====	=====	=====	=====	=====	=====	=====

-ALL is total pathway dose summed for all nuclides.



0Nuclide (j)	t=	0.00E+00	1.00E+00	3.00E+00	1.00E+01	3.00E+01	1.00E+02	3.00E+02	1.00E+03
-----									
Pb-210									
Min		5.97E-04	5.75E-04	5.34E-04	2.87E-04	3.46E-06	2.41E-13	0.00E+00	0.00E+00
Max		6.15E-02	5.96E-02	5.59E-02	4.49E-02	2.40E-02	2.42E-03	0.00E+00	0.00E+00
Avg		9.43E-03	9.07E-03	8.40E-03	6.48E-03	3.16E-03	2.54E-04	0.00E+00	0.00E+00
Std		8.44E-03	8.16E-03	7.63E-03	6.09E-03	3.20E-03	3.14E-04	0.00E+00	0.00E+00
Ra-226									
Min		2.41E-04	3.86E-04	5.24E-04	8.93E-04	9.34E-04	5.57E-05	0.00E+00	0.00E+00
Max		1.65E-02	1.68E-02	1.74E-02	2.06E-02	4.10E-02	4.69E-02	0.00E+00	0.00E+00
Avg		2.76E-03	3.03E-03	3.55E-03	5.03E-03	7.38E-03	7.13E-03	0.00E+00	0.00E+00
Std		2.32E-03	2.38E-03	2.55E-03	3.45E-03	5.67E-03	6.73E-03	0.00E+00	0.00E+00
-ALL									
Min		1.88E-03	1.87E-03	1.84E-03	1.28E-03	9.38E-04	1.51E-04	0.00E+00	0.00E+00
Max		6.58E-02	6.57E-02	6.57E-02	6.55E-02	6.49E-02	4.93E-02	0.00E+00	0.00E+00
Avg		1.22E-02	1.21E-02	1.20E-02	1.15E-02	1.05E-02	7.38E-03	0.00E+00	0.00E+00
Std		9.17E-03	9.15E-03	9.12E-03	9.03E-03	8.73E-03	7.03E-03	0.00E+00	0.00E+00
=====		=====	=====	=====	=====	=====	=====	=====	=====

-ALL is total pathway dose summed for all nuclides.

0Nuclide (j)	t=	0.00E+00	1.00E+00	3.00E+00	1.00E+01	3.00E+01	1.00E+02	3.00E+02	1.00E+03
-----									
Pb-210									
Min		6.03E-04	5.83E-04	5.43E-04	4.24E-04	1.02E-05	6.88E-13	0.00E+00	0.00E+00
Max		1.35E-01	1.31E-01	1.22E-01	9.57E-02	4.71E-02	3.63E-03	0.00E+00	0.00E+00
Avg		1.49E-02	1.44E-02	1.33E-02	1.02E-02	4.94E-03	3.85E-04	0.00E+00	0.00E+00
Std		1.71E-02	1.65E-02	1.53E-02	1.20E-02	5.96E-03	5.03E-04	0.00E+00	0.00E+00
Ra-226									
Min		9.94E-04	1.19E-03	1.55E-03	1.72E-03	1.24E-03	8.12E-05	0.00E+00	0.00E+00
Max		3.46E-02	3.47E-02	3.49E-02	4.44E-02	7.68E-02	7.85E-02	0.00E+00	0.00E+00
Avg		8.15E-03	8.57E-03	9.36E-03	1.16E-02	1.50E-02	1.34E-02	0.00E+00	0.00E+00
Std		5.12E-03	5.21E-03	5.51E-03	7.13E-03	1.11E-02	1.21E-02	0.00E+00	0.00E+00
-ALL									
Min		2.37E-03	2.36E-03	2.34E-03	2.27E-03	2.06E-03	2.22E-04	0.00E+00	0.00E+00
Max		1.41E-01	1.41E-01	1.40E-01	1.36E-01	1.24E-01	8.21E-02	0.00E+00	0.00E+00
Avg		2.31E-02	2.29E-02	2.26E-02	2.18E-02	1.99E-02	1.38E-02	0.00E+00	0.00E+00
Std		1.86E-02	1.85E-02	1.83E-02	1.78E-02	1.66E-02	1.25E-02	0.00E+00	0.00E+00
=====									

-ALL is total pathway dose summed for all nuclides.

0Nuclide (j)	t=	0.00E+00	1.00E+00	3.00E+00	1.00E+01	3.00E+01	1.00E+02	3.00E+02	1.00E+03
-----									
Pb-210									
Min		2.51E-02	2.01E-02	1.29E-02	2.76E-03	2.51E-05	8.05E-13	0.00E+00	0.00E+00
Max		1.02E-01	9.86E-02	9.27E-02	7.45E-02	4.00E-02	4.54E-03	0.00E+00	0.00E+00
Avg		6.83E-02	6.57E-02	6.09E-02	4.70E-02	2.28E-02	1.79E-03	0.00E+00	0.00E+00
Std		2.58E-02	2.50E-02	2.35E-02	1.93E-02	1.08E-02	1.16E-03	0.00E+00	0.00E+00
Ra-226									
Min		5.35E-03	6.14E-03	7.09E-03	8.27E-03	7.86E-03	1.21E-03	0.00E+00	0.00E+00
Max		2.03E-02	2.34E-02	2.94E-02	4.74E-02	8.13E-02	1.14E-01	0.00E+00	0.00E+00
Avg		1.37E-02	1.57E-02	1.95E-02	3.05E-02	4.80E-02	4.76E-02	0.00E+00	0.00E+00
Std		5.16E-03	5.95E-03	7.44E-03	1.20E-02	2.09E-02	2.74E-02	0.00E+00	0.00E+00
-ALL									
Min		3.06E-02	2.63E-02	2.00E-02	1.10E-02	7.90E-03	3.32E-03	0.00E+00	0.00E+00
Max		1.22E-01	1.22E-01	1.22E-01	1.22E-01	1.21E-01	1.18E-01	0.00E+00	0.00E+00
Avg		8.20E-02	8.15E-02	8.04E-02	7.75E-02	7.08E-02	4.94E-02	0.00E+00	0.00E+00
Std		3.09E-02	3.09E-02	3.09E-02	3.12E-02	3.14E-02	2.84E-02	0.00E+00	0.00E+00
=====									

-ALL is total pathway dose summed for all nuclides.

0Nuclide (j)	t=	0.00E+00	1.00E+00	3.00E+00	1.00E+01	3.00E+01	1.00E+02	3.00E+02	1.00E+03
-----									
Pb-210									
Min		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Max		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.30E-05	4.63E-15
Avg		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	6.88E-08	4.24E-17
Std		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	8.66E-07	3.82E-16
Ra-226									
Min		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Max		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.33E-03	1.50E-04
Avg		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	4.66E-06	5.63E-07
Std		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	7.65E-05	8.72E-06
-ALL									
Min		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Max		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.34E-03	1.50E-04
Avg		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	4.73E-06	5.63E-07
Std		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	7.72E-05	8.72E-06
=====									

-ALL is total pathway dose summed for all nuclides.

0Nuclide (j)	t=	0.00E+00	1.00E+00	3.00E+00	1.00E+01	3.00E+01	1.00E+02	3.00E+02	1.00E+03
-----									
DOSE(i,j,t), mrem/yr									
-----									
Pb-210	Min	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
	Max	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	4.77E-06	5.30E-15
	Avg	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	2.17E-08	2.85E-17
	Std	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	2.92E-07	3.25E-16
Ra-226	Min	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
	Max	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	4.87E-04	5.53E-05
	Avg	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.69E-06	2.17E-07
	Std	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	2.81E-05	3.23E-06
-ALL	Min	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
	Max	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	4.92E-04	5.53E-05
	Avg	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.71E-06	2.17E-07
	Std	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	2.84E-05	3.23E-06
=====									

-ALL is total pathway dose summed for all nuclides.

0Nuclide (j)	t=	0.00E+00	1.00E+00	3.00E+00	1.00E+01	3.00E+01	1.00E+02	3.00E+02	1.00E+03
-----									
Pb-210									
Min		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Max		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Avg		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Std		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Ra-226									
Min		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Max		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Avg		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Std		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
-ALL									
Min		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Max		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Avg		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Std		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
=====		=====	=====	=====	=====	=====	=====	=====	=====

-ALL is total pathway dose summed for all nuclides.

0Nuclide (j)	t=	0.00E+00	1.00E+00	3.00E+00	1.00E+01	3.00E+01	1.00E+02	3.00E+02	1.00E+03
-----									
Pb-210									
Min		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Max		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.08E-06	3.87E-16
Avg		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	4.62E-09	2.44E-18
Std		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	6.47E-08	2.48E-17
Ra-226									
Min		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Max		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.11E-04	1.26E-05
Avg		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	3.78E-07	4.46E-08
Std		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	6.38E-06	7.25E-07
-ALL									
Min		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Max		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.12E-04	1.26E-05
Avg		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	3.83E-07	4.46E-08
Std		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	6.44E-06	7.25E-07
=====		=====	=====	=====	=====	=====	=====	=====	=====

-ALL is total pathway dose summed for all nuclides.

0Nuclide (j)	t=	0.00E+00	1.00E+00	3.00E+00	1.00E+01	3.00E+01	1.00E+02	3.00E+02	1.00E+03
DOSE(i,j,t), mrem/yr									
-----									
Pb-210	Min	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
	Max	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.66E-08	5.94E-18
	Avg	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	8.04E-11	3.19E-20
	Std	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.05E-09	3.77E-19
Ra-226	Min	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
	Max	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.70E-06	1.92E-07
	Avg	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	5.87E-09	6.61E-10
	Std	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	9.78E-08	1.11E-08
-ALL	Min	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
	Max	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.71E-06	1.92E-07
	Avg	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	5.95E-09	6.61E-10
	Std	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	9.88E-08	1.11E-08
=====	=====	=====	=====	=====	=====	=====	=====	=====	=====

-ALL is total pathway dose summed for all nuclides.



0Nuclide (j)	t=	0.00E+00	1.00E+00	3.00E+00	1.00E+01	3.00E+01	1.00E+02	3.00E+02	1.00E+03
DOSE(i,j,t), mrem/yr									
-----									
Pb-210									
Min		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Max		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	3.55E-08	1.27E-17
Avg		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.63E-10	6.82E-20
Std		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	2.18E-09	7.90E-19
Ra-226									
Min		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Max		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.37E-06	1.55E-07
Avg		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	5.59E-09	5.60E-10
Std		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	8.08E-08	8.96E-09
-ALL									
Min		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00
Max		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	1.38E-06	1.55E-07
Avg		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	5.75E-09	5.60E-10
Std		0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	0.00E+00	8.20E-08	8.96E-09
=====		=====	=====	=====	=====	=====	=====	=====	=====

-ALL is total pathway dose summed for all nuclides.

Cumulative Probability	Dose (t), mrem/yr							
	t= 0.00E+00	1.00E+00	3.00E+00	1.00E+01	3.00E+01	1.00E+02	3.00E+02	1.00E+03
0.025	2.51E+00	2.50E+00	2.48E+00	2.43E+00	2.23E+00	1.33E+00	0.00E+00	0.00E+00
0.050	2.56E+00	2.55E+00	2.53E+00	2.45E+00	2.30E+00	1.65E+00	0.00E+00	0.00E+00
0.075	2.57E+00	2.56E+00	2.55E+00	2.50E+00	2.33E+00	1.79E+00	0.00E+00	0.00E+00
0.100	2.58E+00	2.58E+00	2.56E+00	2.50E+00	2.36E+00	1.86E+00	0.00E+00	0.00E+00
0.125	2.60E+00	2.60E+00	2.58E+00	2.53E+00	2.39E+00	1.87E+00	0.00E+00	0.00E+00
0.150	2.62E+00	2.61E+00	2.59E+00	2.54E+00	2.42E+00	1.89E+00	0.00E+00	0.00E+00
0.175	2.64E+00	2.63E+00	2.62E+00	2.56E+00	2.43E+00	1.91E+00	0.00E+00	0.00E+00
0.200	2.67E+00	2.66E+00	2.64E+00	2.58E+00	2.44E+00	1.92E+00	0.00E+00	0.00E+00
0.225	2.68E+00	2.67E+00	2.66E+00	2.61E+00	2.46E+00	1.93E+00	0.00E+00	0.00E+00
0.250	2.70E+00	2.69E+00	2.67E+00	2.62E+00	2.48E+00	1.95E+00	0.00E+00	0.00E+00
0.275	2.73E+00	2.72E+00	2.69E+00	2.64E+00	2.50E+00	1.96E+00	0.00E+00	0.00E+00
0.300	2.75E+00	2.74E+00	2.72E+00	2.66E+00	2.52E+00	1.97E+00	0.00E+00	0.00E+00
0.325	2.76E+00	2.75E+00	2.73E+00	2.68E+00	2.54E+00	1.99E+00	0.00E+00	0.00E+00
0.350	2.79E+00	2.77E+00	2.75E+00	2.70E+00	2.56E+00	2.01E+00	0.00E+00	0.00E+00
0.375	2.81E+00	2.80E+00	2.78E+00	2.74E+00	2.58E+00	2.03E+00	0.00E+00	0.00E+00
0.400	2.83E+00	2.82E+00	2.80E+00	2.76E+00	2.60E+00	2.04E+00	0.00E+00	0.00E+00
0.425	2.86E+00	2.85E+00	2.84E+00	2.78E+00	2.62E+00	2.06E+00	0.00E+00	0.00E+00
0.450	2.89E+00	2.88E+00	2.87E+00	2.81E+00	2.64E+00	2.08E+00	0.00E+00	0.00E+00
0.475	2.92E+00	2.91E+00	2.90E+00	2.83E+00	2.67E+00	2.09E+00	0.00E+00	0.00E+00
0.500	2.96E+00	2.95E+00	2.92E+00	2.85E+00	2.69E+00	2.10E+00	0.00E+00	0.00E+00
0.525	2.99E+00	2.98E+00	2.96E+00	2.90E+00	2.72E+00	2.11E+00	0.00E+00	0.00E+00
0.550	3.03E+00	3.02E+00	3.01E+00	2.96E+00	2.76E+00	2.14E+00	0.00E+00	0.00E+00
0.575	3.08E+00	3.07E+00	3.05E+00	2.98E+00	2.80E+00	2.16E+00	0.00E+00	0.00E+00
0.600	3.14E+00	3.13E+00	3.10E+00	3.02E+00	2.83E+00	2.17E+00	0.00E+00	0.00E+00
0.625	3.17E+00	3.17E+00	3.15E+00	3.09E+00	2.87E+00	2.20E+00	0.00E+00	0.00E+00
0.650	3.21E+00	3.20E+00	3.18E+00	3.11E+00	2.92E+00	2.25E+00	0.00E+00	0.00E+00
0.675	3.25E+00	3.24E+00	3.22E+00	3.14E+00	2.94E+00	2.27E+00	0.00E+00	0.00E+00
0.700	3.32E+00	3.31E+00	3.27E+00	3.19E+00	3.02E+00	2.30E+00	0.00E+00	0.00E+00
0.725	3.35E+00	3.33E+00	3.32E+00	3.24E+00	3.04E+00	2.32E+00	0.00E+00	0.00E+00
0.750	3.38E+00	3.37E+00	3.35E+00	3.29E+00	3.09E+00	2.36E+00	0.00E+00	0.00E+00
0.775	3.45E+00	3.44E+00	3.42E+00	3.34E+00	3.12E+00	2.38E+00	0.00E+00	0.00E+00
0.800	3.52E+00	3.51E+00	3.49E+00	3.41E+00	3.21E+00	2.42E+00	0.00E+00	0.00E+00
0.825	3.64E+00	3.62E+00	3.59E+00	3.52E+00	3.30E+00	2.48E+00	0.00E+00	0.00E+00
0.850	3.85E+00	3.84E+00	3.81E+00	3.70E+00	3.44E+00	2.52E+00	0.00E+00	0.00E+00
0.875	4.03E+00	4.01E+00	3.98E+00	3.87E+00	3.60E+00	2.68E+00	0.00E+00	0.00E+00
0.900	4.26E+00	4.25E+00	4.22E+00	4.13E+00	3.79E+00	2.78E+00	0.00E+00	0.00E+00
0.925	4.53E+00	4.51E+00	4.46E+00	4.31E+00	3.99E+00	2.91E+00	0.00E+00	0.00E+00
0.950	4.79E+00	4.77E+00	4.73E+00	4.58E+00	4.26E+00	3.10E+00	0.00E+00	0.00E+00
0.975	6.04E+00	5.94E+00	5.74E+00	5.57E+00	5.02E+00	3.52E+00	0.00E+00	6.79E-13
1.000	1.26E+01	1.25E+01	1.24E+01	1.21E+01	1.11E+01	7.72E+00	1.95E-03	2.19E-04







Repetition	Time of peak mean dose Years	Peak mean dose mrem/yr
1	0.000E+00	3.230E+00
2	0.000E+00	3.219E+00
3	0.000E+00	3.307E+00

Coefficients for peak of mean dose time Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC		SRC		PRCC		SRRC	
	Sig	Coef	Sig	Coef	Sig	Coef	Sig	Coef
Kd of Pb-210 in Contaminated Zone	10	-0.11	9	-0.05	22	0.00	22	0.00
Kd of Pb-210 in Unsaturated Zone 1	13	-0.06	13	-0.03	11	-0.09	11	-0.03
Kd of Pb-210 in Saturated Zone	18	0.02	18	0.01	5	-0.19	5	-0.06
Kd of Ra-226 in Contaminated Zone	8	-0.12	8	-0.07	19	-0.03	19	-0.01
Kd of Ra-226 in Unsaturated Zone 1	17	-0.03	17	-0.02	10	0.12	10	0.04
Kd of Ra-226 in Saturated Zone	6	-0.16	6	-0.08	21	0.01	21	0.00
Plant transfer factor for Pb	3	0.60	3	0.40	3	0.72	3	0.32
Meat transfer factor for Pb	11	0.09	11	0.05	9	0.12	9	0.04
Milk transfer factor for Pb	14	-0.04	14	-0.02	8	-0.14	8	-0.04
Fish transfer factor for Pb	4	-0.29	4	-0.16	16	-0.06	16	-0.02
Plant transfer factor for Ra	2	0.73	2	0.52	2	0.88	2	0.57
Meat transfer factor for Ra	19	-0.01	19	-0.01	20	-0.02	20	-0.01
Milk transfer factor for Ra	9	0.11	10	0.05	14	0.07	14	0.02
Fish transfer factor for Ra	12	-0.09	12	-0.04	6	-0.18	6	-0.06
Well pumping rate	21	0.00	21	0.00	18	0.04	18	0.01
Mass loading for inhalation	7	-0.15	7	-0.07	15	-0.06	15	-0.02
Indoor dust filtration factor	20	-0.01	20	0.00	7	-0.17	7	-0.05
Depth of soil mixing layer	5	-0.16	5	-0.08	4	-0.34	4	-0.12
Depth of roots	1	-0.76	1	-0.57	1	-0.90	1	-0.65
Wet weight crop yield of fruit, grain and non-leafy vegetables	16	-0.04	16	-0.02	17	0.06	17	0.02
Weathering removal constant of all vegetation	15	-0.04	15	-0.02	13	-0.08	13	-0.02
Wet foliar interception fraction of leafy vegetables	22	0.00	22	0.00	12	-0.08	12	-0.03
R-SQUARE		0.78		0.78		0.90		0.90

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak of mean dose time Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC		SRC		PRCC		SRRC	
	Sig	Coeff	Sig	Coeff	Sig	Coeff	Sig	Coeff
Kd of Pb-210 in Contaminated Zone	17	0.02	17	0.02	18	-0.02	18	-0.01
Kd of Pb-210 in Unsaturated Zone 1	15	0.05	15	0.03	12	-0.12	12	-0.04
Kd of Pb-210 in Saturated Zone	21	-0.01	21	0.00	14	-0.04	14	-0.01
Kd of Ra-226 in Contaminated Zone	20	0.01	20	0.00	9	-0.17	9	-0.06
Kd of Ra-226 in Unsaturated Zone 1	22	0.00	22	0.00	7	0.19	7	0.06
Kd of Ra-226 in Saturated Zone	9	-0.17	7	-0.12	8	0.18	8	0.06
Plant transfer factor for Pb	3	0.48	3	0.31	3	0.72	3	0.35
Meat transfer factor for Pb	4	0.28	5	0.17	11	0.13	11	0.04
Milk transfer factor for Pb	11	0.13	11	0.07	16	0.03	16	0.01
Fish transfer factor for Pb	18	0.02	18	0.01	6	-0.20	6	-0.07
Plant transfer factor for Ra	2	0.61	2	0.44	2	0.87	2	0.60
Meat transfer factor for Ra	10	-0.14	10	-0.08	19	-0.02	19	-0.01
Milk transfer factor for Ra	5	0.26	6	0.16	22	0.01	22	0.00
Fish transfer factor for Ra	6	0.24	4	0.17	20	-0.02	20	-0.01
Well pumping rate	16	-0.05	16	-0.03	21	0.02	21	0.01
Mass loading for inhalation	19	-0.02	19	-0.01	5	-0.21	5	-0.07
Indoor dust filtration factor	13	0.07	13	0.04	17	0.03	17	0.01
Depth of soil mixing layer	8	-0.17	9	-0.10	4	-0.37	4	-0.13
Depth of roots	1	-0.71	1	-0.57	1	-0.88	1	-0.62
Wet weight crop yield of fruit, grain and non-leafy vegetables	12	0.10	12	0.06	15	0.04	15	0.01
Weathering removal constant of all vegetation	7	-0.19	8	-0.11	10	-0.14	10	-0.05
Wet foliar interception fraction of leafy vegetables	14	0.06	14	0.03	13	0.05	13	0.02
R-SQUARE		0.72		0.72		0.89		0.89

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.



Coefficients for peak of mean dose time Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC 3		SRC 3		PRCC 3		SRRC 3	
	Sig	Coeff	Sig	Coeff	Sig	Coeff	Sig	Coeff
Kd of Pb-210 in Contaminated Zone	19	-0.02	19	-0.02	6	0.14	6	0.05
Kd of Pb-210 in Unsaturated Zone 1	15	-0.05	15	-0.04	10	-0.08	10	-0.03
Kd of Pb-210 in Saturated Zone	13	-0.08	13	-0.05	15	-0.04	16	-0.01
Kd of Ra-226 in Contaminated Zone	10	0.11	9	0.08	17	-0.04	17	-0.01
Kd of Ra-226 in Unsaturated Zone 1	9	-0.11	10	-0.07	5	0.16	5	0.06
Kd of Ra-226 in Saturated Zone	18	0.03	18	0.02	18	0.03	18	0.01
Plant transfer factor for Pb	3	0.46	3	0.33	3	0.72	3	0.38
Meat transfer factor for Pb	21	-0.01	22	-0.01	16	-0.04	15	-0.01
Milk transfer factor for Pb	22	0.01	21	0.01	19	-0.02	19	-0.01
Fish transfer factor for Pb	17	-0.04	17	-0.03	8	0.12	8	0.04
Plant transfer factor for Ra	2	0.60	2	0.48	2	0.82	2	0.53
Meat transfer factor for Ra	7	-0.14	7	-0.09	12	0.07	12	0.03
Milk transfer factor for Ra	5	-0.15	5	-0.10	11	0.07	11	0.03
Fish transfer factor for Ra	14	-0.07	14	-0.05	7	-0.12	7	-0.05
Well pumping rate	12	-0.08	12	-0.05	13	-0.06	13	-0.02
Mass loading for inhalation	20	-0.02	20	-0.01	14	0.05	14	0.02
Indoor dust filtration factor	8	-0.13	8	-0.09	22	0.00	22	0.00
Depth of soil mixing layer	11	-0.11	11	-0.07	4	-0.28	4	-0.11
Depth of roots	1	-0.62	1	-0.51	1	-0.86	1	-0.64
Wet weight crop yield of fruit, grain and non-leafy vegetables	16	0.05	16	0.03	20	0.02	20	0.01
Weathering removal constant of all vegetation	6	-0.15	6	-0.10	21	-0.01	21	0.00
Wet foliar interception fraction of leafy vegetables	4	0.18	4	0.12	9	0.08	9	0.03
R-SQUARE		0.61		0.61		0.86		0.86

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak All Pathways Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC		SRC		PRCC		SRRC	
	Sig	Coeff	Sig	Coeff	Sig	Coeff	Sig	Coeff
Kd of Pb-210 in Contaminated Zone	10	-0.11	9	-0.05	22	0.00	22	0.00
Kd of Pb-210 in Unsaturated Zone 1	13	-0.06	13	-0.03	11	-0.09	11	-0.03
Kd of Pb-210 in Saturated Zone	18	0.02	18	0.01	5	-0.19	5	-0.06
Kd of Ra-226 in Contaminated Zone	8	-0.12	8	-0.07	19	-0.03	19	-0.01
Kd of Ra-226 in Unsaturated Zone 1	17	-0.03	17	-0.02	10	0.12	10	0.04
Kd of Ra-226 in Saturated Zone	6	-0.16	6	-0.08	21	0.01	21	0.00
Plant transfer factor for Pb	3	0.60	3	0.40	3	0.72	3	0.32
Meat transfer factor for Pb	11	0.09	11	0.05	9	0.12	9	0.04
Milk transfer factor for Pb	14	-0.04	14	-0.02	8	-0.14	8	-0.04
Fish transfer factor for Pb	4	-0.29	4	-0.16	17	-0.06	17	-0.02
Plant transfer factor for Ra	2	0.73	2	0.52	2	0.88	2	0.57
Meat transfer factor for Ra	19	-0.01	19	-0.01	20	-0.02	20	-0.01
Milk transfer factor for Ra	9	0.11	10	0.05	14	0.07	14	0.02
Fish transfer factor for Ra	12	-0.09	12	-0.04	6	-0.18	6	-0.06
Well pumping rate	21	0.00	21	0.00	18	0.04	18	0.01
Mass loading for inhalation	7	-0.15	7	-0.07	15	-0.06	15	-0.02
Indoor dust filtration factor	20	-0.01	20	0.00	7	-0.17	7	-0.05
Depth of soil mixing layer	5	-0.16	5	-0.08	4	-0.34	4	-0.12
Depth of roots	1	-0.76	1	-0.57	1	-0.90	1	-0.65
Wet weight crop yield of fruit, grain and non-leafy vegetables	16	-0.04	16	-0.02	16	0.06	16	0.02
Weathering removal constant of all vegetation	15	-0.04	15	-0.02	13	-0.08	13	-0.02
Wet foliar interception fraction of leafy vegetables	22	0.00	22	0.00	12	-0.08	12	-0.03
R-SQUARE		0.78		0.78		0.90		0.90

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak All Pathways Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC		SRC		PRCC		SRRC	
	Sig	Coef	Sig	Coef	Sig	Coef	Sig	Coef
Kd of Pb-210 in Contaminated Zone	17	0.02	17	0.02	19	-0.02	19	-0.01
Kd of Pb-210 in Unsaturated Zone 1	15	0.05	15	0.03	12	-0.12	12	-0.04
Kd of Pb-210 in Saturated Zone	21	-0.01	21	0.00	15	-0.04	15	-0.01
Kd of Ra-226 in Contaminated Zone	20	0.01	20	0.00	9	-0.17	9	-0.06
Kd of Ra-226 in Unsaturated Zone 1	22	0.00	22	0.00	7	0.19	7	0.06
Kd of Ra-226 in Saturated Zone	9	-0.17	7	-0.12	8	0.18	8	0.06
Plant transfer factor for Pb	3	0.48	3	0.31	3	0.72	3	0.35
Meat transfer factor for Pb	4	0.28	5	0.17	11	0.13	11	0.05
Milk transfer factor for Pb	11	0.13	11	0.07	16	0.03	16	0.01
Fish transfer factor for Pb	18	0.02	18	0.01	6	-0.20	6	-0.07
Plant transfer factor for Ra	2	0.61	2	0.44	2	0.87	2	0.60
Meat transfer factor for Ra	10	-0.14	10	-0.08	18	-0.02	18	-0.01
Milk transfer factor for Ra	5	0.26	6	0.16	22	0.01	22	0.00
Fish transfer factor for Ra	6	0.24	4	0.17	20	-0.02	20	-0.01
Well pumping rate	16	-0.05	16	-0.03	21	0.02	21	0.01
Mass loading for inhalation	19	-0.02	19	-0.01	5	-0.21	5	-0.07
Indoor dust filtration factor	13	0.07	13	0.04	17	0.03	17	0.01
Depth of soil mixing layer	8	-0.17	9	-0.10	4	-0.37	4	-0.13
Depth of roots	1	-0.71	1	-0.57	1	-0.88	1	-0.62
Wet weight crop yield of fruit, grain and non-leafy vegetables	12	0.10	12	0.06	14	0.04	14	0.01
Weathering removal constant of all vegetation	7	-0.19	8	-0.11	10	-0.14	10	-0.05
Wet foliar interception fraction of leafy vegetables	14	0.06	14	0.03	13	0.05	13	0.02
R-SQUARE		0.72		0.72		0.89		0.89

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak All Pathways Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC 3		SRC 3		PRCC 3		SRRC 3	
	Sig	Coeff	Sig	Coeff	Sig	Coeff	Sig	Coeff
Kd of Pb-210 in Contaminated Zone	19	-0.02	19	-0.02	6	0.14	6	0.05
Kd of Pb-210 in Unsaturated Zone 1	15	-0.05	15	-0.04	9	-0.08	9	-0.03
Kd of Pb-210 in Saturated Zone	13	-0.08	13	-0.05	15	-0.04	15	-0.01
Kd of Ra-226 in Contaminated Zone	10	0.11	9	0.08	16	-0.03	16	-0.01
Kd of Ra-226 in Unsaturated Zone 1	9	-0.11	10	-0.07	5	0.16	5	0.06
Kd of Ra-226 in Saturated Zone	18	0.03	18	0.02	18	0.03	18	0.01
Plant transfer factor for Pb	3	0.46	3	0.33	3	0.72	3	0.38
Meat transfer factor for Pb	21	-0.01	22	-0.01	17	-0.03	17	-0.01
Milk transfer factor for Pb	22	0.01	21	0.01	19	-0.02	19	-0.01
Fish transfer factor for Pb	17	-0.04	17	-0.03	8	0.12	8	0.04
Plant transfer factor for Ra	2	0.60	2	0.48	2	0.82	2	0.53
Meat transfer factor for Ra	7	-0.14	7	-0.09	11	0.07	11	0.03
Milk transfer factor for Ra	5	-0.15	5	-0.10	12	0.07	12	0.03
Fish transfer factor for Ra	14	-0.07	14	-0.05	7	-0.12	7	-0.05
Well pumping rate	12	-0.08	12	-0.05	13	-0.06	13	-0.02
Mass loading for inhalation	20	-0.02	20	-0.01	14	0.05	14	0.02
Indoor dust filtration factor	8	-0.13	8	-0.09	22	0.00	22	0.00
Depth of soil mixing layer	11	-0.11	11	-0.07	4	-0.28	4	-0.11
Depth of roots	1	-0.62	1	-0.51	1	-0.86	1	-0.64
Wet weight crop yield of fruit, grain and non-leafy vegetables	16	0.05	16	0.03	20	0.02	20	0.01
Weathering removal constant of all vegetation	6	-0.15	6	-0.10	21	0.00	21	0.00
Wet foliar interception fraction of leafy vegetables	4	0.18	4	0.12	10	0.08	10	0.03
R-SQUARE		0.61		0.61		0.86		0.86

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak External Ground Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC		SRC		PRCC		SRRC	
	Sig	Coeff	Sig	Coeff	Sig	Coeff	Sig	Coeff
Kd of Pb-210 in Contaminated Zone	21	-0.02	21	-0.02	13	-0.08	13	-0.05
Kd of Pb-210 in Unsaturated Zone 1	7	0.08	7	0.08	8	0.10	8	0.07
Kd of Pb-210 in Saturated Zone	19	0.02	19	0.02	16	-0.07	16	-0.04
Kd of Ra-226 in Contaminated Zone	1	0.13	1	0.14	1	0.75	1	0.72
Kd of Ra-226 in Unsaturated Zone 1	11	0.05	10	0.06	20	0.04	20	0.03
Kd of Ra-226 in Saturated Zone	8	0.08	8	0.08	2	0.19	2	0.13
Plant transfer factor for Pb	4	0.09	4	0.09	10	0.08	10	0.05
Meat transfer factor for Pb	14	-0.04	14	-0.04	15	-0.07	15	-0.04
Milk transfer factor for Pb	17	-0.04	17	-0.04	18	0.05	18	0.03
Fish transfer factor for Pb	13	-0.05	13	-0.05	4	-0.15	4	-0.10
Plant transfer factor for Ra	9	0.07	9	0.07	12	-0.08	12	-0.05
Meat transfer factor for Ra	18	0.03	18	0.04	21	-0.04	21	-0.03
Milk transfer factor for Ra	22	0.02	22	0.02	9	-0.09	9	-0.06
Fish transfer factor for Ra	5	0.08	5	0.08	11	0.08	11	0.05
Well pumping rate	6	-0.08	6	-0.08	5	0.12	5	0.08
Mass loading for inhalation	10	-0.06	11	-0.06	6	-0.11	6	-0.07
Indoor dust filtration factor	20	0.02	20	0.02	7	-0.11	7	-0.07
Depth of soil mixing layer	3	0.11	3	0.11	22	-0.03	22	-0.02
Depth of roots	15	-0.04	15	-0.04	17	0.07	17	0.04
Wet weight crop yield of fruit, grain and non-leafy vegetables	16	-0.04	16	-0.04	19	0.04	19	0.03
Weathering removal constant of all vegetation	2	0.13	2	0.13	3	0.19	3	0.12
Wet foliar interception fraction of leafy vegetables	12	-0.05	12	-0.05	14	-0.07	14	-0.05
R-SQUARE		0.09		0.09		0.59		0.59

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak External Ground Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC		SRC		PRCC		SRRC	
	Sig	Coeff	Sig	Coeff	Sig	Coeff	Sig	Coeff
Kd of Pb-210 in Contaminated Zone	15	-0.03	14	-0.04	7	0.10	7	0.07
Kd of Pb-210 in Unsaturated Zone 1	19	0.02	19	0.02	8	-0.10	8	-0.06
Kd of Pb-210 in Saturated Zone	13	0.04	13	0.04	19	0.02	19	0.01
Kd of Ra-226 in Contaminated Zone	5	0.11	5	0.11	1	0.74	1	0.72
Kd of Ra-226 in Unsaturated Zone 1	10	0.08	11	0.07	3	0.17	3	0.11
Kd of Ra-226 in Saturated Zone	11	0.07	9	0.08	13	0.07	13	0.04
Plant transfer factor for Pb	2	0.16	2	0.16	12	0.08	12	0.05
Meat transfer factor for Pb	6	0.10	6	0.10	17	-0.04	17	-0.03
Milk transfer factor for Pb	9	0.08	10	0.08	5	0.12	5	0.08
Fish transfer factor for Pb	12	0.07	12	0.06	22	0.00	22	0.00
Plant transfer factor for Ra	21	-0.01	21	-0.01	10	-0.08	10	-0.05
Meat transfer factor for Ra	1	-0.32	1	-0.33	9	-0.09	9	-0.06
Milk transfer factor for Ra	8	0.09	8	0.09	15	-0.05	15	-0.03
Fish transfer factor for Ra	22	0.00	22	0.00	11	0.08	11	0.05
Well pumping rate	14	0.04	15	0.03	2	-0.18	2	-0.12
Mass loading for inhalation	4	0.12	4	0.12	6	0.10	6	0.07
Indoor dust filtration factor	20	0.02	20	0.02	18	-0.03	18	-0.02
Depth of soil mixing layer	18	0.02	18	0.02	14	-0.06	14	-0.04
Depth of roots	3	-0.13	3	-0.12	21	-0.01	21	-0.01
Wet weight crop yield of fruit, grain and non-leafy vegetables	7	0.10	7	0.10	4	-0.17	4	-0.11
Weathering removal constant of all vegetation	16	-0.03	16	-0.03	16	-0.05	16	-0.03
Wet foliar interception fraction of leafy vegetables	17	0.03	17	0.03	20	-0.02	20	-0.01
R-SQUARE		0.17		0.17		0.58		0.58

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak External Ground Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC 3		SRC 3		PRCC 3		SRRC 3	
	Sig	Coeff	Sig	Coeff	Sig	Coeff	Sig	Coeff
Kd of Pb-210 in Contaminated Zone	6	0.14	4	0.16	11	0.07	11	0.05
Kd of Pb-210 in Unsaturated Zone 1	16	0.05	16	0.04	17	0.03	17	0.02
Kd of Pb-210 in Saturated Zone	17	0.04	17	0.04	21	0.01	21	0.01
Kd of Ra-226 in Contaminated Zone	1	0.23	1	0.25	1	0.75	1	0.71
Kd of Ra-226 in Unsaturated Zone 1	20	-0.01	20	-0.01	6	-0.16	6	-0.10
Kd of Ra-226 in Saturated Zone	13	0.09	12	0.08	8	0.11	8	0.07
Plant transfer factor for Pb	15	-0.05	15	-0.05	20	0.02	20	0.02
Meat transfer factor for Pb	18	0.03	18	0.03	3	-0.24	3	-0.15
Milk transfer factor for Pb	4	-0.16	5	-0.16	10	-0.08	10	-0.05
Fish transfer factor for Pb	14	0.08	14	0.08	13	0.07	13	0.04
Plant transfer factor for Ra	12	0.09	13	0.08	12	0.07	12	0.05
Meat transfer factor for Ra	2	0.20	2	0.19	19	0.03	19	0.02
Milk transfer factor for Ra	21	0.01	21	0.01	16	-0.04	16	-0.02
Fish transfer factor for Ra	7	-0.13	7	-0.14	2	-0.24	2	-0.15
Well pumping rate	8	-0.12	9	-0.11	7	-0.15	7	-0.09
Mass loading for inhalation	9	-0.11	8	-0.13	15	-0.04	15	-0.03
Indoor dust filtration factor	22	0.01	22	0.00	18	-0.03	18	-0.02
Depth of soil mixing layer	3	-0.18	3	-0.17	4	-0.18	4	-0.12
Depth of roots	5	0.15	6	0.14	5	0.16	5	0.10
Wet weight crop yield of fruit, grain and non-leafy vegetables	10	0.11	10	0.10	22	0.00	22	0.00
Weathering removal constant of all vegetation	11	-0.09	11	-0.09	14	-0.05	14	-0.03
Wet foliar interception fraction of leafy vegetables	19	0.03	19	0.03	9	0.08	9	0.05
R-SQUARE		0.20		0.20		0.62		0.62

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak Inhalation Particles Dose		PCC		SRC		PRCC		SRRC	
Coefficient =		1		1		1		1	
Repetition =									
Description of Probabilistic Variable	Sig	Coeff	Sig	Coeff	Sig	Coeff	Sig	Coeff	Sig
Kd of Pb-210 in Contaminated Zone	20	0.03	20	0.01	18	0.04	18	0.01	
Kd of Pb-210 in Unsaturated Zone 1	18	-0.05	19	-0.02	9	-0.16	9	-0.04	
Kd of Pb-210 in Saturated Zone	17	-0.05	18	-0.02	11	-0.15	11	-0.04	
Kd of Ra-226 in Contaminated Zone	7	0.15	7	0.05	15	-0.10	15	-0.03	
Kd of Ra-226 in Unsaturated Zone 1	16	-0.06	15	-0.02	4	-0.31	4	-0.08	
Kd of Ra-226 in Saturated Zone	12	-0.09	12	-0.03	17	-0.06	17	-0.02	
Plant transfer factor for Pb	5	0.19	5	0.07	21	-0.03	21	-0.01	
Meat transfer factor for Pb	19	0.05	17	0.02	10	-0.16	10	-0.04	
Milk transfer factor for Pb	6	-0.18	6	-0.06	22	0.00	22	0.00	
Fish transfer factor for Pb	11	-0.10	11	-0.03	19	0.04	19	0.01	
Plant transfer factor for Ra	9	-0.13	9	-0.04	7	-0.17	7	-0.04	
Meat transfer factor for Ra	21	-0.02	21	-0.01	14	-0.11	14	-0.03	
Milk transfer factor for Ra	15	0.07	16	0.02	8	0.16	8	0.04	
Fish transfer factor for Ra	4	-0.23	4	-0.08	5	-0.29	5	-0.08	
Well pumping rate	14	0.07	14	0.02	12	-0.14	12	-0.04	
Mass loading for inhalation	1	0.88	1	0.60	2	0.92	2	0.60	
Indoor dust filtration factor	3	0.82	3	0.47	3	0.89	3	0.49	
Depth of soil mixing layer	2	-0.83	2	-0.50	1	-0.92	1	-0.60	
Depth of roots	10	-0.12	10	-0.04	13	0.14	13	0.03	
Wet weight crop yield of fruit, grain and non-leafy vegetables	8	-0.14	8	-0.05	16	-0.10	16	-0.02	
Weathering removal constant of all vegetation	22	-0.01	22	0.00	20	-0.04	20	-0.01	
Wet foliar interception fraction of leafy vegetables	13	0.08	13	0.03	6	0.24	6	0.06	
R-SQUARE		0.90		0.90		0.94		0.94	

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.



Coefficients for peak Inhalation Particles Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC		SRC		PRCC		SRRC	
	Sig	Coef	Sig	Coef	Sig	Coef	Sig	Coef
Kd of Pb-210 in Contaminated Zone	18	0.02	18	0.01	17	0.05	17	0.01
Kd of Pb-210 in Unsaturated Zone 1	8	-0.08	8	-0.03	16	-0.05	16	-0.01
Kd of Pb-210 in Saturated Zone	10	-0.06	10	-0.02	13	0.06	13	0.01
Kd of Ra-226 in Contaminated Zone	11	-0.04	11	-0.01	10	0.09	10	0.02
Kd of Ra-226 in Unsaturated Zone 1	17	0.02	17	0.01	20	0.03	20	0.01
Kd of Ra-226 in Saturated Zone	19	-0.01	19	0.00	8	-0.10	8	-0.02
Plant transfer factor for Pb	12	-0.02	12	-0.01	4	-0.17	4	-0.04
Meat transfer factor for Pb	20	0.01	20	0.00	21	-0.01	21	0.00
Milk transfer factor for Pb	6	0.12	6	0.05	7	-0.10	7	-0.02
Fish transfer factor for Pb	22	0.00	22	0.00	19	-0.03	19	-0.01
Plant transfer factor for Ra	21	0.01	21	0.00	11	0.08	11	0.02
Meat transfer factor for Ra	4	-0.13	4	-0.05	14	-0.06	14	-0.01
Milk transfer factor for Ra	13	-0.02	13	-0.01	18	0.04	18	0.01
Fish transfer factor for Ra	7	-0.08	7	-0.04	12	-0.08	12	-0.02
Well pumping rate	14	-0.02	14	-0.01	6	-0.11	6	-0.03
Mass loading for inhalation	3	0.79	3	0.50	2	0.92	2	0.54
Indoor dust filtration factor	2	0.80	2	0.52	3	0.91	3	0.51
Depth of soil mixing layer	1	-0.84	1	-0.60	1	-0.94	1	-0.60
Depth of roots	9	-0.07	9	-0.03	5	0.16	5	0.04
Wet weight crop yield of fruit, grain and non-leafy vegetables	16	0.02	16	0.01	22	0.00	22	0.00
Weathering removal constant of all vegetation	5	0.13	5	0.05	9	0.09	9	0.02
Wet foliar interception fraction of leafy vegetables	15	0.02	15	0.01	15	-0.05	15	-0.01
R-SQUARE		0.87		0.87		0.95		0.95

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak Inhalation Particles Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC 3		SRC 3		PRCC 3		SRRC 3	
	Sig	Coeff	Sig	Coeff	Sig	Coeff	Sig	Coeff
Kd of Pb-210 in Contaminated Zone	4	0.54	4	0.22	6	-0.17	6	-0.04
Kd of Pb-210 in Unsaturated Zone 1	11	0.11	11	0.03	22	0.00	22	0.00
Kd of Pb-210 in Saturated Zone	15	-0.06	15	-0.02	19	-0.03	19	-0.01
Kd of Ra-226 in Contaminated Zone	21	0.02	20	0.01	11	0.09	11	0.02
Kd of Ra-226 in Unsaturated Zone 1	14	-0.07	14	-0.02	18	-0.04	18	-0.01
Kd of Ra-226 in Saturated Zone	19	-0.03	19	-0.01	9	0.10	9	0.02
Plant transfer factor for Pb	13	0.08	13	0.02	12	-0.08	12	-0.02
Meat transfer factor for Pb	8	-0.13	8	-0.04	5	-0.22	5	-0.05
Milk transfer factor for Pb	5	-0.23	5	-0.07	16	-0.06	16	-0.01
Fish transfer factor for Pb	16	-0.05	16	-0.01	17	0.05	17	0.01
Plant transfer factor for Ra	20	-0.02	21	-0.01	21	0.01	21	0.00
Meat transfer factor for Ra	10	0.12	10	0.03	13	0.08	13	0.02
Milk transfer factor for Ra	22	0.01	22	0.00	20	0.01	20	0.00
Fish transfer factor for Ra	12	0.09	12	0.03	4	0.23	4	0.05
Well pumping rate	7	-0.15	7	-0.04	8	0.14	8	0.03
Mass loading for inhalation	1	0.88	1	0.60	2	0.93	2	0.56
Indoor dust filtration factor	3	0.77	3	0.34	3	0.93	3	0.55
Depth of soil mixing layer	2	-0.83	2	-0.40	1	-0.94	1	-0.63
Depth of roots	6	-0.23	6	-0.06	7	0.15	7	0.03
Wet weight crop yield of fruit, grain and non-leafy vegetables	9	0.12	9	0.03	10	-0.10	10	-0.02
Weathering removal constant of all vegetation	17	0.04	17	0.01	15	0.06	15	0.01
Wet foliar interception fraction of leafy vegetables	18	-0.04	18	-0.01	14	0.07	14	0.01
R-SQUARE		0.93		0.93		0.95		0.95

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak Radon (WaterInd.) Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC		SRC		PRCC		SRRC	
	Sig	Coeff	Sig	Coeff	Sig	Coeff	Sig	Coeff
Kd of Pb-210 in Contaminated Zone	0	0.00	0	0.00	0	0.00	0	0.00
Kd of Pb-210 in Unsaturated Zone 1	0	0.00	0	0.00	0	0.00	0	0.00
Kd of Pb-210 in Saturated Zone	0	0.00	0	0.00	0	0.00	0	0.00
Kd of Ra-226 in Contaminated Zone	0	0.00	0	0.00	0	0.00	0	0.00
Kd of Ra-226 in Unsaturated Zone 1	0	0.00	0	0.00	0	0.00	0	0.00
Kd of Ra-226 in Saturated Zone	0	0.00	0	0.00	0	0.00	0	0.00
Plant transfer factor for Pb	0	0.00	0	0.00	0	0.00	0	0.00
Meat transfer factor for Pb	0	0.00	0	0.00	0	0.00	0	0.00
Milk transfer factor for Pb	0	0.00	0	0.00	0	0.00	0	0.00
Fish transfer factor for Pb	0	0.00	0	0.00	0	0.00	0	0.00
Plant transfer factor for Ra	0	0.00	0	0.00	0	0.00	0	0.00
Meat transfer factor for Ra	0	0.00	0	0.00	0	0.00	0	0.00
Milk transfer factor for Ra	0	0.00	0	0.00	0	0.00	0	0.00
Fish transfer factor for Ra	0	0.00	0	0.00	0	0.00	0	0.00
Well pumping rate	0	0.00	0	0.00	0	0.00	0	0.00
Mass loading for inhalation	0	0.00	0	0.00	0	0.00	0	0.00
Indoor dust filtration factor	0	0.00	0	0.00	0	0.00	0	0.00
Depth of soil mixing layer	0	0.00	0	0.00	0	0.00	0	0.00
Depth of roots	0	0.00	0	0.00	0	0.00	0	0.00
Wet weight crop yield of fruit, grain and non-leafy vegetables	0	0.00	0	0.00	0	0.00	0	0.00
Weathering removal constant of all vegetation	0	0.00	0	0.00	0	0.00	0	0.00
Wet foliar interception fraction of leafy vegetables	0	0.00	0	0.00	0	0.00	0	0.00
R-SQUARE		0.00		0.00		0.00		0.00

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak Radon (WaterInd.) Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC		SRC		PRCC		SRRC	
	Sig	Coeff	Sig	Coeff	Sig	Coeff	Sig	Coeff
Kd of Pb-210 in Contaminated Zone	0	0.00	0	0.00	0	0.00	0	0.00
Kd of Pb-210 in Unsaturated Zone 1	0	0.00	0	0.00	0	0.00	0	0.00
Kd of Pb-210 in Saturated Zone	0	0.00	0	0.00	0	0.00	0	0.00
Kd of Ra-226 in Contaminated Zone	0	0.00	0	0.00	0	0.00	0	0.00
Kd of Ra-226 in Unsaturated Zone 1	0	0.00	0	0.00	0	0.00	0	0.00
Kd of Ra-226 in Saturated Zone	0	0.00	0	0.00	0	0.00	0	0.00
Plant transfer factor for Pb	0	0.00	0	0.00	0	0.00	0	0.00
Meat transfer factor for Pb	0	0.00	0	0.00	0	0.00	0	0.00
Milk transfer factor for Pb	0	0.00	0	0.00	0	0.00	0	0.00
Fish transfer factor for Pb	0	0.00	0	0.00	0	0.00	0	0.00
Plant transfer factor for Ra	0	0.00	0	0.00	0	0.00	0	0.00
Meat transfer factor for Ra	0	0.00	0	0.00	0	0.00	0	0.00
Milk transfer factor for Ra	0	0.00	0	0.00	0	0.00	0	0.00
Fish transfer factor for Ra	0	0.00	0	0.00	0	0.00	0	0.00
Well pumping rate	0	0.00	0	0.00	0	0.00	0	0.00
Mass loading for inhalation	0	0.00	0	0.00	0	0.00	0	0.00
Indoor dust filtration factor	0	0.00	0	0.00	0	0.00	0	0.00
Depth of soil mixing layer	0	0.00	0	0.00	0	0.00	0	0.00
Depth of roots	0	0.00	0	0.00	0	0.00	0	0.00
Wet weight crop yield of fruit, grain and non-leafy vegetables	0	0.00	0	0.00	0	0.00	0	0.00
Weathering removal constant of all vegetation	0	0.00	0	0.00	0	0.00	0	0.00
Wet foliar interception fraction of leafy vegetables	0	0.00	0	0.00	0	0.00	0	0.00
R-SQUARE		0.00		0.00		0.00		0.00

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak Radon (WaterInd.) Dose		PCC		SRC		PRCC		SRRC	
Coefficient =		3		3		3		3	
Repetition =									
Description of Probabilistic Variable		Sig	Coeff	Sig	Coeff	Sig	Coeff	Sig	Coeff
Kd of Pb-210 in Contaminated Zone		0	0.00	0	0.00	0	0.00	0	0.00
Kd of Pb-210 in Unsaturated Zone 1		0	0.00	0	0.00	0	0.00	0	0.00
Kd of Pb-210 in Saturated Zone		0	0.00	0	0.00	0	0.00	0	0.00
Kd of Ra-226 in Contaminated Zone		0	0.00	0	0.00	0	0.00	0	0.00
Kd of Ra-226 in Unsaturated Zone 1		0	0.00	0	0.00	0	0.00	0	0.00
Kd of Ra-226 in Saturated Zone		0	0.00	0	0.00	0	0.00	0	0.00
Plant transfer factor for Pb		0	0.00	0	0.00	0	0.00	0	0.00
Meat transfer factor for Pb		0	0.00	0	0.00	0	0.00	0	0.00
Milk transfer factor for Pb		0	0.00	0	0.00	0	0.00	0	0.00
Fish transfer factor for Pb		0	0.00	0	0.00	0	0.00	0	0.00
Plant transfer factor for Ra		0	0.00	0	0.00	0	0.00	0	0.00
Meat transfer factor for Ra		0	0.00	0	0.00	0	0.00	0	0.00
Milk transfer factor for Ra		0	0.00	0	0.00	0	0.00	0	0.00
Fish transfer factor for Ra		0	0.00	0	0.00	0	0.00	0	0.00
Well pumping rate		0	0.00	0	0.00	0	0.00	0	0.00
Mass loading for inhalation		0	0.00	0	0.00	0	0.00	0	0.00
Indoor dust filtration factor		0	0.00	0	0.00	0	0.00	0	0.00
Depth of soil mixing layer		0	0.00	0	0.00	0	0.00	0	0.00
Depth of roots		0	0.00	0	0.00	0	0.00	0	0.00
Wet weight crop yield of fruit, grain and non-leafy vegetables		0	0.00	0	0.00	0	0.00	0	0.00
Weathering removal constant of all vegetation		0	0.00	0	0.00	0	0.00	0	0.00
Wet foliar interception fraction of leafy vegetables		0	0.00	0	0.00	0	0.00	0	0.00
R-SQUARE		0.00		0.00		0.00		0.00	

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak Plant (WaterInd.) Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC		SRC		PRCC		SRRC	
	Sig	Coef	Sig	Coef	Sig	Coef	Sig	Coef
Kd of Pb-210 in Contaminated Zone	8	-0.10	8	-0.05	21	0.00	21	0.00
Kd of Pb-210 in Unsaturated Zone 1	13	-0.07	13	-0.03	13	-0.05	13	-0.01
Kd of Pb-210 in Saturated Zone	19	0.02	19	0.01	5	-0.17	5	-0.05
Kd of Ra-226 in Contaminated Zone	7	-0.13	6	-0.07	19	0.01	19	0.00
Kd of Ra-226 in Unsaturated Zone 1	17	-0.03	17	-0.01	10	0.09	10	0.03
Kd of Ra-226 in Saturated Zone	5	-0.16	5	-0.08	20	0.01	20	0.00
Plant transfer factor for Pb	3	0.60	3	0.39	3	0.72	3	0.33
Meat transfer factor for Pb	11	0.08	11	0.04	12	0.06	12	0.02
Milk transfer factor for Pb	12	-0.07	12	-0.04	4	-0.22	4	-0.07
Fish transfer factor for Pb	4	-0.28	4	-0.16	9	-0.09	9	-0.03
Plant transfer factor for Ra	2	0.73	2	0.52	2	0.87	2	0.57
Meat transfer factor for Ra	18	-0.02	18	-0.01	16	-0.03	16	-0.01
Milk transfer factor for Ra	9	0.10	9	0.05	22	0.00	22	0.00
Fish transfer factor for Ra	10	-0.09	10	-0.04	6	-0.14	6	-0.05
Well pumping rate	22	0.00	22	0.00	18	0.01	18	0.00
Mass loading for inhalation	6	-0.14	7	-0.07	15	-0.03	15	-0.01
Indoor dust filtration factor	20	0.00	20	0.00	7	-0.12	7	-0.04
Depth of soil mixing layer	14	-0.06	14	-0.03	17	-0.02	17	-0.01
Depth of roots	1	-0.76	1	-0.58	1	-0.90	1	-0.66
Wet weight crop yield of fruit, grain and non-leafy vegetables	15	-0.04	16	-0.02	14	0.03	14	0.01
Weathering removal constant of all vegetation	16	-0.04	15	-0.02	8	-0.10	8	-0.03
Wet foliar interception fraction of leafy vegetables	21	0.00	21	0.00	11	-0.09	11	-0.03
R-SQUARE		0.77		0.77		0.90		0.90

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak Plant (WaterInd.) Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC		SRC		PRCC		SRRC	
	Sig	Coef	Sig	Coef	Sig	Coef	Sig	Coef
Kd of Pb-210 in Contaminated Zone	17	0.03	17	0.02	15	-0.04	15	-0.01
Kd of Pb-210 in Unsaturated Zone 1	15	0.06	15	0.04	14	-0.04	14	-0.01
Kd of Pb-210 in Saturated Zone	20	-0.01	20	-0.01	22	0.00	22	0.00
Kd of Ra-226 in Contaminated Zone	21	0.00	21	0.00	5	-0.20	5	-0.07
Kd of Ra-226 in Unsaturated Zone 1	22	0.00	22	0.00	7	0.16	7	0.06
Kd of Ra-226 in Saturated Zone	8	-0.17	7	-0.13	8	0.16	8	0.05
Plant transfer factor for Pb	3	0.47	3	0.31	3	0.73	3	0.36
Meat transfer factor for Pb	4	0.27	5	0.16	9	0.12	9	0.04
Milk transfer factor for Pb	11	0.08	11	0.05	13	-0.05	13	-0.02
Fish transfer factor for Pb	18	0.02	18	0.01	6	-0.20	6	-0.07
Plant transfer factor for Ra	2	0.61	2	0.45	1	0.88	1	0.62
Meat transfer factor for Ra	9	-0.13	9	-0.08	20	0.01	20	0.00
Milk transfer factor for Ra	5	0.26	6	0.16	21	0.01	21	0.00
Fish transfer factor for Ra	6	0.25	4	0.18	19	-0.02	19	-0.01
Well pumping rate	16	-0.04	16	-0.02	17	-0.02	17	-0.01
Mass loading for inhalation	19	-0.01	19	-0.01	4	-0.30	4	-0.11
Indoor dust filtration factor	13	0.07	13	0.04	18	-0.02	18	-0.01
Depth of soil mixing layer	12	-0.08	12	-0.05	11	-0.07	11	-0.02
Depth of roots	1	-0.71	1	-0.57	2	-0.88	2	-0.62
Wet weight crop yield of fruit, grain and non-leafy vegetables	10	0.11	10	0.06	16	0.03	16	0.01
Weathering removal constant of all vegetation	7	-0.20	8	-0.12	10	-0.10	10	-0.03
Wet foliar interception fraction of leafy vegetables	14	0.06	14	0.04	12	0.06	12	0.02
R-SQUARE		0.71		0.71		0.89		0.89

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak Plant (WaterInd.) Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC		SRC		PRCC		SRRC	
	Sig	Coeff	Sig	Coeff	Sig	Coeff	Sig	Coeff
Kd of Pb-210 in Contaminated Zone	19	-0.02	19	-0.02	5	0.13	5	0.05
Kd of Pb-210 in Unsaturated Zone 1	15	-0.05	15	-0.04	10	-0.08	10	-0.03
Kd of Pb-210 in Saturated Zone	11	-0.08	11	-0.05	19	-0.02	19	-0.01
Kd of Ra-226 in Contaminated Zone	10	0.11	9	0.08	17	-0.05	17	-0.02
Kd of Ra-226 in Unsaturated Zone 1	9	-0.11	10	-0.07	7	0.12	7	0.04
Kd of Ra-226 in Saturated Zone	18	0.03	18	0.02	18	0.03	18	0.01
Plant transfer factor for Pb	3	0.46	3	0.33	3	0.73	3	0.39
Meat transfer factor for Pb	21	-0.02	21	-0.01	9	-0.09	9	-0.03
Milk transfer factor for Pb	22	0.00	22	0.00	8	-0.10	8	-0.04
Fish transfer factor for Pb	17	-0.04	17	-0.03	4	0.14	4	0.05
Plant transfer factor for Ra	2	0.60	2	0.48	2	0.83	2	0.54
Meat transfer factor for Ra	7	-0.14	7	-0.09	16	0.05	16	0.02
Milk transfer factor for Ra	5	-0.15	5	-0.10	13	0.06	13	0.02
Fish transfer factor for Ra	13	-0.07	13	-0.05	6	-0.13	6	-0.05
Well pumping rate	12	-0.08	12	-0.05	14	-0.06	14	-0.02
Mass loading for inhalation	20	-0.02	20	-0.01	12	0.06	12	0.02
Indoor dust filtration factor	8	-0.13	8	-0.09	21	0.01	21	0.00
Depth of soil mixing layer	14	-0.06	14	-0.04	15	-0.05	15	-0.02
Depth of roots	1	-0.62	1	-0.51	1	-0.87	1	-0.65
Wet weight crop yield of fruit, grain and non-leafy vegetables	16	0.05	16	0.03	20	-0.01	20	0.00
Weathering removal constant of all vegetation	6	-0.15	6	-0.10	22	0.00	22	0.00
Wet foliar interception fraction of leafy vegetables	4	0.18	4	0.12	11	0.07	11	0.03
R-SQUARE		0.61		0.61		0.87		0.87

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.



Coefficients for peak Meat (WaterInd.) Dose		PCC		SRC		PRCC		SRRC	
Coefficient =		1		1		1		1	
Repetition =									
Description of Probabilistic Variable	Sig	Coef	Sig	Coef	Sig	Coef	Sig	Coef	
Kd of Pb-210 in Contaminated Zone	15	-0.08	15	-0.02	22	0.00	22	0.00	
Kd of Pb-210 in Unsaturated Zone 1	20	0.03	20	0.01	7	0.18	7	0.04	
Kd of Pb-210 in Saturated Zone	22	-0.02	22	-0.01	20	0.03	20	0.01	
Kd of Ra-226 in Contaminated Zone	5	0.31	5	0.10	6	-0.20	6	-0.05	
Kd of Ra-226 in Unsaturated Zone 1	13	-0.10	11	-0.03	16	0.05	16	0.01	
Kd of Ra-226 in Saturated Zone	21	0.03	21	0.01	15	-0.07	15	-0.02	
Plant transfer factor for Pb	4	0.37	4	0.12	12	0.11	12	0.03	
Meat transfer factor for Pb	1	0.94	1	0.78	1	0.95	1	0.73	
Milk transfer factor for Pb	10	-0.14	10	-0.04	19	0.04	19	0.01	
Fish transfer factor for Pb	9	-0.16	8	-0.05	11	-0.13	11	-0.03	
Plant transfer factor for Ra	7	0.20	7	0.06	4	0.47	4	0.13	
Meat transfer factor for Ra	3	0.53	3	0.19	3	0.68	3	0.22	
Milk transfer factor for Ra	19	-0.04	19	-0.01	5	0.31	5	0.08	
Fish transfer factor for Ra	18	-0.05	18	-0.01	13	-0.10	13	-0.02	
Well pumping rate	17	-0.06	17	-0.02	10	0.14	10	0.03	
Mass loading for inhalation	11	-0.11	12	-0.03	21	0.00	21	0.00	
Indoor dust filtration factor	6	-0.27	6	-0.08	9	-0.14	9	-0.03	
Depth of soil mixing layer	2	-0.83	2	-0.42	2	-0.93	2	-0.58	
Depth of roots	16	-0.07	16	-0.02	8	-0.16	8	-0.04	
Wet weight crop yield of fruit, grain and non-leafy vegetables	14	0.09	14	0.02	18	0.04	18	0.01	
Weathering removal constant of all vegetation	12	0.10	13	0.03	14	-0.10	14	-0.02	
Wet foliar interception fraction of leafy vegetables	8	0.16	9	0.05	17	0.05	17	0.01	
R-SQUARE		0.93		0.93		0.94		0.94	

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak Meat (WaterInd.) Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC		SRC		PRCC		SRRC	
	Sig	Coeff	Sig	Coeff	Sig	Coeff	Sig	Coeff
Kd of Pb-210 in Contaminated Zone	21	0.01	20	0.00	9	0.14	9	0.03
Kd of Pb-210 in Unsaturated Zone 1	18	-0.02	18	-0.01	16	-0.06	16	-0.01
Kd of Pb-210 in Saturated Zone	9	-0.12	10	-0.04	10	-0.13	10	-0.03
Kd of Ra-226 in Contaminated Zone	14	-0.10	14	-0.03	11	0.11	11	0.03
Kd of Ra-226 in Unsaturated Zone 1	15	0.10	15	0.03	21	0.01	21	0.00
Kd of Ra-226 in Saturated Zone	8	0.14	5	0.05	19	0.02	19	0.01
Plant transfer factor for Pb	19	0.02	19	0.01	13	0.10	13	0.02
Meat transfer factor for Pb	1	0.94	1	0.79	1	0.95	1	0.72
Milk transfer factor for Pb	12	-0.11	13	-0.03	15	0.07	15	0.02
Fish transfer factor for Pb	17	-0.03	17	-0.01	7	0.18	7	0.04
Plant transfer factor for Ra	4	0.19	4	0.06	5	0.26	5	0.07
Meat transfer factor for Ra	3	0.54	3	0.19	3	0.74	3	0.26
Milk transfer factor for Ra	7	-0.14	8	-0.04	22	0.00	22	0.00
Fish transfer factor for Ra	11	-0.11	9	-0.04	20	-0.02	20	-0.01
Well pumping rate	20	-0.01	21	0.00	17	0.05	17	0.01
Mass loading for inhalation	6	-0.15	7	-0.04	6	-0.25	6	-0.06
Indoor dust filtration factor	16	-0.03	16	-0.01	18	0.03	18	0.01
Depth of soil mixing layer	2	-0.84	2	-0.44	2	-0.92	2	-0.57
Depth of roots	5	-0.16	6	-0.05	8	-0.18	8	-0.04
Wet weight crop yield of fruit, grain and non-leafy vegetables	22	0.01	22	0.00	12	0.11	12	0.03
Weathering removal constant of all vegetation	10	0.11	11	0.03	4	-0.31	4	-0.08
Wet foliar interception fraction of leafy vegetables	13	-0.10	12	-0.03	14	0.09	14	0.02
R-SQUARE		0.92		0.92		0.94		0.94

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak Meat (WaterInd.) Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC 3		SRC 3		PRCC 3		SRRC 3	
	Sig	Coeff	Sig	Coeff	Sig	Coeff	Sig	Coeff
Kd of Pb-210 in Contaminated Zone	21	0.02	21	0.01	21	0.02	21	0.00
Kd of Pb-210 in Unsaturated Zone 1	19	-0.03	19	-0.01	14	-0.07	14	-0.02
Kd of Pb-210 in Saturated Zone	18	-0.03	18	-0.01	6	-0.17	6	-0.04
Kd of Ra-226 in Contaminated Zone	8	-0.13	8	-0.04	13	0.08	13	0.02
Kd of Ra-226 in Unsaturated Zone 1	12	-0.09	12	-0.03	19	0.03	19	0.01
Kd of Ra-226 in Saturated Zone	16	-0.05	15	-0.02	16	-0.06	16	-0.01
Plant transfer factor for Pb	22	0.01	22	0.00	20	0.03	20	0.01
Meat transfer factor for Pb	1	0.93	1	0.79	1	0.95	1	0.76
Milk transfer factor for Pb	4	-0.26	4	-0.09	17	-0.04	17	-0.01
Fish transfer factor for Pb	20	-0.03	20	-0.01	11	0.10	11	0.02
Plant transfer factor for Ra	17	-0.04	17	-0.01	4	0.23	4	0.06
Meat transfer factor for Ra	3	0.61	3	0.24	3	0.72	3	0.25
Milk transfer factor for Ra	15	0.05	16	0.02	5	-0.18	5	-0.04
Fish transfer factor for Ra	7	0.15	7	0.05	18	0.04	18	0.01
Well pumping rate	5	-0.20	5	-0.06	15	0.06	15	0.02
Mass loading for inhalation	11	-0.12	9	-0.04	12	0.08	12	0.02
Indoor dust filtration factor	10	0.12	11	0.04	10	-0.10	10	-0.03
Depth of soil mixing layer	2	-0.87	2	-0.52	2	-0.92	2	-0.55
Depth of roots	6	-0.17	6	-0.05	7	-0.17	7	-0.04
Wet weight crop yield of fruit, grain and non-leafy vegetables	9	-0.12	10	-0.04	22	0.00	22	0.00
Weathering removal constant of all vegetation	14	0.06	14	0.02	8	0.17	8	0.04
Wet foliar interception fraction of leafy vegetables	13	-0.09	13	-0.03	9	0.16	9	0.04
R-SQUARE		0.91		0.91		0.94		0.94

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak Milk (WaterInd.) Dose		PCC		SRC		PRCC		SRRC	
Coefficient =		1		1		1		1	
Repetition =									
Description of Probabilistic Variable	Sig	Coef	Sig	Coef	Sig	Coef	Sig	Coef	
Kd of Pb-210 in Contaminated Zone	10	-0.11	10	-0.03	20	0.02	20	0.01	
Kd of Pb-210 in Unsaturated Zone 1	6	0.16	6	0.05	21	-0.01	21	0.00	
Kd of Pb-210 in Saturated Zone	22	0.00	22	0.00	19	0.03	19	0.01	
Kd of Ra-226 in Contaminated Zone	7	0.15	7	0.05	6	-0.20	6	-0.06	
Kd of Ra-226 in Unsaturated Zone 1	11	-0.09	11	-0.03	9	-0.14	9	-0.04	
Kd of Ra-226 in Saturated Zone	21	0.00	21	0.00	8	-0.17	8	-0.05	
Plant transfer factor for Pb	9	0.12	9	0.04	7	0.20	7	0.05	
Meat transfer factor for Pb	8	-0.14	8	-0.04	15	0.05	15	0.01	
Milk transfer factor for Pb	1	0.94	1	0.80	1	0.93	1	0.70	
Fish transfer factor for Pb	14	-0.09	14	-0.03	14	0.07	14	0.02	
Plant transfer factor for Ra	5	0.34	5	0.10	4	0.35	4	0.10	
Meat transfer factor for Ra	13	0.09	12	0.03	10	0.13	10	0.04	
Milk transfer factor for Ra	3	0.61	3	0.22	3	0.76	3	0.31	
Fish transfer factor for Ra	18	0.07	18	0.02	18	0.05	17	0.01	
Well pumping rate	16	0.07	16	0.02	17	0.05	18	0.01	
Mass loading for inhalation	15	0.08	15	0.02	12	-0.11	12	-0.03	
Indoor dust filtration factor	4	-0.39	4	-0.12	13	-0.09	13	-0.02	
Depth of soil mixing layer	2	-0.83	2	-0.43	2	-0.88	2	-0.51	
Depth of roots	12	-0.09	13	-0.03	5	-0.29	5	-0.08	
Wet weight crop yield of fruit, grain and non-leafy vegetables	20	-0.01	20	0.00	16	0.05	16	0.01	
Weathering removal constant of all vegetation	19	-0.06	19	-0.02	11	-0.12	11	-0.03	
Wet foliar interception fraction of leafy vegetables	17	0.07	17	0.02	22	-0.01	22	0.00	
R-SQUARE		0.92		0.92		0.93		0.93	

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak Milk (WaterInd.) Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC		SRC		PRCC		SRRC	
	Sig	Coeff	Sig	Coeff	Sig	Coeff	Sig	Coeff
Kd of Pb-210 in Contaminated Zone	13	-0.05	12	-0.02	7	0.19	7	0.05
Kd of Pb-210 in Unsaturated Zone 1	5	-0.19	5	-0.06	10	-0.14	10	-0.04
Kd of Pb-210 in Saturated Zone	9	0.08	8	0.03	22	0.00	22	0.00
Kd of Ra-226 in Contaminated Zone	22	0.00	22	0.00	9	-0.17	9	-0.04
Kd of Ra-226 in Unsaturated Zone 1	16	-0.04	16	-0.01	20	-0.02	20	0.00
Kd of Ra-226 in Saturated Zone	20	-0.02	20	-0.01	18	-0.04	18	-0.01
Plant transfer factor for Pb	14	0.05	14	0.01	17	-0.05	17	-0.01
Meat transfer factor for Pb	10	-0.06	10	-0.02	11	-0.11	11	-0.03
Milk transfer factor for Pb	1	0.95	1	0.88	1	0.95	1	0.72
Fish transfer factor for Pb	19	-0.02	19	-0.01	15	0.05	15	0.01
Plant transfer factor for Ra	17	-0.03	17	-0.01	5	0.31	5	0.08
Meat transfer factor for Ra	6	-0.18	6	-0.05	14	0.08	14	0.02
Milk transfer factor for Ra	3	0.54	3	0.19	3	0.76	3	0.29
Fish transfer factor for Ra	18	-0.02	18	-0.01	19	-0.04	19	-0.01
Well pumping rate	7	-0.10	7	-0.03	6	0.22	6	0.06
Mass loading for inhalation	21	-0.01	21	0.00	13	-0.08	13	-0.02
Indoor dust filtration factor	12	0.05	13	0.02	21	0.01	21	0.00
Depth of soil mixing layer	2	-0.76	2	-0.33	2	-0.91	2	-0.55
Depth of roots	4	-0.20	4	-0.06	4	-0.33	4	-0.09
Wet weight crop yield of fruit, grain and non-leafy vegetables	8	-0.08	9	-0.02	8	0.18	8	0.05
Weathering removal constant of all vegetation	11	0.06	11	0.02	12	-0.11	12	-0.03
Wet foliar interception fraction of leafy vegetables	15	-0.04	15	-0.01	16	0.05	16	0.01
R-SQUARE		0.93		0.93		0.94		0.94

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak Milk (WaterInd.) Dose		PCC		SRC		PRCC		SRRC	
Coefficient =		3		3		3		3	
Repetition =									
Description of Probabilistic Variable	Sig	Coeff	Sig	Coeff	Sig	Coeff	Sig	Coeff	Sig
Kd of Pb-210 in Contaminated Zone	19	-0.01	19	0.00	15	-0.09	15	-0.03	
Kd of Pb-210 in Unsaturated Zone 1	7	-0.17	7	-0.05	11	-0.12	11	-0.04	
Kd of Pb-210 in Saturated Zone	5	0.22	5	0.07	20	0.05	20	0.02	
Kd of Ra-226 in Contaminated Zone	20	0.01	20	0.00	8	0.16	8	0.05	
Kd of Ra-226 in Unsaturated Zone 1	21	0.01	21	0.00	21	0.02	21	0.01	
Kd of Ra-226 in Saturated Zone	14	-0.08	14	-0.02	18	-0.05	18	-0.02	
Plant transfer factor for Pb	16	0.07	16	0.02	19	-0.05	19	-0.02	
Meat transfer factor for Pb	10	-0.16	10	-0.05	6	0.18	6	0.06	
Milk transfer factor for Pb	1	0.94	1	0.82	1	0.92	1	0.73	
Fish transfer factor for Pb	12	0.08	12	0.02	22	0.00	22	0.00	
Plant transfer factor for Ra	4	0.29	4	0.09	5	0.24	5	0.07	
Meat transfer factor for Ra	8	-0.17	8	-0.05	10	-0.13	10	-0.04	
Milk transfer factor for Ra	3	0.56	3	0.20	3	0.71	3	0.31	
Fish transfer factor for Ra	18	0.04	18	0.01	13	0.10	13	0.03	
Well pumping rate	6	0.17	6	0.05	12	-0.11	12	-0.03	
Mass loading for inhalation	11	0.13	11	0.04	16	-0.07	16	-0.02	
Indoor dust filtration factor	9	-0.16	9	-0.05	7	-0.17	7	-0.05	
Depth of soil mixing layer	2	-0.84	2	-0.45	2	-0.87	2	-0.54	
Depth of roots	15	-0.07	15	-0.02	4	-0.35	4	-0.11	
Wet weight crop yield of fruit, grain and non-leafy vegetables	22	0.00	22	0.00	17	-0.05	17	-0.02	
Weathering removal constant of all vegetation	13	0.08	13	0.02	14	-0.09	14	-0.03	
Wet foliar interception fraction of leafy vegetables	17	0.05	17	0.01	9	0.15	9	0.05	
R-SQUARE		0.92		0.92		0.91		0.91	

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak Soil Ingestion Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC		SRC		PRCC		SRRC	
	Sig	Coeff	Sig	Coeff	Sig	Coeff	Sig	Coeff
Kd of Pb-210 in Contaminated Zone	6	-0.18	6	-0.04	4	0.23	4	0.03
Kd of Pb-210 in Unsaturated Zone 1	19	-0.03	19	-0.01	13	-0.07	13	-0.01
Kd of Pb-210 in Saturated Zone	13	-0.10	13	-0.02	6	0.11	6	0.01
Kd of Ra-226 in Contaminated Zone	16	0.06	16	0.02	10	-0.09	10	-0.01
Kd of Ra-226 in Unsaturated Zone 1	12	-0.10	11	-0.03	11	-0.09	11	-0.01
Kd of Ra-226 in Saturated Zone	14	-0.08	14	-0.02	19	0.00	19	0.00
Plant transfer factor for Pb	3	0.31	2	0.08	3	0.26	3	0.03
Meat transfer factor for Pb	20	-0.01	20	0.00	18	0.01	18	0.00
Milk transfer factor for Pb	9	-0.14	9	-0.03	9	0.09	9	0.01
Fish transfer factor for Pb	5	-0.21	5	-0.05	14	-0.06	14	-0.01
Plant transfer factor for Ra	18	-0.05	18	-0.01	15	0.05	15	0.01
Meat transfer factor for Ra	21	0.01	21	0.00	12	-0.07	12	-0.01
Milk transfer factor for Ra	10	0.11	10	0.03	22	0.00	22	0.00
Fish transfer factor for Ra	15	-0.07	15	-0.02	7	0.10	7	0.01
Well pumping rate	22	0.00	22	0.00	21	0.00	21	0.00
Mass loading for inhalation	2	-0.31	3	-0.08	2	-0.27	2	-0.03
Indoor dust filtration factor	4	-0.27	4	-0.07	20	0.00	20	0.00
Depth of soil mixing layer	1	-0.97	1	-0.96	1	-0.99	1	-0.99
Depth of roots	11	0.11	12	0.03	8	0.09	8	0.01
Wet weight crop yield of fruit, grain and non-leafy vegetables	17	0.06	17	0.01	17	-0.02	17	0.00
Weathering removal constant of all vegetation	8	-0.14	8	-0.04	5	-0.17	5	-0.02
Wet foliar interception fraction of leafy vegetables	7	0.17	7	0.04	16	-0.03	16	0.00
R-SQUARE		0.95		0.95		0.99		0.99

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak Soil Ingestion Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC		SRC		PRCC		SRRC	
	Sig	Coeff	Sig	Coeff	Sig	Coeff	Sig	Coeff
Kd of Pb-210 in Contaminated Zone	15	-0.07	15	-0.02	2	0.36	2	0.06
Kd of Pb-210 in Unsaturated Zone 1	4	-0.16	4	-0.05	7	-0.13	7	-0.02
Kd of Pb-210 in Saturated Zone	8	0.13	9	0.04	15	-0.08	15	-0.01
Kd of Ra-226 in Contaminated Zone	21	0.04	21	0.01	9	0.12	9	0.02
Kd of Ra-226 in Unsaturated Zone 1	3	0.18	3	0.05	3	0.19	3	0.03
Kd of Ra-226 in Saturated Zone	22	-0.01	22	0.00	16	0.08	16	0.01
Plant transfer factor for Pb	20	0.04	20	0.01	10	-0.11	10	-0.02
Meat transfer factor for Pb	12	0.11	11	0.03	6	0.13	6	0.02
Milk transfer factor for Pb	11	0.11	12	0.03	18	-0.07	18	-0.01
Fish transfer factor for Pb	13	-0.08	13	-0.02	4	0.14	4	0.02
Plant transfer factor for Ra	10	-0.12	10	-0.03	5	0.14	5	0.02
Meat transfer factor for Ra	14	-0.08	14	-0.02	20	-0.02	20	0.00
Milk transfer factor for Ra	17	-0.07	16	-0.02	17	-0.07	17	-0.01
Fish transfer factor for Ra	9	-0.13	5	-0.04	14	-0.08	14	-0.01
Well pumping rate	18	-0.06	18	-0.02	12	-0.10	12	-0.01
Mass loading for inhalation	5	-0.15	6	-0.04	8	-0.12	8	-0.02
Indoor dust filtration factor	19	-0.05	19	-0.01	22	-0.01	22	0.00
Depth of soil mixing layer	1	-0.97	1	-0.97	1	-0.99	1	-0.99
Depth of roots	16	-0.07	17	-0.02	11	-0.10	11	-0.02
Wet weight crop yield of fruit, grain and non-leafy vegetables	2	-0.18	2	-0.05	21	0.02	21	0.00
Weathering removal constant of all vegetation	6	0.15	8	0.04	13	0.09	13	0.01
Wet foliar interception fraction of leafy vegetables	7	-0.14	7	-0.04	19	0.03	19	0.00
R-SQUARE		0.94		0.94		0.98		0.98

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.



Coefficients for peak Soil Ingestion Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC		SRC		PRCC		SRRC	
	Sig	Coeff	Sig	Coeff	Sig	Coeff	Sig	Coeff
Kd of Pb-210 in Contaminated Zone	17	0.03	17	0.01	3	0.22	3	0.03
Kd of Pb-210 in Unsaturated Zone 1	6	0.16	7	0.04	17	0.06	17	0.01
Kd of Pb-210 in Saturated Zone	9	0.14	8	0.04	16	0.09	16	0.01
Kd of Ra-226 in Contaminated Zone	12	-0.12	11	-0.03	9	0.13	9	0.02
Kd of Ra-226 in Unsaturated Zone 1	14	-0.07	15	-0.02	22	0.01	22	0.00
Kd of Ra-226 in Saturated Zone	7	-0.16	6	-0.04	6	-0.15	6	-0.02
Plant transfer factor for Pb	18	0.03	18	0.01	5	-0.19	5	-0.02
Meat transfer factor for Pb	20	-0.02	20	0.00	13	-0.11	13	-0.01
Milk transfer factor for Pb	2	-0.31	2	-0.08	10	-0.12	10	-0.02
Fish transfer factor for Pb	10	0.14	10	0.03	14	-0.10	14	-0.01
Plant transfer factor for Ra	5	-0.17	5	-0.04	11	0.11	11	0.01
Meat transfer factor for Ra	11	0.12	12	0.03	4	0.20	4	0.03
Milk transfer factor for Ra	19	-0.02	19	-0.01	8	-0.13	8	-0.02
Fish transfer factor for Ra	13	0.07	13	0.02	2	0.24	2	0.03
Well pumping rate	3	-0.28	3	-0.07	7	-0.13	7	-0.02
Mass loading for inhalation	16	-0.06	14	-0.02	12	0.11	12	0.01
Indoor dust filtration factor	22	-0.01	22	0.00	21	-0.02	21	0.00
Depth of soil mixing layer	1	-0.97	1	-0.95	1	-0.99	1	-0.99
Depth of roots	8	-0.15	9	-0.04	18	-0.04	18	-0.01
Wet weight crop yield of fruit, grain and non-leafy vegetables	15	0.06	16	0.02	19	0.04	19	0.00
Weathering removal constant of all vegetation	21	0.02	21	0.00	15	0.09	15	0.01
Wet foliar interception fraction of leafy vegetables	4	0.20	4	0.05	20	0.02	20	0.00
R-SQUARE		0.94		0.94		0.98		0.98

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak Water Ingestion Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC		SRC		PRCC		SRRC	
	Sig	Coeff	Sig	Coeff	Sig	Coeff	Sig	Coeff
Kd of Pb-210 in Contaminated Zone	14	0.05	14	0.05	6	-0.12	6	-0.10
Kd of Pb-210 in Unsaturated Zone 1	5	-0.11	5	-0.11	1	-0.40	1	-0.37
Kd of Pb-210 in Saturated Zone	22	-0.01	22	-0.01	13	-0.08	13	-0.07
Kd of Ra-226 in Contaminated Zone	21	0.01	21	0.01	15	0.05	15	0.04
Kd of Ra-226 in Unsaturated Zone 1	15	0.04	15	0.05	16	0.04	16	0.03
Kd of Ra-226 in Saturated Zone	13	-0.06	13	-0.06	22	-0.01	22	-0.01
Plant transfer factor for Pb	18	0.02	18	0.02	2	0.27	2	0.24
Meat transfer factor for Pb	2	0.20	2	0.20	8	-0.10	8	-0.09
Milk transfer factor for Pb	1	0.22	1	0.22	21	-0.02	21	-0.02
Fish transfer factor for Pb	17	-0.02	17	-0.02	18	0.03	18	0.02
Plant transfer factor for Ra	7	-0.10	7	-0.09	17	0.03	17	0.03
Meat transfer factor for Ra	12	-0.06	11	-0.07	14	-0.05	14	-0.04
Milk transfer factor for Ra	10	0.08	10	0.08	12	0.08	12	0.07
Fish transfer factor for Ra	8	-0.09	8	-0.09	7	-0.12	7	-0.10
Well pumping rate	6	0.10	6	0.10	19	0.03	19	0.02
Mass loading for inhalation	19	0.01	19	0.01	11	-0.09	11	-0.07
Indoor dust filtration factor	11	0.07	12	0.06	3	0.22	3	0.19
Depth of soil mixing layer	20	-0.01	20	-0.01	4	-0.15	4	-0.13
Depth of roots	3	0.18	3	0.18	5	0.14	5	0.12
Wet weight crop yield of fruit, grain and non-leafy vegetables	16	-0.02	16	-0.02	10	-0.09	10	-0.08
Weathering removal constant of all vegetation	9	-0.09	9	-0.09	20	0.03	20	0.02
Wet foliar interception fraction of leafy vegetables	4	-0.12	4	-0.11	9	0.10	9	0.09
R-SQUARE		0.14		0.14		0.30		0.30

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak Water Ingestion Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC		SRC		PRCC		SRRC	
	Sig	Coef	Sig	Coef	Sig	Coef	Sig	Coef
Kd of Pb-210 in Contaminated Zone	21	-0.01	21	-0.01	17	-0.05	17	-0.04
Kd of Pb-210 in Unsaturated Zone 1	22	0.00	22	0.00	1	-0.39	1	-0.34
Kd of Pb-210 in Saturated Zone	7	-0.11	7	-0.11	19	-0.05	19	-0.04
Kd of Ra-226 in Contaminated Zone	20	0.01	20	0.01	15	0.07	15	0.06
Kd of Ra-226 in Unsaturated Zone 1	15	-0.02	15	-0.02	20	-0.03	20	-0.02
Kd of Ra-226 in Saturated Zone	12	-0.05	11	-0.06	2	0.28	2	0.24
Plant transfer factor for Pb	8	-0.10	8	-0.10	16	0.06	16	0.05
Meat transfer factor for Pb	1	0.22	1	0.23	3	0.24	3	0.20
Milk transfer factor for Pb	17	-0.01	17	-0.01	21	0.02	21	0.02
Fish transfer factor for Pb	19	0.01	19	0.01	11	0.11	11	0.09
Plant transfer factor for Ra	6	-0.12	6	-0.11	4	-0.20	4	-0.17
Meat transfer factor for Ra	3	0.15	3	0.15	8	0.13	8	0.10
Milk transfer factor for Ra	14	0.04	14	0.04	12	0.09	12	0.07
Fish transfer factor for Ra	11	-0.05	12	-0.06	9	-0.12	10	-0.10
Well pumping rate	16	0.02	16	0.01	18	0.05	18	0.04
Mass loading for inhalation	13	-0.04	13	-0.04	13	-0.09	13	-0.07
Indoor dust filtration factor	2	0.16	2	0.16	22	-0.01	22	0.00
Depth of soil mixing layer	9	0.09	9	0.09	6	-0.13	6	-0.11
Depth of roots	18	-0.01	18	-0.01	14	-0.08	14	-0.06
Wet weight crop yield of fruit, grain and non-leafy vegetables	4	-0.14	5	-0.13	5	-0.17	5	-0.14
Weathering removal constant of all vegetation	5	-0.13	4	-0.13	10	0.12	9	0.10
Wet foliar interception fraction of leafy vegetables	10	-0.08	10	-0.08	7	-0.13	7	-0.10
R-SQUARE		0.14		0.14		0.36		0.36

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak Water Ingestion Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC		SRC		PRCC		SRRC	
	Sig	Coef	Sig	Coef	Sig	Coef	Sig	Coef
Kd of Pb-210 in Contaminated Zone	7	0.12	4	0.14	19	-0.01	19	-0.01
Kd of Pb-210 in Unsaturated Zone 1	22	-0.01	22	-0.01	1	-0.38	1	-0.35
Kd of Pb-210 in Saturated Zone	15	-0.05	15	-0.05	21	0.00	21	0.00
Kd of Ra-226 in Contaminated Zone	21	-0.01	21	-0.01	6	0.12	6	0.11
Kd of Ra-226 in Unsaturated Zone 1	11	-0.07	11	-0.07	10	0.07	10	0.06
Kd of Ra-226 in Saturated Zone	16	0.05	16	0.05	2	0.26	2	0.23
Plant transfer factor for Pb	3	-0.14	5	-0.13	15	0.02	15	0.02
Meat transfer factor for Pb	18	-0.03	18	-0.03	16	-0.02	16	-0.02
Milk transfer factor for Pb	8	-0.10	8	-0.10	9	-0.08	9	-0.07
Fish transfer factor for Pb	17	0.04	17	0.04	4	0.19	4	0.17
Plant transfer factor for Ra	12	-0.07	13	-0.06	18	-0.02	18	-0.01
Meat transfer factor for Ra	19	0.03	19	0.03	8	-0.09	8	-0.08
Milk transfer factor for Ra	2	-0.16	2	-0.16	7	-0.10	7	-0.09
Fish transfer factor for Ra	14	-0.06	12	-0.06	14	0.04	14	0.03
Well pumping rate	6	-0.12	7	-0.12	5	-0.13	5	-0.11
Mass loading for inhalation	4	-0.13	3	-0.15	3	-0.22	3	-0.19
Indoor dust filtration factor	1	-0.17	1	-0.16	20	0.01	20	0.01
Depth of soil mixing layer	9	-0.09	9	-0.08	13	-0.04	13	-0.03
Depth of roots	20	-0.02	20	-0.02	11	0.07	11	0.06
Wet weight crop yield of fruit, grain and non-leafy vegetables	13	-0.07	14	-0.06	12	-0.07	12	-0.06
Weathering removal constant of all vegetation	10	0.07	10	0.07	22	0.00	22	0.00
Wet foliar interception fraction of leafy vegetables	5	0.13	6	0.13	17	0.02	17	0.02
R-SQUARE		0.14		0.14		0.28		0.28

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak Fish Ingestion Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC		SRC		PRCC		SRRC	
	Sig	Coeff	Sig	Coeff	Sig	Coeff	Sig	Coeff
Kd of Pb-210 in Contaminated Zone	14	0.05	14	0.05	6	-0.12	6	-0.10
Kd of Pb-210 in Unsaturated Zone 1	5	-0.11	5	-0.11	1	-0.40	1	-0.37
Kd of Pb-210 in Saturated Zone	22	-0.01	22	-0.01	13	-0.08	13	-0.07
Kd of Ra-226 in Contaminated Zone	21	0.01	21	0.01	15	0.05	15	0.04
Kd of Ra-226 in Unsaturated Zone 1	15	0.04	15	0.05	16	0.04	16	0.03
Kd of Ra-226 in Saturated Zone	13	-0.06	13	-0.06	22	-0.01	22	-0.01
Plant transfer factor for Pb	18	0.02	18	0.02	2	0.27	2	0.24
Meat transfer factor for Pb	2	0.20	2	0.20	9	-0.10	9	-0.09
Milk transfer factor for Pb	1	0.22	1	0.22	21	-0.02	21	-0.02
Fish transfer factor for Pb	17	-0.02	17	-0.02	18	0.03	18	0.03
Plant transfer factor for Ra	7	-0.10	7	-0.09	17	0.03	17	0.03
Meat transfer factor for Ra	12	-0.06	11	-0.07	14	-0.05	14	-0.04
Milk transfer factor for Ra	10	0.08	10	0.08	12	0.08	12	0.07
Fish transfer factor for Ra	8	-0.09	8	-0.09	7	-0.12	7	-0.10
Well pumping rate	6	0.10	6	0.10	20	0.02	20	0.02
Mass loading for inhalation	19	0.01	19	0.01	11	-0.09	11	-0.07
Indoor dust filtration factor	11	0.07	12	0.06	3	0.22	3	0.19
Depth of soil mixing layer	20	-0.01	20	-0.01	4	-0.15	4	-0.13
Depth of roots	3	0.18	3	0.18	5	0.14	5	0.12
Wet weight crop yield of fruit, grain and non-leafy vegetables	16	-0.02	16	-0.02	10	-0.09	10	-0.08
Weathering removal constant of all vegetation	9	-0.09	9	-0.09	19	0.03	19	0.02
Wet foliar interception fraction of leafy vegetables	4	-0.12	4	-0.11	8	0.10	8	0.09
R-SQUARE		0.14		0.14		0.30		0.30

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak Fish Ingestion Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC		SRC		PRCC		SRRC	
	Sig	Coef	Sig	Coef	Sig	Coef	Sig	Coef
Kd of Pb-210 in Contaminated Zone	21	-0.01	21	-0.01	17	-0.05	17	-0.04
Kd of Pb-210 in Unsaturated Zone 1	22	0.00	22	0.00	1	-0.39	1	-0.34
Kd of Pb-210 in Saturated Zone	7	-0.11	7	-0.11	19	-0.05	19	-0.04
Kd of Ra-226 in Contaminated Zone	20	0.01	20	0.01	15	0.07	15	0.06
Kd of Ra-226 in Unsaturated Zone 1	15	-0.02	15	-0.02	20	-0.03	20	-0.02
Kd of Ra-226 in Saturated Zone	12	-0.05	11	-0.06	2	0.28	2	0.24
Plant transfer factor for Pb	8	-0.10	8	-0.10	16	0.06	16	0.05
Meat transfer factor for Pb	1	0.22	1	0.23	3	0.24	3	0.20
Milk transfer factor for Pb	17	-0.01	17	-0.01	21	0.02	21	0.02
Fish transfer factor for Pb	19	0.01	19	0.01	11	0.11	11	0.09
Plant transfer factor for Ra	6	-0.12	6	-0.11	4	-0.20	4	-0.17
Meat transfer factor for Ra	3	0.15	3	0.15	8	0.13	8	0.10
Milk transfer factor for Ra	14	0.04	14	0.04	12	0.09	12	0.07
Fish transfer factor for Ra	11	-0.05	12	-0.06	9	-0.12	10	-0.10
Well pumping rate	16	0.02	16	0.01	18	0.05	18	0.04
Mass loading for inhalation	13	-0.04	13	-0.04	13	-0.09	13	-0.07
Indoor dust filtration factor	2	0.16	2	0.16	22	-0.01	22	0.00
Depth of soil mixing layer	9	0.09	9	0.09	6	-0.13	6	-0.11
Depth of roots	18	-0.01	18	-0.01	14	-0.08	14	-0.06
Wet weight crop yield of fruit, grain and non-leafy vegetables	4	-0.14	5	-0.13	5	-0.17	5	-0.14
Weathering removal constant of all vegetation	5	-0.13	4	-0.13	10	0.12	9	0.10
Wet foliar interception fraction of leafy vegetables	10	-0.08	10	-0.08	7	-0.13	7	-0.10
R-SQUARE		0.14		0.14		0.36		0.36

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak Fish Ingestion Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC		SRC		PRCC		SRRC	
	Sig	Coef	Sig	Coef	Sig	Coef	Sig	Coef
Kd of Pb-210 in Contaminated Zone	7	0.12	4	0.14	19	-0.01	19	-0.01
Kd of Pb-210 in Unsaturated Zone 1	22	-0.01	22	-0.01	1	-0.38	1	-0.35
Kd of Pb-210 in Saturated Zone	15	-0.05	15	-0.05	21	0.00	21	0.00
Kd of Ra-226 in Contaminated Zone	21	-0.01	21	-0.01	6	0.12	6	0.11
Kd of Ra-226 in Unsaturated Zone 1	11	-0.07	11	-0.07	10	0.07	10	0.06
Kd of Ra-226 in Saturated Zone	16	0.05	16	0.05	2	0.26	2	0.23
Plant transfer factor for Pb	3	-0.13	5	-0.13	15	0.02	15	0.02
Meat transfer factor for Pb	18	-0.03	18	-0.03	16	-0.02	16	-0.02
Milk transfer factor for Pb	8	-0.10	8	-0.10	9	-0.08	9	-0.07
Fish transfer factor for Pb	17	0.05	17	0.04	4	0.19	4	0.17
Plant transfer factor for Ra	12	-0.07	13	-0.06	18	-0.02	18	-0.01
Meat transfer factor for Ra	19	0.03	19	0.03	8	-0.09	8	-0.08
Milk transfer factor for Ra	2	-0.16	2	-0.16	7	-0.10	7	-0.09
Fish transfer factor for Ra	14	-0.06	12	-0.06	14	0.04	14	0.03
Well pumping rate	6	-0.12	7	-0.12	5	-0.13	5	-0.11
Mass loading for inhalation	4	-0.13	3	-0.15	3	-0.22	3	-0.19
Indoor dust filtration factor	1	-0.17	1	-0.16	20	0.01	20	0.01
Depth of soil mixing layer	9	-0.09	9	-0.08	13	-0.04	13	-0.03
Depth of roots	20	-0.02	20	-0.02	11	0.07	11	0.06
Wet weight crop yield of fruit, grain and non-leafy vegetables	13	-0.06	14	-0.06	12	-0.07	12	-0.06
Weathering removal constant of all vegetation	10	0.07	10	0.07	22	0.00	22	0.00
Wet foliar interception fraction of leafy vegetables	5	0.13	6	0.13	17	0.02	17	0.02
R-SQUARE		0.14		0.14		0.28		0.28

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak Radon (WaterDep.) Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC		SRC		PRCC		SRRC	
	Sig	Coeff	Sig	Coeff	Sig	Coeff	Sig	Coeff
Kd of Pb-210 in Contaminated Zone	0	0.00	0	0.00	0	0.00	0	0.00
Kd of Pb-210 in Unsaturated Zone 1	0	0.00	0	0.00	0	0.00	0	0.00
Kd of Pb-210 in Saturated Zone	0	0.00	0	0.00	0	0.00	0	0.00
Kd of Ra-226 in Contaminated Zone	0	0.00	0	0.00	0	0.00	0	0.00
Kd of Ra-226 in Unsaturated Zone 1	0	0.00	0	0.00	0	0.00	0	0.00
Kd of Ra-226 in Saturated Zone	0	0.00	0	0.00	0	0.00	0	0.00
Plant transfer factor for Pb	0	0.00	0	0.00	0	0.00	0	0.00
Meat transfer factor for Pb	0	0.00	0	0.00	0	0.00	0	0.00
Milk transfer factor for Pb	0	0.00	0	0.00	0	0.00	0	0.00
Fish transfer factor for Pb	0	0.00	0	0.00	0	0.00	0	0.00
Plant transfer factor for Ra	0	0.00	0	0.00	0	0.00	0	0.00
Meat transfer factor for Ra	0	0.00	0	0.00	0	0.00	0	0.00
Milk transfer factor for Ra	0	0.00	0	0.00	0	0.00	0	0.00
Fish transfer factor for Ra	0	0.00	0	0.00	0	0.00	0	0.00
Well pumping rate	0	0.00	0	0.00	0	0.00	0	0.00
Mass loading for inhalation	0	0.00	0	0.00	0	0.00	0	0.00
Indoor dust filtration factor	0	0.00	0	0.00	0	0.00	0	0.00
Depth of soil mixing layer	0	0.00	0	0.00	0	0.00	0	0.00
Depth of roots	0	0.00	0	0.00	0	0.00	0	0.00
Wet weight crop yield of fruit, grain and non-leafy vegetables	0	0.00	0	0.00	0	0.00	0	0.00
Weathering removal constant of all vegetation	0	0.00	0	0.00	0	0.00	0	0.00
Wet foliar interception fraction of leafy vegetables	0	0.00	0	0.00	0	0.00	0	0.00
R-SQUARE		0.00		0.00		0.00		0.00

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.



Coefficients for peak Radon (WaterDep.) Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC		SRC		PRCC		SRRC	
	Sig	Coeff	Sig	Coeff	Sig	Coeff	Sig	Coeff
Kd of Pb-210 in Contaminated Zone	0	0.00	0	0.00	0	0.00	0	0.00
Kd of Pb-210 in Unsaturated Zone 1	0	0.00	0	0.00	0	0.00	0	0.00
Kd of Pb-210 in Saturated Zone	0	0.00	0	0.00	0	0.00	0	0.00
Kd of Ra-226 in Contaminated Zone	0	0.00	0	0.00	0	0.00	0	0.00
Kd of Ra-226 in Unsaturated Zone 1	0	0.00	0	0.00	0	0.00	0	0.00
Kd of Ra-226 in Saturated Zone	0	0.00	0	0.00	0	0.00	0	0.00
Plant transfer factor for Pb	0	0.00	0	0.00	0	0.00	0	0.00
Meat transfer factor for Pb	0	0.00	0	0.00	0	0.00	0	0.00
Milk transfer factor for Pb	0	0.00	0	0.00	0	0.00	0	0.00
Fish transfer factor for Pb	0	0.00	0	0.00	0	0.00	0	0.00
Plant transfer factor for Ra	0	0.00	0	0.00	0	0.00	0	0.00
Meat transfer factor for Ra	0	0.00	0	0.00	0	0.00	0	0.00
Milk transfer factor for Ra	0	0.00	0	0.00	0	0.00	0	0.00
Fish transfer factor for Ra	0	0.00	0	0.00	0	0.00	0	0.00
Well pumping rate	0	0.00	0	0.00	0	0.00	0	0.00
Mass loading for inhalation	0	0.00	0	0.00	0	0.00	0	0.00
Indoor dust filtration factor	0	0.00	0	0.00	0	0.00	0	0.00
Depth of soil mixing layer	0	0.00	0	0.00	0	0.00	0	0.00
Depth of roots	0	0.00	0	0.00	0	0.00	0	0.00
Wet weight crop yield of fruit, grain and non-leafy vegetables	0	0.00	0	0.00	0	0.00	0	0.00
Weathering removal constant of all vegetation	0	0.00	0	0.00	0	0.00	0	0.00
Wet foliar interception fraction of leafy vegetables	0	0.00	0	0.00	0	0.00	0	0.00
R-SQUARE		0.00		0.00		0.00		0.00

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak Radon (WaterDep.) Dose		PCC		SRC		PRCC		SRRC	
Coefficient =		3		3		3		3	
Repetition =									
Description of Probabilistic Variable		Sig	Coeff	Sig	Coeff	Sig	Coeff	Sig	Coeff
Kd of Pb-210 in Contaminated Zone		0	0.00	0	0.00	0	0.00	0	0.00
Kd of Pb-210 in Unsaturated Zone 1		0	0.00	0	0.00	0	0.00	0	0.00
Kd of Pb-210 in Saturated Zone		0	0.00	0	0.00	0	0.00	0	0.00
Kd of Ra-226 in Contaminated Zone		0	0.00	0	0.00	0	0.00	0	0.00
Kd of Ra-226 in Unsaturated Zone 1		0	0.00	0	0.00	0	0.00	0	0.00
Kd of Ra-226 in Saturated Zone		0	0.00	0	0.00	0	0.00	0	0.00
Plant transfer factor for Pb		0	0.00	0	0.00	0	0.00	0	0.00
Meat transfer factor for Pb		0	0.00	0	0.00	0	0.00	0	0.00
Milk transfer factor for Pb		0	0.00	0	0.00	0	0.00	0	0.00
Fish transfer factor for Pb		0	0.00	0	0.00	0	0.00	0	0.00
Plant transfer factor for Ra		0	0.00	0	0.00	0	0.00	0	0.00
Meat transfer factor for Ra		0	0.00	0	0.00	0	0.00	0	0.00
Milk transfer factor for Ra		0	0.00	0	0.00	0	0.00	0	0.00
Fish transfer factor for Ra		0	0.00	0	0.00	0	0.00	0	0.00
Well pumping rate		0	0.00	0	0.00	0	0.00	0	0.00
Mass loading for inhalation		0	0.00	0	0.00	0	0.00	0	0.00
Indoor dust filtration factor		0	0.00	0	0.00	0	0.00	0	0.00
Depth of soil mixing layer		0	0.00	0	0.00	0	0.00	0	0.00
Depth of roots		0	0.00	0	0.00	0	0.00	0	0.00
Wet weight crop yield of fruit, grain and non-leafy vegetables		0	0.00	0	0.00	0	0.00	0	0.00
Weathering removal constant of all vegetation		0	0.00	0	0.00	0	0.00	0	0.00
Wet foliar interception fraction of leafy vegetables		0	0.00	0	0.00	0	0.00	0	0.00
R-SQUARE		0.00		0.00		0.00		0.00	

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak Plant (WaterDep.) Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC		SRC		PRCC		SRRC	
	Sig	Coeff	Sig	Coeff	Sig	Coeff	Sig	Coeff
Kd of Pb-210 in Contaminated Zone	14	0.05	14	0.05	6	-0.12	6	-0.10
Kd of Pb-210 in Unsaturated Zone 1	5	-0.11	5	-0.11	1	-0.40	1	-0.37
Kd of Pb-210 in Saturated Zone	22	-0.01	22	-0.01	13	-0.08	13	-0.07
Kd of Ra-226 in Contaminated Zone	21	0.01	21	0.01	15	0.05	15	0.04
Kd of Ra-226 in Unsaturated Zone 1	15	0.04	15	0.05	16	0.04	16	0.03
Kd of Ra-226 in Saturated Zone	13	-0.06	13	-0.06	22	-0.01	22	-0.01
Plant transfer factor for Pb	18	0.02	18	0.02	2	0.27	2	0.24
Meat transfer factor for Pb	2	0.20	2	0.20	8	-0.10	8	-0.09
Milk transfer factor for Pb	1	0.22	1	0.22	21	-0.02	21	-0.02
Fish transfer factor for Pb	17	-0.02	17	-0.02	18	0.03	18	0.02
Plant transfer factor for Ra	7	-0.10	7	-0.09	17	0.03	17	0.03
Meat transfer factor for Ra	12	-0.06	11	-0.07	14	-0.05	14	-0.04
Milk transfer factor for Ra	10	0.08	10	0.08	12	0.08	12	0.07
Fish transfer factor for Ra	8	-0.09	8	-0.09	7	-0.12	7	-0.10
Well pumping rate	6	0.10	6	0.10	19	0.03	19	0.02
Mass loading for inhalation	19	0.01	19	0.01	11	-0.09	11	-0.07
Indoor dust filtration factor	11	0.07	12	0.06	3	0.22	3	0.19
Depth of soil mixing layer	20	-0.01	20	-0.01	4	-0.15	4	-0.13
Depth of roots	3	0.18	3	0.18	5	0.14	5	0.12
Wet weight crop yield of fruit, grain and non-leafy vegetables	16	-0.02	16	-0.02	10	-0.09	10	-0.08
Weathering removal constant of all vegetation	9	-0.09	9	-0.09	20	0.03	20	0.02
Wet foliar interception fraction of leafy vegetables	4	-0.12	4	-0.11	9	0.10	9	0.09
R-SQUARE		0.14		0.14		0.30		0.30

-Rank is set to zero if the dose is zero or the correlation matrix is singular.

-R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak Plant (WaterDep.) Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC		SRC		PRCC		SRRC	
	Sig	Coef	Sig	Coef	Sig	Coef	Sig	Coef
Kd of Pb-210 in Contaminated Zone	21	-0.01	21	-0.01	17	-0.05	17	-0.04
Kd of Pb-210 in Unsaturated Zone 1	22	0.00	22	0.00	1	-0.39	1	-0.34
Kd of Pb-210 in Saturated Zone	7	-0.11	7	-0.11	19	-0.05	19	-0.04
Kd of Ra-226 in Contaminated Zone	20	0.01	20	0.01	15	0.07	15	0.06
Kd of Ra-226 in Unsaturated Zone 1	15	-0.02	15	-0.02	20	-0.03	20	-0.02
Kd of Ra-226 in Saturated Zone	12	-0.05	11	-0.06	2	0.28	2	0.24
Plant transfer factor for Pb	8	-0.10	8	-0.10	16	0.06	16	0.05
Meat transfer factor for Pb	1	0.22	1	0.23	3	0.24	3	0.20
Milk transfer factor for Pb	17	-0.01	17	-0.01	21	0.02	21	0.02
Fish transfer factor for Pb	19	0.01	19	0.01	11	0.11	11	0.09
Plant transfer factor for Ra	6	-0.12	6	-0.11	4	-0.20	4	-0.17
Meat transfer factor for Ra	3	0.15	3	0.15	8	0.13	8	0.10
Milk transfer factor for Ra	14	0.04	14	0.04	12	0.09	12	0.07
Fish transfer factor for Ra	11	-0.05	12	-0.06	9	-0.12	10	-0.10
Well pumping rate	16	0.02	16	0.01	18	0.05	18	0.04
Mass loading for inhalation	13	-0.04	13	-0.04	13	-0.09	13	-0.07
Indoor dust filtration factor	2	0.16	2	0.16	22	-0.01	22	0.00
Depth of soil mixing layer	9	0.09	9	0.09	6	-0.13	6	-0.11
Depth of roots	18	-0.01	18	-0.01	14	-0.08	14	-0.06
Wet weight crop yield of fruit, grain and non-leafy vegetables	4	-0.14	5	-0.13	5	-0.17	5	-0.14
Weathering removal constant of all vegetation	5	-0.13	4	-0.13	10	0.12	9	0.10
Wet foliar interception fraction of leafy vegetables	10	-0.08	10	-0.08	7	-0.13	7	-0.10
R-SQUARE		0.14		0.14		0.36		0.36

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak Plant (WaterDep.) Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC 3		SRC 3		PRCC 3		SRRC 3	
	Sig	Coeff	Sig	Coeff	Sig	Coeff	Sig	Coeff
Kd of Pb-210 in Contaminated Zone	7	0.12	4	0.14	19	-0.01	19	-0.01
Kd of Pb-210 in Unsaturated Zone 1	22	-0.01	22	-0.01	1	-0.38	1	-0.35
Kd of Pb-210 in Saturated Zone	15	-0.05	15	-0.05	21	0.00	21	0.00
Kd of Ra-226 in Contaminated Zone	21	-0.01	21	-0.01	6	0.12	6	0.11
Kd of Ra-226 in Unsaturated Zone 1	11	-0.07	11	-0.07	10	0.07	10	0.06
Kd of Ra-226 in Saturated Zone	16	0.05	16	0.05	2	0.26	2	0.23
Plant transfer factor for Pb	3	-0.13	5	-0.13	15	0.02	15	0.02
Meat transfer factor for Pb	18	-0.03	18	-0.03	16	-0.02	16	-0.02
Milk transfer factor for Pb	8	-0.10	8	-0.10	9	-0.08	9	-0.07
Fish transfer factor for Pb	17	0.04	17	0.04	4	0.19	4	0.17
Plant transfer factor for Ra	12	-0.07	13	-0.06	18	-0.02	18	-0.01
Meat transfer factor for Ra	19	0.03	19	0.03	8	-0.09	8	-0.08
Milk transfer factor for Ra	2	-0.16	2	-0.16	7	-0.10	7	-0.09
Fish transfer factor for Ra	14	-0.06	12	-0.06	14	0.04	14	0.03
Well pumping rate	6	-0.12	7	-0.12	5	-0.13	5	-0.11
Mass loading for inhalation	4	-0.13	3	-0.15	3	-0.22	3	-0.19
Indoor dust filtration factor	1	-0.17	1	-0.16	20	0.01	20	0.01
Depth of soil mixing layer	9	-0.09	9	-0.08	13	-0.04	13	-0.03
Depth of roots	20	-0.02	20	-0.02	11	0.07	11	0.06
Wet weight crop yield of fruit, grain and non-leafy vegetables	13	-0.07	14	-0.06	12	-0.07	12	-0.06
Weathering removal constant of all vegetation	10	0.07	10	0.07	22	0.00	22	0.00
Wet foliar interception fraction of leafy vegetables	5	0.13	6	0.13	17	0.02	17	0.02
R-SQUARE		0.14		0.14		0.28		0.28

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak Meat (WaterDep.) Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC		SRC		PRCC		SRRC	
	Sig	Coeff	Sig	Coeff	Sig	Coeff	Sig	Coeff
Kd of Pb-210 in Contaminated Zone	14	0.05	14	0.05	6	-0.12	6	-0.10
Kd of Pb-210 in Unsaturated Zone 1	5	-0.11	5	-0.11	1	-0.40	1	-0.37
Kd of Pb-210 in Saturated Zone	22	-0.01	22	-0.01	13	-0.08	13	-0.07
Kd of Ra-226 in Contaminated Zone	21	0.01	21	0.01	15	0.05	15	0.04
Kd of Ra-226 in Unsaturated Zone 1	15	0.04	15	0.05	16	0.04	16	0.03
Kd of Ra-226 in Saturated Zone	13	-0.06	13	-0.06	22	-0.01	22	-0.01
Plant transfer factor for Pb	18	0.02	18	0.02	2	0.27	2	0.24
Meat transfer factor for Pb	2	0.20	2	0.20	9	-0.10	9	-0.09
Milk transfer factor for Pb	1	0.22	1	0.22	21	-0.02	21	-0.02
Fish transfer factor for Pb	17	-0.02	17	-0.02	18	0.03	18	0.03
Plant transfer factor for Ra	7	-0.10	7	-0.09	17	0.03	17	0.03
Meat transfer factor for Ra	12	-0.06	11	-0.07	14	-0.05	14	-0.04
Milk transfer factor for Ra	10	0.08	10	0.08	12	0.08	12	0.07
Fish transfer factor for Ra	8	-0.09	8	-0.09	7	-0.12	7	-0.10
Well pumping rate	6	0.10	6	0.10	20	0.02	20	0.02
Mass loading for inhalation	19	0.01	19	0.01	11	-0.09	11	-0.07
Indoor dust filtration factor	11	0.07	12	0.06	3	0.22	3	0.19
Depth of soil mixing layer	20	-0.01	20	-0.01	4	-0.15	4	-0.13
Depth of roots	3	0.18	3	0.18	5	0.14	5	0.12
Wet weight crop yield of fruit, grain and non-leafy vegetables	16	-0.02	16	-0.02	10	-0.09	10	-0.08
Weathering removal constant of all vegetation	9	-0.09	9	-0.09	19	0.03	19	0.02
Wet foliar interception fraction of leafy vegetables	4	-0.12	4	-0.11	8	0.10	8	0.09
R-SQUARE		0.14		0.14		0.30		0.30

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak Meat (WaterDep.) Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC		SRC		PRCC		SRRC	
	Sig	Coef	Sig	Coef	Sig	Coef	Sig	Coef
Kd of Pb-210 in Contaminated Zone	21	-0.01	21	-0.01	17	-0.05	17	-0.04
Kd of Pb-210 in Unsaturated Zone 1	22	0.00	22	0.00	1	-0.39	1	-0.34
Kd of Pb-210 in Saturated Zone	7	-0.11	7	-0.11	19	-0.05	19	-0.04
Kd of Ra-226 in Contaminated Zone	20	0.01	20	0.01	15	0.07	15	0.06
Kd of Ra-226 in Unsaturated Zone 1	15	-0.02	15	-0.02	20	-0.03	20	-0.02
Kd of Ra-226 in Saturated Zone	12	-0.05	11	-0.06	2	0.28	2	0.24
Plant transfer factor for Pb	8	-0.10	8	-0.10	16	0.06	16	0.05
Meat transfer factor for Pb	1	0.22	1	0.23	3	0.24	3	0.20
Milk transfer factor for Pb	17	-0.01	17	-0.01	21	0.02	21	0.02
Fish transfer factor for Pb	19	0.01	19	0.01	11	0.11	11	0.09
Plant transfer factor for Ra	6	-0.12	6	-0.11	4	-0.20	4	-0.17
Meat transfer factor for Ra	3	0.15	3	0.15	8	0.13	8	0.10
Milk transfer factor for Ra	14	0.04	14	0.04	12	0.09	12	0.07
Fish transfer factor for Ra	11	-0.05	12	-0.06	9	-0.12	10	-0.10
Well pumping rate	16	0.02	16	0.01	18	0.05	18	0.04
Mass loading for inhalation	13	-0.04	13	-0.04	13	-0.09	13	-0.07
Indoor dust filtration factor	2	0.16	2	0.16	22	-0.01	22	0.00
Depth of soil mixing layer	9	0.09	9	0.09	6	-0.13	6	-0.11
Depth of roots	18	-0.01	18	-0.01	14	-0.08	14	-0.06
Wet weight crop yield of fruit, grain and non-leafy vegetables	4	-0.14	5	-0.13	5	-0.17	5	-0.14
Weathering removal constant of all vegetation	5	-0.13	4	-0.13	10	0.12	9	0.10
Wet foliar interception fraction of leafy vegetables	10	-0.08	10	-0.08	7	-0.13	7	-0.10
R-SQUARE		0.14		0.14		0.36		0.36

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak Meat (WaterDep.) Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC		SRC		PRCC		SRRC	
	Sig	Coef	Sig	Coef	Sig	Coef	Sig	Coef
Kd of Pb-210 in Contaminated Zone	7	0.12	4	0.14	19	-0.01	19	-0.01
Kd of Pb-210 in Unsaturated Zone 1	22	-0.01	22	-0.01	1	-0.38	1	-0.35
Kd of Pb-210 in Saturated Zone	15	-0.05	15	-0.05	21	0.00	21	0.00
Kd of Ra-226 in Contaminated Zone	21	-0.01	21	-0.01	6	0.12	6	0.11
Kd of Ra-226 in Unsaturated Zone 1	11	-0.07	11	-0.07	10	0.07	10	0.06
Kd of Ra-226 in Saturated Zone	16	0.05	16	0.05	2	0.26	2	0.23
Plant transfer factor for Pb	3	-0.14	5	-0.13	15	0.02	15	0.02
Meat transfer factor for Pb	18	-0.03	18	-0.03	16	-0.02	16	-0.02
Milk transfer factor for Pb	8	-0.10	8	-0.10	9	-0.08	9	-0.07
Fish transfer factor for Pb	17	0.04	17	0.04	4	0.19	4	0.17
Plant transfer factor for Ra	12	-0.07	13	-0.06	18	-0.02	18	-0.01
Meat transfer factor for Ra	19	0.03	19	0.03	8	-0.09	8	-0.08
Milk transfer factor for Ra	2	-0.16	2	-0.16	7	-0.10	7	-0.09
Fish transfer factor for Ra	14	-0.06	12	-0.06	14	0.04	14	0.03
Well pumping rate	6	-0.12	7	-0.12	5	-0.13	5	-0.11
Mass loading for inhalation	4	-0.13	3	-0.15	3	-0.22	3	-0.19
Indoor dust filtration factor	1	-0.17	1	-0.16	20	0.01	20	0.01
Depth of soil mixing layer	9	-0.09	9	-0.08	13	-0.04	13	-0.03
Depth of roots	20	-0.02	20	-0.02	11	0.07	11	0.06
Wet weight crop yield of fruit, grain and non-leafy vegetables	13	-0.07	14	-0.06	12	-0.07	12	-0.06
Weathering removal constant of all vegetation	10	0.07	10	0.07	22	0.00	22	0.00
Wet foliar interception fraction of leafy vegetables	5	0.13	6	0.13	17	0.02	17	0.02
R-SQUARE		0.14		0.14		0.28		0.28

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.



Coefficients for peak Milk (WaterDep.) Dose		PCC		SRC		PRCC		SRRC	
Coefficient =		1		1		1		1	
Repetition =									
Description of Probabilistic Variable		Sig	Coeff	Sig	Coeff	Sig	Coeff	Sig	Coeff
Kd of Pb-210 in Contaminated Zone		14	0.05	14	0.05	6	-0.12	6	-0.10
Kd of Pb-210 in Unsaturated Zone 1		5	-0.11	5	-0.11	1	-0.40	1	-0.37
Kd of Pb-210 in Saturated Zone		22	-0.01	22	-0.01	13	-0.08	13	-0.07
Kd of Ra-226 in Contaminated Zone		21	0.01	21	0.01	15	0.05	15	0.04
Kd of Ra-226 in Unsaturated Zone 1		15	0.04	15	0.05	16	0.04	16	0.03
Kd of Ra-226 in Saturated Zone		13	-0.06	13	-0.06	22	-0.01	22	-0.01
Plant transfer factor for Pb		18	0.02	18	0.02	2	0.27	2	0.24
Meat transfer factor for Pb		2	0.20	2	0.20	8	-0.10	8	-0.09
Milk transfer factor for Pb		1	0.22	1	0.22	21	-0.02	21	-0.02
Fish transfer factor for Pb		17	-0.02	17	-0.02	18	0.03	18	0.02
Plant transfer factor for Ra		7	-0.10	7	-0.09	17	0.03	17	0.03
Meat transfer factor for Ra		12	-0.06	11	-0.07	14	-0.05	14	-0.04
Milk transfer factor for Ra		10	0.08	10	0.08	12	0.08	12	0.07
Fish transfer factor for Ra		8	-0.09	8	-0.09	7	-0.12	7	-0.10
Well pumping rate		6	0.10	6	0.10	19	0.03	19	0.02
Mass loading for inhalation		19	0.01	19	0.01	11	-0.09	11	-0.07
Indoor dust filtration factor		11	0.07	12	0.06	3	0.22	3	0.19
Depth of soil mixing layer		20	-0.01	20	-0.01	4	-0.15	4	-0.13
Depth of roots		3	0.18	3	0.18	5	0.14	5	0.12
Wet weight crop yield of fruit, grain and non-leafy vegetables		16	-0.02	16	-0.02	10	-0.09	10	-0.08
Weathering removal constant of all vegetation		9	-0.09	9	-0.09	20	0.03	20	0.02
Wet foliar interception fraction of leafy vegetables		4	-0.12	4	-0.11	9	0.10	9	0.09
R-SQUARE			0.14		0.14		0.30		0.30

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak Milk (WaterDep.) Dose		PCC		SRC		PRCC		SRRC	
Coefficient =		2		2		2		2	
Repetition =									
Description of Probabilistic Variable	Sig	Coef	Sig	Coef	Sig	Coef	Sig	Coef	
Kd of Pb-210 in Contaminated Zone	21	-0.01	21	-0.01	17	-0.05	17	-0.04	
Kd of Pb-210 in Unsaturated Zone 1	22	0.00	22	0.00	1	-0.39	1	-0.34	
Kd of Pb-210 in Saturated Zone	7	-0.11	7	-0.11	19	-0.05	19	-0.04	
Kd of Ra-226 in Contaminated Zone	20	0.01	20	0.01	15	0.07	15	0.06	
Kd of Ra-226 in Unsaturated Zone 1	15	-0.02	15	-0.02	20	-0.03	20	-0.02	
Kd of Ra-226 in Saturated Zone	12	-0.05	11	-0.06	2	0.28	2	0.24	
Plant transfer factor for Pb	8	-0.10	8	-0.10	16	0.06	16	0.05	
Meat transfer factor for Pb	1	0.22	1	0.23	3	0.24	3	0.20	
Milk transfer factor for Pb	17	-0.01	17	-0.01	21	0.02	21	0.02	
Fish transfer factor for Pb	19	0.01	19	0.01	11	0.11	11	0.09	
Plant transfer factor for Ra	6	-0.12	6	-0.11	4	-0.20	4	-0.17	
Meat transfer factor for Ra	3	0.15	3	0.15	8	0.13	8	0.10	
Milk transfer factor for Ra	14	0.04	14	0.04	12	0.09	12	0.07	
Fish transfer factor for Ra	11	-0.05	12	-0.06	9	-0.12	10	-0.10	
Well pumping rate	16	0.02	16	0.01	18	0.05	18	0.04	
Mass loading for inhalation	13	-0.04	13	-0.04	13	-0.09	13	-0.07	
Indoor dust filtration factor	2	0.16	2	0.16	22	-0.01	22	0.00	
Depth of soil mixing layer	9	0.09	9	0.09	6	-0.13	6	-0.11	
Depth of roots	18	-0.01	18	-0.01	14	-0.08	14	-0.06	
Wet weight crop yield of fruit, grain and non-leafy vegetables	4	-0.14	5	-0.13	5	-0.17	5	-0.14	
Weathering removal constant of all vegetation	5	-0.13	4	-0.13	10	0.12	9	0.10	
Wet foliar interception fraction of leafy vegetables	10	-0.08	10	-0.08	7	-0.13	7	-0.10	
R-SQUARE		0.14		0.14		0.36		0.36	

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak Milk (WaterDep.) Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC		SRC		PRCC		SRRC	
	Sig	Coef	Sig	Coef	Sig	Coef	Sig	Coef
Kd of Pb-210 in Contaminated Zone	7	0.12	4	0.14	19	-0.01	19	-0.01
Kd of Pb-210 in Unsaturated Zone 1	22	-0.01	22	-0.01	1	-0.38	1	-0.35
Kd of Pb-210 in Saturated Zone	15	-0.05	15	-0.05	21	0.00	21	0.00
Kd of Ra-226 in Contaminated Zone	21	-0.01	21	-0.01	6	0.12	6	0.11
Kd of Ra-226 in Unsaturated Zone 1	11	-0.07	11	-0.07	10	0.07	10	0.06
Kd of Ra-226 in Saturated Zone	16	0.05	16	0.05	2	0.26	2	0.23
Plant transfer factor for Pb	3	-0.13	5	-0.13	15	0.02	15	0.02
Meat transfer factor for Pb	18	-0.03	18	-0.03	16	-0.02	16	-0.02
Milk transfer factor for Pb	8	-0.10	8	-0.10	9	-0.08	9	-0.07
Fish transfer factor for Pb	17	0.04	17	0.04	4	0.19	4	0.17
Plant transfer factor for Ra	12	-0.07	13	-0.06	18	-0.02	18	-0.01
Meat transfer factor for Ra	19	0.03	19	0.03	8	-0.09	8	-0.08
Milk transfer factor for Ra	2	-0.16	2	-0.16	7	-0.10	7	-0.09
Fish transfer factor for Ra	14	-0.06	12	-0.06	14	0.04	14	0.03
Well pumping rate	6	-0.12	7	-0.12	5	-0.13	5	-0.11
Mass loading for inhalation	4	-0.13	3	-0.15	3	-0.22	3	-0.19
Indoor dust filtration factor	1	-0.17	1	-0.16	20	0.01	20	0.01
Depth of soil mixing layer	9	-0.09	9	-0.08	13	-0.04	13	-0.03
Depth of roots	20	-0.02	20	-0.02	11	0.07	11	0.06
Wet weight crop yield of fruit, grain and non-leafy vegetables	13	-0.07	14	-0.06	12	-0.07	12	-0.06
Weathering removal constant of all vegetation	10	0.07	10	0.07	22	0.00	22	0.00
Wet foliar interception fraction of leafy vegetables	5	0.13	6	0.13	17	0.02	17	0.02
R-SQUARE		0.14		0.14		0.28		0.28

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak Pb-210 Dose		PCC		SRC		PRCC		SRRC	
Coefficient =		1		1		1		1	
Repetition =									
Description of Probabilistic Variable	Sig	Coeff	Sig	Coeff	Sig	Coeff	Sig	Coeff	Sig
Kd of Pb-210 in Contaminated Zone	8	-0.13	10	-0.07	17	-0.05	17	-0.01	
Kd of Pb-210 in Unsaturated Zone 1	17	-0.06	17	-0.03	21	0.01	21	0.00	
Kd of Pb-210 in Saturated Zone	20	-0.03	20	-0.02	13	-0.08	13	-0.02	
Kd of Ra-226 in Contaminated Zone	7	-0.14	7	-0.08	16	-0.06	16	-0.02	
Kd of Ra-226 in Unsaturated Zone 1	9	0.13	8	0.08	7	0.14	7	0.04	
Kd of Ra-226 in Saturated Zone	5	-0.22	5	-0.12	8	-0.11	8	-0.03	
Plant transfer factor for Pb	1	0.79	1	0.73	1	0.93	1	0.75	
Meat transfer factor for Pb	11	0.12	11	0.07	5	0.15	5	0.05	
Milk transfer factor for Pb	16	-0.06	16	-0.03	19	0.02	19	0.01	
Fish transfer factor for Pb	3	-0.32	3	-0.19	15	0.07	15	0.02	
Plant transfer factor for Ra	12	-0.12	12	-0.06	10	-0.09	10	-0.03	
Meat transfer factor for Ra	10	-0.12	9	-0.08	12	0.08	12	0.02	
Milk transfer factor for Ra	6	0.16	6	0.09	4	0.20	4	0.06	
Fish transfer factor for Ra	21	-0.02	21	-0.01	6	-0.15	6	-0.04	
Well pumping rate	19	-0.05	19	-0.03	14	0.07	14	0.02	
Mass loading for inhalation	13	-0.12	13	-0.06	20	0.02	20	0.01	
Indoor dust filtration factor	18	-0.05	18	-0.03	9	-0.09	9	-0.03	
Depth of soil mixing layer	4	-0.30	4	-0.17	3	-0.53	3	-0.19	
Depth of roots	2	-0.66	2	-0.47	2	-0.87	2	-0.55	
Wet weight crop yield of fruit, grain and non-leafy vegetables	14	-0.10	14	-0.05	22	0.00	22	0.00	
Weathering removal constant of all vegetation	22	0.00	22	0.00	11	-0.08	11	-0.02	
Wet foliar interception fraction of leafy vegetables	15	-0.08	15	-0.04	18	0.02	18	0.01	
R-SQUARE		0.74		0.74		0.91		0.91	

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak Pb-210 Dose		PCC		SRC		PRCC		SRRC	
Coefficient =		2		2		2		2	
Repetition =									
Description of Probabilistic Variable	Sig	Coeff	Sig	Coeff	Sig	Coeff	Sig	Coeff	Sig
Kd of Pb-210 in Contaminated Zone	10	-0.10	9	-0.06	6	0.17	6	0.05	
Kd of Pb-210 in Unsaturated Zone 1	13	0.08	13	0.05	7	-0.16	7	-0.05	
Kd of Pb-210 in Saturated Zone	16	-0.03	16	-0.02	4	-0.26	4	-0.07	
Kd of Ra-226 in Contaminated Zone	22	-0.01	22	-0.01	9	-0.15	9	-0.04	
Kd of Ra-226 in Unsaturated Zone 1	14	-0.04	15	-0.02	14	0.09	14	0.02	
Kd of Ra-226 in Saturated Zone	21	0.02	18	0.01	19	0.05	19	0.01	
Plant transfer factor for Pb	1	0.74	1	0.63	1	0.94	1	0.76	
Meat transfer factor for Pb	11	0.09	11	0.05	22	0.03	22	0.01	
Milk transfer factor for Pb	12	0.09	12	0.05	5	0.21	5	0.06	
Fish transfer factor for Pb	20	0.02	21	0.01	8	-0.15	8	-0.04	
Plant transfer factor for Ra	18	0.02	19	0.01	15	0.09	15	0.02	
Meat transfer factor for Ra	4	-0.22	4	-0.12	20	0.05	20	0.01	
Milk transfer factor for Ra	3	0.22	3	0.13	21	-0.03	21	-0.01	
Fish transfer factor for Ra	15	-0.04	14	-0.02	17	0.07	17	0.02	
Well pumping rate	8	0.13	8	0.07	12	-0.09	12	-0.03	
Mass loading for inhalation	19	-0.02	20	-0.01	18	-0.06	18	-0.02	
Indoor dust filtration factor	9	0.11	10	0.06	11	0.10	11	0.03	
Depth of soil mixing layer	7	-0.16	7	-0.09	3	-0.68	3	-0.25	
Depth of roots	2	-0.68	2	-0.52	2	-0.88	2	-0.50	
Wet weight crop yield of fruit, grain and non-leafy vegetables	5	0.18	6	0.10	13	0.09	13	0.03	
Weathering removal constant of all vegetation	6	-0.18	5	-0.10	16	-0.08	16	-0.02	
Wet foliar interception fraction of leafy vegetables	17	0.03	17	0.02	10	0.11	10	0.03	
R-SQUARE		0.72		0.72		0.93		0.93	

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak Pb-210 Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC 3		SRC 3		PRCC 3		SRRC 3	
	Sig	Coeff	Sig	Coeff	Sig	Coeff	Sig	Coeff
Kd of Pb-210 in Contaminated Zone	20	-0.02	20	-0.02	14	0.08	14	0.02
Kd of Pb-210 in Unsaturated Zone 1	10	-0.09	10	-0.05	22	0.01	22	0.00
Kd of Pb-210 in Saturated Zone	14	-0.06	13	-0.04	11	0.11	11	0.03
Kd of Ra-226 in Contaminated Zone	15	0.04	15	0.03	20	0.01	20	0.00
Kd of Ra-226 in Unsaturated Zone 1	9	-0.10	9	-0.06	16	0.06	16	0.02
Kd of Ra-226 in Saturated Zone	22	0.00	22	0.00	7	0.16	7	0.05
Plant transfer factor for Pb	1	0.72	1	0.63	1	0.94	1	0.78
Meat transfer factor for Pb	18	-0.03	19	-0.02	8	0.15	8	0.04
Milk transfer factor for Pb	19	-0.03	18	-0.02	4	0.30	4	0.09
Fish transfer factor for Pb	13	-0.06	14	-0.04	6	0.20	6	0.06
Plant transfer factor for Ra	21	-0.01	21	0.00	17	-0.06	17	-0.02
Meat transfer factor for Ra	3	-0.21	3	-0.13	9	0.13	9	0.04
Milk transfer factor for Ra	5	-0.19	5	-0.12	18	-0.04	18	-0.01
Fish transfer factor for Ra	12	-0.06	12	-0.04	19	0.02	19	0.01
Well pumping rate	16	-0.04	17	-0.02	5	-0.26	5	-0.07
Mass loading for inhalation	17	0.04	16	0.03	21	0.01	21	0.00
Indoor dust filtration factor	8	-0.14	8	-0.08	13	0.11	13	0.03
Depth of soil mixing layer	7	-0.14	7	-0.09	3	-0.53	3	-0.18
Depth of roots	2	-0.64	2	-0.50	2	-0.87	2	-0.50
Wet weight crop yield of fruit, grain and non-leafy vegetables	11	0.08	11	0.05	10	0.12	10	0.03
Weathering removal constant of all vegetation	4	-0.19	4	-0.12	15	-0.07	15	-0.02
Wet foliar interception fraction of leafy vegetables	6	0.18	6	0.11	12	0.11	12	0.03
R-SQUARE		0.65		0.65		0.92		0.92

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak Ra-226 Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC		SRC		PRCC		SRRC	
	Sig	Coef	Sig	Coef	Sig	Coef	Sig	Coef
Kd of Pb-210 in Contaminated Zone	11	-0.09	10	-0.05	14	-0.07	14	-0.02
Kd of Pb-210 in Unsaturated Zone 1	15	-0.05	15	-0.03	10	-0.09	10	-0.02
Kd of Pb-210 in Saturated Zone	19	0.02	19	0.01	6	-0.19	6	-0.05
Kd of Ra-226 in Contaminated Zone	10	-0.09	8	-0.05	16	0.07	16	0.02
Kd of Ra-226 in Unsaturated Zone 1	6	-0.10	6	-0.06	17	0.07	17	0.02
Kd of Ra-226 in Saturated Zone	7	-0.09	7	-0.05	19	-0.04	19	-0.01
Plant transfer factor for Pb	3	0.31	3	0.17	3	0.41	3	0.13
Meat transfer factor for Pb	14	0.05	13	0.03	11	0.09	11	0.02
Milk transfer factor for Pb	13	-0.06	14	-0.03	5	-0.22	5	-0.06
Fish transfer factor for Pb	4	-0.22	4	-0.12	9	-0.10	9	-0.03
Plant transfer factor for Ra	1	0.81	1	0.67	1	0.93	1	0.71
Meat transfer factor for Ra	17	0.04	17	0.02	12	-0.08	12	-0.02
Milk transfer factor for Ra	8	0.09	9	0.05	15	0.07	15	0.02
Fish transfer factor for Ra	9	-0.09	11	-0.04	20	-0.03	20	-0.01
Well pumping rate	20	0.01	20	0.01	22	-0.01	22	0.00
Mass loading for inhalation	5	-0.15	5	-0.07	18	-0.04	18	-0.01
Indoor dust filtration factor	22	0.00	22	0.00	8	-0.12	8	-0.03
Depth of soil mixing layer	12	-0.06	12	-0.03	4	-0.30	4	-0.09
Depth of roots	2	-0.72	2	-0.51	2	-0.91	2	-0.61
Wet weight crop yield of fruit, grain and non-leafy vegetables	21	0.00	21	0.00	21	0.01	21	0.00
Weathering removal constant of all vegetation	16	-0.05	16	-0.02	13	-0.08	13	-0.02
Wet foliar interception fraction of leafy vegetables	18	0.03	18	0.01	7	-0.13	7	-0.04
R-SQUARE		0.77		0.77		0.92		0.92

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.

Coefficients for peak Ra-226 Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC		SRC		PRCC		SRRC	
	Sig	Coeff	Sig	Coeff	Sig	Coeff	Sig	Coeff
Kd of Pb-210 in Contaminated Zone	14	0.07	12	0.05	16	-0.04	16	-0.01
Kd of Pb-210 in Unsaturated Zone 1	18	0.03	18	0.02	12	-0.11	12	-0.03
Kd of Pb-210 in Saturated Zone	22	0.01	20	0.01	21	-0.01	21	0.00
Kd of Ra-226 in Contaminated Zone	21	0.01	21	0.01	10	-0.14	10	-0.04
Kd of Ra-226 in Unsaturated Zone 1	20	0.01	22	0.01	9	0.16	9	0.05
Kd of Ra-226 in Saturated Zone	7	-0.19	7	-0.15	7	0.18	7	0.06
Plant transfer factor for Pb	6	0.25	6	0.15	3	0.44	3	0.15
Meat transfer factor for Pb	3	0.31	4	0.19	11	0.12	11	0.04
Milk transfer factor for Pb	10	0.11	10	0.06	17	-0.02	17	-0.01
Fish transfer factor for Pb	19	0.02	19	0.01	5	-0.22	5	-0.07
Plant transfer factor for Ra	1	0.69	1	0.55	1	0.92	1	0.72
Meat transfer factor for Ra	11	-0.09	11	-0.06	20	-0.02	20	-0.01
Milk transfer factor for Ra	5	0.25	5	0.15	13	0.09	13	0.03
Fish transfer factor for Ra	4	0.27	3	0.20	14	-0.08	14	-0.03
Well pumping rate	13	-0.08	14	-0.04	19	-0.02	19	-0.01
Mass loading for inhalation	17	-0.03	17	-0.02	4	-0.27	4	-0.09
Indoor dust filtration factor	16	0.04	16	0.02	15	-0.06	15	-0.02
Depth of soil mixing layer	9	-0.13	9	-0.08	8	-0.17	8	-0.05
Depth of roots	2	-0.67	2	-0.52	2	-0.89	2	-0.60
Wet weight crop yield of fruit, grain and non-leafy vegetables	12	0.08	13	0.05	18	0.02	18	0.01
Weathering removal constant of all vegetation	8	-0.19	8	-0.11	6	-0.19	6	-0.06
Wet foliar interception fraction of leafy vegetables	15	0.05	15	0.03	22	0.01	22	0.00
R-SQUARE		0.71		0.71		0.91		0.91

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.



Coefficients for peak Ra-226 Dose  
 Coefficient =  
 Repetition =

Description of Probabilistic Variable	PCC 3		SRC 3		PRCC 3		SRRC 3	
	Sig	Coeff	Sig	Coeff	Sig	Coeff	Sig	Coeff
Kd of Pb-210 in Contaminated Zone	20	-0.01	20	-0.01	4	0.22	4	0.08
Kd of Pb-210 in Unsaturated Zone 1	16	-0.04	16	-0.03	10	-0.09	10	-0.03
Kd of Pb-210 in Saturated Zone	13	-0.08	13	-0.05	21	0.02	21	0.01
Kd of Ra-226 in Contaminated Zone	5	0.13	5	0.09	17	-0.03	17	-0.01
Kd of Ra-226 in Unsaturated Zone 1	9	-0.11	9	-0.07	5	0.18	5	0.07
Kd of Ra-226 in Saturated Zone	15	0.05	15	0.03	20	0.02	20	0.01
Plant transfer factor for Pb	3	0.30	3	0.20	3	0.44	3	0.17
Meat transfer factor for Pb	22	-0.01	22	-0.01	12	-0.07	12	-0.02
Milk transfer factor for Pb	21	0.01	21	0.01	8	-0.12	8	-0.04
Fish transfer factor for Pb	17	-0.04	17	-0.02	13	0.06	13	0.02
Plant transfer factor for Ra	1	0.69	1	0.60	1	0.88	1	0.67
Meat transfer factor for Ra	10	-0.10	10	-0.06	18	0.03	18	0.01
Milk transfer factor for Ra	6	-0.13	6	-0.08	15	-0.04	15	-0.01
Fish transfer factor for Ra	14	-0.07	14	-0.05	6	-0.16	6	-0.06
Well pumping rate	12	-0.09	12	-0.06	9	-0.12	9	-0.04
Mass loading for inhalation	19	-0.02	19	-0.02	11	0.08	11	0.03
Indoor dust filtration factor	7	-0.12	7	-0.08	19	-0.03	19	-0.01
Depth of soil mixing layer	11	-0.09	11	-0.06	7	-0.13	7	-0.05
Depth of roots	2	-0.59	2	-0.45	2	-0.86	2	-0.61
Wet weight crop yield of fruit, grain and non-leafy vegetables	18	0.02	18	0.02	22	-0.01	22	0.00
Weathering removal constant of all vegetation	8	-0.12	8	-0.08	16	-0.04	16	-0.01
Wet foliar interception fraction of leafy vegetables	4	0.17	4	0.11	14	0.05	14	0.02
R-SQUARE		0.63		0.63		0.87		0.87

-Rank is set to zero if the dose is zero or the correlation matrix is singular.  
 -R-SQUARE varies between 0 and 1 and is called the coefficient of determination; it provides a measure of the variation in the dependent variable (Dose) explained by regression on the independent variables.