

EFFLUENT AND WASTE DISPOSAL

SEMI-ANNUAL REPORT

1st & 2nd QUARTER, 1980

FACILITY: Indian Point Station

LICENSEE: Consolidated Edison Company of New York, Inc.

This information is provided in accordance with the requirements of Regulatory Guide 1.21. The numbered sections of this report reference corresponding sections of the subject Regulatory Guide, pages 1.21-10 to 12. The Power Authority of the State of New York has chosen to issue a separate semi-annual report. The Semi-Annual Effluent and Waste Disposal Report for Indian Point Units 1 and 2 covers discharges for the first and second quarters of 1980. The liquid discharges have been proportioned to Units 1 and 2 based on the total volume of site waste processed at Con Edison. The air-borne releases are for Units 1 and 2 only.

A. Supplemental Information

1. Regulatory Limits

Indian Point Units 1, 2, and 3 are presently subject to specifications on radioactive waste releases that are set forth in Sections 2.4 and 3.4 of Appendix B to Docket's 50-3, 50-247, or 50-286 entitled "Environmental Technical Specification Requirements for Once-Through Cooling" (ETSR). The "percent of limit" reported in Table 2A of this document, is the percent of the applicable quarterly limit specified in the ETSR.

2. Maximum Permissible Concentrations

a. Fission and Activation Gases

The quarterly limits for those specifications stated in the ETSR have been used to calculate the percent of technical specification limit. The isotopic concentration values as reported for elevated releases (Table 1B) and the isotopic concentration values as reported for ground releases (Table 1C) are used in conjunction with K,L,M,N, values found in Table 2.4-5 of the ETSR to determine the $\overline{K}, \overline{L}, \overline{M}, \overline{N}$ values for Unit 1 stack releases and Unit 2 vent release points respectively. The K,L,M,N values represent the gamma-beta dose factors.

b&c. Iodines and Particulates

The applicable quarterly limits for Iodine-131 and particulates with half-lives greater than 8 days in Section 2.4.2.b.3 of the ETSR have been used as the

8009030386

maximum permissible concentrations for the purpose of calculating the percent of technical specification limit in Table 1A of this document.

d. Liquid Effluents

All liquid discharges from Indian Point are made through a common discharge canal with a minimum of 100,000 gpm dilution water. The isotopic content, excluding tritium and dissolved noble gas, of continuous and batch mode discharges for each calendar quarter have been added, and a weighted average fraction of MPC has been calculated for this isotopic mixture as described in 10CFR20. The percent of applicable limit reported in Table 2A of this document is the percent of MPC concentration of the time averaged diluted concentration for each calendar quarter.

The first and second quarter continuous releases are for Units 1 and 2 only. The batch releases discharged through the common site processing facility have been apportioned according to the volume transferred from the respective units to the processing facility.

The tritium limit has been established in the same manner as the other isotopes in liquid effluents.

Since there is no limit stated for dissolved noble gases in 10CFR20, we have adapted the conservative MPC of 2×10^{-4} uci/ml as suggested by K.F. Eckerman.

3. Average Energy

The average energy ($\bar{E}$) of the radionuclide mixture in releases of fission and activation gases for the first quarter was $\bar{E} \gamma$ 4.29 E-2 Mev/Dis., and $\bar{E} \beta$ of 1.64 E-1 Mev/Dis. The corresponding values for the second quarter were $\bar{E} \gamma$ 4.66 E-2 Mev/Dis., and $\bar{E} \beta$ of 1.52 E-1 Mev/Dis.

4. Measurements and Approximations of Total Radioactivity

a. Fission and Activation Gases

Analysis of effluent gases has been performed in compliance with the requirements of Table 2.4-2 of the ETSR. In the case of isolated tanks (batch releases) the total activity discharged is based on an isotopic analysis of each batch and the volume of gas in that batch corrected to standard temperature and pressure.

Vapor containment ventilation discharges have been generally treated as batch releases. At least one complete isotopic concentration analysis of containment air is performed per month and this is applied to a gross analysis of the ventilation air performed prior to each discharge. This information is combined with the volume of air in each discharge to calculate the radionuclide composition of these discharges.

The continuous discharges are based on the isotopic content determined from weekly samples of ventilation air. This information is combined with total air volume discharged by this route. The accumulation of batch and containment ventilation releases is then used to determine total discharges.

b&c. Iodines and Particulates

Iodine-131 and particulate releases are quantified by collecting a continuous sample of ventilation air on a potassium-iodide impregnated activated charcoal cartridge and a glass-fiber filter paper. These samples are changed weekly as required in Table 2.4-2 of the ETSR, and the concentration of isotopes found by analysis of these samples is combined with the volume of air discharged during the sampling period to calculate the amount of activity discharged.

For other iodine isotopes the ratio of each isotope to Iodine-131 is determined for a monthly 24 hour sample. These ratios are then used, along with the total monthly discharge of Iodine-131, to calculate the amount of these isotopes discharged in this monthly period.

d. Liquid Effluents

A proportional composite sample of each batch discharge is taken and an isotopic analysis is performed in compliance with requirements specified in Table 2.4-1 of the ETSR. This isotopic concentration data is combined with information of volume discharged to determine the amount of each isotope discharged in the period.

Samples of continuous discharges have been taken and analyzed in compliance with Table 2.4-1 of the ETSR. This concentration data is combined with the volume discharged to calculate the total activity discharged.

5. Batch Releases

1980

1st Qtr. 2nd Qtr.

a. Liquid

Number of Batch Releases	55	53
Total Time Period Batch Release (Min.)	9495	5923
Maximum Time Period Batch Release (Min.)	743	360
Average Time Period Batch Releases (Min.)	173	112
Minimum Time Period Batch Releases (Min.)	31	33
Average Stream Flow (cfs)	18725	20885

b. Gaseous

Number of Batch Releases	127	106
Total Time Period Batch Releases (Min.)	51601	19058
Maximum Time Period Batch Releases (Min.)	1440	1026
Average Time Period Batch Releases (Min.)	406	180
Minimum Time Period Batch Releases (Min.)	5	70

6. Abnormal Releases

a. Liquid - None

b. Gaseous - None

Effluent and Waste Disposal

Semi-Annual Report

B - Gaseous Effluents

First Half - 1980

TABLE 1A

EFFLUENT AND WASTE DISPOSAL SEMIANNUAL REPORT (1980)

GASEOUS EFFLUENTS—SUMMATION OF ALL RELEASES

	Unit	Quarter 1st	Quarter 2nd	Est. Total Error, %
--	------	----------------	----------------	------------------------

A. Fission & activation gases

1. Total release	Ci	8.02 E+2	8.51 E+2	±5.0 E+1
2. Average release rate for period	μCi/sec	1.02 E+2	1.08 E+2	
3. Percent of Technical specification limit	%	2.46 E0	2.41 E0	

B. Iodines

1. Total iodine-131	Ci	1.90 E-2	2.71 E-2	±5.0 E+1
2. Average release rate for period	μCi/sec	2.42 E-3	3.45 E-3	
3. Percent of technical specification limit	%	6.06 E-1	5.66 E-1	

C. Particulates

1. Particulates with half-lives >8 days	Ci	6.60 E-4	8.99 E-4	±5.0 E+1
2. Average release rate for period	μCi/sec	8.39 E-5	1.14 E-4	
3. Percent of technical specification limit	%	6.06 E-1	5.66 E-1	
4. Gross alpha radioactivity	Ci	2.60 E-7	5.97 E-7	

D. Tritium

1. Total release	Ci	6.78 E 0	1.91 E 0	±5.0 E+1
2. Average release rate for period	μCi/sec	8.62 E-1	2.43 E-1	
3. Percent of technical specification limit	%	 E 	 E 	

EFFLUENT AND WASTE DISPOSAL SEMI-ANNUAL REPORT (1980)

GASEOUS EFFLUENTS—ELEVATED RELEASE

CONTINUOUS MODE

BATCH MODE

Nuclides Released	Unit	1stQuarter	2ndQuarter	Quarter	Quarter
-------------------	------	------------	------------	---------	---------

1. Fission gases

krypton-85	Ci	2.14 E+1	1.13 E-1	. E	. E
krypton-85m	Ci	4.53 E-1	1.05 E-1	. E	. E
krypton-87	Ci	. E	. E	. E	. E
krypton-88	Ci	2.48 E-1	4.04 E-1	. E	. E
xenon-133	Ci	1.94 E+1	2.28 E+1	. E	. E
xenon-135	Ci	7.04 E-2	3.30 E-1	. E	. E
xenon-135m	Ci	. E	3.29 E-1	. E	. E
xenon-138	Ci	. E	. E	. E	. E
Others (specify) xenon-131m	Ci	. E	3.66 E0	. E	. E
xenon-133m	Ci	2.85 E-1	5.29 E-1	. E	. E
	Ci	. E	. E	. E	. E
unidentified	Ci	. E	. E	. E	. E
Total for period	Ci	4.18 E+1	2.83 E+1	. E	. E

2. Iodines

iodine-131	Ci	1.67 E-2	2.20 E-2	. E	. E
iodine-133	Ci	7.22 E-5	4.31 E-4	. E	. E
iodine-135	Ci	8.66 E-4	2.41 E-3	. E	. E
Total for period	Ci	1.76 E-2	2.48 E-2	. E	. E

3. Particulates

strontium-89	Ci	<2.53 E-6	<5.44 E-6	. E	. E
strontium-90	Ci	<1.19 E-6	<1.29 E-6	. E	. E
cesium-134	Ci	3.00 E-5	2.58 E-5	. E	. E
cesium-137	Ci	1.02 E-4	8.96 E-5	. E	. E
barium-lanthanum-140	Ci	<1.24 E-5	<2.83 E-5	. E	. E
Others (specify) I-131	Ci	5.54 E-6	1.27 E-5	. E	. E
Co-58	Ci	1.73 E-5	5.29 E-5	. E	. E
Co-60	Ci	6.03 E-5	1.07 E-4	. E	. E
Mn-54	Ci	4.72 E-6	9.07 E-7	. E	. E
Cr-51	Ci	-	7.68 E-6		
Fe-55	Ci	2.12 E-5	1.71 E-5		
P-32	Ci	1.85 E-6	3.87 E-6		
Ni-63	Ci	2.73 E-5	3.43 E-4		

TABLE 1C

EFFLUENT AND WASTE DISPOSAL SEMI-ANNUAL REPORT (1980)

GASEOUS EFFLUENTS—GROUND-LEVEL RELEASES

Nuclides Released	Unit	CONTINUOUS MODE		BATCH MODE	
		1st Quarter	2nd Quarter	1st Quarter	2nd Quarter

1. Fission gases

krypton-85	Ci	8.38 E+1	1.45 E 0	1.32 E+1	4.44 E 0
krypton-85m	Ci	1.73 E 0	1.67 E 0	8.25 E-1	8.74 E-1
krypton-87	Ci	. E	7.43 E -2	1.53 E-1	2.26 E-1
krypton-88	Ci	3.33 E 0	3.13 E 0	1.01 E 0	1.07 E 0
xenon-133	Ci	1.30 E+2	2.90 E+2	4.90 E+2	4.75 E+2
xenon-135	Ci	1.43 E+1	1.89 E+1	1.02 E+1	8.73 E 0
xenon-135m	Ci	1.39 E-1	. E	8.30 E-1	1.06 E 0
xenon-138	Ci	. E	. E	2.25 E-1	3.85 E-3
Others (specify) xenon-131m	Ci	2.08 E-1	3.86 E 0	4.40 E 0	6.18 E 0
xenon-133m	Ci	6.52 E-1	7.80 E-1	5.04 E 0	4.81 E 0
xenon-137	Ci	. E	. E	. E	2.77 E-3
unidentified	Ci	. E	. E	. E	. E
Total for period	Ci	2.34 E+2	3.20 E+2	5.26 E+2	5.03 E+2

2. Iodines

iodine-131	Ci	2.27 E-3	5.10 E-3	. E	. E
iodine-133	Ci	1.72 E-3	1.62 E-3	. E	. E
iodine-135	Ci	4.27 E-3	3.60 E-3	. E	. E
Total for period	Ci	8.26 E-3	1.03 E-2	. E	. E

3. Particulates

strontium-89	Ci	< 3.67 E-6	< 3.99 E-6	. E	. E
strontium-90	Ci	< 8.88 E-7	< 7.54 E-7	. E	. E
cesium-134	Ci	4.87 E-5	1.43 E-5	. E	. E
cesium-137	Ci	7.90 E-5	4.25 E-5	. E	. E
barium-lanthanum-140	Ci	< 1.94 E-5	< 2.08 E-5	. E	. E
Others (specify) I-131	Ci	1.27 E-5	7.86 E-6	. E	. E
Co-58	Ci	2.86 E-5	2.97 E-5	. E	. E
Co-60	Ci	9.02 E-5	5.13 E-5	. E	. E
Mn-54	Ci	2.72 E-6	1.27 E-6	. E	. E
Cr-51	Ci	9.38 E-6			
Fe-55	Ci	5.20 E-5	1.04 E-5		
P-32	Ci	2.95 E-6	2.44 E-6		
Ni-63	Ci	5.45 E-6	1.63 E-5		

Effluent and Waste Disposal

Semi-Annual Report

C - Liquid Effluent

First Half - 1980

TABLE 2A

EFFLUENT AND WASTE DISPOSAL SEMIANNUAL REPORT (1980)

LIQUID EFFLUENTS—SUMMATION OF ALL RELEASES

Unit	Quarter 1	Quarter 2	Est. Total Error, %
------	--------------	--------------	------------------------

A. Fission and activation products

1. Total release (not including tritium, gases, alpha)	Ci	4.85 E-1	1.80 E-1	± 2.5 E+1
2. Average diluted concentration during period	μCi/ml	3.38 E-9	5.84 E-10	
3. Percent of applicable limit	%	1.71 E-1	1.80 E-2	

B. Tritium

1. Total release	Ci	2.75 E+1	1.11 E+2	± 2.5 E+1
2. Average diluted concentration during period	μCi/ml	1.91 E-7	3.62 E-7	
3. Percent of applicable limit	%	6.39 E-3	1.20 E-2	

C. Dissolved and entrained gases

1. Total release	Ci	4.03 E-3	3.40 E-3	± 2.5 E+1
2. Average diluted concentration during period	μCi/ml	2.8 E-11	1.10 E-11	
3. Percent of applicable limit	%	1.4 E-5	5.52 E-6	

D. Gross alpha radioactivity

1. Total release	Ci	<1.69 E-3	<1.10 E-3	± 2.5 E+1
------------------	----	-----------	-----------	-----------

E. Volume of waste released (prior to dilution)	liters	2.59 E+7	1.27 E+7	± 1.0 E+1
---	--------	----------	----------	-----------

F. Volume of dilution water used during period	liters	1.43 E+11	3.08 E+11	± 1.0 E+1
--	--------	-----------	-----------	-----------

TABLE 2B

EFFLUENT AND WASTE DISPOSAL SEMIANNUAL REPORT (1980)

LIQUID EFFLUENTS

Nuclides Released	Unit	CONTINUOUS MODE		BATCH MODE	
		1 Quarter	2 Quarter	1 Quarter	2 Quarter
strontium-89	Ci	< 8.42 E-4	< 3.05 E-4	< 7.91 E-5	< 1.85 E-4
strontium-90	Ci	< 3.34 E-4	< 2.66 E-4	< 5.45 E-4	< 5.21 E-4
cesium-134	Ci	7.02 E-3	6.05 E-4	1.22 E-2	1.91 E-2
cesium-137	Ci	3.22 E-2	5.75 E-4	3.30 E-2	5.53 E-2
iodine-131	Ci	5.69 E-2	3.76 E-3	8.87 E-3	1.06 E-2
cobalt-58	Ci	2.49 E-3	2.39 E-3	7.62 E-3	4.63 E-3
cobalt-60	Ci	4.02 E-2	6.91 E-4	1.99 E-2	1.36 E-2
iron-59	Ci	6.04 E-3	1.70 E-3	3.62 E-4	5.17 E-4
zinc-65	Ci	5.84 E-3	1.53 E-3	5.04 E-4	7.67 E-4
manganese-54	Ci	2.04 E-3	9.54 E-4	2.88 E-3	2.40 E-3
chromium-51	Ci	1.89 E-2	4.19 E-3	3.22 E-3	4.57 E-3
zirconium-niobium-95	Ci	< 3.87 E-3	< 1.34 E-3	< 1.71 E-3	< 5.88 E-4
molybdenum-99	Ci	< 1.15 E-2	< 3.81 E-3	< 1.61 E-3	< 1.96 E-3
technetium-99m	Ci	< 2.38 E-2	< 2.76 E-4	< 2.10 E-4	< 2.45 E-4
barium-lanthanum-140	Ci	< 9.12 E-3	< 3.14 E-3	< 1.74 E-3	< 2.17 E-3
cerium-141	Ci	< 1.97 E-3	< 7.38 E-4	< 5.64 E-4	< 6.14 E-4
FE 55	Ci	1.37 E-1	1.21 E-2	5.24 E-3	7.85 E-3
NI 63	Ci	1.16 E-2	6.64 E-3	6.50 E-3	6.28 E-3
P 32	Ci	5.15 E-3	1.92 E-3	1.38 E-3	1.08 E-3
	Ci	. E	. E	. E	. E
	Ci	. E	. E	. E	. E
unidentified	Ci	. E	. E	. E	. E
Total for period (above)	Ci	3.77 E-1	4.69 E-2	1.08 E-1	1.33 E-1
xenon-133	Ci	1.87 E-3	7.85 E-4	7.26 E-4	1.87 E-3
xenon-135	Ci	1.25 E-3	5.06 E-4	1.74 E-4	2.41 E-4

Effluent and Waste Disposal

Semi-Annual Report

D - Solid Waste

First Half - 1980

TABLE 3

EFFLUENT AND WASTE DISPOSAL SEMI-ANNUAL REPORT (1980)

SOLID WASTE AND IRRADIATED FUEL SHIPMENTS

A. SOLID WASTE SHIPPED OFFSITE FOR BURIAL OR DISPOSAL (Not irradiated fuel)

1. Type of waste	Unit	1st & 2nd Qtr 6-month Period	Est. Total Error, %
a. Spent resins, filter sludges, evaporator bottoms, etc.	m ³ Ci	2.88 E+2 1.12 E+2	±1.0 E+2
b. Dry compressible waste, contaminated equip, etc.	m ³ Ci	2.33 E+2 8.7 E+1	±1.0 E+2
c. Irradiated components, control rods, etc.	m ³ Ci	. E . E	. E
d. Other (describe)	m ³ Ci	. E . E	. E

2. Estimate of major nuclide composition (by type of waste)

a.	¹³⁷ Cs	46 %	± 1.0 E+2
	Cs ¹³⁴	22 %	± 1.0 E+2
	Co ⁵⁸	9 %	± 1.0 E+2
	Co ⁶⁰ , I ¹³¹ , Mn ⁵⁴ , Fe ⁵⁵ , Ni ⁶³	23 %	± 1.0 E+2
b.	Same as "a".	%	. E
		%	. E
c.		%	. E
		%	. E
		%	. E
d.		%	. E
		%	. E
		%	. E

3. Solid Waste Disposition

Number of Shipments

42

Mode of Transportation

Truck

Destination

Barnwell, S.C.

B. IRRADIATED FUEL SHIPMENTS (Disposition)

Number of Shipments

None

Mode of TransportationDestination

Effluent and Waste Disposal

Semi-Annual Report

E-- Radiological Impact on Man

First Half - 1980

RADIOLOGICAL IMPACT EVALUATION

Doses from noble gas immersion, inhalation, ground deposition, and vegetation ingestion were evaluated for the nearest residence likely to be occupied in the critical sector for each pathway and were combined to provide a conservative determination of the maximum individual offsite radiation dose from these pathways. Doses were also evaluated for an individual ingesting milk and meat from a cow located about 8.9 miles to the ESE. In all cases, these evaluations were performed using the models presented in Regulatory Guide 1.109. Noble gas releases were considered to be either batch or continuous, while all iodine and particulate release were considered to be continuous. Noble gas release from containment pressure reliefs and purges were treated as a continuous release, because of the large number of such releases. Other releases (e.g. gas decay tank releases) were considered as batch and were evaluated using actual meteorological conditions existing during the release period. Continuous releases were evaluated using average meteorological conditions based on the six month release period. Estimates of relative deposition per unit area were obtained from the numerical approximation presented in the NRC computer program XOQDOQ for ground releases. Values of atmospheric dispersion factors (X/Q) were computed using the Sagendorf straight line air flow model, assuming a ground level release, and the subroutine POLYN (from XOQDOQ) for computing standard deviations in the horizontal direction (σ_y) and vertical direction (σ_z).

Integrated doses for the population within 50 miles of Indian Point from gaseous effluents were computed based on linear interpolation of 1970 - 2010 population data contained in the Indian Point Unit No. 3 FSAR.

NUREG-0017, "Calculation of Release of Radioactive Materials in Gaseous and Liquid effluents from Pressurized Water Reactors", assumes an annual release of 8.0Ci/yr of Carbon-14 per reactor. Therefore, to be consistent with NUREG-0017, a release of 4.0 Curies of Carbon-14 for Unit No. 2 was assumed for the six month period in addition to the radioactive materials measured in Indian Point gaseous effluents.

Indian Point Units 1 and 2

RADIOLOGICAL IMPACT ON MAN

(Reference regulatory guide 1.21, page 12)

A, Maximum Individual Doses

<u>Pathways</u> (Gaseous)	<u>Total Body</u> (mr)	<u>Skin</u> (mr)	<u>Thyroid</u> (mr)	<u>Bone</u> (mr)
Noble Gas Immersion				
a) Batch Releases	.119	.300		
b) Continuous Releases	.0588	.156		
Inhalation*	6.07×10^{-3}	-	.160	2.36×10^{-2}
Ground Deposition	3.99×10^{-3}	4.69×10^{-3}	-	-
Milk Ingestion*	4.24×10^{-3}	-	2.28×10^{-2}	1.97×10^{-2}
Meat Ingestion***	6.46×10^{-4}	-	7.43×10^{-4}	3.22×10^{-3}
Vegetable Ingestion***	2.18×10^{-2}	-	2.77×10^{-1}	1.07×10^{-1}

* Infants are critical age group

** Adults are critical age group

*** Children are critical age group

Pathways
(Liquid)

All

See Attached "LADTAP" printout
Attachment I

B. Population Doses

<u>Pathways</u> (Gaseous)	<u>Total Body</u> (man-rem)	<u>Thyroid</u> (man-thyroid rem)
Noble Gas Immersion		
a) Batch Release	2.59	
b) Continuous Release	2.46	
Inhalation	.226	5.05
Ground Deposition	.098	
Totals	5.374	5.05

Pathways
(Liquid)

All See attached "LADTAP" printout
Attachment I

C. Average Doses to Individuals

1. Liquid-Total Body
 1.09×10^{-4} mr
2. Gaseous-Total Body
 2.83×10^{-4} mr

* * * AS LOW AS REASONABLY ACHIEVABLE * * *

ADULT DOSES

DOSE (MREM PER YEAR INTAKE)

PATHWAY	SKIN	BONE	LIVER	TOTAL BODY	THYROID	KIDNEY	LUNG	GI-LLI
FISH		7.90E-01	1.96E-01	1.34E-01	8.52E-03	5.10E-02	1.88E-02	8.29E-02
INVERTEBRATE		3.92E-02	2.22E-02	7.77E-03	6.87E-04	2.40E-03	1.60E-02	2.14E-02
ALGAE		1.67E-06	1.36E-07	8.41E-08	1.08E-08	9.88E-09	1.19E-08	1.93E-07
DRINKING		6.06E-13	8.07E-13	6.93E-13	2.31E-12	4.78E-13	4.83E-13	4.81E-13
SHORELINE	8.91E-04	7.60E-04	7.60E-04	7.60E-04	7.60E-04	7.60E-04	7.60E-04	7.60E-04
SWIMMING	0.0	6.97E-06	6.97E-06	6.97E-06	6.97E-06	6.97E-06	6.97E-06	6.97E-06
BOATING	0.0	6.97E-06	6.97E-06	6.97E-06	6.97E-06	6.97E-06	6.97E-06	6.97E-06
TOTAL	8.91E-04	8.30E-01	2.19E-01	1.42E-01	9.99E-03	5.42E-02	3.56E-02	1.05E-01

	USAGE (KG/YR,HR/YR)	DILUTION	TIME(HR)	SHOREWIDTH FACTOR=0.2
FISH	21.0	5.0	25.00	
INVERTEBRATE	5.0	5.0	25.00	
ALGAE	0.0	5.0	25.00	
DRINKING	0.0	500.0	112.00	
SHORELINE	50.0	5.0	1.00	
SWIMMING	50.0	5.0	1.00	
BOATING	100.0	5.0	1.00	

TEENAGER DOSES

DOSE (MREM PER YEAR INTAKE)

PATHWAY	SKIN	BONE	LIVER	TOTAL BODY	THYROID	KIDNEY	LUNG	GI-LLI
FISH		6.26E-01	1.86E-01	7.96E-02	7.54E-03	3.89E-02	2.10E-02	6.29E-02
INVERTEBRATE		3.05E-02	1.73E-02	5.74E-03	6.03E-04	1.83E-03	1.23E-02	1.61E-02
ALGAE		1.68E-06	1.41E-07	8.17E-08	1.25E-08	9.88E-09	1.29E-08	1.97E-07
DRINKING		7.76E-13	8.51E-13	5.77E-13	2.56E-12	4.78E-13	4.32E-13	3.92E-13
SHORELINE	1.19E-03	1.02E-03	1.02E-03	1.02E-03	1.02E-03	1.02E-03	1.02E-03	1.02E-03
SWIMMING	0.0	1.39E-05	1.39E-05	1.39E-05	1.39E-05	1.39E-05	1.39E-05	1.39E-05
BOATING	0.0	6.97E-06	6.97E-06	6.97E-06	6.97E-06	6.97E-06	6.97E-06	6.97E-06
TOTAL	1.19E-03	6.58E-01	2.04E-01	8.64E-02	9.18E-03	4.17E-02	3.43E-02	8.00E-02

	USAGE (KG/YR,HR/YR)	DILUTION	TIME(HR)	SHOREWIDTH FACTOR=0.2
FISH	16.0	5.0	25.00	
INVERTEBRATE	3.8	5.0	25.00	
ALGAE	0.0	5.0	25.00	
DRINKING	0.0	500.0	112.00	
SHORELINE	67.0	5.0	1.00	
SWIMMING	100.0	5.0	1.00	
BOATING	100.0	5.0	1.00	

CHILD DOSES

DOSE (MREM PER YEAR INTAKE)

PATHWAY	SKIN	BONE	LIVER	TOTAL BODY	THYROID	KIDNEY	LUNG	GI-LLI
FISH		3.49E-01	1.48E-01	3.18E-02	7.77E-03	1.68E-02	1.61E-02	2.71E-02
INVERTEBRATE		1.51E-02	8.60E-03	2.69E-03	6.44E-04	8.17E-04	5.57E-03	7.18E-03
ALGAE		1.72E-03	1.66E-04	8.40E-05	3.00E-05	9.88E-06	1.55E-05	1.97E-04
DRINKING		1.61E-09	1.61E-09	8.86E-10	5.99E-09	4.78E-10	7.32E-10	6.35E-10
SHORELINE	2.49E-04	2.13E-04	2.13E-04	2.13E-04	2.13E-04	2.13E-04	2.13E-04	2.13E-04
SWIMMING	0.0	3.49E-06	3.49E-06	3.49E-06	3.49E-06	3.49E-06	3.49E-06	3.49E-06
BOATING	0.0	6.97E-06	6.97E-06	6.97E-06	6.97E-06	6.97E-06	6.97E-06	6.97E-06
TOTAL	2.49E-04	3.66E-01	1.57E-01	3.47E-02	8.67E-03	1.78E-02	2.19E-02	3.47E-02

	USAGE (KG/YR,HR/YR)	DILUTION	TIME(HR)	SHOREWIDTH FACTOR=0.2
FISH	0.9	5.0	25.00	
INVERTEBRATE	1.7	5.0	25.00	
ALGAE	0.0	5.0	25.00	
DRINKING	0.0	500.0	112.00	
SHORELINE	14.0	5.0	1.00	
SHIMMING	25.0	5.0	1.00	
BOATING	100.0	5.0	1.00	

INFANT DOSES

DOSE (MREM PER YEAR INTAKE)

PATHWAY	SKIN	BONE	LIVER	TOTAL BODY	THYROID	KIDNEY	LUNG	GI-LLI
FISH		3.48E-02	2.38E-02	2.17E-03	1.36E-03	1.21E-03	2.75E-03	1.96E-03
INVERTEBRATE		1.01E-03	6.38E-04	1.67E-04	9.09E-05	4.80E-05	3.44E-04	4.22E-04
ALGAE		1.79E-06	2.31E-07	8.70E-08	7.22E-08	9.88E-09	2.34E-08	1.97E-07
DRINKING		2.88E-12	3.21E-12	1.23E-12	1.40E-11	4.78E-13	1.16E-12	9.00E-13
SHORELINE	5.34E-05	4.56E-05	4.56E-05	4.56E-05	4.56E-05	4.56E-05	4.56E-05	4.56E-05
SHIMMING	0.0	1.39E-12	1.39E-12	1.39E-12	1.39E-12	1.39E-12	1.39E-12	1.39E-12
BOATING	0.0	3.49E-08	3.49E-08	3.49E-08	3.49E-08	3.49E-08	3.49E-08	3.49E-08
TOTAL	5.34E-05	3.59E-02	2.45E-02	2.38E-03	1.49E-03	1.31E-03	3.14E-03	2.43E-03

	USAGE (KG/YR,HR/YR)	DILUTION	TIME(HR)	SHOREWIDTH FACTOR=0.2
FISH	0.5	5.0	25.00	
INVERTEBRATE	0.1	5.0	25.00	
ALGAE	0.0	5.0	25.00	
DRINKING	0.0	500.0	112.00	
SHORELINE	3.0	5.0	1.00	
SHIMMING	0.0	5.0	1.00	
BOATING	0.5	5.0	1.00	

* * * SELECTED LOCATION * * *

LOCATION IS DOWNSTREAM

ADULT DOSES

DOSE (MREM PER YEAR INTAKE)

PATHWAY	SKIN	BONE	LIVER	TOTAL BODY	THYROID	KIDNEY	LUNG	GI-LLI
FISH		5.58E-01	1.40E-01	9.53E-02	5.96E-03	3.64E-02	1.34E-02	5.85E-02
INVERTEBRATE		2.77E-02	1.58E-02	5.54E-03	4.80E-04	1.71E-03	1.14E-02	1.52E-02
ALGAE		1.18E-06	9.59E-08	5.95E-08	7.53E-09	7.04E-09	8.49E-09	1.40E-07
DRINKING		4.40E-11	5.80E-11	4.97E-11	2.21E-10	3.46E-11	3.45E-11	3.48E-11
SHORELINE	6.36E-04	5.43E-04	5.43E-04	5.43E-04	5.43E-04	5.43E-04	5.43E-04	5.43E-04
SWIMMING	0.0	4.95E-06	4.95E-06	4.95E-06	4.95E-06	4.95E-06	4.95E-06	4.95E-06
BOATING	0.0	4.95E-06	4.95E-06	4.95E-06	4.95E-06	4.95E-06	4.95E-06	4.95E-06
TOTAL	6.36E-04	5.87E-01	1.56E-01	1.01E-01	6.99E-03	3.87E-02	2.54E-02	7.43E-02

	USAGE (KG/YR,HR/YR)	DILUTION	TIME(HR)	SHOREWIDTH FACTOR=0.2
FISH	21.0	7.0	31.00	
INVERTEBRATE	5.0	7.0	31.00	
ALGAE	0.0	7.0	31.00	
DRINKING	0.0	7.0	19.00	
SHORELINE	50.0	7.0	7.00	
SWIMMING	50.0	7.0	7.00	
BOATING	100.0	7.0	7.00	

LOCATION IS DOWNSTREAM

TEENAGER DOSES

DOSE (MREM PER YEAR INTAKE)

PATHWAY	SKIN	BONE	LIVER	TOTAL BODY	THYROID	KIDNEY	LUNG	GI-LLI
FISH		4.43E-01	1.32E-01	5.67E-02	5.27E-03	2.78E-02	1.50E-02	4.44E-02
INVERTEBRATE		2.15E-02	1.24E-02	4.09E-03	4.22E-04	1.30E-03	8.75E-03	1.15E-02
ALGAE		1.19E-06	1.00E-07	5.78E-08	8.76E-09	7.04E-09	9.24E-09	1.39E-07
DRINKING		5.62E-11	6.12E-11	4.14E-11	2.46E-10	3.46E-11	3.09E-11	2.83E-11
SHORELINE	8.52E-04	7.27E-04	7.27E-04	7.27E-04	7.27E-04	7.27E-04	7.27E-04	7.27E-04
SWIMMING	0.0	9.90E-06	9.90E-06	9.90E-06	9.90E-06	9.90E-06	9.90E-06	9.90E-06
BOATING	0.0	4.95E-06	4.95E-06	4.95E-06	4.95E-06	4.95E-06	4.95E-06	4.95E-06
TOTAL	8.52E-04	4.65E-01	1.45E-01	6.15E-02	6.44E-03	2.98E-02	2.45E-02	5.66E-02

	USAGE (KG/YR,HR/YR)	DILUTION	TIME(HR)	SHOREWIDTH FACTOR=0.2
FISH	16.0	7.0	31.00	
INVERTEBRATE	3.8	7.0	31.00	
ALGAE	0.0	7.0	31.00	
DRINKING	0.0	7.0	19.00	
SHORELINE	67.0	7.0	7.00	
SWIMMING	100.0	7.0	7.00	
BOATING	100.0	7.0	7.00	

LOCATION IS DOWNSTREAM

CHILD DOSES

DOSE (MREM PER YEAR INTAKE)

PATHWAY	SKIN	BONE	LIVER	TOTAL BODY	THYROID	KIDNEY	LUNG	GI-LLI
FISH		2.48E-01	1.06E-01	2.26E-02	5.43E-03	1.20E-02	1.15E-02	1.91E-02
INVERTEBRATE		1.07E-02	6.14E-03	1.91E-03	4.50E-04	5.83E-04	3.98E-03	5.11E-03

ALGAE		1.22E-03	1.18E-04	5.94E-05	2.09E-05	7.04E-06	1.11E-05	1.39E-04
DRINKING		1.16E-07	1.15E-07	6.38E-08	5.83E-07	3.46E-08	5.24E-08	4.57E-08
SHORELINE	1.78E-04	1.52E-04	1.52E-04	1.52E-04	1.52E-04	1.52E-04	1.52E-04	1.52E-04
SWIMMING	0.0	2.47E-06	2.47E-06	2.47E-06	2.47E-06	2.47E-06	2.47E-06	2.47E-06
BOATING	0.0	4.95E-06	4.95E-06	4.95E-06	4.95E-06	4.95E-06	4.95E-06	4.95E-06
TOTAL	1.78E-04	2.60E-01	1.12E-01	2.47E-02	6.06E-03	1.27E-02	1.56E-02	2.45E-02

	USAGE (KG/YR,HR/YR)	DILUTION	TIME(HR)	SHOREWIDTH FACTOR=0.2
FISH	6.9	7.0	31.00	
INVERTEBRATE	1.7	7.0	31.00	
ALGAE	0.0	7.0	31.00	
DRINKING	0.0	7.0	19.00	
SHORELINE	14.0	7.0	7.00	
SWIMMING	25.0	7.0	7.00	
BOATING	100.0	7.0	7.00	

LOCATION IS DOWNSTREAM

INFANT DOSES

DOSE (MREM PER YEAR INTAKE)

PATHWAY	SKIN	BONE	LIVER	TOTAL BODY	THYROID	KIDNEY	LUNG	GI-LLI
FISH		2.47E-02	1.70E-02	1.55E-03	9.48E-04	8.68E-04	1.96E-03	1.39E-03
INVERTEBRATE		7.19E-04	4.56E-04	1.19E-04	6.36E-05	3.43E-05	2.46E-04	3.00E-04
ALGAE		1.26E-06	1.64E-07	6.16E-08	5.05E-08	7.04E-09	1.67E-08	1.39E-07
DRINKING		2.08E-10	2.31E-10	8.89E-11	1.37E-09	3.46E-11	8.28E-11	6.46E-11
SHORELINE	3.82E-05	3.26E-05	3.26E-05	3.26E-05	3.26E-05	3.26E-05	3.26E-05	3.26E-05
SWIMMING	0.0	9.90E-13	9.90E-13	9.90E-13	9.90E-13	9.90E-13	9.90E-13	9.90E-13
BOATING	0.0	2.47E-08	2.47E-08	2.47E-08	2.47E-08	2.47E-08	2.47E-08	2.47E-08
TOTAL	3.82E-05	2.55E-02	1.75E-02	1.70E-03	1.04E-03	9.35E-04	2.24E-03	1.72E-03

	USAGE (KG/YR,HR/YR)	DILUTION	TIME(HR)	SHOREWIDTH FACTOR=0.2
FISH	0.5	7.0	31.00	
INVERTEBRATE	0.1	7.0	31.00	
ALGAE	0.0	7.0	31.00	
DRINKING	0.0	7.0	19.00	
SHORELINE	3.0	7.0	7.00	
SWIMMING	0.0	7.0	7.00	
BOATING	0.5	7.0	7.00	

* * * FISH CONSUMPTION POPULATION DOSE * * *

MAN-REM

SPORTFISH HARVEST

-----DOSE (MAN-REM)-----

PATHWAY	AGE GROUP	USAGE	BONE	LIVER	TOTAL BODY	THYROID	KIDNEY	LUNG	GI-LLI
FISH	ADULT	7.24E+04	1.49E+00	4.48E-01	3.07E-01	1.24E-02	1.23E-01	4.54E-02	1.53E-01
FISH	TEENAGER	1.16E+04	2.51E-01	8.99E-02	3.78E-02	2.30E-03	1.97E-02	1.07E-02	2.44E-02
FISH	CHILD	7.00E+03	2.08E-01	1.03E-01	2.10E-02	3.32E-03	1.19E-02	1.14E-02	1.47E-02
FISH	TOTAL	9.10E+04	1.95E+00	6.40E-01	3.66E-01	1.80E-02	1.55E-01	6.74E-02	1.93E-01

DILUTION CATCH TIME(HR)-INCLUDES FOOD PROCESSING TIME OF 1.68E+02 HR POPULATION=1.59E+04
 7.00E+00 9.10E+04 1.68E+02

AVERAGE INDIVIDUAL CONSUMPTION (KG/YR) ADULT=6.90E+00 TEEN=5.20E+00 CHILD=2.20E+00

* * * FISH CONSUMPTION POPULATION DOSES * * *

MAN-REM

COMMERCIAL HARVEST

-----DOSE (MAN-REM)-----

PATHWAY	AGE GROUP	USAGE	BONE	LIVER	TOTAL BODY	THYROID	KIDNEY	LUNG	GI-LLI
FISH	ADULT	8.65E+07	5.57E+00	1.84E+00	1.26E+00	4.05E-02	5.18E-01	1.91E-01	5.65E-01
FISH	TEENAGER	1.38E+07	9.43E-01	3.71E-01	1.54E-01	7.50E-03	8.28E-02	4.48E-02	8.98E-02
FISH	CHILD	8.36E+06	8.07E-01	4.28E-01	8.56E-02	1.08E-02	5.01E-02	4.80E-02	5.41E-02
FISH	TOTAL	1.09E+08	7.32E+00	2.64E+00	1.50E+00	5.88E-02	6.51E-01	2.84E-01	7.09E-01

DILUTION CATCH TIME(HR)-INCLUDES FOOD PROCESSING TIME OF 2.40E+02 HR POPULATION=1.90E+07
 7.00E+00 1.55E+05 2.40E+02

AVERAGE INDIVIDUAL CONSUMPTION (KG/YR) ADULT=6.90E+00 TEEN=5.20E+00 CHILD=2.20E+00

NEPA DOSES

NOTE--TOATL NEPA DOSE MUST INCLUDE SPORT CATCH, DOSES BELOW ARE FOR COMMERCIAL CATCH ONLY

-----DOSE (MAN-REM)-----

PATHWAY	AGE GROUP	USAGE	BONE	LIVER	TOTAL BODY	THYROID	KIDNEY	LUNG	GI-LLI
FISH	ADULT	1.23E+05	2.25E+00	7.44E-01	5.11E-01	1.64E-02	2.10E-01	7.72E-02	2.29E-01
FISH	TEENAGER	1.97E+04	3.81E-01	1.50E-01	6.25E-02	3.04E-03	3.35E-02	1.81E-02	3.63E-02
FISH	CHILD	1.19E+04	3.27E-01	1.73E-01	3.47E-02	4.38E-03	2.03E-02	1.94E-02	2.19E-02
FISH	TOTAL	1.55E+05	2.96E+00	1.07E+00	6.08E-01	2.38E-02	2.63E-01	1.15E-01	2.87E-01

* * * INVERTEBRATE CONSUMPTION POPULATION DOSES * * *

MAN-REM

SPORTFISH HARVEST

-----DOSE (MAN-REM)-----

PATHWAY	AGE GROUP	USAGE	BONE	LIVER	TOTAL BODY	THYROID	KIDNEY	LUNG	GI-LLI
INVER	ADULT	7.94E+03	3.45E-02	2.40E-02	8.26E-03	4.66E-04	2.65E-03	1.78E-02	2.25E-02
INVER	TEENAGER	1.26E+03	5.64E-03	3.92E-03	1.28E-03	8.52E-05	4.21E-04	2.85E-03	3.54E-03
INVER	CHILD	7.94E+02	4.01E-03	2.75E-03	8.40E-04	1.27E-04	2.65E-04	1.82E-03	2.22E-03
INVER	TOTAL	1.00E+04	4.41E-02	3.07E-02	1.04E-02	6.78E-04	3.33E-03	2.25E-02	2.82E-02

DILUTION CATCH TIME(HR)-INCLUDES FOOD PROCESSING TIME OF 1.68E+02 HR POPULATION=1.20E+04
 7.00E+00 1.00E+04 1.68E+02

AVERAGE INDIVIDUAL CONSUMPTION (KG/YR) ADULT=1.00E+00 TEEN=7.50E-01 CHILD=3.30E-01

* * * INVERTEBRATE CONSUMPTION POPULATION DOSES * * *

MAN-REM

COMMERCIAL HARVEST

-----DOSE (MAN-REM)-----

PATHWAY	AGE GROUP	USAGE	BONE	LIVER	TOTAL BODY	THYROID	KIDNEY	LUNG	GI-LLI
INVER	ADULT	1.25E+07	2.11E-02	1.63E-02	5.55E-03	2.50E-04	1.80E-03	1.22E-02	1.50E-02
INVER	TEENAGER	1.99E+06	3.46E-03	2.66E-03	8.57E-04	4.55E-05	2.87E-04	1.95E-03	2.37E-03
INVER	CHILD	1.25E+06	2.49E-03	1.86E-03	5.65E-04	6.80E-05	1.80E-04	1.25E-03	1.49E-03
INVER	TOTAL	1.58E+07	2.71E-02	2.08E-02	6.98E-03	3.63E-04	2.27E-03	1.54E-02	1.89E-02

DILUTION CATCH TIME(HR)-INCLUDES FOOD PROCESSING TIME OF 2.40E+02 HR POPULATION=1.90E+07
 7.00E+00 1.00E+03 2.40E+02

AVERAGE INDIVIDUAL CONSUMPTION (KG/YR) ADULT=1.00E+00 TEEN=7.50E-01 CHILD=3.30E-01

NEPA DOSES

NOTE--TOATL NEPA DOSE MUST INCLUDE SPORT CATCH, DOSES BELOW ARE FOR COMMERCIAL CATCH ONLY

-----DOSE (MAN-REM)-----

PATHWAY	AGE GROUP	USAGE	BONE	LIVER	TOTAL BODY	THYROID	KIDNEY	LUNG	GI-LLI
INVER	ADULT	7.94E+02	3.07E-03	2.37E-03	8.09E-04	3.63E-05	2.63E-04	1.77E-03	2.19E-03
INVER	TEENAGER	1.26E+02	5.04E-04	3.87E-04	1.25E-04	6.63E-06	4.18E-05	2.84E-04	3.45E-04
INVER	CHILD	7.94E+01	3.63E-04	2.72E-04	8.23E-05	9.90E-06	2.63E-05	1.82E-04	2.16E-04
INVER	TOTAL	1.00E+03	3.94E-03	3.03E-03	1.02E-03	5.29E-05	3.31E-04	2.24E-03	2.75E-03

* * * POPULATION WATER CONSUMPTION DOLLARS * * *

-----DOSE (MAN-REM)-----

PATHWAY	AGE GROUP	USAGE	BONE	LIVER	TOTAL BODY	THYROID	KIDNEY	LUNG	GI-LLI
DRINKING	ADULT	2.44E+02	7.69E-09	2.54E-09	2.11E-09	6.95E-11	8.11E-10	3.15E-10	1.98E-10
DRINKING	TEENAGER	3.64E+01	1.33E-09	4.81E-10	2.76E-10	8.19E-12	1.21E-10	6.51E-11	3.12E-11
DRINKING	CHILD	5.20E+01	3.14E-09	1.37E-09	4.76E-10	2.24E-11	1.73E-10	1.73E-10	5.54E-11
DRINKING	TOTAL	3.33E+02	1.22E-08	4.39E-09	2.86E-09	1.00E-10	1.10E-09	5.54E-10	2.84E-10

POPULATION=1.00E+00 DILUTION=1.00E+02 TRANSIT TIME=1.00E+06 HR (INCLUDING 24 HR FOR TREATMENT FACILITY)

AVERAGE INDIVIDUAL CONSUMPTION (L/YR)	ADULT=3.70E+02	TEEN=2.60E+02	CHILD=2.60E+02
---------------------------------------	----------------	---------------	----------------

-----CUMULATIVE TOTAL-----

PATHWAY	AGE GROUP	USAGE	BONE	LIVER	TOTAL BODY	THYROID	KIDNEY	LUNG	GI-LLI
DRINKING	CUMUL TOTAL	3.33E+02	1.22E-08	4.39E-09	2.86E-09	1.00E-10	1.10E-09	5.54E-10	2.84E-10

HYDROSPHERE TRITIUM DOSE

[illegible]

* * * RECREATION POPULATION DOSES * * *

DOSE(MAN-REM)

PATHWAY	AGE GROUP	USAGE	SKIN	TOTAL BODY	THYROID
SHORELINE	TOTAL POPUL	1.66E+07	2.11E-01	1.80E-01	1.80E-01

LOCATION- DOWNSTREAM

DILUTION=0.70E+01 TRANSIT TIME=0.40E+01 HR SHF=0.2

DOSE(MAN-REM)

PATHWAY	AGE GROUP	USAGE	SKIN	TOTAL BODY	THYROID
SHIMMING	TOTAL POPUL	1.66E+07	0.0	1.65E-03	1.65E-03

LOCATION- DOWNSTREAM

DILUTION=0.70E+01 TRANSIT TIME=0.40E+01 HR

DOSE(MAN-REM)

PATHWAY	AGE GROUP	USAGE	SKIN	TOTAL BODY	THYROID
BOATING	TOTAL POPUL	1.66E+07	0.0	8.24E-04	8.24E-04

LOCATION- DOWNSTREAM

DILUTION=0.70E+01 TRANSIT TIME=0.40E+01 HR

* * * JOSE TO BIOTA * * *

MRADS PER YEAR

ILLUTION= 7.00E+00 TRANSIT TIME= 4.00E+00 HR

	INTERNAL	EXTERNAL	TOTAL
FISH	1.88E+00	4.76E-01	2.35E+00
INVERTEBRATE	4.87E-01	9.52E-01	1.44E+00
ALGAE	8.42E+00	8.70E-04	8.43E+00
MUSKRAT	1.45E+01	3.18E-01	1.48E+01
RACCOON	4.02E-01	2.38E-01	6.40E-01
HERON	1.07E+01	3.17E-01	1.10E+01
DUCK	1.43E+01	4.76E-01	1.48E+01

* * * COST-BENEFIT ANALYSIS * * *

NUCLIDE		RELEASE CI/YR	MAN-REM DOSE		MAN-REM PER CURIE	
			TOTAL BODY	THYROID	TOTAL BODY	THYROID
1H	3	1.37E+02	1.08E-03	1.08E-03	7.91E-06	7.91E-06
38SR	89	1.41E-03	3.07E-05	1.62E-08	2.18E-02	1.15E-05
38SR	90	1.67E-03	7.51E-03	2.01E-09	4.50E+00	1.20E-06
55CS	134	3.89E-02	5.67E-01	1.55E-02	1.46E+01	3.98E-01
55CS	137	1.21E-01	1.08E+00	7.17E-02	8.96E+00	5.93E-01
53I	131	8.01E-02	3.61E-04	7.70E-02	4.51E-03	9.62E-01
27CO	58	1.50E-02	4.90E-04	3.89E-04	3.26E-02	2.59E-02
27CO	60	7.44E-02	9.41E-02	9.26E-02	1.26E+00	1.24E+00
26FE	59	8.62E-03	5.71E-04	1.77E-04	6.63E-02	2.05E-02
30ZN	65	8.64E-03	9.04E-03	3.88E-04	1.05E+00	4.49E-02
25MN	54	8.27E-03	2.28E-03	6.84E-04	2.76E-01	8.27E-02
24CR	51	3.09E-02	1.29E-05	1.25E-05	4.18E-04	4.04E-04
40ZR	95	7.51E-03	1.33E-04	1.33E-04	1.77E-02	1.77E-02
42MO	99	1.89E-02	2.41E-05	2.31E-05	1.28E-03	1.22E-03
43TC	99M	2.45E-02	8.45E-06	8.45E-06	3.45E-04	3.45E-04
56BA	140	1.62E-02	4.83E-05	3.64E-05	2.98E-03	2.25E-03
58CE	141	3.89E-03	4.17E-06	4.16E-06	1.07E-03	1.07E-03
26FE	55	1.62E-01	1.58E-02	2.31E-08	9.75E-02	1.43E-07
28NI	63	3.10E-02	8.76E-04	0.0	2.83E-02	0.0
15P	32	9.53E-03	2.85E-01	1.35E-07	2.99E+01	1.42E-05
TOTAL			2.07E+00	2.60E-01		

TABLE 4A

HOURS AT EACH WIND SPEED AND DIRECTION ^a

PERIOD OF RECORD: January 1 - March 31, 1980

STABILITY CLASS: A

ELEVATION: 10 Meters

Wind Direction	Wind Speed (mph) at 10m Level						TOTAL
	1-3	4-7	8-12	13-18	19-24	>24	
N	0	11	15	0	0	0	26
NNE	1	22	6	0	0	0	29
NE	2	17	11	1	0	0	21
ENE	0	1	0	0	0	0	1
E	0	0	0	0	0	0	0
ESE	0	0	0	0	0	0	0
SE	0	0	0	0	0	0	0
SSE	1	0	0	0	0	0	1
S	8	8	0	0	0	0	16
SSW	10	9	0	0	0	0	19
SW	7	7	0	0	0	0	14
WSW	0	0	1	0	0	0	1
W	0	4	0	0	0	0	4
WNW	1	0	0	0	0	0	1
NW	1	5	3	1	0	0	10
NNW	0	12	22	5	0	0	39

VARIABLE

Total	31	86	58	7	0	0	182
-------	----	----	----	---	---	---	-----

Periods of calm (hours): 0

Hours of missing data:

Total hours of missing data for all stability classes this quarter = 52.

^a In the table, record the total number of hours of each category of wind direction for each calendar quarter. Provide similar tables separately for each atmospheric stability class and elevation.

TABLE 4A

HOURS AT EACH WIND SPEED AND DIRECTION ^a

PERIOD OF RECORD: January 1 - March 31, 1980

STABILITY CLASS: B

ELEVATION: 10 Meters

Wind Direction	Wind Speed (mph) at 10m Level						TOTAL
	1-3	4-7	8-12	13-18	19-24	>24	
N	1	8	7	0	0	0	16
NNE	1	7	2	0	0	0	10
NE	0	11	7	1	0	0	19
ENE	0	1	0	0	0	0	1
E	0	0	0	0	0	0	0
ESE	0	0	0	0	0	0	0
SE	0	0	0	0	0	0	0
SSE	0	0	0	0	0	0	0
S	1	1	0	0	0	0	2
SSW	1	1	0	0	0	0	2
SW	1	2	0	0	0	0	3
WSW	1	0	0	0	0	0	1
W	0	2	0	0	0	0	2
WNW	1	4	1	0	0	0	6
NW	0	12	2	1	0	0	15
NNW	1	7	9	5	0	0	22
VARIABLE							
Total	8	56	28	7	0	0	99
Periods of calm (hours):	0						
Hours of missing data:							

^a In the table, record the total number of hours of each category of wind direction for each calendar quarter. Provide similar tables separately for each atmospheric stability class and elevation.

TABLE 4A

HOURS AT EACH WIND SPEED AND DIRECTION ^a

PERIOD OF RECORD: January 1 - March 31, 1980

STABILITY CLASS: C

ELEVATION: 10 Meters

Wind Direction	Wind Speed (mph) at 10m Level						TOTAL
	1-3	4-7	8-12	13-18	19-24	>24	
N	4	10	9	0	0	0	23
NNE	3	13	2	0	0	0	18
NE	1	9	16	1	0	0	27
ENE	0	7	1	0	0	0	8
E	1	2	0	0	0	0	3
ESE	0	0	0	0	0	0	0
SE	0	0	0	0	0	0	0
SSE	1	1	0	0	0	0	2
S	2	0	0	0	0	0	2
SSW	2	1	0	0	0	0	3
SW	4	2	0	0	0	0	6
WSW	0	1	0	0	0	0	1
W	2	1	0	0	0	0	3
WNW	0	1	0	0	0	0	1
NW	0	5	3	2	0	0	10
NNW	0	6	13	4	0	0	23

VARIABLE

Total	20	59	44	7	0	0	130
Periods of calm (hours):		0					
Hours of missing data:							

^a In the table, record the total number of hours of each category of wind direction for each calendar quarter. Provide similar tables separately for each atmospheric stability class and elevation.

TABLE 4A

HOURS AT EACH WIND SPEED AND DIRECTION ^a

PERIOD OF RECORD: January 1 - March 31, 1980

STABILITY CLASS: D

ELEVATION: 10 Meters

Wind Direction	Wind Speed (mph) at 10m Level						TOTAL
	1-3	4-7	8-12	13-18	19-24	>24	
N	15	118	52	1	0	0	186
NNE	10	56	13	2	0	0	81
NE	12	101	74	20	0	0	207
ENE	11	33	5	1	0	0	50
E	7	7	0	0	0	0	14
ESE	3	1	0	0	0	0	4
SE	2	2	0	0	0	0	4
SSE	0	0	0	0	0	0	0
S	11	4	0	0	0	0	15
SSW	18	5	0	0	0	0	23
SW	14	14	5	0	0	0	33
WSW	10	6	0	0	0	0	16
W	16	9	0	0	0	0	25
WNW	9	23	2	0	0	0	34
NW	8	52	24	1	0	0	85
NNW	9	85	82	9	0	0	185

VARIABLE

Total	155	516	257	34	0	0	962
-------	-----	-----	-----	----	---	---	-----

Periods of calm (hours): 0

Hours of missing data:

^a In the table, record the total number of hours of each category of wind direction for each calendar quarter. Provide similar tables separately for each atmospheric stability class and elevation.

TABLE 4A

HOURS AT EACH WIND SPEED AND DIRECTION ^a

PERIOD OF RECORD: January 1 - March 31, 1980

STABILITY CLASS: E

ELEVATION: 10 Meters

Wind Direction	Wind Speed (mph) at 10m Level						TOTAL
	1-3	4-7	8-12	13-18	19-24	>24	
N	12	21	4	0	0	0	37
NNE	15	14	3	0	0	0	32
NE	32	41	12	10	0	0	95
ENE	28	23	1	0	0	0	52
E	24	7	0	0	0	0	31
ESE	7	0	0	0	0	0	7
SE	15	0	0	0	0	0	15
SSE	12	0	0	0	0	0	12
S	34	2	0	0	0	0	36
SSW	63	7	4	0	0	0	74
SW	42	11	4	0	0	0	57
WSW	31	3	0	0	0	0	34
W	32	1	1	0	0	0	34
WNW	14	6	0	0	0	0	20
NW	10	17	0	0	0	0	27
NNW	8	10	2	1	0	0	21

VARIABLE

Total	379	163	31	11	0	0	584
Periods of calm (hours):	1						
Hours of missing data:							

^a In the table, record the total number of hours of each category of wind direction for each calendar quarter. Provide similar tables separately for each atmospheric stability class and elevation.

TABLE 4A

HOURS AT EACH WIND SPEED AND DIRECTION ^a

PERIOD OF RECORD: January 1 - March 31, 1980

STABILITY CLASS: F

ELEVATION: 10 Meters

Wind Direction	Wind Speed (mph) at 10m Level						TOTAL
	<u>1-3</u>	<u>4-7</u>	<u>8-12</u>	<u>13-18</u>	<u>19-24</u>	<u>>24</u>	
N	6	0	0	0	0	0	6
NNE	7	0	0	0	0	0	7
NE	10	2	0	0	0	0	12
ENE	12	0	0	0	0	0	12
E	8	5	0	0	0	0	13
ESE	0	0	0	0	0	0	0
SE	2	0	0	0	0	0	2
SSE	2	2	1	0	0	0	5
S	12	6	0	0	0	0	18
SSW	15	<u>1</u>	0	0	0	0	16
SW	8	0	0	0	0	0	8
WSW	10	0	0	0	0	0	10
W	9	1	0	0	0	0	10
WNW	9	0	0	1	0	0	10
NW	1	0	1	0	0	0	2
NNW	2	0	0	0	0	0	2

VARIABLE

Total	113	17	2	1	0	0	133
Periods of calm (hours):	2						
Hours of missing data:							

^a In the table, record the total number of hours of each category of wind direction for each calendar quarter. Provide similar tables separately for each atmospheric stability class and elevation.

TABLE 4A

HOURS AT EACH WIND SPEED AND DIRECTION ^a

PERIOD OF RECORD: January 1 - March 31, 1980

STABILITY CLASS: G

ELEVATION: 10 Meters

Wind Direction	Wind Speed (mph) at 10m Level						TOTAL
	1-3	4-7	8-12	13-18	19-24	>24	
N	2	0	0	0	0	0	2
NNE	2	0	0	0	0	0	2
NE	4	5	1	0	0	0	10
ENE	2	2	0	0	0	0	4
E	3	0	0	0	0	0	3
ESE	0	0	0	0	0	0	0
SE	0	0	0	0	0	0	0
SSE	1	0	0	0	0	0	1
S	4	3	0	0	0	0	7
SSW	3	1	0	0	0	0	4
SW	1	0	0	0	0	0	1
WSW	0	0	0	0	0	0	0
W	2	0	0	0	0	0	2
WNW	1	0	0	0	0	0	1
NW	0	0	0	0	0	0	0
NNW	2	0	0	0	0	0	2

VARIABLE

Total	27	11	1	0	0	0	39
Periods of calm (hours):	0						
Hours of missing data:							

^a In the table, record the total number of hours of each category of wind direction for each calendar quarter. Provide similar tables separately for each atmospheric stability class and elevation.

TABLE 4A

HOURS AT EACH WIND SPEED AND DIRECTION^a

PERIOD OF RECORD: April 1 - June 30, 1980

STABILITY CLASS: A

ELEVATION: 10 Meters

Wind Direction	Wind Speed (mph) at 10m Level						TOTAL
	1-3	4-7	8-12	13-18	19-24	>24	
N	13	46	5	1	0	0	65
NNE	5	45	4	0	0	0	54
NE	1	21	6	0	0	0	28
ENE	1	0	0	0	0	0	1
E	0	0	0	0	0	0	0
ESE	0	0	0	0	0	0	0
SE	0	0	0	0	0	0	0
SSE	0	0	0	0	0	0	0
S	14	37	0	0	0	0	51
SSW	27	43	0	0	0	0	70
SW	17	25	0	0	0	0	42
WSW	9	12	0	0	0	0	21
W	11	18	0	0	0	0	29
WNW	4	12	0	0	0	0	16
NW	5	26	2	0	0	0	33
NNW	5	19	8	3	0	0	35

VARIABLE

Total	112	304	25	4	0	0	445
-------	-----	-----	----	---	---	---	-----

Periods of calm (hours): 0

Hours of missing data: Total hours of missing data for all stability classes this quarter = 87.

^a In the table, record the total number of hours of each category of wind direction for each calendar quarter. Provide similar tables separately for each atmospheric stability class and elevation.

TABLE 4A

HOURS AT EACH WIND SPEED AND DIRECTION ^a

PERIOD OF RECORD: April 1 - June 30, 1980

STABILITY CLASS: B

ELEVATION: 10 Meters

Wind Direction	Wind Speed (mph) at 10m Level						TOTAL
	1-3	4-7	8-12	13-18	19-24	>24	
N	1	7	4	0	0	0	12
NNE	3	5	0	0	0	0	8
NE	1	8	1	0	0	0	10
ENE	1	3	0	0	0	0	4
E	1	0	0	0	0	0	1
ESE	0	0	0	0	0	0	0
SE	0	0	0	0	0	0	0
SSE	1	0	0	0	0	0	1
S	8	2	0	0	0	0	10
SSW	5	4	0	0	0	0	9
SW	6	3	1	0	0	0	10
WSW	2	2	0	0	0	0	4
W	3	4	2	0	0	0	9
WNW	3	1	0	0	0	0	4
NW	4	3	1	0	0	0	8
NNW	2	4	1	0	0	0	7
VARIABLE							
Total	41	46	10	0	0	0	97
Periods of calm (hours):	0						
Hours of missing data:							

^a In the table, record the total number of hours of each category of wind direction for each calendar quarter. Provide similar tables separately for each atmospheric stability class and elevation.

TABLE 4A

HOURS AT EACH WIND SPEED AND DIRECTION ^a

PERIOD OF RECORD: April 1 - June 30, 1980

STABILITY CLASS: C

ELEVATION: 10 Meters

Wind Direction	Wind Speed (mph) at 10m Level						TOTAL
	1-3	4-7	8-12	13-18	19-24	>24	
N	3	8	2	0	0	0	13
NNE	4	6	0	0	0	0	10
NE	4	6	2	0	0	0	12
ENE	1	1	0	0	0	0	2
E	2	0	0	0	0	0	2
ESE	0	0	0	0	0	0	0
SE	1	0	0	0	0	0	1
SSE	0	0	0	0	0	0	0
S	8	2	0	0	0	0	10
SSW	4	2	0	0	0	0	6
SW	3	2	0	0	0	0	5
WSW	6	0	0	0	0	0	6
W	3	3	0	0	0	0	6
WNW	2	1	0	0	0	0	3
NW	1	4	0	0	0	0	5
NNW	0	4	1	0	0	0	5

VARIABLE

Total	42	39	5	0	0	0	86
-------	----	----	---	---	---	---	----

Periods of calm (hours): 0

Hours of missing data:

^a In the table, record the total number of hours of each category of wind direction for each calendar quarter. Provide similar tables separately for each atmospheric stability class and elevation.

TABLE 4A

HOURS AT EACH WIND SPEED AND DIRECTION^a

PERIOD OF RECORD: April 1 - June 30, 1980

STABILITY CLASS: D

ELEVATION: 10 Meters

Wind Direction	Wind Speed (mph) at 10m Level						TOTAL
	1-3	4-7	8-12	13-18	19-24	>24	
N	6	23	8	1	0	0	38
NNE	17	44	7	0	0	0	68
NE	23	41	11	0	0	0	75
ENE	16	14	2	0	0	0	32
E	14	4	0	0	0	0	18
ESE	8	0	0	0	0	0	8
SE	10	2	0	0	0	0	12
SSE	12	3	0	0	0	0	15
S	36	11	0	0	0	0	47
SSW	40	19	0	0	0	0	59
SW	17	13	0	0	0	0	30
WSW	15	5	0	0	0	0	20
W	11	15	4	0	0	0	30
WNW	7	14	2	0	0	0	23
NW	4	16	5	0	0	0	25
NNW	5	13	4	0	0	0	22
VARIABLE							
Total	241	237	43	1	0	0	522
Periods of calm (hours):	0						
Hours of missing data:							

^a In the table, record the total number of hours of each category of wind direction for each calendar quarter. Provide similar tables separately for each atmospheric stability class and elevation.

TABLE 4A

HOURS AT EACH WIND SPEED AND DIRECTION ^a

PERIOD OF RECORD: April 1 - June 30, 1980

STABILITY CLASS: E

ELEVATION: 10 Meters

Wind Direction	Wind Speed (mph) at 10m Level						TOTAL
	1-3	4-7	8-12	13-18	19-24	>24	
N	24	34	0	0	0	0	58
NNE	25	24	1	0	0	0	50
NE	26	35	2	0	0	0	63
ENE	26	16	1	0	0	0	43
E	29	2	0	0	0	0	31
ESE	14	1	0	0	0	0	15
SE	13	1	0	0	0	0	14
SSE	9	1	0	0	0	0	10
S	55	6	0	0	0	0	61
SSW	72	45	3	0	0	0	120
SW	49	18	3	0	0	0	70
WSW	25	1	0	0	0	0	26
W	22	5	1	0	0	0	28
WNW	17	18	1	0	0	0	36
NW	9	16	1	0	0	0	26
NNW	11	12	0	0	0	0	23
VARIABLE							
Total	426	235	13	0	0	0	674
Periods of calm (hours):	3						
Hours of missing data:							

^a In the table, record the total number of hours of each category of wind direction for each calendar quarter. Provide similar tables separately for each atmospheric stability class and elevation.

TABLE 4A

HOURS AT EACH WIND SPEED AND DIRECTION ^a

PERIOD OF RECORD: April 1 - June 30, 1980

STABILITY CLASS: F

ELEVATION: 10 Meters

Wind Direction	Wind Speed (mph) at 10m Level						TOTAL
	1-3	4-7	8-12	13-18	19-24	>24	
N	18	0	0	0	0	0	18
NNE	16	0	0	0	0	0	16
NE	28	9	1	0	0	0	38
ENE	30	7	0	0	0	0	37
E	8	4	0	0	0	0	12
ESE	5	0	0	0	0	0	5
SE	9	0	0	0	0	0	9
SSE	6	0	0	0	0	0	6
S	14	1	0	0	0	0	15
SSW	13	1	1	0	0	0	15
SW	16	0	0	0	0	0	16
WSW	4	0	0	0	0	0	4
W	11	0	0	0	0	0	11
WNW	6	0	0	0	0	0	6
NW	3	0	0	0	0	0	3
NNW	7	0	0	0	0	0	7
VARIABLE							
Total	194	22	2	0	0	0	218
Periods of calm (hours):		2					
Hours of missing data:							

^a In the table, record the total number of hours of each category of wind direction for each calendar quarter. Provide similar tables separately for each atmospheric stability class and elevation.

TABLE 4A

HOURS AT EACH WIND SPEED AND DIRECTION ^a

PERIOD OF RECORD: April 1 - June 30, 1980

STABILITY CLASS: G

ELEVATION: 10 Meters

Wind Direction	Wind Speed (mph) at 10m Level						TOTAL
	1-3	4-7	8-12	13-18	19-24	>24	
N	4	0	0	0	0	0	4
NNE	5	0	0	0	0	0	5
NE	8	5	0	0	0	0	13
ENE	8	3	0	0	0	0	11
E	0	1	0	0	0	0	1
ESE	1	0	0	0	0	0	1
SE	0	0	0	0	0	0	0
SSE	1	0	0	0	0	0	1
S	1	0	0	0	0	0	1
SSW	0	0	0	0	0	0	0
SW	4	0	0	0	0	0	4
WSW	0	0	0	0	0	0	0
W	3	0	0	0	0	0	3
WNW	2	0	0	0	0	0	2
NW	2	0	0	0	0	0	2
NNW	2	0	0	0	0	0	2
VARIABLE							
Total	41	9	0	0	0	0	50
Periods of calm (hours):	0						
Hours of missing data:							

^a In the table, record the total number of hours of each category of wind direction for each calendar quarter. Provide similar tables separately for each atmospheric stability class and elevation.