


Harry G. Woodbury
Executive Vice President

Consolidated Edison Company of New York, Inc.
4 Irving Place, New York, N Y 10003
Telephone (212) 460-6001

September 25, 1973

3
~~50-206~~
50-262
~~50-206~~

Mr. George W. Knighton
Chief, Environmental Projects
Branch #1
Directorate of Licensing
United States Atomic Energy Commission
Washington, D.C. 20545



Dear Mr. Knighton:

At the July 25-26, 1973 meeting at Oak Ridge, Tennessee to discuss mathematical modelling of the impact of Indian Point on striped bass populations, considerable interest was expressed by ORNL personnel in the preliminary findings of the Texas Instruments river-wide ichthyoplankton and beach seining programs.

These preliminary findings indicate that many of the striped bass eggs spawned in the estuary concentrate and remain in deep "holes" in the river bottom in the vicinity of Storm King Mountain and in adjacent "holes" below it. Evidence of this can be seen in the attached table which gives a comparison of striped bass eggs and larvae collected in the Tucker Trawl and the Epibenthic Sled. In addition, beach seining in the latter half of June has yielded striped bass larvae with total body lengths of 15-30 mm. This indicates that striped bass larvae are heading to the beaches (where they are behaviorally non-entrainable) at a much earlier age than has been previously assumed.

These findings can have a significant bearing on the development of models to predict the impact of Indian Point on striped bass populations. Therefore I suggest that a second meeting be held with the same parties present as the July 25-26, 1973 meeting in order to discuss in detail how this data should be incorporated into striped bass models. If you agree, this meeting might be held at our offices in New York in late

7644

8111230269 730925
PDR ADOCK 05000286
A PDR

B-2015

October, say October 25-26, 1973. By that time we should have a report from Texas Instruments providing complete results of the May and June ichthyoplankton survey work and the June and July beach seining. Data presented in this report should provide a good basis for discussion at this meeting and copies of this report will be sent to you in advance of the meeting. Please advise me whether you agree to holding this meeting and whether the date and place suggested are suitable.

Very truly yours,

Harry G. Woodbury
Harry G. Woodbury

rsh/klg

cc: Dr. Mary Jane Oestmann
Myron Karman, Esq.
Nicholas A. Robinson, Esq.
Angus Mac Beth, Esq.
J. Bruce Mac Donald, Esq.
Harry H. Voigt, Esq.
Dr. Webb Van Winkle

Table 1. Counts of striped bass eggs and larvae taken by the Ichthyoplankton Survey, May 15 and May 16, 1973 between river mile 54 and river mile 43 inclusive.



COUNTS **								
GEAR	SAMPLE #	RIVER DEPTH (ft)	R.M.*	SAMPLE DEPTH (ft)	EGGS	LARVAE 1	LARVAE 2	LARVAE 3
Tucker								
Trawl	300	56	54	47	0	0	0	0
	301	55	54	28	19	3	0	0
	302	65	54	26	0	0	0	0
	309	73	51	60	7	0	0	0
	310	73	51	41	0	2	0	0
	311	70	51	14	0	2	0	0
	Average				4.3	1.2	0.0	0.0
Epibenthic								
Sled	303	66	54	66	304	18	0	0
	304	65	54	65	35	6	0	0
	305	65	54	65	126	26	0	0
	306	76	51	76	984	4	0	0
	307	76	51	76	698	22	1	0
	308	76	51	76	770	25	0	0
	312	80	48	80	441	35	0	0
	313	80	48	80	256	15	0	0
	315	120	47	120	280	10	0	7
	316	120	47	120	67	0	15	0
	317	120	47	120	60	10	0	0
	318	125	45	125	601	0	74	1
	319	125	45	125	37	31	0	0
	320	125	45	125	800	118	0	0
	321	35	43	35	15	0	8	0
	322	35	43	35	0	5	7	0
	323	35	43	35	61	9	0	0
	Average				325.6	19.1	5.8	0.5

*R.M. = river mile

** Larvae 1 = Yolk Sack Larvae
Larvae 2 = Post Yolk Sack Larvae

Larvae 3 = Juvenile

Regulatory Docket File



10 11 12 13
14 15 16 17
18 19 20 21

22 23 24 25
26 27 28 29
30 31 32 33