


- LEGEND**
- ★ HAR Site
 - Main Dam
 - ▲ Sears No. 1 and Groce No. 1 Wells (Bain and Brown, 1981)
 - 8-km (5-mi.) Radius from Site
 - 40-km (25-mi.) Radius from Site
 - Paleozoic Faults
 - Mesozoic Faults (dashed where inferred, dotted where concealed) (Ebasco, 1975, CP&L, 1983, NCGS, 1985, Wooten et al., 1996, Harding Lawson, 1997, NCGS, 2006, and this study)
 - 49 Cretaceous and Cenozoic Faults (numbers refer to descriptions in Table 2.5.1-201; from Parker, 1979, and Prowell, 1983)
 - Paleozoic Folds (dashed where inferred)
 - Mesozoic Folds (dashed where inferred) (NCGS, 2006)
 - Dikes (dashed where inferred) (CP&L, 1983, NCGS, 1985, NCGS, 2006, and Technos, this study)

Geologic Units (NCGS, 1985) (See Figure 2.5.1-230, Sheet 2 for Unit Descriptions)

Tec	TRcs	CZfv
Tpy	PPg	CZfv1
Tt	PPmg	CZg
Kb	PzZg	CZig
Kc	PzZq	CZiv
Km	PzZu	CZmd
Jd	CZam	CZph
TRc	CZbg	CZv
TRcc	CZc	CZve
TRcp	CZfg	CZmv

Note N1: This fault included in NCGS (2006) unpublished mapping appears to represent variations in metamorphic grade adjacent to the pluton rather than a fault contact.



Progress Energy Carolinas
**Shearon Harris Nuclear Power Plant
 Units 2 and 3
 Part 2, Final Safety Analysis Report**
 New Hill, North Carolina

Site Vicinity Geologic Map
 (40-km [25-mi.] Radius)

FIGURE 2.5.1-230 (SHEET 1 OF 2) Rev. 1

DESCRIPTION OF GEOLOGIC UNITS

Pliocene-Quaternary	Tt	TERRACE DEPOSITS AND UPLAND SEDIMENT - Gravel, clayey sand, and sand and minor iron-oxide cemented sandstone.	Paleozoic	PPg	GRANITIC ROCK (Pennsylvanian to Permian, 265-325 m.y.) - Megacrystic to equigranular.	Late Proterozoic - Cambrian	CZmd	METAMUDSTONE AND META-ARGILLITE - Bedding plane and axial planar cleavage common; interbedded with metasandstone, metaconglomerate, and metavolcanic rock.
	Tec	CASTLE HAYNE FORMATION - Comfort Member and New Hanover Member, undivided. Comfort Member: Bryozoan-echinoid skeletal limestone, locally dolomitized, solution cavities common. New Hanover Member: Phosphate-pebble conglomerate, micritic, thin; restricted to basal part of Castle Hayne Formation in southeastern counties.		PPmg	FOLIATED TO MASSIVE GRANITIC ROCK (Pennsylvanian to Permian, 270-320 m.y.) - Megacrystic to equigranular.		CZph	PHYLLITE AND SCHIST - Locally laminated and pyritic; includes phyllonite, sheared fine-grained metasediment, and metavolcanic rock.
Tertiary	Tpy	YORKTOWN FORMATION AND DUPLIN FORMATION, UNDIVIDED - Yorktown Formation: Fossiliferous clay with varying amounts of fine-grained sand, bluish gray, shell material commonly concentrated in lenses; mainly in area north of Neuse River. Duplin Formation: Shelly, medium- to coarse-grained sand, sandy marl, and limestone, bluish gray; mainly in area south of Neuse River.	Paleozoic	PzZg	METAMORPHOSED GABBRO AND DIORITE - Foliated to massive.	Late Proterozoic - Cambrian	CZve	METAVOLCANIC EPICLASTIC ROCK - Metamorphosed argillite, mudstone, volcanic sandstone, conglomerate, and volcanic rock.
	Kb	BLACK CREEK FORMATION - Clay gray to black, lignitic; contains thin beds and laminae of fine-grained micaceous sand and thick lenses of cross-bedded sand. Glauconitic, fossiliferous clayey sand lenses in upper part.		PzZu	META-ULTRAMAFIC ROCK - Metamorphosed dunite and peridotite; serpentinite, soapstone, and other altered ultramafic rock. Only larger bodies shown.		CZmv	MAFIC METAVOLCANIC ROCK - Metamorphosed basalt flows and tuffs, dark green to black; interbedded with felsic and intermediate metavolcanic rock and metamudstone.
Cretaceous	Km	MIDDENDORF FORMATION - Sand, sandstone, and mudstone, gray to pale gray with an orange cast, mottled; clay balls and iron-cemented concretions common, beds laterally discontinuous, cross bedding common.	Late Proterozoic - Cambrian	CZam	AMPHIBOLITE - Metamorphosed mafic extrusive and intrusive rock; includes hornblende gneiss, thin layers of mica schist, and small non-layered masses of metadiorite and metagabbro.	Late Proterozoic - Cambrian	Source: NCGS (1985)	
	Kc	CAPE FEAR FORMATION - Sandstone and sandy mudstone, yellowish gray to bluish gray, mottled red to yellowish orange, indurated, graded and laterally continuous bedding, blocky clay, faint cross-bedding, feldspar and mica common.		CZbg	BIOTITE GNEISS AND SCHIST - Inequigranular and megacrystic; in places contains garnet; interlayered and gradational with mica schist and amphibolite; includes small masses of granitic rock.		CZfc	FELSIC MICA GNEISS - Interlayered with graphitic mica schist and mica-garnet schist, commonly with kyanite; minor hornblende gneiss.
	Jd	DIABASE - Dikes, gray to black.		CZc	VOLCANIC METACONGLOMERATE - Includes metagraywacke and metamudstone.		CZfg	FELSIC METAVOLCANIC ROCK - Metamorphosed dacitic to rhyolitic flows and tuffs, light gray to greenish gray; interbedded with mafic and intermediate metavolcanic rock, meta-argillite, and metamudstone.
Jurassic	TRc	CHATHAM GROUP (undivided) - Conglomerate, fanglomerate, sandstone, and mudstone.	Late Proterozoic - Cambrian	CZfv	FELSIC METAVOLCANIC ROCK - Metamorphosed dacitic to rhyolitic flows and tuffs, light gray to greenish gray; interbedded with mafic and intermediate metavolcanic rock, meta-argillite, and metamudstone.	Late Proterozoic - Cambrian	CZg	METAMORPHOSED GRANITIC ROCK (Late Proterozoic to late Cambrian, 520-650m.y.) - Megacrystic, well-foliated, locally contains hornblende.
	TRcs	SANFORD FORMATION - Conglomerate, fanglomerate, sandstone, and mudstone.		CZig	INJECTED GNEISS - Biotite gneiss and schist intruded by numerous sills and dikes of granite, pegmatite, and apatite; minor hornblende gneiss.		CZiv	INTERMEDIATE METAVOLCANIC ROCK - Metamorphosed andesitic tuffs and flows, medium to dark grayish green; minor felsic and mafic metavolcanic rock.
Triassic	TRcc	CUMNOCK FORMATION - Sandstone and mudstone, gray to black; coal beds and carbonaceous shale. Grades into Pekin and Sanford formations.	Late Proterozoic - Cambrian	CZlg	LINEATED-FELSIC MICA GNEISS - White to pink with strong lineation of muscovite-biotite streaks and prismatic quartz aggregates; planar foliation and layering weak; minor mica schist and hornblende gneiss.	Late Proterozoic - Cambrian	Progress Energy Carolinas	
	TRcp	PEKIN FORMATION - Conglomerate, sandstone, and mudstone.		CZiv	INTERMEDIATE METAVOLCANIC ROCK - Metamorphosed andesitic tuffs and flows, medium to dark grayish green; minor felsic and mafic metavolcanic rock.		Shearon Harris Nuclear Power Plant Units 2 and 3	
							Part 2, Final Safety Analysis Report	
						New Hill, North Carolina		
						Site Vicinity Geologic Map (40-km [25-mi.] Radius)		
						FIGURE 2.5.1-230 (SHEET 2 OF 2) Rev. 1		