

NUREG/CR-6112  
BNL-NUREG-52394

---

---

# Impact of Reduced Dose Limits on NRC Licensed Activities

## Major Issues in the Implementation of ICRP/NCRP Dose Limit Recommendations

### Final Report

---

---

Manuscript Completed: March 1995  
Date Published: May 1995

Prepared by  
C. B. Meinhold

G. E. Powers, NRC Project Manager

Brookhaven National Laboratory  
Upton, NY 11973

Prepared for  
Division of Regulatory Applications  
Office of Nuclear Regulatory Research  
U.S. Nuclear Regulatory Commission  
Washington, DC 20555-0001  
NRC Job Code L1285

**MASTER**

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED  
*ML*

...the ... of ...

## **DISCLAIMER**

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, make any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

## **DISCLAIMER**

**Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.**

## Abstract


This report summarizes information required to estimate, at least qualitatively, the potential impacts of reducing occupational dose limits below those given in 10 CFR 20 (Revised).

For this study, a questionnaire was developed and widely distributed to the radiation protection community. The resulting data together with data from existing surveys and sources were used to estimate the impact of three dose-limit options; 10 mSv yr<sup>-1</sup> (1 rem yr<sup>-1</sup>), 20 mSv yr<sup>-1</sup> (2 rem yr<sup>-1</sup>), and a combination of an annual limit of 50 mSv yr<sup>-1</sup> (5 rem yr<sup>-1</sup>) coupled with a cumulative limit, in rem, equal to age in years. Due to the somewhat small number of responses and the lack of data in some specific areas, a working committee of radiation protection experts from a variety of licensees was employed to ensure the exposure data were representative.

The following overall conclusions were reached:

- (1) Although 10 mSv yr<sup>-1</sup> is a reasonable limit for many licensees, such a limit could be extraordinarily difficult to achieve and potentially destructive to the continued operation of some licensees, such as nuclear power, fuel fabrication, and medicine.
- (2) Twenty mSv yr<sup>-1</sup> as a limit is possible for some of these groups, but for others it would prove difficult.
- (3) Fifty mSv yr<sup>-1</sup> and age in 10s of mSv appear reasonable for all licensees, both in terms of the lifetime risk of cancer and severe genetic effects to the most highly exposed workers, and the practicality of operation. In some segments of the industry, this acceptability is based on the adoption of a "grandfather clause" for those people exceeding or close to exceeding the cumulative limit at this time.

Detailed information for fuel fabrication, waste management, manufacturing, well logging, and industrial radiography is sparse and such data would be useful for a full understanding of the potential impact of any reduction in the dose limits.



# Contents

	Page
Abstract .....	iii
Executive Summary .....	v
Foreword .....	xi
Acknowledgments .....	xii
1 Introduction .....	1
2 Historical Background and Literature Survey .....	2
2.1 1928 to 1977 .....	2
2.2 1977 to 1987 .....	2
2.3 1987 to 1994 .....	3
2.4 Background Summary .....	5
3 Data Gathering .....	6
3.1 Existing Surveys .....	6
3.1.1 1992 Edison Electric Institute (EEI) Report on Dose Limits and Guidelines .....	6
3.1.2 Department of Energy Report (DOE) on the Implications of the BEIR V Report .....	6
3.1.3 Nuclear Regulatory Commission (NRC) Radiation Exposure Information and Reporting System (REIRS) .....	6
3.1.4 Environmental Protection Agency (EPA) Report on Occupational Exposure to Ionizing Radiation in the United States .....	7
3.2 Survey Performed for this Report .....	7
3.2.1 Questionnaire Design .....	7
3.2.1.1 Options for Potential Dose Limits .....	7
3.2.1.2 Impacts of Reduced Dose Limits .....	7
3.2.1.3 1989 Dose Experience .....	8
3.2.2 Questionnaire Distribution .....	8
3.2.3 Working Committee on the Impact of Reduced Dose Limits .....	8
3.3 Comments Received to the Draft NUREG Report .....	8
4 Survey Results .....	9
4.1 Edison Electric Institute (EEI) Report .....	9
4.1.1 Administrative Control Levels .....	9
4.1.2 Annual Reported Doses for 1985 and 1989 .....	9
4.1.3 Cumulative Dose Administrative Guidelines .....	9
4.1.4 Cumulative Reported Doses for 1989 .....	10
4.1.5 Projected Cumulative Doses for 1994 .....	10

	Page
4.1.6 Effects of Changing the Annual Dose Guidance .....	10
4.1.7 Effects of Establishing a Cumulative Dose Limit .....	11
4.2 Department of Energy (DOE) Report .....	11
4.2.1 Cost Impact .....	11
4.2.2 Annual Reported Doses, 1978 to 1988 .....	12
4.2.3 Lifetime Cumulative Exposure Limits .....	13
4.2.4 Impact on Facility Operations .....	13
4.3 Selected 1990 Data from NRC REIRS .....	14
4.4 Information Obtained from the 1984 EPA Report .....	18
4.4.1 Male and Female Workers in the Nuclear Industry .....	18
4.4.2 Correlation of Radiation Dose with Age .....	18
4.4.3 Males .....	18
4.4.4 Females .....	19
4.5 Responses to the Request for Comment .....	19
4.5.1 Nuclear Power Reactor and Nuclear Power Reactor Contractors .....	19
4.5.1.1 Organization of Licensees .....	19
4.5.1.2 Three Separate Responses were Received from Three of the Nuclear Power Plant Sites .....	21
4.5.2 Test and Measurement Including Industrial Radiography .....	22
4.5.3 Manufacturing and Distribution Including Cyclotron Produced Radiopharmaceuticals .....	22
4.5.4 Fuel Fabrication, UF <sub>6</sub> Production .....	22
5 Questionnaire Results Obtained in this Survey .....	23
5.1 Medical/Dental and Veterinary Practice .....	23
5.2 Nuclear Power Reactors .....	25
5.3 Nuclear Power Reactor Contractors .....	29
5.4 Test and Measurements Including Industrial Radiography .....	31
5.5 Universities .....	33
5.6 Manufacturing and Distribution, Including Cyclotron Produced Radiopharmaceuticals .....	34
5.7 Waste Management .....	36
5.8 Fuel Fabrication, UF <sub>6</sub> Production .....	37
5.9 Well Logging .....	39
5.10 Others (R&D, Regulatory) .....	40
6 High Dose Groups Within an Industry .....	41
6.1 Introduction .....	41
6.2 NRC-Sponsored Study on High Dose Group Workers .....	41
6.2.1 Analysis of Dose Data Obtained in the Study .....	44
6.2.1.1 Pressurized Water Reactor Data .....	44
6.2.1.2 Boiling Water Reactor Data .....	44

	Page
6.2.1.3 Contractor Data .....	44
<b>7 Costs Associated With Dose Reduction Modifications in the Nuclear Power Industry .....</b>	<b>46</b>
7.1 Introduction .....	46
7.2 Costs (and the Related Dose Saved) of Selected Modifications Which Might be Employed to Reduce Exposure .....	46
7.3 Estimated Impacts .....	51
<b>8 Summary .....</b>	<b>55</b>
8.1 Medical/Dental/Veterinary .....	55
8.1.1 1 Rem Yr <sup>-1</sup> .....	55
8.1.2 2 Rem Yr <sup>-1</sup> .....	55
8.1.3 5 Rem Yr <sup>-1</sup> and Cumulative Dose in Rem Less Than Age in Years Limit .....	55
8.2 Nuclear Power Reactor Plants and Their Contractors .....	55
8.2.1 1 Rem Per Yr <sup>-1</sup> .....	55
8.2.2 2 Rem Yr <sup>-1</sup> .....	56
8.2.3 5 Rem Yr <sup>-1</sup> and Cumulative Dose in Rem Less Than Age in Years Limit .....	56
8.3 Test and Measurement Including Industrial Radiography .....	57
8.3.1 1 Rem Yr <sup>-1</sup> .....	57
8.3.2 2 Rem Yr <sup>-1</sup> .....	57
8.3.3 5 Rem Yr <sup>-1</sup> and Cumulative Dose in Rem Less Than Age in Years Limit .....	57
8.4 Universities not Including Medical, Dental, or Veterinary Schools .....	57
8.4.1 1 Rem Yr <sup>-1</sup> Limit .....	57
8.4.2 2 Rem Yr <sup>-1</sup> Limit .....	58
8.4.3 5 Rem Yr <sup>-1</sup> and Cumulative Dose in Rem Less Than Age in Years Limit .....	58
8.5 Manufacturing and Distribution Including Cyclotron-Produced Radiopharmaceuticals .....	58
8.5.1 1 Rem Yr <sup>-1</sup> Limit .....	58
8.5.2 2 Rem Yr <sup>-1</sup> Limit .....	58
8.5.3 5 Rem Yr <sup>-1</sup> and Cumulative Dose in Rem Less Than Age in Years Limit .....	58
8.6 Waste Management .....	58
8.6.1 1 Rem Yr <sup>-1</sup> Limit .....	58
8.6.2 2 Rem Yr <sup>-1</sup> Limit .....	58
8.6.3 5 Rem Yr <sup>-1</sup> and Cumulative Dose in Rem Less Than Age in Years Limit .....	58
8.7 Fuel Fabrication, UF <sub>6</sub> Production .....	58
8.8 Well Logging .....	59
8.9 General Conclusions .....	59
<b>9 References .....</b>	<b>61</b>
<b>Appendix A .....</b>	<b>64</b>

## Figures

	Page
4.1 Annual Site Doses for Utility Personnel (UT) and Total Plant Workers (TO) .....	9
4.2 Annual Site Doses for Utility Personnel (UT) and Total Plant Workers (TO) for 1985 and 1989 ...	9
4.3 Cumulative Site Doses for Utility and Contractor Personnel for 1989 .....	10
4.4 Projected Cumulative Site Doses for 1994 Utility and Contractor Personnel .....	10
4.5 Average Annual Dose Equivalent for DOE Workers with Measurable Exposure, 1978-1988 ....	12
4.6 Number of DOE Employees .....	12
4.7 Percent of Male Radiation Workers in Various Sectors .....	18
4.8 Percent of Female Radiation Workers in Various Sectors .....	18
7.1 Person-Rem Values .....	52
7.2 Total Number of Reactors and Collective Dose .....	53

## Tables

4.1 Annual Exposure Data* 1990 .....	15
4.2 Annual Exposure Information for Industrial Radiographers* 1989 .....	16
4.3 Annual Exposure Information for Fuel Fabricators* 1989 .....	16
4.4 Annual Exposure Information for Manufacturers and Distributors* 1989 .....	16
4.5 Summary of Annual Whole Body Distributions By Year and Reactor Type 1989 .....	17
4.6 Mean Annual Dose Equivalent for U.S. Radiation Worker .....	19
5.1 Impacts on Medical/Dental and Veterinary Practice .....	23
5.2 1989 Exposure Experience .....	24
5.3 Impacts in Nuclear Power Reactors .....	25
5.4 1989 Exposure Experience .....	26
5.5 1993 Exposure Experience (estimated) .....	26
5.6 Impacts in Nuclear Power Reactor Contractors .....	29
5.7 1989 Exposure Experience .....	30
5.8 1993 Exposure Experience .....	30
5.9 Impacts in Test and Measurements Including Industrial Radiography .....	31
5.10 1989 Exposure Experience .....	32
5.11 1993 Exposure Experience .....	32
5.12 Impacts in Universities .....	33
5.13 1989 Exposure Experience .....	33
5.14 Impacts in Manufacturing and Distribution .....	34
5.15 1989 Exposure Experience .....	35
5.16 1993 Exposure Experience .....	35
5.17 Impacts in Waste Management .....	36
5.18 1989 Exposure Experience .....	37
5.19 Impacts in Fuel Fabrication, UF <sub>6</sub> Production .....	37
5.20 1989 Exposure Experience .....	38
5.21 Impacts in Well Logging .....	39
5.22 1989 Exposure Experience .....	39
5.23 Impacts in Others (R&D, Regulatory) .....	40
5.24 1989 Exposure Experience .....	40
6.1 Whole-Body Dose Data for PWR Plants for 1988 .....	41
6.2 Whole-Body Dose Data for BWR Plants for 1988 .....	42
6.3 Whole-Body Dose Data for Various Worker Groups at PWR Plants for 1988 .....	43
6.4 Whole-Body Dose Data for Various Worker Groups at BWR Plants for 1988 .....	43
7.1 Estimated Costs and Dose Savings for Modifications at Nuclear Power Plants .....	46

## Executive Summary

The revised Nuclear Regulatory Commission (NRC) regulations 10 CFR 20 were based largely on the 1977 recommendations of the International Commission on Radiation Protection (ICRP), as interpreted and promulgated by the Environmental Protection Agency (EPA) in 1987. Since then, the United Nations Scientific Committee on the Effects of Atomic Radiation (UNSCEAR), the National Research Council Committee on the Biological Effects of Ionizing Radiation (BEIR), and the International Commission on Radiological Protection (ICRP) have published new information indicating that the risk associated with exposure to ionizing radiation is somewhat greater than that used by the ICRP and others in 1977. This increase reflects additional cancers found in the Japanese survivors of the atomic bombings, new dosimetry, and the adoption of a projection model which accounts for the excess cancer cases that are expected to occur in those survivors who are still alive.

The ICRP recommended a dose limit of 100 mSv in 5 years (10 rem in five years) in its 1990 recommendations. The National Council on Radiation Protection and Measurements (NCRP) in 1987 recommended an annual limit of 50 mSv yr<sup>-1</sup> (5 rem yr<sup>-1</sup>) and suggested that no individual should exceed a cumulative dose equal to his/her age in 10s of mSv (age in rem). This suggestion has been raised to the level of a recommendation in the 1993 Recommendations of the NCRP. Many countries in the world are drafting new regulations adopting the ICRP system.

This study was requested by NRC to obtain a preliminary estimate of the potential impacts to NRC licensees of any reduction in the dose limits. In general, the past in-depth reviews of the impact of lowering dose limits were based on an assumption that there would be no reduction in the source terms, no improvement in equipment (remote tooling and surveillance), nor any increase in the productivity of radiation workers.

Four approaches were used in this study. The first was the development and distribution of a questionnaire designed to solicit and evaluate information on the potential impacts of decreased dose limits from a wide variety of licensees. The second approach was the review and analysis of previous surveys on dose impacts and other data collections. These surveys were conducted by the Edison Electric Institute (EEI) Health Physics Committee, the Department of Energy (DOE), Office of Health and Safety, and the Brookhaven National Laboratory (BNL) ALARA Center. The data collections are those of the NRC Radiation Exposure Information Reporting System (REIRS) and Environmental Protection Agency (EPA) 1984 Report on Occupational Exposure.

The third approach was to use a working committee to validate and extend the data obtained from the questionnaire, and also review and comment on this report. This committee was composed of radiation protection experts from various sectors of NRC licensees, together with individuals from Nuclear Management and Resources Council (NUMARC), DOE, NRC, and the BNL ALARA Center. The fourth approach was to incorporate the comments made to the draft NUREG/CR-6112.

Where possible, the data for 1989 was used as the basis for this report to allow meaningful intercomparisons. The BNL High Dose Group Study was based on 1988 data, and the EPA Report was based on data of 1984 and earlier. Although the data for 1990 and 1993 suggests a reduction in individual and collective dose has taken place, the overall conclusions drawn from the 1989 study remain valid.

Examples of costs associated with reducing the source term in nuclear power plants were obtained from the NUREG/CR-4373, "Compendium of Cost-Effectiveness Evaluations of Modifications for dose Reduction at Nuclear Power Plants," (Baum and Matthews, 1985).

From the information given in this report and that offered by the working committee, several tentative conclusions can be drawn.

The analysis suggests there would be minimal impact on collective doses, on costs of modifying facilities, or on annual radiation-protection costs under the combined limit of 50 mSv yr<sup>-1</sup> (5 rem yr<sup>-1</sup>) and cumulative dose in 10s of mSv (rem) equal to age in years. The lifetime risk associated with this limit - to an individual maximally exposed - would be slightly less than that incurred by a similar individual controlled by the ICRP's limit of 100 mSv in 5 years (10 rem in 5 years). However, a "grandfather clause" allowing up to 20 mSv yr<sup>-1</sup> (2 rem yr<sup>-1</sup>) after exceeding the age limit will be required for perhaps less than 1000 workers.

A 20 mSv yr<sup>-1</sup> (2 rem yr<sup>-1</sup>) limit would appear achievable, although some tasks, particularly those in medicine and in certain parts of the nuclear power industry, might prove extremely difficult to maintain. Extensive modifications, such as steam generation, maintenance, and refueling including the installation and use of robots and partial/full system decontamination, would be required for many tasks in nuclear power plants. Depending upon the extent of the modifications, the collective dose might go up or down. That is, extensive use of robots, source term reductions, and facility modifications might lower collective doses. Less ambitious modifications, less decontamination, and the use of fewer robots might keep the collective doses at about the same level while reducing individual doses; making no changes and allowing the same tasks to be performed would necessarily result in higher collective doses. The working committee suggested that with this annual limit, there could be a potential impact on safety since some discretionary inspection and maintenance might be constrained.

For a 10 mSv yr<sup>-1</sup> (1 rem yr<sup>-1</sup>) limit, the risk to the most highly exposed individual would be lower than for other options, i.e. equivalent to that of fatal accidents in United States industries, but the impacts are expected to be quite serious for many of the industries which responded to the questionnaire. There are tasks, again in medicine, which under present procedures could be prohibitively expensive. For industries with large source terms, facility modifications and radiation protection costs would be extremely large (see Section 7). For these reasons, collective dose may increase substantially.

One additional issue must be kept in mind when assessing the impact of lower dose limits. That is, for licensees to ensure that doses do not exceed the regulated dose limits, they routinely use administrative limits. For example, with a regulatory limit of a 50 mSv yr<sup>-1</sup> (5 rem yr<sup>-1</sup>), an administrative limit of a 40 mSv yr<sup>-1</sup> (4 rem yr<sup>-1</sup>) might be used. At 20 mSv yr<sup>-1</sup> (2 rem yr<sup>-1</sup>) limit, a 15 mSv yr<sup>-1</sup> (1.5 rem yr<sup>-1</sup>) administrative limit might be used, and so on.

## Foreword

On May 21, 1991, the Nuclear Regulatory Commission (NRC) published a revision to 10 CFR Part 20, "Standards for Protection Against Radiation." The rule went into effect in June 1991, and all licensees were required to implement the regulations on or before January 1, 1994.

The revised 10 CFR Part 20 is based on the recommendations of the International Commission on Radiological Protection (ICRP) in Publication 26 (ICRP 1977). In 1991, ICRP published revised recommendations in Publication 60. These recommendations were based upon revised dosimetry and epidemiology, including the information presented in reports such as the 1988 United Nations Scientific Committee on the Effects of Atomic Radiation (UNSCEAR). In this revision, ICRP reduced its recommended dose limit to 100 mSv (10 rem) in 5 years, with the additional limitation that no more than 50 mSv (5 rem) is received in any one year.


In 1991, the National Council on Radiation Protection and Measurements (NCRP) recommended a lifetime limit of 10 mSv (1 rem) times age in years (NCRP 91). NCRP is currently reexamining its recommendations based on ICRP 60.

As a result of these recommendations, in 1988, the NRC staff initiated a study by Brookhaven National Laboratory to analyze the potential impacts of reduced dose limits on its licensees and to provide a technical base upon which to base future regulatory decisions. This NUREG summarizes the results of that study, which included a survey of radiation protection experts. Even though the information presented is not complete for certain categories of licensees due to unavailability, the conclusions for those where data was available are considered valid.

In view of the small number of licensees who responded to the survey, the NRC staff decided to publish a draft of this report for public comment in the hopes that additional data and expert opinion would result, so that a more extensive technical base could be developed. Licensees, agreement states, and all other interested parties were encouraged to submit comments and relevant data on this draft report.

This NUREG incorporates the information from the comments received.

NUREG/CR-6112 is not a substitute for NRC regulations, and compliance is not required. The approaches and/or methods described in this NUREG/CR are provided for information only. Publication of the report does not necessarily constitute NRC approval or agreement with the information cited therein.



John E. Glenn, Chief  
Radiation Protection & Health Effects Branch  
Division of Regulatory Applications  
Office of Nuclear Regulatory Research

## **Acknowledgments**

This report is the result of efforts of many individuals. In addition to the author, the staff of the Brookhaven National Laboratory (BNL) ALARA Center, John Baum, Tasneem Khan, Bruce Dionne, and Casper Sun made major contributions to the report.

The working committee of Larry Brennecke, Thomas McLeod, Thomas Gaines, George O'Bannion, Howard Elson, Frank Rescek, Frank Roddy, Robert Robinson, Alan Roecklein, Anthony Weadock, George Powers, Ralph Andersen, Jay Maisler, Tasneem Khan, and Bruce Dionne provided data, insightful comments and suggestions, and helpful editorial suggestions.

Alan Roecklein and the Project Manager, George Powers, gave us the necessary oversight, advice, and support required to complete this phase of the work.

Finally, the patience and precision of Karen Wagner in preparing and editing the report is gratefully acknowledged.

# Impact of Reduced Dose Limits on NRC Licensed Activities

## 1 Introduction

The revised Nuclear Regulatory Commission (NRC) regulations, 10 CFR 20, (NRC, 1991) impose an annual effective dose equivalent limit of 50 mSv (5 rem) on occupationally exposed workers. This requirement corresponds to that given in the Environmental Protection Agency's (EPA's) 1987 Radiation Protection Guidance for Occupational Exposure-Recommendations (EPA, 1987) approved by the President. Both of these agencies based their requirements largely on the 1977 recommendations of the International Commission on Radiological Protection (ICRP) given in their Publication 26 (ICRP, 1977).

In the late 1980s, the Radiation Effects Research Foundation (RERF) updated the data on their life-span study of the Japanese atomic bomb survivors to account for the increase in cancer incidence as a function of dose associated with a revision in the dosimetry (Shimizu et al., 1987; 1988). Another increase in the risk factors resulted from a potential increase in the risk associated with further epidemiological support for the multiplicative or relative risk projection model. The National Council on Radiation Protection and Measurements (NCRP) modified their basic recommendations to reflect this preliminary data in 1987 (NCRP, 1987). The NCRP also noted the substantial decrease in the frequency of fatal industrial accidents that had been the basis for the risk-based dose limit given by ICRP in 1977. This frequency decreased from about  $1 \times 10^{-4} \text{ yr}^{-1}$  in the early 1970s to about  $4 \times 10^{-5}$  in the mid 1980s.

Shortly thereafter, the United Nations Scientific Committee on the Effects of Atomic Radiation (UNSCEAR) and the National Research Council Committee on the Biological Effects of Ionizing Radiation (BEIR) produced the 1988 UNSCEAR Report (UNSCEAR, 1988) and the 1990 BEIR V Report, (NASBEIR, 1990) respectively.

Using the preliminary information from the 1988 UNSCEAR report, the ICRP began a major revision to its recommendations, beginning with a detailed review of the data. The revised estimate of the lifetime fatal cancer risk for low dose or low dose-rate exposure given in ICRP Publication 60 is  $\sim 4 \times 10^{-2} \text{ Sv}^{-1}$  ( $\sim 4 \times$

$10^4 \text{ rem}^{-1}$ ) for adults, and about  $5 \times 10^{-2} \text{ Sv}^{-1}$  ( $5 \times 10^{-4} \text{ rem}^{-1}$ ) for the total population (ICRP, 1991). Although the ICRP has changed its criteria for selecting dose limits, this increased estimate of the risk of fatal cancer alone from 1.25 to  $4 \times 10^{-2} \text{ Sv}^{-1}$  ( $1.25$  to  $4 \times 10^{-4} \text{ rem}^{-1}$ ) given in ICRP Publication 26 (ICRP, 1977) suggested that an annual limit of 50 mSv (5.0 rem) over a working lifetime was unlikely to be considered acceptable. Their solution, given in Publication 60, was to recommend an occupational limit of 100 mSv in 5 years ( $20 \text{ mSv yr}^{-1}$ ) [ $10 \text{ rem in 5 years (2 rem yr}^{-1})$ ] with an additional limit of 50 mSv (5 rem) in any year.

The International Atomic Energy Agency (IAEA) and the Commission of European Communities (CEC) already have begun to revise their basic safety standards to conform with ICRP's new recommendations.

In light of these developments, in 1988 the NRC requested that a preliminary study be made to analyze the potential impacts of reduced dose limits on its licensees, and to provide a technical base for making future regulatory decisions on limits. This report summarizes the results of a review on the impact of reduced dose limits to NRC licensees.

## 2 Historical Background and Literature Survey

### 2.1 1928 to 1977

The first widely accepted dose-limiting recommendations were based on keeping exposures below the threshold for observable effects (Mutscheller, 1925). By the end of the second world war, these limits, which by then reflected concern over leukemia and genetic effects, were expressed as 300 mrem/week to tissues at a depth of 5 cm or more in the body, and 600 mrem/week to the surface of the body (NBS, 1954; ICRP, 1954). These values were equivalent to the later limits of 15 rem yr<sup>-1</sup> to most of the individual organs with the exception of the blood-forming tissues, the gonads, and the lens of the eye (NCRP, 1971; ICRP, 1959a), and 30 rem yr<sup>-1</sup> to the skin (NRC, 1960; ICRP, 1964).

After the second world war, there was much public concern over world-wide fallout from nuclear tests (Divine, 1978). Mueller and others were convinced that for genetic effects at least, there was a linear no-threshold response (Mueller, 1927; Lea, 1947). The National Academy of Sciences-National Research Council (NAS-NRC, 1956) and the British Medical Research Council (MRC, UK, 1956) formed expert committees to examine the radiobiological evidence. The basic consideration was the need to restrict the genetic damage to both exposed individuals and to the general population. Based heavily on the dose-effect relationship for genetic effects seen in *Drosophila* and on the observed genetic burden seen in humans, assumed to be partly due to the natural radiation background (Haldane, 1948), the next set of limits reflected: 1) a need to limit cumulative dose, and 2) a need to restrict the cumulative dose to workers in their reproductive years below that for older workers. The resulting limits for whole-body penetrating radiation were (age - 18) 5 rem cumulative dose and 3 rem/quarter (NCRP, 1957; ICRP, 1959b).

By the early 60s, the data from the Japanese survivors of the atomic bombs began to emerge (UNSCEAR, 1962). This data, together with that from the early radiologists and British spondylitic patients, suggested that the incidence of leukemia increased as a result of radiation.

A decade later, it was apparent that the incidence of certain solid tumors also increased in the Japanese survivors, the British spondylitic patients, and women

with mastitis who had been treated with X rays (UNSCEAR, 1972; BEIR, 1972).

Consequently, the International Commission on Radiological Protection (ICRP), the National Council on Radiation Protection and Measurements (NCRP), and the Federal Radiation Council (FRC) all re-emphasized the need to keep exposure as low as practical, practicable, or reasonably achievable.

### 2.2 1977 to 1987

In the middle 70s, the United Nations Scientific Committee on the Effects of Atomic Radiation (UNSCEAR, 1977) felt there was sufficient information from the Japanese atomic bomb survivors to estimate the risks to individual organs. This led to the adoption by the ICRP in 1977 of the effective dose equivalent concept<sup>1</sup>, with its attendant w<sub>T</sub> values (weighting factors representing the proportion of the stochastic risk from individual tissues relative to the risk to the whole body when the body is irradiated uniformly). In addition, the ICRP "justified" the 50 mSv yr<sup>-1</sup> (5 rem yr<sup>-1</sup>) limit on the basis that the average dose would be less than 10 mSv yr<sup>-1</sup> (1 rem yr<sup>-1</sup>) and, as UNSCEAR had done, assumed that the risk from low dose, low dose-rate exposure was 2.5 times less than that seen in Japanese atomic bomb survivors. The first of these two criteria led ICRP in 1977 to eliminate the (age - 18) 5 rem recommendation.

Perhaps the greatest significance of the 1977 ICRP Publication 26 was the development of the close relationship between risk and dose limits. Simply put, an average excess risk of fatal cancer and severe genetic effects of 1 x 10<sup>-2</sup> Sv<sup>-1</sup> (1 x 10<sup>-4</sup> rem<sup>-1</sup>) was judged to be "acceptable" by the ICRP.

At the time that ICRP published their recommended occupational limit of 50 mSv yr<sup>-1</sup> (5 rem yr<sup>-1</sup>) (ICRP, 1977), several different sets of limits were being recommended or used in the United States.

<sup>1</sup> The concept originated in ICRP Publication 26 (ICRP, 1977) although the term "effective dose equivalent" was not introduced until 1978 (statement from the 1978 Stockholm Meeting of the ICRP Annals of the ICRP, Vol. 2, No. 1. (1978).

The NCRP was recommending a limit of 5 rem yr<sup>-1</sup> and (age - 18)5 rem (NCRP, 1987); the Federal Radiation Council (FRC) was recommending 3 rem/quarter and (age-18) 5 rem (FRC, 1960); both the Nuclear Regulatory Commission (NRC) and the Occupational Safety and Health Administration (OSHA) were enforcing 3 rem/quarter and (age -18) 5 rem, and the Department of Energy (DOE) were enforcing 3 rem/quarter and 5 rem yr<sup>-1</sup>. During this period, the Natural Resources Defense Council (NRDC) petitioned both the EPA and the NRC to lower occupational exposure limits in the United States. The federal agencies' response to the petition eventually led to several reports on the impact of lowering the Annual Dose Equivalent limit from 5 rem to 0.5 rem.

The earliest report was prepared for Stone and Webster Engineering Corporation by Warman et al., 1978. Their basic conclusion was that a decrease in the dose limit to about 2 rem yr<sup>-1</sup> would exponentially increase both collective dose and the number of additional workers needed. Below 2 rem yr<sup>-1</sup>, the increase per unit dose reduction would be even greater. These results were based on the dose distribution of Pressurized Water Reactor (PWR) and Boiling Water Reactor (BWR) workers in 1976. The basic assumptions were that the dose received by workers that was above any new dose limit would have to be received by additional workers, and that the dose rates existing at the power plants at that time would be representative of future dose rates. All workers were assumed to be productive 90% of the time.

A more detailed analysis was made by the Atomic Industrial Forum (AIF) a few years later in which the impacts were analyzed by tasks (AIF, 1978). The overall conclusion, taken from a statement in the report, was "While exposure and costs do increase, manpower is considered the most significant concern." Again, it is important to recognize that AIF assumed (1) "that there will be no significant design improvements made leading to the reduction of exposure or to improved operation or maintenance", and (2) "that work in a radiation environment at commercial nuclear power plants will not be performed significantly differently at lowered exposure limits than it is at present limits."

The DOE conducted a similar study for their facilities (DOE, 1979). Rather than employ the models used in the AIF study, DOE relied on a detailed questionnaire and a review committee. However, their conclusions

were no different than those of the two reports discussed above, except that the impacts occurred at slightly lower doses because DOE was then using a 5 rem yr<sup>-1</sup> limit. The DOE report recommended that the concept of As Low As Reasonably Achievable (ALARA) should have greater attention than a reduction in dose limit. Also, there was more emphasis on potential facility modifications and reduction of source terms.

Fortunately, since these reports were issued, extraordinary strides in reducing exposure using the ALARA principle and restrictive administrative limits have significantly reduced collective dose without increasing the average annual dose to workers. In fact, the combination of improvements in productivity, design, and source-term reduction has decreased the average individual dose at both NRC licensees and DOE facilities over the past decade. This was most clearly demonstrated in the Naval Nuclear Propulsion Program (Schmitt and Brice, 1984), and in the commercial nuclear power industry (Brooks, 1988).

### 2.3 1987 to 1994

Today, the weight of new radiobiological evidence on dose limits is as important as it was in the early 1950s. The incorporation of (age -18) 5 rem into the recommendations and limits at that time was accepted with little difficulty (except, perhaps, in uranium mining and fuel fabrication). The most recent evidence from the Japanese survivors, reviewed by UNSCEAR (UNSCEAR, 1988) and the National Academy of Sciences (NAS) Committee on the Biological Effects of Ionizing Radiation (BEIR, 1990), suggests that the risks of fatal cancer and severe genetic effects may be up to 4 times greater than those estimated in 1977.

Reacting to the emerging information from the Radiation Effects Research Foundation (RERF) in Japan (Preston and Pierce, 1981), the ICRP issued a statement in 1987 following its meeting in Como, Italy (ICRP, 1987). The Commission suggested that: (1) revised dosimetry could increase the cancer risk/unit dose by a factor of 1.4, (2) the observed increase in the incidence of solid tumors in "younger" members of the exposed population might lead to a combined increase of a factor of 2, and (3) the relative risk projection model could increase the risk factor even further. The Commission also noted that a new

## Historical Background

set of basic recommendations would be available in 1990.

Most workers seem to have been adequately protected under the (age - 18) 5 rem dose limit. The average annual exposure to monitored workers with measurable exposure was about 230 mrem (EPA, 1984). Using the 1990 ICRP risk estimates of  $5 \times 10^{-2} \text{ Sv}^{-1}$  ( $\sim 4 \times 10^{-4} \text{ rem}^{-1}$ ) for fatal cancer for those aged 18-65, the lifetime of fatal cancer risk to an individual receiving the annual exposure of 2.3 mSv (230 mrem) is predicted to be  $\sim 1 \times 10^{-4}$ . This figure is comparable to the risk of accidental death in U.S. industry.

However, for a worker receiving 50 mSv (5 rem) in one year, these same risk estimates project a lifetime risk of attributable fatal cancer and severe genetic effects at  $2.5 \times 10^{-3}$ . Such an annual level of risk is comparable to that associated with the upper range of risk in mining, construction, and agriculture, including deep-sea fishing. For those few workers who may receive annual doses near the dose limit over much of their working lives, the cumulative level of risk may be unacceptable.

Consequently, the National Radiation Protection Board (NRPB) in England issued interim guidance in November 1987 (NRPB, 1987) in which they recommended that "... occupational workers exposure should be so controlled as not to exceed an average effective dose equivalent of 15 mSv yr<sup>-1</sup>."

This NRPB Guidance is, in fact, quite similar to the 1987 recommendation of the NCRP in its Report 91 (NCRP, 1987) in which the Council stated "...the community of radiation users is encouraged to control their operations in the workplace in such a manner as to ensure, in effect, that the numerical value of the individual worker's lifetime effective dose equivalent in tens of mSv (rem) does not exceed the value of his or her age in years." Both approaches would lead to lifetime doses below 750 mSv (75 rem).

Both guidances reflected an expectation that risk estimates would increase and safe industries would continue to become safer.

In general agreement with other countries, the Federal Republic of Germany stated that before changing annual dose limits it will await completion of international discussion following the issuance of the 1990 ICRP recommendations. However, the German au-

thorities made a rather dramatic change in their recommendations (Kaul et al., 1989):

"Under the present conditions, the German Commission on Radiological Protection (SSK) recommends that the rule of minimization be applied more strictly and that in the future, in adherence to the annual dose limit of the Radiological Protection Ordinance of 50 mSv, a total dose of 400 mSv during a whole working lifetime shall not be exceeded (occupational lifetime dose)."

A comprehensive report on the impacts of dose-limit reduction was produced in 1988 for the Electrical Power Research Institute (EPRI) (Le Surf, 1988). The author suggested that although there have been significant reductions in both individual and collective doses in the U. S. nuclear power industry, basic and fundamental changes are needed if this industry is to comply with lower limits. He points out that other countries have successfully reduced exposure in three ways: first, by changing the philosophy of radiation protection, emphasizing line responsibility and training; second, by introducing aggressive measures to reduce the source term; and third, by incorporating similar approaches to prevent the buildup of radiation fields. The NRC established an ALARA Center at Brookhaven National Laboratory (BNL), which maintains a database for these issues (Khan et al., 1992; Baum and Khan, 1992; Khan et al., 1991b).

In January 1991, the ICRP issued its Publication 60, "The 1990 Recommendations of the International Commission on Radiological Protection" (ICRP, 1991) recommending a limit of 100 mSv in 5 years, with the caveat that no more than 50 mSv be allowed in any one year. The Commission's intention was to limit the lifetime effective dose to  $\sim 1 \text{ Sv}$  (100 rem) and the average annual effective dose equivalent to 20 mSv (2 rem).

The most recent NCRP recommendations given in its Report 116, "Limitation of Exposure to Ionizing Radiation," raise the guidance given in NCRP Report 91, "Recommendation on Limits for Exposure to Ionizing Radiation," on a lifetime dose in 10s of mSv equal to age in years (lifetime dose in rem equal to age in years) to the level of a recommendation. The NCRP Report 116 also maintains the recommendation of 50 mSv yr<sup>-1</sup> (5 rem yr<sup>-1</sup>).

It is important to know that UNSCEAR reaffirmed the ICRP risk estimate in both their 1993 (UNSCEAR, 1993) and 1994 (UNSCEAR, 1994) reports.

The IAEA has revised of the Basic Safety Standards as has the CEC. The European Community is expected to have a new set of requirements based on ICRP 60 in place by the middle of this decade, with many other nations following soon after.

## **2.4 Background Summary**

In general, past in-depth reviews of the impact of lowering dose limits were based on an assumption that there would be no reduction in the source terms, no improvement in equipment (remote tooling and surveillance), nor any increase in the productivity of radiation workers. However, reductions in dose limits led to the realization that all of these assumptions may be incorrect. It is essential that any review of the impact of lowering dose limits addresses the financial impact of lowering collective doses, not simply the redistribution of existing exposure.

## 3 Data Gathering

In this study we proposed to use existing surveys and to obtain opinions on the impacts of reductions in the dose limit from as broad a spectrum of users as possible without resorting to an intensive site-by-site assessment. In addition to reviewing such surveys, such as the EEI, the DOE, and recent NRC-sponsored studies on dose reduction, there was a widespread distribution of a questionnaire to elicit the responders' opinion and to obtain specific data to assist in our overall assessment of the impact. Data from the NRC's Radiation Exposure Information and Reporting System (REIRS) and the 1984 EPA Report on Occupational Exposure were used to validate the survey data. In addition, data and information supplied in the comments received on the draft NUREG/CR-6112 will be reviewed.

### 3.1 Existing Surveys

#### 3.1.1 1992 Edison Electric Institute (EEI) Report on Dose Limits and Guidelines

Questionnaires were sent to all members of the EEI Health Physics Committee addressing the following topics: 1) current practices and experience on administrative dose-control levels, 2) cumulative dose guidelines and experience, 3) projected impacts associated with lifetime dose limits, and 4) effects of a reduced annual dose limit and of establishing a cumulative dose limit. Twenty-seven individuals replied, representing 23 nuclear utilities. These responses covered 43 Pressurized Water Reactors, 18 Boiling Water Reactors, and a High Temperature Gas Cooled Reactor, encompassing more than half the nuclear power plants (62 out of 108 units in 1989). They obtained dose data for > 14,500 and > 12,500 individuals with doses > 500 mrem in 1985 and 1989, respectively. For these two years, the number of personnel at U. S. power reactors with doses > 500 mrem was about 27,000 and 25,000, respectively.

The responses were stored in a computer database and published as graphs and tables, with the authors of the report using their best judgment to interpret the utilities' responses. The full survey is reported in the EEI Nuclear Report, "Utility Response to Questionnaire on Dose Limits" (EEI, 1991); Section 4.1 gives a brief summary.

#### 3.1.2 Department of Energy Report (DOE) on the Implications of the BEIR V Report

In response to a request by the Secretary of Energy, the Office of Health reviewed the implications of the BEIR V report for the Department of Energy (DOE). A questionnaire was developed by a DOE Internal Review Committee to survey DOE contractors to estimate costs for additional personnel, programmatic upgrades, and engineering modifications that would be needed to comply with an anticipated reduction in the dose limits.

The questionnaire was sent to the Albuquerque, Chicago, Idaho, Nevada, Oak Ridge, Richland, San Francisco, and Savannah River Field Offices on January 30, 1990, for distribution to their contractors. Thirty-seven contractor sites responded, which operate the following types of nuclear facilities: accelerators, fuel/uranium enrichment, fuel fabrication, fuel processing, maintenance and support, hot cells, reactors (test, research, and production types), research and development, fusion, waste processing/storage, weapons fabrication and testing, tritium production, and radiography. Two significant contributors to DOE's collective dose, the Rocky Flats plant in Golden, Colorado, and Los Alamos National Laboratory in Los Alamos, New Mexico did not respond.

The scope and findings of the survey are given in the "Final Report to the Secretary of Energy; Implication of the BEIR V Report to the Department of Energy" (DOE, 1990). The results are summarized in Section 4.2 of this report.

#### 3.1.3 Nuclear Regulatory Commission (NRC) Radiation Exposure Information and Reporting System (REIRS)

The NRC established a radiation exposure information and reporting system (REIRS) and publishes data from six of the seven categories of NRC licensees subject to the reporting requirements of 10 CFR 20.407. Selected data from NUREG 0713 Vol 12 (Raddatz and Hagemeyer, 1993), which presents data for 1990, are given in Section 4.3 of this report; it serves as one element of the process of ensuring that the survey responses provide a realistic picture of the

exposure statistics. It should be noted that the REIRS data contains information from NRC licensees only. Companies that are licensed by agreement states do not report their exposures to the NRC, so the data for industrial radiography, manufacturing, and distribution of specified quantities of by-products and low-level waste do not reflect the total United States exposure.

### 3.1.4 Environmental Protection Agency (EPA) Report on Occupational Exposure to Ionizing Radiation in the United States

Because the U. S. nuclear industry is spread over many diverse sectors, it is very difficult to get a complete, comprehensive picture of the radiation exposure of all radiation workers. Fortunately, the Environmental Protection Agency (EPA) made a study which covers almost every sector (EPA, 1984). We analyzed their data to gain detailed information for one year over the entire U. S. nuclear industry. Although the study is several years old, it is by far the most detailed of its kind and its main conclusions are useful to the current effort. Section 4 presents our analysis.

## 3.2 Survey Performed for this Report

A questionnaire designed to elicit response from a wide variety of radiation users was developed (Appendix A).

A working group of technical experts (see 3.2.3) reviewed the data from the questionnaire and obtained additional data where needed.

### 3.2.1 Questionnaire Design

Three classes of information were judged to be important: The responders' estimate of the impact as a function of several dose limiting options; their organization's preliminary data on exposures; and lastly, their comments and suggestions. The questionnaire also solicited information about the respondent's organization and asked if the respondent could become a member of a working group to review and assess the results of the questionnaire.

### 3.2.1.1 Options for Potential Dose Limits

Four dose-limit options were proposed, each reflecting a rational response to the new risk estimates. The first option considered was  $2 \text{ rem yr}^{-1}$ , which was the basic recommendation in the widely circulated draft of the ICRP revision to its Publication 26 (the final recommendation was 100 mSv in five years, and less than 50 mSv in any one year).

The second option was  $1 \text{ rem yr}^{-1}$ , based on the UNSCEAR 1988 risk estimate being about 4 times the UNSCEAR 1977 risk estimate. Therefore, it might be prudent to reduce the  $5 \text{ rem yr}^{-1}$  limit to about  $1 \text{ rem yr}^{-1}$  to account for this difference. In addition, the age-related approach suggested in NCRP 91 could result in  $1 \text{ rem yr}^{-1}$  if the regulatory agency is concerned about the record-keeping of cumulative dose limits. Furthermore, perhaps this is the lowest level that could be imposed and still permit widespread use of radiation and radioactive materials.

The third option was age in rem and  $5 \text{ rem yr}^{-1}$ , which simply escalates the "guidance" given in NCRP Report 91 to a regulatory limit. It allows up to  $5 \text{ rem yr}^{-1}$  which permits the continued operation of previously designed facilities without significant modifications, but ensures that the lifetime risk to any individual will be less than 100 rem.

Fourth, a limit of age in rem and  $2 \text{ rem yr}^{-1}$  was given because a regulatory agency may want to regulate the rate of exposure more closely than option 3. In addition, this limit option appears to be closer to the ICRP's recommended limit of 100 mSv in five years, and has the advantage of restricting exposure in the early years of working life more than does option 3.

These four options are not intended as suggestions for new regulatory limits, but merely as the most probable ones which a regulator might consider.

### 3.2.1.2 Impacts of Reduced Dose Limits

Previous studies on the impacts of reduced dose limits usually cite increased costs and increased collective dose. The questionnaire asked that costs be broken down between those required for modifying the facility, and operating costs. The first are expected to be one-time costs, and the latter recurring costs.

## Data Gathering

### 3.2.1.3 1989 Dose Experience

To allow BNL to make a less subjective assessment, six items of related data were requested. The first three were the number of employees with exposure in excess of 5 rem, 2 rem, and 1 rem in 1989, data clearly related to the potential limits given in the options. The fourth item was a request for information on the number of employees whose current lifetime dose in rem exceeds their age, which would highlight any need for "grandfathering". The number of employees with measurable dose was requested to judge the weight that should be given to the specific data in the questionnaire. The annual collective dose also was requested, which, when taken with the above data, could provide information on the dose distribution, and assist in evaluating the answers about the impact on collective dose.

### 3.2.2 Questionnaire Distribution

The questionnaire and an explanation of its intended use was published in the July 1990 issue of the Health Physics Society Newsletter, which is distributed to the nearly 6,000 members of the society. The society is composed of scientists, engineers, and professionals concerned with radiation protection throughout the United States, so it was felt that virtually all categories of radiation users would have access to it. A letter describing the questionnaire and its availability was published in the newsletter of the American Association of Physicists in Medicine. The majority of medical physicists and medical health physicists belong to this society, so this category of radiation users was given a unique opportunity to participate.

### 3.2.3 Working Committee on the Impact of Reduced Dose Limits

From the inception of this study, we recognized that the questionnaire alone could not ensure that all occupational exposure practices were adequately assessed. In addition, the questionnaire might elicit subjective information which, while helpful, could lead to misinterpretation of the actual impact, particularly where there were few responses from a particular industry or practice. Therefore, a working committee was assembled composed of individuals with experience and knowledge in radiation protection from a wide variety of industries and practices. The membership included: from medical activities (Larry Brennecke and Thomas McLeod); from industrial

radiography (Thomas M. Gaines); from well logging (George O'Bannion); from the university community (Howard K. Elson); from nuclear power plants (Frank Rescek); from nuclear plant contractors (Frank Roddy); from fuel fabricators (Robert Robinson); from NUMARC (Ralph Andersen and Jay Maisler); from the Nuclear Regulatory Commission (George Powers and Alan Roecklein); and from the Department of Energy (Anthony Weadock). Bruce Dionne and Tasneem Khan of the BNL ALARA Center also participated.

The working committee met on March 27, 1991 to review the data from the questionnaires. They also reviewed the study by the DOE on the implications of BEIR V to the DOE, and the BNL ALARA Center study on high-dose worker groups at nuclear plants (both are discussed elsewhere in this report). Additional data was received from the participants during the meeting, and areas requiring more information were identified.

After this meeting, questionnaires were mailed to additional radiographers, fuel-fabrication workers, and nuclear-plant contractors.

A letter in the October 1991 issue of the Health Physics Society Newsletter summarized the information from the responses received up to that point. This letter specifically requested comments and suggestions. Because there were no responses, a follow-up letter was published in the March 1992 issue. Only two responses were received by the end of May.

A second meeting of the working group was held in July 1992 when several specific comments and suggestions were made (see Chapter 5).

## 3.3 Comments Received to the Draft NUREG Report

The NUREG/CR-6112 Draft Report was distributed for public comment in early 1994. Seven sets of comments were received. This additional information included comments and data from the nuclear power plant community, the fuel fabrication community, the radio-pharmaceutical production community, and the industrial radiography community. The comments have been addressed in Section 4.5, and much of the data and detailed comments are included in Section 5.

## 4 Survey Results

### 4.1 Edison Electric Institute (EEI) Report

#### 4.1.1 Administrative Control Levels

In the EEI Report (EEI, 1991) twenty-seven people reported administrative control levels: six use a 5 rem annual "limit"; eight have adopted a 4.5 to 4.9 rem yr<sup>-1</sup> value; eleven use an annual control level of approximately 4 rem yr<sup>-1</sup> (which was the guideline published by the Institute of Nuclear Power Operations (INPO) in 1988); and two have adopted progressive levels of 2.5 rem yr<sup>-1</sup>.

#### 4.1.2 Annual Reported Doses for 1985 and 1989

Figure 4.1 (taken from the EEI Report) shows the number of workers from 11 sites with annual doses greater than 0.5, 1.0, 2.0, 3.0, 4.0, and 5.0 rem in 1985 and 1989. The data include both utility personnel (UT) and total personnel (TO) which includes contractors. Figure 4.2 (also from the same report) shows the percentage of utility personnel and total personnel with annual doses greater than these dose values for the same two years. The contractor doses are only those reported by the individual utilities and may not reflect their total dose (i.e. the sum of doses received at two or more sites).




Figure 4.1 Annual Site Doses for Utility Personnel (UT) and Total Plant Workers (TO) (11 Responders)




Figure 4.2 Annual Site Doses for Utility Personnel (UT) and Total Plant Workers (TO) for 1985 and 1989

There is a clear decrease in the number and percentage of both utility and total personnel above each dose value in 1989 relative to 1985. No person exceeded 5 rem yr<sup>-1</sup>. About 8% (967) of the people at 11 sites had annual doses greater than 2 rem yr<sup>-1</sup> in 1989.


#### 4.1.3 Cumulative Dose Administrative Guidelines

The survey showed that 13 of the 26 responders had established some form of a cumulative dose guideline, the most common being age times 1 rem. Four have a review or reference level based on age, or a cumulative lifetime value, for which individual doses would be tracked and intervention would occur. Ten responders had not established a cumulative guideline in 1989 but most were in the process of adopting one. We noted that seven responders had adopted a cumulative-dose exemption procedure to exceed, which typically required the approval of a Vice President, Director, or Plant Manager. The report stated that "...it is likely that in a few years most nuclear utilities will have in place some form of lifetime or cumulative dose guidance". In its December 1991 guidelines, INPO urged utilities to strive to meet the NCRP recommendation of a lifetime dose not to exceed the workers age in rem.

## Survey Results

### 4.1.4 Cumulative Reported Doses for 1989

Figure 4.3 (reproduced from the EEI Report) shows the number of personnel, from 19 responder sites in 1989, with cumulative doses in the categories 25-50, 50-75, 75-100, 100-150, and > 150 rem for utility and contractor personnel. The EEI Report does not show how many individuals exceed a lifetime dose of their age in rem, but rather, the number of workers younger and older than 50 that appeared in each cumulative dose interval.




**Figure 4.3 Cumulative Site Doses for Utility and Contractor Personnel for 1989 (19 Responders)**

Of this total worker population, less than 50 utility and contractor personnel younger than 50 had lifetime exposures greater than 50 rem. Other findings on cumulative doses were: 1) no utility worker had lifetime doses greater than 75 rem, and 2) several contractor personnel had lifetime doses greater than 75 rem, and a couple had more than 150 rem in 1989.

#### 4.1.5 Projected Cumulative Doses for 1994

Figure 4.4 (reproduced from the EEI Report) shows the projected number of personnel from 14 responder sites anticipated to have cumulative doses in the same dose categories listed in Section 4.1.4. These numbers are for both utility and contractor personnel projected from past data trends out to 1994.



**Figure 4.4 Projected Cumulative Site Doses for 1994 Utility and Contractor Personnel (14 Responders)**

If these projections are realistic, less than 17 workers younger than 50 would have cumulative doses greater than 50 rem in 1994. Also, no utility or contractor personnel are expected to have a lifetime dose greater than 100 rem. The authors of the EEI report extrapolated this data to the entire nuclear industry "if we assume that the 15 responders represent one-fourth of the industry, we might expect about 600 workers with lifetime doses over 50 rem in 1994, with about one-fourth of them (i.e., 150, probably all contractors) over 75 rem and one-tenth of them (i.e., 60 contractors) over 100 rem."

#### 4.1.6 Effects of Changing the Annual Dose Guidance

The EEI questionnaire asked: "If all utilities adopted Uniform Site Annual Whole Body Dose Equivalent Administrative Limits (or guidance values), set at the following values, what difficulties, additional costs, collective dose increases, and ALARA effects do you see occurring: 4 rem, 3 rem, 2.5 rem, 2 rem, 1 rem, 0.5 rem?" The responses to this question were varied, and complicated by the fact that a similar question was asked: "If NRC lowered the 10 CFR 20 annual committed effective dose equivalent limit to the following values, what do you see occurring: 4, 3, 2.5, 2, 1, 0.5 rem?" The following conclusions were drawn from the responses:

## Survey Results

1. None of the seventeen responders felt that an annual dose limit of 4 rem would affect operations significantly. (Ten felt the effect would be minimal; seven said very minor.)
  2. According to seven responders, an annual limit of 3 rem is achievable, but the contractor's workforce would have to be expanded.
  3. At 2 rem yr<sup>-1</sup>, two of five responders felt the limit was achievable. One responder felt the limit was possibly achievable, and two felt it would significantly affect operations. An example given was the lack of a qualified labor pool to work outages.
  4. At 1 rem yr<sup>-1</sup>, all responders felt operations would be "...extremely difficult, if not impossible."
4. At a cumulative limit of 1 x age, only 1 responder predicted little effect on contractor personnel; 20 responders felt there would be impacts. The same impacts as those listed in 3. would occur, but to a greater degree.
  5. At the level of age times 0.5, most responders expected substantial effects on utility personnel and all but two see substantial effects for contractor personnel.

### 4.1.7 Effects of Establishing a Cumulative Dose Limit

The questionnaire asked, "If a cumulative or lifetime effective whole body dose limit were imposed by the NRC, what difficulties, additional costs, collective dose increases, and ALARA effects ... do you see occurring at 3 x age, 2 x age, 1.5 x age, 1 x age and 0.5 x age?" Because many of the 21 responders already had adopted a 1 x age administrative guideline and had experience with its effects, the responses were more consistent than those on other questions about anticipated effects:

1. Most responders felt that minimal impact would occur for utility personnel with a cumulative limit of 3 x age, 2 x age, and 1.5 x age; at a level of 0.5 x age, most saw substantial effects.
2. The majority of responders felt that minimal impact would occur for contractor personnel at 3 x age, and about half felt that there would be minimal impact at 2 x age.
3. At a cumulative limit of 1 x age, 11 responders saw minimal impact on the numbers of utility personnel; the 10 other responders mentioned impacts, such as scheduling problems, lack of critical plant specialists, increased personnel and associated dose for certain jobs, and additional costs, e.g., source term reduction modifications/operations, radiation protection, and salaries.

## 4.2 Department of Energy (DOE) Report

### 4.2.1 Cost Impact

Based on responses from 37 DOE contractors (~ 60%), the projected costs for all sites combined for a 20 mSv (2 rem) annual limit without a doubling of the neutron quality factor, and with a doubling of the neutron quality factor are as follows:

## Survey Results

	<u>Neutron Quality Factor of 10</u>	<u>Neutron Quality Factor of 20</u>
Personnel Costs	\$11M	\$15M
Modification Costs:		
Initial	\$279M	\$369M
Annual	\$ 3M	\$ 4M
Radiation Protection Costs:		
Initial	\$ 13M	\$ 17M
Annual	\$ 5M	\$ 7M
Increased Collective Dose	103 person-rem	243 person-rem

As noted in Section 3.1.2, the estimates do not include the Rocky Flats plants and Los Alamos National Laboratory, which have significant neutron exposures and collective doses. In addition, the costs associated with more restrictive Annual Limits on Intakes (ALI) for intakes of radioactive materials and the use of committed effective dose equivalent are not fully represented.

### 4.2.2 Annual Reported Doses, 1978 to 1988

Figure 4.5 (reproduced from the DOE report) shows a downward trend in the average annual dose equivalent for DOE personnel with measurable exposures from 1985 to 1988.




Figure 4.5 Average Annual Dose Equivalent for DOE Workers with Measurable Exposure, 1978-1988

The average dose per worker, with measurable exposure, was typically less than 2 mSv yr<sup>-1</sup> (200 mrem yr<sup>-1</sup>), which is well below both the DOE annual limit of 50 mSv yr<sup>-1</sup> (5 rem yr<sup>-1</sup>) and the proposed 20 mSv yr<sup>-1</sup> (2 rem yr<sup>-1</sup>). The recent decreases are attributable to DOE's continuing ALARA efforts and changes in its mission.

Figure 4.6 (taken from the DOE Report) shows the total number of DOE employees and visitors exceeding 2.0, 1.0, and 0.5 rem annually from 1978 to 1988.




Figure 4.6 Number of DOE Employees and annual dose 1978-1988

In 1988, the total number of DOE personnel and visitors exceeding ~ 2.0, 1.0, and 0.5 rem was 35, 548, and 1,862, respectively. If the decreasing trend in annual doses continues, a very small percentage (< 1%) of DOE workers will exceed 2.0 rem yr<sup>-1</sup>. Following the survey, the DOE issued its Radiological Control Manual in June 1992, establishing an administrative limit of 2.0 rem yr<sup>-1</sup>.

#### 4.2.3 Lifetime Cumulative Exposure Limits

The DOE survey on the impact of BEIR V asked the respondents to identify which workers might exceed or come within 10% of exceeding a cumulative lifetime limit of age in years times 1 rem. Respondents also were asked to estimate costs associated with implementing a cumulative dose limit (these data are not typically maintained at DOE contractor facilities).

Facility responses were summarized as follows:

- Few DOE contractor facilities responded to this question, because most did not maintain records on lifetime cumulative exposure; 57 workers were identified as having exceeded or being within 10% of exceeding the lifetime exposure limit.

- The current occupational categories for the 57 workers identified were as follows:

21% - Managers/Administrators  
 14% - Operators (plant/system/utility)  
 14% - Engineers  
 11% - Science Technicians  
 7% - Pipefitters

The remaining occupational categories represented less than 5% of the total.

- Total costs identified by the respondents for implementing a cumulative lifetime exposure limit of age in years x 1 rem are as follows (rounded to the nearest million):

Initial costs     \$1M  
 Annual costs     \$2M

The DOE Radiological Control Manual dated June 1992, established a requirement for a special control level of less than 10 mSv yr<sup>-1</sup> (1 rem yr<sup>-1</sup>) when a

worker's cumulative lifetime dose exceeds age in years.

#### 4.2.4 Impact on Facility Operations

The DOE survey on the impact of BIER V asked the respondents to identify those operations at their facility that would have to be discontinued if the proposed limits of 2.0, 1.0, and 0.5 rem were adopted. Two options were to be assumed, the current neutron quality factor (QF), and the proposed doubling of the neutron QF.

The responses from 60% of the DOE facilities (not including Rocky Flats and Los Alamos National Laboratory) are summarized below. We note that this summary does not identify all significant operational impacts.

##### 2.0 Rem Impact on Operations

Respondents identified typically little or no effect, both for the current neutron QF and assuming a neutron QF of 20. Previous internal reviews at Rocky Flats and the Los Alamos National Laboratory, however, identified that plutonium operations would be affected and will require significant modifications at a 2.0 rem limit, coupled with a neutron QF equal to 20.

##### 1.0 Rem Impact on Operations

With the current neutron QF, respondents from one research reactor facility identified the need to operate at a 25 percent reduction in power level.

Assuming a neutron QF of 20, the following additional operations would be discontinued:

- A heat source program and radiography operations at one facility.
- Plutonium metal production at one facility.

##### 0.5 Rem Impact on Operations

The impact of the proposed 0.5 rem on operations was severe, both with the current neutron QF and assuming a neutron QF of 20. Specific operations that would be discontinued or require a change in mission, in addition to the above, include the following:

## Survey Results

- Overall fuel and high-level waste processing operations - several respondents identified the need to construct new facilities, with extensive use of robots, to continue processing fuel and to carry out high-level waste operations.
- Respondents from one research reactor identified the need to operate at a 50% reduction in power level.
- The sampling, retrieval, and recovery of transuranic waste would be discontinued at one facility.
- Plutonium scrap recovery would be discontinued at one facility.
- A calorimetry program would be discontinued at one facility.

### 4.3 Selected 1990 Data from NRC REIRS

Table 4.1 gives the annual exposure data for 6 licensee categories for 1990. Additional data for 1989 are given for industrial radiographers in Table 4.2, for fuel fabricators in Table 4.3, for manufacturers and distributors in Table 4.4, and for nuclear power reactors in Table 4.5. A similar set of tables is provided for 1991.

The 1989 data gives a better measure of "verification" of the survey results while the 1991 data is provided to reflect any change in the dose distributions.

Table 4.1 Annual Exposure Data\* 1990

License Category	Number of Licensees Reporting	Number of Monitored Individuals	Number of Workers with Measurable Doses	Collective Dose (person-rems or person-cSv)	Average Individual Dose (rems or cSv)	Average Measurable Dose per Worker (rems or cSv)
Industrial Radiography	258	6,523	4,458	2,120	0.33	0.48
Manufacturing and Distribution	55	4,195	2,345	770	0.17	0.33
Low-Level Waste Disposal	2	925	119	35	0.04	0.29
Independent Spent Fuel Storage	2	190	102	33	0.17	0.33
Fuel Fabrication and Processing	10	13,756	3,233	287	0.02	0.09
Commercial Light Water Reactors***	116	189,254**	100,104**	36,607	0.19	0.37
Totals	443	214,568**	110,204**	39,739	0.19	0.36

\* Taken from Table 3.1 from NRC NUREG 0713 Vol 12 (Raddatz and Hagemeyer, 1993).

\*\* These figures are adjusted to account for the multiple counting of transient reactor workers.

\*\*\* Includes all LWRs that reported, although all may not have been in commercial operation for a full year, and excludes the gas-cooled reactor.

Survey Results

**Table 4.2 Annual Exposure Information for Industrial Radiographers\* 1989**

Type of Licenses	Number of Licenses	Number of Monitored Individuals	Workers with Measurable Doses	Collective Dose (person-rems or person-cSv)	Average Measurable Dose (rems or cSv)
Single location	66	832	304	41	0.13
Multiple locations	192	5,691	4,154	2,079	0.50
Total	258	6,523	4,458	2,120	0.48

\* Taken from Table 3.4 of NRC NUREG 0713 Vol 12 (Raddatz and Hagemeyer, 1993).

**Table 4.3 Annual Exposure Information for Fuel Fabricators\* 1989**

Type of License	Number of Licenses	Number of Monitored Individuals	Workers with Measurable Doses	Collective Dose (person-rems or person-cSv)	Average Measurable dose (rems or cSv)
Uranium Fuel Fab	8	11,583	2,992	243	0.08

\* Taken from Table 3.6 of NRC NUREG 0713 Vol 12 (Raddatz and Hagemeyer, 1993).

**Table 4.4 Annual Exposure Information for Manufacturers and Distributors\* 1989**

Type of Licenses	Number of Licenses	Number of Monitored Individuals	Workers with Measurable Doses	Collective Dose (person-rems or person-cSv)	Average Measurable Dose (rems or cSv)
M & D-"A"-Broad	10	3,091	1,862	6,551	0.35
M & D-Limited	45	1,104	410	38	0.09
Total	55	4,195	2,272	693	0.31

\* Taken from Table 3.5 of NRC Report 0713 Vol 12 (Raddatz and Hagemeyer, 1993).

**Table 4.5 Summary of Annual Whole Body Distributions By Year and Reactor Type  
1989**

Reactor Type	Number of Individuals with Whole Body Doses in the Ranges (rems or cSv)											Total Number Monitored	Number With Meas. Exposure	Total Collective Dose
	Not Measurable	Meas. <0.10	0.10-0.25	0.25-0.50	0.50-0.75	0.75-1.0	1.0-2.0	2.0-3.0	3.0-4.0	4.0-5.0	5-12.0			
BWRs	39,102	17,210	7,336	5,992	3,717	2,493	4,162	625	41	1		80,679	41,577	15,780
PWRs	54,572	29,791	13,030	10,747	5,759	3,384	4,712	607	43	-		122,645	68,073	20,812
Total	93,674	47,001	20,366	16,739	9,476	5,877	8,874	1,232	84	1		203,324	109,650	36,592

\* Adopted from Appendix F of NRC NUREG 0713 Vol 12 (Raddatz and Hagemeyer, 1993).

## Survey Results

### 4.4 Information Obtained from the 1984 EPA Report

#### 4.4.1 Male and Female Workers in the Nuclear Industry

The EPA report (EPA, 1984) shows that the number of male and female workers employed in radiation related work are roughly the same, about 600,000 women and slightly over 700,000 men. Figures 4.7 and 4.8 give the proportion of all male and female workers in various sectors, medicine, industry, the nuclear fuel cycle, government, and miscellaneous fields, including those in nuclear power operations. The data have been separated into male and female subgroups because of the different kinds of activities that they pursue.




Figure 4.7 Percent of Male Radiation Workers in Various Sectors

Figures 4.7 and 4.8 show that males and females carrying out very different kinds of tasks. Most female radiation workers are employed in medicine and dentistry, whereas the male radiation workers are fairly evenly split among all the various sectors, with industry being the largest. Further analysis indicated that the males employed in medicine are performing different functions than the females.

It also is noteworthy that the mean age of all male radiation workers is slightly higher (36 years) than that for females (31 years).




Figure 4.8 Percent of Female Radiation Workers in Various Sectors

#### 4.4.2 Correlation of Radiation Dose with Age

Two of the dose limit options include lifetime limits on dose. This approach has been questioned because some experts feel that the older radiation workers (age 40 or older), because of their greater experience, may be required for tasks which expose them to high doses. This would imply that the older workers would have had higher annual doses.

To assess this view, we checked for a correlation between age and radiation dose for occupational workers. The age group data given in the EPA report (EPA, 1984) were transformed to mean ages for each group and compared with the mean annual dose to each group. The data were weighted by the number of workers in each group. Table 4.6 shows the mean annual dose equivalent for all U. S. radiation workers by sex and age.

#### 4.4.3 Males

In Table 4.6, the relationship between age and mean annual exposure indicates that males aged 22 to 42 had the higher exposures. For those aged 42 to 67 there is a downward trend of exposure with age.

#### 4.4.4 Females

Table 4.6 shows a somewhat different picture. As women radiation workers grow older they seem to receive more radiation, although the mean annual dose is low compared with males of all ages. As shown in Figure 4.8, the majority of female radiation workers are employed in the medicine and dentistry, which probably accounts for the mean annual dose for women being about 60 mrem in contrast to the significantly higher mean annual dose for men, who work primarily in industry (Figure 4.7).

Table 4.6 Mean Annual Dose Equivalent for U.S. Radiation Worker

Age	Males		Females	
	Mean Annual Dose (mrem)	Number of Workers	Mean Annual Dose (mrem)	Number of Workers
19	100	8,035	40	25,090
22	210	84,336	50	168,534
27	180	147,742	50	158,986
32	160	157,869	60	94,237
27	160	104,636	60	57,865
42	170	69,220	60	38,649
47	150	52,934	60	24,760
52	130	39,650	60	17,608
57	130	30,781	60	12,360
62	100	14,489	50	5,389
67	90	5,958	60	1,485

#### 4.5 Responses to the Request for Comment

There were seven responses to the draft NUREG/CR-6112, "Impact of Reduced Dose Limits on NRC Licenses Activities - Major Issues in the Implementation of ICRP/NCRP Dose Limit Recommendations," which was distributed for comment. It is interesting to note that, although there were only seven, they were extremely valuable in updating the report and in enforcing the validity of the recommendations.

Discussion of the responses including resolution of the comments are grouped by practice or industry types as done in Section 5.

#### 4.5.1 Nuclear Power Reactor and Nuclear Power Reactor Contractors

##### 4.5.1.1 Organization of Licensees

An extensive response was received from an organization of nuclear power plant licensees. A major issue raised by the authors deals with the omission of an evaluation of ICRPs Publication 60 recommendation of 100 mSv in 5 years (10 rem in 5 years). The working party outlined in Section 3.2.3 discussed this topic in some detail, and the general consensus was that such a scheme would be very difficult to legislate and, therefore, be an unlikely candidate for regulation. For example, the National Radiological Protection Board in the U.K. has recommended a limit of 20 mSv yr<sup>-1</sup> (2 rem yr<sup>-1</sup>), and several other European countries

## Survey Results

have formally adopted  $20 \text{ mSv yr}^{-1}$  in their regulations. The authors of the CEC Draft Safety Standards recognize this issue by giving the member countries the option of either  $100 \text{ mSv}$  in 5 years ( $10 \text{ rem}$  in 5 years) or  $20 \text{ mSv yr}^{-1}$  ( $2 \text{ rem yr}^{-1}$ ).

The authors made several observations which they summarize as follows: "...there are aspects of lower annual dose limits which create impacts that potentially make such limits inappropriate and inefficient as regulatory limits for the purpose of controlling cumulative lifetime dose consistent with ICRP and NCRP recommendations." The detailed comments on this document have been added to the comments section of Section 5.2

The authors expanded on the concept presented in the draft report that the need for administrative guidelines below the dose limit further exacerbates the issue. The authors note, for example, "The difference in potential impacts associated with a typical administrative dose guideline currently used in the nuclear power industry,  $4 \text{ rem}$  ( $0.8 \times 5 \text{ rem}$ ) and the guideline inferred by the higher of the two annual dose limit options in the draft NUREG, e.g.,  $1.6 \text{ rem}$  ( $0.8 \times 2 \text{ rem}$ ), are significant. The NRC reports that in 1989, the year surveyed for the draft NUREG, 34 workers received doses in excess of  $4 \text{ rem}$ , approximately 2,000 received doses in excess of  $2 \text{ rem}$ , and more than 10,000 workers received doses in excess of  $1 \text{ rem}$ . This would indicate that the magnitude of potential impacts will increase markedly if regulatory dose limits reduce from  $5 \text{ rem per year}$  to  $2 \text{ rem per year}$ , entailing use of an administrative dose guideline of  $1.6 \text{ rem per year}$ ." This issue is emphasized in the Summary and Executive Summary sections of this report.

The authors raise a number of issues related to dollar/person-rem issues. Their concerns have been addressed by adding an extended summary to Section 7 outlining the limitations and uncertainties in such estimates.

The impact of reduced exposure limits for the embryo-fetus was raised by the authors. Since the information on this group of exposed individuals has not been addressed in any of the studies cited, a note to that effect has been added to the summary section of this report.

The authors noted that there are many instances in "high dose" tasks where non-uniform exposure takes place. In such situations "multi badging takes place" and the maximum reading is assigned as a whole-body dose. Under the EDE concept, this would certainly overestimate the exposure, but such overestimates are unlikely to be as important in assessing the impact of reduced dose limits for nuclear power workers as they are for medical x-ray workers whose TEDE is likely to be about 50% of their badge reading.

The authors collected extensive data on workers who exceeded their age (or came close to it) in 1993, and on the workers who exceeded  $2 \text{ rem}$  in 1993. The estimated total number of workers in each of these categories are shown in Table 5.5. They noted that "maintenance, mechanics, and fuel handlers make up more than half of the workers with lifetime dose greater than age. The remainder include managers and supervisors, health physics technicians, welders, engineers, in-service inspectors, and others. Utility and non-utility employees make up about half of the workers with lifetime dose greater than age." They further noted that, "mechanics, management, and health physics technicians are predominantly utility workers, while fuel handlers, welders, engineers, and in-service inspectors are mostly non-utility employees. The predominantly non-utility work groups include outage workers with specific skills and experience qualifications that lead to a higher demand for their work specialties at multiple facilities during the year (e.g., to support refueling outages). Their replacement with less experienced workers would be difficult." And "maintenance mechanics, steam generator workers, and engineers make up nearly 60% of the workers whose 1993 annual doses were greater than  $2 \text{ rem}$ . The remainder include insulators, health physics technicians, fuel handlers, carpenters, welders, in-service inspectors, deconners, and others. Overall, 88% of the workers with 1993 annual doses greater than  $2 \text{ rem}$  are non-utility workers. Maintenance workers consist of 21% utility workers and 79% non-utility workers. For the balance of workers in other work groups, 95% of the workers are non-utility workers. This is in contrast to the population of workers with lifetime dose greater than age, who are more evenly divided between utility and non-utility workers."

The authors also provided age distribution data for those who exceed either of these thresholds. For lifetime greater than age in rem, the average age is 44 years, ranging from 24 to 75 years. For those exceeding 2 rem per year, the average age is 36 years with the average lifetime dose being 14 rem.

The authors raised the issue of worker employability, which would arise with a 2 rem per year limit. They noted, "a lower annual dose limit may severely impact the employability of some non-utility workers who are routinely employed at several outages during the course of a year because of their specialized skill and experienced qualifications. Some utility workers may also be impacted, in particular, workers who are employed at multiple outage within the utility during the year and utility workers."

A similar issue raised by the authors was the need for additional workers. For example, they note, "steam generator work, control rod drive maintenance, in-service inspections, and the reactor head work are jobs with higher potential for impacts that may result in the use of increased numbers of workers because these jobs may involve higher individual doses to workers" And "workers such as insulators, in-service inspectors, pipe fitters, mechanics, and welders are among those expected to be constrained by lower dose limits potentially necessitating an overall increase in the available numbers of workers. The increased numbers of workers indicated for health physics technician and decontaminators may be indicative of the potential need to provide more extensive support (in the form of health physics coverage and decontamination services) for jobs where workers are expected to be constrained by lower limits, e.g., steam generator work and controlled rod drive maintenance."

The authors note that there could be an important impact on outage scheduling, specifically, "for example, job preparation activity such as decontamination and installation of temporary shielding may be expanded in response to lower dose limits with the effect of extending the overall schedule for the job. Also, contingencies may emerge that lead to an expanded work scope and the need to bring in additional workers due to dose considerations that may result in delays and extend the job

schedule." And "Jobs with higher potential for such impacts include steam generator work, reactor head work, plant modifications, primary system maintenance, and in-service inspection. Some of these jobs may be critical to time of completion of the overall outage schedule. Therefore, scheduled impacts on these jobs could lead to significant cost increases, e.g., replacement power costs. For such critical jobs, the number of in-processed and available workers for the outage may be increased as a contingency, which also leads to additional costs impacts."

#### **4.5.1.2 Three Separate Responses were Received from Three of the Nuclear Power Plant Sites**

One of these provided information on about 60,000 transient workers from data obtained from the National Index System used by 19 utilities. This data is given in Table 5.8 in Section 5.3. The authors noted, "For the most flexibility we can obtain low radiological risk while allowing for major outages by retaining the annual regulatory limit of 5 rems TEDE and by capping lifetime dose equivalent in rems at the workers age." They did indicate the need for allowing 2 rem/yr for those who exceeded the lifetime limit. The second response expressed concern over the dated material in some sections of the draft report. They noted, "We believe that the results of this report would likely be altered if recent plant dose trends and ALARA data were considered." It would appear that this conclusion is not true as evidenced by the extensive survey discussed above based on 1993 information. The conclusions one might reach from this data are just about the same as those one might reach from the data in the draft report that went out for comment.

The authors also raised the question about the current risk estimates. The data in the draft report is based on the latest UNSCEAR, ICRP, and NCRP reports. There is no later estimate available at this time.

The third response raised many of the questions raised in the extensive nuclear industry response discussed above. One additional interesting issue the authors raised is the concern over the ability to obtain realistic historical dose estimates. As

## Survey Results

with any others, they felt the grandfather clause would be very important.

### 4.5.2 Test and Measurement Including Industrial Radiography

A response was received from a professional association concerned with industrial radiographic workers. The authors provided update information which has been included in Section 5.4 in both Table 5.11 and within the comments section. They also noted that the comments given in Section 5.2 and 5.3 generally apply to radiographic procedures. They also deplored the lack of input from industrial radiography organizations, although one of the working party members was selected on the basis of his extensive experience in this field.

### 4.5.3 Manufacturing and Distribution Including Cyclotron Produced Radiopharmaceuticals

A response from a concerned medical society questioned the validity of the NCRP and UNSCEAR risk estimates. This issue was coupled with a concern over the potential increase in costs for preparing a radiopharmaceutical if the limits were reduced. The issues raised in this regard are included in the comments to Section 5.6.

A second response in this category of radiation users was received from a trade organization concerned with this aspect of radiation protection. The authors supplied additional detailed information collected from radiopharmaceutical suppliers, which has been incorporated in Section 5.6 both in the tables and in the comments section. The authors noted, "We agree with most of the findings of the draft NUREG/CR-6112 as shown in the attached comments. The current occupational dose limit of 5 rem/yr coupled with an age in rem limit could be viable provided that a grandfather clause is provided to exclude a few long-term employees. It is premature to implement a 2 rem/yr limit, that it might be considered as a design goal for new operations or a criteria for routinely conducting audits to show the doses are ALARA. A 1 rem/yr limit is unlikely to be viable in the near future."

### 4.5.4 Fuel Fabrication, UF<sub>6</sub> Production

Part of the extensive response received from the nuclear power plant licensee organization dealt with this topic. The authors had received information from fuel fabrication facilities and other uranium use companies. The conclusion to their general comments was, "The potential impacts of lower dose limits on the nuclear fuel fabrication industry could be significant. The employability of some workers might be significantly restricted, especially in the case of a lifetime dose limit. The number of workers would likely be increased, resulting in a larger number of exposed workers, potentially involving an increase in collective dose. One-time costs associated with implementing lower dose limits are estimated at \$100,000 to \$1,000,000 per facility and possibly more for some facilities. Increased annual cost per facility related to changes in radiation protection programs are estimated at \$50,000 to \$100,000 per year. If staffing increases are needed, the increased annual cost per facility would be much higher."

They also provided a list of potential impacts which have been included in Section 5.8. The authors wanted to particularly emphasize the impact of combining internal and external exposures. For example, they note specifically the issue of complexity involved in the transition from ICRP 2 to ICRP 26 dose methods, especially as related to licensees with sources of internal exposure, should be highlighted and emphasized in the executive summary. In particular, the expected increase in reported individual dose to fuel fabrication industry workers, as a product of this complexity in dose methods, and not reflective of an increase in actual worker doses should be explained. The discussion in the final report should acknowledge the need to further assess dose trends among fuel fabrication licensees using data developed under the revised Part 20 to make valid estimates of potential benefits of lower limits and the associated impacts.

## 5 Questionnaire Results Obtained in this Survey

The data given here should be taken as an indication of the issues. A small number of responses was received from each industry. As noted in Section 3.2.3., the validity of the conclusions of the report depend on the working groups' detailed evaluation, for which the responses to the questionnaire provided a framework. The responses and the data given by the working committee are presented by practice or industry type, and are summarized for each in two tables. This data has been supplemented with an addition of information contained in the comments to the draft NUREG/CR-6112.

The first table, for each industry class, gives information on impacts, and the second on exposure experience. In addition, all comments from the questionnaire or from the members of the working party are given.

### 5.1 Medical/Dental and Veterinary Practice

There were 20 responses from medical institutions and one from veterinary practice.

Table 5.1 Impacts on Medical/Dental and Veterinary Practice

Impacts	Possible Dose Limit			
	2 rem y <sup>-1</sup>	1 rem y <sup>-1</sup>	5 rem y <sup>-1</sup> age in rem	2 rem y <sup>-1</sup> age in rem
<i>Collective dose:</i>				
No Change	17	15	19	19
Increase	-	2	-	-
% Decrease	3	3	1	1
<i>Facility Modification:</i>				
None required	13	8	19	14
Modifications required	7	11	-	3
Costs (individual responses)	\$2K to \$150K	\$4K to \$300K	-	\$24K to \$100K
<i>Radiation Protection Cost:</i>				
No increase	11	8	19	12
Will increase	9	11	-	6
Cost/yr (individual responses)	\$3K to \$100K	\$1K to \$100K	-	\$16K to \$100K

## Questionnaire Results

**Table 5.2 1989 Exposure Experience**

<b>Number of Employees with Annual Doses:</b>
>5 rem <u>11</u> >2 rem <u>14</u> >1 rem <u>47</u>
Number of Employees with Lifetime Dose Greater Than Age in Rem: <u>3</u>
Number of Employees with Measurable Dose: <u>4370</u>
Annual Collective Dose: <u>613 rem</u>

### Medical/Dental and Veterinary Practice Comments

1. Our cardiac catheterization and angiography areas are the biggest person-rem inflator. All recommended shield/safeguards are in place, but with patient volume, exposures are still high. Limiting annual doses to 1-2 rem would be unattainable.
2. This is a cardiac catheterization lab and reported dose is outside the apron. Facility modifications may not be possible and very much depends on the willingness of the cardiologist.
3. Bad idea to require  $D < \text{current}$ . Means increased therapy room shielding at no benefit. Waste of patient resources and care.
4. Film readings outside the apron. Six cardiologists will require \$8,000 in ceiling-suspended shielding in 3 cath labs just to keep the badge readings down. This dose limit is not justified.
5. There is an urgent need for guidance on computation of effective dose equivalent.
6. This change will be of no impact in the medical field.
7. Why bother unless there is clear evidence of harm at annual doses less than 5 rem? RERF database is hardly applicable to medical workers.
8. The 1 rem yr<sup>-1</sup> limit might have some problems for radiologists/cardiologists performing fluoroscopy on collar badge readings. These groups are provided with two dosimeters.
9. In the design of medical facility, there would be a significant increase in construction costs and essentially no benefit to patients or personnel. Must consider badge position.
10. Data are whole-body exposure, when two badges worn, data for that worn under Pb apron used.
11. Increase in Radiation Protection cost for purchase of additional lead glasses and thyroid shields and possible use of double badging for specific groups of workers.
12. Current dose limits are ambiguous when applied to a diagnostic radiology department since the film badge measurement dose is typically 5 to 10 times the EDE due to apron, glasses, etc.
13. Although our actual exposures are low, a change in the "general public" levels would require modifications at our vaults. The expense would be non-trivial and the benefit would be trivial. I don't believe any of the current evidence warrants changing the current limits.
14. Above are whole-body doses only. If "head and neck" dose limits is reduced to 1 rem yr<sup>-1</sup> our cardiology physicians would have to limit the number of cardiac cath cases.
15. Might as well do away with  $< \text{age}$ . 1 rem too low for special procedures.
16. The data showing 11 people over 5 rem yr<sup>-1</sup> and 14 over 2 rem from 20 sources may be lower than the real numbers. In my experience, a significant number of personnel using fluoroscopy do not wear the dosimeter that is provided.

17. Let's set a 10 mSv annual BRC/de minimum dose as soon as possible, so we can focus efforts on the real health hazards of radiation and stop wasting time, money, and personal efforts on trivia.
18. Personnel dosimetry data reported by medical institutions for radiation producing devices personnel, such as cardiologist, invasive radiologist, etc., should be viewed with some suspicion as these individuals may be badged at more than one institution and may fail to properly use such devices.
19. How much reduction in personnel exposure can be realized by proper radiation safety instructions furnished to non-radiology personnel is difficult to assess. How much radiation safety instructions are furnished to physicians, nurses, operating room personnel, etc., outside the radiology departments

is variable. Additionally, application of these instructions to properly reduce exposure is also variable. Uniform instructions and operational application of proper technique might reduce exposure at very little additional cost.

20. With regard to the frequently reported partial-body exposures to personnel performing medical procedures, there exists a need for guidance as to the proper assessment to a whole-body dose equivalent. Without such equivalency, the "outside the apron" dose is discounted as insignificant or overstated as impossible to do anything about.

## 5.2 Nuclear Power Reactors

There were seventeen responses from nuclear power stations.

Table 5.3 Impacts in Nuclear Power Reactors

Impacts	Possible Dose Limit			
	2 rem y <sup>-1</sup>	1 rem y <sup>-1</sup>	5 rem y <sup>-1</sup> age in rem	2 rem y <sup>-1</sup> age in rem
<i>Collective dose:</i>				
No change	6	1	17	6
Increase	10	15	-	11
Decrease	1	1	-	-
% Increase	2 to 20	5 to 30	-	2 to 20
<i>Facility Modifications:</i>				
None required	9	3	16	9
Modifications required	8	14	1	7
Costs (individual responses)	\$25K to \$10M	\$25K to \$50M	\$25K	\$50K to \$10M
<i>Radiation Protection Cost:</i>				
No increase	8	1	16	7
Will increase	9	16	1	10
Costs/yr (individual responses)	\$5K to \$.5M	\$5K to \$1M	\$100K	\$5K to \$750K

Questionnaire Results

Table 5.4 1989 Exposure Experience

Number of Employees with Annual Doses:		
>5 rem <u>0</u>	>2 rem <u>331</u>	>1 rem <u>3,101</u>
Number of Employees with Lifetime Dose Greater Than Age in Rem: <u>178</u>		
Number of Employees with Measurable Dose: <u>24,098</u>		
Annual Collective Dose: <u>10,915</u>		

Table 5.5 1993 Exposure Experience (estimated)

Number of Employees with Annual Doses:		
>5 rem <u>0</u>	>2 rem <u>1000</u>	>1 rem <u>      </u>
Number of Employees with Lifetime Dose Greater Than Age in Rem: <u>500</u>		
Number of Employees with Measurable Dose: <u>      </u>		
Annual Collective Dose: <u>      </u>		

Nuclear Power Reactor Comments

1. Two rem yr<sup>-1</sup> is a challenge but achievable with management support. 1 rem yr<sup>-1</sup> will require major modifications and increase in personnel (especially for older facilities >15 years).
2. LWR's will not be able to operate with a 1 rem yr<sup>-1</sup> limit.
3. Facility modifications should not be necessary; specialized tasks or maintenance evolutions may result in higher doses for a few individuals (i.e., 10-15), more frequent TLD processing may be required, outage contractors may be unavailable for work due to dose restrictions.
4. A limit of cumulative < age, 3 rem yr<sup>-1</sup> not to exceed 10 rem in 5 years is workable. We need flexibility.
5. We are attempting to limit HP Techs to <1 rem for 1990; it could have been done in 1989. A few

(contractor) employees have > rem than years may be put out of work. Initial approach to  $\leq 1$  rem yr<sup>-1</sup> will probably be to hire more people.

6. Costs are extremely difficult to assess.
7. The nuclear power facilities have not provided an informed, representative response to the questionnaire.
8. We recognize that the current regulatory limits do not provide a total lifetime dose limitation other than the defacto limit of 5 rem yr<sup>-1</sup> and, therefore, theoretically allows significant lifetime dose. If the regulatory limits need updating, the annual dose limits should not be changed, and a lifetime dose limit should be instituted equivalent to the NCRP recommendation of age = rem, with the proviso that persons who have already exceeded this limit be provided a special annual limit of 1-2 rem.

## Questionnaire Results

9. The use of administrative dose limits established below regulatory dose limits should be considered.
10. Two  $\text{rem yr}^{-1}$  limit would be difficult and costly, but achievable for utility workers. However, for contract personnel it would be very difficult and exceedingly costly.
11. For those individuals who would exceed the lifetime limit of age in rem, a  $2 \text{ rem yr}^{-1}$  limit would be necessary in order to maintain their employment within the industry.
12. The number of the more highly exposed contractor staff working our outages ranges from 50-100, each receiving 1-2 rem per outage. Since the contractor staff works up to four to five outages per year, each of the more highly exposed workers becomes restricted by year's end under the current administrative dose limits of approximately  $4 \text{ rem yr}^{-1}$ . [Note that most of the contractor staff do not have a "high" lifetime dose (e.g.,  $0.2\text{-}0.5 \times$  age in rem), as their employment has not always been in the higher dose work activities.
13. If lower regulatory dose limits were instituted, the contractor companies would be forced to hire more "temporary" staff, perform more training, charge higher rates, and, as a result, increase the financial cost. More importantly, this would result in increased collective dose due to using a larger and less skilled workforce. Likewise, we would incur an increase in our company Health Physics and support staff's dose since we would be supporting a larger, less skilled radiation worker force. In addition, the use of more "temporary, less skilled" workers also increased the probability of personnel error, which is a decrease in nuclear safety for both the co-workers and the general public.
14. In the process of setting new regulatory dose limits, it is important to understand the dose limitation system typically in use at nuclear facilities restricts actual doses to approximately 80 percent of the regulatory limits; i.e., "administrative limits" are set by the utilities well below the "regulatory limits." The use of administrative dose limits provides a "safety margin" designed to help the worker avoid exceeding regulatory limits. If the NRC regulatory limit were  $2 \text{ rem yr}^{-1}$ , nuclear facilities would essentially be required to set administrative limits in the range of  $1.5 \text{ rem yr}^{-1}$ .
15. In addition to regulatory and administrative dose limits, the nuclear industry has achieved successes in steadily reducing individual, collective, and lifetime accumulated dose to As Low As Reasonably Achievable (ALARA). In light of the entire system of dose limitation and ALARA practices, we believe current annual dose limits under the revised 10 CFR 20 provide appropriate and adequate worker protection. In addition, an ALARA cost/benefit analysis has not been performed, which indicates that reductions in the individual's annual dose justify the expected increase in collective dose.
16. If reduced dose limits must be instituted, we believe that the important parameter to control should be lifetime dose, not annual dose. A modified lifetime limit similar to the National Council on Radiation Protection and Measurements (NCRPs) recommendation would be appropriate. The modification would be to allow a  $1\text{-}2 \text{ rem yr}^{-1}$  provision for persons who are approaching or have already exceeded this limit. We believe that the International Commission on Radiological Protection (ICRP) recommendation of 10 rem in five years, with a yearly limit of 5 rem, would unnecessarily restrict our operational flexibility.
17. It is noted that the dose risk models of BEIR V do not make a distinction between the risk for chronic exposures based on annual dose rates which vary from  $2\text{-}5 \text{ rem yr}^{-1}$ , i.e., risk associated with chronic exposure is primarily a function of total dose. Therefore, risk associated with current regulatory dose limits could be reduced by use of the NCRP recommendation for lifetime dose with a  $5 \text{ rem yr}^{-1}$  cap, while simultaneously allowing us the operational flexibility necessary to operate efficiently.
18. Provision should be made to permit exposures in excess of the limits, i.e. special planned exposures. This may be particularly true if NRC mandated backfits occur.
19. With an annual limit of 2 rem or less, some discretionary safety related inspections might have to be constrained.

## Questionnaire Results

20. Lower annual dose limits would impact nuclear power reactor licensees more than the longer-term limits recommended by the ICRP and NCRP, without any substantial improvement to worker health and safety. For example, the use of a one year term for limiting dose is incongruent with the typical 18 month fuel cycle at nuclear power plants. The majority of worker doses are received during refueling outages. Therefore, during one year at continuous power operations, relatively little dose may be received by utility workers. In the following year, workers may approach, and be constrained by the lower dose limits because of work performed during a refueling outage. In this case, the average dose for the two year period, which is a more valid reflection of lifetime risk, would be well below the lower annual dose limit, but the limit would unnecessarily impact the ability to perform outage work. Contract workers attempting to work in several outages during the course of a year may be severely constrained, even though the potential annual dose would likely be below 5 rem (based on current dose experience) and the cumulative lifetime dose received over the entire working lifetime would also be within ICRP and NCRP recommendations. The potential impacts imposed in this case would be without substantial benefit to worker health and safety.
21. Lower annual dose limits would deny some operational flexibility. For example, such limits would be in conflict with the trend to alternate between major and minor outage scopes to provide for relatively short duration refueling outages in alternate outage years. The major outage scope could be constrained by the lower annual dose limits, and may be economically unacceptable due to associated cost impacts. Also, significant large-scale projects that occur very infrequently would be disproportionately impacted by lower annual dose limits. Examples of such projects include steam generator or recirculation piping replacement projects.
22. Unscheduled outages, e.g., due to unanticipated plant shutdown, if occurring late in the year may be significantly impacted by the lack of remaining allowable dose among key workers who received doses approaching the lower annual dose limit during a routine refueling outage earlier in the year. Similarly, a task involving moderate dose that emerges in a routine outage (e.g., a stuck reactor vessel stud) could result in significant schedule impacts due to constraints of lower annual dose limits on critical workers with limited remaining allowable dose. Loss of operational flexibility in these outage scenarios could result in additional individual and collective dose, and induce resource impacts greater than those associated with longer-term limits.

### 5.3 Nuclear Power Reactor Contractors

There were three responses from power reactor contractors.

Table 5.6 Impacts in Nuclear Power Reactor Contractors

Impacts	Possible Dose Limit			
	2 rem y <sup>-1</sup>	1 rem y <sup>-1</sup>	5 rem y <sup>-1</sup> age in rem	2 rem y <sup>-1</sup> age in rem
<i>Collective dose:</i>				
No change	1	-	3	2
Increase	2	3	-	1
Decrease	-	-	-	-
% Increase	50 and 100	50 to 1000	-	100
<i>Facility Modifications:</i>				
None required	1	1	2	2
Modifications required	1	1	1	1
Costs (individual responses)	up to \$50M	\$100M	\$1M	\$20-50M
<i>Radiation Protection Cost:</i>				
No Increase	-	-	3	2
Will Increase	2	2	-	1
Costs/yr (individual responses)	\$.2M and \$10M	\$.3M and \$25M	-	\$10M

Questionnaire Results

**Table 5.7 1989 Exposure Experience**

Number of Employees with Annual Doses:		
>5 rem <u>  1  </u>	>2 rem <u> 448 </u>	>1 rem <u>1,871</u>
Number of Employees with Lifetime Dose Greater Than Age in Rem: <u> 56 </u>		
Number of Employees with Measurable Dose: <u> 5,292 </u>		
Annual Collective Dose: <u> 1,718 </u>		

**Table 5.8 1993 Exposure Experience\***

Number of Employees with Annual Doses:		
>5 rem <u>      </u>	>2 rem <u> 516 </u>	>1 rem <u>1,812</u>
Number of Employees with Lifetime Dose Greater Than Age in Rem: <u>      </u>		
Number of Employees with Measurable Dose: <u>      </u>		
Annual Collective Dose: <u>      </u>		

\* Transient workers at 19 nuclear utilities

Nuclear Power Reactor Contractors Comments

1. The dose limits in the new 10 CFR 20 we are prepared to meet. Dose limits on the order of 1 rem yr<sup>-1</sup> per person would be catastrophic. AIF study 10 years ago showed this.
2. Only 5-10 members of the work force annually accumulate exposures greater than 1 rem. They are the most skilled and efficient. If limited, the collective dose will increase.
3. The general population is young and usually change jobs in 5-7 years, thereby not accumulating a large lifetime dose.
4. Can meet a 100 rem/lifetime plus a "grandfather clause" with 5 rem yr<sup>-1</sup> limit.
5. A "grandfather clause" would be necessary if a lifetime limit is adopted.

6. Utilities that perform their own outage maintenance will have many of the same difficulties as the contractors.

## 5.4 Test and Measurements Including Industrial Radiography

There were nine responses from test and measurement groups.

**Table 5.9 Impacts in Test and Measurements Including Industrial Radiography**

Impacts	Possible Dose Limit			
	2 rem y <sup>-1</sup>	1 rem y <sup>-1</sup>	5 rem y <sup>-1</sup> age in rem	2 rem y <sup>-1</sup> age in rem
<i>Collective dose:</i>				
No change	8	5	9	8
Increase	1	1	-	1
Decrease	-	2	-	-
% Increase	-	20	-	-
<i>Facility Modifications:</i>				
None required	9	7	9	9
Modifications required	-	2	-	-
Costs (individual responses)	-	\$20K to \$400K	-	-
<i>Radiation Protection Cost:</i>				
No increase	6	3	9	5
Will increase	3	3	-	1
Cost/yr (individual responses)	\$15K to \$25K	\$30K to \$50K	-	\$25K

Questionnaire Results

Table 5.10 1989 Exposure Experience

<p><b>Number of Employees with Annual Doses:</b></p> <p>&gt;5 rem <u>  0  </u>    &gt;2 rem <u> 10 </u>    &gt;1 rem <u> 21 </u></p> <p>Number of Employees with Lifetime Dose Greater Than Age in Rem: <u>  0 </u></p> <p>Number of Employees with Measurable Dose: <u> 285 </u></p> <p>Annual Collective Dose: <u> 109 </u></p>
---

Table 5.11 1993 Exposure Experience

<p><b>Number of Employees with Annual Doses:</b></p> <p>&gt;5 rem <u>      </u>    &gt;2 rem <u> 139 </u>    &gt;1 rem <u> 178 </u></p> <p>Number of Employees with Lifetime Dose Greater Than Age in Rem: <u>      </u></p> <p>Number of Employees with Measurable Dose: <u> 229 </u></p> <p>Annual Collective Dose: <u> 412 </u></p>
--

Tests and Measurements Comments

1. We have large NDT x-ray facilities, but radiation protection practices effectively limit the monthly dose to 25 to 50 mrem.
2. It is anticipated that any reduction in dose limit below the existing 5 rem/yr limit will increase (1) collective work force dose, (2) size of work force, (3) cost to modify facilities, (4) cost to purchase equipment, (5) cost of radiation protection, and (6) overall operating cost of business. Accurate estimates are not available at this time, but should be considered by the NRC prior to the development of a proposed rule to reduce the dose limits.
3. We also anticipate that any reduction in dose limit below the existing 5 rem/yr limit will decrease average worker training, skill, experience, and productivity and will cause a corresponding increased risk of accidental acute overexposure.

In addition, some types of radiography work will be difficult to do with reduced dose limits.

4. The NRC 10 CFR Part 34 radiography regulations recently adopted reduced dose limits and additional ALARA requirements. Radiography licensees will need a few years to evaluate the impact of these changes. It is too soon to provide an accurate estimate of the impact of additional reductions in dose limits.

## 5.5 Universities

There were four responses from universities.

**Table 5.12 Impacts in Universities**

Impacts	Possible Dose Limit			
	2 rem y <sup>-1</sup>	1 rem y <sup>-1</sup>	5 rem y <sup>-1</sup> age in rem	2 rem y <sup>-1</sup> age in rem
<i>Collective dose:</i>				
No change	4	4	4	4
Increase	-	-	-	-
Decrease	-	-	-	-
<i>Facilities Modifications:</i>				
None required	4	3	4	4
Modifications required	-	1	-	-
Costs (individual responses)	-	\$80K	-	-
<i>Radiation Protection Cost:</i>				
No increase	3	3	4	4
Will increase	1	1	-	-
Costs yr (individual responses)	\$80K	\$80K	-	-

**Table 5.13 1989 Exposure Experience**

<p><b>Number of Employees with Annual Doses:</b></p> <p>&gt;5 rem <u>  0  </u>    &gt;2 rem <u>  1-2  </u>    &gt;1 rem <u>  3-4  </u></p> <p>Number of Employees with Lifetime Dose Greater Than Age in Rem: <u>  0  </u></p> <p>Number of Employees with Measurable Dose: <u>  850  </u></p> <p>Annual Collective Dose: <u>  255  </u></p>
--

Questionnaire Results

Universities Comments

1. The kinds of activities carried out in this university environment should not weigh heavily in setting dose limits for high hazard work environments.

## 5.6 Manufacturing and Distribution, Including Cyclotron Produced Radiopharmaceuticals

There were six responses from this group.

**Table 5.14 Impacts in Manufacturing and Distribution**

Impacts	Possible Dose Limit			
	2 rem y <sup>-1</sup>	1 rem y <sup>-1</sup>	5 rem y <sup>-1</sup> age in rem	2 rem y <sup>-1</sup> age in rem
<i>Collective dose:</i>				
No change	4	1	3	1
Increase	2	5	1	2
Decrease	-	-	-	-
% Increase	8-13	18-25	6	13-17
<i>Facility Modifications:</i>				
None required	2	-	3	2
Modifications required	4	6	1	2
Costs (individual responses)	\$25K-\$44M	\$10K-\$127M	\$10M	\$60K-\$53M
<i>Radiation Protection Cost:</i>				
No increase	3	-	4*	2
Will increase	3	6	1	3
Cost/yr (individual responses)	\$30K- \$700K	\$10K to \$1.4M	\$100K	\$800K

\* 3-no increase, 1-not sure

Table 5.15 1989 Exposure Experience

<b>Number of Employees with Annual Doses:</b>		
>5 rem	<u>0</u>	>2 rem <u>20</u> >1 rem <u>72</u>
Number of Employees with Lifetime Dose Greater Than Age in Rem: <u>6</u>		
Number of Employees with Measurable Dose: <u>117</u>		
Annual Collective Dose: <u>86</u>		

Table 5.16 1993 Exposure Experience

<b>Number of Employees with Annual Doses:</b>		
>5 rem	<u>0</u>	>2 rem <u>97</u> >1 rem <u>327</u>
Number of Employees with Lifetime Dose Greater Than Age in Rem: <u>18</u>		
Number of Employees with Measurable Dose: <u>1693</u>		
Annual Collective Dose: <u>707</u>		

Manufacturing and Distribution Comments

- 1 or 2 rem yr<sup>-1</sup> will almost certainly increase the cost of radiopharmaceuticals produced in cyclotrons such as this facility.
- Special exposure limits may be needed for workers who produce isotopes with cyclotrons.
- We have several people with >50% extremity limit. If extremities are lowered by 2/5 as above, then we would have large expenses, ~\$50,000 for equipment.
- At a recent accelerator meeting, the subject of the economic effect of a 2 rem yr<sup>-1</sup> dose limit was informally discussed. The consensus was that positive ion cyclotrons, now commonly used, will not be economically feasible for radiopharmaceutical production with a 2 rem yr<sup>-1</sup> limit. Manufacturers are assuming that this limit will be in effect within several years and all new production machines will almost certainly be negative ion. The approximate cost of a negative ion machine is about \$5M, and there is a company that can convert the Cyclotron Corporation CS-30, a common production machine, to negative ion for a reported \$2.5M. I would expect 2 or 3 replacements and an equal number of conversions (if this proves feasible).
- The consideration of lowering the dose limits to 2 rem or lower could prevent nuclear pharmacists and scientists from carrying out the tasks vital to the nuclear medicine industry. It would force radiopharmaceutical companies to expend more resources on additional employees necessary to operate at current levels. Unfortunately, radiopharmaceutical companies have no choice but to pass on the additional costs to the consumer, raising the overall cost of nuclear medicine services. For many manufacturers the 2 rem yr<sup>-1</sup> limit would require major facility changes to production equipment and shielding. Under the 1 rem yr<sup>-1</sup>

**Questionnaire Results**

limit, the radiopharmaceutical industry would incur substantial cost necessary to maintain the flexibility and rapid handling of short lived radionuclides. This need would increase the costs both to the manufacturer and the user with no significant benefit.

- 6. NRC would also run the risk of states developing two sets of standards that are completely different. The first set would be for byproduct material regulated by the NRC and would reflect scientific claims not yet validated, about the effects of radiation. The second would be set by the states and reflect the use of radiation currently not

covered under NRC jurisdiction. The medical use of fluoroscopy, especially with cardiovascular procedures often gives worker film badge readings approaching the 5 rem y<sup>-1</sup> limit. This is also true of staff operating accelerators for radionuclide production and radiopharmaceutical synthesis. In order to stop large increases in health care costs from limitation of worker doses for no valid reason, such double standards may occur.

**5.7 Waste Management**

There were three responses, two from U.S. operators, the other from an operator outside the country; the latter indicated with an asterisk.

**Table 5.17 Impacts in Waste Management**

Impacts	Possible Dose Limit			
	2 rem y <sup>-1</sup>	1 rem y <sup>-1</sup>	5 rem y <sup>-1</sup> age in rem	2 rem y <sup>-1</sup> age in rem
<i>Collective dose:</i>				
No change	2	1	2	2
Increase	1*	1	-	-
Decrease	-	1*	1*	1*
% Increase	-	10	-	-
<i>Facility Modifications:</i>				
None required	1	1	2	1
Modifications Required	2*	2*	1*	2*
Costs (individual responses)	-	-	-	-
<i>Radiation Protection Cost:</i>				
No increase	-	-	2	-
Will increase	3	3	1*	3
Cost/yr (individual responses)	\$5K to \$1.2M*	\$10K to \$1.2M*	\$1.2M*	\$1.2M*

\* non-U.S.

Table 5.18 1989 Exposure Experience

<b>Number of Employees with Annual Doses:</b>			
>5 rem	<u>0</u>	>2 rem	<u>7</u>
>1 rem	<u>24</u>		
Number of Employees with Lifetime Dose Greater Than Age in Rem: <u>0</u>			
Number of Employees with Measurable Dose: <u>142</u>			
Annual Collective Dose: <u>77.3</u>			

Waste Management Comments

1. Radioactive waste management is generally changing from shallow land burial to engineered-at-grade disposal. Because of this basic change, we have used a design goal that the average radiation worker should not exceed 500 mrem yr<sup>-1</sup>.

2. These totals do not include exposure associated with waste processing services provided at generator locations.

**5.8 Fuel Fabrication, UF<sub>6</sub> Production**

There were two responses in this category.

Table 5.19 Impacts in Fuel Fabrication, UF<sub>6</sub> Production

Impacts	Possible Dose Limit			
	2 rem yr <sup>-1</sup>	1 rem yr <sup>-1</sup>	5 rem yr <sup>-1</sup> age in rem	2 rem yr <sup>-1</sup> age in rem
<i>Collective Dose:</i>				
No change	1	1	1	1
Increase	-	-	-	-
Decrease	-	-	-	-
% Increase	-	-	-	-
<i>Facility Modification:</i>				
None required	1	1	1	1
Modifications required	1	1	-	-
Cost (individual responses)	\$3.75M	\$6.25M	-	-
<i>Radiation Protection Cost:</i>				
No increase	1	1	-	-
Will increase	1	1	-	-
Cost/yr (individual responses)	\$0.45M	\$0.7M	-	-

## Questionnaire Results

**Table 5.20 1989 Exposure Experience**

<p><b>Number of Employees with Annual Doses:</b> &gt;5 rem <u>0</u>   &gt;2 rem <u>91</u>   &gt;1 rem <u>96</u> Number of Employees with Lifetime Dose Greater Than Age in Rem: <u>75</u> Number of Employees with Measurable Dose: <u>817</u> Annual Collective Dose: <u>545</u></p>
---

### Fuel Fabrication Comments

1. We are concerned that any reduction in the occupational dose may be a lever to lower the already ultra-conservative public dose limits - to what benefit?
2. The addition of external and internal exposure will increase these doses by a factor of ~10.
3. Implementation of lower dose limits would necessitate major modifications related primarily to modifying existing or installing new containments, upgrading ventilation systems, and re-engineering work processes, potentially involving complete refurbishment, or even replacement of the facility.
4. The effects of significant restructuring of radiation protection programs that occurred with implementation of new dose concepts in the revised Part 20 (i.e., combining internal and external dose as the effective dose equivalent) need to be evaluated.
5. The conservative factors and assumptions that underlie prior exposure records and current dose monitoring records raise significant issues.
6. There are technical difficulties and substantial resource burdens involved in reformatting prior exposure data (e.g., MPC-hours) in terms of less conservative, more realistic effective dose equivalent data.

## 5.9 Well Logging

These data came primarily from a member of the working group on the basis of a personal survey. One additional response to the questionnaire is included. The data are for 1988.

Table 5.21 Impacts in Well Logging

Estimated Impacts	Possible Dose Limit			
	2 rem y <sup>-1</sup>	1 rem y <sup>-1</sup>	5 rem y <sup>-1</sup> age in rem	2 rem y <sup>-1</sup> age in rem
<i>Collective dose:</i>				
No change	2	2	2	2
Increase	-	-	-	-
Decrease	-	-	-	-
<i>Facility Modifications:</i>				
None required	2	2	2	2
Modifications required	-	-	-	-
Cost (individual responses)	-	-	-	-
<i>Radiation Protection Cost:</i>				
No increase	2	2	2	2
Will increase	-	-	-	-
Cost/yr (individual responses)	-	-	-	-

Table 5.22 1988 Exposure Experience

<b>Number of Employees with Annual Doses:</b>			
>5 rem	<u>4</u>	>2 rem	<u>9</u>
>1 rem	<u>193</u>		
Number of Employees with Lifetime Dose Greater Than Age in Rem: <u>0</u>			
Number of Employees with Measurable Dose: <u>3,378</u>			
Annual Collective Dose: <u>-</u>			

Questionnaire Results

Well logging Comments

1. Approximately a third of the dose received by well loggers is from neutrons (QF=5). Few well logging technicians are over 40 years of age and the average tenure of a technician is about 10 years. Therefore, the 5 rem yr<sup>-1</sup> and lifetime would seem to be achievable, although more data are needed.

**5.10 Others (R&D, Regulatory)**

There are two responses included in this section. Although they have little in common, the impacts are quite similar, so a single presentation is considered acceptable.

**Table 5.23 Impacts in Others (R&D, Regulatory)**

Impacts	Possible Dose Limit			
	2 rem y <sup>-1</sup>	1 rem y <sup>-1</sup>	5 rem y <sup>-1</sup> age in rem	2 rem y <sup>-1</sup> age in rem
<i>Collective dose:</i>				
No change	2	2	2	2
Increase	-	-	-	-
Decrease	-	-	-	-
<i>Facility Modifications:</i>				
None required	2	2	2	2
Modifications required	-	-	-	-
Cost (individual responses)	-	-	-	-
<i>Radiation Protection Cost:</i>				
No increase	2	2	2	2
Will increase	-	-	-	-
Cost/yr (individual responses)	-	-	-	-

**Table 5.24 1989 Exposure Experience**

<b>Number of Employees with Annual Doses:</b>			
>5 rem	<u>  0  </u>	>2 rem	<u>  0  </u>
		>1 rem	<u>  0  </u>
Number of Employees with Lifetime Dose Greater Than Age in Rem: <u>  0  </u>			
Number of Employees with Measurable Dose: <u>  33  </u>			
Annual Collective Dose: <u>  2 rem  </u>			

## 6 High Dose Groups Within an Industry

### 6.1 Introduction

The data given in the tables in Section 5 do not reveal the potential impacts of lowered doses to selected categories of workers receiving higher annual doses than the average. Some indications of the importance of this issue appear in the comments of Section 5, particularly for the medical and nuclear-power communities. In medicine, particularly cardiology, angiography, and interventional radiology, reduction of dose limits might impact the availability of specialized medical attention.

### 6.2 NRC-Sponsored Study on High Dose Group Workers

In 1989, the NRC sponsored a BNL study of the distribution of dose as a function of special work groups in the nuclear-power industry (Khan et al., 1991a). Information was obtained from responses to a questionnaire addressing the following:

- a). What proportion of workers were getting higher than average dose;
- b). What was the magnitude of these doses;
- c). Are there any special, highly skilled work groups that are chronically getting the higher doses;
- d). Is there a shortage of skilled workers who are receiving higher than average doses?

Twenty-two nuclear power sites and six nuclear power contractor organizations responded. Among the power plant organizations responding, thirteen were pressurized water reactor (PWR) sites and nine were boiling water reactor (BWR) sites.

Table 6.1 shows the whole-body dose data for one year for the PWR plants in this group; Table 6.2 shows the data for BWR plants. The data cover the total number of persons monitored at the plant, including contractors.

Table 6.1 Whole-Body Dose Data for PWR Plants for 1988

Plant (units)	Total Number monitored	Number of Persons with Annual Whole-Body Dose				Average Dose Per worker (rem)
		> 1 rem		> 2 rem		
		Persons	%	Persons	%	
PW1 (3)	3,841	237	6.2	24	0.6	0.11
PW2 (2)	4,446	606	13.6	164	3.7	0.24
PW3 (2)	2,234	8	0.4	0	0	0.06
PW4 (1)	2,519	53	2.1	2	0.1	0.26
PW5 (2)	2,943	93	3.2	6	0.2	0.19
PW6 (2)	759	-	-	0	0	0.10
PW7 (2)	3,290	481	14.6	80	2.4	0.33
PW8 (3)	374	166	4.4	10	0.3	0.11
PW9 (2)	1,446	76	5.3	5	0.3	0.32
PW10 (1)	1,975	272	13.8	60	3.0	0.50
PW12 (1)	1,984	18	0.9	18	0.9	0.23
PW13 (1)	1,279	28	2.2	1	0.1	0.21

High Dose Groups

Table 6.2 Whole-Body Dose Data for BWR Plants for 1988

Plant (units)	Total Numbered Monitored	Number of Persons with Annual Whole-Body Dose				Average Dose per worker (rem)
		> 1 rem		> 2 rem		
		Persons	%	Persons	%	
BW1 (2)	1,684	28	1.7	5	0.3	0.33
BW3 (1)	4,887	68	1.4	7	0.1	0.19
BW4 (1)	2,265	302	13.3	63	2.8	0.51
BW5 (2)	2,616	316	12.1	22	0.8	0.28
BW6 (2)	3,957	1,073	27.1	326	8.2	0.45
BW7 (2)	3,727	569	15.3	69	1.9	0.29
BW8 (3)	10,322	862	8.4	201	1.9	0.28
BW9 (1)	3,215	612	19	148	4.6	0.52

Both tables show that the average dose per worker is only a small fraction of the present annual whole-body dose limit. In addition, only a small percentage of workers (from 0.1 to 8%) are getting doses greater than 20 mSv (2 rem) annually.

The PWR data for 1988 (Table 6.3) show that workers had annual doses above 20 mSv, and 76

have lifetime doses (in rem) greater than their age. Such workers are maintenance technicians, welders, riggers, millwrights, and assorted contract personnel. Most of the 76 persons from the high-dose groups in the dose greater than age category were maintenance technicians and other contract personnel.

Table 6.3 Whole-Body Dose Data for Various Worker Groups at PWR Plants for 1988

Work Group	Number with dose		
	Annual		Lifetime
	>1 rem	>2 rem	>age
Maintenance Techs	178	23	20
Boiler Makers	26	5	2
Welders	119	24	0
Health Physics Techs	127	10	6
Pipe fitters	75	11	0
Riggers	255	61	5
Millwrights	237	49	2
Fuel Handlers	39	11	0
Decon Workers	36	7	0
Other Contract Personnel	181	85	41
<b>Total</b>	<b>1273</b>	<b>286</b>	<b>76</b>

Table 6.4 Whole-Body Dose Data for Various Worker Groups at BWR Plants for 1988

Work Group	Number with Dose		
	Annual		Lifetime
	> 1 rem	> 2 rem	> age
Pipe fitters	83	23	0
Health Physics Techs	188	8	7
Millwrights	1154	418	1
Boiler Makers	15	2	0
Riggers	19	1	0
Maintenance Techs	277	18	54
I & C Techs	85	13	0
Quality Assurance	28	2	2
Radwaste Handlers	18	3	1
Other Contract Personnel	277	100	2
<b>Total</b>	<b>2144</b>	<b>588</b>	<b>67</b>

## High Dose Groups

For BWRs (Table 6.4). 588 workers are getting annual doses above 20 mSv. However, the number of persons whose lifetime dose is greater than their age is less than for PWRs, 67 workers. Almost all workers getting doses greater than 20 mSv yr<sup>-1</sup> are in two groups, millwrights and other contract personnel. The preponderant proportion of the 67 persons with lifetime dose greater than age are maintenance technicians.

### 6.2.1 Analysis of Dose Data Obtained in the Study

Table 6.1 shows that for some PWRs nearly 15% of the persons monitored are likely to receive > 1 rem yr<sup>-1</sup>. Because the number monitored implies anyone who is issued a radiation badge, and therefore, typically includes all visitors, engineering, and management personnel, the number with annual dose greater than 1 rem as a proportion of the actual radiation workforce is likely to be even higher than 15%. Table 6.2 shows the proportion of persons getting annual doses greater than 1 rem, which may range up to 27% for BWR plants. Contract personnel are included in these sets of numbers for the two plant types. However, the data do not reflect the total doses to transient workers getting dose at several sites.

Correlations with other factors were made for plants listed in Tables 6.1 and 6.2 in which more than 10% of the workers had doses greater than 1 rem yr<sup>-1</sup>. We found no correlations with power rating, vendor, multiple vs. single plant sites, utilities with several plants vs. those with one or two nuclear units, and the date the plant went into service.

Only three PWR plants and three BWR plants reported that more than 2% of people were getting doses above 2 rem yr<sup>-1</sup>. Again, since this value is based on all who were issued a radiation badge, the number with dose greater than 2 rem yr<sup>-1</sup> as a proportion of the actual radiation workforce will be greater. Every plant reported an average dose per worker of less than 0.5 rem yr<sup>-1</sup>.

#### 6.2.1.1 Pressurized Water Reactor Data

Typically, between 100-200 people per reactor unit at PWR plants had doses above 1 rem yr<sup>-1</sup> is. However, PWR2 (Table 6.1) with 2 units reported over 300 persons per unit with doses above 1 rem yr<sup>-1</sup>. The

number of health physics technicians and maintenance technicians with doses above 1 rem yr<sup>-1</sup> per plant are generally in double digits. The same is true for welders, millwrights, and riggers, except for 2 sites with more than 100 people getting doses above 1 rem yr<sup>-1</sup>.

Compared to 1 rem yr<sup>-1</sup>, there is a five-fold decrease in the total number of persons getting doses above 2 rem yr<sup>-1</sup>. This is also reflected in the doses to individual work groups. The number of workers in any work group having more than 2 rem yr<sup>-1</sup> is appreciably less than the number of workers with more than 1 rem yr<sup>-1</sup>. The number with lifetime dose greater than age is a further factor of 4 lower than the number with annual dose above 2 rem.

The average dose to each work group is generally less than 1 rem yr<sup>-1</sup>, except at PWR 7 with two units, where boiler makers, welders, riggers, and electrical technicians are getting higher doses, and at PWR 10, where welders are getting an average dose of slightly more than 1 rem. The craft workers receiving average annual doses >2 rem are typically millwrights, pipe fitters, maintenance, and inspection & control technicians.

#### 6.2.1.2 Boiling Water Reactor Data

Typically, 700 persons at BWRs get annual whole-body doses greater than 1 rem, which is higher than for PWRs. Up to about 100 per unit get annual doses greater than 2 rem. Up to 30 have lifetime doses greater than age.

The craft workers receiving annual average whole-body doses greater than 1 rem are typically millwrights, health physics technicians, maintenance technicians, pipe fitters, and instrumentation and control technicians.

#### 6.2.1.3 Contractor Data

Significantly more persons with higher doses were expected from the nuclear power plant contractors. However, although the numbers were larger than for nuclear power plant workers, they are not significantly different. In fact, one PWR contractor showed some of the lowest dose data.

Despite the good results for one contractor, both PWR and BWR contractors reported hundreds of people

with doses greater than 1 rem yr<sup>-1</sup>. One major PWR contractor reported over 300 people with dose over 2 rem yr<sup>-1</sup>; however, in all other cases, the number was less than 60. Once again, the lifetime dose less than age was less frequently exceeded; only 2 contractors reported double digit figures (14 for one, 51 for the other).

The average dose for each craft can be used to determine the work groups that are receiving the higher doses. For contractors, the groups that get an average annual dose greater than 1 rem included maintenance technicians, riggers, electrical technicians, station men, radwaste handlers, and quality-assurance technicians.

## 7 Costs Associated With Dose Reduction Modifications in the Nuclear Power Industry

### 7.1 Introduction

One of the more difficult aspects of projecting the impacts of dose reduction is the estimate of costs. The ALARA Center at Brookhaven National Laboratory has been compiling and evaluating the cost and resulting dose reduction for a wide variety of reactor plant modifications since the early 1980s. The data selected for presentation indicate the basis for many of the cost estimates given in Tables 5.3 and 5.5. NUREG/CR 4373 (Baum, 1985) describes the approach taken to obtain the listed values and gives additional examples.

### 7.2 Costs (and the Related Dose Saved) of Selected Modifications Which Might be Employed to Reduce Exposure

The following list is taken from NUREG/CR 4373 and contains examples of items with a cost-effectiveness of \$10 per person-Sv (\$1,000 per person-rem) or less. Examples of less cost-effective modifications also can be found in this report.

**Table 7.1 Estimated Costs and Dose Savings for Modifications at Nuclear Power Plants  
(Baum, 1985)**

Modification	Dose Saved person-Sv (person-rem) *	Capital Cost (\$)**
PWR Refueling Machine (New Plant, on Critical Path)	.9 (90)	220000
PWR Reactor Vessel Head Multi-Stud Tensioner-/Detensioner (Two Reactor Site, on Critical Path)	16 (1,600)	940000
PWR Reactor Vessel Head Multi-Stud Tensioner-/Detensioner (Single Reactor Site, on Critical Path)	7.9 (790)	940000
PWR Integrated Head Assembly (New Plant, on Critical Path)	1.2 (120)	75000
Multi-Stud Tensioners/Detensioners for PWR Reactor Pressure Vessel (on Critical Path)	2.4 (240)	600000
PWR Reactor Vessel Head Tensioner/Detensioner (on Critical Path 25% of Time)	3.6 (360)	340000
Steam Generator Channel Head Decontamination (Not on Critical Path)	37 (3,700)	2145191

\* Dose savings accumulated over the useful period for the item (typically 30 years).

\*\* In 1984 dollars. Includes the cost of replacement power for modifications that affect critical path time.

Table 7.1 Continued

Modification	Dose Saved person-Sv (person rem)*	Capital Cost (\$)**
Reactor Cavity Decontamination Using the WEPA Cleaning System	.48 (48)	89,000
BWR Control-Rod-Drive-Handling Tool (on Critical Path 25% of Time)	9.4 (940)	325,000
PWR Reactor Vessel Head Tensioner/Detensioner (on Critical Path 25% of Time)	4.2 (420)	349,000
PWR Reactor Vessel Head Tensioner (on Critical Path 25% of Time)	9.6 (960)	349,000
Shredder-Compactor for Dry Active Waste	2.6 (260)	450,000
Robotics System for Remote Inspections of BWR Moisture Separator and Feedwater Pump Areas (Three Reactor Site)	21 (2,100)	66,700
PWR Quick Opening Fuel Transfer Tube Closure (New Plant, on Critical Path)	.15 (15)	1,500
Remote Readout Near PWR Seal Table	.59 (59)	89,000
PWR Steam Generator Manway Tensioner/Detensioner and Handling Device (on Critical Path 25% of Time)	4.4 (440)	500,000
Photographic Technique for PWR Steam Generator Tube Plugging Inspections	16 (1,600)	5,000
PWR Steam Generator Manway Tensioner/Detensioner	.9 (90)	133,000
Robotic Inspection of PWR Ice Condenser Area	1.5 (150)	100,000
Solid Radioactive Waste Handling Using High Integrity Containers	.51 (51)	150,000
Robotics System for Inspections in BWR Moisture Separator and Feedwater Areas (Single Reactor Site)	7 (700)	65,900
Robotic Mechanism for Surveillance of BWR High Pressure Feedwater Heater Rooms (Three Reactor Site)	1.2 (120)	22,400

Costs

Table 7.1 Continued

Modification	Dose Saved Person-Sv (person-rem)*	Capital Cost (\$)**
Portable Robotic System for Smoke Detector Inspection (Three Reactor Site)	1.4 (140)	20,000
Robotic Mechanism for Surveillance of BWR High Pressure Feedwater Heater Rooms (Single Reactor Site)	.39 (39)	20,800
Portable Robotics System for Smoke Detector Inspection (Single Reactor Site)	58 (5,800)	20,000
BWR-CRD Scram Discharge Line Flange for Hydro-lazing the Header	2.95 (295)	4,000
Portable Shielding System for the PWR Steam Generator Channel Heads	14.9 (1,490)	50,000
Shielding for CVCS Demineralizers (Option B)	.30 (30)	1,300
Clean Seal Cooling Water Supply for BWR Recirculation Pump	5.95 (595)	25,000
PWR Power Level Monitor Using <sup>16</sup> N Detectors	2.4 (240)	15,000
Cobalt Replacement in PWR Reactor Coolant Pumps (Three Loop Operating Plant, Pumps Replaced for Other Reasons)	5.6 (560)	30,000
Shielding for CVCS Demineralizers (Option A)	.51 (51)	2,600
Replacement of PWR Steam Generators with Low-Cobalt (<0.03%) Tubing (Three-Loop Operating Plant, Replacement Needed for Other Reasons)	35 (3,500)	198,000
Replacement of PWR Control-Rod-Drive Mechanisms with Low-Cobalt Parts (Three-Loop Operating Plant, Replacement Needed for Other Reasons)	8.1 (810)	50,000
Replacement of PWR Steam Generators with Those Having Low-Cobalt (0.015%) Tubing (Three-Loop Operating Plant, Replacement Needed for Other Reasons)	47 (4,700)	300,000
Replacement of PWR Control-Rod-Drive Mechanisms with Low Cobalt Parts (Four-Loop Operating Plant, Replacement Needed for Other Reasons)	7.7 (770)	50,000

Table 7.1 Continued

Modification	Dose Saved person-Sv (person-rem)*	Capital Cost (\$)**
Replacement of PWR Steam Generators Using Low-Cobalt (<0.03%) Tubing (Four-Loop Operating Plant, Replacement Needed for Other Reasons)	37 (3,700)	264,000
Replacement of PWR Steam Generators with Those Having Low-Cobalt (0.015%) Tubing (Four-Loop Operating Plant, Replacement Needed for Other Reasons)	50 (5,000)	400,000
Temporary Shielding for PWR Reactor Vessel Head	.88 (88)	1,500
Low-Cobalt Specifications for PWR Fuel Assembly Nozzles (New Plant)	.93 (93)	10,230
Cobalt Replacement in PWR Reactor Coolant Pumps (Four-Loop Plant, Pumps Replaced for Other Reasons)	3.2 (320)	30,000
TV Robot Inspection of PWR Vessel Head (Single Reactor Site)	2.7 (270)	19,000
Reduce Cobalt Impurity in New PWR Steam Generator Tubing (Sizewell 'B' Plant)	2700	330,000
Handling Equipment for PWR Steam Generator Manway Covers	.45 (45)	5,600
Mock-Up Training for PWR Steam Jobs	29 (2,900)	60,000
Installation of Viewing Windows in BWR Plants	2.24 (224)	25,000
PWR Reactor Pressure-Vessel Head Laydown Shield	.9 (90)	15,000
BWR Control Rod Drive Disassembly and Decontamination Tank	2.68 (268)	35,000
Permanent Shield for PWR Reactor Vessel Head (Three Reactor Site)	8.9 (890)	185,000
Electropolishing Tank for BWR Control Rod Drives	2.99 (299)	40,000
Relocation of Instrument Readout at PWR Spent-Fuel Pit Heat Exchanger	.13 (13)	2,500

Costs

Table 7.1 Continued

Modification	Dose Saved person-Sv (person-rem)*	Capital Cost (\$)**
Helium Leak Detection for BWR Condenser Tubes	1.8 (180)	25,000
Relocation of Fuel Sipping Cans	.3 (30)	5,000
Shielding for PWR Reactor Upper Internals (Two Reactor Site)	.84 (84)	19,500
Ultrasonic Testing of PWR Pressurizer Surge Line (Three Reactor Site)	.17 (17)	8,000
PWR Reactor Vessel Head Shielding (No Critical Path Expense)	2 (200)	65,321
PWR Steam Generator Tube Inspection and Repair Robot	23 (2,300)	450,000
BWR Pipe Insulation Improvements for In-Service Inspections	3.9 (390)	100,000
TV Monitor for BWR Cleanup Heat Exchanger Room	2.5 (25)	7,000
BWR Control Rod Drive Handling Tool	9.4 (940)	325,000
PWR Reactor Vessel Head Shielding (On Critical Path)	2 (200)	95,321
Acoustic Emission Instrumentation for ISI of the Reactor Vessel and Reactor Coolant Piping	13 (1,300)	450,000
Decontamination of a BWR Recirculation System	9 (900)	750,000
Air-Cooled Anticontamination Suit, Radio Dosimetry, and Radio Communications	1.5 (150)	56,000
Shielding for PWR Reactor Upper Internals (Single Reactor Site)	.41 (41)	19,500

\* Dose savings accumulated over the useful period for the item (typically 30 years).

\*\* In 1984 dollars. Includes the cost of replacement power for modifications that affect critical path time.

### 7.3 Estimated Impacts

During the 1980s, considerable efforts were made by the nuclear industry to reduce collective and individual doses at nuclear power plants. This effort was stimulated by several factors, including anticipated lowering of dose limits to conform with the 1977 ICRP recommendations, the reassessments of risks based on new dosimetry and epidemiological data on the Japanese survivors of the World War II atomic weapons, and anticipated further restrictions on annual and lifetime dose limits.

These pressures led the U.S. utilities to expend significant sums on dose control modifications of the type illustrated in Table 7.1. The judgements on cost-effectiveness were generally based on a valuation of the dose avoided, that was in the range of a few hundred thousand dollars to about \$2.6 million per person-Sv saved (Baum, 1991). Figure 7.1 summarizes the values employed at nuclear power plants in 1991-1992 (Kindred, 1992).

These high monetary values of the cost or value of dose savings were based primarily on the costs of hiring additional workers that were necessitated by lower administrative dose limits. For example, a worker hired at a cost of \$53,000<sup>2</sup> per year who might be permitted only 40 mSv (4 rem) (typical administrative limit) exposure per year leads to a cost of dose avoided of \$53,000/0.04 Sv = \$1,325,000 person-Sv (\$13,250 per person-rem). Not all workers would be near the administrative limits and a worker's productivity may not drop to zero when the limit is reached, so the adopted value of cost for dose avoided for a particular job or plant is usually less (e.g. average = \$434,300 per person-Sv (\$7,343 per person-rem on Figure 7.1).

Figure 7.2 shows the total number of reactors and total collective dose for commercial nuclear plants from 1973 through 1989 (Hinson, 1992). While the number of reactors increased from 68 in 1980 to 112 in 1992, the collective dose decreased from about 540 person-Sv (54,000 person-rem) to about 280 person-Sv (28,000 person-rem); or collective dose per reactor decreased from about 7.94 person-Sv (794 person-rem yr<sup>-1</sup>) in 1980 to about 2.5 person-Sv yr<sup>-1</sup> (250 person-rem yr<sup>-1</sup>) in 1991. Assuming this reduction was at an average cost of \$700,000 per person-Sv (\$7,000 per person-rem), the cost is about 544 (794-250) person rem per reactor per year x \$7,000 per

person rem = \$3,808,000 per reactor per year. Many dose reduction efforts in the past did not require the \$700,000 per person-Sv (\$7,000 per person-rem) expenditure. However, because many of the less costly modifications have already been implemented, it is likely that future reductions will require the higher expenditure. Thus, for the nuclear power industry one can anticipate that the impact of any lower dose rates are likely to be proportional to the product of the collective dose being received above the new limit and about \$700,000 per person-Sv (\$7,000 per person-rem).

Table 4.5 shows that there were 8,845, 1,290, 121, and 11 persons in 1989 who received between 1.0-2.0, 2.0-3.0, 3.0-4.0, and 4.0-5.0 rem, respectively. The collective dose above 1 rem yr<sup>-1</sup> received by these individuals is estimated as 16,420 person-rem, assuming that the average dose for each group is equal to the midpoint for that dose range (e.g. average dose for the 1.0-2.0 dose range is 1.5 rem).

If a 10 mSv (1 rem yr<sup>-1</sup>) limit were imposed, it would require a collective dose reduction of 164.2 person-Sv (16,420 per person-rem). This would cost:

$$\frac{164.2 \frac{\text{person-Sv}}{\text{yr}} \times \frac{700,000}{\text{person-Sv}}}{112 \text{ reactors}} = \frac{\$1,026,000}{\text{reactor year}}$$

For a 20 mSv yr<sup>-1</sup> (2 rem yr<sup>-1</sup>) limit, the required collective dose reduction would be about 33.69 person-Sv yr<sup>-1</sup> (3,369 person-rem yr<sup>-1</sup>). This would cost:

$$\frac{33.69 \frac{\text{person-Sv}}{\text{yr}} \times \frac{700,000}{\text{person-Sv}}}{112 \text{ reactors}} = \frac{\$210,000}{\text{reactor year}}$$

The impact of imposing an "age x 1" limit on workers cumulative effective dose is difficult to judge from the limited data. Two estimates are made to indicate a likely range.

<sup>2</sup> Fully loaded cost for operating and maintenance personnel expressed in 1984 dollars including all fringe benefits, but not including overhead and general and administrative expenses (Ball, et al, 1984).

Costs

# Values of Dose Avoided

Compiled by G.W.Kindred (1992)




Figure 7.1 Person-Rem Values

Total Number of Reactors and Collective Dose




Figure 7.2 Total Number of Reactors and Collective Dose

The first estimate is based on the number of workers who exceed the age limit in the high dose groups in the survey of 22 reactors (Section 6). There were 76 individuals in the 13 PWRs and 67 in the 9 BWRs who exceeded the age limit, a total of 143. Assuming each of these workers were replaced at an annual cost of \$53,000, and that these replacements were sufficient to provide the crews needed to work under new limits, the annual cost per reactor would be about:

$$\frac{143 \times \$53,000}{22} = \frac{\$344,500}{\text{reactor year}}$$

Since the replaced workers would be useful for other work (not high dose), this cost estimate is an upper limit.

The second estimate assumes that workers currently exceeding their age limit would be given an exception to the age rule (a "grandfathering" clause) and would stay below either 10 mSv (1 rem) yr<sup>-1</sup> or 2 rem yr<sup>-1</sup> limit. The cost was estimated by considering the cost of implementing these limits and the number of workers affected for each limit in the sample survey of Section 6.

The number of workers exceeding 10 mSv yr<sup>-1</sup> (1 rem yr<sup>-1</sup>) and 20 mSv yr<sup>-1</sup> (2 rem yr<sup>-1</sup>) were 3,417 and 874, respectively. Assuming the replacement worker and dose reduction costs per worker are the same for those exceeding the age limit and those exceeding the 10 mSv (1 rem) and 20 mSv (2 rem), annual limits, the costs for an age limit with "grandfathering", can be estimated from the ratios of workers in the various groups and the earlier cost estimates.

Compared to the 10 mSv (1 rem) yr<sup>-1</sup> cost estimates:

$$\frac{143}{3417} \times \$1,026,000 \approx \frac{\$43,000}{\text{reactor year}}$$

Compared to the 20 mSv (2 rem) yr<sup>-1</sup> cost estimates:

$$\frac{143}{874} \times \$210,000 \approx \frac{\$34,000}{\text{reactor year}}$$

These two estimates are nearly equal and can be rounded to about \$40,000 per reactor per year for a 50 mSv yr<sup>-1</sup> limit with an "age x 1" (rem or 10 mSv) cumulative limit.

## Costs

In summary, the estimated cost impacts on nuclear power plant operations for the three dose limit options considered are (rounded to one significant figure):

<u>Option</u>	<u>Estimated Cost Per Year Per Reactor</u>
10 mSv (1 rem) yr <sup>-1</sup> limit	≈ \$1,000,000
20 mSv (2 rem) yr <sup>-1</sup> limit	≈ \$ 200,000
Age 10's of mSv (Age x 1) with a	
50 mSv (5 rem) yr <sup>-1</sup> limit	≈ \$ 40,000 with "grandfathering" ≈ \$ 300,000 without "grandfathering"

It should be recognized that the general approach to estimating impacts is limited and prone to longer uncertainties. As noted above, "many of the less costly modifications have already been implemented (and) it is likely that future reductions will require the higher expenditure." In light of this, the \$/person-rem values currently being used by the industry may be viewed as representing the lower end of the range of expenditures that would be needed to reduce doses to comply with lower dose limits. This is especially likely to be the case for higher dose jobs that would be most impacted by lower dose limits because these jobs have already been the primary focus for industry dose reduction efforts. In addition, present actions are primarily aimed at keeping worker doses ALARA, including minimization of collective and individual doses which involves much discretion and flexibility in application, with due consideration of limitations on resources available for dose reduction. This is the essence of "reasonably achievable," i.e., social and economic factors being taken into account.

On the other hand, dose reduction actions required in response to lower dose limits in regulation would suffer from significantly less ability to exercise discretion and flexibility because achieving reduced doses to comply with regulatory limits would be mandatory, regardless of limitations on available resources. Therefore, the relative cost impact of such expenditures could be much greater than in the past.

The variation between plants in values of dose avoided, as given in Figure 7.1 and in the collective dose, suggest that the impacts of reduced dose limits must be plant specific. This would mean that the values obtained in Section 7.2 may be more realistic than those obtained by the generalized approach given in Section 7.3.

## 8 Summary

As in Section 5, the conclusions will be given by practice and industry type, followed by a general conclusion. In general, the options used in the questionnaire will guide this presentation except that the 20 mSv yr<sup>-1</sup> (2 rem yr<sup>-1</sup>), coupled with the age limitation option, will not be used, because it differs very little with a 20 mSv yr<sup>-1</sup> (2 rem yr<sup>-1</sup>) limit.

### 8.1 Medical/Dental/Veterinary

#### 8.1.1 1 Rem Yr<sup>-1</sup>

Although several issues raised in the comments reflected a general feeling that there was no biological need for reducing the dose limit, UNSCEAR (UNSCEAR, 1988), BEIR (NAS BEIR, 1990), and ICRP (ICRP, 1991) indicated differently. Overall, the estimated costs were moderate, even with the most severe limitation of 10 mSv yr<sup>-1</sup> (1 rem yr<sup>-1</sup>). However, selected occupational groups within the medical community would be severely impacted, specifically, cardiologists and interventional radiologists. In fact, one comment suggested that their exposure may be underestimated due to the lack of compliance with personal monitoring procedures.

#### 8.1.2 2 Rem Yr<sup>-1</sup>

The vast majority of respondents considered that 20 mSv yr<sup>-1</sup> (2 rem yr<sup>-1</sup>) was attainable, although there were clearly costs associated with this option. Significantly, the assessment of dose was raised by several respondents. It is still too often the case that the exposure to the badge worn on the collar by an individual wearing a lead apron is used for determining compliance with dose limits. UNSCEAR (UNSCEAR, 1988) suggests that in diagnostic radiology the dosimeter usually overestimates the effective dose by about 2-4. Although this issue needs to be addressed, it gives some support to the suggestion that the impact of lowering the doses would not be severe if the dose were assessed appropriately. Anticipated NCRP guidance on the effective dose equivalent from partial body exposure may resolve some of these issues.

There was a comment that better training of selected medical personnel could reduce their exposure at little additional cost.

#### 8.1.3 5 Rem Yr<sup>-1</sup> and Cumulative Dose in Rem Less Than Age in Years Limit

The respondents were unanimous in their assessment that 50 mSv yr<sup>-1</sup> (5 rem yr<sup>-1</sup>), coupled with the restriction on the cumulated dose in 10s of mSv (or in rem) not exceeding their age in years, is without serious impact. We note that Figure 4.15 indicates that women workers in medicine appear to receive more radiation as they grow older. However, due to low average exposure to medical workers, the age limitation on dose should not have an important impact.

This category does not include those medical workers whose job consists of making cyclotron-produced radiopharmaceuticals; they are treated in Section 8.5 on manufacturing and distribution.

### 8.2 Nuclear Power Reactor Plants and Their Contractors

There are many sources of information on this category of workers. For this report, the following studies were reviewed: The EEI study, which was based on 27 responses to a survey; the BNL high-dose group study, which was based on 22 power plant site responses; the questionnaire results, based on 18 responses; the NRC REIRS data, which provides dose distribution data on nuclear power and contractor workers; the 1984 EPA report, which examined the available dosimetric data from a variety of view points, such as cumulative exposure as a function of age and sex, and the extensive comments received on the draft NUREG/CR-6112. The data given in NUREG/CR-4373 were used to evaluate the cost estimates.

#### 8.2.1 1 Rem Per Yr<sup>-1</sup>

A 10 mSv yr<sup>-1</sup> (1 rem yr<sup>-1</sup>) dose limit would have enormous impacts in the nuclear power industry, even to the point of being impossible without unreasonable costs for most existing facilities. The REIRS data for 1990 given in Table 4.5 indicate that nearly 10% of the LWR workers with measurable exposure exceeded 10 mSv yr<sup>-1</sup> (1 rem yr<sup>-1</sup>), as does Table 5.4 from the questionnaire. Tables 6.1 and 6.2 indicate that in the 22 nuclear power plant sites participating in the high-dose worker study, nearly 6,000 workers had annual exposures exceeding 10 mSv (1 rem) in 1988.

## Summary

As expected, this is even more critical for craft workers, as shown in Tables 6.3 and 6.4 from the BNL high-dose group study. For the 22 plant sites, there are 3,500 craft workers in high-dose groups with an annual exposure over 10 mSv (1 rem). For example, there were 1,400 millwrights, 455 maintenance techs, and 315 health physics techs all in excess of 10 mSv yr<sup>-1</sup> (1 rem yr<sup>-1</sup>).

This should not be taken to mean that the next generation of nuclear power plants cannot be designed to operate with exposure below 10 mSv yr<sup>-1</sup> (1 rem yr<sup>-1</sup>), but with the current plants it is unlikely to be economically feasible. For example, the EEI Report found that at this level, "all responders felt operations would be extremely difficult, if not impossible" (EEI 1991).

### 8.2.2 2 Rem Yr<sup>-1</sup>

A limit of 20 mSv yr<sup>-1</sup> (2 rem yr<sup>-1</sup>) would also appear to be very difficult to achieve for the nuclear power industry, although just over 1000 workers exceed 20 mSv yr<sup>-1</sup> (2 rem yr<sup>-1</sup>), (Table 4.5 from the REIRS data and Table 5.5). Figures 4.1 and 4.2 show that the utility personnel (UT) would not be as severely affected as the contractors (TO). The high-dose group study also indicates that the craft groups would, again, have the highest percentage of people exceeding 20 mSv yr<sup>-1</sup> (2 rem yr<sup>-1</sup>). The responses to the questionnaire indicate that the impact at 20 mSv yr<sup>-1</sup> (2 rem yr<sup>-1</sup>) would be about half that at 10 mSv yr<sup>-1</sup> (1 rem yr<sup>-1</sup>), but still several million dollars per plant in capital costs, nearly half a million dollars per plant in annual costs, and a 2 to 100% increase in collective dose.

The greatest diversity was seen here among responders. For utilities which do not perform their own major maintenance, the impact is not too great. For utilities that do, and for contractors supplying skilled craft workers, the impact is far greater. Questions were raised in the EEI Report, the questionnaires, and re-emphasized in the comments to the draft NUREG/CR-6112 about the availability of skilled personnel at this dose limit. Even utilities who felt they could live with a 20 mSv yr<sup>-1</sup> (2 rem yr<sup>-1</sup>) limit noted that additional personnel would be needed. For this dose limit option the issue is practicality. Unlike the 10 mSv yr<sup>-1</sup> (1 rem yr<sup>-1</sup>) limit, it would be possible, but expensive, both in capital cost and in increased collective dose. Many more skilled craft workers would be needed to work on vital safety systems, yet the supply is already

limited. At such a dose limit, there might be potentially serious impacts on safety since some discretionary inspections and maintenance might be constrained. In general, such a dose limit would require an extensive change in scheduling and in the way modifications are made and maintenance is done. System decontamination, remote tooling, and robots would be essential. This can be summed up by one of the comments from Section 5.2, "Two rem/yr would be difficult and costly but achievable for utility workers. However, for contract personnel, it would be very difficult and exceedingly costly."

As discussed in the comments to Section 5.2, the impact should be addressed with the understanding that the plants use administrative dose limits of ~80% of the regulatory limit.

### 8.2.3 5 Rem Yr<sup>-1</sup> and Cumulative Dose in Rem Less Than Age in Years Limit

The REIRS data as given in Table 4.5 for 1990 indicates that none of the LWR workers exceeded 50 mSv yr<sup>-1</sup> (5 rem yr<sup>-1</sup>). The REIRS does not contain cumulative dose data. This study's questionnaire data indicates virtually no impact under this limit; although, the comment given in Section 5.2 notes that some sort of grandfathering is needed - "For those individuals who would exceed the lifetime limit of age in rem, a 2 rem/year limit would be necessary to maintain their employment in the industry." The need for "grandfathering" also is shown in Table 5.5 which indicates that in 1993, ~500 workers exceeded their age in rem, and in Tables 6.3 and 6.4 on the high-dose groups where 143 workers are shown to exceed their lifetime limit. The data in the EEI Report suggests a concern about contractor availability.

There is a hidden aspect of the age in rem limit. It is noteworthy that, in effect, the worker will have to average less than 15 mSv yr<sup>-1</sup> (1.5 rem yr<sup>-1</sup>) over the working lifetime. This is somewhat ameliorated by data from the EPA Report that suggests that for males (most workers in the nuclear industry) the average exposure received each year decreases with age after age 42 (Table 4.6). For female workers (primarily in medicine) there is no decrease, although their mean annual dose is less than half that of male workers.

The 50 mSv yr<sup>-1</sup> (5.0 rem yr<sup>-1</sup>) and age in 10s of mSv (rem) limit together with a "grandfather clause," which

permits 20 mSv yr<sup>-1</sup> (2 rem yr<sup>-1</sup>) after exceeding the age limits, seems acceptable, because it would have very little impact on either the industry or the individual worker. Such a "grandfathering" exception would have to be closely controlled, since the risk to such workers could conceivably be in excess of the risk of accidental death of workers in more hazardous industries in the United States.

### 8.3 Test and Measurement Including Industrial Radiography

The data were obtained from responses to the questionnaire, from the REIRS, and from discussions in the working committee. Table 4.2, from the REIRS data, shows a substantial difference in the dose received between single versus multiple locations. The data given in Table 5.9 seems to reflect the status for multiple locations. The protection problems are more variable with multiple locations and the potential for unintended exposure is greater.

As can be seen in the comments section, there is concern that the NRC 10 CFR Part 34 regulation needs to be assessed before adequate judgement can be made.

In Section 4.5.5, the authors noted that the higher doses now being reported in the fuel fabrication industry reflect a change in the reporting requirements rather than an increase in exposure. They also pointed out that there is a need to assess dose trends using the requirements of the revised Part 20 before the impact of lower doses can be assessed in this industry.

There is an expectation expressed in a response to the draft NUREG/CR-6112, as given in the comments of Section 5.4, that any reduction in the dose limits will increase collective dose, the size of the workforce, the cost of facilities and equipment, and the cost of radiation protection.

#### 8.3.1 1 Rem Yr<sup>-1</sup>

At a 10 mSv yr<sup>-1</sup> (1 rem yr<sup>-1</sup>) limit, there will be moderate increases in collective dose and in cost for both modification and operating radiation protection programs (Table 5.9). There is concern that worker

training and productivity would suffer. The responders indicated (Table 5.10) that about 10% of the employees with measurable dose received exposures in excess of 10 mSv yr<sup>-1</sup> (1 rem yr<sup>-1</sup>), while the data reported in the response to the draft NUREG/CR-6112, as given in Table 5.11, suggest that >75% of the employees exceeded 20 mSv yr<sup>-1</sup> (2 rem yr<sup>-1</sup>).

#### 8.3.2 2 Rem Yr<sup>-1</sup>

Although most thought they could operate with this option, the comments received on draft NUREG/CR-6112 suggested there might be impacts like those given for 1 rem yr<sup>-1</sup>. The data for 1989 (Table 5.10) indicates that 4% of the workers exceed 20 mSv yr<sup>-1</sup> (2 rem yr<sup>-1</sup>), while the data reported in the response to the draft NUREG/CR-6112, as given in Table 5.11, suggest that greater than 60% exceeded 20 mSv yr<sup>-1</sup> (2 rem yr<sup>-1</sup>).

#### 8.3.3 5 Rem Yr<sup>-1</sup> and Cumulative Dose in Rem Less Than Age in Years Limit

For this option, there would be no impact expected in collective dose, facility modification, or radiation protection. This suggests that many of the higher annual doses were inadvertent, and that the same individuals were unlikely to receive such exposure very often during their working lifetime.

### 8.4 Universities not Including Medical, Dental, or Veterinary Schools

The data here were obtained from the questionnaires and the working committee. Although there were few respondents, the working committee felt the impacts were unlikely to differ very much from those reflected in the questionnaire survey results.

#### 8.4.1 1 Rem Yr<sup>-1</sup> Limit

Although there was no projected increases in collective dose, one respondent suggested that there would be some costs for facility modification and some increase in radiation protection costs.

## Summary

### 8.4.2 2 Rem Yr<sup>-1</sup> Limit

Here, there apparently would be no impact either on the collective dose or on the facility modification; however, some increase in radiation protection costs was reflected by one of the respondents.

### 8.4.3 5 Rem Yr<sup>-1</sup> and Cumulative Dose in Rem Less Than Age in Years Limit

No impact was seen for this dose limitation option.

## 8.5 Manufacturing and Distribution Including Cyclotron-Produced Radiopharmaceuticals

This data came from responses to the questionnaire as given in Table 5.11 and 5.12, from Table 5.13 with information received in comments on the draft NUREG/CR-6112, from the REIRS data given in Table 4.4, and from the working committee. Material submitted by one medical respondent which dealt with cyclotron-produced radiopharmaceuticals, was included in this category, and the average measurable dose (calculated from Table 5.12 from the questionnaire results) is more than double that given in Table 4.4 from the REIRS Report. This difference may be due to the inclusion of cyclotron workers in Table 5.12, which are not necessarily included in Table 4.4 (they may not be operated by NRC licensees). In a comment received on the Draft Report, the authors raised the issue of the NRC and the states ending up with different regulations.

### 8.5.1 1 Rem Yr<sup>-1</sup> Limit

This group of workers is one of the more highly impacted groups, with the respondents and the working committee suggesting there would be substantial increases in collective dose, in facility modifications, and in annual radiation protection costs.

### 8.5.2 2 Rem Yr<sup>-1</sup> Limit

Here the impact was substantially reduced; however, there still will be important costs both in terms of collective dose, facility modification, and radiation protection. One respondent specifically noted, "...If extremities are lowered by 2/5, as above, we would have

large expenses..." Also, there is concern about the feasibility of operating positive ion cyclotrons under this option.

### 8.5.3 5 Rem Yr<sup>-1</sup> and Cumulative Dose in Rem Less Than Age in Years Limit

The respondents felt there would be no impact for this dose limitation, although one comment noted special exposure limits may be needed for workers who produce isotopes with cyclotrons.

## 8.6 Waste Management

The data are very sparse because there were only two U.S. and one non-U.S. respondents. The REIRS data (Table 4.1) reflects low exposure for those reporting to the NRC (only 119 workers with measurable exposure). Exposure which occurs at the generator site is not included.

### 8.6.1 1 Rem Yr<sup>-1</sup> Limit

Collective dose and radiation protection costs are expected to increase under this dose limit. Table 5.14 indicates that 20% of workers with measurable dose exceeded 1 rem yr<sup>-1</sup>.

### 8.6.2 2 Rem Yr<sup>-1</sup> Limit

Collective dose, facility modification costs, and radiation protection costs are all expected to increase slightly.

### 8.6.3 5 Rem Yr<sup>-1</sup> and Cumulative Dose in Rem Less Than Age in Years Limit

Collective dose, facility modification costs, and radiation protection costs are not expected to increase.

## 8.7 Fuel Fabrication, UF<sub>6</sub> Production

Again, the data are relatively sparse, but over 817 employees with measurable dose were included in the responses. It is extremely important to note that most dose records for this category of workers, i.e. that are given in the REIRS report and in the questionnaire, do not include internal exposure as required under the

revised 10 CFR Part 20. The impact of any change in limits is expected to be severe. From the comments Section of 5.8, the addition of external and internal exposure can be expected to increase reported individual exposures by a factor of 10.

## 8.8 Well Logging

The data from the questionnaire came primarily from a member of the working group, based on a personal survey.

For each exposure limit option, there would be no impact on collective dose, facility modification costs, nor radiation protection. The comment given in Section 5.9 is particularly important: "few well logging technicians are over 40 years of age, and the average tenure of a technician is about ten years. Therefore, the 5 rem yr<sup>-1</sup> and lifetime limit would seem to be achievable, although more data are needed."

## 8.9 General Conclusions

There would be little impact on collective doses, facility modification costs, or annual radiation protection costs under the combined 50 mSv yr<sup>-1</sup> (5 rem yr<sup>-1</sup>) and cumulative dose in 10s of mSv (rem) equal to age in years limit. We point out that the lifetime risk associated with this option - to an individual maximally exposed - would be slightly less than that incurred by a similar individual controlled by the ICRP's limit of 100 mSv in 5 years (10 rem in 5 years). However, a "grandfather" clause allowing up to 2 rem yr<sup>-1</sup> after exceeding the age limit would be needed for several hundred workers in order for them to continue as radiation workers.

A 20 mSv yr<sup>-1</sup> (2 rem yr<sup>-1</sup>) limit would appear achievable, although some tasks, particularly in medicine, fuel fabrication, power reactor maintenance, and, perhaps, industrial radiography might prove extremely difficult to perform. In addition, extensive modifications would be required for many tasks, including the use of robots and remote tools. Depending upon the extent of the modifications made, the collective dose could increase or decrease. That is, extensive remote tooling and facility modifications might lower collective dose. Less ambitious modifications, and less use of remote tooling might keep the collective dose at about the same level while reducing individual doses; lastly, making no changes and allowing the

same tasks to be performed would result in higher collective doses. With this annual limit there may be a potent impact on safety, because some discretionary inspection and maintenance might be constrained.

There has been a suggestion that for a 10 mSv yr<sup>-1</sup> (1 rem yr<sup>-1</sup>) limit, the risk to the most highly exposed individual would be lower than for other options (i.e. equivalent to that in safe industries), but the impacts are expected to be quite serious for many of the industries which responded to the questionnaire. Some tasks in nuclear power, fuel fabrication, and medicine could not be performed under present procedures. For industries with large source terms, facility modifications and radiation protection costs are expected to be extremely large. From a trade-off between the costs of facility modifications and radiation detriment, collective dose may increase substantially.

This summary has focused on the high-dose issues. Many respondents to the questionnaire, however, felt that a 10 mSv yr<sup>-1</sup> (1 rem yr<sup>-1</sup>) limit was entirely feasible. This diversity in potential exposure led the ICRP to recommend applying dose constraints. Such annual dose constraints would be imposed by regulating authorities on specific licensees, based on their source terms, potential for exposure, and costs incurred. Exceeding such constraints would lead to regulatory action. Such a procedure assures that those licensees who can keep below 50, 20, 10 mSv (5,2,1 rem), do so, while recognizing that some operators can not. These latter must have the ability to use the full dose limit.

Two additional issues must be kept in mind when assessing the impact of lower dose limits. The first is the need for licensees to have rigorous administrative limits below the regulated dose limits. For example, with a regulatory limit of a 50 mSv yr<sup>-1</sup> (5 rem yr<sup>-1</sup>), an administrative limit of a 40 mSv yr<sup>-1</sup> (4 rem yr<sup>-1</sup>) might be imposed. With a 20 mSv yr<sup>-1</sup> (2 rem yr<sup>-1</sup>) limit, a 15 mSv yr<sup>-1</sup> (1.5 rem yr<sup>-1</sup>) administrative limit might be used, and so on.

There is one worker group, transportation workers who frequently handle packages containing radioactive materials, which has not been traditionally included under individual dose limitation, which deserve brief mention here, because the reduction in occupational dose limits could affect them.

Exposures to transportation workers have been controlled by limiting both the quantity of radioactive ma-

## Summary

terials (to reduce ingestion of radionuclides) and the external dose rate (to reduce external dose to those workers and to passersby). The bases for the radiation for limits on packages are now relatively obsolete, since they were established on the  $5 \text{ mSv yr}^{-1}$  ( $500 \text{ mrem yr}^{-1}$ ) limit for the public and  $15 \text{ mSv yr}^{-1}$  ( $1.5 \text{ rem yr}^{-1}$ ) limit for workers. Published data suggests that actual experience in terms of doses to these workers shows that there is not an issue of impact here, but the IAEA might be advised to review their basic criteria documents for transportation of radioactive material against today's risk estimates and any new dose limits suggested.

The one additional important group that has not been addressed is the pregnant employee. It is interesting to note, however, that the regulations given in 10 CFR 20 Revised do not differ significantly from those of the 1991 recommendations of the NCRP (NCRP, 1991). The NCRP recommendation is a monthly limit of  $0.5 \text{ mSv}$  ( $50 \text{ mrem}$ ). The 10 CFR 20 Revised requirement is that the fetus receive no more than  $5 \text{ mSv}$  ( $500 \text{ mrem}$ ) over the entire gestation period coupled with effort to avoid substantial variation in monthly doses.

## 9 References

- Atomic Industrial Forum, Inc., "Study of the Effects of Reduced Occupational Radiation Exposure Limits on the Nuclear Power Industry," AIF NESP-017, 1978.
- Ball, J.R., S. Cohen, E.Z. Ziegler, "A Handbook for Cost Estimating," NUREG/CR-3971, ANL/EES-TM-265, Argonne National Laboratory, Argonne, IL 60439, October 1984.
- Baum, J.W., "Valuation of Dose Avoided at U.S. Nuclear Power Plants, Nuclear Plant Journal, Vol. 9, No. 2, pp. 40-47, March-April 1991.
- Baum, J.W. and T. A. Khan, BNL ALARA Center Experience with an Information Exchange System on Dose Control at Nuclear Power Plants, Presented at the NEA Workshop on Work Management in Occupational Dose Control, Paris, France, February 4-6 1992.
- Baum, J.W. and G. R. Matthews, "Compendium of Cost-Effectiveness Evaluations of Modifications for Dose Reduction at Nuclear Power Plants," NUREG/CR-4373, BNL-NUREG-51915, Brookhaven National Laboratory, Upton, NY 11973, December 1985 (Pg 73).
- Brooks, B.G., "Occupational Radiation Exposure at Commercial Nuclear Power Reactors and Other Facilities 1985," NUREG-0713, Vol. 7, U.S. Nuclear Regulatory Commission, Office of Nuclear Regulatory Research, Washington, D.C., 1988.
- Department of Energy, "Study of Anticipated Impact on DOE Programs from Proposed Reductions to the External Occupational Radiation Exposure Limit," DOE/EV-0045, U.S. Department of Energy, Assistant Secretary for Environment, Division of Operational and Environmental Safety, Washington, D.C. 20585, 1979.
- Department of Energy, Final Report to the Secretary of Energy: Implications of the BEIR V Report to the Department of Energy, DOE/EH-0158T, 1990.
- Divine, Robert A., Blowing on the Wind, The Nuclear Test Ban Debate, Oxford University Press, 1978.
- Edison Electric Institute Health Physics Committee, "Utility Responses to Questionnaire on Dose Limits and Guidance," EEI Nuclear Report # HPC-91-001, 1991.
- Environmental Protection Agency, "Occupational Exposure to Ionizing Radiation in the United States: A Comprehensive Review for the Year 1980 and a Summary of Trends for the Years 1960-1985," EPA Report 520/1-84-005 EPA, Washington, DC, 1984.
- Environmental Protection Agency, "Radiation Protection Guidance to Federal Agencies for Occupational Exposure," EPA, *Federal Register* 52, No. 17, 2822; with corrections published in the *Federal Registers* of Friday, January 30, and Wednesday, February 4, 1987, 1987.
- Federal Radiation Council, "Background Material for the Development of Radiation Protection Standards, FRC: Report No. 1, 1960.
- Haldane, J.B.S., "The Formal Genetics of Man," (Croonian Lecture), Proc. Roy. Soc., Ser. B, 135, 147, 1948.
- Hinson, C.S., NRC, Memorandum to NRC Director, F.J. Congel, "LWR Occupational Dose Data for 1991," October 28, 1992.
- International Commission on Radiological Protection, "Recommendations of the International Commission on Radiological Protection," ICRP, (Revised December 1, 1954), British Journal of Radiology, Supplement No. 6, 1954.
- International Commission on Radiological Protection, "Recommendations of the ICRP (adopted September 9, 1958)," ICRP Publication 1, Pergamon Press, London, 1959a.
- International Commission on Radiological Protection, "Report of Committee II on Permissible Dose for Internal Radiation," ICRP Publication 2, Pergamon Press, London, 1959b.
- International Commission on Radiological Protection, "Recommendations of the International Commission on Radiological Protection (As amended 1959 and revised 1962)," ICRP Publication 6, Pergamon Press, London, 1964.
- International Commission on Radiological Protection, "Recommendations of the International Commission on Radiological Protection," ICRP Publication 26, Pergamon Press, Oxford, 1977.

## References

- International Commission for Radiation Protection, "Protection of the Patient in Nuclear Medicine," Annals of the ICRP, Volume 17, No. 4, Pergamon Press, Oxford, 1987.
- International Commission on Radiological Protection, "1990 Recommendations of the International Commission on Radiological Protection," ICRP Publication 60, Annals of the ICRP, Vol 21, No. 1-3 Pergamon Press, New York, 1991.
- Kaul, A., F. Kossel, K. Martignoni, and J. Nitschke, "Limitation of Occupational Radiation Risk by Radiation Protection Legislation in the Federal Republic of Germany," J. Radiol. Prot., Vol. 9, No. 2, pp. 85-92, 1989.
- Khan, T.A., J.W. Baum, and B.J. Dionne, "Dose-Reduction Techniques for High-Dose Worker Groups in Nuclear Power Plants, NUREG/CR 5139, BNL-NUREG-52278, Brookhaven National Laboratory, Upton, NY 11973, March 1991a.
- Khan T.A., J.W. Baum, and B.J. Dionne, ACE-ALARA Center's Information Service, Radiation Protection Management Journal, Volume 8, No. 4, pp 24-34, July-August 1991b.
- Khan T.A., D.S. Vulin, H. Liang, and J.W. Baum, "Data Base on Dose Reduction Research Projects for Nuclear Power Plants," NUREG/CR-4409, BNL-NUREG-5193, Vol. 4, August 1992.
- Kindred, G.W., Perry Nuclear Plant, Available on the ACE On-Line Information System or from the BNL ALARA Center, Bldg. 703M, Upton, NY 11973, 1992.
- Lea, D.E., Actions of Radiations on Living Cells, The MacMillan Company, New York, 1947.
- Le Surf, J.E., "Implications of Reduction in Federal Radiation Exposure Limits," prepared for Electric Power Research Institute (EPRI), Palo Alto, CA 943-03, 1988.
- Medical Research Council, "The Hazards to Man of Nuclear and Allied Radiations," MRC, UK, Cmd 9780, H.M.S. Office, 1956.
- Muller, H.J., "Artificial Transmutation of the Gene," Science, 46, 84-87, 1927.
- Mutscheller, A.M., Am. J. Roentgenol. Radium Therapy Nucl. Med. 13, 65 (1925).
- National Academy of Sciences National Research Council, "The Biological Effects of Atomic Radiation-Summary Reports," NAS-NRC, 1956.
- National Academy of Sciences National Research Council Committee on the Biological Effects of Ionizing Radiation, "The Effects on Populations of Exposure to Low Levels of Ionizing Radiation," BEIR V Report, National Academy of Sciences, National Research Council, Washington, DC, 1972.
- National Academy of Sciences National Research Council Committee on the Biological Effects of Ionizing Radiation, "Health Effects of Exposure to Low Levels of Ionizing Radiation," BEIR V Report, NAS BEIR, National Academy of Sciences, National Academy Press, Washington, DC, 1990.
- National Bureau of Standards, Permissible Dose from External Sources of Ionizing Radiation, Handbook, 59, NBS, 1954.
- National Committee on Radiation Protection and Measurements, "Maximum Permissible Radiation Exposure to Man-A Preliminary Statement of the NCRP (January 8, 1957), Amer. J. Roentgen. 77, 910 (1957) and Radiology 68, 260 (1957).
- National Council on Radiation Protection and Measurements, "Basic Radiation Protection Criteria," NCRP Report No. 39, Washington, D.C. 20014, January 15, 1971.
- National Council on Radiation Protection and Measurements, Recommendations on Limits for Exposure to Ionizing Radiation, NCRP Report No. 91 (National Council on Radiation Protection and Measurements, Bethesda, Maryland), 1987.
- National Radiation Protection Board, "Interim Guidance on the Implications of Recent Revisions of Risk Estimates and the ICRP 1987 Como Statement," NRPB-GS9, NRPB, Chilton, Didcot, Oxon OX11 0RQ, 1987.
- Nuclear Regulatory Commission, Basic Standards for Radiation Protection, NRC, Title 10, Code of Federal Regulations, Part 20, 1960.

## References

Nuclear Regulatory Commission, Standards for Protection Against Radiation, NRC, Title 10, Code of Federal Regulations, Part 20, Federal Register 56-(98): 23390-23470, 1991.

Preston, D.L. and D.A. Pierce, "The Effect of Changes in Dosimetry on Cancer Mortality Risk Estimates in the Atomic Bomb Survivors," Technical Report RERF TR-9-87, Radiation Effects Research Foundation, Hiroshima, Japan, 1981.

Raddatz, C.T. and D. Hagemeyer, "Occupational Radiation Exposure at Commercial Nuclear Power Reactors and Other Facilities 1990," NUREG-0713, Vol. 12, January 1993.

Report of the United Nations Scientific Committee on the Effects of Atomic Radiation, General Assembly Official Records, Seventeenth Session, Supplement No. 16 (A/5216), United Nations, New York, 1962.

Schmitt, C.H. and Brice, J.F., "Occupational Exposure from U.S. Naval Nuclear Propulsion Plans and Their Support Facilities," U.S. Department of the Navy Report NT-84-2, Department of the Navy, Washington, D.C., 1984.

Shimizu, Y., H. Kato, and W.J. Schull, Life Span Study Report 11, Part II: Cancer Mortality in the Years 1950-1985 Based on the Recently Revised Doses (DS86). RERF TR/5-88 (1988).

Shimizu, Y., H. Kato, W.J. Schull et al., Life Span Study Report 11, Part I: Comparison of Risk Coefficients for Site-Specific Cancer Mortality Based on the DS86 and T65DR Shielded Kerma and Organ Doses. RERF TR/12-87 (1987).

United Nations Scientific Committee on the Effects of Atomic Radiation, General Assembly, Official Records: Seventeenth Session, Supplement No. 16 (A/5216), United Nations, New York, 1962.

United Nations Scientific Committee on the Effects of Atomic Radiation, Ionizing Radiation: Levels and Effects, Vol. II: Effects, UNSCEAR, United Nations, New York, 1972.

United Nations Scientific Committee on the Effects of Atomic Radiation, "Sources and Effects of Ionizing Radiation," UNSCEAR, 1977 Report to the General Assembly, United Nations, New York, 1977.

United Nations Scientific Committee on the Effects of Atomic Radiation, "Sources, Effects, and Risks of Ionizing Radiation," UNSCEAR, 1988 Report to the General Assembly, United Nations, New York, 1988.

United Nations Scientific Committee on the Effects of Atomic Radiation, "Sources and Effects of Ionizing Radiation," UNSCEAR, 1993 Report to the General Assembly, United Nations, New York, 1993.

United Nations Scientific Committee on the Effects of Atomic Radiation, "Sources and Effects of Ionizing Radiation," UNSCEAR, 1994 Report to the General Assembly, United Nations, New York, 1994.

Warman E.A., L.M. Wainio, and E.A.B. Eastman, "Estimated Additional Number of Workers and Additional Cumulative Exposure Due to Reducing Annual Dose Limits per Individual," RP-29, Stone and Webster Engineering Corp., P.O. Box 2325, Boston, MA 02108. April 19, 1978.

# Appendix A

## Questionnaire on the Impact of Reduced Dose Limits at Your Facility

Possible Dose Limit				
Estimated Impacts:	2 rem y <sup>-1</sup>	1 rem y <sup>-1</sup>	Cumulative < age and 5 rem y <sup>-1</sup>	Cumulative < age and 2 rem y <sup>-1</sup>
Will collective dose change? Y, N; Up, Down; Estimate				
Facilities Modifications needed? Y, N; and est. costs				
Increase Rad. Protection? Y, N; and est. costs				

Your 1989 experience:

**Number of Employees with Annual Doses:**  
 >5 rem \_\_\_\_ >2 rem \_\_\_\_ >1 rem \_\_\_\_

Number of Employees with Lifetime Dose Greater Than Age in Rem: \_\_\_\_\_

Number of Employees with Measureable Dose: \_\_\_\_\_

Annual Collective Dose: \_\_\_\_\_

Comments and Suggestions:

---



---



---

Would you be willing and available to participate in a working group to review and assess the results of this questionnaire?  
 Yes \_\_\_\_ No \_\_\_\_

Please indicate which of the following apply to your organization:

- |   |   |  |
|---|---|--|
| <input type="checkbox"/> Medical/Dental               | <input type="checkbox"/> Manufacturing/Distribution                 | <input type="checkbox"/> Mining and Milling          |
| <input type="checkbox"/> Veterinary                   | <input type="checkbox"/> Naval Nuclear Propulsion Program           | <input type="checkbox"/> Fuel Fabrication            |
| <input type="checkbox"/> University                   | <input type="checkbox"/> Other Military & their Contractors         | <input type="checkbox"/> Waste Management & Disposal |
| <input type="checkbox"/> Other Research & Development | <input type="checkbox"/> Nuclear Power Reactors & their Contractors | <input type="checkbox"/> Research and Test Reactors  |
| <input type="checkbox"/> Well Logging                 | <input type="checkbox"/> Test & Measurements                        | <input type="checkbox"/> Other (Specify) _____       |

Please provide additional description of your organization where the primary category is insufficient, e.g., x-ray therapy, radiopharmacy, reactor refueling, etc.: \_\_\_\_\_

Name & Title: \_\_\_\_\_

Company: \_\_\_\_\_

Address: \_\_\_\_\_

Telephone: \_\_\_\_\_

Please fold and return with any additional comments to Charles B. Meinhold, Radiological Sciences Division, Brookhaven National Laboratory, Building 703M, Upton, New York 11973. Telephone: (516) 282-4425, FAX (516) 282-5810.

BIBLIOGRAPHIC DATA SHEET

(See instructions on the reverse)

1. REPORT NUMBER  
(Assigned by NRC. Add Vol., Supp., Rev.,  
and Addendum Numbers, if any.)

NUREG/CR-6112  
BNL/NUREG-52394

2. TITLE AND SUBTITLE

Impact of Reduced Dose Limits on NRC Licensed Activities  
Major Issues in the Implementation of ICRP/NCRP Dose Limit  
Recommendations  
  
Final Report

3. DATE REPORT PUBLISHED

MONTH YEAR

May 1995

4. FIN OR GRANT NUMBER

L1285

5. AUTHOR(S)

C.B. Meinhold

6. TYPE OF REPORT

Technical

7. PERIOD COVERED (Inclusive Dates)

10/01/89 to 3/01/95

8. PERFORMING ORGANIZATION - NAME AND ADDRESS (If NRC, provide Division, Office or Region, U.S. Nuclear Regulatory Commission, and mailing address; if contractor, provide name and mailing address.)

Brookhaven National Laboratory  
Upton, NY 11973

9. SPONSORING ORGANIZATION - NAME AND ADDRESS (If NRC, type "Same as above"; if contractor, provide NRC Division, Office or Region, U.S. Nuclear Regulatory Commission, and mailing address.)

Division of Regulatory Applications  
Office of Nuclear Regulatory Research  
U.S. Nuclear Regulatory Commission  
Washington, DC 20555-0001

10. SUPPLEMENTARY NOTES

11. ABSTRACT (200 words or less)

This report summarizes information required to estimate, at least qualitatively, the potential impacts of reducing occupational dose limits below those given in 10 CFR 20 (Revised). The following overall conclusions were reached: (1) Although  $10 \text{ mSv yr}^{-1}$  is a reasonable limit for many licensees, such a limit could be extraordinarily difficult to achieve and potentially destructive to the continued operation of some licensees, such as nuclear power, fuel fabrication, and medicine, (2) Twenty  $\text{mSv yr}^{-1}$  as a limit is possible for some of these groups, but for others it would prove difficult, (3) Fifty  $\text{mSv yr}^{-1}$  and age in 10s of  $\text{mSv}$  appear reasonable for all licensees, both in terms of the lifetime risk of cancer and severe genetic effects to the most highly exposed workers, and the practicality of operation. In some segments of the industry, this acceptability is based on the adoption of a "grandfather clause" for those people exceeding or close to exceeding the cumulative limit at this time.

12. KEY WORDS/DESCRIPTORS (List words or phrases that will assist researchers in locating the report.)

Radiation Doses-Occupational Exposure, Biological Radiation Effects-Lifetime, Dosimetry, Radiation Effects-Dose Rate, Dose-Response Relations, Dose Rates-Genetic Effects, Threshold Dose, Radiation Monitoring, Dose Limits, Risk Assessment

13. AVAILABILITY STATEMENT

Unlimited

14. SECURITY CLASSIFICATION

(This Page)

Unclassified

(This Report)

Unclassified

15. NUMBER OF PAGES

16. PRICE