

IPRenewal NPEmails

From: Stuyvenberg, Andrew
Sent: Thursday, June 30, 2011 3:33 PM
To: Gray, Dara F
Cc: IPRenewal NPEmails
Subject: FW: nmfs questions re. IP thermal
Attachments: temps_sns-2.docx

Dara -

Attached is a version of the shortnose sturgeon technical information from Julie that contains full references should anyone in the Entergy staff need them.

Best,
Drew

-----Original Message-----

From: Julie Crocker [<mailto:Julie.Crocker@Noaa.Gov>]
Sent: Thursday, June 30, 2011 2:31 PM
To: Stuyvenberg, Andrew
Cc: Balsam, Briana; Imboden, Andy; Logan, Dennis; Julie Williams; Mark Murray-Brown; IPRenewal NPEmails
Subject: Re: nmfs questions re. IP thermal

should have thought about that! Attached is a version with the full citations at the end. I also noticed an error when doing that - both Zeigweid et al. papers are from 2008, so I corrected that with 2008a and 2008b to differentiate between the two papers.

Julie

Hearing Identifier: IndianPointUnits2and3NonPublic_EX
Email Number: 2720

Mail Envelope Properties (AF843158D8D87443918BD3AA953ABF782B680FF596)

Subject: FW: nmfs questions re. IP thermal
Sent Date: 6/30/2011 3:33:29 PM
Received Date: 6/30/2011 3:33:31 PM
From: Stuyvenberg, Andrew

Created By: Andrew.Stuyvenberg@nrc.gov

Recipients:
"IPRenewal NPEmails" <IPRenewal.NPEmails@nrc.gov>
Tracking Status: None
"Gray, Dara F" <DGray@entergy.com>
Tracking Status: None

Post Office: HQCLSTR02.nrc.gov

Files	Size	Date & Time
MESSAGE	756	6/30/2011 3:33:31 PM
temps_sns-2.docx	15781	

Options
Priority: Standard
Return Notification: No
Reply Requested: No
Sensitivity: Normal
Expiration Date:
Recipients Received:

The temperature preference for shortnose sturgeon is not known (Dadswell et al. 1984) but shortnose sturgeon have been found in waters with temperatures as low as 2 to 3°C (Dadswell et al. 1984) and as high as 34°C (Heidt and Gilbert 1978). Foraging is known to occur at temperatures greater than 7°C (Dadswell 1979). Temperatures above 28°C are thought to be stressful to shortnose sturgeon. In the Altamaha River, temperatures of 28-30°C during summer months are thought to trigger movements to deep cool water refuges. Ziegewald et al. (2008a) conducted studies to determine critical and lethal thermal maxima for young-of-the-year (YOY) shortnose sturgeon acclimated to temperatures of 19.5 and 24.1°C. Lethal thermal maxima were 34.8°C (± 0.1) and 36.1°C (± 0.1) for fish acclimated to 19.5 and 24.1°C, respectively. The study also used thermal maximum data to estimate upper limits of safe temperature, final thermal preferences, and optimum growth temperatures for YOY shortnose sturgeon. Visual observations suggest that fish exhibited similar behaviors with increasing temperature regardless of acclimation temperature. As temperatures increased, fish activity appeared to increase; approximately 5–6°C prior to the lethal endpoint, fish began frantically swimming around the tank, presumably looking for an escape route. As fish began to lose equilibrium, their activity level decreased dramatically, and at about 0.3°C before the lethal endpoint, most fish were completely incapacitated. Estimated upper limits of safe temperature (ULST) ranged from 28.7 to 31.1°C and varied with acclimation temperature and measured endpoint. Upper limits of safe temperature (ULST) were determined by subtracting a safety factor of 5°C from the lethal and critical thermal maxima data. Final thermal preference and thermal growth optima were nearly identical for fish at each acclimation temperature and ranged from 26.2 to 28.3°C. Thermal maxima ranged from 33.7 to 36.1°C and varied with acclimation temperature and designated endpoint. Ziegewald et al. (2008b) used data from laboratory experiments to examine the individual and interactive effects of salinity, temperature, and fish weight on the survival of young-of-year shortnose sturgeon. Survival in freshwater declined as temperature increased, but temperature tolerance increased with body size. The temperature lethal to 50% of the test fish after 48 h ranged from 28.2°C to 30.7°C. The authors conclude that temperatures above 29°C substantially reduce the probability of survival for young-of-year shortnose sturgeon. However, previous studies indicate that juvenile sturgeons achieve optimum growth at temperatures close to their upper thermal survival limits (Mayfield and Cech 2004; Allen et al. 2006; Ziegewald et al. 2008a), suggesting that shortnose sturgeon may seek out a narrow temperature window to maximize somatic growth without substantially increasing maintenance metabolism. For shortnose sturgeon, dissolved oxygen (DO) also seems to play a role in temperature tolerance, with increased stress levels at higher temperatures with low DO versus the ability to withstand higher temperatures with elevated DO (Niklitchek 2001).

References

- Allen PJ, Nicholl M, Cole S, Vlazny A, Cech JJ Jr. 2006. Growth of larval to juvenile green sturgeon in elevated temperature regimes. *Trans Am Fish Soc* 135:89–96
- Dadswell, M.J. 1979. Biology and population characteristics of the shortnose sturgeon, *Acipenser brevirostrum* LeSueur 1818 (Osteichthyes: Acipenseridae), in the Saint John River estuary, New Brunswick, Canada. *Canadian Journal of Zoology* 57:2186-2210.
- Dadswell, M.J., B.D. Taubert, T.S. Squiers, D. Marchette, and J. Buckley. 1984. Synopsis of biological data on shortnose sturgeon, *Acipenser brevirostrum* Lesueur 1818. NOAA Technical Report, NOAA Fisheries 14, National Marine Fisheries Service. October 1984 45 pp.

Heidt, A.R., and R.J. Gilbert. 1978. The shortnose sturgeon in the Altamaha River drainage, Georgia. Pages 54-60 in R.R. Odum and L. Landers, editors. Proceedings of the rare and endangered wildlife symposium. Georgia Department of Natural Resources, Game and Fish Division, Technical Bulletin WL 4, Athens, Georgia.

Mayfield RB, Cech JJ Jr. 2004. Temperature effects on green sturgeon bioenergetics. *Trans Am Fish Soc* 133:961–970

Niklitschek, J. E. 2001. Bioenergetics modeling and assessment of suitable habitat for juvenile Atlantic and shortnose sturgeons (*Acipenser oxyrinchus* and *A. brevirostrum*) in the Chesapeake Bay. Dissertation. University of Maryland at College Park, College Park.

Ziegeweid, J.R., C.A. Jennings, and D.L. Peterson. 2008a. Thermal maxima for juvenile shortnose sturgeon acclimated to different temperatures. *Environmental Biology of Fish* 3: 299-307.

Ziegeweid, J.R., C.A. Jennings, D.L. Peterson and M.C. Black. 2008b. Effects of salinity, temperature, and weight on the survival of young-of-year shortnose sturgeon. *Transactions of the American Fisheries Society* 137:1490-1499.