



Figure 2.5.1-323 Earthquakes by Depth and Major Plate Boundary Structures in the Northeastern Area of the North America-Caribbean Plate Boundary



Source: Reference 639


Figure 2.5.1-324 GPS Site Velocities with Respect to North America




Source: [Reference 780](#)

Figure 2.5.1-325 Kinematic Illustrations Showing Interactions of Septentrional and Northern Hispaniola Faults at Depth

(A) EAST OF 70° LONGITUDE




(B) WEST OF 70° LONGITUDE



Note: Northern Hispaniola Thrust fault is equivalent to North Hispaniola Subduction Zone.

Source: [Reference 638](#)


Figure 2.5.1-326 Focal Mechanisms for Major Earthquakes in the North Hispaniola Subduction Zone



Notes:
EPGF = Enriquillo-Plantain Garden fault zone
NHF-TB = Northern Hispaniola fold-thrust belt
SF = Septentrional fault


Source: [Reference 638](#)

Figure 2.5.1-327 Damage Zones for Major Earthquakes in the Northeastern Caribbean, 1615-1992



Source: [Reference 591](#)


Figure 2.5.1-328 Bathymetry, Structural Features, and GPS Vectors relative to North America, Northeastern Caribbean



Source: [Reference 585](#)


Turkey Point Units 6 & 7
COL Application
Part 2 — FSAR

Figure 2.5.1-329 Timeline of Regional Tectonic and Geologic Events



Sources: [References 307](#), [368](#), and [639](#)


Figure 2.5.1-330 North America Stress Provinces



Base Source: [Reference 822](#)
Source of world stress data: [Reference 731](#)

Turkey Point Units 6 & 7
COL Application
Part 2 — FSAR

Figure 2.5.1-331 Site Vicinity Geologic Map



Base sources: [Reference 435](#)

Source of geologic information: [Reference 827](#)

Turkey Point Units 6 & 7
COL Application
Part 2 — FSAR


Figure 2.5.1-332 Site Stratigraphy

ERATHEM	SYSTEM	SERIES	HYDRO- GEOLOGIC UNIT		STRATIGRAPHIC UNIT	LITHOLOGY	APPROXIMATE TOP ELEVATION (ft NAVD 88)	APPROXIMATE THICKNESS (ft)	
CENOZOIC	QUATERNARY	HOLOCENE			organic muck	organic soil and silt	0	3	
		PLEISTOCENE	Surficial aquifer system	Biscayne aquifer	Miami Limestone	sandy, oolitic limestone	-3	25	
					Key Largo Limestone	well indurated, vuggy, coralline limestone	-28	22	
					Fort Thompson Formation	poor/well indurated fossiliferous limestone	-50	65	
	TERTIARY	PLIOCENE			Semi-confining unit	Tamiami Formation	sand and silt with calcareneitic limestone	-115	105
		MIOCENE	Intermediate confining unit		Hawthorn Group	Peace River Formation	silty calcareous sand and silt	-220	235
						Arcadia Formation	calcareous wackestone with indurated limestones, sandstone, and sand	-455	>160
								drilling ended at -616.5 ft	

Note: see [Figures 2.5.1-338, 2.5.1-339, 2.5.1-340, and 2.5.1-341](#) for site geologic cross sections.

Turkey Point Units 6 & 7
COL Application
Part 2 — FSAR


Figure 2.5.1-333 Vegetated Depressions Identified Within Site from Photographs Taken Before Construction of the Cooling Canal System



Note: Reconnaissance mapping performed using 1940s 1:40,000 scale panchromatic stereo aerial photography (Reference 386), but shown on 2004 imagery (Reference 435) of the Units 6 & 7 site for reference.

Turkey Point Units 6 & 7
COL Application
Part 2 — FSAR


Figure 2.5.1-334 Site Geologic Map



Base sources: [References 829](#), and [435](#)
Source of geologic information: [Reference 827](#)

Turkey Point Units 6 & 7
COL Application
Part 2 — FSAR

Figure 2.5.1-335 Site Area Geologic Map




Base sources: [Reference 435](#)

Source of geologic information: [Reference 219](#)


Turkey Point Units 6 & 7
COL Application
Part 2 — FSAR

Figure 2.5.1-336 Locations of Geologic Cross Sections



Turkey Point Units 6 & 7
COL Application
Part 2 — FSAR

Figure 2.5.1-337 Surficial Deposits Map




Base sources: [Reference 829](#)

Source of geologic information: [References 715](#) and [830](#)

Turkey Point Units 6 & 7
COL Application
Part 2 — FSAR


Figure 2.5.1-338 Geologic Cross Section A-A'



Note: The location of this cross section is shown on [Figure 2.5.1-336](#).

Turkey Point Units 6 & 7
COL Application
Part 2 — FSAR


Figure 2.5.1-339 Geologic Cross Section B-B'



Note: The location of this cross section is shown on [Figure 2.5.1-336](#).

Turkey Point Units 6 & 7
COL Application
Part 2 — FSAR


Figure 2.5.1-340 Geologic Cross Section C-C'



Note: The location of this cross section is shown on [Figure 2.5.1-336](#).

Turkey Point Units 6 & 7
COL Application
Part 2 — FSAR


Figure 2.5.1-341 Geologic Cross Section D-D'



Note: The location of this cross section is shown on [Figure 2.5.1-336](#).


Turkey Point Units 6 & 7
COL Application
Part 2 — FSAR

Figure 2.5.1-342 Isopach of the Site: Key Largo Limestone



Turkey Point Units 6 & 7
COL Application
Part 2 — FSAR

Figure 2.5.1-343 Structure Contour Map: Top of Fort Thompson Formation



Turkey Point Units 6 & 7
COL Application
Part 2 — FSAR

Figure 2.5.1-344 Isopach of the Site: Fort Thompson Formation






Figure 2.5.1-345 Geologic Hazards for Coastal Zones of Cuba



Modified from: [Reference 742](#)


Figure 2.5.1-346 Interpreted Seismic Line across the Edge of the Little Bahama Bank



Note: Sequence G (the shallow-water carbonate platform sampled at Site 627) is offset, while sequences A-F thicken across the fault trace, suggesting syn-sedimentary movement.

Modified from: [Reference 785](#)

Figure 2.5.1-347 Initiation of the Greater Antilles Arc and Collision with the Caribbean Oceanic Plateau




Notes:



A. Present-day distribution of Cretaceous to Recent island arc and late Cretaceous oceanic plateau crust in the Caribbean.

B. Mid-Cretaceous (Cenomanian) reconstruction of the Caribbean island arc and oceanic plateau

Source: [Reference 833](#)

Figure 2.5.1-348 Tsunami Sediments



C. Typical tsunami deposit	Typical storm deposit
 <ul style="list-style-type: none"> • mudcap • lamina sets may be separated by thin mud or heavy mineral lamina • often normally graded • rip up clasts • 5-25 cm thick • abrupt lower contact 	 <ul style="list-style-type: none"> • mudcap rare • may have foresets, troughs, climbing ripples • planar stratification • many laminae and laminasets • 25-200 cm thick • abrupt lower contact

Notes:

A and B. Comparison of typical inundation distances, sediment-transport distances, and maximum water levels (indicated by height of wrack line) for deposition by tsunamis (A) and coastal storms (B)

C. Composite characteristics of typical sandy tsunami and storm deposits

Source: [Reference 890](#)