


**LEGEND**

- ★ HAR Site
- Main Dam
- ▲ Sears No. 1 and Groce No. 1 Wells (Bain and Brown, 1981)
- 8-km (5-mi.) Radius from Site
- 40-km (25-mi.) Radius from Site
- Paleozoic Faults
- Mesozoic Faults (dashed where inferred, dotted where concealed) (Ebasco, 1975, CP&L, 1983, NCGS, 1985, Wooten et al., 1996, Harding Lawson, 1997, NCGS, 2006, and this study)
- 49 Cretaceous and Cenozoic Faults (numbers refer to descriptions in Table 2.5.1-201; from Parker, 1979, and Prowell, 1983)
- Paleozoic Folds (dashed where inferred)
- Mesozoic Folds (dashed where inferred) (NCGS, 2006)
- Dikes (dashed where inferred) (CP&L, 1983, NCGS, 1985, NCGS, 2006, and Technos, this study)

Geologic Units (NCGS, 1985) (See Figure 2.5.1-230, Sheet 2 for Unit Descriptions)

Tec	TRcs	CZfv
Tpy	PPg	CZfv1
Tt	PPmg	CZg
Kb	PzZg	CZig
Kc	PzZq	CZiv
Km	PzZu	CZmd
Jd	CZam	CZph
TRc	CZbg	CZv
TRcc	CZc	CZve
TRcp	CZfg	CZmv

Note N1: This fault included in NCGS (2006) unpublished mapping appears to represent variations in metamorphic grade adjacent to the pluton rather than a fault contact.



Progress Energy Carolinas  
**Shearon Harris Nuclear Power Plant  
 Units 2 and 3  
 Part 2, Final Safety Analysis Report**  
 New Hill, North Carolina

Site Vicinity Geologic Map  
 (40-km [25-mi.] Radius)

FIGURE 2.5.1-230 (SHEET 1 OF 2) Rev. 1

**DESCRIPTION OF GEOLOGIC UNITS**

Pliocene-Quaternary	<b>Tt</b>	TERRACE DEPOSITS AND UPLAND SEDIMENT - Gravel, clayey sand, and sand and minor iron-oxide cemented sandstone.	Paleozoic	<b>PPg</b>	GRANITIC ROCK (Pennsylvanian to Permian, 265-325 m.y.) - Megacrystic to equigranular.	Late Proterozoic - Cambrian	<b>CZmd</b>	METAMUDSTONE AND META-ARGILLITE - Bedding plane and axial planar cleavage common; interbedded with metasandstone, metaconglomerate, and metavolcanic rock.
	<b>Tec</b>	CASTLE HAYNE FORMATION - Comfort Member and New Hanover Member, undivided. Comfort Member: Bryozoan-echinoid skeletal limestone, locally dolomitized, solution cavities common. New Hanover Member: Phosphate-pebble conglomerate, micritic, thin; restricted to basal part of Castle Hayne Formation in southeastern counties.		<b>PPmg</b>	FOLIATED TO MASSIVE GRANITIC ROCK (Pennsylvanian to Permian, 270-320 m.y.) - Megacrystic to equigranular.		<b>CZph</b>	PHYLLITE AND SCHIST - Locally laminated and pyritic; includes phyllonite, sheared fine-grained metasediment, and metavolcanic rock.
Tertiary	<b>Tpy</b>	YORKTOWN FORMATION AND DUPLIN FORMATION, UNDIVIDED - Yorktown Formation: Fossiliferous clay with varying amounts of fine-grained sand, bluish gray, shell material commonly concentrated in lenses; mainly in area north of Neuse River. Duplin Formation: Shelly, medium- to coarse-grained sand, sandy marl, and limestone, bluish gray; mainly in area south of Neuse River.	Paleozoic	<b>PzZg</b>	METAMORPHOSED GABBRO AND DIORITE - Foliated to massive.	Late Proterozoic - Cambrian	<b>CZve</b>	METAVOLCANIC EPICLASTIC ROCK - Metamorphosed argillite, mudstone, volcanic sandstone, conglomerate, and volcanic rock.
				<b>PzZu</b>	META-ULTRAMAFIC ROCK - Metamorphosed dunite and peridotite; serpentinite, soapstone, and other altered ultramafic rock. Only larger bodies shown.		<b>CZmv</b>	MAFIC METAVOLCANIC ROCK - Metamorphosed basalt flows and tuffs, dark green to black; interbedded with felsic and intermediate metavolcanic rock and metamudstone.
Cretaceous	<b>Kb</b>	BLACK CREEK FORMATION - Clay gray to black, lignitic; contains thin beds and laminae of fine-grained micaceous sand and thick lenses of cross-bedded sand. Glauconitic, fossiliferous clayey sand lenses in upper part.	Late Proterozoic - Cambrian	<b>CZam</b>	AMPHIBOLITE - Metamorphosed mafic extrusive and intrusive rock; includes hornblende gneiss, thin layers of mica schist, and small non-layered masses of metadiorite and metagabbro.			
	<b>Km</b>	MIDDENDORF FORMATION - Sand, sandstone, and mudstone, gray to pale gray with an orange cast, mottled; clay balls and iron-cemented concretions common, beds laterally discontinuous, cross bedding common.		<b>CZbg</b>	BIOTITE GNEISS AND SCHIST - Inequigranular and megacrystic; in places contains garnet; interlayered and gradational with mica schist and amphibolite; includes small masses of granitic rock.			
	<b>Kc</b>	CAPE FEAR FORMATION - Sandstone and sandy mudstone, yellowish gray to bluish gray, mottled red to yellowish orange, indurated, graded and laterally continuous bedding, blocky clay, faint cross-bedding, feldspar and mica common.		<b>CZc</b>	VOLCANIC METACONGLOMERATE - Includes metagraywacke and metamudstone.			
Jurassic	<b>Jd</b>	DIABASE - Dikes, gray to black.	Late Proterozoic - Cambrian	<b>CZfg</b>	FELSIC MICA GNEISS - Interlayered with graphitic mica schist and mica-garnet schist, commonly with kyanite; minor hornblende gneiss.			
	<b>TRc</b>	CHATHAM GROUP (undivided) - Conglomerate, fanglomerate, sandstone, and mudstone.		<b>CZfv</b>	FELSIC METAVOLCANIC ROCK - Metamorphosed dacitic to rhyolitic flows and tuffs, light gray to greenish gray; interbedded with mafic and intermediate metavolcanic rock, meta-argillite, and metamudstone.			
Triassic	<b>TRcs</b>	SANFORD FORMATION - Conglomerate, fanglomerate, sandstone, and mudstone.	Late Proterozoic - Cambrian	<b>CZg</b>	METAMORPHOSED GRANITIC ROCK (Late Proterozoic to late Cambrian, 520-650m.y.) - Megacrystic, well-foliated, locally contains hornblende.			
	<b>TRcc</b>	CUMNOCK FORMATION - Sandstone and mudstone, gray to black; coal beds and carbonaceous shale. Grades into Pekin and Sanford formations.		<b>CZig</b>	INJECTED GNEISS - Biotite gneiss and schist intruded by numerous sills and dikes of granite, pegmatite, and apatite; minor hornblende gneiss.			
	<b>TRcp</b>	PEKIN FORMATION - Conglomerate, sandstone, and mudstone.		<b>CZlg</b>	LINEATED-FELSIC MICA GNEISS - White to pink with strong lineation of muscovite-biotite streaks and prismatic quartz aggregates; planar foliation and layering weak; minor mica schist and hornblende gneiss.			
			Late Proterozoic - Cambrian	<b>CZiv</b>	INTERMEDIATE METAVOLCANIC ROCK - Metamorphosed andesitic tuffs and flows, medium to dark grayish green; minor felsic and mafic metavolcanic rock.			

Source: NCGS (1985)

Progress Energy Carolinas  
**Shearon Harris Nuclear Power Plant  
 Units 2 and 3**  
**Part 2, Final Safety Analysis Report**  
 New Hill, North Carolina

Site Vicinity Geologic Map  
 (40-km [25-mi.] Radius)  
 FIGURE 2.5.1-230 (SHEET 2 OF 2) Rev. 1