


In the Matter of: Progress Energy Florida, Inc.
(Levy County Nuclear Power Plant, Units 1 and 2)

	ASLBP #: 09-879-04-COL-BD01	Identified: 10/31/2012
	Docket #: 05200029 05200030	Withdrawn:
	Exhibit #: NRC050-00-BD01	Stricken:
	Admitted: 10/31/2012	
	Rejected:	
Other:		

Library of Congress Cataloging in Publication Data

Oceanic whitecaps and their role in air-sea exchange processes.



(Oceanic sciences library)

"Published in association with the Galway University Press."

"Proceedings of the 1983 Galway Whitecap Workshop"—CIP foreword.

Bibliography: p.

Includes index.

1. Ocean waves—Congresses. 2. Ocean-atmosphere interaction—Congresses. I. Monahan, Edward C., 1936— II. Mac Niocaill, Gearóid.
- III. Galway Whitecap Workshop (1983) IV. Series.

GC206.027 1986 551.47'02 86-6521

ISBN 90-277-2251-X

Published by D. Reidel Publishing Company
P.O. Box 17, 3300 AA Dordrecht, Holland

Sold and distributed in the U.S.A. and Canada
by Kluwer Academic Publishers,
190 Old Derby Street, Hingham, MA 02043, U.S.A.

In all other countries, sold and distributed
by Kluwer Academic Publishers Group,
P.O. Box 322, 3300 AH Dordrecht, Holland

This work is related to Department of the Navy Grant N00014-83-G-0110
issued by the Office of Naval Research.

The United States Government has a royalty-free license throughout the world
in all copyrightable material contained herein.

A MODEL OF MARINE AEROSOL GENERATION VIA WHITECAPS AND WAVE DISRUPTION

E. C. MONAHAN, D. E. SPIEL & K. L. DAVIDSON

University College, Galway,

BDM Corporation, Monterey, CA 93940,

Dept. of Meteorology, Naval Postgraduate School, Monterey, CA 93940

1. Introduction

We have, over the past several years, as one element in the development of a time-dependent model of the aerosol population of the marine atmospheric boundary layer, attempted to define, in terms of aerosol droplet radius (r) and 10m-elevation wind speed (U), a model of open-ocean sea-surface aerosol generation. This source function is represented by the expression $dF(r, U)/dr$, which states the rate of production of marine aerosol droplets, per unit area of the sea surface, per increment of droplet radius. In the initial modeling efforts only the indirect aerosol production mechanisms associated with the bursting of whitecap bubbles (see Figure 1) were considered. The model for sea surface aerosol generation by the indirect mechanisms, first introduced in our Canberra SSAG-1 (Monahan, et al, 1979) and Manchester SSAG-2 (Monahan, 1980) papers, is given by Equation 1, where W is the

$$(1) \quad dF_0/dr = W \tau^{-1} dE/dr$$

instantaneous fraction of the sea surface covered by whitecaps, τ is the time constant characterizing the exponential whitecap decay (measured in seconds), and dE/dr is the differential whitecap aerosol productivity, i.e. the number of droplets per increment droplet radius produced during the decay of a unit area of whitecap (expressed in $m^{-2} \mu m^{-1}$). The necessary expression for $W(U)$ was obtained from shipboard photographic observations of whitecaps (Monahan, 1971; Toba and Chaen, 1973),




Fig. 1. Schematic representation of the relative importance of various marine aerosol production mechanisms. The relative widths of the shaded columns, at any particular wind speed, are meant to indicate the relative significance of the direct and indirect aerosol production mechanisms represented by the various columns. Note that spilling and plunging waves both form whitecaps.

while values for τ and dE/dr were derived from measurements made using the University College, Galway, whitecap simulation tank.

The specific expression for $W(U)$ used in the latest model of dF_0/dr , SSAG-3 (Monahan, et al, 1982), and given in Equation 2, was extracted

$$W(U) = 3.84 \times 10^{-6} U^{3.41} \quad (2)$$