

## Statement on Tissue Reactions

Approved by the Commission on April 21, 2011

(1) The Commission issued new recommendations on radiological protection in 2007 (ICRP, 2007), which formally replaced the Commission's 1990 Recommendations (ICRP, 1991a). The revised recommendations included consideration of the detriment arising from non-cancer effects of radiation on health. These effects, previously called deterministic effects, are now referred to as tissue reactions because it is increasingly recognised that some of these effects are not determined solely at the time of irradiation but can be modified after radiation exposure. Previously, the Commission had reviewed various aspects of non-cancer health effects of low linear-energy-transfer (LET) ionising radiation in *Publication 41* (ICRP, 1984), high LET radiation in *Publication 58* (ICRP, 1990), the skin in *Publication 59* (ICRP, 1991b), and the skin and the eye in *Publication 85* (ICRP, 2000).

(2) The Commission has now reviewed recent epidemiological evidence suggesting that there are some tissue reaction effects, particularly those with very late manifestation, where threshold doses are or might be lower than previously considered. For the lens of the eye, the threshold in absorbed dose is now considered to be 0.5 Gy.

(3) For occupational exposure in planned exposure situations the Commission now recommends an equivalent dose limit for the lens of the eye of 20 mSv in a year, averaged over defined periods of 5 years, with no single year exceeding 50 mSv.

(4) Although uncertainty remains, medical practitioners should be made aware that the absorbed dose threshold for circulatory disease may be as low as 0.5 Gy to the heart or brain. Doses to patients of this magnitude could be reached during some complex interventional procedures, and therefore particular emphasis should be placed on optimisation in these circumstances.

(5) The Commission continues to recommend that optimisation of protection be applied in all exposure situations and for all categories of exposure. With the recent evidence, the Commission further emphasises that protection should be optimised not only for whole body exposures, but also for exposures to specific tissues, particularly the lens of the eye, and to the heart and the cerebrovascular system.

## REFERENCES

- ICRP, 1984. Nonstochastic effects of irradiation. ICRP Publication 41. Ann. ICRP 14 (3).
- ICRP, 1990. RBE for deterministic effects. ICRP Publication 58. Ann. ICRP 20 (4).
- ICRP, 1991a. 1990 Recommendations of the International Commission on Radiological Protection. ICRP Publication 60. Ann. ICRP 21 (1-3).
- ICRP, 1991b. Radiation effects on the skin. ICRP Publication 59. Ann. ICRP 22 (2).
- ICRP, 2000. Radiopathology of skin and eye and radiation risk. ICRP Publication 85. Ann. ICRP 30 (2).
- ICRP, 2007. The 2007 Recommendations of the International Commission on Radiological Protection. ICRP Publication 103. Ann. ICRP 37 (2-4).