

Delaware River Basin Commission

2012 Delaware River and Bay Water Quality Assessment



March 2012



This report provides an assessment of waters in the Delaware River and Bay for support of various designated uses in accordance with Section 305(b) of the Clean Water Act and identifies impaired waters, which consist of waters that do not meet Delaware River Basin Commission's (DRBC) Water Quality Regulations (18 CFR 410). It assesses data compiled from October 1, 2006 through September 30, 20011 (a five-year data window) into the 2012 Delaware River and Bay Water Quality Assessment (2012 Assessment). The assessment methodology used to develop the 2012 Assessment was noticed in the Federal Register on August 12, 2011 and published on DRBC's web site.

Table of Contents

Introduction and Overview	1
Background.....	1
Delaware River Basin.....	2
Delaware River Water Quality Assessment.....	4
Water Quality Standards	4
Designated Uses	5
Criteria	8
Assessment Methods	8
Assessment Units	9
Data Window	9
Data Sets.....	9
Analytical Parameters supporting Designated Uses	9
Data Requirements.....	9
Assessment Results	19
Aquatic Life.....	19
Public Water Supply	34
Contact Recreation.....	38
Fish Consumption	38
Shellfish Consumption.....	42
Assessment Summary.....	45
Recommendations for Future Action.....	46
Appendix A: Descriptions of DRBC Monitoring Programs	47
Appendix B: Other DRBC Water Pollution Control Programs.....	51
Appendix C: Fish Tissue Assessment Supplement.....	56
Appendix D: Toxic Pollutants Water Quality Assessment Details	67
Appendix E: Taste and Odor Water Quality Assessment Details	86
Appendix F: Public Participation Procedures	88

List of Tables

Table 1: Approximate Geographical Statistics for the Delaware River Basin.....	4
Table 2: Delaware River Water Quality Management (WQM) Zones	7
Table 3: Designated Uses by DRBC Water Quality Management Zones	8
Table 4: Aquatic Life data requirements and assessment criteria	11
Table 5: Public Water Supply data requirements and assessment criteria.....	14
Table 6: Primary Contact Recreation data requirements and assessment criteria.....	16
Table 7: Secondary Contact Recreation data requirements and assessment criteria	17
Table 8: Fish Consumption Data requirements and assessment criteria	18
Table 9: Shellfish Consumption data requirements and assessment criteria	19
Table 10: Aquatic Life Designated Use Assessment Results.....	20
Table 11: DO Assessment Results.....	21
Table 12: pH Assessment Results	24
Table 13: Turbidity Assessment Results	25
Table 14: Temperature Assessment Results	28
Table 15: TDS Assessment Results.....	30
Table 16: Alkalinity Assessment Results	31
Table 17: Summary of Biological Assessment Results.....	34
Table 18: Public Water Supply Designated Use Assessment Results	35
Table 19: Hardness Assessment Results.....	35
Table 20: Chlorides Assessment Results.....	36
Table 21: Primary and Secondary Contact Recreation Assessment Results	38
Table 22: Fish Consumption Advisory Summary	40
Table 23: Shellfish Consumption Designated Use Assessment Result	44
Table 24: Summary of the 2012 Assessment	45

List of Figures

Figure 1: Delaware River Basin.....	3
Figure 2: Delaware River Water Quality Management Zones / Assessment Units.....	6
Figure 3: Zone 6 Shellfish Management Assessment Units.....	7
Figure 4: DO Observations Compared to Criteria in Zone 3	23
Figure 5: pH Observations Compared to Criteria in Zone 1E.....	25
Figure 6: Turbidity Compared to Criteria in Zone 5.....	27
Figure 7: Water Temperature Observations Compared to Criteria in Zone 2	29
Figure 8: Shellfish Consumption Classifications designated by New Jersey and Delaware for the Delaware Bay (DRBC WQM zone 6)	43

Introduction and Overview

The 2012 Delaware River and Bay Water Quality Assessment (2012 Assessment) reports the extent to which waters of the Delaware River and Bay are attaining designated uses in accordance with Delaware River Basin Commission's Water Quality Regulations (18 CFR 410, DRBC WQR) for the period October 1, 2006 through September 30, 2011. The designated water uses to be protected within the Delaware Basin are as follows:

- 1) Agricultural, industrial, and public water supplies after reasonable treatment, except where natural salinity precludes such uses;
- 2) Wildlife, fish and other aquatic life;
- 3) Recreation;
- 4) Navigation;
- 5) Controlled and regulated waste assimilation to the extent that such use is compatible with other uses; and
- 6) Such other uses as may be provided by the Commission's Comprehensive Plan (2001).

The assessment involves comparison of several key water quality parameters with applicable DRBC water quality criteria. DRBC regulations designate public water supply, agricultural, and industrial uses for the Delaware River. Since the public water supply use is assessed and protective of the other uses, agricultural and industrial uses are not assessed separately for this report. For each designated use in each assessment unit, a number of water quality parameters, relevant to the use, are compared to the existing, applicable water quality criteria.

Background

This assessment follows previous similar efforts performed beginning in 1996 and published under the DRBC Water Quality Assessment Reports/305(b) banner of the DRBC web site at:
<http://www.state.nj.us/drbc/about/public/publications/index.html>

DRBC's water quality assessment report has been developed every even numbered year since its initiation. In the past several cycles, we referred to the report as an "Integrated List" water quality assessment report. This name referred to the reports developed by states for EPA which *integrated* the water quality assessment function, under section 305(b) of the Clean Water Act, together with the listing function, under section 303(d) of the Clean Water Act, into a single document. In the case of DRBC, this name was a mismatch. While DRBC does perform a water quality assessment function consistent with Section 305(b) of the Clean Water Act, only the states list water bodies not meeting standards, as per section 303(d). The DRBC does not list, but provides its assessment to the states for consideration in their listing determinations.

Pennsylvania, New Jersey, Delaware, and New York consider this assessment, in the context of their own EPA approved assessment and listing methodologies, to determine whether sections of the mainstem Delaware River should be listed on the state 303(d) list. Because their methodologies differ, listing decisions for shared waters are not automatically consistent.

Delaware River Basin

The Delaware River is the longest un-dammed river east of the Mississippi, extending from the confluence of its East and West branches at Hancock, N.Y. to the mouth of the Delaware Bay. The Delaware River is fed by 216 tributaries, the largest being the Schuylkill and Lehigh Rivers in Pennsylvania. In all, the basin contains approximately 13,500 square miles, draining parts of Pennsylvania (50.3 percent of the basin's total land area); New Jersey (23.3%); New York (18.5%); and Delaware (7.9%) (Figure 1).

Approximately 15 million people, or about 5% of the U.S. population, rely on the waters of the Delaware River Basin for drinking and industrial use, and the Delaware Bay is only a one to two hour drive away for about 20% of the people living in the United States. Yet the basin drains only four-tenths of one percent of the total continental U.S. land area. The population of the Delaware River Basin in 2010 stood at approximately 8.26 million people. Table 1 provides additional geographical statistics for the Delaware River Basin. The Delaware Bay and tidal reach of the Delaware River have been included in the National Estuary Program, a partnership initiative authorized by Section 320 of the Clean Water Act designed to protect estuarine systems of national significance.

Three reaches of the Delaware River have been included in the National Wild and Scenic Rivers System. One section extends 73 miles from the confluence of the river's East and West branches at Hancock, NY, downstream to Milrift, PA; the second is a 40-mile stretch from just south of Port Jervis, NY, downstream to the Delaware Water Gap near Stroudsburg, PA. The Lower Delaware Wild and Scenic Rivers Act, signed into law on November 1, 2000, adds approximately 65 miles of the Delaware and selected tributaries to the national system, linking the Delaware Water Gap and Washington Crossing, PA, just upstream of Trenton, N.J. Almost the entire non-tidal Delaware River (the portion north of the "fall line" at Trenton, NJ) is included in the National Wild and Scenic Rivers System. In addition, 35.4 miles of the Maurice River and its tributaries in New Jersey and approximately 190 miles of the White Clay Creek and its tributaries in Pennsylvania and Delaware have been included in the national system. Most recently, on December 22, 2006, President George W. Bush signed into law the Musconetcong Wild and Scenic Rivers Act, which designates 24.2 miles of the Musconetcong River (a tributary of the Delaware River located in New Jersey) as a component of the National Wild and Scenic Rivers System.

There are numerous economic benefits from the river. The Delaware River Port Complex (including docking facilities in Pennsylvania, New Jersey, and Delaware) is the largest freshwater port in the world. According to testimony submitted to a U.S. House of Representatives subcommittee in 2005, the port complex generates \$19 billion in annual economic activity. It is one of only 14 strategic ports in the nation transporting military supplies and equipment by vessel to support our troops overseas. The Delaware River and Bay is home to the third largest petrochemical port as well as five of the largest east coast refineries. Nearly 42 million gallons of crude oil are moved on the Delaware River on a daily basis. There are approximately 3,000 deep draft vessel arrivals each year and it is the largest receiving port in the United States for Very Large Crude Carriers (tank ships greater than 125,000 deadweight tons). It is the largest North American port for steel, paper, and meat imports as well as the largest importer of cocoa beans and fruit on the east coast. Over 65% of Chilean and other South American fruits imported into the United States arrive at terminal facilities in the tri-state port complex. Wilmington, Delaware

is home to the largest U.S. banana importing port, handling over one million tons of this cargo annually from Central America. According to Rear Admiral Sally Brice-O'Hara, District Commander of the Fifth Coast Guard District, "The port is critical not only to the region, but also to the nation."

Figure 1: Delaware River Basin



In addition, Dr. Gerald Kauffman of the University of Delaware has estimated that the Delaware River Basin provides \$25 billion annually in economic activity, including recreation, water quality, water supply, and hunting and fishing, \$21 billion annually in ecosystem goods and services (natural capital), and \$10 billion in annual wages.

Table 1: Approximate Geographical Statistics for the Delaware River Basin

Total Basin Land Area (mi ²) ^{a,b}	12,700
Population (2010)	8.26 million
Major River Basins (HUC 8) ^c	13
River Miles (Named) ^a	9,080
Border (Shared) River Miles ^a	339
Square Miles of Public Lakes and Reservoirs ^c	140
Square Miles of Estuary/Bay ^c	783
Square Miles of Wetlands ^c	480

^aDRBC GIS files

^bTotal Basin area minus area of Estuary and Bay

^cNational Hydrographic Dataset

Delaware River Water Quality Assessment

Water Quality Standards

Water quality standards provide a description of water body uses to be protected, as well as water quality criteria necessary to protect those uses. DRBC's water quality standards program derives its authority from Section 3.2 of the Delaware River Basin Compact (1961) which directs the Commission to adopt "a comprehensive plan...for the immediate and long range development and uses of the water resources of the basin" and to adopt "a water resources program, based upon the comprehensive plan, which shall include a systematic presentation of the quantity and quality of water resources needs of the area..."; and Section 5.2 which allows the Commission to "assume jurisdiction to control future pollution and abate existing pollution in the waters of the basin, whenever it determines...that the effectuation of the comprehensive plan so requires."

Designated Uses

Water uses are paramount in determining stream quality criteria, which, in turn, are the basis for determining discharge effluent quality requirements. Water quality standards require that all surface waters of the Basin be maintained in a safe and satisfactory condition for the following six (6) uses:

- 1) Agricultural, industrial and public water supplies after reasonable treatment, except where natural salinity precludes such uses;
- 2) Wildlife, fish and other aquatic life;
- 3) Recreation;
- 4) Navigation;
- 5) Controlled and regulated waste assimilation to the extent that such use is compatible with other uses; and
- 6) Such other uses as may be provided by the Commission's Comprehensive Plan.

The Delaware River and Bay consists of a non-tidal and tidal Zone. Zones C1-8 and intrastate streams (Zones E, W1, W2, N1 and N2) are not assessed in this report as they are assessed in the Integrated Reports of the Basin States. The non-tidal main stem consists of five Water Quality Management (WQM) Zones: 1A, 1B, 1C, 1D, and 1E (Figure 2). These Zones form the boundaries for the DRBC's assessment units (AUs) in the non-tidal Zone. The Zones as defined by river mile (RM) are included in Table 2. The designated uses applicable to the non-tidal AUs include aquatic life, fish consumption, primary contact recreation, and drinking water (Table 3).

The tidal Delaware River consists of AU 2, 3, 4, and 5 (Figure 2) and extends from RM 133.4 to RM 48.2 (Table 2). Assessment unit 6 (Delaware Bay) includes multiple units that are defined in part by shellfish management areas issued by the states of Delaware and New Jersey (Figure 3). The uses designated in the estuary and bay are indicated in Table 3. Shellfish consumption only applies to WQM Zone 6.

Figure 2: Delaware River Water Quality Management Zones / Assessment Units




Table 2: Delaware River Water Quality Management (WQM) Zones

WQM Zone	Location (as River Mile)
1A	330.7 – 289.9
1B	289.9 – 254.75
1C	254.75 – 217.0
1D	217.0 – 183.66
1E	183.66 – 133.4
2	133.4 – 108.4
3	108.4 – 95.0
4	95.0 – 78.8
5	78.8 – 48.2
6	48.2 – 0.0

Figure 3: Zone 6 Shellfish Management Assessment Units



Table 3: Designated Uses by DRBC Water Quality Management Zones

Designated Water Use	Water Quality Management Zone									
	1A	1B	1C	1D	1E	2	3	4	5	6
Aquatic Life	X	X	X	X	X	X	X	X	X	X
Public Water Supply	X	X	X	X	X	X	X			
Recreation										
Primary & Secondary	X	X	X	X	X	X		X ^A	X	X
Secondary only								X ^A	X ^A	
Fish Consumption	X	X	X	X	X	X	X	X	X	X
Shellfish Consumption										X

^A Primary recreation below RM 81.8; Secondary recreation above RM 81.8

Criteria

Sections 3.20, 3.30, and 3.40 of DRBC's Water Quality Regulations define the "Water Quality Objectives." From this point on, the objectives will be referred to as "Water Quality Criteria" (WQC) for the tidal and non-tidal river. Criteria are Zone-based and define the water quality necessary to protect the designated uses in those Zones. For the water quality assessments, monitored data are compared against the Zone standards for determining use attainment.

Zones 1, 2 and 3 of the Delaware River are given the designated use of "public water supplies after reasonable treatment." It is the general policy of DRBC that all ground water of the Basin, as well as surface sources of drinking water, should be maintained to support drinking water (18 CFR Part 410, 3.10.3.B, 3.40.4). In Zones 2 and 3, there is additional definition of the permissible levels of specific toxicants in waters designated for both drinking water and fish consumption (due to the bioaccumulation of certain substances even at very low ambient levels).

Assessment Methods

Because DRBC's role is to assess shared waters in the Basin, coordination with the Basin States is important. The Integrated Listing process defines a list of waters for which TMDLs must be prepared (i.e., 303(d) list). However, the regulatory responsibility for preparing a 303(d) list rests with the States. The DRBC did public notice the methodology for the 2012 Delaware River and Bay Water Quality Assessment in the Federal Register on August 11, 2011.

Assessment Units

As noted in the previous section, the non-tidal assessment units include WQM Zones 1A, 1B, 1C, 1D, and 1E (Figure 2). The designated uses assessed in Zones 1A through 1E include aquatic life, public water supply, primary recreation, and fish consumption (Table 3). WQM Zones 2, 3, 4, and 5 make up the tidal portion of the Delaware River Basin. Fish consumption, aquatic life, and recreation apply to all the tidal Zones. In the estuary, the public water supply use is only applicable to WQM Zones 2 and 3. The Delaware Bay consists of WQM Zone 6. The assessed designated uses for the Bay include aquatic life, primary recreation, fish consumption, and shellfish consumption.

Data Window

This assessment considers all readily available data collected in the 5-year period from October 1, 2006 through September 30, 2011.

Data Sets

This assessment considers all readily available data. To obtain the data, DRBC queried the EPA STORET database, the USGS NWIS database, the NOAA PORTS database, as well as internal DRBC databases. We also published a data solicitation in the Federal Register. The majority of the data considered is from the following monitoring programs and/or data sets:

- USGS continuous real time monitors via NWIS;
- USGS surface water monitoring programs via NWIS;
- DRBC / NPS Scenic Rivers Monitoring Program (SRMP);
- DRBC Biological Monitoring Program;
- DRBC Lower Delaware Monitoring Program;
- DRBC Boat Run monitoring program;
- NOAA PORTS continuous data;
- PAWQN Monitoring program via STORET;
- NY Department of Environmental Conservation, Division of Water, via STORET;
- NJDEP Bureau of Freshwater and Biological Monitoring via STORET;
- Delaware Department Of Natural Resources And Environmental Control via STORET;
- NJDEP Bureau of Marine Water Monitoring via STORET;
- Philadelphia Water Department Monitoring programs.

Other data sets contained in STORET were considered as well, but represented a small subset of the overall available data.

Analytical Parameters supporting Designated Uses

Data Requirements

This section looks at the general approach for each designated use assessed relative to DRBC water quality standards and other supporting evidence. The tables below also describe the

parameter-specific data requirements. It should be noted, however, that assessments might also be made using less robust data than indicated by the data requirements, when the weight of evidence is compelling.

Listed below are cases where insufficient data (ID) are available and the uses cannot be assessed against DRBC criteria. Such data would fail to support the designated use, but the assessment may be identified as “ID” rather than “not supported” when the following conditions exist:

- a) The number of samples per AU over an assessment period or season was below data requirements
- b) Background level was not specified in DRBC WQR and cannot reasonably be determined for a particular AU
- c) The parameter was not monitored in an AU
- d) The parameter was analyzed in a matrix other than surface water

Aquatic Life

Aquatic life is to be protected in all DRBC WQM Zones (Table 3). The assessment is based upon these water quality parameters: dissolved oxygen (DO), pH, turbidity, temperature, total dissolved solids (TDS), alkalinity, and in Zone 1, biological monitoring results (Table 4). In addition, toxic pollutants with criteria based on chronic and freshwater conditions are used to support aquatic life in Zones 2 through 5. For protection of aquatic life, Zone 6 was assessed as a whole unit.

Table 4: Aquatic Life data requirements and assessment criteria

Parameter	AU	Criteria	Assessment Method	Data Requirements
DO	All	Meet all Zone specific instantaneous minimum, minimum 24-hour average, spawning, and seasonal criteria listed in DRBC Water Quality Regulations, Sections 3.20 and 3.30	For instantaneous minimums, less than 1 observation plus 1 confirmatory observation per AU fail the criteria. For 24-hour averages, less than one 24-hour average plus one confirmatory 24-hour average fail the criteria.	For instantaneous minimums, at least 20 measurements over the assessment period. For 24-hour averages, at least 20 daily averages over the assessment period.
Temperature	1A-1E	Not to exceed Zone specific increases above ambient temperature	Estimate ambient temperature using data or models. Less than 1 observation plus 1 confirmatory observation per AU fail the criteria, considered in conjunction with the ambient temperature variability or model standard error.	At least 20 samples per AU over the assessment period
	2-6	Not to exceed Zone specific maximum temperatures listed in DRBC Water Quality Regulations, Sections 3.30 and 4.30	Less than 1 daily average plus 1 confirmatory daily average per AU fail the criteria	At least 20 samples per AU over the assessment period
pH	All	Meet Zone specific pH criteria range listed in DRBC Water Quality Regulations, Sections 3.20 and 3.30	Less than 1 observation plus 1 confirmatory observation per AU fail the criteria, unless evidence shows that pH violation are the result of natural conditions and biological communities are not impaired	At least 20 samples per AU over the assessment period
Turbidity	1A-1E, 2-3	Not to exceed Zone specific criteria listed in DRBC Water Quality Regulations, Sections 3.20 and 3.30	Less than 1 observation plus 1 confirmatory observation per AU fail the criteria	At least three samples in a 30-day period (AU 3) At least 20 samples per AU over the assessment period

TDS	1A-1E, 2-4	Not to exceed Zone specific TDS criteria listed in the DRBC Water Quality Regulations, Sections 3.20, 3.30 and 4.20.2	Less than 1 observation plus 1 confirmatory observation per AU fail the criteria	At least 20 samples per AU over the assessment period
Alkalinity	1E, 2-6	Meet Zone specific criteria range in DRBC Water Quality Regulations, Sections 3.20 and 3.30	Less than 1 observation plus 1 confirmatory observation per AU fail the criteria	At least 20 samples per AU over the assessment period
Toxic Pollutants	2-5	Not to exceed criteria noted in DRBC Water Quality Regulations, Table 5	No more than one (1) exceedence in an AU over a three year window	Available data
	1, 6	Not to exceed EPA recommended CCC criteria	No more than one (1) exceedence in an AU over a three year window	Available data
Biological Monitoring	1A – 1E	6-metric IBI not to fall below 75.6 unit threshold	No more than 30% of samples per AU below the threshold in the assessment period	At least 2 years of data with multiple sites per AU

Public Water Supply

The public water supply use is designated for WQM Zones 1A through 1E, 2, and 3. The parameters used for determining public water supply use support are:

- TDS;
- chlorides;
- toxic substances (human health criteria for systemic toxicants and carcinogens in Zones 2 and 3 only);
- hardness;
- odor;
- phenol;
- sodium (Na); and
- turbidity.

Since this particular use relates to human health, the assessment also takes into account information on actual impacts to the use such as frequent or extended closures of drinking water facilities due to recurring or chronic water quality concerns. Data requirements are shown in Table 5.

Contact Recreation

In the DRBC Water Quality Regulations, the "Recreation" designated use includes all water-contact sports, and thus corresponds to "primary contact" recreation. Some waters, however,

are designated as "Recreation - secondary contact" which restricts activities to where the probability of significant contact or water ingestion is minimal, encompassing but not limited to:

1. boating,
2. fishing,
3. those other activities involving limited contact with surface waters incident to shoreline recreation.

Criteria protective of the primary contact designated use are also protective of secondary contact uses. Criteria protective of secondary contact uses are not protective of primary contact uses. Contact recreation data requirements are shown in Tables 6 (Primary Contact) and 7 (Secondary Contact).

Primary

Primary contact recreation applies to Zones 1A-1E, 2, 4 below RM 81.8, and 5 and 6. The parameter used for determining primary contact recreation in Zones 1A-1E is fecal coliform. In addition to fecal coliform, enterococcus bacteria is used to assess primary contact recreation in the tidal Zones 2, 4, 5, and 6. Zone 4 is only assessed against primary contact standards below RM 81.8. The criteria are based on a geometric mean, with samples taken at a certain frequency and location as to permit valid interpretation.

Secondary

DRBC WQM Zones 3 and 4 above RM 81.8 are restricted to secondary contact recreation. Fecal coliform and enterococcus bacteria are used to assess secondary contact recreation. Zone 4 is assessed against secondary contact standards above RM 81.8. The criteria are based on a geometric mean, with samples taken at a certain frequency and location as to permit valid interpretation.

Table 5: Public Water Supply data requirements and assessment criteria

Parameter	AU	Criteria	Assessment Method	Data Requirements
TDS	1A-1E, 2-3	Not to exceed Zone specific TDS criteria listed in the DRBC Water Quality Regulations, Sections 3.20, 3.30 and 4.20.2	Less than 1 observation plus 1 confirmatory observation per AU fail the criteria	At least 20 samples per AU over the assessment period
Hardness	2-3	Not to exceed Zone specific 30-day average criteria listed in DRBC Water Quality Regulations, Section 3.30.2 and 3.30.3	Less than 1 observation plus 1 confirmatory observation per AU fail the criteria	At least three samples in a 30-day period At least 20 samples per AU over the assessment period
Chlorides	2-3	Not to exceed Zone specific criteria listed in DRBC Water Quality Regulations, Section 3.30.2 and 3.30.3	Less than 1 observation plus 1 confirmatory observation per AU fail the criteria	At least two samples in a 15-day period (AU 2) At least three samples in a 30-day period (AU 3) At least 20 samples per AU over the assessment period
Odor	1A-1E, 2-3	Not to exceed Zone specific criteria listed in DRBC Water Quality Regulations, Sections 3.20 and 3.30	Less than 1 observation plus 1 confirmatory observation per AU fail the criteria	Available data
Phenols	1A-1E, 2-3	Not to exceed Zone specific criteria listed in DRBC Water Quality Regulations, Section 3.20 and 3.30	Less than 1 observation plus 1 confirmatory observation per AU fail the criteria	At least 20 samples per AU over the assessment period
Na	3 at or above RM 98	Not to exceed 30-day average criteria listed in DRBC Water Quality Regulations, Section 3.30.3	Less than 1 observation plus 1 confirmatory observation per AU fail the criteria	At least three samples in a 30-day period (AU 3) At least 20 samples per AU over the assessment period
Turbidity	1A-1E, 2-3	Not to exceed Zone specific criteria listed in DRBC Water Quality Regulations, Sections 3.20 and 3.30	Less than 1 observation plus 1 confirmatory observation per AU fail the criteria	At least three samples in a 30-day period (AU 3) At least 20 samples per AU over the assessment period

Systemic Toxicants	2-3	Not to exceed criteria listed in DRBC Water Quality Regulations, Section 3.30, Table 7	No more than one (1) exceedence in an AU over a three year window	Available data
Carcinogens	2-3	Not to exceed criteria listed in DRBC Water Quality Regulations, Section 3.30, Table 6	No more than one (1) exceedence in an AU over a three year window	Available data
Drinking Water Closures	1A-1E, 2-3	No frequent or extended closures of drinking water facilities due to recurring or chronic water quality concerns	No closures affecting an AU over over the assessment period	Administrative closures for drinking water supply over the assessment period. Information from one or more drinking water intake facility per AU.

Table 6: Primary Contact Recreation data requirements and assessment criteria

Parameter	AU ^A	Criteria	Assessment Method	Data Requirements
Fecal coliform	1A-1E,2,4 (below RM 81.8),5,6	Not to exceed Zone specific Fecal coliform criteria listed in the DRBC Water Quality Regulations, Sections 3.20 and 3.30	Geometric mean of samples per AU during each assessment year	At least 5 samples per AU during each assessment year
Enterococcus	2,4 (below RM 81.8)	Not to exceed Zone and sub-Zone specific Enterococcus criteria listed in the DRBC Water Quality Regulations, Section 3.30	Geometric mean of samples per AU during each assessment year	At least 5 samples per AU during each assessment year
	5,6	Not to exceed Zone and sub-Zone specific Enterococcus criteria listed in the DRBC Water Quality Regulations, Section 3.30	Geometric mean of samples per AU during each assessment year	At least 5 samples per AU during each assessment year

^AWQM Zone 4 is assessed for the parameters below RM 81.8.

Table 7: Secondary Contact Recreation data requirements and assessment criteria

Parameter	AU ^A	Criteria	Assessment Method	Data Requirements
Fecal coliform	3,4 (above RM 81.8)	Not a single geometric mean to exceed 770 / 100 ml	Geometric mean of samples per AU during each assessment year	At least 5 samples per AU during each assessment year
Enterococcus	3,4 (above RM 81.8)	Not a single geometric mean to exceed 88 / 100 ml	Geometric mean of samples per AU during each assessment year	At least 5 samples per AU during each assessment year

^AWQM Zone 4 is assessed for the parameters above RM 81.8.

Fish Consumption

The fish consumption designated use applies to all DRBC WQM Zones. The assessment criterion is based primarily on the presence of the Basin states' fish consumption advisories in the mainstem Delaware River and Bay for the 2010 to 2011 assessment period. The presence of fish consumption advisories results in an assessment of "not supporting the designated use". Fish Consumption data requirements are shown in Table 8.

Table 8: Fish Consumption Data requirements and assessment criteria

Parameter	AU	Criteria	Assessment Method	Data Requirements
Fish Consumption Advisory	1A-1E, 2-6	Not a single fish advisory listed for an AU	Count of the number of fish consumption advisories per AU listed over the assessment period	NY, NJ, DE, and PA fish consumption advisories for the general population based upon the Basin states' water quality or fish tissue data

Shellfish Consumption

Shellfish consumption designated use only applies to DRBC WQM zone 6 (RM 48.2 to the mouth of the Delaware Bay). New Jersey and Delaware assess this use in their coastal waters, using procedures developed by the FDA National Shellfish Sanitation Program (NSSP). Both states use total coliform (as most probable number) as the assessment tool and compare it against federal shellfish standards.

In both states, waters classified for shellfishing may be opened for that use all year round. In some cases, the AU is opened seasonally (typically in winter). In other cases, harvesting may be prohibited due to administrative closures based upon proximity to sewer outfalls. In still other cases, waters may be open to harvesting, but with special treatment of the shellfish, such as transplantation to cleaner waters for a period of time prior to the harvesting. Finally, some waters are closed to shellfish harvesting due to existing water quality concerns. Shellfish Consumption data requirements are shown in Table 9.

Table 9: Shellfish Consumption data requirements and assessment criteria

Parameter	AU ^A	Criterion	Assessment Method	Data Requirements
Shellfish Consumption Classifications	6	No prohibitions and/or year-round closures in an AU. Shellfish waters with special conditions and temporal windows are assessed as supporting but with conditions	Determine the number of shellfish harvesting prohibitions, year-round closures, and limiting conditions per AU listed over the assessment period	DE and NJ shellfish consumption and harvesting advisories, prohibitions, closures, and limiting conditions per AU over the assessment period

^A WQM zone 6 is subdivided into multiple units based on Shellfish Management Directives.

Assessment Results

The following sections provide the 2012 assessment results for the designated uses:

- Aquatic Life;
- Public Water Supply;
- Contact Recreation;
- Fish Consumption; and
- Shellfish Consumption.

Meaningful assessment continues to be hampered by the requirement to indicate 1 exceedance plus 1 confirmatory exceedance as not meeting criteria and subsequently not supporting a designated use.

Aquatic Life

The Aquatic Life Assessment results are presented in Table 10 below. The composite aquatic life assessment for 2012 yields a result of “Not supporting” for all assessment units. It is important to note, however, that this result is largely driven by the requirement to categorize as not meeting criteria any assessment unit with 1 exceedance plus 1 confirmatory exceedance.

Table 10: Aquatic Life Designated Use Assessment Results

Zone (Assessment Unit)	DO	pH	Turbidity	Temperature	TDS	Alkalinity	Toxic Pollutants	Biological Assessment	2012 Composite	2010 Composite
1A	+	+	- ^A	N/A ^C	+	N/A	- ^F	+	NS ^E	NS
1B	- ^A	- ^A	- ^A	N/A ^C	+	N/A	- ^F	+	NS ^E	NS ^E
1C	+	- ^A	+	N/A ^C	+	N/A	- ^F	+	NS ^E	ID
1D	+	- ^A	- ^A	N/A ^C	+	N/A	- ^F	+	NS ^E	NS ^E
1E	+	-	- ^A	N/A ^C	+	- ^A	+ ^F	-	NS ^E	NS
2	- ^A	- ^A	- ^A	- ^{A, B}	+	+	+	NC	NS ^E	NS
3	- ^A	+	+	- ^{A, B}	+	- ^A	+	NC	NS ^E	NS ^E
4	+	+	+	- ^{A, B}	N/A ^D	+	+	NC	NS ^E	NS ^E
5	- ^A	+	-	- ^{A, B}	N/A	+	-	NC	NS ^E	NS ^E
6	- ^A	- ^A	- ^A	- ^{A, B}	N/A	- ^A	-	NC	NS ^E	NS ^E

Notes:

A – Rate of criteria exceedance is below the historical threshold of 10%.

B – Temperature criteria exceedance may be driven, in part, by meteorologic and atmospheric conditions. The proportion of temperature exceedance caused by controllable anthropogenic inputs is unknown at this time.

C – Temperature criteria in Zone 1A through 1E are expressed relative to ambient temperature, but ambient temperature is not defined. We interpret these criteria to be applicable to thermal mixing zones. Therefore, Zones 1A through 1E lack a surface water quality standard for temperature.

D – Criteria expressed relative to background, but background is undefined.

NC – No criteria developed.

E – Based primarily on fewer than 10% exceedances of criteria

F – The Assessment Unit meets (+) or does not meet (-) EPA's 2006 National Recommended Water Quality Criteria, where numerical criteria have not yet been adopted

NS – The assessment does not support the designated use

N/A – The parameter is not applicable in this assessment unit

Dissolved Oxygen

Dissolved oxygen (DO) refers to the concentration of oxygen gas incorporated in water. Oxygen enters water both by direct absorption from the atmosphere, which is enhanced by turbulence, and as a by-product of photosynthesis from algae and aquatic plants. Sufficient DO is essential to growth and reproduction of aerobic aquatic life. Oxygen levels in water bodies can be depressed by the discharge of oxygen-depleting materials (measured in aggregate as biochemical oxygen demand, BOD, from wastewater treatment facilities), from the decomposition of organic matter including algae generated during nutrient-induced blooms, and from the oxidation of ammonia and other nitrogen-based compounds.

Table 11 below shows the assessment results for DO for all Zones. All criteria were met in Zones 1E and Zone 4. In Zones 1A, 1C, and 1D, all instantaneous minima criteria were met. Since only daytime spot measurements were made in Zones 1A through 1D, attainment of the 24-hour mean criteria was presumed since all measurements were above (met) that criteria. All seasonal mean criteria were met in Zones 2 through 5. The majority of observations met criteria in Zones 1B, 2, 3, and 5.


Table 11: DO Assessment Results

Zone	% Observations Meeting Daily Mean Criteria	% Meeting Seasonal Criteria	% Meeting Instantaneous Minimum Criteria	Primary Data Source(s)	Notes
1A	100% (presumed)	N/A	100%	<ul style="list-style-type: none"> • SRMP Monitoring 	Daytime spot measurements only
1B	98.3% (presumed)	N/A	98.3%	<ul style="list-style-type: none"> • SRMP Monitoring • NYS Dept. of EnCon, Division of Water 	Daytime spot measurements only
1C	100% (presumed)	N/A	100%	<ul style="list-style-type: none"> • SRMP Monitoring 	Daytime spot measurements only
1D	100% (presumed)	N/A	100%	<ul style="list-style-type: none"> • SRMP / Lower Delaware Monitoring • NJDEP Bureau of Freshwater and Biological Monitoring 	Daytime spot measurements only
1E	100%	N/A	100%	<ul style="list-style-type: none"> • USGS 01463500 Delaware River at Trenton NJ 	
2	98%	100%	N/A	<ul style="list-style-type: none"> • USGS 01467029 Delaware River div at Delran NJ 	No data before September 2008

3	99.6%	100%	N/A	<ul style="list-style-type: none"> • USGS 01467200 Delaware R at Ben Franklin Bridge at Philadelphia 	
4	100%	100%	N/A	<ul style="list-style-type: none"> • USGS 01477050 Delaware River at Chester, PA 	
5	96%	100%	N/A	<ul style="list-style-type: none"> • USGS 01482800 Delaware River at Reedy Island Jetty, DE 	
6	90.9% (presumed)	N/A	99.2%	<ul style="list-style-type: none"> • DRBC Boat Run • Delaware Department Of Natural Resources And Environmental Control • NJDEP Bureau of Marine Water Monitoring 	Daytime spot measurements only

Determining whether 24-hour criteria were met is most appropriately accomplished by comparing the daily mean DO from continuous monitors, which record data hourly or sub-hourly, and comparing these computed results to the criteria. Where only daytime spot measurements are available, we presumed that if the measured value is less than the 24-hour mean criterion, then the 24-hour mean is also likely to be below (not meet) the criterion. Zones 1A, 1B, 1C, 1D and 6 currently lack continuous water quality meters, and were assessed using daytime spot measurements only. Figure 4 below shows a comparison of daily mean and seasonal mean DO observations at the USGS monitor at the Ben Franklin Bridge (Zone 3) to the 24-hour mean and seasonal criteria.

Figure 4: DO Observations Compared to Criteria in Zone 3



As shown in Table 11, the vast majority of the measurements meet criteria. The DO assessment in particular highlights the flaw associated with making a determination of not meeting criteria on the sole basis of one exceedance plus one confirmatory exceedance. Sparse, periodic exceedances may indeed negatively impact aquatic life, but more work is needed to determine the frequency and duration of exceedance that would cause this impact.

pH

The pH of surface waters has long been recognized as both a natural and human-induced constraint to the aquatic life of fresh and salt water bodies, both through direct effects of pH and through indirect effects on the solubility, concentration, and ionic state of other important chemicals (e.g., metals, ammonia). Among natural waters, both highly alkaline waters and highly acidic waters (like the NJ Pinelands) are known to severely restrict the species of plants and animals that can thrive in particular lakes and streams. Likewise, human alteration of the pH regimen for a water body can alter both the quality of that water and the aquatic life inhabiting that system. Table 12 below shows the assessment results for pH for each Zone.


Table 12: pH Assessment Results

Zone	% Observation Days Meeting Criteria	Primary Data Source(s)	Notes
1A	100%	<ul style="list-style-type: none"> • SRMP Monitoring • USGS field monitoring 	Daytime spot measurements only
1B	95.2%	<ul style="list-style-type: none"> • SRMP Monitoring • USGS field monitoring • NYS Dept. of EnCon, Division of Water 	Daytime spot measurements only
1C	97.6%	<ul style="list-style-type: none"> • SRMP Monitoring • USGS field monitoring 	Daytime spot measurements only
1D	95.4%	<ul style="list-style-type: none"> • SRMP / Lower Delaware Monitoring • NJDEP Bureau of Freshwater and Biological Monitoring 	Daytime spot measurements only
1E	77.4%	<ul style="list-style-type: none"> • USGS 01463500 Delaware River at Trenton NJ 	
2	99.9%	<ul style="list-style-type: none"> • USGS 01467029 Delaware River div at Delran NJ 	No data before September 2008
3	100%	<ul style="list-style-type: none"> • USGS 01467200 Delaware R at Ben Franklin Bridge at Philadelphia 	
4	100%	<ul style="list-style-type: none"> • USGS 01477050 Delaware River at Chester, PA 	
5	100%	<ul style="list-style-type: none"> • USGS 01482800 Delaware River at Reedy Island Jetty, DE 	
6	97.9%	<ul style="list-style-type: none"> • DRBC Boat Run • Delaware Department Of Natural Resources And Environmental Control 	Daytime spot measurements only

As shown in Table 12, pH criteria were met in Zones 1A, 3, 4, and 5. pH criteria were mostly met, with the exception of Zone 1E, where daily pH maximum values routinely exceeded the maximum criterion of 8.5, as shown in Figure 5. In Zones 1A through 1D, pH assessment is hampered by the lack of continuous monitors. Like DO, pH has a diel cycle, with the lowest pH values expected in the early morning hours or pre-dawn, and the highest pH values expected in the mid to late afternoon. Monitoring programs that rely on spot measurements are far more likely to capture daytime high values, and miss pre-dawn low values.

As noted in previous assessments, we believe our existing criteria fail to recognize that pH swings outside the range between 6.5 and 8.5 could occur due to natural conditions. We are in the process of developing recommended revised criteria. Yet extreme swings in pH can indicate excessive plant and algae growth resulting from nutrient stimulation. As a result, diel pH cycles may also be part of the nutrient criteria development process.

Figure 5: pH Observations Compared to Criteria in Zone 1E



Turbidity

According to Standard Methods (2005), “Turbidity in water is caused by suspended and colloidal matter such as clay, silt, finely divided organic and inorganic matter, and plankton and other microscopic organisms. Turbidity is an expression of the optical property that causes light to be scattered and absorbed rather than transmitted with no change in direction or flux level through the sample.” From an observational perspective, water with low turbidity appears clear, while higher turbidity waters are cloudy or muddy. Table 13 below shows the assessment results for each Zone relative to DRBC’s turbidity criteria.

Table 13: Turbidity Assessment Results

Zone	% Observation Meeting Max Criteria	% Meeting 30-day Average Criteria	Primary Data Source(s)	Notes
1A	95.2%	Insufficient Data to Assess	<ul style="list-style-type: none"> SRMP Monitoring USGS field monitoring 	Spot measurements only
1B	99.0%	Insufficient Data to Assess	<ul style="list-style-type: none"> SRMP Monitoring USGS field monitoring NYS Dept. of EnCon, Division of Water 	Spot measurements only
1C	100%	Insufficient Data to Assess	<ul style="list-style-type: none"> SRMP Monitoring USGS field monitoring 	Spot measurements only
1D	99.1%	Insufficient Data to Assess	<ul style="list-style-type: none"> SRMP / Lower Delaware Monitoring USGS field monitoring 	Daytime spot measurements only

			<ul style="list-style-type: none"> NJDEP Bureau of Freshwater and Biological Monitoring 	
1E	98.4%	98.9%	<ul style="list-style-type: none"> USGS 01463500 Delaware River at Trenton NJ 	
2	97.8%	97.9%	<ul style="list-style-type: none"> USGS 01467029 Delaware River div at Delran NJ 	No continuous monitor data before September 2008
3	100%	100%	<ul style="list-style-type: none"> USGS 01467200 Delaware R at Ben Franklin Bridge at Philadelphia 	
4	100%	Insufficient Data to Assess	<ul style="list-style-type: none"> DRBC Boat Run 	No Turbidity on USGS Monitor, spot measurements only
5	37%	0%	<ul style="list-style-type: none"> USGS 01482800 Delaware River at Reedy Island Jetty, DE 	No USGS data before April 2009
6	99.7%	Insufficient Data to Assess	<ul style="list-style-type: none"> DRBC Boat Run Delaware Department Of Natural Resources And Environmental Control 	Spot measurements only


DRBC's turbidity criteria include numerical limits for each zone, which should not be exceeded, "unless exceeded due to natural conditions." In Zone 1E, where both discharge and turbidity are monitored and recorded at the USGS monitor at Trenton (01463500), we compared flow on days when criteria were exceeded to the historical record from 1986 to the present (representing the current flow management regime). Of the 26 days where turbidity criteria were exceeded, 16 days had mean discharge in excess of the 95th percentile of mean daily flow values. The remaining 10 days with turbidity above criteria had mean daily flow below the 90th percentile of flow, and in fact, August 21 and 22, 2007 and June 5, 2008 all registered flow values below the median flow. It is reasonable to conclude that high turbidity events coinciding with high flow are due to natural conditions, but high turbidity events at lower flows are more problematic. One possibility is that localized intense storms send pulses of turbid water into the mainstem. These localized events may push turbidity over criteria without concurrent elevated mainstem flows. The uncertainty of this assessment is compounded by the role that landscape modification may play in mobilizing sediment. Because we cannot attribute all the high turbidity events to high flow, we conclude that some high turbidity events constitute of violation of criteria.

In previous assessment cycles, only spot measurements were available for turbidity assessment in Zone 5. In April 2009, USGS added turbidity to the monitor at Reedy Island Jetty (01482800). In reviewing these data, it became clear that turbidity in Zone 5 is largely not meeting criteria. In fact, no rolling 30-day mean during the data period met the 30-day mean criteria, and only 37% of observation days met the instantaneous maximum criteria. Figure 6 below shows the daily minimum and maximum observed

turbidity at Reedy Island, as well as the computed daily mean and the rolling 30-day mean (where 30 days of uninterrupted data are available) compared to criteria. However, this mismatch between measured turbidity and criteria may indicate a problem with the criteria rather than a water pollution problem. Zone 5 of the estuary spans the well documented estuary turbidity maximum (ETM) for the Delaware. ETM's are naturally occurring features of most estuaries, and typically occur near the toe of the salt wedge. Again, however, it is unclear how anthropogenic drivers, such as vessel traffic and dredging impact the natural turbidity regime in the ETM, and what level of turbidity supports an aquatic life use.

As a result, we recommend additional coordination with physical oceanographers and estuarine ecologists to determine whether the existing turbidity criterion in Zone 5 is relevant and protective, or whether revision of the current criteria is warranted.

Figure 6: Turbidity Compared to Criteria in Zone 5



Temperature

Water temperature is an important factor for the health and survival of native fish and aquatic communities. Temperature can affect embryonic development; juvenile growth; adult migration; competition with non-native species; and the relative risk and severity of disease. Estuary Temperature Criteria are expressed in DRBC regulations by day of year in Zones 2, 3, and 4. In Zones 5 and 6, a single maximum water temperature is specified. Table 14 below shows that water temperature criteria were mostly met, with the greatest number of exceedances occurring in Zone 2, and decreasing exceedances moving down-estuary toward the mouth of the Bay. The decreasing frequency of exceedances may be due to the shift from urbanized to rural watershed, the heat buffering capacity of the ocean boundary, and the change in criteria format in Zones 5 and 6.


As noted in previous assessments, criteria in Zones 1A through 1E are clearly oriented toward determining compliance of thermal mixing zones for point discharges. Currently, DRBC has no ambient surface water temperature standards in Zones 1A through 1E. In previous assessments, we investigated approaches for assessing surface waters in the non-tidal river against the thermal mixing zone criteria. None of these approaches was workable. DRBC continues work on development of ambient temperature criteria in the non-tidal river, as well as clarifying language regarding the application of the existing criteria.

Table 14: Temperature Assessment Results

Zone	% Observation Days Meeting Criteria	Primary Data Source(s)	Notes
1A	Criteria applicable to Heat Dissipation Areas only for Zone 1 AU's		
1B			
1C			
1D			
1E			
2	92.2%	<ul style="list-style-type: none"> • Newbold (NOAA PORTS) • Burlington (NOAA PORTS) • Delran (USGS NWIS) 	
3	93.7%	<ul style="list-style-type: none"> • Philadelphia (NOAA PORTS) • Ben Franklin Bridge (USGS NWIS) • Tacony Palmyra (NOAA PORTS) 	
4	94.5%	<ul style="list-style-type: none"> • Marcus Hook (NOAA PORTS) • Chester (USGS NWIS) • Fort Mifflin (USGS NWIS) 	
5	98.8%	<ul style="list-style-type: none"> • Reedy Island (USGS NWIS) • Reedy Point (NOAA PORTS) 	
6	99.7%	<ul style="list-style-type: none"> • Brandywine Shoal (NOAA PORTS) • Lewes (NOAA PORTS) • Ship John Shoal (NOAA PORTS) 	

Figure 7 below shows the comparison of water temperature observations in Zone 2 to the day-of-year temperature criteria. Note that observations in Zone 2 include continuous monitor data from a USGS monitor at Delran, and two different NOAA PORTS stations.

Figure 7: Water Temperature Observations Compared to Criteria in Zone 2



As also noted in previous assessments, atmospheric temperatures and meteorological conditions are strong drivers of water temperature. DRBC previously demonstrated that water temperatures are strongly linked to air temperatures, and that a notable increase in air temperatures is observable between the temperature gradient period (1961-1966) and the current period. At present, we lack the tools to determine which portion of the exceedance is attributable to potentially controllable anthropogenic thermal inputs, and which portion is due to meteorological drivers beyond our control.

Total Dissolved Solids (TDS)

Total Dissolved Solids (TDS) reflects the concentration of solids in a water sample capable of passing through a filter (typically 2 um) and dried. As an analytical parameter, TDS represents the collective mass of individual constituents, including cations, anions, and dissolved organic material. Studies have shown that high TDS concentrations negatively impact aquatic life and cause shifts in biological communities. In freshwater, TDS is frequently used as an indicator of the anthropogenic burden.

Table 15 below shows the TDS criteria in Zones 1A through 3 were met, with no Zone showing more than 1 exceedance. The TDS criteria in Zone 4 is expressed only as a percentage above background, and background in Zone 4 has not been defined. DRBC has no TDS criteria in Zones 5 and 6, presumably because TDS in marine waters is naturally high.

Table 15: TDS Assessment Results

Zone	% Observations Meeting Criteria	Primary Data Source(s)	Notes
1A	100%	<ul style="list-style-type: none"> • SRMP Monitoring 	
1B	100%	<ul style="list-style-type: none"> • SRMP Monitoring • NYS Dept. of EnCon, Division of Water 	
1C	99.6%	<ul style="list-style-type: none"> • SRMP Monitoring 	1 exceedance only, no confirmatory exceedance
1D	100%	<ul style="list-style-type: none"> • SRMP / Lower Delaware Monitoring 	
1E	98.4%	<ul style="list-style-type: none"> • SRMP / Lower Delaware Monitoring 	1 exceedance only, no confirmatory exceedance
2	100%	<ul style="list-style-type: none"> • DRBC Boat Run • PWD Boat based monitoring 	
3	98.9%	<ul style="list-style-type: none"> • DRBC Boat Run • PWD Boat based monitoring 	1 exceedance only, no confirmatory exceedance
4	N/A		Criteria relative only to background, background not defined
5	No Criteria		
6			

Alkalinity

According to Standard Methods (2005), “alkalinity of a water is its acid-neutralizing capacity. It is the sum of all the titratable bases.” As shown in Table 16 below, attainment of alkalinity criteria exceeded 96% in all Zones where criteria have been adopted.

Table 16: Alkalinity Assessment Results

Zone	% Observations Meeting Criteria	Primary Data Source(s)	Notes
1A	No Criteria		
1B			
1C			
1D			
1E	96.8%	<ul style="list-style-type: none"> • SRMP / Lower Delaware Monitoring 	
2	100%	<ul style="list-style-type: none"> • DRBC Boat Run • PWD Boat based monitoring 	1 observation at criteria but not below
3	97.9%	<ul style="list-style-type: none"> • DRBC Boat Run • PWD Boat based monitoring 	
4	100%	<ul style="list-style-type: none"> • DRBC Boat Run • PWD Boat based monitoring 	
5	100%	<ul style="list-style-type: none"> • DRBC Boat Run 	
6	98.6%	<ul style="list-style-type: none"> • DRBC Boat Run 	

Toxic Pollutants

The Delaware River Basin Commission (DRBC) updated stream quality objectives for human health and aquatic life in the tidal portion of the Delaware River Basin from the head of tide at Trenton, NJ to the Delaware Bay (Zones 2 through 5) and adopted stream quality objectives for toxic pollutants to uniformly apply in the estuary to the mouth of the Delaware Bay (Zone 6) in 2011. The changes in criteria reflect new scientific information and harmonize DRBC criteria with basin states’ criteria. The updated criteria are used in the 2012 assessment for Zones 2 through 6. As described in Methodology for the 2012 Delaware River and Bay Water Quality Assessment Report, the DRBC compared observations in Zones 1A through 1E (where DRBC has not adopted numerical criteria) to EPA criteria, as a method of implementing DRBC’s narrative standard.

Data Quantity and Quality

Water quality monitoring data from multiple organizations (DRBC, DNREC, NYSDEC, NJDEP, and USGS) were included in the 2012 assessment. Toxic pollutants data reviewed was collected using EPA approved methods or equivalent methods. The level of monitoring varied by Zone with more monitoring for toxics reported in Zones 2 through 6 than Zone 1. The level of monitoring also varied by toxic pollutant with >600 data points for dissolved copper and > 50 data points for methyl mercury including submitted QA data were reviewed as part of the assessment while cadmium and aluminum

had approximately a dozen data points each. DRBC toxics pollutants monitored during the timeframe of the assessment are listed in Appendix D.

Use of Freshwater or Marine Stream Quality Objectives

DRBC regulations include aquatic life toxics criteria for fresh and marine waters. As a policy, freshwater criteria will apply in all areas of the estuary upstream of the Delaware Memorial Bridges. In the main stem Delaware River below the Delaware Memorial Bridges and above Liston Point (RM 48.2, the downstream limit of Zone 5), the more stringent of the freshwater or marine criteria will apply. Downstream from Liston Pt., the marine criteria are used.

- A supplemental assessment was done in 2012. Site-specific paired salinity measured between RM 69.7 and 48.2 concurrently with toxic analytical parameters confirmed that when exceedances of freshwater objectives occurred ambient conditions were < 5 ppt salinity and when exceedances of marine objectives occurred ambient conditions were ≥ 5 ppt salinity.

Dissolved Metals

For criteria expressed as the dissolved form of the metal, assessment of monitoring data is as follows:

- In assessment Zones with dissolved metals data collected, direct comparison to DRBC dissolved criteria is the preferred assessment.
- In assessment Zones with only total metals data collected (as noted in Appendix D), comparison of total metals data to estimated total metals criteria using conversion factors listed in "Revised Procedure for Converting Total Recoverable Water Quality Criteria for Metals to Dissolved Criteria" <http://www.state.nj.us/drbc/regs/critmetals.pdf>.

Hardness Dependent Stream Quality Objectives

Some criteria require hardness values to compute the actual criteria numeric value. In these cases, toxics data from ambient water are compared to stream quality objectives using hardness values listed in DRBC Water Quality Regulations (i.e., 74 mg/L as CaCO₂).

- An additional comparison was conducted as part of the 2012 assessment whereby exceedances of DRBC regulatory hardness dependent values were confirmed with site-specific paired hardness measured concurrently with toxic analytical parameters.

Exceedances in Zones 5 and 6

- Data showed exceedances of DRBC acute marine stream quality objective for **copper** in Zones 5 and 6 as well as exceedances of the DRBC chronic freshwater stream quality objective for copper in Zone 5. Assessment of metals in ambient water is complicated by factors such as field sampling and analytical issues with contamination, the applicability of DRBC's freshwater or marine criteria, and the influence of other water quality attributes that influence the partitioning and toxicity of copper. The DRBC has scheduled additional data collection using enhanced analytical methods, modified collection procedures and changes in the spatial scale of sampling in segments of the Delaware River which have exhibited apparent copper exceedances. The information collected will provide additional data to determine metal concentrations in ambient water and the impact of metals on water quality. The survey targets copper, zinc and nickel. Coordination among basin states and agencies should continue to ensure the

use of the most appropriate methods and procedures for the conduct of monitoring studies in the Basin, and the harmonization of water quality criteria and assessment methodologies.

- Data showed exceedances of DRBC stream quality objectives for **methyl mercury** in Zones 5 and 6. The methylmercury criterion is fish tissue residue based as recommended by USEPA. No exceedances were observed in the fish species monitored by the DRBC in tidal and non-tidal waters (channel catfish, white perch, smallmouth bass and white sucker). DNREC database includes Atlantic croaker, tautog, and striped bass with exceedances for striped bass in Zones 5 and 6. DRBC staff is soliciting data on methyl mercury in fish tissue sampled from the Delaware River. In addition, comments are requested on implementation procedures for the tissue-based methylmercury criterion.

Exceedances in Zone 1

Data showed exceedances of acute EPA criteria for **cadmium** in Zone 1B and exceedances of chronic EPA criteria for **aluminum** in Zones 1A, 1B and 1D.

PADEP database showed exceedances of EPA water quality criteria based on fish tissue residue for **methyl mercury** in smallmouth bass and American eel for Zones 1C , 1D and West Branch .

Biological Assessment

Biological assessment results indicate reference-quality invertebrate communities in Zones 1A, 1B, 1C, and 1D. This includes “attaining” scores in the thermally altered upper mainstem Delaware River between Hancock and Callicoon (i.e., the upper portion of Zone 1A). Only a single sample in Zone 1C fell below the impairment threshold; such rarity of low scores in these zones is consistent with the definition of this threshold (i.e., 10th percentile of the reference distribution defined by 2001 to 2006 data). Below the Lehigh River in Zone 1E, however, 41% of the invertebrate samples fell below the impairment threshold (7 of 17 samples). For the interim methodology, DRBC has defined “impairment” as greater than 30% of sampling falling below the threshold (see Table 17). Therefore, Zone 1E is indicated as not meeting the biocriteria for the assessment period.

Table 17: Summary of Biological Assessment Results

Zone	Years of Data	Stations per Assessment Unit	% of samples in time window w/ 6-metric IBI < 75.6
1A	2 (2007, 2008)	4	0%
1B		3	0%
1C		4	12.5%
1D		5	0%
1E		7	41%

Public Water Supply

The public water supply designated use is assessed through evaluation of TDS, hardness, chlorides, odor, phenols, sodium, turbidity, systemic toxicants, carcinogens, and administrative drinking water closures. Table 18 below shows the Public Water Supply assessment results for the 2012 assessment. Additional detail on each evaluation is provided in the subsequent sections.

Table 18: Public Water Supply Designated Use Assessment Results

Zone (AU)	TDS	Hardness	Chlorides	Odor	Phenols	Sodium	Turbidity	Systemic Toxicants	Carcinogens	Drinking water closures	2012 Assessment	2010 Assessment
1A	+	N/A	N/A	ID	ID	N/A	- ^A	+	+	+	NS ^B	S
1B	+	N/A	N/A	ID	+	N/A	- ^A	+	+	+	NS ^B	S
1C	+	N/A	N/A	ID	ID	N/A	+	+	+	+	+	NS
1D	+	N/A	N/A	ID	ID	N/A	- ^A	+	+	+	NS ^B	S
1E	+	N/A	N/A	ID	ID	N/A	- ^A	+	+	+	NS ^B	NS
2	+	+	+	ID	ID	N/A	- ^A	+	+	+	NS ^B	S
3	+	+	+	ID	ID	+	+	+	+	+	+	S

Notes:

A – Rate of criteria exceedance is below the historical threshold of 10%.

B – Based primarily on fewer than 10% exceedances of criteria

ID – Insufficient Data

TDS

A description of TDS and assessment against the TDS criteria are presented under the Aquatic Life designated use in the previous section.

Hardness

Hardness is an integrated measure of divalent metallic cations. Measuring hardness in source water provides an indication of whether water softening will be desirable either in drinking water processing or in the finished drinking water at the point of use. Table 19 below shows that hardness criteria were met in all samples in Zones 2 and 3, where hardness criteria have been adopted.

Table 19: Hardness Assessment Results

Zone	% Observations Meeting Criteria	Primary Data Source(s)	Notes
1A	No Criteria		
1B			
1C			
1D			
1E			
2	100% (presumed)	• DRBC Boat Run	No individual observation exceeded criteria, therefore, attainment of 30-day mean criteria is presumed
3	100% (presumed)	• DRBC Boat Run	

4	
5	
6	Use not applicable in this Zone

Chlorides

Chloride is one of the major inorganic ions in water and wastewater, and can impart a salty taste to drinking water at elevated concentrations. Chloride criteria are expressed as a 15-day mean in Zone 2 and a 30-day mean in Zone 3. Although sequential daily measurements are not made as part of routine surface water monitoring programs, no individual observation exceeded the numerical criteria (as shown in Table 20 below), therefore 100% attainment of the criteria is presumed.

Table 20: Chlorides Assessment Results

Zone	% Observations Meeting Criteria	Primary Data Source(s)	Notes
1A	No Criteria		
1B			
1C			
1D			
1E			
2	100% (presumed)	<ul style="list-style-type: none"> • DRBC Boat Run • PWD Boat based monitoring 	No individual observation exceeded criteria, therefore, attainment of 15-day mean criteria is presumed
3	100% (presumed)	<ul style="list-style-type: none"> • DRBC Boat Run • PWD Boat based monitoring 	No individual observation exceeded criteria, therefore, attainment of 30-day mean criteria is presumed
4	Use not applicable in this Zone		
5			
6			

Odor

No odor data was indicated in any of the data sets reviewed. Therefore, no assessment against odor criteria was made.

Phenols

Phenols were analyzed only in Zone 1B by New York State Department of Environmental Conservation. All results were Non-Detect. Therefore, the criterion is met in Zone 1B but indicated as "insufficient data" in all other zones with a drinking water designated use.

Sodium

A criterion for sodium exists only in Zone 3. A review of all available data shows that all values were below the 30-day mean criterion of 100 mg/L. Although the spacing of the data did not support computing a 30-day mean, since all values were below 100 mg/L, attainment of this criterion is presumed.

Turbidity

A detailed discussion of the turbidity assessment is provided in the Aquatic Life section of this report.

Systemic Toxicants

Systemic toxicants affect the entire body or many organs rather than a specific site. For example, cyanide is a systemic toxicant that can affect every cell and organ in the body by interrupting oxygen exchange by cells. Stream quality objectives for systemic toxicants are established if a reference dose (RfD) exists in EPA's Integrated Risk Information System (IRIS). Drinking water is supported in Zones 1, 2, and 3 (Table 18, with additional detail in Appendix D).

Carcinogens

Carcinogens are substances that act directly in causing cancer. This may be due to the ability of the substance such as dioxins/furans to damage the genome or to disrupt cellular metabolic processes. Stream quality objectives for carcinogenic toxicants are established if a cancer potency factor (CPF) is available and the substance is classified as a carcinogen in EPA's Integrated Risk Information System (IRIS). Drinking water is supported in Zones 1, 2, and 3 (Table 18, with additional detail in Appendix D)

Emerging Contaminants

Emerging contaminants are unregulated substances that have entered the environment through human activities. Current regulatory approaches are inadequate to address these contaminants and the increasing public concern over their environmental and human health implications. Emerging contaminants have historically not been routinely monitored therefore limited information is available on past trends. A pilot survey of emerging contaminants in the main stem of the tidal Delaware River ambient waters in 2007, 2008 and 2009 detected pharmaceuticals, personal care products, perfluorinated compounds, hormones, sterols, nonyl phenols and polybrominated diphenyl ethers at levels comparable to similar compounds and concentrations measured in occurrence studies of ambient water in other urban areas (DRBC Draft Report February 2012. Contaminants of Emerging Concern in the Tidal Delaware River: Pilot Monitoring Survey 2007 – 2009). Assessment priorities in the tidal River include further characterization of persistent and bioaccumulative perfluorinated compounds and a more comprehensive evaluation of potential ecological effects from pharmaceuticals in the estuary. Benchmark values for environmental safety are needed and in some cases water quality criteria may need to be derived for some emerging contaminants to facilitate future water quality assessment.

Drinking Water Closures

For the Assessment Period (Oct 2006 – Sept 2011), there were no administrative closures to drinking water intakes as a result of water quality issues or violations.

Contact Recreation

The DRBC water quality regulations sub-divide Zone 4 for bacteria criteria. The upper portion of Zone 4, above River Mile 81.8, is designated as secondary contact recreation only, while the lower portion of Zone 4, below River Mile 81.8, is designated for both primary and secondary contact recreation.

Primary contact recreation is supported in all applicable Zones, except Zone 4 below RM 81.8, where there is insufficient data. Secondary contact recreation is supported in Zones 3 and 4. As shown in Table 21, primary and secondary contact uses were supported in all applicable Zones, except for the lower portion of Zone 4, where insufficient data were available.

Table 21: Primary and Secondary Contact Recreation Assessment Results

AU	Fecal Coliform		Enterococcus		2012 Assessment	2010 Assessment
	Primary	Secondary	Primary	Secondary		
1A	+	+	N/A	N/A	S	S
1B	+	+	N/A	N/A	S	S
1C	+	+	N/A	N/A	S	S
1D	+	+	N/A	N/A	S	S
1E	+	+	N/A	N/A	S	S
2	+	+	+	+	S	S
3	N/A	+	N/A	+	S	S
4 (> RM 81.8)	N/A	+	N/A	+	S	S
4 (< RM 81.8)	ID	ID	ID	ID	ID	ID
5	+	+	+	+	S	S
6	+	+	+	+	S	S

Fish Consumption

The fish consumption designated use applies to all DRBC WQM Zones. The assessment criterion is based primarily on the presence of the Basin states' fish consumption advisories in the mainstem Delaware River and Bay for the 2010 to 2011 assessment period. The presence of fish consumption advisories results in an assessment of "not supporting the designated use".

The following fish advisories reports were used:

- Delaware Fish Consumption Advisories:
http://www.dnrec.delaware.gov/fw/Fisheries/Documents/Delaware_Fish_Advisory_Chart.pdf - January 30, 2012)
- New Jersey Fish Consumption Advisories: <http://www.state.nj.us/dep/dsr/njmainfish.htm>
- Pennsylvania Fish Consumption Advisories:
http://www.portal.state.pa.us/portal/server.pt/community/fish_consumption/10560/fish_advisory/554001

- New York Fish Consumption Advisories:
http://www.health.ny.gov/environmental/outdoors/fish/health_advisories/

Table 22 below provides a summary of the consumption advisories issued by the states. For each assessment unit, between 2 and 8 advisories were issued. There is no assessment unit without an advisory, so the use is not supported in any zone.

Table 22: Fish Consumption Advisory Summary

Fish Species	Contaminant	Fish Consumption Advisory – General Population									
		1A	1B	1C	1D	1E	2	3	4	5	6
DELAWARE											
All Finfish	PCBs, Dioxins, Mercury, Chlorinated Pesticides									NC (state line to C&D canal)	
Weakfish (all sizes), Bluefish (>14 in)	PCBs									1/month (C&D Canal to head of Bay)	1/month
White Perch, American Eel, Channel Catfish, White Catfish, Bluefish (>14 in)	PCBs, Mercury									1/year (C&D Canal to head of Bay)	1/year
Striped Bass	PCBs, Mercury									2/year (C&D Canal to head of Bay)	2/year
NEW JERSEY											
Smallmouth Bass	Not listed			1/week	1/week ^b	1/week					
White Sucker	Not listed			1/month		1/month					
Largemouth Bass	Not listed										
American Eel	Not listed				1/month	1/year	1/year	1/year			
Channel Catfish	Not listed				4/year	1/year	1/year	1/year			
White Catfish	Not listed			1/week ^a		1/month	1/month	1/month			
Carp	Not listed										
Sunfish	Not listed										
Striped Bass	Not listed				4/year	4/year	4/year	4/year			
White Perch	Not listed					4/year	4/year	4/year			
All Finfish	Not listed								NC (state line to C&D)		

										canal)	
Bluefish - larger than 24 inches or 6 lbs.	Not listed										NC
Bluefish - smaller than 24 inches or 6 lbs.	Not listed										1/year
Striped Bass, White Perch, American Eel, Channel Catfish, White Catfish	Not listed										1/year
Weakfish	Not listed										1/week
American Eel (Zone 6 tributaries)	Not listed										1/month

NEW YORK^c

No advisories for the mainstem Delaware River.

PENNSYLVANIA

Smallmouth Bass	Mercury	2/month	2/month	2/month	2/month ^b						
American Eel	Mercury	2/month	2/month	2/month	2/month	2/month					
	PCBs						NC	NC	NC		
Carp	PCBs						NC	NC	NC		
White Perch, Channel Catfish, Flathead Catfish, and Striped Bass	PCBs						1/month	1/month	1/month		
Number of Advisories		2	2	4	4	6	8	8	8	5	8

Shellfish Consumption

Shellfish consumption, as a DRBC designated use, only applies to DRBC WQM zone 6. The state of Delaware classifies its designated shellfish waters within Delaware Bay as falling into the following two categories:

- Approved
- Prohibited

New Jersey classifies shellfish waters as falling into the following categories:

- Unrestricted
- Special Restricted
- Seasonal (Nov to Apr)
- Prohibited

Figure 8 indicates the current DE and NJ classifications for shellfish in zone 6. Table 23 lists the current DE and NJ classifications and the 2012 Assessment results, with the 2010 Assessment results given for comparison (*note: 2012 and 2010 designations are identical*).

For the current 2012 assessment, approved harvesting areas were considered to be supporting (S) the use. Prohibited waters were considered to be not supporting (NS) the use. AUs classified as special restricted and seasonally restricted are considered to be supported, but with special conditions (SS). Note, however, that the states of DE and NJ do not list all prohibited or provisionally approved waters as impaired waters, as not all restrictions on shellfish harvesting are due to water quality issues (see the respective state Integrated Assessment reports for further information).

In total for the 2012 assessment, 637 mi² are in full support (90% of zone 6), 33 mi² are supporting with special conditions (5%), and 40 mi² are not supporting the shellfish consumption use (5%).

Figure 8: Shellfish Consumption Classifications designated by New Jersey and Delaware for the Delaware Bay (DRBC WQM zone 6)



Table 23: Shellfish Consumption Designated Use Assessment Result

State	Sub-Assessment Unit within Zone 6	Area (mi ²)	DE / NJ Shellfish Classification	2012 Assessment	2010 Assessment
Delaware	6de1	306	Approved	S	S
	6de2	6	Prohibited	NS	NS
	6de3	5	Prohibited	NS	NS
	6de4	5	Prohibited	NS	NS
	6de5	1	Prohibited	NS	NS
	6de6	4	Prohibited	NS	NS
	6de7	17	Prohibited	NS	NS
New Jersey	6nj1	331	Approved	S	S
	6nj2	1	Prohibited	NS	NS
	6nj3	1	Prohibited	NS	NS
	6nj4	3	Seasonal (Nov-Apr)	SS	SS
	6nj5	4	Seasonal (Nov-Apr)	SS	SS
	6nj6	3	Special Restricted	SS	SS
	6nj7	1	Seasonal (Nov-Apr)	SS	SS
	6nj8	1	Seasonal (Nov-Apr)	SS	SS
	6nj9	1	Special Restricted	SS	SS
	6nj10	3	Seasonal (Nov-Apr)	SS	SS
	6nj11	0.2	Seasonal (Nov-Apr)	SS	SS
	6nj12	2	Special Restricted	SS	SS
	6nj13	0.2	Seasonal (Nov-Apr)	SS	SS
	6nj14	15	Special Restricted	SS	SS

S = "Supports": The assessment unit supports the designated use

SS = "Supports – Special": The assessment unit supports the designated use, but with special conditions

NS = "Not Supporting": The assessment unit does not support the designated use

ID = "Insufficient Data": Insufficient or unreliable data is present

Assessment Summary

Table 24 below shows the summary of assessments for Aquatic Life, Public Water Supply, Recreation, and Fish Consumption. Meaningful assessment continues to be hampered by the requirement to indicate one exceedance plus one confirmatory exceedance as not meeting criteria and subsequently not supporting a designated use. Turbidity exceedances not associated with high flow events during this assessment cycle resulted in Not Supporting indications for public water supply for several zones which were indicated as Supporting in the previous assessment.

Assessment of Shellfish applies only to Zone 6 and utilizes shellfish-specific assessment units. The Shellfish assessment summary is provided in Table 23 in the previous section.

Table 24: Summary of the 2012 Assessment

Zone (AU)	Aquatic Life		Drinking Water		Recreation		Fish Consumption	
	2012	2010	2012	2010	2012	2010	2012	2010
1A	NS ^A	NS	NS ^A	S	S	S	NS	NS
1B	NS ^A	NS ^A	NS ^A	S	S	S	NS	NS
1C	NS ^A	ID	S	S	S	S	NS	NS
1D	NS ^A	NS ^A	NS ^A	S	S	S	NS	NS
1E	NS ^A	NS	NS ^A	S	S	S	NS	NS
2	NS ^A	NS	NS ^A	S	S	S	NS	NS
3	NS ^A	NS ^A	S	S	S	S	NS	NS
4	NS ^A	NS ^A	N/A	N/A	ID/S	ID/S	NS	NS
5	NS ^A	NS ^A	N/A	N/A	S	S	NS	NS
6	NS ^A	NS ^A	N/A	N/A	S	S	NS	NS

Notes:

A – Based primarily on fewer than 10% exceedances of criteria

Recommendations for Future Action

Completion of the assessment for this cycle was hampered by several difficulties. We recommend addressing the following issues:

- Current guidance from EPA indicates that the proportion of allowable exceedances must be adopted as part of criteria in order to be considered during assessment. Where the criteria do not already include this information, EPA has indicated that assessors must indicate an assessment unit as having not met criteria when one exceedance plus one confirmatory exceedance are found. However, this approach is logically flawed and not supportable because it fails to recognize inherent measurement and analytical uncertainty, and imposes a wholly different standard than that which was in place at the time the criteria were developed. Prior to the next assessment cycle, DRBC should develop and adopt in Article 4, criteria implementation tables to assist in future assessment cycles
- DRBC should finalize adoption of new pH criteria before the next assessment cycle.
- DRBC should finalize adoption of surface water temperature criteria in Zones 1A through 1E.
- DRBC should continue work on better defining the linkage between atmospheric and meteorological drivers, in order to estimate the proportion of temperature exceedances attributable to potentially controllable anthropogenic activities.
- DRBC should continue to work with other sponsors to ensure that continuous monitoring resources are established in Zones 1A through 1D.
- The DRBC has scheduled additional data collection using enhanced analytical methods, modified collection procedures and changes in the spatial scale of sampling in segments of the Delaware River which have exhibited apparent copper exceedances. The information collected will provide additional data to determine metal concentrations in ambient water and the impact of metals on water quality. The survey targets copper, zinc and nickel. Coordination among basin states and agencies should continue to ensure the use of the most appropriate methods and procedures for the conduct of monitoring studies in the Basin, and the harmonization of water quality criteria and assessment methodologies.

Appendix A: Descriptions of DRBC Monitoring Programs

The surface water quality monitoring program utilized by the DRBC consists of the following programs:

- The upper and middle non-tidal portions of the River (RM 330.7 to 209.5) are monitored through the *Scenic Rivers Monitoring Program*, a joint NPS and DRBC effort;
- The lower non-tidal portions (RM 209.5 to 133.4) are monitored through the *Lower Delaware Monitoring Program*;
- The Estuary, or tidal portion of the Delaware River (RM 133.4 to the mouth of the Delaware Bay), is monitored through the *Delaware River Boat Run Monitoring Program*, a joint effort between the DNREC and DRBC; and
- DRBC Ambient Water Monitoring of the Delaware River for Chronic Toxicity, which is included as an additional study under the Boat Run Monitoring Program.
- The *Biological Monitoring Program* collects macroinvertebrate samples throughout the non-tidal River (RM 300.7 to 133.4) for assessment of Aquatic Life Use

In addition, data obtained from other agencies' monitoring efforts are used to supplement data obtained through the DRBC sampling efforts. The other data sources include:

- DNREC Dioxins and Furans in Fish from the Delaware River Study,
- Pennsylvania Department of Environmental Protection (PADEP) Water Quality Network (WQN),
- New Jersey Department of Environmental Protection (NJDEP) Ambient Surface Water Monitoring Network (from STORET),
- New York State Department of Environmental Conservation (NYSDEC) Ambient Water Quality Monitoring Program (from STORET),
- United States Geological Survey (USGS) National Water Information System (NWIS),
- DRBC/USGS Cooperative Monitoring Program (continuous monitors),
- National Oceanic Atmospheric Administration (NOAA) Physical Oceanographic Real-Time System (PORTS) data, and
- EPA National Coastal Assessment Programs.

The DRBC water quality monitoring programs and the DNREC dioxin and furan study are described below. For information on quality objectives and criteria and sample design, refer to the following DRBC Quality Assurance Project Plans (QAPPs):

- Scenic Rivers Monitoring Program QAPP, Revision 1 (2006)
- Ambient Water Monitoring of the Delaware River for Chronic Toxicity QAPP, June 13, 2006
- Lower Delaware Water Quality Monitoring Program QAPP (2004)
- Delaware River Boat Run Monitoring Program QAPP (2004)
- Delaware River Biomonitoring Program QAPP (2007)

Scenic Rivers Monitoring Program (SRMP)

In 1984, the SRMP began monitoring approximately a 121-mile reach of the Delaware River, from RM 330.7 to RM 209.5, which contains two portions of the National Wild and Scenic Rivers System and numerous high quality tributaries that drain portions of New York, New Jersey, and Pennsylvania. The DRBC and NPS collect water quality measurements for the following purposes:

1. To convert reach-wide EWQ targets to ICP and/or BCP targets;
2. To support water quality models for SPW implementation; and
3. To gather sufficient water quality information to implement DRBC SPW regulations using a site-specific statistical approach to define and assess possible changes to existing water quality.

There are 47 sampling locations; however, for the 2010 Assessment, only data from Interstate Control Points (ICP) along the main stem Delaware River are utilized. Tributary boundary sites are not used.

Lower Delaware Monitoring Program (LDMP)

In 1999, DRBC began monitoring to characterize the existing water quality of the Lower Non-tidal Delaware River, the reach extending from Trenton, NJ, (RM 134) to the Delaware Water Gap (RM 210). This monitoring network was established because little data existed to characterize water quality in this reach, portions of which have been included in the National Wild and Scenic Rivers System. In 2004, DRBC completed a five-year effort to define existing water quality and to develop a water quality management strategy that protects and improves the water quality of the Lower Delaware region. Based on LDMP monitoring results, the Lower Delaware was declared by DRBC in 2005 as "Significant Resource Waters."

Program objectives include:

- Establishing EWQ for future comparison;
- Assessing attainment of water quality standards;
- Setting geographic and water quality priorities to maintain or improve EWQ; and
- Long-term monitoring so that DRBC can consistently perform its 305b assessment, evaluate trends, prioritize agency management activities, and assess effectiveness of strategy implementation.

Sampling is conducted at 9 Delaware River ICP sites and 15 tributary sites. Only the results for the ICP sites are used in the assessment.

Estuary Boat Run Program (Boat Run)

The Boat Run consists of monitoring of the tidal portion of the Delaware River from the head of tide at Trenton, NJ, (RM 133.4) to the mouth of the Delaware Bay, delineated as a line from Cape May, NJ, to Lewes, DE. The goals of the program are to provide accurate, precise, and defensible estimates of the surface water quality of the Delaware Estuary and to allow assessment of water quality standards compliance.

Sampling occurs 8 to 12 times per year at up to 22 locations, depending on funding. The samples are analyzed for routine and bacterial parameters, nutrients, heavy metals, sodium and biotic ligand model parameters, chlorophyll-a, dissolved silica, productivity, and volatile organics.

Delaware River Chronic Toxicity Study

The Toxic Advisory Committee (TAC) for the DRBC recommended and the DRBC Commissioners asked the DRBC staff to study and characterize the nature and extent of ambient chronic toxicity in the Delaware Estuary (Zone 2 through 5). As part of that ongoing effort, ambient toxicity surveys were conducted in 2007 and 2008. The surveys used ambient water to measure potential chronic toxicity in the tidal Delaware River (RM 50 to RM 131). The objective was to assess if chronic lethal or sublethal toxicity, as measured in laboratory experiments, was present in river water samples. Ambient toxicity at sixteen fixed stations in the main-stem of the tidal Delaware River with salinities from 0 to 15 parts per thousand (ppt) was assessed using six species: *Pimephales promelas*, *Americanasys bahia*, and *Menidia beryllina* in 7-day tests; *Ceriodaphnia dubia* in a test conducted for a maximum of 8-days; *Pseudokirchneriella subcapitata* in a 96-hour test; and *Hyalella azteca* in a 10-day water-only test. Survival, growth, and when possible, reproduction were measured in the toxicity tests. Sampling in two different years indicated, based on the measured endpoints, that the ambient samples from the mainstem of the Delaware River were not chronically toxic to the tested species. The sampling was not designed to characterize any potential near-field toxicity issues immediately surrounding point source discharges or contaminated sites. The surveys did identify tributaries that warrant further assessment for potential impairment from chronic lethal or sublethal toxicity.

Biological Monitoring Program

DRBC's biological monitoring of the non-tidal Delaware River (RM 330.7 to 133.4) began in 2001 using benthic macroinvertebrates as the monitoring endpoint. For many years, DRBC has assessed the Aquatic Life Use of the non-tidal river using physical and chemical parameters. The biological monitoring program seeks to complement this physical/chemical monitoring with measurements of the diversity and health of the aquatic life community itself.

The initial years of data collection were designed to characterize the spatial and temporal variation in invertebrate communities at 25 fixed monitoring stations within riffle habitats (see Appendix A-1 for station locations; see Biomonitoring QAPP and Silldorff and Limbeck 2009 for details of the monitoring design). Using these initial data, DRBC has worked with the Biological Advisory Subcommittee to the WQAC in the analysis of the data and in the development of an interim assessment methodology based on these macroinvertebrate collections. Data from macroinvertebrate collections during 2007 and 2008

were then interpreted relative to the newly developed interim assessment methodology for the 2012 Integrated Assessment.

Appendix B: Other DRBC Water Pollution Control Programs

DRBC's water pollution control program is carried out through a series of interdependent steps and provides a rational approach to protecting and restoring water quality in the basin. The waters of the Basin are protected for designated uses with water quality criteria (WQC) that specify what levels of individual parameters are appropriate, based upon a review of the current scientific understanding about the needs of those uses. DRBC's monitoring programs provide a mechanism to evaluate how those WQC are being met, and assessment of those monitored data provide the link to how well the designated uses are being protected. The identified impairment of interstate waters in the Basin leads to the development of total maximum daily loads (TMDLs), issuing of permits and other mechanisms to reduce loading of pollutants in order to improve water quality to levels that meet the criteria. In addition, DRBC has other layers of protection (i.e., Special Protection Waters) that aim to maintain existing water quality where it is better than the water quality criteria. The following are examples of how the Commission takes a multi-faceted approach to water quality regulation.

Special Protection Waters

Currently, portions of the Delaware River are designated by DRBC as "Special Protection Waters" (SPW) and have associated with them a variety of specific pollution prevention and reduction requirements driven by a "no measurable change" policy toward water quality. Designated reaches of SPW fall into two categories:

(1) Outstanding Basin Waters

- The Upper Delaware Scenic and Recreational River from Hancock, NY, to Milrift, NY (Delaware River between RM 330.7 and 258.4)
- Portions of intrastate tributaries located within the established boundary of the Upper Delaware Scenic and Recreational River Corridor
- The Middle Delaware Scenic and Recreational River from Milrift, NY, to the Delaware Water Gap (Delaware River between RM 250.1 and 209.5)
- Portions of tributaries located within the established boundaries of the Delaware Water Gap National Recreation Area

(2) Significant Resource Waters

- The Delaware River from Milrift, NY, to Milford, PA (RM 258.4 to 250.1)
- The Delaware River from the Delaware Water Gap to Trenton, NJ (RM 209.5 to 133.4).

SPW regulations take a watershed approach to antidegradation of water quality. The regulations apply to the drainage area of the designated waters. Policies provide an up-front approach to reducing or eliminating new pollutant loadings, through requirements made in the docket (permit) review process, for the purpose of maintaining "Existing Water Quality" (EWQ) in designated waters. This is accomplished, in part, by looking at the cumulative impacts of point and non-point sources as they may affect the designated waters, either through direct discharge or through tributary loading. EWQ is

defined in the regulations by numerical tables (DRBC WQR 2008). Numerical criteria for SPW EWQ are defined as (a) an annual or seasonal mean of the available water quality data, (b) two-tailed upper and lower 95 percent confidence limits around the mean, and (c) the 10th and 90th percentiles of the dataset from which the mean was calculated.

Estuary CBOD Allocations

The Commission determined that the 1964 carbonaceous biochemical oxygen demand (CBOD₂₀) of the effluent load to Zones 2, 3, 4, and 5 exceeded the waste assimilative capacity of those Zones to meet the stream quality objectives based upon numerical modeling study conducted in the late 1960s. In accordance with the regulations, the assimilative capacity of each Delaware Estuary Zone minus a reserve was originally allocated in 1968 among the individual dischargers based upon the concept of uniform reduction of raw waste in a Zone (Zones 2, 3, 4, and 5). Since 1968, the wasteload allocations for individual dischargers have been updated and documented by the Commission.

Pollutant Minimization Plans

In 2005, DRBC established requirements for the development of Pollutant Minimization Plans (PMP) for selected toxic pollutants including polychlorinated biphenyls (PCBs). These plans are currently being required for selected point and non-point discharges of PCBs in the Delaware Estuary. The goal of this program is to work toward meeting water quality standards and to eliminate fish consumption advisories due to PCBs. Because of the limited ability of dischargers to reduce their PCB loadings quickly enough to fully comply in the short term with the numeric limits that are based on water quality standards, this non-numeric approach allows the Commission to require dischargers to take actions in reducing PCB loadings to the Estuary. Pollutant Minimization Plans require biennial PCB sampling and submission of an annual report summarizing PCB loading reduction efforts. The Commission in cooperation with the states of New Jersey, Delaware and Pennsylvania has established a workgroup to include supplementary monitoring requirements via NPDES permits in order to better evaluate these efforts. Additionally, dischargers who were not initially required to develop a PMP have been required as part of their NPDES renewal or at the direction of the Commission to develop a PMP, perform monitoring and submit annual reports. The DRBC also has the authority to require PMPs for contaminated sites to further reduce non-point sources of PCB loadings to the Estuary.

More detailed descriptions of PCB efforts are provided at the DRBC web site, including the PCB Model Calibration Report (http://www.state.nj.us/drbc/library/documents/PCBhomolog_model-rpt0511.pdf), and extensive information on the PMP program (<http://www.state.nj.us/drbc/programs/quality/pmp.html>).

Point Source Control Program

DRBC uses a variety of programs to regulate point source pollutant loadings that would impact the Delaware River. These consist of docket review, pollutant allocations (including Pollutant Minimization Plans, PMPs), SPW regulations, and basin-wide minimum treatment standards and interstate cooperative agreements.

Section 3.8 of the compact states that "No project having substantial effect on the water resources of the Basin shall hereafter be undertaken by any person, corporation or governmental authority unless it

shall have been first submitted to and approved by the Commission". Projects are reviewed for potential impacts to the waters of the basin and for consistency with the Comprehensive Plan (http://www.state.nj.us/drbc/cp_wo_2.pdf), which consists of the statements of policies and programs that the commission determines are necessary to govern the proper development and use of the Delaware River Basin (DRBC Rules of Practice and Procedure (<http://www.state.nj.us/drbc/regs/rules.pdf>), 2002)/.

In addition, it is the policy of the Commission that there be no measurable change in existing water quality except towards natural conditions in Special Protection Waters (SPW). The DRBC implements both point source and non-point source controls through its SPW regulations. All new or expanding wastewater treatment projects must demonstrate that the new or incremental increase in the facility's load will not cause a measurable change in existing water quality at the relevant water quality control point for several parameters.

Article 4 of DRBC's Water Quality Regulations (<http://www.state.nj.us/drbc/regs/WQregs.pdf>) identifies basin-wide minimum treatment standards for wastewater discharges. These include:

- Removal of total suspended solids;
- Minimum secondary treatment for biodegradable wastes;
- BOD treatment requirements;
- Disinfection requirements;
- Color standards; Dissolved substance standards;
- pH standards;
- Ammonia standards;
- Temperature standards

DRBC maintains cooperative agreements with all four Basin states, which provide that all NPDES permits for projects that lie within the Basin must comply with DRBC standards as well as state standards.

Nonpoint Source Control Program

DRBC regulates non-point pollution as part of the anti-degradation requirements of SPW. Under DRBC SPW regulations, the service areas of all new or expanding wastewater discharge or water withdrawal project sponsors located in the drainage areas of SPW must submit for approval a Non-point Source Pollution Control Plan with their application. The plan must control the new or increased non-point source loads generated within the portion of the project sponsor's service area that is also located within the drainage area of SPW. The plans must document the Best Management Practices to be applied to the project site and/or service area. Non-point source pollution from runoff of developed areas in SPW may not be subject to antidegradation constraints if they are associated with an existing, non-expanding facility, such as a wastewater treatment plant that is not expanding its service area.

Non-point sources of PCBs may also be regulated, on a project-specific basis, by PMPs that the DRBC has begun requiring assistance in reducing PCB loadings into the Delaware River.

Coordination with Other Agencies

The nature of DRBC's water quality management activities relies on interstate coordination and cooperation. For instance, the agency maintains agreements with all four Basin states regarding permit review. Additionally, all new or amended DRBC regulations are ruled on by the Commission, which has representation by the four states and federal government. The Scenic Rivers Monitoring Program (SRMP) and Estuary Boat Run also rely on cooperation between DRBC and other agencies. The SRMP is a partnership between DRBC and the National Park Service (NPS), while the Boat Run is a partnership between DRBC and the Delaware Department of Natural Resources and Environmental Conservation (DNREC).

Integrated Resource Plans

In 1998, DRBC amended its Southeastern Pennsylvania Ground Water Protected Area Regulations to include watershed-based ground water withdrawal limits for sub-basins that lie entirely or partially within the protected area. As required by the Regulations, those withdrawal limits may be revised by the Commission to be more protective of streams designated by the State of Pennsylvania as either "high quality" or "exceptional value," or "wild," "scenic," or "pastoral," or to correspond to more stringent requirements in "integrated resource plans" adopted and implemented by all municipalities in the sub-basin. Integrated Resource Plans (IRPs) must assess water resources and existing uses of water; estimate future water demands and resource requirements; evaluate supply-side and demand-side alternatives to meet water withdrawal needs; assess options for wastewater discharge to subsurface formations and streams; consider storm water and floodplain management; assess the capacity of the sub-basin to meet present and future demands for withdrawal and non-withdrawal uses such as instream flows; identify potential conflicts and problems; incorporate public participation; and outline plans and programs including land use ordinances to resolve conflicts and meet needs. The development of IRPs helps focus and coordinate planning tools to consider the multiple uses of water resources and the interrelationships of water quality and quantity to meet various needs.

Integrated Resource Management

In 2001, DRBC began a multi-stakeholder process to develop a “forward-looking” Water Resources Plan for the Delaware River Basin (Basin Plan). In September 2004, the Governors of the Basin States and representatives of six federal agencies, signed a resolution showing their support for the Basin Plan. The Basin Plan is a unified framework of desired outcomes, goals, objectives, and milestones for protecting, preserving, and enhancing water resources. The central theme of the Basin Plan is a watershed-based approach to the achievement of integrated resource management. The Basin Plan sets a direction for water resource policy and management through 2030 and calls for the active involvement of a broad range of governmental and non-governmental entities in addition to DRBC.

Among the concepts included in the Basin Plan are the integration of water resources considerations into land use planning and management, the development of analytical tools to evaluate water resources impacts of municipal land use plans, the implementation of TMDLs to meet water quality standards for the protection of designated uses, and the use of regulatory and non-regulatory approaches to maintaining and improving water quality where it is better than criteria.

Appendix C: Fish Tissue Assessment Supplement

Fish Tissue Screening Values

Certain chemicals tend to concentrate (“bioaccumulate”) in fish to levels thousands of times greater than the levels in the water itself. The resulting concentrations in fish and the attendant health risks to those individuals who consume the fish, such as recreational and subsistence anglers, are of concern to government agencies and the public. The DRBC has developed fish tissue screening values (FTSV) for carcinogens and systemic toxicants at a risk level of one in a million (10^6) for fish tissue concentrations for specific bioaccumulative toxic pollutants following USEPA’s “Guidance for Assessing Chemical Contaminant Data for Use in Fish Advisories – Volume 1, 2 and 3 (USEPA 2000b) for establishing fish tissue thresholds. (<http://water.epa.gov/scitech/swguidance/fishshellfish/techguidance/guidance.cfm>)

DRBC FTSV are based on estimated body weight and fish ingestion rates for human populations. Carcinogenic chemical screening values include a factor estimating the human cancer potency of the chemical. Non-carcinogenic (systemic toxicant) chemical screening values include a reference dose which is an estimate of daily exposure to the chemical that will not cause harm in humans. Screening values are defined as concentrations of target analytes in fish or shellfish tissue that are of potential public health concern and that are used as threshold values against which levels of contamination in similar tissue collected from the ambient environment can be compared. Exceedance of these SVs should be taken as an indication that more intensive site-specific monitoring and/or evaluation of human health risk should be conducted. Field data, greater than the screening levels, are worthy of further evaluation. Possible further evaluation would include additional data collection, detailed risk analysis, and potential risk management action. It is important to note that fish tissue screening values are not intended to replace formal risk analysis. Rather, they help the assessor to decide whether a detailed risk analysis is even warranted and how to prioritize several analyses if screening values are exceeded at more than one location.

Assessment of Fish Tissue Data Collected Between 2004 and 2007

DRBC FTSVs for carcinogens and systemic toxicants are listed in Tables C1 and C2, respectively. The bioconcentrations factors (BCF), cancer potency factors and DRBC human health criteria (fish ingestion only) used to derive the FTSV are also listed in the tables. Comparable screening values from the EPA, DNREC and NJDEP are included in the tables. Fish tissue data collected from the Delaware River were compared to the FTSV. Concentrations in fish tissue higher than the FTSV are noted in Tables C1 and C2. Fish tissue samples from the Delaware River have the carcinogens arsenic, aldrin, chlordane, DDT, dieldrin and PCBs at concentrations higher than the FTSV for carcinogens. While concentrations of other carcinogens heptachlor, heptachlor epoxide, alpha- and beta-BHC, and toxaphene were below the FTSV. A brief summary of the carcinogenic parameters with concentrations higher than the FTSV are described below. None of the systemic toxicants measured (cadmium, mercury, nickel, selenium, zinc, aldrin, gamma-BHC, chlordane, DDT, dieldrin, endosulfan, endosulfan sulphate, endrin, endrin aldehyde, heptachlor, heptachlor epoxide and PBDE) had concentrations higher than the systemic FTSV.

Mercury

Although concentrations of mercury as wet weight in fish fillet from the Delaware River do not exceed a residue based water quality criteria of 300 ppb methylmercury assuming methyl mercury is approximately 80% of total mercury measured in the fish tissue based on USEPA, 2000. Guidance for Assessing Chemical Contaminant Data for Use in Fish Advisories – Volume 1, 2 and 3 (USEPA 2000b) for establishing fish tissue thresholds.

(<http://water.epa.gov/scitech/swguidance/fishshellfish/techguidance/guidance.cfm>) (Figure C1), mercury is worth noting because the assessment is based on a residue based criteria not a FTSV. If calculated based on dry weight, mercury concentrations would exceed the criteria. Furthermore, the DRBC methylmercury criterion is fish tissue residue based as recommended by USEPA. Although no exceedances were observed in the fish species monitored by the DRBC in tidal and non-tidal waters (channel catfish, white perch, smallmouth bass and white sucker). PADEP database includes exceedances in smallmouth bass and American eel in Zone 1 and West Branch and DNREC database includes Atlantic croaker, tautog, and striped bass with exceedances for striped bass in Zones 5 and 6. DRBC staff is soliciting data on methyl mercury in fish tissue sampled from the Delaware River. In addition, comments are requested on implementation procedures for the tissue-based methylmercury criterion.

Arsenic

Concentrations of arsenic as wet weight in white perch and channel catfish from the tidal Delaware River (Zone 1) exceed a FTSV of 2.67 ppb inorganic arsenic assuming an adjustment factor of 10% to estimate inorganic arsenic from measured total arsenic. Concentrations of arsenic in smallmouth bass and white sucker from the non-tidal Delaware River were below the FTSV (Figure C2).

Aldrin

Concentrations of aldrin as wet weight in white perch and channel catfish from the tidal Delaware River (Zones 2 through 5) exceed a FTSV of 0.24 ppb. Concentrations of aldrin in smallmouth bass and white sucker from the non-tidal Delaware River were below the FTSV (Figure C3).

Chlordane

Concentrations of chlordane (sum of all chlordanes) as wet weight in channel catfish from the tidal Delaware River (Zones 2 through 5) exceed a FTSV of 11.43 ppb. Concentrations of chlordane in white perch from the tidal river as well as smallmouth bass and white sucker from the non-tidal river were below the FTSV (Figure C4).

DDT

Concentrations of DDT and metabolites as wet weight in channel catfish, white perch, white sucker and smallmouth bass from the tidal and non-tidal Delaware River (Zones 1 through 5) exceed a FTSV of 11.76 ppb. Concentrations are highest in the tidal species (Figure C5).

Dieldrin

Concentrations of dieldrin as wet weight in channel catfish, white perch, white sucker and smallmouth bass from the tidal and non-tidal Delaware River (Zones 1 through 5) exceed a FTSV of 0.25 ppb. Concentrations are higher in the tidal species (Figure C6).

PCB

Concentrations of PCB as wet weight in channel catfish, white perch, white sucker and smallmouth bass from the tidal and non-tidal Delaware River (Zones 1 through 5) exceed a cancer FTSV of 1,500, pg/g (1.5 ppb). The calculated DRBC FTSV for total PCB at 0.52 ppb is lower than EPA, DNREC or NJDEP screening levels. The DRBC FTSV for PCB is calculated from values used to derive the current DRBC water quality criteria. Revised PCB criteria for the Delaware Estuary have been proposed. When updated water quality criteria are adopted, the FTSV for PCB will be reviewed. For this assessment, an EPA screening value at 1.5 ppb was used to assess the fish tissue data. Median PCB concentrations are 10-100x screening values. (Figure C7).

DxFs

Concentrations of dioxin and furans as wet weight in channel catfish, white perch, white sucker and smallmouth bass from the tidal and non-tidal Zones 1 through 5 had concentrations higher than the systemic FTSV exceed a cancer screening value of 0.019 pg/g (0.000019 ppb) (Figure C8). EPA recommends basing the fish consumption screening value for DxFs on Toxic Equivalents (TEQs) related to 2,3,7,8-TCDD toxicity. To calculate the TEQ of a dioxin mixture, the concentration of each toxic compound is multiplied with its Toxic Equivalency Factor (TEF) and then added together. Median DxF TEQs are approximately 100x screening values.

PBDE

Environmental monitoring programs conducted worldwide during the past decade have shown increasing levels of some BDE congeners in contrast to a general decline in the occurrence of dioxins, PCBs and chlorinated pesticides. PBDEs, an emerging and unregulated compounds, have been observed in whole or fillet fish tissue at concentration from non-detect to 1,300 ppb total PBDE ww in U.S. waterways (Wenning et al, 2011). PBDE congeners with oral reference dose listed in EPA-IRIS (BDE-47, BDE-99, BDE-153 and BDE-209) were not measured in fish tissue from the Delaware River at concentrations higher than the DRBC calculated systemic FTSV (Table C2). FTSVs for carcinogenic effects are not available for PBDE. Although BDE-209 has suggestive evidence of carcinogenic potential, an oral

slope factor is not listed in IRS. There is insufficient data currently available to determine if BDE-47, BDE-99, and BDE-153 are potential carcinogens.

References

- Cavallo, GJ and TJ Fikslin. 2010. A TEQ Analysis of PCBs and Dioxin/Furans (DxF's) in Fish Tissue from the Delaware River. Gregory J. Cavallo and Thomas J. Fikslin. 2010 AWRA Water Resources Conference Presentation, Philadelphia, PA, November 4, 2010
- DNREC and DHSS. 2005. Technical Procedures for Evaluating Human Health Risks Associated with the Consumption of Chemically Contaminated Fish and Shellfish, March, 2005 (Revision 1),
- Faulds, A., N. Connelly, B. Knuth, J. Benowitz, J. Matassino, K.P. Norton. 2004. Patterns of Sport-fish Consumption at Six Pennsylvania Sites Along the Tidal Portion of the Delaware River with Special Emphasis on Shore Anglers. Report prepared for the Pennsylvania Coastal Zone Management Program. Project Number: CZ1:02PD.09. March 31, 2004.
- Hardy, M. 2004. A comparison of the fish bioconcentration factors for brominated flame retardants with their nonbrominated analogues. Environmental Toxicology and Chemistry 23:656-661.
- KCA Research Division. 1994. Fish Consumption Patterns of Delaware Recreational Fisherman and their Households. Prepared for the State of Delaware Department of Natural Resources and Environmental Control. April 13, 1994.
- NJDEP. 2010 Integrated Water Quality Monitoring and Assessment Methods
- USEPA's Integrated Risk Information System (IRIS). <http://www.epa.gov/IRIS/>
- USEPA, 2000. Guidance for Assessing Chemical Contaminant Data for Use in Fish Advisories – Volume 1, 2 and 3 (USEPA 2000b) for establishing fish tissue thresholds. (<http://water.epa.gov/scitech/swguidance/fishshellfish/techguidance/guidance.cfm>)
- USEPA. 2001. Update of ambient Water Quality Criteria for Cadmium. EPA-822-R-01-001
- USEPA. 2002. National Recommended Water Quality Criteria 2002. Human Health Criteria Calculation Matrix. EPA-822-R-02-012
- USEPA. 2003. Technical Summary of Information Available on the Bioaccumulation of Arsenic in Aquatic Organisms. EPA-822-R-03-032. <http://water.epa.gov/scitech/swguidance/standards/criteria/aqlife/pollutants/arsenic/fs.cfm>
- Wenning, R.J., L. Martello and A. Prusak-Daniel. 2011. "Dioxins, PCBs and PBDE in Aquatic Organism". in Environmental Contaminants in Biota: Interpreting Tissue Concentrations, 2nd Ed, Editors: WN Beyer and JP Meador.

Table C1										29-Mar-12	
DRBC Fish Tissue Screening Values - Carcinogens			DRAFT								
PARAMETER	BCF ^a	CANCER POTENCY	DRBC	DRBC	risk level 10 ⁻⁶	risk level 10 ⁻⁵	NJDEP ^f	risk level 10 ⁻⁶	concentrations	in	comments
		Fish Ingestion	Fish Ingestion Only	risk level 10 ⁻⁵	EPA-SV	DNREC ^e	Fish Tissue	DRBC	fish tissue		
		Only ^c	Regulatory	derived	ave adult	Based Toxics	Fish Tissue Screening Value	Fish Tissue Screening Value	ww		
	FACTOR		Only ^c	EPA-SV ^d	FTSV	Based Toxics	FTSV ^g	FTSV			
	oral slope factor ^b	mg/kg/day	ug/L	ug/L	ppb	ppb	ppb	ppb			
Arsenic	44.00	1.50E+00	0.061	0.061	26	2.6	36		2.67	yes ^h	1
Aldrin	4670.0	1.70E+01	0.000050	0.000050					0.24	yes	2
Chlordane	14100.0	3.50E-01	0.000811	0.00081	114	11.4	42	11	11.43	yes ⁱ	1
DDT	53600.0	3.40E-01	0.000219	0.00022	117	11.7	159	86	11.76	yes	1
DDE	53600.0	3.40E-01	0.000219	0.00022					11.76		
DDD	53600.0	2.40E-01	0.000311	0.00031					16.67		
Dieldrin	4670.0	1.60E+01	0.000054	0.000054	2.5	0.25	3		0.25	yes	1
Heptachlor	11200.0	4.50E+00	0.000079	0.000079					0.89	no	2
Heptachlor epoxide	11200.0	9.10E+00	0.000039	0.000039	4.39	0.44	6		0.44	no	1
alpha - BHC (HCH)	130.0	6.30E+00	0.004884	0.004900					0.63	no	2
beta - BHC (HCH)	130.0	1.80E+00	0.017094	0.017000					2.22	no	2
PCBs (Total) BCF	31200.0	7.70E+00	0.0000448	0.000045	1.5		27	8	0.52	yes	3
Toxaphene	13100.0	1.10E+00	0.000278	0.00028	36.3	3.63			3.64	no	1
Dioxin/furans					0.000000256	2.56E-08			0.00019	yes	4
2,3,7,8-TCDD	5000	156000	0.000000005	0.000000005					0.0003	yes	2
2,3,7,8-TCDD TEQs							0.0004		0.0004		5

- a) USEPA, 2002. National Recommended Water Quality Criteria 2002. Human Health Criteria Calculation Matrix
 b) EPA - IRIS
 c) Calculations use consumption rate of 17.5 grams per day and body weight of 70 kg
 d) USEPA 2000 (EPA 823-B-00-007 and EPA 823-B-00-007)
 e) DNREC and DHSS, Technical Procedures for Evaluating Human Health Risks Associated with the Consumption of Chemically Contaminated Fish and Shellfish 2005
 f) NJDEP, 2010 Integrated Water Quality Monitoring and Assessment Methods
 g) DRBC fish tissue screening value = (RL/CSF)/BW/CR : RL-risk level, CSF-oral cancer potency factor(mg/kg-d), BW-body weight (kg), CR-mean daily consumption rate (kg/g)
 h) one tenth of measured total arsenic is estimated to be organic arsenic on which the FTSV is based EPA-822-R-03-032
 i) sum of all chlordane
 Comments:
 1) DRBC FTSV, EPA SV and DRBC WQ criteria are consistent.
 2) EPA SV is not available and the derived DRBC FTSV is used.
 3) EPA SV is used.
 4) EPA SV is not available and a basin state SV is used.
 5) DRBC FTSV is TEQ based. Basin state SV used.
 Concentrations are based on wet weight.

Table C2										29-Mar-12	
DRBC Fish Tissue Screening Values - Systemic Toxicants											
PARAMETER	BCF ^a	REFERENCE	RSC	Fish Ingestion	Fish Ingestion	EPA-SV ^d	EPA-SV	DNREC ^e	NJDEP ^f	DRBC fish tissue	concentrations
		DOSE ^b		Only ^c	Only	recreational			fish tissue	in	comment
		(MG/KG/DAY)		(UG/L)	DRBC Regulatory Value (ug/L)	fishers ppm	ppb	ave adult ppb	toxics ppb	screening value (FTSV) ^g ppb	fish tissue ww higher than FTSV
Cadmium	64.0	1.00E-03	0.25	15.63	16	4.0	4,000	2,161		4,000	no 1
Mercury (methylmercury)	-			0.3 mg/kg (ppm) fish tissue		0.4	400	300	180	300	no ^h 1
Nickel	47.0	2.00E-02		1702.13	1700					80,000	no 2
Selenium	4.8	5.00E-03		4167	4200	20.0	20,000	10,803		20,000	no 1
Zinc	47.0	3.00E-01		25532	26000					1,200,000	no 2
Aldrin	4,670	3.00E-05		0.026	0.026					120	no 2
gamma - BHC (Lindane)HCH	130	3.00E-04	0.2	1.8	1.8	1.2	1,200			1,200	no 1
Chlordane	14,100	5.00E-04		0.14	0.14	2.0	2,000			2,000	no ^j 1
DDT	53,600	5.00E-04		0.037	0.037	2.0	2,000	1,080	86,000	2,000	no 1
Dieldrin	4,670	5.00E-05		0.043	0.043	0.2	200	108		200	no 1
Endosulfan (alpha)	270	6.00E-03		88.89	89	24.0	24,000	12,963		24,000	no 1
Endosulfan (beta)	270	6.00E-03		88.89	89	24.0	24,000	12,963		24,000	no 1
Endosulfan sulphate	270	6.00E-03		88.89	89					24,000	no 2
Endrin	3,970	3.00E-04		0.060	0.060	1.2	1,200	648		1,200	no 1
Endrin aldehyde	3,970	3.00E-04		0.060	0.060					1,200	no 2
Heptachlor	11,200	5.00E-04		0.18	0.18					2,000	no 2
Heptachlor epoxide	11,200	1.30E-05		0.0046	0.0046	0.052	52	28		52	no 1
PBDE-47	26,050	1.00E-04		0.0154	0.0150					400	no
PBDE-99		1.00E-04								400	no
PBDE-153		2.00E-04								800	no
PBDE-209		7.00E-03								28,000	no

- a) Cadmium from USEPA, 2001 (EPA-822-R-01-001); PBDE-47 from Hardy, 2004; all other BCF from USEPA, 2002 (EPA-822-R-02-012)
 b) EPA - IRIS
 c) Calculations use consumption rate of 17.5 grams per day and body weight of 70 kg
 d) USEPA 2000 (EPA 823-B-00-007)
 e) DNREC and DHSS, Technical Procedures for Evaluating Human Health Risks Associated with the Consumption of Chemically Contaminated Fish and Shellfish 2005
 f) NJDEP, 2010 Integrated Water Quality Monitoring and Assessment Methods
 g) DRBC fish tissue screening value = (RFD/BW)/CR : RFD-oral reference dose (mg/kg-d), BW-body weight (kg), CR-mean daily consumption rate (kg/g)
 h) Total mercury measured while the tissue based criteria is for methyl mercury. Exceeds FTSV as dry weight.
 i) sum of all chlordane
 BCF - bioconcentration factor, SV - screening value; FTSV- fish tissue screening value; MCL = maximum contaminant level
 Comments:
 1) DRBC FTSV, EPA SV and DRBC WQ criteria are consistent.
 2) EPA SV is not available and the derived DRBC FTSV is used.




Fig C1. Mercury




Fig C2. Arsenic

Aldrin in Fish Fillet 2004 to 2007




Fig C3. Aldrin

Sum of Chlordanes in Fish Fillet 2004 to 2007




Fig C4. Chlordanes

DDT and metabolites in Fish Fillet 2004 to 2007




Fig C5. DDT

Dieldrin in Fish Fillet 2004 to 2007




Fig C6. Dieldrin




Fig C7. Total PCB




Fig C8. Total dioxin/furans TEQ

Appendix D: Toxic Pollutants Water Quality Assessment Details

Table D1: Human Health Objectives (Toxics MCLs) Assessment Results

Parameter	Maximum Contaminant Level (µg/l)	2012 Assessment/Zones monitored
Metals		
Arsenic	10	No exceedance /Zones 1C,1D, 1E and 2
Barium	2000	No exceedance /Zones 1C and 1E
Beryllium	4	No exceedance /Zones 1C,1D, 1E and 2
Chromium (trivalent)	100	NE (as total)/ Zones 1C,1D,1E,2,3
Copper	1300	No exceedance / Zones 1B,1C,1D,1E, 2,3
Lead	15	No exceedance / Zones 1B,1C,1D,1E, 2,3
Selenium	50	NE/Zones 1C,1D,1E
Pesticides/PCBs		
alpha-BHC	0.2	NM
beta-BHC	0.2	NM
gamma - BHC (Lindane)	2	NE/Zone 1E
2,4-Dichloro-phenoxyacetic acid (2,4-D)	70	NM
Methoxychlor	40	NE/Zone 1E
Toxaphene	3	NE/Zone 1E

Parameter	Maximum Contaminant Level (µg/l)	2012 Assessment/Zones monitored
Dioxin (2,3,7,8-TCDD)	0.00003	NM
2,4,5 Trichloro-phenoxypropionic acid (2,4,5-TP-Silvex)	50	NM
Volatile Organic Compounds (VOCs)		
Benzene	5	NE /Zones 1C,1D,1E, 2,3,4,5,6
Carbon Tetrachloride	5	NE/ Zones 2,3,4,5,6
1,2-Dichloroethane	5	NE /Zones 1C,1D,1E, 2,3,4,5,6
1,1-Dichloroethylene	7	NE/Zones 1C,1D,1E, 2,3,4,5,6
[1,2 - trans – Dichloroethene] 1,2 - trans - Dichloroethylene	100	NE/Zones 1C,1D,1E
Dichloromethane (methylene chloride)	5	NE/Zones 1B, 2,3,4,5,6
Tetrachloroethylene (PCE)	5	NE/Zones 1B,1C,1D,1E, 2,3,4,5,6
Toluene	1000	NE/Zones 1B,1C,1D,1E, 2,3
Total Trihalomethanes	80	NE/Zones 1C,1D,1E
1,1,1-Trichloroethane	200	NE/Zones 1C,1D,1E, 2,3,4,5,6
1,1,2-Trichloroethane	5	NE/Zones 1C,1D,1E, 2,3,4,5,6
Trichloroethylene	5	NE/Zones 1B,1C,1D,1E, 2,3,4,5,6
Vinyl Chloride	2	NE/Zones 1B,1C,1D,1E, 2,3,4,5,6
Polycyclic Aromatic Hydrocarbons (PAHs)		
Benzo(a)Pyrene	0.2	NM

Parameter	Maximum Contaminant Level (µg/l)	2012 Assessment/Zones monitored
Other Compounds		
Asbestos	7 million fibers/L	NM
Bis(2-Ethylhexyl) Phthalate	6	NM
Fluoride	4,000	NE/Zones/1B,1C,1D,1E,2
Nitrate	10,000	NE Zones 1A,1B,1C,1D,1E,2,3,4,5,6
Pentachlorophenol	1	NM

Table D2: Aquatic Life Objectives Assessment Results

Parameter	Freshwater Objectives (µg/l)		2012	Marine Objectives (µg/l)		2012
	Acute	Chronic		Acute	Chronic	
	Metals					
Aluminum ^{a,b}	750	87	E chronic/Zones 1A, 1B, 1D NM /Zones 1C + 1E	NA	NA	NM
Arsenic (trivalent) ^c	340	150	NE / Zones 1C,1D, 1E and 2)	69	36	NM
Cadmium ^c	0.651*EXP(1.0166* LN(hardness)-3.924)	0.651*EXP(0.7409* LN(hardness)-4.719)	NE/Zone 1C, 1E, 2 E acute/Zone 1B	40	8.8	NM
Chromium (trivalent) ^c	0.277*EXP(0.819* LN(hardness)+3.7256)	0.277*EXP(0.819* LN(hardness)+0.6848)	NE (as total) / Zones 1C,1D,1E,2,3,4,5	NA	NA	NE as total Zones 5,6
Chromium (hexavalent) ^c	16	11	NE / Zones 1D,1E,2,3,4,5	1,100	50	NE Zones 5,6
Copper ^c	0.908*EXP(0.9422* LN(hardness)-1.7)	0.908*EXP(0.8545* LN(hardness)-1.702)	NE / Zones 1D,1E,2,3,4,5	4.8	3.1	E acute/ Zones 5,6
Lead ^c	38	5.4	NE / Zones 1B,1C,1D,1E, 2,3,4,5	210	8.1	One E / Zone 5,6
Mercury ^c	1.4	0.77	NE/Zones 1B, 1C, 1D, 1E, 2	1.8	0.94	NM

Parameter	Freshwater Objectives (µg/l)		2012	Marine Objectives (µg/l)		2012
	Acute	Chronic		Acute	Chronic	
Nickel ^c <i>RM 197.8 site USGC reported as <50 no method listed</i>	0.846*EXP(0.846* LN(hardness)+2.255)	0.846*EXP(0.846* LN(hardness)+0.0584)	NE all Zones	64	22	NE Zones 5 to 6
Selenium ^a	20	5.0	NE/Zones 1C,1D,1E	290	71	NM
Silver ^c	0.85*EXP(1.72* LN(hardness)-6.59)	NA	NE/Zones 1C,1D,1E	1.9	NA	NM
Zinc ^c	0.95*EXP(0.8473* LN(hardness)+0.884)	0.95*EXP(0.8473* LN(hardness)+0.884)	NE All Zones (estimated from total in 1B and 1C)	90	81	NE Zones 5 to 6
		Pesticides/PCBs				
Aldrin	3	NA	NE/Zone 1E	1.3	NA	NM
gamma - BHC (Lindane)	0.95	NA	NE/Zone 1E	0.16	NA	NM
Chlordane	2.4	0.0043	NE/Zone 1E	0.09	0.004	NM
Chlorpyrifos (Dursban)	0.083	0.041	NE/Zones 1C,1E	0.011	0.0056	NM
DDT and metabolites (DDE & DDD) ^d	1.1	0.001	Single E/Zone 1E (no impairment)	0.13	0.001	NM
Dieldrin	0.24	0.056	NE/Zones 1C,1E	0.71	0.0019	NM
Endosulfan ^e	0.22	0.056	NE/Zones 1C, 1E	0.034	0.0087	NM
Endrin	0.086	0.036	NE / Zone 1E	0.037	0.0023	NM

Parameter	Freshwater Objectives (µg/l)			Marine Objectives (µg/l)		2012
	Acute	Chronic		Acute	Chronic	
Heptachlor	0.52	0.0038	NE / Zone 1E	0.053	0.0036	NM
Heptachlor Epoxide	0.52	0.0038	NE / Zone 1E	0.053	0.0036	NM
Parathion	0.065	0.013	NM	NA	NA	NM
PCBs (Total)	1.0	0.014	Not assessed ongoing TMDL	5.0	0.03	TMDL
Toxaphene	0.73	0.0002	DL>C	0.21	0.0002	NM
		Other Compounds				
Cyanide (free)	22	5.2	NM	1	1	NM
Pentachlorophenol	$e^{(1.005 * \text{pH}-4.83)}$	$e^{(1.005 * \text{pH}-5.29)}$	NM	13	7.9	NM
		Indicator Parameters				
Whole Effluent Toxicity	0.3 Toxic Units _{acute}	1.0 Toxic Units _{chronic}	NE ^f	0.3 TU _a	1.0 TU _c	NE ^f

^aTotal recoverable criterion

^b Aluminum criteria listed should be restricted to waters with pH between 6.5 and 9.0.

^cDissolved Criterion

^dThis criterion applies to DDT and its metabolites (i.e., the total concentration of DDT and its metabolites should not exceed this value).

^eThis value was derived from data for endosulfan and is most appropriately applied to the sum of alpha-endosulfan and beta-endosulfan.

Criteria for cadmium, chromium(trivalent), copper, nickel, silver and zinc are hardness dependent and are expressed as the dissolved form (see Section 3.10.3.C.2. for form of metal).

Multiple exceedances of EPA criteria for cadmium in Zone 1B at a single site (RM 254).

^fSampling in 2007, 2008 and 2009 indicated, based on the measured endpoints, that the samples from sites tested in the main-stem of the Delaware River and from the majority of its tributaries were not chronically toxic to the tested species (*Pimephales promelas*, *Americamysis bahia*, *Menidia beryllina*, and *Ceriodaphnia dubia* in 7-d tests;

Pseudokirchneriella subcapitata in a 96-h test; and *Hyalella azteca* in a 10-d water-only test). The surveys identified tributaries that warrant further assessment for potential impairment from toxicity.

For 1 of the 3 test species, in 2 separate years of sampling, 2 sites (Assunpink Creek and Red Lion Creek) indicated chronic toxicity in both screening tests and confirmatory tests. Integrated Environmental Assessment and Management : Volume 7, Number 3, pp. 466–477.

NA = not available; NE = no exceedances; E = exceedances

NM = not monitored

DL>C = detection limit is greater than DRBC criteria

WB – West Branch of Delaware River (total mercury in PADEP database)

Table D3: Human Health Objectives (Carcinogens) Assessment Results

PARAMETER	FRESHWATER OBJECTIVES (µg/l)		2012 assessment	MARINE OBJECTIVES (µg/l)	2012 assessment
	FISH & WATER INGESTION	FISH INGESTION ONLY			
Metals					
Arsenic	*	NA	No exceedance/ Zones 1C,1D, 1E and 2)	NA	NM
Pesticides/PCBs					
Aldrin	0.000049	0.000050	NE/Zone 1E	0.000050	NM
alpha – BHC	0.0026	0.0049	NM	0.0049	NM
beta – BHC	0.0091	0.017	NM	0.017	NM
Chlordane	0.00080	0.00081	NE/Zone 1E	0.00081	NM
DDD	0.00031	0.00031	Single E/Zone 1E (no impairment)	0.00031	NM
DDE	0.00022	0.00022	Single E/Zone 1E (no impairment)	0.00022	NM
DDT	0.00022	0.00022	Single E/Zone 1E (no impairment)	0.00022	NM
Dieldrin	0.000052	0.000054	NE/Zones 1C,1E	0.000054	NM
Heptachlor	0.000079	0.000079	NE / Zone 1E	0.000079	NM

PARAMETER	FRESHWATER OBJECTIVES (µg/l)		2012 assessment	MARINE OBJECTIVES (µg/l)	2012 assessment
	FISH & WATER INGESTION	FISH INGESTION ONLY			
Heptachlor Epoxide	0.000039	0.000039	NE / Zone 1E	0.000039	NM
PCBs (Total)	0.0000444	0.0000448	Not assessed ongoing TMDL	0.0000079	Not assessed ongoing TMDL
Toxaphene	0.00028	0.00028	DL>C	0.00028	NM
Volatile Organic Compounds (VOCs)					
Acrylonitrile	0.051	0.25	DL>C/Zones 1C,1D,1E	0.25	NM
Benzene	0.61	14	NE /Zones 1C,1D,1E, 2,3,4,5	14	NE /Zones 5,6
Benzidine	0.000086	0.00020	NM	0.00020	NM
Bromoform(tribromomethane)	4.3	140	NE /Zones 1A,1B,1C,1D, 1E, 2,3,4,5	140	NE /Zones 5,6
Bromodichloromethane	0.55	17	NE/Zones 1C,1D,1E	17	NM
Carbon Tetrachloride	0.23	1.6	NE/Zones 2,3,4,5	1.6	NE /Zones 5,6
Chlorodibromomethane	0.40	13	NE/Zones 1B,2,3,4,5	13	NE/Zones 5,6
Chloroform	5.7	470	NE/Zones	470	NE/Zones

PARAMETER	FRESHWATER OBJECTIVES (µg/l)		2012 assessment	MARINE OBJECTIVES (µg/l)	2012 assessment
	FISH & WATER INGESTION	FISH INGESTION ONLY			
			1B,2,3,4,5		5,6
3,3 - Dichlorobenzidine	0.021	0.028	NM	0.028	NM
1,2 - Dichloroethane	0.38	37	NE /Zones 1C,1D,1E, 2,3,4,5	37	NE /Zones 5,6
1,2 - Dichloropropane	0.50	15	NM	15	NM
1,3 - Dichloropropene	0.34	21	NE/1C,1D,1E,2,3,4,5	21	NE/Zones 5,6
Dichloromethane (Methylene chloride)	*	590	NE/1B,2,3,4,5	590	NE/5,6
Tetrachloroethylene	0.69	3.3	NE/Zones 1B,1C,1D,1E, 2,3,4,5	3.3	NE/Zones 5,6
1,1,2,2 - Tetrachloroethane	0.17	4.0	NM	4.0	NM
1,1,2 - Trichloroethane	0.59	16	NE/Zones 1C,1D,1E, 2,3,4,5,	16	NE/Zones 5,6
Trichloroethylene	2.5	30	NE/Zones 1B,1C,1D,1E, 2,3,4,5	30	NE/Zones 5,6
Vinyl Chloride	0.025	2.4	NE/Zones 1B,1C,1D,1E, 2,3,4,5	2.4	NE/Zones 5,6
Polycyclic Aromatic Hydrocarbons (PAHs)					
Benz[a]anthracene	0.0038	0.18	NM	0.18	NM

PARAMETER	FRESHWATER OBJECTIVES (µg/l)		2012 assessment	MARINE OBJECTIVES (µg/l)	2012 assessment
	FISH & WATER INGESTION	FISH INGESTION ONLY			
Benzo[b]fluoranthene	0.038	0.18	NM	0.18	NM
Benzo[k]fluoranthene	0.38	1.8	NM	1.8	NM
Benzo[a]pyrene	0.0038	0.018	NM	0.018	NM
Chrysene	3.8	18	NM	18	NM
Dibenz[a,h]anthracene	0.0038	0.018	NM	0.018	NM
Indeno[1,2,3-cd]pyrene	0.038	0.18	NM	0.18	NM
Other Compounds					
Bis (2-chloroethyl) ether	0.03	0.53	NM	0.53	NM
Bis (2-ethylhexyl) phthalate	1.2	2.2	NM	2.2	NM
2,4 - Dinitrotoluene	0.11	3.4	NM	3.4	NM
1,2 - Diphenylhydrazine	0.036	0.2	NM	0.2	NM
Hexachlorobenzene	0.00028	0.00029	NM	0.00029	NM
Hexachlorobutadiene	0.44	18	NE/Zones 1C,1D,1E	18	NM
Hexachloroethane	1.4	3.3	NE/Zones 1C,1D,1E	3.3	NM
Isophorone	35	960	NM	960	NM

PARAMETER	FRESHWATER OBJECTIVES (µg/l)		2012 assessment	MARINE OBJECTIVES (µg/l)	2012 assessment
	FISH & WATER INGESTION	FISH INGESTION ONLY			
N-Nitrosodi-N-butylamine	0.0063	14	NM	14	NM
N-Nitrosodi-N-methylamine	0.00069	3.0	NM	3.0	NM
N-Nitrosodiethylamine	0.0008	1.24	NM	1.24	NM
N-Nitrosodi-N-phenylamine	3.3	6	NM	6	NM
N-Nitrosodi-N-propylamine	0.0050	0.51	NM	0.51	NM
N-Nitrosopyrrolidine	0.016	34	NM	34	NM
Pentachlorophenol	0.27	3.0	NM	3.0	NM
Dioxin (2,3,7,8 – TCDD)	0.000000005	0.0000000051	NM	0.0000000051	NM
2,4,6 - Trichlorophenol	1.4	2.4	NM	2.4	NM

* The MCL for this compound applies in Zones 2 and 3 and is listed in Table 3.

NA = not available

Table D4: Human Health Objectives (Systemic Toxicants) Assessment Results

PARAMETER	FRESHWATER OBJECTIVES ($\mu\text{g/l}$)		2012	MARINE OBJECTIVES ($\mu\text{g/l}$)	2012
	FISH & WATER INGESTION	FISH INGESTION ONLY		FISH INGESTION ONLY	
	Metals				
Antimony	5.6	640	NM	640	NM
Arsenic	*	NA	No exceedance (monitored Zones 1C,1D, 1E and 2)	NA	NM
Beryllium	*	420	No exceedance /Zones 1C,1D, 1E and 2)	420	NM
Cadmium	3.4	16	NE Zone 1B, 1C, 1D and 1E	16	NM
Chromium (trivalent)	*	380,000	NE as total Zones 1C,1D,1E,2,3,4,5	380,000	NE as total Zones 5,6
Chromium (hexavalent)	92	NA	NE Zones 1D,1E,2,3,4,5	NA	NE Zones 5,6
Chromium (total)	NA	750	NE Zones 1C,1D,1E,2,3,4,5	750	NE Zones 5,6
Mercury	0.050	0.051	NE / Zones 1B,1C,1D, 1E and 2	0.051	NM
Methylmercury	0.3 mg/kg fish tissue	0.3 mg/kg fish tissue	E/Zones WB, 1C, 1D; NE/Zones 1A, 1B, 1E, 2,3,4	0.3 mg/kg fish tissue	E/Zone 5 , 6
Nickel	500	1,700	NE all Zones	1,700	NE Zones 5,6

PARAMETER	FRESHWATER OBJECTIVES (µg/l)		2012	MARINE OBJECTIVES (µg/l)	2012
	FISH & WATER INGESTION	FISH INGESTION ONLY		FISH INGESTION ONLY	
Selenium	170	4,200	NE/Zones 1C,1D,1E	4,200	NM
Silver	170	40,000	NE/Zones 1C,1D,1E	40,000	NM
Thallium	0.24	0.47	NE/Zone 1D	0.47	NM
Zinc	7,400	26,000	NE All Zones	26,000	NE Zones 5,6
	Pesticides/PCBs				
Aldrin	0.025	0.025	NE/Zone 1E	0.025	NM
gamma - BHC (Lindane)	0.98	1.8	NE/Zone 1E	1.8	NM
Chlordane	0.14	0.14	NE/Zone 1E	0.14	NM
DDT and Metabolites (DDD and DDE)	0.037	0.037	Single E/Zone 1E (no impairment)	0.037	NM
Dieldrin	0.041	0.043	NE/Zones 1C,1E	0.043	NM
alpha -Endosulfan	62	89	NE/Zones 1C, 1E	89	NM
beta- Endosulfan	62	89	NE/Zones 1C, 1E	89	NM
Endosulfan Sulfate	62	89	NE /Zones 1C,1E	89	NM
Endrin	0.059	0.060	NE / Zone 1E	0.060	NM
Endrin Aldehyde	0.29	0.30	NE /Zones 1E (single sample)	0.30	NM

PARAMETER	FRESHWATER OBJECTIVES (µg/l)		2012	MARINE OBJECTIVES (µg/l)	2012
	FISH & WATER INGESTION	FISH INGESTION ONLY		FISH INGESTION ONLY	
Heptachlor	0.18	0.18	NE / Zone 1E	0.18	NM
Heptachlor Epoxide	0.0046	0.0046	NE / Zone 1E	0.0046	NM
Total PCBs	0.00839	0.00849	Not assessed ongoing TMDL	0.00149	Not assessed ongoing TMDL
	Volatile Organic Compounds (VOCs)				
Acrolein	6.1	9.3	NM	9.3	NM
Benzene	*	3,100	NE /Zones 1C,1D,1E, 2,3,4,5	3,100	NE /Zones 5,6
Bromoform (tribromomethane)	650	9,600	NE /Zones 1A,1B,1C,1D, 1E, 2,3,4,5	9,600	NE /Zones 5,6
Bromodichloromethane	680	NA	NE/Zones 1C,1D,1E	NA	NM
Dibromochloromethane	680	21,000	NE/Zones 1C, 1D, 1E	21,000	NM
Carbon Tetrachloride	*	150	NE/ Zones 2,3,4,5	150	NE/ Zones 5,6
Chloroform	68	2,100	NE/Zones 1B,2,3,4,5	2,100	NE/Zones 5,6
Chlorobenzene	130	1,600	NM	1,600	NM
1,1 - Dichloroethylene	*	7,100	NE/Zones	7,100	NE/Zones

PARAMETER	FRESHWATER OBJECTIVES (µg/l)		2012	MARINE OBJECTIVES (µg/l)	2012
	FISH & WATER INGESTION	FISH INGESTION ONLY		FISH INGESTION ONLY	
			1C,1D,1E, 2,3,4,5		5,6
1,2 - trans - Dichloroethylene	140	10,000	NE/Zones 1C,1D,1E	10,000	NM
1,3 - Dichloropropene	1,000	63,000	NE/1C,1D,1E,2,3,4,5	63,000	NE/Zones 5,6
Ethylbenzene	530	2,100	NE/2,3,45	2,100	NE/5,6
Methyl Bromide	47	1,500	NE/2,3,45	1,500	NE/5,6
Methylene Chloride	*	260,000	NE/ 1B,2,3,45	260,000	NE/5,6
1,1,2 – Trichloroethane	*	3,600	NE/Zones 1C,1D,1E, 2,3,4,5,	3,600	NE/Zones 5,6
Tetrachloroethylene	*	1,300	NE/Zones 1B,1C,1D,1E, 2,3,4,5	1,300	NE/Zones 5,6
Toluene	1,300	15,000	NE/Zones 1B,1C,1D,1E, 2,3,4,5	15,000	NE/Zones 5,6
	Polycyclic Aromatic Hydrocarbons (PAHs)				
Anthracene	8,300	40,000	NM	40,000	NM
Fluoranthene	130	140	NM	140	NM
Fluorene	1,100	5,300	NM	5,300	NM
Pyrene	830	4,000	NM	4,000	NM
	Other Compounds				

PARAMETER	FRESHWATER OBJECTIVES (µg/l)		2012	MARINE OBJECTIVES (µg/l)	2012
	FISH & WATER INGESTION	FISH INGESTION ONLY			
Acenaphthene	670	990	NM	990	NM
Benzidine	59	140	NM	140	NM
Bis (2-chloroisopropyl) ether	1,400	65,000	NM	65,000	NM
Bis (2-ethylhexyl) phthalate	*	620	NM	620	NM
Butylbenzyl phthalate	1,500	1,900	NM	1,900	NM
2 - Chloronaphthalene	1,000	1,600	NM	1,600	NM
2 - Chlorophenol	81	150	NM	150	NM
Cyanide	140	140	NM	140	NM
Dibutyl Phthalate	2,000	4,500	NM	4,500	NM
1,2 - Dichlorobenzene	420	1,300	NE/Zones 2,3,4,5	1,300	NE/Zones 5,6
1,3 - Dichlorobenzene	420	1,300	NE/Zones 2,3,4,5	1,300	NE/Zones 5,6
1,4 - Dichlorobenzene	63	190	NE/Zones 2,3,4,5	190	NE/Zones 5,6
2,4 - Dichlorophenol	77	290	NM	290	NM
Diethyl Phthalate	17,000	44,000	NM	44,000	NM
Dimethyl Phthalate	270,000	1,100,000	NM	1,100,000	NM

PARAMETER	FRESHWATER OBJECTIVES (µg/l)		2012	MARINE OBJECTIVES (µg/l)	2012
	FISH & WATER INGESTION	FISH INGESTION ONLY		FISH INGESTION ONLY	
2,4 - Dimethylphenol	380	850	NM	850	NM
2,4 - Dinitrophenol	69	5,300	NM	5,300	NM
2,4 - Dinitrotoluene	68	2,100	NM	2,100	NM
Hexachlorobenzene	0.35	0.36	NM	0.36	NM
Hexachlorocyclopentadiene	40	1,100	NM	1,100	NM
Hexachloroethane	20	46	NE/Zones 1C,1D,1E	46	NM
Isophorone	6,700	180,000	NM	180,000	NM
2-Methyl-4,6-dinitrophenol	13	280	NM	280	NM
Nitrobenzene	17	690	NM	690	NM
Pentachlorobenzene	1.4	1.5	NM	1.5	NM
Pentachlorophenol	*	11,000	NM	11,000	NM
Phenol	10,000	860,000	NE/Zone 1B	860,000	NM
1,2,4,5-Tetrachlorobenzene	0.97	1.1	NM	1.1	NM
1,2,4 - Trichlorobenzene	35	70	NE/Zones 1C,1D,1E	70	NM
2,4,5-Trichlorophenol	1,800	3,600	NM	3,600	NM
Vinyl Chloride	*	10,000	NE/Zones 1B,1C,1D,1E, 2,3,4,5	10,000	NE/Zones 5,6

* The MCL for this compound applies in Zones 2 and 3 and is listed in Table 3.

NA = not available

For this assessment cycle, where DRBC has not adopted numeric toxics criteria (Zones 1A through 1E), the DRBC narrative toxics standard is implemented by comparing measured toxics concentrations to USEPA's most recent National Recommended Water Quality Criteria for each parameter where an appropriate Criteria Maximum Concentration (CMC) and Criterion Continuous Concentration (CCC) are listed. EPA's most recent recommended criteria can be found at

<http://www.epa.gov/waterscience/criteria/wqctable/#cmc>

Some criteria require hardness values to compute the actual criteria numeric value. In these cases, multiple sources of hardness information may be used. Where multiple sources of hardness data are available, the assessment will consider the weight of evidence for multiple derivations of the criteria. Sources of hardness data could include:

- Site-specific paired hardness measured concurrently with toxic analytical parameter;
- Median site-specific hardness measured at other times;
- Hardness values listed in DRBC Water Quality Regulations.

For criteria expressed as the dissolved form of the metal, assessment of monitoring data is as follows:

- In assessment Zones with dissolved metals data collected, direct comparison to DRBC dissolved criteria;
- In assessment Zones with only total metals data collected (as noted in Table 5), comparison of total metals data to estimated total metals criteria using conversion factors listed in "Revised Procedure for Converting Total Recoverable Water Quality Criteria for Metals to Dissolved Criteria" <http://www.state.nj.us/drbc/regs/critmetals.pdf>.

The methylmercury criterion is fish tissue residue based as recommended by USEPA. No exceedances were observed in the fish species monitored by the DRBC in tidal and non-tidal waters (channel catfish, white perch, smallmouth bass and white sucker). PADEP database includes exceedances in smallmouth bass and American eel in Zone 1 and West Branch. DNREC database includes Atlantic croaker, tautog, and striped bass with exceedances for striped bass in Zones 5 and 6. DRBC staff is soliciting data on methyl mercury in fish tissue sampled from the Delaware River. In addition, comments are requested on implementation procedures for the tissue-based methylmercury criterion.

Appendix E: Taste and Odor Water Quality Assessment Details

Table E1: Taste and Odor as Human Health Objectives Assessment Results

PARAMETER	STREAM QUALITY OBJECTIVE ($\mu\text{g/l}$)	2012 assessment
Phenol	300	NE/Zone 1B
2 - Chlorophenol	0.1	NM
2,4 - Dichlorophenol	0.3	NM
2,4 - Dimethylphenol	400	NM
4 - Chloro - 3 - methylphenol	3.0 mg/l	NM
Pentachlorophenol	30	NM
Acenaphthene	20	NM
Chlorobenzene	20	NM
Hexachlorocyclopentadiene	1.0	NM
Nitrobenzene	30	NM

Appendix F: Public Participation Procedures

The table below highlights specific dates in the public participation and coordination process associated with this Assessment Report.

Table F1: Public Participation Milestones

<u>Date</u>	<u>Action</u>
July 6, 2011	Draft Assessment Methodology published on DRBC's web site.
August 12, 2011	Publication of a notice in the Federal Register regarding publication of the draft assessment methodology, including link to the Methodology on the DRBC web site.
August 31, 2011	Comments on draft Methodology due to DRBC.
September 21, 2011	Presentation on Draft Assessment Methodology to NJ Water Monitoring Council (NJWMC).
March 2, 2012	Draft Assessment report e-mailed to Water Quality Advisory Committee (WQAC), Monitoring Advisory Committee (MAC), Assessment leads in Pennsylvania, New Jersey, Delaware, and New York, and commenters on the Draft Assessment Methodology.
March 21, 2012	Presentation on Assessment Results to the Water Quality Advisory Committee.
April 1, 2012	Final Assessment Report e-mailed to USEPA, Region 3.