

SEP 20 1982

NUREG/ CR-2384  
PNL-3912

ASLBP LIBRARY

---

---

# Age-Specific Inhalation Radiation Dose Commitment Factors for Selected Radionuclides

---

---

Prepared by D. L. Strenge, R. A. Peloquin, D. A. Baker

**Pacific Northwest Laboratory**  
Operated by  
Battelle Memorial Institute

Prepared for  
**U.S. Nuclear Regulatory  
Commission**

## NOTICE

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, or any of their employees, makes any warranty, expressed or implied, or assumes any legal liability of responsibility for any third party's use, or the results of such use, of any information, apparatus, product or process disclosed in this report, or represents that its use by such third party would not infringe privately owned rights.

### Availability of Reference Materials Cited in NRC Publications

Most documents cited in NRC publications will be available from one of the following sources:

1. The NRC Public Document Room, 1717 H Street, N.W.  
Washington, DC 20555
2. The NRC/GPO Sales Program, U.S. Nuclear Regulatory Commission,  
Washington, DC 20555
3. The National Technical Information Service, Springfield, VA 22161

Although the listing that follows represents the majority of documents cited in NRC publications, it is not intended to be exhaustive.

Referenced documents available for inspection and copying for a fee from the NRC Public Document Room include NRC correspondence and internal NRC memoranda; NRC Office of Inspection and Enforcement bulletins, circulars, information notices, inspection and investigation notices; Licensee Event Reports; vendor reports and correspondence; Commission papers; and applicant and licensee documents and correspondence.

The following documents in the NUREG series are available for purchase from the NRC/GPO Sales Program: formal NRC staff and contractor reports, NRC-sponsored conference proceedings, and NRC booklets and brochures. Also available are Regulatory Guides, NRC regulations in the *Code of Federal Regulations*, and *Nuclear Regulatory Commission Issuances*.

Documents available from the National Technical Information Service include NUREG series reports and technical reports prepared by other federal agencies and reports prepared by the Atomic Energy Commission, forerunner agency to the Nuclear Regulatory Commission.

Documents available from public and special technical libraries include all open literature items, such as books, journal and periodical articles, and transactions. *Federal Register* notices, federal and state legislation, and congressional reports can usually be obtained from these libraries.

Documents such as theses, dissertations, foreign reports and translations, and non-NRC conference proceedings are available for purchase from the organization sponsoring the publication cited.

Single copies of NRC draft reports are available free upon written request to the Division of Technical Information and Document Control, U.S. Nuclear Regulatory Commission, Washington, DC 20555.

Copies of industry codes and standards used in a substantive manner in the NRC regulatory process are maintained at the NRC Library, 7920 Norfolk Avenue, Bethesda, Maryland, and are available there for reference use by the public. Codes and standards are usually copyrighted and may be purchased from the originating organization or, if they are American National Standards, from the American National Standards Institute, 1430 Broadway, New York, NY 10018.

---

---

# Age-Specific Inhalation Radiation Dose Commitment Factors for Selected Radionuclides

---

---

Manuscript Completed: July 1982  
Date Published: August 1982

Prepared by  
D. L. Streng, R. A. Peloquin, D. A. Baker

Pacific Northwest Laboratory  
Richland, WA 99352

Prepared for  
Division of Fuel Cycle and Material Safety  
Office of Nuclear Material Safety and Safeguards  
U.S. Nuclear Regulatory Commission  
Washington, D.C. 20555  
NRC FIN B2218



## ABSTRACT

Inhalation dose commitment factors are presented for selected radionuclides for exposure of individuals in four age groups: infant, child, teen and adult. Radionuclides considered are  $^{35}\text{S}$ ,  $^{36}\text{Cl}$ ,  $^{45}\text{Ca}$ ,  $^{67}\text{Ga}$ ,  $^{75}\text{Se}$ ,  $^{85}\text{Sr}$ ,  $^{109}\text{Cd}$ ,  $^{113}\text{Sn}$ ,  $^{125}\text{I}$ ,  $^{133}\text{Ba}$ ,  $^{170}\text{Tm}$ ,  $^{169}\text{Yb}$ ,  $^{182}\text{Ta}$ ,  $^{192}\text{Ir}$ ,  $^{198}\text{Au}$ ,  $^{201}\text{Tl}$ ,  $^{204}\text{Tl}$ , and  $^{236}\text{Pu}$ . The calculational method is based on the human metabolic model of ICRP as defined in Publication 2 (ICRP 1959) and as used in previous age-specific dose factor calculations by Hoenes and Soldat (1977). Dose commitment factors are presented for the following organs of reference: total body, bone, liver, kidney, thyroid, lung and lower large intestine.

## ACKNOWLEDGMENTS

This work was performed for the Nuclear Regulatory Commission, Uranium Fuel Licensing Branch.

The authors would like to acknowledge the help of D. L. Haggard in providing radionuclide data and a computer program for calculation of effective absorbed energies. The help of J. D. Schwab in data input preparation is also acknowledged.

## CONTENTS

ABSTRACT . . . . .	iii
ACKNOWLEDGMENTS . . . . .	iv
INTRODUCTION . . . . .	1
METHOD OF CALCULATION . . . . .	2
RESULTS . . . . .	5
REFERENCES . . . . .	10
APPENDIX A--RADIOACTIVE DECAY DATA . . . . .	A-1
APPENDIX B--METABOLIC PARAMETERS . . . . .	B-1

LIST OF TABLES

	<u>Page</u>
Table 1. Mass and Radius of Body Organs for the Four Age Groups...	3
Table 2. Radioiodine Age Dependent Biological Half Times.....	3
Table 3. Infant Inhalation Dose Commitment Factors (mrem/50 yrs per pCi inhaled in first yr).....	6
Table 4. Child Inhalation Dose Commitment Factors (mrem/50 yrs per pCi inhaled in first yr).....	7
Table 5. Teen Inhalation Dose Commitment Factors (mrem/50 yrs per pCi inhaled in first yr).....	8
Table 6. Adult Inhalation Dose Commitment Factors (mrem/50 yrs per pCi inhaled in first yr).....	9
Table A-1 Emission Data for <sup>35</sup> S.....	A-2
Table A-2 Emission Data for <sup>36</sup> Cl.....	A-2
Table A-3 Emission Data for <sup>45</sup> Ca.....	A-2
Table A-4 Emission Data for <sup>67</sup> Ga.....	A-2
Table A-5 Emission Data for <sup>75</sup> Se.....	A-3
Table A-6 Emission Data for <sup>85</sup> Sr.....	A-3
Table A-7 Emission Data for <sup>109</sup> Cd.....	A-4
Table A-8 Emission Data for <sup>113m</sup> In.....	A-4
Table A-9 Emission Data for <sup>113</sup> Sn .....	A-5
Table A-10 Emission Data for <sup>125</sup> I.....	A-5
Table A-11 Emission Data for <sup>133</sup> Ba.....	A-6
Table A-12 Emission Data for <sup>170</sup> Tm.....	A-7
Table A-13 Emission Data for <sup>169</sup> Yb.....	A-7
Table A-14 Emission Data for <sup>182</sup> Ta.....	A-9
Table A-15 Emission Data for <sup>192</sup> Ir.....	A-10
Table A-16 Emission Data for <sup>198</sup> Au.....	A-11
Table A-17 Emission Data for <sup>201</sup> Tl.....	A-12
Table A-18 Emission Data for <sup>204</sup> Tl.....	A-12



Table A-19	Emission Data for $^{236}\text{Pu}$ .....	A-13
Table B-1	Metabolic Parameters.....	B-2



## INTRODUCTION

This report presents inhalation dose commitment factors derived in support of a project for the Nuclear Regulatory Commission's Office of Nuclear Material Safety and Safeguards (NMSS). The overall project involves review of radiological contingency plans submitted by NMSS licensees in response to orders issued February 11, 1981. Dose commitment factors have been calculated for radionuclides which fuel cycle and materials licensees are authorized to possess in significant quantities. This report considers 18 radionuclides for which dose factors are not already available in NUREG-0172 (Hoenes and Soldat 1977). The methods employed in NUREG-0172 were followed for the present calculations to ensure compatibility of results with the previous work.

## METHOD OF CALCULATION

The inhalation dose commitment factors presented here relate chronic one-year radionuclide intake to a 50-year dose commitment. The units of the factors are: mrem in 50 years per pCi inhaled during the first year. The dose received by an individual is dependent on several metabolic parameters that change with age. Therefore four age groups were considered in estimating the dose commitment factors. The age groups considered are "infant" (0 to 1 year old), "child" (1 to 11 years old), "teen" (11 to 17 years old) and "adult" (17 years and older).

The calculations require the following types of data:

- radioactive decay data
- metabolic parameters
- organ mass and diameter

The basic radioactive decay data as used in the present calculations is tabulated in Appendix A with appropriate references. The radionuclide specific metabolic parameters are given in Appendix B. The remaining data is presented in this section.

The radioactive decay data is used to estimate the effective absorbed energy in each organ of interest. The radiations considered are:

- photons (x-rays and gamma rays)
- beta particles ( $e^-$  and  $e^+$ )
- conversion electrons (and Auger electrons)
- alpha particles

The methods and equations defined in ICRP Publication 2 (1959) are used to estimate the effective absorbed energy for these radiation types. These effective absorbed energies are presented in Appendix B for each age group and organ of interest for each radionuclide. The effective energies for photon radiation are dependent on organ radius. The organ sizes and masses used in this study are given in Table 1 and are taken from Hoenes and Soldat (1977). The radius of each organ at a given age is assumed to be proportional to the cube root of the organ mass relative to the adult.

$$(X)_{\text{age}} = \left( \frac{X}{(\text{mass})^{1/3}} \right)_{\text{adult}} \cdot (\text{mass})_{\text{age}}^{1/3}$$

The mass of the contents of the lower large intestine (GI-LLI) is assumed to be proportional to the total body mass.

The chemical, radiological and biological parameters for each radionuclide used to calculate dose commitment factors are presented in Appendix B. These parameters include:

- physical half-life (days)
- biological half-time (days)
- effective half-life (days)

TABLE 1. Mass and Radius of Body Organs for the Four Age Groups\*

Organ	Infant		Child		Teenager		Adult	
	Mass (g)	Radius (cm)	Mass (g)	Radius (cm)	Mass (g)	Radius (cm)	Mass (g)	Radius (cm)
Bone	770	2.4	1,640	3	4,900	4	7,000	5
Liver	200	5	530	7	1,200	9	1,700	10
Total Body	7,700	14	16,400	20	49,000	27	70,000	30
Thyroid	2	1.4	5	2	15	2.7	20	3
Kidney	55	4	100	5	210	6	300	7
Lung	110	5	300	7	580	8	1,000	10
GI-LLI	16	2.4	35	3	100	4	150	5

\*Taken from Hoenes and Soldat (1977).

- fraction reaching organ of reference
- fraction not absorbed before reaching the LLI
- effective absorbed energy (MeV per disintegration)

For the radionuclides considered in this report, the biological half-time, effective half-life, fraction reaching the organ and the fraction not absorbed are assumed to be independent of age except for radioiodine. The variation of biological half-time with age for radioiodine is given in Table 2.

TABLE 2. Radioiodine Age Dependent Biological Half Times

Element	Organ	Half-life (days)			
		Infant	Child	Teenager	Adult
Iodine	Total Body and Thyroid	20	20	50	100

Equations for calculating inhalation dose commitment factors were derived by Hoenes and Soldat (1977) from those given by the ICRP (1959) for body burden and maximum permissible concentration (MPC). These equations use the data and parameters described above. The reader is referred to the Hoenes and Soldat report (1977) for details of the equations.

The model employed for age-dependence assumes an infant to be of uniform size for one year. Then the child immediately grows to the size of a four-year old and remains at this size for 10 years. At 11 years the child immediately becomes the size of a fourteen year old and remains at this size for seven

years at which time the size of an adult is assumed. Hoenes and Soldat present equations for each age period and indicate how each contribution is summed to get the appropriate dose conversion factors.

## RESULTS

The calculated 50-year dose commitment factors for chronic one-year inhalation are presented in Tables 3 through 6. Each table lists dose commitment factors for one age group, six organs and the selected radionuclides. The factors give the 50-year dose commitment (mrem) normalized to unit intake (1 pCi) over a one-year period. The basic radionuclide specific input data used to generate the factors in Tables 3 through 6 are given in Appendix A and Appendix B.

TABLE 3. Infant Inhalation Dose Commitment Factors (mrem/50 yrs per pCi inhaled in first yr)

Radionuclide	Bone	Liver	Total Body	Thyroid	Kidney	Lungs	GI-LLI
<sup>35</sup> S	5.4 x 10 <sup>-5</sup>	0	2.2 x 10 <sup>-5</sup>	0	0	0	1.9 x 10 <sup>-7</sup>
<sup>36</sup> Cl	0	0	4.8 x 10 <sup>-5</sup>	0	0	0	8.7 x 10 <sup>-7</sup>
<sup>45</sup> Ca	2.8 x 10 <sup>-4</sup>	0	6.5 x 10 <sup>-6</sup>	0	0	4.0 x 10 <sup>-4</sup>	7.2 x 10 <sup>-6</sup>
<sup>67</sup> Ga	5.1 x 10 <sup>-6</sup>	3.5 x 10 <sup>-6</sup>	5.8 x 10 <sup>-7</sup>	0	1.2 x 10 <sup>-6</sup>	1.6 x 10 <sup>-5</sup>	1.5 x 10 <sup>-6</sup>
<sup>75</sup> Se	0	2.0 x 10 <sup>-6</sup>	8.3 x 10 <sup>-7</sup>	0	1.9 x 10 <sup>-6</sup>	4.9 x 10 <sup>-4</sup>	3.4 x 10 <sup>-6</sup>
<sup>85</sup> Sr	2.7 x 10 <sup>-5</sup>	0	5.4 x 10 <sup>-6</sup>	0	0	3.0 x 10 <sup>-4</sup>	4.8 x 10 <sup>-6</sup>
<sup>109</sup> Cd+D	0	2.6 x 10 <sup>-4</sup>	1.0 x 10 <sup>-5</sup>	0	2.0 x 10 <sup>-4</sup>	6.2 x 10 <sup>-4</sup>	8.0 x 10 <sup>-6</sup>
<sup>113</sup> Sn+D	6.0 x 10 <sup>-5</sup>	1.6 x 10 <sup>-6</sup>	3.6 x 10 <sup>-6</sup>	1.3 x 10 <sup>-6</sup>	0	7.8 x 10 <sup>-4</sup>	1.2 x 10 <sup>-6</sup>
<sup>125</sup> I	9.6 x 10 <sup>-6</sup>	9.3 x 10 <sup>-6</sup>	6.8 x 10 <sup>-6</sup>	4.0 x 10 <sup>-3</sup>	1.1 x 10 <sup>-5</sup>	0	1.2 x 10 <sup>-7</sup>
<sup>133</sup> Ba	1.9 x 10 <sup>-4</sup>	1.7 x 10 <sup>-6</sup>	1.3 x 10 <sup>-5</sup>	0	8.9 x 10 <sup>-9</sup>	9.1 x 10 <sup>-4</sup>	7.7 x 10 <sup>-6</sup>
<sup>170</sup> Tm	5.7 x 10 <sup>-4</sup>	0	1.8 x 10 <sup>-5</sup>	0	4.7 x 10 <sup>-5</sup>	1.1 x 10 <sup>-3</sup>	2.7 x 10 <sup>-5</sup>
<sup>169</sup> Yb	2.5 x 10 <sup>-5</sup>	0	1.4 x 10 <sup>-6</sup>	0	6.9 x 10 <sup>-6</sup>	3.3 x 10 <sup>-4</sup>	1.2 x 10 <sup>-5</sup>
<sup>182</sup> Ta	6.7 x 10 <sup>-5</sup>	1.6 x 10 <sup>-4</sup>	2.2 x 10 <sup>-5</sup>	0	5.7 x 10 <sup>-5</sup>	1.3 x 10 <sup>-3</sup>	1.8 x 10 <sup>-5</sup>
<sup>192</sup> Ir	0	2.1 x 10 <sup>-5</sup>	3.2 x 10 <sup>-6</sup>	0	2.3 x 10 <sup>-5</sup>	1.1 x 10 <sup>-3</sup>	2.0 x 10 <sup>-5</sup>
<sup>198</sup> Au	0	4.4 x 10 <sup>-8</sup>	3.4 x 10 <sup>-8</sup>	0	1.2 x 10 <sup>-7</sup>	8.5 x 10 <sup>-3</sup>	3.0 x 10 <sup>-5</sup>
<sup>201</sup> Tl	7.9 x 10 <sup>-7</sup>	4.7 x 10 <sup>-7</sup>	4.0 x 10 <sup>-7</sup>	0	2.3 x 10 <sup>-6</sup>	1.4 x 10 <sup>-5</sup>	4.2 x 10 <sup>-6</sup>
<sup>204</sup> Tl	0	1.2 x 10 <sup>-6</sup>	8.0 x 10 <sup>-7</sup>	0	7.7 x 10 <sup>-6</sup>	1.6 x 10 <sup>-3</sup>	2.0 x 10 <sup>-5</sup>
<sup>236</sup> Pu	8.1 x 10 <sup>-1</sup>	5.3 x 10 <sup>-1</sup>	3.6 x 10 <sup>-2</sup>	0	1.3 x 10 <sup>-1</sup>	3.8 x 10 <sup>-1</sup>	4.7 x 10 <sup>-5</sup>



TABLE 4. Child Inhalation Dose Commitment Factors (mrem/50 yrs per pCi inhaled in first yr)

Radionuclide	Bone	Liver	Total Body	Thyroid	Kidney	Lungs	GI-LLI
<sup>35</sup> S	$3.1 \times 10^{-5}$	0	$1.2 \times 10^{-5}$	0	0	0	$1.8 \times 10^{-7}$
<sup>36</sup> Cl	0	0	$2.5 \times 10^{-5}$	0	0	0	$8.5 \times 10^{-7}$
<sup>45</sup> Ca	$2.1 \times 10^{-4}$	0	$4.9 \times 10^{-6}$	0	0	$1.8 \times 10^{-4}$	$7.1 \times 10^{-6}$
<sup>67</sup> Ga	$2.4 \times 10^{-6}$	$1.5 \times 10^{-6}$	$3.2 \times 10^{-7}$	0	$7.0 \times 10^{-7}$	$6.5 \times 10^{-6}$	$1.0 \times 10^{-6}$
<sup>75</sup> Se	0	$1.2 \times 10^{-6}$	$5.7 \times 10^{-7}$	0	$1.4 \times 10^{-6}$	$2.4 \times 10^{-4}$	$3.9 \times 10^{-6}$
<sup>85</sup> Sr	$1.2 \times 10^{-5}$	0	$3.2 \times 10^{-6}$	0	0	$1.5 \times 10^{-4}$	$5.5 \times 10^{-6}$
<sup>109</sup> Cd+D	0	$1.9 \times 10^{-4}$	$8.0 \times 10^{-6}$	0	$1.7 \times 10^{-4}$	$3.0 \times 10^{-4}$	$8.1 \times 10^{-6}$
<sup>113</sup> Sn+D	$3.8 \times 10^{-5}$	$8.9 \times 10^{-7}$	$2.3 \times 10^{-6}$	$7.1 \times 10^{-7}$	0	$3.6 \times 10^{-4}$	$1.3 \times 10^{-6}$
<sup>125</sup> I	$4.6 \times 10^{-6}$	$4.0 \times 10^{-6}$	$3.7 \times 10^{-6}$	$1.8 \times 10^{-3}$	$6.4 \times 10^{-6}$	0	$1.3 \times 10^{-7}$
<sup>133</sup> Ba	$1.1 \times 10^{-4}$	$1.1 \times 10^{-6}$	$1.0 \times 10^{-5}$	0	$5.4 \times 10^{-9}$	$5.2 \times 10^{-4}$	$8.3 \times 10^{-6}$
<sup>170</sup> Tm	$4.1 \times 10^{-4}$	0	$1.2 \times 10^{-5}$	0	$3.5 \times 10^{-6}$	$4.5 \times 10^{-4}$	$2.6 \times 10^{-5}$
<sup>169</sup> Yb	$1.2 \times 10^{-5}$	0	$7.7 \times 10^{-7}$	0	$4.0 \times 10^{-6}$	$1.4 \times 10^{-4}$	$1.3 \times 10^{-5}$
<sup>182</sup> Ta	$4.6 \times 10^{-5}$	$1.1 \times 10^{-4}$	$1.7 \times 10^{-5}$	0	$4.6 \times 10^{-5}$	$6.4 \times 10^{-4}$	$2.0 \times 10^{-5}$
<sup>192</sup> Ir	0	$1.1 \times 10^{-5}$	$2.1 \times 10^{-6}$	0	$1.6 \times 10^{-5}$	$5.6 \times 10^{-4}$	$2.1 \times 10^{-5}$
<sup>198</sup> Au	0	$1.7 \times 10^{-8}$	$1.7 \times 10^{-8}$	0	$6.5 \times 10^{-8}$	$3.3 \times 10^{-5}$	$3.0 \times 10^{-5}$
<sup>201</sup> Tl	$3.7 \times 10^{-7}$	$1.9 \times 10^{-7}$	$2.1 \times 10^{-7}$	0	$1.3 \times 10^{-6}$	$5.4 \times 10^{-6}$	$4.2 \times 10^{-6}$
<sup>204</sup> Tl	0	$4.6 \times 10^{-7}$	$3.7 \times 10^{-7}$	0	$4.2 \times 10^{-6}$	$8.0 \times 10^{-4}$	$2.0 \times 10^{-5}$
<sup>236</sup> Pu	$7.5 \times 10^{-1}$	$4.8 \times 10^{-1}$	$3.4 \times 10^{-2}$	0	$1.2 \times 10^{-1}$	$1.8 \times 10^{-1}$	$4.7 \times 10^{-5}$

TABLE 5. Teen Inhalation Dose Commitment Factors (mrem/50 yrs per pCi inhaled in first yr)

Radionuclide	Bone	Liver	Total Body	Thyroid	Kidney	Lungs	GI-LLI
<sup>35</sup> S	1.2 x 10 <sup>-5</sup>	0	4.2 x 10 <sup>-6</sup>	0	0	0	1.9 x 10 <sup>-7</sup>
<sup>36</sup> Cl	0	0	8.5 x 10 <sup>-6</sup>	0	0	0	8.8 x 10 <sup>-7</sup>
<sup>45</sup> Ca	7.0 x 10 <sup>-5</sup>	0	1.8 x 10 <sup>-6</sup>	0	0	9.3 x 10 <sup>-5</sup>	7.4 x 10 <sup>-6</sup>
<sup>67</sup> Ga	8.3 x 10 <sup>-7</sup>	7.2 x 10 <sup>-7</sup>	1.2 x 10 <sup>-7</sup>	0	3.5 x 10 <sup>-7</sup>	3.5 x 10 <sup>-6</sup>	2.8 x 10 <sup>-5</sup>
<sup>75</sup> Se	0	7.0 x 10 <sup>-7</sup>	2.6 x 10 <sup>-7</sup>	0	8.8 x 10 <sup>-7</sup>	1.3 x 10 <sup>-4</sup>	4.9 x 10 <sup>-6</sup>
<sup>85</sup> Sr	5.0 x 10 <sup>-6</sup>	0	1.3 x 10 <sup>-6</sup>	0	0	8.8 x 10 <sup>-5</sup>	6.9 x 10 <sup>-6</sup>
<sup>109</sup> Cd+D	0	1.0 x 10 <sup>-4</sup>	3.4 x 10 <sup>-6</sup>	0	6.7 x 10 <sup>-5</sup>	1.6 x 10 <sup>-4</sup>	8.6 x 10 <sup>-6</sup>
<sup>113</sup> Sn+D	1.5 x 10 <sup>-5</sup>	4.7 x 10 <sup>-7</sup>	9.7 x 10 <sup>-7</sup>	2.9 x 10 <sup>-7</sup>	0	2.0 x 10 <sup>-4</sup>	1.5 x 10 <sup>-6</sup>
<sup>125</sup> I	1.6 x 10 <sup>-6</sup>	2.0 x 10 <sup>-6</sup>	2.5 x 10 <sup>-6</sup>	1.3 x 10 <sup>-3</sup>	3.2 x 10 <sup>-6</sup>	0	1.4 x 10 <sup>-7</sup>
<sup>133</sup> Ba	4.7 x 10 <sup>-5</sup>	8.0 x 10 <sup>-7</sup>	3.3 x 10 <sup>-6</sup>	0	2.8 x 10 <sup>-9</sup>	2.9 x 10 <sup>-4</sup>	9.7 x 10 <sup>-6</sup>
<sup>170</sup> Tm	1.8 x 10 <sup>-4</sup>	0	5.1 x 10 <sup>-6</sup>	0	2.0 x 10 <sup>-5</sup>	2.3 x 10 <sup>-4</sup>	2.8 x 10 <sup>-5</sup>
<sup>169</sup> Yb	6.1 x 10 <sup>-6</sup>	0	4.3 x 10 <sup>-7</sup>	0	3.0 x 10 <sup>-6</sup>	7.6 x 10 <sup>-5</sup>	1.4 x 10 <sup>-5</sup>
<sup>182</sup> Ta	2.0 x 10 <sup>-4</sup>	6.2 x 10 <sup>-5</sup>	8.3 x 10 <sup>-6</sup>	0	2.8 x 10 <sup>-5</sup>	3.6 x 10 <sup>-4</sup>	2.3 x 10 <sup>-5</sup>
<sup>192</sup> Ir	0	6.1 x 10 <sup>-6</sup>	8.7 x 10 <sup>-7</sup>	0	8.9 x 10 <sup>-6</sup>	3.1 x 10 <sup>-4</sup>	2.3 x 10 <sup>-5</sup>
<sup>198</sup> Au	0	8.1 x 10 <sup>-9</sup>	6.4 x 10 <sup>-9</sup>	0	3.2 x 10 <sup>-8</sup>	1.7 x 10 <sup>-5</sup>	2.9 x 10 <sup>-5</sup>
<sup>201</sup> Tl	1.3 x 10 <sup>-7</sup>	9.0 x 10 <sup>-8</sup>	7.7 x 10 <sup>-8</sup>	0	6.6 x 10 <sup>-7</sup>	2.9 x 10 <sup>-6</sup>	4.3 x 10 <sup>-6</sup>
<sup>204</sup> Tl	0	2.0 x 10 <sup>-7</sup>	1.3 x 10 <sup>-7</sup>	0	2.0 x 10 <sup>-6</sup>	4.1 x 10 <sup>-4</sup>	2.1 x 10 <sup>-5</sup>
<sup>236</sup> Pu	2.6 x 10 <sup>-1</sup>	2.1 x 10 <sup>-1</sup>	1.2 x 10 <sup>-2</sup>	0	5.4 x 10 <sup>-2</sup>	9.5 x 10 <sup>-2</sup>	4.8 x 10 <sup>-5</sup>

TABLE 6. Adult Inhalation Dose Commitment Factors (mrem/50 yrs per pCi inhaled in first yr)

Radionuclide	Bone	Liver	Total Body	Thyroid	Kidney	Lungs	GI-LLI
<sup>35</sup> S	$6.8 \times 10^{-6}$	0	$2.6 \times 10^{-6}$	0	0	0	$1.8 \times 10^{-7}$
<sup>36</sup> Cl	0	0	$5.9 \times 10^{-6}$	0	0	0	$8.3 \times 10^{-7}$
<sup>45</sup> Ca	$4.9 \times 10^{-5}$	0	$1.1 \times 10^{-6}$	0	0	$5.4 \times 10^{-5}$	$6.9 \times 10^{-6}$
<sup>67</sup> Ga	$6.0 \times 10^{-7}$	$5.3 \times 10^{-7}$	$8.9 \times 10^{-8}$	0	$2.6 \times 10^{-7}$	$2.2 \times 10^{-6}$	$2.0 \times 10^{-5}$
<sup>75</sup> Se	0	$4.3 \times 10^{-7}$	$1.6 \times 10^{-7}$	0	$5.9 \times 10^{-7}$	$8.6 \times 10^{-5}$	$5.4 \times 10^{-6}$
<sup>85</sup> Sr	$4.0 \times 10^{-6}$	0	$9.7 \times 10^{-5}$	0	0	$6.0 \times 10^{-5}$	$7.6 \times 10^{-6}$
<sup>109</sup> Cd+D	0	$4.9 \times 10^{-5}$	$1.6 \times 10^{-6}$	0	$4.7 \times 10^{-5}$	$9.1 \times 10^{-5}$	$8.2 \times 10^{-6}$
<sup>113</sup> Sn+D	$8.2 \times 10^{-6}$	$2.7 \times 10^{-7}$	$5.6 \times 10^{-7}$	$1.7 \times 10^{-7}$	0	$1.2 \times 10^{-4}$	$1.5 \times 10^{-6}$
<sup>125</sup> I	$1.1 \times 10^{-6}$	$1.4 \times 10^{-6}$	$2.3 \times 10^{-6}$	$1.2 \times 10^{-3}$	$2.4 \times 10^{-6}$	0	$1.4 \times 10^{-7}$
<sup>133</sup> Ba	$9.5 \times 10^{-6}$	$4.2 \times 10^{-7}$	$2.5 \times 10^{-6}$	0	$2.1 \times 10^{-9}$	$1.9 \times 10^{-4}$	$1.0 \times 10^{-5}$
<sup>170</sup> Tm	$8.3 \times 10^{-5}$	0	$2.4 \times 10^{-6}$	0	$9.8 \times 10^{-6}$	$1.3 \times 10^{-4}$	$2.6 \times 10^{-5}$
<sup>169</sup> Yb	$3.6 \times 10^{-6}$	0	$2.1 \times 10^{-7}$	0	$1.5 \times 10^{-6}$	$4.7 \times 10^{-5}$	$1.4 \times 10^{-5}$
<sup>182</sup> Ta	$1.1 \times 10^{-5}$	$3.3 \times 10^{-5}$	$4.4 \times 10^{-6}$	0	$1.5 \times 10^{-5}$	$2.3 \times 10^{-4}$	$2.4 \times 10^{-5}$
<sup>192</sup> Ir	0	$3.8 \times 10^{-6}$	$5.7 \times 10^{-7}$	0	$5.4 \times 10^{-6}$	$2.0 \times 10^{-4}$	$2.4 \times 10^{-5}$
<sup>198</sup> Au	0	$5.7 \times 10^{-9}$	$4.5 \times 10^{-8}$	0	$2.3 \times 10^{-8}$	$1.0 \times 10^{-5}$	$2.7 \times 10^{-5}$
<sup>201</sup> Tl	$9.0 \times 10^{-8}$	$6.5 \times 10^{-8}$	$5.6 \times 10^{-8}$	0	$4.7 \times 10^{-7}$	$1.8 \times 10^{-6}$	$4.0 \times 10^{-6}$
<sup>204</sup> Tl	0	$1.4 \times 10^{-7}$	$8.8 \times 10^{-8}$	0	$1.4 \times 10^{-6}$	$2.4 \times 10^{-4}$	$1.9 \times 10^{-5}$
<sup>236</sup> Pu	$1.6 \times 10^{-1}$	$1.2 \times 10^{-1}$	$7.0 \times 10^{-3}$	0	$3.2 \times 10^{-2}$	$7.2 \times 10^{-2}$	$4.6 \times 10^{-5}$

## REFERENCES

Hoenes, G.R., and J.K. Soldat. 1977. Age-specific Radiation Dose Commitment Factors for a One-Year Chronic Intake. NUREG-0172. Pacific Northwest Laboratory for U.S. Nuclear Regulatory Commission, Washington, D.C.

ICRP. 1959. Report of Committee II on Permissible Dose for Internal Radiation. International Commission on Radiological Protection, ICRP Publication 2, Pergammon Press, Oxford.

APPENDIX A

RADIOACTIVE DECAY DATA



## APPENDIX A--RADIOACTIVE DECAY DATA

The basic decay data for each radionuclide is given in Tables A-1 through A-11 as follows.

<u>Table</u>	<u>Radionuclide</u>
A-1	$^{35}\text{S}$
A-2	$^{36}\text{Cl}$
A-3	$^{45}\text{Ca}$
A-4	$^{67}\text{Ga}$
A-5	$^{75}\text{Se}$
A-6	$^{85}\text{Sr}$
A-7	$^{109}\text{Cd}$
A-8	$^{113}\text{In}$ (daughter of $^{113}\text{Sn}$ )
A-9	$^{113}\text{Sn}$
A-10	$^{125}\text{I}$
A-11	$^{133}\text{Ba}$
A-12	$^{170}\text{Tm}$
A-13	$^{169}\text{Yb}$
A-14	$^{182}\text{Ta}$
A-15	$^{192}\text{Ir}$
A-16	$^{198}\text{Au}$
A-17	$^{201}\text{Tl}$
A-18	$^{204}\text{Tl}$
A-19	$^{236}\text{Pu}$

The source of data is provided by each radiation type. The references are listed at the end of this appendix. The tables only include significant radiations and are not intended to serve as a complete list of radiations.

TABLE A-1. Emission Data for  $^{35}\text{S}$

Beta Particles (Martin 1976)

<u>Maximum Energy (MeV)</u>	<u>Intensity (%)</u>
0.1674	100

TABLE A-2. Emission Data for  $^{36}\text{Cl}$

Beta Particles (Kocher 1981)

<u>Maximum Energy (MeV)</u>	<u>Intensity (%)</u>
0.710	99

TABLE A-3. Emission Data for  $^{45}\text{Ca}$

Beta Particles (Martin 1976)

<u>Maximum Energy (MeV)</u>	<u>Intensity (%)</u>
0.2587	100

TABLE A-4. Emission Data for  $^{67}\text{Ga}$

Photons (Kocher 1981)

<u>Type</u>	<u>Energy (MeV)</u>	<u>Intensity (%)</u>
Zn X-Rays	0.0086	48.5
	0.0096	6.6
Gamma	0.091	2.9
	0.093	35.7
	0.185	19.7
	0.209	2.2
	0.300	16.0
	0.394	4.5
	0.888	0.14

Electrons (Kocher 1981)

<u>Energy (MeV)</u>	<u>Intensity (%)</u>
0.0010	165
0.0075	60
0.0816	0.2
0.0837	26.8
0.0921	3.3
0.0932	1.1
0.1749	0.33



TABLE A-5. Emission Data for <sup>75</sup>Se

Photons (Martin 1976)		
Type	Energy MeV	Intensity (%)
As X-rays	0.011	48.
	0.012	7.3
Gamma	0.066	1.0
	0.097	3.2
	0.121	16.7
	0.136	56.8
	0.199	1.4
	0.265	59.5
	0.280	25.1
	0.304	1.4
	0.401	11.8

Electrons (Martin 1976)		
	Energy MeV	Intensity (%)
	0.0125	4.4
	0.0229	0.9
	0.0242	0.1
	0.0542	0.36
	0.0850	2.5
	0.0952	0.33
	0.109	0.62
	0.124	1.5
	0.134	0.16
	0.253	0.38
	0.268	0.18

TABLE A-6. Emission Data for <sup>85</sup>Sr

Photons (Kocher 1981)		
Type	Energy MeV	Intensity (%)
Rb X-Rays	0.0017	1.6
	0.0133	17.7
	0.0134	33.0
	0.015	8.7
Gamma	0.5139	99.3

Electrons (Kocher 1981)		
	Energy MeV	Intensity (%)
	0.0017	108.2
	0.0114	29.1
	0.4988	0.63

TABLE A-7. Emission Data for  $^{109}\text{Cd}^*$

Photons (Kocher 1981)		
Type	Energy MeV	Intensity (%)
Ag X-Rays	0.003	11.1
	0.0220	28.5
	0.0222	54.0
	0.0249	17.4
Gamma	0.0880	3.7

Electrons (Kocher 1981)		
	Energy MeV	Intensity (%)
	0.0026	166
	0.0185	20.5
	0.0625	41.7
	0.0842	44.0
	0.0873	8.9
	0.0879	1.6

\*Includes radiations from  $^{109\text{m}}\text{Ag}$ .

TABLE A-8. Emission Data for  $^{113\text{m}}\text{In}$

Photons (Martin 1976)		
Type	Energy (MeV)	Intensity (%)
In X-rays	0.0240	6.8
	0.0242	12.9
	0.0273	4.3
Gamma	0.392	64.9

Electrons (Martin 1976)		
	Energy (MeV)	Intensity (%)
	0.0028	29.7
	0.020	4.2
	0.364	28.2
	0.387	5.5
	0.391	1.1
	0.392	0.25

TABLE A-9. Emission Data for  $^{113}\text{Sn}$

<u>Type</u>	Photons (Martin 1976)	
	<u>Energy (MeV)</u>	<u>Intensity (%)</u>
In X-rays	0.0240	20.7
	0.0242	39.0
	0.0273	13.0
Gamma	0.2550	2.12
Electrons (Martin 1976)		
	<u>Energy (MeV)</u>	<u>Intensity (%)</u>
	0.020	12.8
	0.0028	85.0

TABLE A-10. Emission Data for  $^{125}\text{I}$

<u>Type</u>	Photons (Martin 1976)	
	<u>Energy (MeV)</u>	<u>Intensity (%)</u>
Te X-rays	0.0038	15.
	0.0272	39.8
	0.0275	74.2
	0.031	25.8
Gamma	0.0353	6.7
Electrons (Martin 1976)		
	<u>Energy (MeV)</u>	<u>Intensity (%)</u>
	0.0032	156.
	0.0037	80.
	0.0227	20.
	0.0305	10.5
	0.0345	2.2
	0.0353	0.7

TABLE A-11. Emission Data for  $^{133}\text{Ba}$

<u>Type</u>	<u>Photons (Kocher 1981)</u>	
	<u>Energy (MeV)</u>	<u>Intensity (%)</u>
Cs X-Rays	0.0043	17
	0.0306	34.2
	0.0310	63.4
Gamma	0.035	22.8
	0.0532	2.1
	0.0796	2.6
	0.0810	33.0
	0.1606	0.60
	0.2231	0.44
	0.2764	6.9
	0.3028	17.8
	0.3560	60
	0.3839	8.7

<u>Electrons (Kocher 1981)</u>	
<u>Energy (MeV)</u>	<u>Intensity (%)</u>
0.0036	136
0.0172	10.5
0.0255	14.1
0.0436	3.9
0.0450	48
0.0474	1.6
0.0519	0.40
0.0739	0.56
0.0753	7.4
0.0798	1.5
0.0808	0.39
0.2404	0.32
0.2669	0.67
0.3200	1.28
0.3479	0.15
0.3503	0.21

TABLE A-12. Emission Data for  $^{170}\text{Tm}$

Photons (Lederer and Shirley 1978; Kocher 1981)

<u>Type</u>	<u>Energy (MeV)</u>	<u>Intensity (%)</u>
Yb X-rays	0.0514	1.3
	0.0524	2.3
	0.0593	0.9
	0.0610	0.2
Gamma	0.0786	0.39
	0.0843	3.2

Beta Particles (Lederer and Shirley 1978)

<u>Maximum Energy (MeV)</u>	<u>Intensity (%)</u>
0.968	76.
0.884	24.

TABLE A-13. Emission Data for  $^{169}\text{Yb}$

Photons (Kocher 1981)

<u>Type</u>	<u>Energy (MeV)</u>	<u>Intensity (%)</u>
Tm X-Rays	0.0072	51
	0.0498	52.8
	0.0507	93
	0.0575	38.3
Gamma	0.0084	0.33
	0.0208	0.21
	0.0631	43.7
	0.0936	2.7
	0.1098	17.4
	0.1182	1.9
	0.1305	11.1
	0.1772	21.4
	0.1980	34.9
	0.2403	0.12
0.2611	1.8	
0.3077	10.8	

TABLE A.13. (contd)

Electrons (Kocher 1981)

<u>Energy (MeV)</u>	<u>Intensity (%)</u>
0.0037	39.9
0.0057	160
0.0061	95
0.0106	9.3
0.0184	2.1
0.0203	0.68
0.0342	8.4
0.0409	10
0.0504	34.9
0.0530	7.1
0.0588	1.3
0.0608	1.6
0.0626	0.43
0.0711	6.0
0.0835	1.45
0.0913	0.42
0.0997	5.6
0.1075	1.3
0.1081	1.4
0.1093	0.37
0.1159	0.42
0.1178	10.3
0.1204	5.2
0.1282	1.25
0.1300	0.35
0.1386	12.7
0.1671	1.8
0.1749	0.42
0.1878	2.1
0.1956	0.47
0.2483	0.52

TABLE A-14. Emission Data for  $^{182}\text{Ta}$

Photons (Lederer and Shirley 1978; Kocher 1981)

Type	Energy (MeV)	Intensity (%)
W X-rays	0.0580	10.4
	0.0593	18.1
	0.0672	7.7
Gamma	0.0691	1.6
	0.0317	0.9
	0.0427	0.3
	0.0657	2.9
	0.0677	43.7
	0.0847	2.4
	0.100	13.5
	0.110	0.1
	0.114	1.8
	0.116	0.4
	0.152	6.7
	0.156	2.5
	0.179	2.9
	0.198	1.4
	0.222	7.2
	0.229	3.5
	0.264	3.4
	0.928	0.6
	0.960	0.3
	1.002	2.0
1.113	0.4	
1.121	35.1	
1.157	1.0	
1.189	15.7	
1.221	25.9	
1.231	11.0	
1.257	1.5	
1.274	0.7	
1.289	1.4	
1.343	0.2	
1.374	0.2	
1.387	0.1	

Beta Particles (Lederer and Shirley 1978)

Maximum Energy (MeV)	Intensity (%)
0.540	45
0.408	24
0.246	31

TABLE A-15. Emission Data for  $^{192}\text{Ir}$

Photons (Lederer and Shirley 1978)		
Type	Energy (MeV)	Intensity (%)
Os X-rays	0.0615	1.2
	0.0630	2.0
	0.0713	0.7
	0.0734	0.17
Pt X-rays	0.0650	2.5
	0.0670	4.3
	0.0760	1.5
	0.0780	0.4
Gamma	0.136	0.2
	0.201	0.5
	0.206	3.2
	0.283	0.3
	0.296	28.7
	0.308	29.7
	0.317	82.9
	0.374	0.7
	0.416	0.7
	0.421	0.1
	0.468	48.1
	0.485	3.2
	0.489	0.4
	0.589	4.6
	0.604	8.3
0.612	5.4	
0.885	0.3	
1.061	0.1	

Beta Particles (Lederer and Shirley 1978)

Maximum Energy (MeV)	Intensity (%)
0.672	46
0.536	41
0.240	8



TABLE A-16. Emission Data for  $^{198}\text{Au}$

<u>Type</u>	Photons (Martin 1976)	
	<u>Energy (MeV)</u>	<u>Intensity (%)</u>
Hg X-rays	0.010	1.3
	0.0690	0.8
	0.0708	1.4
	0.080	0.6
Gamma	0.412	95.5
	0.676	1.1
	1.090	0.23

Beta Particles (Martin 1976)	
<u>Maximum Energy (MeV)</u>	<u>Intensity (%)</u>
0.285	1.3
0.961	98.7

Electrons (Martin 1976)	
<u>Energy (MeV)</u>	<u>Intensity</u>
0.0076	2.1
0.329	2.9
0.397	1.0
0.408	0.27
0.411	0.1

TABLE A-17. Emission Data for  $^{201}\text{Tl}$

Photons (Kocher 1981)		
<u>Type</u>	<u>Energy (MeV)</u>	<u>Intensity (%)</u>
Hg X-Rays	0.010	44
	0.0689	27.4
	0.0708	46.5
	0.0803	20.5
Gamma	0.0306	0.22
	0.0322	0.22
	0.1353	2.65
	0.1674	10

Electrons (Kocher 1981)		
<u>Type</u>	<u>Energy (MeV)</u>	<u>Intensity (%)</u>
	0.0076	73
	0.0158	8.1
	0.0174	7.0
	0.0270	2.5
	0.0286	2.2
	0.0522	7.5
	0.0538	3.3
	0.0843	15.4
	0.1205	1.3
	0.1318	0.4
	0.1526	2.62
	0.1639	0.61
	0.1666	0.2

TABLE A-18. Emission Data for  $^{204}\text{Tl}$

Photons (Lederer and Shirley 1978)		
<u>Type</u>	<u>Energy (MeV)</u>	<u>Intensity %</u>
Hg X-rays	0.0689	0.41
	0.0708	0.71
	0.0802	0.24
	0.0825	0.04

Beta Particles (Lederer and Shirley 1978)	
<u>Maximum Energy (MeV)</u>	<u>Intensity (%)</u>
0.7635	97.4

TABLE A-19. Emission Data for  $^{236}\text{Pu}$

Photons (Kocher 1981)		
<u>Type</u>	<u>Energy (MeV)</u>	<u>Intensity (%)</u>
U X-Rays	0.0136	13
Electrons (Kocher 1981)		
	<u>Energy (MeV)</u>	<u>Intensity (%)</u>
	0.0099	10.2
	0.0259	23.3
	0.0421	8.5
Alpha Particles (Kocher 1981)		
	<u>Energy (MeV)</u>	<u>Intensity (%)</u>
	5.614	0.18
	5.722	31.8
	5.770	68.1

## APPENDIX A--REFERENCES

- Kocher, D. C. 1981. Radioactive Decay Data Tables. DOE/TIC-11026. U.S. Department of Energy, National Technical Information Service, Springfield, Virginia.
- Lederer, C. M., and V. S. Shirley. 1978. Table of Isotopes. Seventh Edition. John Wiley and Sons, Inc. New York, New York.
- Martin, M. J. 1976. Nuclear Decay Data for Selected Radionuclides. ORNL-5114. Oak Ridge National Laboratory, Oak Ridge, Tennessee.

APPENDIX B

METABOLIC PARAMETERS



## APPENDIX B--METABOLIC PARAMETERS

This appendix presents the metabolic parameters used to calculate the age-specific dose commitment factors for each radionuclide. The data presented here include parameters for both inhalation and ingestion uptake even though only inhalation dose commitment factors are calculated. The ingestion parameter values are included for completeness with respect to corresponding tables in the original report on age-specific dose commitment factors (Hoenes and Soldat 1977).

The data are listed in Table B-1 and include the following parameters.

- Half-life      ● radiological decay half-time (days)
- F-W            ● fraction of radionuclide reaching the organ of reference by ingestion,  $f_w$
- F-\* for GI     ● fraction of radionuclide not absorbed before reaching the lower-large intestine. F-\* is calculated by
- $$F-* = 1 - f_w(\text{Total Body})$$
- and
- $$F-* \geq 0.05$$
- F-A            ● fraction of radionuclide reaching the organ of reference by inhalation,  $f_a$
- F-2'           ● fraction of radionuclides transferred from the blood to the organ of reference,  $f_2'$
- TBIOL         ● biological removal half-time for the radionuclide and organ of reference and each age group (days)
- EPSILON       ● effective absorbed energy for the radionuclide and organ of reference and each age group (MeV).

The values in the column in Table B-1 labeled F-A or F-2' represent either  $f_a$  or  $f_2'$  depending on the organ of reference and the material solubility. For radionuclides assumed to be soluble ( $^{35}\text{S}$ ,  $^{36}\text{Cl}$ , and  $^{125}\text{I}$ ) the value represents  $f_a$  for all organs. For radionuclides assumed to be insoluble the value represents  $f_a$  for lung and GI-LLI and  $f_2'$  for other organs.

The values for the parameter EPSILON were calculated using a small program prepared by N. M. Robinson and modified for the present calculations by D. L. Strenge.

TABLE B-1. Metabolic Parameters

ORGAN	F-W (F-* FOR GI) OR F-2'	INFANT		CHILD		TEEN		ADULT	
		TBIOL (DAY)	EPSILON (MEV)	TBIOL (DAY)	EPSILON (MEV)	TBIOL (DAY)	EPSILON (MEV)	TBIOL (DAY)	EPSILON (MEV)
S35	HALF LIFE = 8.740E 01 DAYS								
BONE	3.00E-02	6.00E 02	2.80E-01	6.00E 02	2.80E-01	6.00E 02	2.80E-01	6.00E 02	2.80E-01
LIVER	0.	0.	0.	0.	0.	0.	0.	0.	0.
TOTAL BODY	1.00E 00	9.00E 01	5.60E-02	9.00E 01	5.60E-02	9.00E 01	5.60E-02	9.00E 01	5.60E-02
THYROID	0.	0.	0.	0.	0.	0.	0.	0.	0.
KIDNEY	0.	0.	0.	0.	0.	0.	0.	0.	0.
LUNG INGES	0.	0.	0.	0.	0.	0.	0.	0.	0.
LUNG INHAL	0.	0.	0.	0.	0.	0.	0.	0.	0.
GI-LLI INGES	5.00E-02	0.	5.60E-02	0.	5.60E-02	0.	5.60E-02	0.	5.60E-02
GI-LLI INHAL	5.00E-02	0.	5.60E-02	0.	5.60E-02	0.	5.60E-02	0.	5.60E-02
CL36	HALF LIFE = 1.100E 08 DAYS								
BONE	0.	0.	0.	0.	0.	0.	0.	0.	0.
LIVER	0.	0.	0.	0.	0.	0.	0.	0.	0.
TOTAL BODY	1.00E 00	7.50E-01	2.58E-01	2.90E 01	2.58E-01	2.90E 01	2.58E-01	2.90E 01	2.58E-01
THYROID	0.	0.	0.	0.	0.	0.	0.	0.	0.
KIDNEY	0.	0.	0.	0.	0.	0.	0.	0.	0.
LUNG INGES	0.	0.	0.	0.	0.	0.	0.	0.	0.
LUNG INHAL	0.	0.	0.	0.	0.	0.	0.	0.	0.
GI-LLI INGES	5.00E-02	0.	2.58E-01	0.	2.58E-01	0.	2.58E-01	0.	2.58E-01
GI-LLI INHAL	5.00E-02	0.	2.58E-01	0.	2.58E-01	0.	2.58E-01	0.	2.58E-01
CA45	HALF LIFE = 1.650E 02 DAYS								
INSOLUBLE FOR INHALATION ONLY									
BONE	5.40E-01	9.00E-01	1.80E 04	4.38E-01	1.80E 04	4.38E-01	1.80E 04	4.38E-01	1.80E 04
LIVER	0.	0.	0.	0.	0.	0.	0.	0.	0.
TOTAL BODY	6.00E-01	1.00E 00	1.64E 04	8.76E-02	1.64E 04	8.76E-02	1.64E 04	8.76E-02	1.64E 04
THYROID	0.	0.	0.	0.	0.	0.	0.	0.	0.
KIDNEY	0.	0.	0.	0.	0.	0.	0.	0.	0.
LUNG INGES	0.	0.	0.	0.	0.	0.	0.	0.	0.
LUNG INHAL	0.	0.	0.	0.	0.	0.	0.	0.	0.
GI-LLI INGES	4.00E-01	0.	1.20E 02	8.76E-02	1.20E 02	8.76E-02	1.20E 02	8.76E-02	1.20E 02
GI-LLI INHAL	1.00E 00	6.20E-01	0.	8.76E-02	0.	8.76E-02	0.	8.76E-02	0.



TABLE B-1. (contd)

GAG7	INSOLUBLE FOR INHALATION ONLY									
	HALF LIFE = 3.261E 00 DAYS					INSOLUBLE FOR INHALATION ONLY				
BONE	3.00E-04	3.00E-01	1.20E 01	2.17E-01	1.20E 01	2.17E-01	1.20E 01	2.25E-01	1.20E 01	2.31E-01
LIVER	2.50E-04	2.50E-01	4.80E 00	6.21E-02	4.80E 00	6.92E-02	4.80E 00	7.58E-02	4.80E 00	7.90E-02
TOTAL BODY	1.00E-03	1.00E 00	6.00E 00	9.09E-02	6.00E 00	1.06E-01	6.00E 00	1.21E-01	6.00E 00	1.26E-01
THYROID	0.	0.	0.	0.	0.	0.	0.	0.	0.	0.
KIDNEY	2.00E-05	2.00E-02	9.00E 00	5.84E-02	9.00E 00	6.21E-02	9.00E 00	6.57E-02	9.00E 00	6.92E-02
LUNG INGES	0.	0.	0.	0.	0.	0.	0.	0.	0.	0.
LUNG INHAL	0.	1.20E-01	1.20E 02	6.21E-02	1.20E 02	6.92E-02	1.20E 02	7.26E-02	1.20E 02	7.90E-02
GI-LLI INGES	1.00E 00	0.	0.	5.23E-02	0.	5.46E-02	0.	5.84E-02	0.	6.21E-02
GI-LLI INHAL	1.00E 00	6.20E-01	0.	5.23E-02	0.	5.46E-02	0.	5.84E-02	0.	6.21E-02
SE75	INSOLUBLE FOR INHALATION ONLY									
HALF LIFE = 1.185E 02 DAYS										
BONE	0.	0.	0.	0.	0.	0.	0.	0.	0.	0.
LIVER	6.00E-02	7.00E-02	2.40E 01	6.80E-02	2.40E 01	8.70E-02	2.40E 01	1.05E-01	2.40E 01	1.14E-01
TOTAL BODY	9.00E-01	1.00E 00	1.10E 01	1.46E-01	1.10E 01	1.88E-01	1.10E 01	2.28E-01	1.10E 01	2.42E-01
THYROID	0.	0.	0.	0.	0.	0.	0.	0.	0.	0.
KIDNEY	4.00E-02	4.00E-02	1.10E 01	5.80E-02	1.10E 01	6.80E-02	1.10E 01	8.20E-02	1.10E 01	9.60E-02
LUNG INGES	0.	0.	0.	0.	0.	0.	0.	0.	0.	0.
LUNG INHAL	0.	1.20E-01	1.20E 02	6.80E-02	1.20E 02	8.70E-02	1.20E 02	9.60E-02	1.20E 02	1.14E-01
GI-LLI INGES	1.00E-01	0.	0.	4.10E-02	0.	4.80E-02	0.	5.80E-02	0.	6.80E-02
GI-LLI INHAL	1.00E 00	6.20E-01	0.	4.10E-02	0.	4.80E-02	0.	5.80E-02	0.	6.80E-02
SR85	INSOLUBLE FOR INHALATION ONLY									
HALF LIFE = 6.484E 01 DAYS										
BONE	2.10E-01	7.00E-01	1.80E 04	1.50E-01	1.80E 04	1.50E-01	1.80E 04	1.75E-01	1.80E 04	1.98E-01
LIVER	0.	0.	0.	0.	0.	0.	0.	0.	0.	0.
TOTAL BODY	3.00E-01	1.00E 00	1.30E 04	2.07E-01	1.30E 04	2.65E-01	1.30E 04	3.19E-01	1.30E 04	3.39E-01
THYROID	0.	0.	0.	0.	0.	0.	0.	0.	0.	0.
KIDNEY	0.	0.	0.	0.	0.	0.	0.	0.	0.	0.
LUNG INGES	0.	0.	0.	0.	0.	0.	0.	0.	0.	0.
LUNG INHAL	0.	1.20E-01	1.20E 02	9.69E-02	1.20E 02	1.24E-01	1.20E 02	1.37E-01	1.20E 02	1.62E-01
GI-LLI INGES	7.00E-01	0.	0.	5.84E-02	0.	6.76E-02	0.	8.25E-02	0.	9.69E-02
GI-LLI INHAL	1.00E 00	6.20E-01	0.	5.84E-02	0.	6.76E-02	0.	8.25E-02	0.	9.69E-02

TABLE B-1. (contd)

ORGAN	F-W (F-* FOR GI) OR F-2'	INFANT		CHILD		TEEN		ADULT	
		TBIOL (DAY)	EPSILON (MEV)	TBIOL (DAY)	EPSILON (MEV)	TBIOL (DAY)	EPSILON (MEV)	TBIOL (DAY)	EPSILON (MEV)
CD109+D	INSOLUBLE FOR INHALATION ONLY								
	HALF LIFE = 4.640E 02 DAYS								
BONE	0.	0.	0.	0.	0.	0.	0.	0.	0.
LIVER	1.90E-03	7.50E-01	1.04E-01	2.00E 02	1.06E-01	2.00E 02	1.07E-01	2.00E 02	1.08E-01
TOTAL BODY	2.50E-03	1.00E 00	2.00E 02	1.09E-01	2.00E 02	1.10E-01	2.00E 02	1.11E-01	2.00E 02
THYROID	0.	0.	0.	0.	0.	0.	0.	0.	0.
KIDNEY	2.50E-04	1.00E-01	3.00E 02	1.02E-01	3.00E 02	1.04E-01	3.00E 02	1.05E-01	3.00E 02
LUNG INGES	0.	0.	0.	0.	0.	0.	0.	0.	0.
LUNG INHAL	0.	1.20E-01	1.04E-01	1.20E 02	1.06E-01	1.20E 02	1.07E-01	1.20E 02	1.08E-01
GI-LLI INGES	1.00E 00	0.	9.77E-02	0.	9.95E-02	0.	1.02E-01	0.	1.04E-01
GI-LLI INHAL	1.00E 00	6.20E-01	0.	9.77E-02	0.	9.95E-02	0.	1.02E-01	0.
SN113+D	INSOLUBLE FOR INHALATION ONLY								
	HALF LIFE = 1.150E 02 DAYS								
BONE	2.00E-02	3.00E-01	1.00E 02	7.68E-01	1.00E 02	7.68E-01	1.00E 02	7.81E-01	1.00E 02
LIVER	5.00E-04	1.00E-02	7.00E 01	1.92E-01	7.00E 01	2.10E-01	7.00E 01	2.24E-01	7.00E 01
TOTAL BODY	5.00E-02	1.00E 00	3.50E 01	2.56E-01	3.50E 01	2.87E-01	3.50E 01	3.15E-01	3.50E 01
THYROID	5.00E-06	1.00E-04	7.00E 01	1.57E-01	7.00E 01	1.64E-01	7.00E 01	1.71E-01	7.00E 01
KIDNEY	0.	0.	0.	0.	0.	0.	0.	0.	0.
LUNG INGES	0.	0.	0.	0.	0.	0.	0.	0.	0.
LUNG INHAL	0.	1.20E-01	1.20E 02	1.92E-01	1.20E 02	2.10E-01	1.20E 02	2.17E-01	1.20E 02
GI-LLI INGES	9.50E-01	0.	0.	1.46E-02	0.	1.58E-02	0.	1.76E-02	0.
GI-LLI INHAL	1.00E 00	6.20E-01	0.	1.46E-02	0.	1.58E-02	0.	1.76E-02	0.
I125	HALF LIFE = 6.020E 01 DAYS								
BONE	7.00E-02	5.30E-02	1.40E 01	1.69E-01	1.40E 01	1.69E-01	1.40E 01	1.71E-01	1.40E 01
LIVER	1.20E-01	9.00E-02	7.00E 00	4.53E-02	7.00E 00	5.09E-02	7.00E 00	5.57E-02	7.00E 00
TOTAL BODY	1.00E 00	7.50E-01	2.00E 01	6.45E-02	2.00E 01	7.14E-02	5.00E 01	7.65E-02	1.00E 02
THYROID	3.00E-01	2.30E-01	2.00E 01	3.23E-02	2.00E 01	3.48E-02	5.00E 01	3.75E-02	1.00E 02
KIDNEY	4.00E-02	3.00E-02	7.00E 00	4.21E-02	7.00E 00	4.53E-02	7.00E 00	4.81E-02	7.00E 00
LUNG INGES	0.	0.	0.	0.	0.	0.	0.	0.	0.
LUNG INHAL	0.	0.	0.	0.	0.	0.	0.	0.	0.
GI-LLI INGES	5.00E-02	0.	0.	3.64E-02	0.	3.86E-02	0.	4.21E-02	0.
GI-LLI INHAL	5.00E-02	5.00E-01	0.	3.64E-02	0.	3.86E-02	0.	4.21E-02	0.

TABLE B-1. (contd)

ORGAN	F-W (F-* FOR GI) OR F-2'	INFANT		CHILD		TEEN		ADULT	
		TBIOL (DAY)	EPSILON (MEV)	TBIOL (DAY)	EPSILON (MEV)	TBIOL (DAY)	EPSILON (MEV)	TBIOL (DAY)	EPSILON (MEV)
BA133	HALF LIFE = 3.840E 03 DAYS								
	INSOLUBLE FOR INHALATION ONLY								
BONE	3.50E-02	7.00E-01	3.83E-01	6.50E 01	3.83E-01	6.50E 01	4.01E-01	6.50E 01	4.18E-01
LIVER	3.00E-05	6.00E-04	1.27E-01	9.75E 02	1.50E-01	9.75E 02	1.71E-01	9.75E 02	1.80E-01
TOTAL BODY	5.00E-02	1.00E 00	2.15E-01	6.50E 01	2.57E-01	6.50E 01	2.96E-01	6.50E 01	3.10E-01
THYROID	0.	0.	0.	0.	0.	0.	0.	0.	0.
KIDNEY	5.00E-06	1.00E-04	1.15E-01	8.50E 00	1.27E-01	8.50E 00	1.39E-01	8.50E 00	1.50E-01
LUNG INGES	1.00E-05	0.	1.27E-01	6.50E 03	1.50E-01	6.50E 03	1.61E-01	6.50E 03	1.80E-01
LUNG INHAL	0.	1.20E-01	1.27E-01	1.20E 02	1.50E-01	1.20E 02	1.61E-01	1.20E 02	1.80E-01
GI-LLI INGES	9.50E-01	0.	9.36E-02	0.	1.02E-01	0.	1.15E-01	0.	1.27E-01
GI-LLI INHAL	1.00E 00	6.20E-01	9.36E-02	0.	1.02E-01	0.	1.15E-01	0.	1.27E-01
TM170	HALF LIFE = 1.286E 02 DAYS								
	INSOLUBLE FOR INHALATION ONLY								
BONE	6.50E-05	6.50E-01	1.16E 00	1.00E 03	1.16E 00	1.00E 03	1.16E 00	1.00E 03	1.16E 00
LIVER	0.	0.	0.	0.	0.	0.	0.	0.	0.
TOTAL BODY	1.00E-04	1.00E 00	3.26E-01	6.75E 02	3.27E-01	6.75E 02	3.27E-01	6.75E 02	3.28E-01
THYROID	0.	0.	0.	0.	0.	0.	0.	0.	0.
KIDNEY	2.00E-06	2.00E-02	3.25E-01	3.35E 02	3.25E-01	3.35E 02	3.25E-01	3.35E 02	3.25E-01
LUNG INGES	0.	0.	0.	0.	0.	0.	0.	0.	0.
LUNG INHAL	0.	1.20E-01	3.25E-01	1.20E 02	3.25E-01	1.20E 02	3.25E-01	1.20E 02	3.26E-01
GI-LLI INGES	1.00E 00	0.	3.25E-01	0.	3.25E-01	0.	3.25E-01	0.	3.25E-01
GI-LLI INHAL	1.00E 00	6.20E-01	3.25E-01	0.	3.25E-01	0.	3.25E-01	0.	3.25E-01
YB169	HALF LIFE = 3.197E 01 DAYS								
	INSOLUBLE FOR INHALATION ONLY								
BONE	5.80E-05	5.80E-01	7.15E-01	1.00E 03	7.15E-01	1.00E 03	7.36E-01	1.00E 03	7.54E-01
LIVER	0.	0.	0.	0.	0.	0.	0.	0.	0.
TOTAL BODY	1.00E-04	1.00E 00	2.38E-01	6.85E 02	2.72E-01	6.85E 02	3.04E-01	6.85E 02	3.15E-01
THYROID	0.	0.	0.	0.	0.	0.	0.	0.	0.
KIDNEY	5.00E-06	5.00E-02	1.64E-01	6.85E 02	1.73E-01	6.85E 02	1.81E-01	6.85E 02	1.89E-01
LUNG INGES	0.	0.	0.	0.	0.	0.	0.	0.	0.
LUNG INHAL	0.	1.20E-01	1.73E-01	1.20E 02	1.89E-01	1.20E 02	1.97E-01	1.20E 02	2.11E-01
GI-LLI INGES	1.00E 00	0.	1.50E-01	0.	1.56E-01	0.	1.64E-01	0.	1.73E-01
GI-LLI INHAL	1.00E 00	6.20E-01	1.50E-01	0.	1.56E-01	0.	1.64E-01	0.	1.73E-01

TABLE B-1. (contd)

ORGAN	F-H (F-* FOR GI)	F-A OR F-2'	INFANT		CHILD		TEEN		ADULT	
			TBIOL (DAY)	EPSILON (MEV)	TBIOL (DAY)	EPSILON (MEV)	TBIOL (DAY)	EPSILON (MEV)	TBIOL (DAY)	EPSILON (MEV)
TA182	HALF LIFE = 1.150E 02 DAYS									
INSOLUBLE FOR INHALATION ONLY										
BONE	2.00E-05	2.00E-01	3.00E 02	8.66E-01	3.00E 02	8.66E-01	3.00E 02	9.25E-01	3.00E 02	9.80E-01
LIVER	3.00E-05	3.00E-01	4.00E 02	3.07E-01	4.00E 02	3.69E-01	4.00E 02	4.25E-01	4.00E 02	4.57E-01
TOTAL BODY	1.00E-04	1.00E 00	2.40E 02	5.61E-01	2.40E 02	6.97E-01	2.40E 02	8.27E-01	2.40E 02	8.75E-01
THYROID	0.	0.	0.	0.	0.	0.	0.	0.	0.	0.
KIDNEY	3.00E-06	3.00E-02	4.00E 02	2.74E-01	4.00E 02	3.07E-01	4.00E 02	3.38E-01	4.00E 02	3.69E-01
LUNG INGES	0.	0.	0.	0.	0.	0.	0.	0.	0.	0.
LUNG INHAL	0.	1.20E-01	1.20E 02	3.07E-01	1.20E 02	3.69E-01	1.20E 02	3.99E-01	1.20E 02	4.57E-01
GI-LLI INGES	1.00E 00	0.	0.	2.20E-01	0.	2.41E-01	0.	2.74E-01	0.	3.07E-01
GI-LLI INHAL	1.00E 00	6.20E-01	0.	2.20E-01	0.	2.41E-01	0.	2.74E-01	0.	3.07E-01
IR192	HALF LIFE = 7.420E 01 DAYS									
INSOLUBLE FOR INHALATION ONLY										
BONE	0.	0.	0.	0.	0.	0.	0.	0.	0.	0.
LIVER	2.30E-02	2.30E-01	2.70E 01	2.99E-01	2.70E 01	3.43E-01	2.70E 01	3.84E-01	2.70E 01	4.03E-01
TOTAL BODY	1.00E-01	1.00E 00	2.00E 01	4.75E-01	2.00E 01	5.66E-01	2.00E 01	6.53E-01	2.00E 01	6.84E-01
THYROID	0.	0.	0.	0.	0.	0.	0.	0.	0.	0.
KIDNEY	4.50E-03	4.50E-02	5.00E 01	2.77E-01	5.00E 01	2.99E-01	5.00E 01	3.21E-01	5.00E 01	3.43E-01
LUNG INGES	0.	0.	0.	0.	0.	0.	0.	0.	0.	0.
LUNG INHAL	0.	1.20E-01	1.20E 02	2.99E-01	1.20E 02	3.43E-01	1.20E 02	3.64E-01	1.20E 02	4.03E-01
GI-LLI INGES	9.00E-01	0.	0.	2.39E-01	0.	2.53E-01	0.	2.77E-01	0.	2.99E-01
GI-LLI INHAL	1.00E 00	6.20E-01	0.	2.39E-01	0.	2.53E-01	0.	2.77E-01	0.	2.99E-01
AU198	HALF LIFE = 2.700E 00 DAYS									
INSOLUBLE FOR INHALATION ONLY										
BONE	0.	0.	0.	0.	0.	0.	0.	0.	0.	0.
LIVER	4.00E-03	4.00E-02	3.00E 02	3.98E-01	3.00E 02	4.19E-01	3.00E 02	4.40E-01	3.00E 02	4.49E-01
TOTAL BODY	1.00E-01	1.00E 00	1.20E 02	4.85E-01	1.20E 02	5.31E-01	1.20E 02	5.74E-01	1.20E 02	5.89E-01
THYROID	0.	0.	0.	0.	0.	0.	0.	0.	0.	0.
KIDNEY	3.00E-03	3.00E-02	2.80E 02	3.86E-01	2.80E 02	3.98E-01	2.80E 02	4.09E-01	2.80E 02	4.19E-01
LUNG INGES	0.	0.	0.	0.	0.	0.	0.	0.	0.	0.
LUNG INHAL	0.	1.20E-01	1.20E 02	3.98E-01	1.20E 02	4.19E-01	1.20E 02	4.30E-01	1.20E 02	4.49E-01
GI-LLI INGES	9.00E-01	0.	0.	3.67E-01	0.	3.74E-01	0.	3.86E-01	0.	3.98E-01
GI-LLI INHAL	1.00E 00	6.20E-01	0.	3.67E-01	0.	3.74E-01	0.	3.86E-01	0.	3.98E-01

TABLE B-1. (contd)

ORGAN	F-W (F-* FOR GI)	F-A OR F-2'	INFANT		CHILD		TEEN		ADULT	
			TBIOL (DAY)	EPSILON (MEV)	TBIOL (DAY)	EPSILON (MEV)	TBIOL (DAY)	EPSILON (MEV)	TBIOL (DAY)	EPSILON (MEV)
TL201	HALF LIFE = 3.040E 00 DAYS									
INSOLUBLE FOR INHALATION ONLY										
BONE	2.50E-02	5.50E-02	7.00E 00	2.37E-01	7.00E 00	2.37E-01	7.00E 00	2.43E-01	7.00E 00	2.48E-01
LIVER	2.00E-02	4.00E-02	5.00E 00	5.74E-02	5.00E 00	6.14E-02	5.00E 00	6.52E-02	5.00E 00	6.70E-02
TOTAL BODY	4.50E-01	1.00E 00	5.00E 00	7.38E-02	5.00E 00	8.28E-02	5.00E 00	9.16E-02	5.00E 00	9.49E-02
THYROID	0.	0.	0.	0.	0.	0.	0.	0.	0.	0.
KIDNEY	2.30E-02	5.00E-02	7.00E 00	5.53E-02	7.00E 00	5.74E-02	7.00E 00	5.94E-02	7.00E 00	6.14E-02
LUNG INGES	2.70E-03	0.	6.00E 00	5.74E-02	6.00E 00	6.14E-02	6.00E 00	6.33E-02	6.00E 00	6.70E-02
LUNG INHAL	0.	1.20E-01	1.20E 02	5.74E-02	1.20E 02	6.14E-02	1.20E 02	6.33E-02	1.20E 02	6.70E-02
GI-LLI INGES	5.50E-01	0.	0.	5.18E-02	0.	5.32E-02	0.	5.53E-02	0.	5.74E-02
GI-LLI INHAL	1.00E 00	6.20E-01	0.	5.18E-02	0.	5.32E-02	0.	5.53E-02	0.	5.74E-02
TL204	HALF LIFE = 1.377E 03 DAYS									
INSOLUBLE FOR INHALATION ONLY										
BONE	0.	0.	0.	0.	0.	0.	0.	0.	0.	0.
LIVER	2.00E-02	4.00E-02	5.00E 00	2.45E-01	5.00E 00	2.45E-01	5.00E 00	2.45E-01	5.00E 00	2.46E-01
TOTAL BODY	4.50E-01	1.00E 00	5.00E 00	2.46E-01	5.00E 00	2.46E-01	5.00E 00	2.46E-01	5.00E 00	2.46E-01
THYROID	0.	0.	0.	0.	0.	0.	0.	0.	0.	0.
KIDNEY	2.30E-02	5.00E-02	7.00E 00	2.45E-01	7.00E 00	2.45E-01	7.00E 00	2.45E-01	7.00E 00	2.45E-01
LUNG INGES	2.70E-03	0.	6.00E 00	2.45E-01	6.00E 00	2.45E-01	6.00E 00	2.45E-01	6.00E 00	2.46E-01
LUNG INHAL	0.	1.20E-01	1.20E 02	2.45E-01	1.20E 02	2.45E-01	1.20E 02	2.45E-01	1.20E 02	2.46E-01
GI-LLI INGES	5.50E-01	0.	0.	2.45E-01	0.	2.45E-01	0.	2.45E-01	0.	2.45E-01
GI-LLI INHAL	1.00E 00	6.20E-01	0.	2.45E-01	0.	2.45E-01	0.	2.45E-01	0.	2.45E-01
PU236	HALF LIFE = 1.041E 03 DAYS									
INSOLUBLE FOR INHALATION ONLY										
BONE	1.35E-05	4.50E-01	3.65E 04	2.88E 02	3.65E 04	2.88E 02	3.65E 04	2.88E 02	3.65E 04	2.88E 02
LIVER	1.35E-05	4.50E-01	1.46E 04	5.76E 01	1.46E 04	5.76E 01	1.46E 04	5.76E 01	1.46E 04	5.76E 01
TOTAL BODY	3.00E-05	1.00E 00	6.50E 04	5.76E 01	6.50E 04	5.76E 01	6.50E 04	5.76E 01	6.50E 04	5.76E 01
THYROID	0.	0.	0.	0.	0.	0.	0.	0.	0.	0.
KIDNEY	6.00E-07	2.00E-02	3.20E 04	5.76E 01	3.20E 04	5.76E 01	3.20E 04	5.76E 01	3.20E 04	5.76E 01
LUNG INGES	0.	0.	0.	0.	0.	0.	0.	0.	0.	0.
LUNG INHAL	0.	1.20E-01	1.20E 02	5.76E 01	1.20E 02	5.76E 01	1.20E 02	5.76E 01	1.20E 02	5.76E 01
GI-LLI INGES	1.00E 00	0.	0.	5.76E-01	0.	5.76E-01	0.	5.76E-01	0.	5.76E-01
GI-LLI INHAL	1.00E 00	6.20E-01	0.	5.76E-01	0.	5.76E-01	0.	5.76E-01	0.	5.76E-01



DISTRIBUTION

<u>No. of Copies</u>		<u>No. of Copies</u>
<u>OFFSITE</u>		<u>ONSITE</u>
	S. Acharya U.S. Nuclear Regulatory Commission Washington, DC 20555	<u>Pacific Northwest Laboratory</u> D.A. Baker (5) K.E. Byers G.R. Hoenes J. Mishima W.E. Kennedy B.A. Napier R.A. Peloquin L.C. Schwendiman J.K. Soldat D.L. Streng (25) B.E. Vaughan E.C. Watson
	E.F. Branagan U.S. Nuclear Regulatory Commission Washington, DC 20555	Technical Information (5) Publishing Coordination (2)
	T.R. Decker U.S. Nuclear Regulatory Commission Washington, DC 20555	
	F.D. Fisher U.S. Nuclear Regulatory Commission Washington, DC 20555	
	F. Swanberg U.S. Nuclear Regulatory Commission Washington, DC 20555	
2	DOE Technical Information Center	
205	U.S. Nuclear Regulatory Commission Division of Technical Information and Document Control 7920 Norfolk Avenue Bethesda, MD 20014	
	R.B. McPherson Battelle Memorial Institute 505 King Avenue Columbus, OH 43201	





<b>NRC FORM 335</b> (11-81)		<b>U.S. NUCLEAR REGULATORY COMMISSION</b> <b>BIBLIOGRAPHIC DATA SHEET</b>		<b>1. REPORT NUMBER (Assigned by DDC)</b> NUREG/CR-2384 PNL-3912	
<b>4. TITLE AND SUBTITLE (Add Volume No., if appropriate)</b> Age-Specific Inhalation Radiation Dose Commitment Factors for Selected Radionuclides				<b>2. (Leave blank)</b>	
<b>7. AUTHOR(S)</b> D.L. Streng , R. A. Peloquin, D. A. Baker				<b>3. RECIPIENT'S ACCESSION NO.</b>	
<b>9. PERFORMING ORGANIZATION NAME AND MAILING ADDRESS (Include Zip Code)</b> Pacific Northwest Laboratory Richland, Washington 99352				<b>5. DATE REPORT COMPLETED</b> MONTH   YEAR July   1982	
<b>12. SPONSORING ORGANIZATION NAME AND MAILING ADDRESS (Include Zip Code)</b> Division of Fuel Cycle and Material Safety Office of Nuclear Material Safety and Safeguards U. S. Nuclear Regulatory Commission Washington, D. C. 20555				<b>6. (Leave blank)</b>	
<b>13. TYPE OF REPORT</b> Topical Report				<b>7. (Leave blank)</b>	
<b>15. SUPPLEMENTARY NOTES</b>				<b>8. (Leave blank)</b>	
<b>16. ABSTRACT (200 words or less)</b> Inhalation dose commitment factors are presented for selected radionuclides for exposure of individuals in four age groups: infant, child, teen and adult. Radionuclides considered are <sup>35</sup> S, <sup>36</sup> Cl, <sup>45</sup> Ca, <sup>67</sup> Ga, <sup>75</sup> Se, <sup>85</sup> Sr, <sup>109</sup> Cd, <sup>113</sup> Sn, <sup>125</sup> I, <sup>133</sup> Ba, <sup>170</sup> Tm, <sup>169</sup> Yb, <sup>182</sup> Ta, <sup>192</sup> Ir, <sup>198</sup> Au, <sup>201</sup> Tl, <sup>204</sup> Tl, and <sup>236</sup> Pu. The calculational method is based on the human metabolic model of ICRP as defined in Publication 2 (ICRP 1959) and as used in previous age-specific dose factor calculations by Hoenes and Soldat (1977). Dose commitment factors are presented for the following organs of reference: total body, bone, liver, kidney, thyroid, lung and lower large intestine.				<b>9. (Leave blank)</b>	
<b>17. KEY WORDS AND DOCUMENT ANALYSIS</b> airborne radioactivity, inhalation exposure, dose		<b>17a. DESCRIPTORS</b>			
<b>17b. IDENTIFIERS/OPEN-ENDED TERMS</b>					
<b>18. AVAILABILITY STATEMENT</b> Unlimited		<b>19. SECURITY CLASS (This report)</b> Unclassified		<b>21. NO. OF PAGES</b>	
<b>20. SECURITY CLASS (This page)</b> Unclassified		<b>22. PRICE</b> S			

