

CATEGORY 1

REGULATORY INFORMATION DISTRIBUTION SYSTEM (RIDS)

ACCESSION NBR: 9605060364 DOC. DATE: 95/12/31 NOTARIZED: NO DOCKET #
 FACIL: 50-397 WPPSS Nuclear Project, Unit 2, Washington Public Powe 05000397
 AUTH. NAME AUTHOR AFFILIATION
 WEBRING, R.L. Washington Public Power Supply System
 RECIP. NAME RECIPIENT AFFILIATION

*See
 Enviro
 Reports*

ZELLER, J.J. Washington, State of

SUBJECT: "Operational Ecological Monitoring Program for Nuclear Plant
 2,1995 Annual Rept." W/960430 ltr.

DISTRIBUTION CODE: C001D COPIES RECEIVED: LTR 1 ENCL 1 SIZE: 50
 TITLE: Licensing Submittal: Environmental Rept Amdt & Related Correspondence

NOTES:

	RECIPIENT		COPIES		RECIPIENT		COPIES	
	ID	CODE/NAME	LTR	ENCL	ID	CODE/NAME	LTR	ENCL
	PD4-2	PD	1	1	CLIFFORD, J		1	1
INTERNAL:	ACRS		6	6	<u>FILE CENTER</u>	01	1	1
	NUDOCS-ABSTRACT		1	1	<u>OGC/HDS3</u>		1	0
	RGN4	DRSS/RPB	1	1				
EXTERNAL:	LITCO AKERS, D		1	1	NOAC		1	1
	NRC PDR		1	1				


NOTE TO ALL "RIDS" RECIPIENTS:
 PLEASE HELP US TO REDUCE WASTE! CONTACT THE DOCUMENT CONTROL DESK,
 ROOM OWEN 5D-5 (EXT. 415-2083) TO ELIMINATE YOUR NAME FROM
 DISTRIBUTION LISTS FOR DOCUMENTS YOU DON'T NEED!

TOTAL NUMBER OF COPIES REQUIRED: LTR 15 ENCL 14

C
A
T
E
G
O
R
Y

1

D
O
C
U
M
E
N
T



WASHINGTON PUBLIC POWER SUPPLY SYSTEM

P.O. Box 968 • 3000 George Washington Way • Richland, Washington 99352-0968 • (509) 372-5000

April 30, 1996
GO2-96-093

Mr. Jason J. Zeller, Manager
Energy Facility Site Evaluation Council
P.O. Box 43172
Olympia, WA 98501-3172

Dear Mr. Zeller:

Subject: **SUPPLY SYSTEM NUCLEAR PLANT NO. 2
ECOLOGICAL MONITORING PROGRAM ANNUAL REPORT FOR 1995**

Reference: Letter dated March 27, 1996, JV Parrish (Supply System) to JJ Zeller (EFSEC),
"Nonradiological Environmental Monitoring Program"

Enclosed, please find five (5) copies of the subject report which is submitted per Council Resolution No. 266. In the referenced letter the Supply System proposed to discontinue the monitoring activities because ten years of data have not disclosed any environmental impact attributable to plant operation. If you have questions concerning the report or the monitoring program, please contact W.A. Kiel at (509) 377-4490.

Sincerely,




R.L. Webring (Mail Drop PE08)
Vice President, Operations Support/PIO

Enclosures

cc (w/encl):

L Russell (WDOE-Kenn)
RL Dirkes (PNNL)
JW Clifford (NRC NRR)
LJ Callan (NRC RIV)
NRC Document Control Desk (Docket No. 50-397)

9605060364 951231
PDR ADOCK 05000397
R PDR



Cool
11

MARCH 7, 1995

DOCKET NO(S). 50-397, STN 50-528,
STN 50-529, AND STN 50-530

SEE ATTACHED LIST

SUBJECT: SEMIANNUAL RADIOACTIVE EFFLUENT RELEASE REPORT FOR PALO VERDE NUCLEAR
GENERATING STATION AND WPPSS NUCLEAR PROJECT, UNIT 2

The following documents concerning our review of the subject facility are transmitted for your information.

✓	DESCRIPTION OF DOCUMENT	DATED
	Notice of Receipt of Application	
	Draft/Final Environmental Statement	
	Notice of Availability of Draft/Final Environmental Statement	
	Safety Evaluation Report, or Supplement No. _____	
	Environmental Assessment and Finding of No Significant Impact	
	Notice of Issuance of Environmental Assessment	
	Notice of Consideration of Issuance of Facility Operating License or Amendment to Facility Operating License	
	Biweekly Notice; Applications and Amendments to Operating Licenses Involving No Significant Hazards Conditions See Page(s) _____	
	Exemption	
	Construction Permit No. CPPR- _____, Amendment No. _____	
	Facility Operating License No. _____, Amendment No. _____	
	Order	
	Monthly Operating Report for _____ transmitted by Letter	
	Annual/Semi-Annual Report: UNP-2 CORRECTION TO REPORT	9/22/94
	PALO VERDE 1, 2, 3 EFFLUENT RELEASE REPORT FOR JAN-JUNE transmitted by Letter	8/31/94
	Other _____	

Office of Nuclear Reactor Regulation

PROJECT DIRECTORATE IV-2

Enclosures:
As Stated

cc:

OFFICE▶	LA:DRPW				
SURNAME▶	FOSTER-CURSEEN				
DATE▶	3/7/95				



10-10-68

10-10-68

10-10-68

10-10-68

July 29, 1994

DOCKET NO(S): 50-397

Mr. Jerry Leitch
 Radiation Program Manager, Region 10
 Environmental Protection Agency
 1200 E 15th Avenue
 Seattle, WA 98101

SUBJECT: WNP-2 Operational Ecological Monitoring Program Annual Report
 and Radiological Environ Monitoring Program Annual Report

The following documents concerning our review of the subject facility are transmitted for your information.

	DESCRIPTION OF DOCUMENT	DATED
✓	Notice of Receipt of Application	
	Draft/Final Environmental Statement	
	Notice of Availability of Draft/Final Environmental Statement	
	Safety Evaluation Report, or Supplement No. _____	
	Environmental Assessment and Finding of No Significant Impact	
	Notice of Issuance of Environmental Assessment	
	Notice of Consideration of Issuance of Facility Operating License or Amendment to Facility Operating License	
	Biweekly Notice; Applications and Amendments to Operating Licenses Involving No Significant Hazards Conditions See Page(s) _____	
	Exemption	
	Construction Permit No. CPPR- _____, Amendment No. _____	
	Facility Operating License No. _____, Amendment No. _____	
	Order	
	Monthly Operating Report for _____ transmitted by Letter	
X	Annual/Semi-Annual Report: Operational Ecological Monitoring Program	4/29/94
	Annual Radiological Monitoring Rept. _____ transmitted by Letter	4/28/94
	Other _____	

Office of Nuclear Reactor Regulation
 Project Directorate IV-2

Enclosures:
 As Stated

cc:

OFFICE▶	...LA:DRP <i>gjc</i>						
SURNAME▶	DFoster-Curseen						
DATE▶	7/29/94						



March 24, 1994

DOCKET NO(S). 50-397

Dr. William C. ^{unn}Cyraningham
 FDA Research Chemist
 NIST Building 235/B125
 Gaithersburg, MD 20899

SUBJECT: WASHINGTON PUBLIC POWER SUPPLY SYSTEM

The following documents concerning our review of the subject facility are transmitted for your information.

✓	DESCRIPTION OF DOCUMENT	DATED
	Notice of Receipt of Application	
	Draft/Final Environmental Statement	
	Notice of Availability of Draft/Final Environmental Statement	
	Safety Evaluation Report, or Supplement No. _____	
	Environmental Assessment and Finding of No Significant Impact	
	Notice of Issuance of Environmental Assessment	
	Notice of Consideration of Issuance of Facility Operating License or Amendment to Facility Operating License	
	Biweekly Notice; Applications and Amendments to Operating Licenses Involving No Significant Hazards Conditions See Page(s) _____	
	Exemption	
	Construction Permit No. CPPR- _____, Amendment No. _____	
	Facility Operating License No. _____, Amendment No. _____	
	Order	
	Monthly Operating Report for _____ transmitted by Letter	
	Annual/Semi-Annual Report: _____ transmitted by Letter	
XX	Other ANNUAL ENVIRONMENTAL OPERATING REPORT	2/25/94

Office of Nuclear Reactor Regulation


Project Directorate V

Enclosures:
 As Stated

cc:

ACKS-1

OFFICE▶	LA:PDV						
SURNAME▶	DFoster						
DATE▶	3/24/94						



March 24, 1994

DOCKET NO(S). 50-397

Chief, Branch of Federal Activities
 Division of Habitat Conservation
 400 ARLSQ
 U.S. Fish and Wildlife Service
 1849 C Street, N.W.
 Washington, D.C. 20240

SUBJECT: WASHINGTON PUBLIC POWER SUPPLY SYSTEM

The following documents concerning our review of the subject facility are transmitted for your information.

	DESCRIPTION OF DOCUMENT	DATED
	Notice of Receipt of Application	
	Draft/Final Environmental Statement	
	Notice of Availability of Draft/Final Environmental Statement	
	Safety Evaluation Report; or Supplement No. _____	
	Environmental Assessment and Finding of No Significant Impact	
	Notice of Issuance of Environmental Assessment	
	Notice of Consideration of Issuance of Facility Operating License or Amendment to Facility Operating License	
	Biweekly Notice; Applications and Amendments to Operating Licenses Involving No Significant Hazards Conditions See Page(s) _____	
	Exemption	
	Construction Permit No: CPPR- _____, Amendment No. _____	
	Facility Operating License No. _____, Amendment No. _____	
	Order	
X	ANNUAL Operating Report for <u>WPPSS</u> transmitted by Letter	2/25/94
	Annual/Semi-Annual Report: _____ transmitted by Letter	
XX	Other <u>RADIOACTIVE EFFLUENT RELEASE RET JAN-DEC 1993 & ODCM</u>	2/28/94

Office of Nuclear Reactor Regulation
 PROJECT DIRECTORATE V

Enclosures:
 As Stated

cc:

ACRS-1

OFFICE▶	LA:PDV <i>dfc</i>					
SURNAME▶	DFoster- Curseen					
DATE▶	3/24/94					



2014

1

2014

2

2014

3

4

5

Distribution

✓ Docket File
PDV R/F

~~DFoster-Curseen~~

March 23, 1994

DOCKET NO(S). 50-397

Mr. Jerry Leitch
Radiation Program Manager, Region 9
(A1-002)
Environmental Protection Agency
1200 Sixth Avenue
Seattle, WA 98101

SUBJECT: TRANSMITTAL OF ANNUAL REPORTS FOR WPPSS NUCLEAR PROJECT, UNIT 2

The following documents concerning our review of the subject facility are transmitted for your information.

✓	DESCRIPTION OF DOCUMENT	DATED
	Notice of Receipt of Application	
	Draft/Final Environmental Statement	
	Notice of Availability of Draft/Final Environmental Statement	
	Safety Evaluation Report, or Supplement No. _____	
	Environmental Assessment and Finding of No Significant Impact	
	Notice of Issuance of Environmental Assessment	
	Notice of Consideration of Issuance of Facility Operating License or Amendment to Facility Operating License	
	Biweekly Notice; Applications and Amendments to Operating Licenses Involving No Significant Hazards Conditions See Page(s) _____	
	Exemption	
	Construction Permit No. CPPR- _____, Amendment No. _____	
	Facility Operating License No. _____, Amendment No. _____	
	Order	
	Operating Report for WPPSS _____ transmitted by Letter	2/25/94
	Annual/Semi-Annual Report: UNIT 2 RADIOACTIVE EFFLUENT RELEASE... _____ transmitted by Letter	2/28/94
	Other _____	

Office of Nuclear Reactor Regulation
Project Directorate V

Enclosures:
As Stated

cc:

ACRS-1

OFFICE▶	LA:PDV					
SURNAME▶	D. Foster-Curseen					
DATE▶	3/23/94					



1950


20

1950

1950

OPERATIONAL ECOLOGICAL MONITORING PROGRAM FOR NUCLEAR PLANT 2

WASHINGTON PUBLIC POWER SUPPLY SYSTEM



1995 ANNUAL REPORT

PREPARED BY ENVIRONMENTAL SCIENCES DEPARTMENT

9605060364





TABLE OF CONTENTS

<u>Section</u>	<u>Page</u>
EXECUTIVE SUMMARY	i
ACKNOWLEDGEMENTS	ii
TABLES	iii
FIGURES	v
1.0 INTRODUCTION	1-1
1.1 BACKGROUND	1-1
1.2 THE SITE	1-1
2.0 WATER QUALITY	2-1
2.1 INTRODUCTION	2-1
2.2 MATERIALS AND METHODS	2-2
2.2.1 Sample Collection	2-3
2.2.2 Analysis Methods	2-4
2.3 RESULTS AND DISCUSSION	2-6
2.2.1 Temperature	2-6
2.2.2 Dissolved Oxygen	2-7
2.2.3 pH and Alkalinity	2-8
2.2.4 Hardness	2-9
2.2.5 Conductivity	2-10
2.2.6 Turbidity	2-10
2.2.7 Metals (Total)	2-12
2.2.8 Oil and Grease	2-16
2.2.9 Total Phosphorus and Inorganic Phosphate	2-16
2.2.10 Sulfate	2-18
2.2.11 Total Dissolved Solids	2-18
2.2.12 VOCs and Semi-VOCs	2-19

3.0 SOIL AND VEGETATION STUDIES	3-1
3.1 INTRODUCTION	3-1
3.2 MATERIALS AND METHODS	3-3
3.2.1 Herbaceous Canopy Cover	3-3
3.2.2 Herbaceous Phytomass	3-3
3.2.3 Soil Chemistry	3-3
3.3 RESULTS AND DISCUSSION	3-4
3.2.1 Herbaceous Cover	3-7
3.2.2 Herbaceous Phytomass	3-12
3.2.3 Soil Chemistry	3-18
4.0 REFERENCES	4-1



EXECUTIVE SUMMARY

The Ecological Monitoring Program is comprised of several elements which are intended to determine the effects of the operation of the Supply System's Nuclear Plant No. 2 on the environment. These program elements include: plant effluent and Columbia River water quality; vegetation cover and phytomass in selected plots; and soil chemistry at established sampling locations. The results of the 1995 monitoring efforts may be summarized as follows:

- Plant cooling water discharges had no discernible effect on Columbia River water quality.
- No measurable effects of plant cooling tower drift were observed on vegetation cover, phytomass, or soil chemistry. A dramatic increase in cover and phytomass was observed at most stations. This is in direct correlation with the record precipitation recorded during the growing season.

ACKNOWLEDGEMENTS

This report, prepared by the Washington Public Power Supply System, describes the soil and vegetation studies, and water quality programs for WNP-2.

Project Team

Terry E. Northstrom

Supervisor, Environmental Sciences

Deborah C. Singleton

Environmental Scientist

Richard E. Welch

Environmental Scientist

LIST OF TABLES

<u>Number</u>	<u>Title</u>	<u>Page</u>
2-1.	Summary of Water Quality Parameters, Stations, and Sampling Frequencies, 1995	2-2
2-2.	Summary of Water Quality Parameters, EPA and Standard Methods Numbers	2-4
2-3	Summary of Temperature ($^{\circ}\text{C}$) Measurements	2-6
2-4.	Summary of Dissolved Oxygen (mg/L) Measurements	2-6
2-5.	Summary of pH Measurements	2-7
2-6.	Summary of Alkalinity (mg/L as CaCO_3) Measurements	2-8
2-7.	Summary of Total Hardness (mg/L as CaCO_3) Measurements	2-8
2-8.	Summary of Conductivity ($\mu\text{S/cm}$) Measurements	2-9
2-9.	Summary of Turbidity (NTU) Measurements	2-10
2-10.	Summary of Copper ($\mu\text{g/L}$) Measurements	2-11
2-11.	Summary of Nickel ($\mu\text{g/L}$) Measurements	2-12
2-12.	Summary of Zinc ($\mu\text{g/L}$) Measurements	2-12
2-13.	Summary of Iron ($\mu\text{g/L}$) Measurements	2-13
2-14.	Summary of Lead ($\mu\text{g/L}$) Measurements	2-13
2-15.	Summary of Cadmium ($\mu\text{g/L}$) Measurements	2-14
2-16.	Summary of Chromium ($\mu\text{g/L}$) Measurements	2-14
2-17.	Summary of Oil and Grease (mg/L) Measurements	2-15
2-18.	Summary of Total Phosphorus (mg/L as P) Measurements	2-16
2-19.	Summary of Inorganic Phosphate (mg/L as P) Measurements	2-16



LIST OF TABLES (Continued)

<u>Number</u>	<u>Title</u>	<u>Page</u>
2-20.	Summary of Sulfate (mg/L) Measurements	2-17
2-21.	Summary of Quarterly Total Dissolved Solid (mg/L) Measurements	2-17
2-22.	Summary of Volatile Organic Compound Measurements	2-18
2-23.	Summary of Semivolatile Organic Compound Measurements	2-18
3-1.	Vascular Plants Observed During 1995	3-4
3-2.	Herbaceous Cover for Fifteen Sampling Stations - 1995	3-8
3-3.	Mean Frequency Values (%) by Species for Each Sampling Station	3-9
3-4.	Mean Herbaceous Cover for 1975 through 1995	3-10
3-5.	Herbaceous Phytomass Measurements for 1995	3-14
3-6.	Comparison of Herbaceous Phytomass (g/m ²) for 1975 through 1995	3-15
3-7.	Summary of Soil Chemistry for 1995	3-16

LIST OF FIGURES

<u>Number</u>	<u>Title</u>	<u>Page</u>
1-1.	WNP-2 Location Map	1-2
2-1.	Location of Sampling Stations	2-1
2-2.	Schematic of River Sample Locations for Water Quality	2-5
3-1.	Soil and Vegetation Sampling Location Map	3-2
3-2.	Layout of Vegetation and Soil Sampling Plots	3-3.
3-3.	Mean Herbaceous Cover for 1975 through 1995.	3-13
3-4.	Mean Herbaceous Cover and Total Precipitation	3-13
3-5.	Phytomass at Grassland and Shrub Stations for 1975 through 1995	3-15

1.0 INTRODUCTION

1.1 BACKGROUND

The Site Certification Agreement (SCA) for WNP-2 was approved on May 17, 1972, by the State of Washington and the Washington Public Power Supply System (Supply System). The SCA requires that environmental monitoring be conducted during the preoperational and operational phases of site development and use. The objective of the monitoring program is to provide an environmental measurement history for evaluation by the Supply System and the Washington State Energy Facility Site Evaluation Council (EFSEC) and to identify significant effects of plant operation on the environment. Since 1972, several revisions of the monitoring program have been approved by EFSEC in the form of SCA attachments and EFSEC resolutions Nos. 193, 194, 214, 239, and 266.

Most of the studies, analyses, and reports for the preoperational (1973-1984) environmental program of the SCA were performed by outside laboratories for the Supply System. The aquatic studies were in reports by Battelle Pacific Northwest Laboratories for the period of September 1974 through August 1978 (Battelle 1976, 1977, 1978, 1979a, 1979b) and by Beak Consultants, Inc. for the period of August 1978 through March 1980 (Beak 1980). The terrestrial program was performed and reports were prepared by Battelle from 1974 until 1979 (Rickard 1976, 1977, 1979a, 1979b) and then by Beak from 1980 to 1982 (Beak 1981, 1982a, 1982b).


Since 1983, Supply System scientists have been responsible for the entire operational environmental monitoring program. Using the data acquired during 1984, the first comprehensive operational environmental annual report was prepared by Supply System scientists (Supply System 1985) and has since continued annually (Supply System 1986 through 1994). A few studies and reports were completed by Supply System personnel prior to the annual reports, including animal studies (Schleder 1982, 1983, 1984) and terrestrial monitoring (Northstrom 1984).

This report presents the results of the Ecological Monitoring Program for the period of January through December 1995.

1.2 THE SITE

The Supply System leases the WNP-2 site (441 hectares or 1089 acres) from the U. S. Department of Energy. WNP-2 lies within the boundaries of the Columbia Basin between the Cascade Range in Washington and Blue Mountains in Oregon and comprises approximately two-thirds of the area lying east of the Cascades. Approximately 5 km (3.25 miles) to the east, the site is bounded by the Columbia River. The plant communities within the region are described as shrub-steppe communities consisting of various layers of perennial grasses overlaid by a discontinuous layer of shrubs. In general, moisture relations do not support arborescent species, except along streambanks. In August 1984, a range fire destroyed much of the shrub cover on the Hanford site and temporarily modified the shrub-steppe associations which were formerly present.

Figure 1-1 WNP-2 Location Map



2.0 WATER QUALITY

2.1 INTRODUCTION


The water quality sampling stations are located near the west bank of the Columbia River at river mile 352. Sampling was limited to the main channel on the Benton County side. Near the site, the river averages 370 meters (1200 feet) wide with a water surface elevation of 105 meters (345 feet) above sea level and ranges to 7.3 meters (24 feet) deep. Sampling stations have been established in the river both upstream and downstream from the plant intake and discharge structures. The river level in this area fluctuates considerably during a 24-hour period and from day to day in response to release patterns at the Priest Rapids Dam (river mile 397).

2.2 MATERIALS AND METHODS

Columbia River Surface water was sampled monthly from January through December 1995. Samples were collected near river mile 352 from four stations numbered 1, 7, 11, and 8 (Figures 2-1 and 2-2). Station 1 is upstream of the WNP-2 intake and discharge and represents the control. Station 7 is in the center of the mixing zone approximately 45 meters (150 feet) downstream of the discharge and provides a measure of near field blowdown effects. Station 11, at 91 meters (300 feet) downstream from the discharge, represents the extremity of the mixing zone. Substations 11M and 11B sample water from middle and bottom depths, respectively. Station 8 is approximately 568 meters (1870 feet) downstream from the discharge and represents a location where the blowdown is well mixed in the Columbia River. With the exception of Substations 11M and 11B, Columbia River samples were analyzed for temperature, dissolved oxygen (DO), pH, conductivity, turbidity, total alkalinity, total hardness, total phosphorus, inorganic phosphate, sulfate, total copper, total iron, total zinc, total nickel, total lead, total cadmium and total chromium. The samples from substations (11M and 11B) were analyzed for total copper only.

Plant blowdown was sampled monthly during 1995. Blowdown samples were analyzed for temperature, pH, conductivity, turbidity, total phosphorus, inorganic phosphate, sulfate, oil and grease, total copper, total iron, total zinc, total nickel, total lead, total cadmium, and total chromium. Volatile organic compounds (VOCs) and semi-volatile organic compounds were analyzed on a quarterly basis.

Figure 2-1. Location of Sampling Stations



The evaporation/percolation pond (storm drain pond) is located approximately 1500 feet northeast of the plant. The pond is a collection point for water from various locations within the controlled area. Water and sediment were sampled monthly and semiannually, respectively. Monthly water samples were analyzed for pH, conductivity, total iron, total copper, total nickel, total zinc, total lead, total cadmium, total chromium, and oil and grease. In addition, quarterly water samples were analyzed for total dissolved solids and VOCs and semi-VOCs. Semiannual sediment samples were analyzed for the same total metals as the monthly water samples, excluding iron. A summary of water quality parameters, stations and sampling frequencies is presented in Table 2-1.

Table 2-1. Summary of Water Quality Parameters, Stations, and Sampling Frequencies, 1995

Parameter	I	7**	11**	11M & 11B**	8**	Plant Blowdown	Pond
Temperature	M	M	M	—	M	M	—
Dissolved Oxygen	M	M	M	—	M	—	—
pH	M	M	M	—	M	M	M
Turbidity	M	M	M	—	M	M	—
Total Alkalinity	M	M	M	—	M	—	—
Filtrable Residue (TDS)	—	—	—	—	—	—	Q
Conductivity	M	M	M	—	—	M	M
Iron (Total)	M	M	M	—	M	M	M
Copper (Total)	M	M	M	M	M	M	M
Nickel (Total)	M	M	M	—	M	M	M
Zinc (Total)	M	M	M	—	M	M	M
Lead (Total)	M	M	M	—	M	M	M
Cadmium (Total)	M	M	M	—	M	M	M
Chromium (Total)	M	M	M	—	M	M	M
Sulfate	M	M	M	—	M	M	—
Orthophosphorus	M	M	M	—	M	M	—
Total Phosphorus	M	M	M	—	M	M	—
Oil and Grease	—	—	—	—	M	M	M
Hardness	M	M	M	—	M	—	—
Organics (VOCs and semi-Vocs)	—	—	—	—	—	Q	Q

Symbols Key

Q= Quarterly

M= Monthly

**= Samples collected only if the plant is operating

2.2.1 Sample Collection

Columbia River water samples were collected by boat approximately 300 feet from the Benton County shore. Temperature was determined in situ with portable instruments. Water for total metal, conductivity, pH, sulfate, total phosphorus, inorganic phosphate, turbidity, total alkalinity and total hardness analyses was collected in 2.8 liter polypropylene cubitainers and stored in a cooler until delivered to the Supply System's Environmental and Analytical Support Laboratory (EASL). Water for total copper analysis from substations 11M and 11B was collected in one-liter polypropylene cubitainers with an all-Teflon pump and Tygon tubing. Water for dissolved oxygen measurements was collected in 300 ml (Biological Oxygen Demand) bottles.

Blowdown temperature was determined in situ. Water for pH, conductivity, turbidity, total phosphorus, inorganic phosphate and total metals analysis was collected in 2.8 liter polypropylene cubitainers. Water for oil and grease and semivolatile organics analysis was collected in one-liter clear and amber glass bottles, respectively. Water for volatile organics analysis was collected in 40 ml glass bottles.

Evaporation/percolation pond water for pH, conductivity and total metals was collected in 2.8 liter polypropylene cubitainers. Water for total dissolved solids analysis was collected in 500 ml plastic bottles. Water for oil and grease, VOCs and semi-VOCs was collected as described under blowdown sampling. All samples were stored in a cooler until delivered to the laboratory for analysis.

River water quality samples collected during the annual plant maintenance outage (April through June) consisted of station 1 (control) samples only.

2.2.2 Analysis Methods

Field temperature measurements were made using a Fisher NIST-traceable thermometer. Temperature was recorded to within 0.1°C after the probe had been allowed to equilibrate for a minimum of one minute.

Total metals, sulfate, conductivity, pH, dissolved oxygen, inorganic phosphate, turbidity, total alkalinity, total hardness, VOCs and semi-VOCs, total phosphorus, and oil and grease, were determined by Supply System laboratory personnel. Analyses for total dissolved solids and some total metals were performed by an offsite laboratory. Sample holding times followed those recommended by the U.S. Environmental Protection Agency (EPA 1983). Table 2-2 lists the approved EPA and Standard Methods used.




Table 2-2. Summary of Water Quality Parameters, EPA and Standard Method Method Numbers

Parameter	EPA Method Number	Standard Methods Method Number
Water Temperature (°C)	170.1	
Turbidity (NTU)	180.1	
Conductivity ($\mu\text{S}/\text{cm}$) at 25°C	120.1	
Dissolved Oxygen (mg/L) Probe	360.1	
Dissolved Oxygen (mg/L) Modified Winkler	360.2	
pH (Standard Unit)	150.1	
Total Alkalinity (mg/L as CaCO_3)	310.1	
Total Hardness (mg/L as CaCO_3)	130.2, 6010	2340B
Oil and Grease (mg/L)	412.2	
Total Phosphorus (mg/L as P)	365.2	4500-P
Inorganic Phosphate (mg/L as P)	300, 365.2	
Sulfate (mg/L as SO_4)	300, 375.4	
Total Copper ($\mu\text{g}/\text{L}$ as Cu)	200.7, 220.1, 220.2	
Total Iron ($\mu\text{g}/\text{L}$ as Fe)	200.7, 236.1, 236.2	
Total Nickel ($\mu\text{g}/\text{L}$ as Ni)	200.7, 249.1, 249.2	
Total Zinc ($\mu\text{g}/\text{L}$ as Zn)	200.7, 289.1, 289.2	
Total Lead ($\mu\text{g}/\text{L}$ as Pb)	200.7, 239.1, 239.2	
Total Cadmium ($\mu\text{g}/\text{L}$ as Cd)	200.7, 212.1, 212.2	
Total Chromium ($\mu\text{g}/\text{L}$ as Cr)	200.7, 218.1, 218.2	
Filterable Residue: TDS (mg/L)	160.1	
Volatile Organics ($\mu\text{g}/\text{L}$)	8240	
Semivolatile Organics ($\mu\text{g}/\text{L}$)	8270	



Figure 2-2. Schematic of River Sample Locations for Water Quality



2.3 RESULTS AND DISCUSSION

The evaporation/percolation pond is a discharge to ground and is not related to potential effects of the blowdown on the Columbia River. There was no plant blowdown during the February sampling event. For all sampling periods, significant interstation differences could not be detected for any of the measured parameters. The water quality monitoring results for both the river and the pond are presented in the following subsections.

2.3.1 Temperature

Columbia River surface temperatures varied seasonally with a temperature of 3.2°C at all stations on January 30 and a maximum of 19.8°C at station 1 on September 19 (Table 2-3). Blowdown temperatures ranged from 19.8°C in January to 41.7°C in July.

Table 2-3. Summary of Temperature (°C) Measurements for 1995

Sample Date	1	7	11	8	Plant Blowdown
01/30/95	3.2	3.2	3.2	3.2	19.8
02/28/95	3.5	3.5	3.5	3.5	--
03/30/95	5.5	5.5	5.5	5.5	37.1*
04/26/95	8.6	--	--	--	--
05/30/95	13.1	--	--	--	--
06/29/95	16.0	--	--	--	--
07/25/95	18.5	18.5	18.5	18.5	41.7*
08/31/95	19.0	19.0	19.0	19.0	40.6*
09/19/95	19.8	19.7	19.7	19.7	39.1*
10/24/95	13.8	13.8	13.8	13.8	35.5*
11/09/95	11.7	11.7	11.6	11.7	34.0*
12/20/95	6.8	6.8	6.8	6.8	24.2*

*Measurements taken from cooling tower 2B-top deck.

2.3.2 Dissolved Oxygen (DO)

DO measurements for each sample station are presented in Table 2-4. Columbia River DO concentrations ranged from 9.4 mg/L at Stations 1, 7 and 11 in September to 14.0 mg/L at Station 1 in February.

Table 2-4. Summary of Dissolved Oxygen (mg/L) Measurements

Sample Date	1	7	11	8
01/30/95	13.6	13.6	13.5	13.5
02/28/95	14.0	13.9	13.9	13.9
03/30/95	13.4	13.4	13.5	13.4
04/26/95	13.5	--	--	--
05/30/95	11.9	--	--	--
06/29/95	10.8	--	--	--
07/25/95	10.1	10.2	10.2	10.2
08/31/95	10.0	10.0	9.9	10.0
09/19/95	9.4	9.4	9.4	9.5
10/24/95	10.2	10.5	10.1	10.3
11/09/95	11.0	10.8	10.9	10.8
12/20/95	12.1	12.1	12.0	12.2

DO concentrations were inversely related to river temperature as would be expected from solubility laws. DO levels were never below the water quality standard for Class A waters (WDOE 1992) indicating good water quality with respect to dissolved oxygen throughout the year.

2.3.3 pH and Alkalinity

Columbia River pH values ranged from 7.47 at Station 1 in January to 8.12 at Station 1 in May. The pH water quality standard for Class A waters is from 6.5 to 8.5 (WDOE 1992). Blowdown pH values ranged from 7.86 in December, to 8.47 in September. Pond pH values ranged from 7.09 in January to 9.19 in April. Columbia River alkalinities ranged from 49 to 62 mg/L as calcium carbonate. Results for pH and alkalinity are listed in Tables 2-5 and 2-6.

Table 2-5. Summary of pH Measurements

Sample Date	1	7	11	8	Plant Blowdown	Pond
01/30/95	7.47	7.51	7.48	7.48	8.11	7.09
02/22/95	7.57	7.53	7.59	7.65	-	7.92
03/30/95	7.48	7.52	7.54	7.53	8.15	8.12
04/26/95	8.01	-	-	-	-	9.19
05/30/95	8.12	-	-	-	-	7.81
06/27/95	7.94	-	-	-	-	8.09
07/25/95	7.83	7.76	7.81	7.82	8.32	7.92
08/31/95	7.94	7.78	8.07	7.91	8.46	8.09
09/19/95	7.87	7.92	7.97	7.85	8.47	8.07
10/24/95	7.62	7.71	7.63	7.73	8.21	8.16
11/07/95	7.64	7.64	7.65	7.64	8.37	8.04
12/20/95	7.71	7.65	7.87	7.62	7.86	8.13




Table 2-6. Summary of Alkalinity Measurements

Sample Date	1	7	11	8
01/30/95	61	62	62	62
02/28/95	57	60	58	57
03/30/95	55	55	56	55
04/26/95	57	--	--	--
05/30/95	57	--	--	--
06/29/95	49	--	--	--
07/25/95	54	55	55	55
08/31/95	57	58	58	58
09/19/95	57	57	56	57
10/24/95	54	54	54	54
11/09/95	56	56	55	56
12/20/95	61	60	61	61

2.3.4 Hardness

Hardness ranged from 55 to 85 mg/L as calcium carbonate. This data is presented in Table 2-7.

Table 2-7. Summary of Total Hardness Measurements

Sample Date	1	7	11	8
01/30/95	72	74	72	73
02/28/95	72	73	73	72
03/30/95	71	72	71	71
04/26/95	85	--	--	--
05/30/95	58	--	--	--
06/29/95	55	--	--	--
07/25/95	59	60	61	60
08/31/95	66	65	66	67
09/19/95	65	64	63	65
10/24/95	63	64	63	63
11/09/95	67	69	67	66
12/20/95	71	70	69	71

2.3.5 Conductivity

Columbia River conductivity measurements ranged from 112 $\mu\text{S}/\text{cm}$ at 25°C at station 1 in June to 153 $\mu\text{S}/\text{cm}$ at 25°C at station 7 in January and February. Blowdown measurements ranged from 1060 $\mu\text{S}/\text{cm}$ at 25°C to 1570 $\mu\text{S}/\text{cm}$ at 25°C. Storm drain pond values ranged from 68 to 575 $\mu\text{S}/\text{cm}$ at 25°C. Conductivity measurements are listed in Table 2-8.

Table 2-8. Summary of Conductivity Measurements

Sample Date	1	7	11	8	Plant Blowdown	Pond
01/30/95	151	153	151	152	1140	68
02/28/95	150	153	148	148	--	575
03/30/95	149	150	149	149	1270	431
04/26/95	146	--	--	--	--	326
05/30/95	148	--	--	--	--	273
06/29/95	112	--	--	--	--	198
07/25/95	128	127	127	130	1480	223
08/31/95	138	140	140	140	1550	226
09/19/95	138	138	138	139	1570	305
10/24/95	125	124	131	133	1340	455
11/09/95	127	129	128	129	1320	433
12/20/95	146	147	147	147	1060	421

2.2.7 Turbidity

In the Columbia River, measured turbidities were low and ranged from 1.0 nephelometric turbidity units (NTU) to 4.5 NTU. Blowdown values ranged from 9 to 40 NTU. Turbidity results are listed in Table 2-9.

Table 2-9. Summary of Turbidity Measurements

Sample Date	1	7	11	8	Plant Blowdown
01/30/95	1.5	1.1	1.4	1.0	10
02/28/95	4.0	4.3	4.3	4.2	--
03/30/95	3.6	3.8	3.5	3.6	40
04/26/95	4.5	--	--	--	--
05/30/95	4.5	--	--	--	--
06/29/95	1.0	--	--	--	--
07/25/95	2.0	2.0	1.9	1.8	9
08/31/95	1.3	1.5	1.4	1.6	12
09/19/95	1.6	1.6	1.7	1.5	16
10/24/95	2.0	1.9	1.9	2.1	15
11/09/95	1.4	1.5	1.5	1.3	25
12/20/95	2.3	2.4	2.2	2.4	38



2.3.7 Metals (Total)

Columbia River cadmium concentrations were below the respective method detection limit ($1.4\mu\text{g/L}$) at all stations during all periods. River copper concentrations ranged from $<1.9\mu\text{g/L}$ to $4.6\mu\text{g/L}$. Zinc concentrations ranged from $<5.0\mu\text{g/L}$ to $14.4\mu\text{g/L}$ and iron concentrations ranged from $33\mu\text{g/L}$ to $218\mu\text{g/L}$. Nickel concentrations were generally below the detection limit of $2.0\mu\text{g/L}$. The highest nickel reading of $2.6\mu\text{g/L}$ was recorded at station 7 in January.

Blowdown cadmium concentrations were below the detection limit for all stations and periods, except October ($5.1\mu\text{g/L}$). Nickel and lead concentrations were fairly low, ranging from $<2.0\mu\text{g/L}$ to $7.0\mu\text{g/L}$ and $<1.0\mu\text{g/L}$ to $4.7\mu\text{g/L}$, respectively. Blowdown copper, zinc and iron concentrations were substantially higher than river concentrations and ranged from $45\mu\text{g/L}$ to $110\mu\text{g/L}$, $47\mu\text{g/L}$ to $108\mu\text{g/L}$, and $360\mu\text{g/L}$ to $2010\mu\text{g/L}$, respectively. Chromium concentrations ranged from $<0.5\mu\text{g/L}$ to $5.0\mu\text{g/L}$.

Evaporation/percolation pond water cadmium and nickel concentrations were below their respective detection limits for all periods. Lead concentrations ranged from $<1.0\mu\text{g/L}$ to $19.0\mu\text{g/L}$. Chromium concentrations ranged from $<0.5\mu\text{g/L}$ to $1.6\mu\text{g/L}$. Copper concentrations ranged from $<1.9\mu\text{g/L}$ to $35\mu\text{g/L}$ and zinc concentrations ranged from $27\mu\text{g/L}$ to $774\mu\text{g/L}$. Iron concentrations ranged from a low of $20\mu\text{g/L}$ in November to a high of $196\mu\text{g/L}$ in January. With the exception of lead and nickel, measurable levels for all other metal constituents were observed in the storm drain pond sediment samples.

Total metal results are listed in Tables 2-10 through 2-16.

Table 2-10. Summary of Copper ($\mu\text{g/L}$) Measurements

Sample Date	1	7	11	11M	11B	8	Plant Blowdown	Pond	Pond Sediment ($\mu\text{g/g}$)
01/30/95	<1.9	2.1	<1.9	<1.9	<1.9	<1.9	54	8.5	--
02/28/95	<1.9	2.4	2.4	<1.9	<1.9	2.1	--	2.4	--
03/30/95	<1.9	2.0	<1.9	<1.9	<1.9	<1.9	51	<1.9	--
04/26/95	<1.9	--	--	--	--	--	--	35	--
05/30/95	2.6	--	--	--	--	--	--	17	--
06/29/95	2.9	--	--	--	--	--	--	13	71.4
07/25/95	2.0	3.0	2.0	4.6	2.4	2.0	110	6.0	--
08/31/95	<1.9	2.0	2.0	<1.9	<1.9	2.0	58	5.0	--
09/19/95	2.0	<1.9	<1.9	<1.9	<1.9	<1.9	45	8.0	--
10/24/95	<1.9	<1.9	<1.9	<1.9	<1.9	<1.9	54	<1.9	--
11/09/95	<1.9	<1.9	<1.9	<1.9	<1.9	<1.9	69	3.0	--
12/20/95	<1.9	<1.9	<1.9	<1.9	<1.9	<1.9	57	2.0	101

Table 2-11. Summary of Nickel ($\mu\text{g/L}$) Measurements

Sample Date	1	7	11	8	Plant Blowdown	Pond	Pond Sediment ($\mu\text{g/g}$)
01/30/95	<2.0	2.6	<2.0	<2.0	4.2	<2.0	—
02/28/95	<2.0	<2.0	<2.0	<2.0	—	<2.0	—
03/30/95	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	—
04/26/95	<2.0	—	—	—	—	<2.0	—
05/30/95	<2.0	—	—	—	—	<2.0	—
06/29/95	<2.0	—	—	—	—	<2.0	<0.4
07/25/95	<2.0	<2.0	<2.0	<2.0	3.0	<2.0	—
08/31/95	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	—
09/19/95	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	—
10/24/95	<2.0	<2.0	<2.0	<2.0	<2.0	<2.0	—
11/09/95	<2.0	<2.0	<2.0	<2.0	5.0	<2.0	—
12/20/95	<2.0	<2.0	<2.0	<2.0	7.0	<2.0	<0.4

Table 2-12. Summary of Zinc ($\mu\text{g/L}$) Measurements

Sample Date	1	7	11	8	Plant Blowdown	Pond	Pond Sediment ($\mu\text{g/g}$)
01/30/95	6.9	7.5	<5.0	8.7	54	124	—
02/28/95	6.2	7.8	9.4	8.4	—	27	—
03/30/95	<5.0	<5.0	<5.0	<5.0	47	53	—
04/26/95	14.1	—	—	—	—	67	—
05/30/95	7.7	—	—	—	—	262	—
06/29/95	6.8	—	—	—	—	75	258
07/25/95	6.0	6.9	7.4	7.0	108	134	—
08/31/95	5.9	6.0	8.4	5.7	68	435	—
09/19/95	<5.0	8.6	<5.0	<5.0	53	774	—
10/24/95	5.4	5.4	6.0	5.4	65	41	—
11/09/95	<5.0	<5.0	<5.0	<5.0	57	44	—
12/20/95	6.6	7.2	5.4	5.0	68	62	740

Table 2-13. Summary of Iron ($\mu\text{g/L}$) Measurements

Sample Date	1	7	11	8	Plant Blowdown	Pond
01/30/95	41	44	46	49	906	196
02/28/95	207	206	216	218	-	29
03/30/95	35	33	37	33	360	67
04/26/95	241	-	-	-	-	186
05/30/95	149	-	-	-	-	70
06/29/95	196	-	-	-	-	144
07/25/95	103	101	107	100	435	-
08/31/95	70	73	72	71	603	38
09/19/95	56	70	69	61	589	71
10/24/95	83	87	69	75	624	24
11/09/95	74	87	83	75	1250	20
12/20/95	135	134	131	124	2010	28

Table 2-14. Summary of Lead ($\mu\text{g/L}$) Measurements

Sample Date	1	7	11	8	Plant Blowdown	Pond	Pond Sediment ($\mu\text{g/g}$)
01/30/95	2.7	<1.0	<1.0	<1.0	<9	<1.0	-
02/28/95	<1.0	3.3	<1.0	<1.0	-	<1.0	-
03/30/95	<1.0	<1.0	<1.0	<1.0	4.7	<1.0	-
04/26/95	<1.0	-	-	-	-	<1.0	-
05/30/95	<1.0	-	-	-	-	19	-
06/29/95	<1.0	-	-	-	-	14	<0.14*
07/25/95	2.0	8.0	2.0	4.0	1.0	2.0	-
08/31/95	<1.0	<1.0	<1.0	<1.0	<1.0	<1.0	-
09/19/95	<1.0	<1.0	<1.0	<1.0	<1.0	<1.0	-
10/24/95	<1.0	<1.0	<1.0	<1.0	1.0	<1.0	-
11/09/95	1.0	<1.0	<1.0	<1.0	2.0	<1.0	-
12/20/95	<1.0	<1.0	<1.0	<1.0	2.0	<1.0	14.2

*Detection limit for solid sample.



Table 2-15. Summary of Cadmium ($\mu\text{g/L}$) Measurements

Sample Date	1	7	11	8	Plant Blowdown	Pond	Pond Sediment ($\mu\text{g/g}$)
01/30/95	<1.4	<1.4	<1.4	<1.4	<1.4	<1.4	-
02/28/95	<1.4	<1.4	<1.4	<1.4	-	<1.4	-
03/30/95	<1.4	<1.4	<1.4	<1.4	<1.4	<1.4	-
04/26/95	<1.4	-	-	-	-	<1.4	-
05/30/95	<1.4	-	-	-	-	<1.4	-
06/29/95	<1.4	-	-	-	-	<1.4	3.6
07/25/95	<1.4	<1.4	<1.4	<1.4	<1.4	<1.4	-
08/31/95	<1.4	<1.4	<1.4	<1.4	<1.4	<1.4	-
09/19/95	<1.4	<1.4	<1.4	<1.4	<1.4	<1.4	-
10/24/95	<1.4	<1.4	<1.4	<1.4	5.1	<1.4	-
11/09/95	<1.4	<1.4	<1.4	<1.4	<1.4	<1.4	-
12/20/95	<1.4	<1.4	<1.4	<1.4	<1.4	<1.4	10.4

Table 2-16. Summary of Chromium ($\mu\text{g/L}$) Measurements

Sample Date	1	7	11	8	Plant Blowdown	Pond	Pond Sediment ($\mu\text{g/g}$)
01/30/95	<0.5	<0.5	<0.5	<0.5	1.1	<0.5	-
02/28/95	<0.5	<0.5	<0.5	<0.5	-	<0.5	-
03/30/95	<0.5	<0.5	<0.5	<0.5	5.0	<0.5	-
04/26/95	<0.5	-	-	-	-	<0.5	-
05/30/95	<0.5	-	-	-	-	<0.5	-
06/29/95	<0.5	-	-	-	-	<0.5	8.8
07/25/95	0.8	0.9	0.8	0.9	2.5	0.8	-
08/31/95	<0.5	<0.5	<0.5	<0.5	<0.5	1.6	-
09/19/95	<0.5	<0.5	<0.5	<0.5	2.1	<0.5	-
10/24/95	<0.5	<0.5	<0.5	<0.5	1.5	<0.5	-
11/09/95	0.8	0.8	0.7	0.9	2.8	0.9	-
12/20/95	0.9	1.0	1.0	0.9	5.0	0.9	11.2

2.3.8 Oil and Grease

Blowdown and pond oil and grease values were below the detection limit of 1.0 mg/L for all periods sampled. Oil and grease measurements are summarized in the following table.

Table 2-17. Summary of Oil and Grease (mg/L) Measurements

Sample Date	Plant Blowdown	Pond
01/30/95	<1.0	<1.0
02/28/95	-	<1.0
03/30/95	<1.0	<1.0
04/26/95	-	<1.0
05/30/95	-	<1.0
06/29/95	-	<1.0
07/29/95	<1.0	<1.0
08/31/95	<1.0	<1.0
09/19/95	<1.0	<1.0
10/24/95	<1.0	<1.0
11/09/95	<1.0	<1.0
12/20/95	<1.0	<1.0

2.3.9 Total Phosphorus and Inorganic Phosphate

Columbia River total phosphorus concentrations ranged from <0.01 to 0.02 mg/L as P. Blowdown values ranged from 2.4 to 4.7 mg/L as P. Columbia River inorganic phosphate concentrations were at or below 0.1 mg/L for all stations and periods, except Station II in March (0.2 mg/L). Blowdown inorganic phosphate measurements ranged from 0.8 to 1.6 mg/L as P. Total phosphorus and inorganic phosphate measurements are summarized in Tables 2-18 and 2-19.



Table 2-18. Summary of Total Phosphorus (mg/L as P) Measurements

Sample Date	1	7	11	8	Plant Blowdown
01/30/95	0.02	0.02	0.02	0.01	2.4
02/28/95	0.02	0.02	0.02	0.02	--
03/30/95	0.03	0.02	0.02	0.02	4.1
04/26/95	0.02	--	--	--	--
05/30/95	0.01	--	--	--	--
06/29/95	<0.01	--	--	--	--
07/25/95	0.01	0.01	0.01	0.01	4.4
08/31/95	0.01	0.01	0.01	0.01	4.4
09/19/95	0.01	0.02	0.02	0.02	4.4
10/24/95	0.01	0.02	0.02	0.02	4.1
11/09/95	0.01	0.02	0.01	0.01	4.7
12/20/95	0.03	0.02	0.02	0.02	3.3

Table 2-19. Summary of Inorganic Phosphate (mg/L as P) Measurements

Sample Date	1	7	11	8	Plant Blowdown
01/30/95	<0.1	<0.1	<0.1	<0.1	1.0
02/28/95	<0.1	<0.1	0.1	<0.1	--
03/30/95	<0.1	<0.1	0.2	<0.1	1.0
04/26/95	<0.1	--	--	--	--
05/30/95	<0.1	--	--	--	--
06/29/95	<0.1	--	--	--	--
07/25/95	<0.1	<0.1	<0.1	<0.1	0.8
08/31/95	<0.1	<0.1	<0.1	<0.1	0.9
09/19/95	<0.1	<0.1	<0.1	<0.1	0.8
10/24/95	<0.1	<0.1	<0.1	<0.1	1.3
11/09/95	<0.1	<0.1	<0.1	<0.1	1.2
12/20/95	<0.1	<0.1	<0.1	<0.1	1.6

2.3.10 Sulfate

Individual Columbia River sulfate measurements ranged from 9.3 to 10.8 mg/L. Blowdown measurements ranged from 419 to 775 mg/L. The results are presented in Table 2-20.

Table 2-20. Summary of Sulfate (mg/L) Measurements

Sample Date	1	7	11	8	Plant Blowdown
01/30/95	10.1	10.8	10.2	10.1	471
02/28/95	10.2	10.2	10.2	10.2	—
03/30/95	10.5	10.6	10.6	10.5	443
04/26/95	10.3	—	—	—	—
05/30/95	8.6	—	—	—	—
06/29/95	7.3	—	—	—	—
07/25/95	9.0	9.0	9.2	9.0	660
08/31/95	8.9	8.9	8.8	8.8	700
09/19/95	9.1	9.1	9.1	9.1	775
10/24/95	9.2	9.4	9.2	9.2	595
11/09/95	9.3	10.5	10.1	9.3	525
12/20/95	9.2	9.2	9.2	9.2	419

2.3.11 Total Dissolved Solids

The quarterly total dissolved solids (TDS) measurements of the pond ranged from 130 mg/L to 260 mg/L. This data is presented in Table 2-21.

Table 2-21. Summary of Quarterly Total Dissolved Solid (mg/L) Measurements

Sample Date	Conc.
03/30/95	250
06/29/95	130
09/19/95	230
12/20/95	260



2.3.12 VOCs and Semi-VOCs

Blowdown volatile and semivolatile concentrations were below their respective detection limits for all compounds during all periods.

Evaporation/percolation pond semivolatile organic compound concentrations were below their respective detection limits for all compounds during all periods, except bis (2-ethylhexyl) phthalate in March (480 $\mu\text{g/L}$) and September (24 $\mu\text{g/L}$). Limit of detection for bis(2-ethylhexyl) phthalate is 10 $\mu\text{g/L}$. Volatile organic concentrations were below their respective detection limits for all compounds during all periods, except freon 113 in September (24 $\mu\text{g/L}$) and December (88 $\mu\text{g/L}$) and chloroform in December (11 $\mu\text{g/L}$). Limits of detection for freon 113 and chloroform are 10 $\mu\text{g/L}$ and 5 $\mu\text{g/L}$, respectively. A list of the volatile and semivolatile organic compounds analyzed are presented in Tables 2-22 and 2-23, respectively.

Table 2-22. Summary of Volatile Organic Compounds

Chloromethane	Vinyl chloride	trans-1,3-Dichloropropene
Trichlorofluoromethane	Bromomethane	Dibromochloromethane
Freon 113	Chloroethane	Toluene
1,1-Dichloroethene	Carbon disulfide	2-Hexanone
Acetone	Methylene chloride	Ethylbenzene
cis-1,2-Dichloroethene	trans-1,2-Dichloroethene	Styrene
1,1-Dichloroethane	Chloroform	1,4-Dichlorobenzene
1,2-Dichloroethane	2-Butanone	1,1,2,2-Tetrachloroethane
1,1,1-Trichloroethane	Carbon tetrachloride	Bromoform
Benzene	Trichloroethene	4-Methyl-2-pentanone
1,2-Dichloropropane	Vinyl acetate	Tetrachloroethene
Bromodichloromethane	2-Chloroethylvinylether	Chlorobenzene
cis 1,3-Dichloropropene	1,1,2-Trichloroethane	Total Xylenes
1,3-Dichlorobenzene	1,2-Dichlorobenzene	

Table 2-23. Summary of Semivolatile Organic Compounds

<u>Acids</u>	<u>Base Neutrals</u>	
Phenol	2-Chloronaphthalene	2,4-Dinitrotoluene
2-Chlorophenol	2-Nitroaniline	Diethylphthalate
2-Methylphenol	Dimethylphthalate	Fluorene
4-Methylphenol	Acenaphthalene	4-Chlorophenyl-phenylether
		4-Nitroaniline
2-Nitrophenol	2,6-Dinitrotoluene	n-Nitrosodiphenylamine
2,4-Dimethylphenol	3-Nitroaniline	4-Bromophenyl-phenylether
2,4-Dichlorophenol	Acenaphthene	Hexachlorobenzene
		bis (2-Chloroethyl)ether
Benzoic Acid	Dibenzofuran	Anthracene
4-Chloro-3-methylphenol	Phenanthrene	Di-n-butylphthalate
2,4,6-Trichlorophenol	1,3-Dichlorobenzene	Fluoranthene
2,4,5-Trichlorophenol	1,4-Dichlorobenzene	Pyrene
2,4-Dinitrophenol	Benzyl Alcohol	
4-Nitrophenol	1,2-Dichlorobenzene	

Table 3-23. Summary of Semivolatile Organic Compounds

<u>Acids</u>	<u>Base Neutrals</u>	
Pentachlorophenol	n-Nitroso-di-n-propylamine	Benzoflanthracene
	Hexachloroethane	Nitrobenzene
	Isophorone	bis (2-Chloroethoxy)methane
	1,2,4-Trichlorobenzene	Naphthalene
	4-Chloroaniline	Dibenzo[a,h]anthracene
	Benzo[g,h,i]perylene	Hexachlorobutadiene
	2-Methylnaphthalene	Hexachlorocyclopentadiene
	Benzo[b]fluoranthene	3,3-Dichlorobenzidine
	Benzo[k]fluoranthene	Chrysene
	Benzo[a]pyrene	bis (2-Ethylhexyl)phthalate
	Indeno[1,2,3-cd]pyrene	

3.0 SOIL AND VEGETATION STUDIES


3.1 INTRODUCTION

The objective of the soil and vegetation studies is to identify any significant effects or impacts of plant cooling tower operation upon the plant communities surrounding WNP-2. Vegetation and soil sampling is conducted at the peak of the cheatgrass growth cycle known as the purple stage (Klemmedson 1964). Cheatgrass (*Bromus tectorum*) is the predominant species within all fifteen of the sampling plots with a mean frequency >98% and cover often approaching 50%.

Cheatgrass fruits turn purple shortly after reaching viability and then brown when mature. The purple stage of development correlates well with the peak productivity of many associated species and serves as a marker for initiation of annual sampling and comparison of phytomass productivity between years. The program includes the measurement of herbaceous canopy cover, herbaceous phytomass and soil chemistry. Soil chemical parameters measured include pH, carbonate, bicarbonate, sulfate, chloride, sodium, copper, zinc and conductivity. Fifteen sampling stations are located within a five mile radius of the plant. The stations consist of eight grassland (G01-G08) and seven shrub sites (S01-S07). The location of each station is illustrated in Figure 3-1.



Figure 3-1. Soil and Vegetation Sampling Location Map




800234.18 Map
 July 1993

3.2 MATERIALS AND METHODS

3.2.1 Herbaceous Canopy Cover

At each of the fifteen stations fifty microplots (20 cm x 50 cm) were placed at 1-meter intervals on alternate sides of the herbaceous transect (fig. 3-2). Canopy cover was estimated for each species occurring within a microplot using Daubenmire's (1968) cover classes. Data were recorded on standard data sheets. To assure the quality of the sampling, three randomly selected microplots were sampled twice. The entire transect was resampled if cover estimates for any major species (>50% frequency) differed by more than one cover class.

Figure 3-2. Layout of Vegetation and Soil Sampling Plots



3.2.2 Herbaceous Phytomass

Phytomass sampling was conducted concurrently with cover sampling. Phytomass sampling plots were randomly located within an area adjacent to the permanent transects or plots (Figure 3-2). At each station, all live herbaceous vegetation rooted in the designated microplot (20 x 50 cm) was clipped to ground level and placed in paper bags. Each bag was stapled shut and labeled with station code, plot number, date and personnel initials.

Sampling bags were transported to the laboratory, opened, and placed in a drying oven until a consistent weight was obtained. Following drying, the bags were removed singularly from the oven and their contents immediately weighed to the nearest 0.1 g. Laboratory quality assurance consisted of independently reworking 10 percent of the phytomass samples to assess data validity and reliability.

3.2.3 Soil Chemistry

At each of the fifteen grassland and shrub stations, two soil samples were collected from the top 15 cm of soil with a clean stainless steel trowel. The soil samples are randomly selected and taken from the phytomass sampling plot. The samples were placed in 250 ml sterile plastic cups with lids, labeled and refrigerated at 4°C. Nine parameters were analyzed in each sample, including pH, bicarbonate, carbonate, conductivity, sulfate, chloride, copper, zinc, and sodium. Aliquots of soil for trace metal analysis were microwave digested according to Gilman (1989). Preservation times and conditions, when applicable, followed EPA procedures (1983).

Laboratory quality control comprised 10-20% of the sample analysis load. Routine quality control samples included internal laboratory check standards, reagent blanks, and prepared EPA or NIST controls.

3.3 RESULTS AND DISCUSSION

During the 1995 season, 62 plant taxa were observed in the study areas. Table 3-1 lists the vascular plants observed during 1995 field studies.

Table 3-1. Vascular Plants Observed During 1995

<u>Scientific Name</u>	<u>Common Name</u>
APIACEAE	Pasley Family
<i>Cymopterus terebinthinus</i> (Hook.) T.&G. var. <i>terebinthinus</i>	Turpentine cymopterus
ASTERACEAE	Aster Family
<i>Achillea millefolium</i> L.	Yarrow
<i>Antennaria dimorpha</i> (Nutt.) T. & G.	Low pussy-toes
<i>Artemisia tridentata</i> Nutt.	Big Sagebrush
<i>Balsamorhiza careyana</i> Gray	Carey's balsamroot
<i>Chrysothamnus nauseosus</i> (Pall.) Britt	Gray rabbitbrush
<i>Chrysothamnus viscidiflorus</i> (Hook.) Nutt	Green rabbitbrush
<i>Crepis atrabarba</i> Heller	Slender hawksbeard
<i>Franseria acanthicarpa</i> Hook.	Bur ragweed
<i>Layia glandulosa</i> (Hook.) H & A	White daisy tidytips
<i>Tragopogon dubius</i> Scop.	Yellow salsify
<i>Aster canescens</i> Pursh	Hoary aster
BORAGINACEAE	Borage Family
<i>Amsinckia lycopsoides</i> Lehm.	Tarweed fiddleneck
<i>Cryptantha circumscissa</i> (H&A) Johnst.	Matted cryptantha
<i>Cryptantha leucophaea</i> (Dougl.) Pays	NA
<i>Cryptantha pterocarya</i> (Torr.) Greene	Winged cryptantha
BRASSICAEAE	Mustard Family
<i>Descurainia pinnata</i> (Walt.) Britt.	Western tansymustard
<i>Draba verna</i> L.	Spring draba
<i>Erysimum asperum</i> (Nutt.) DC.	Prairie rocket
<i>Sisymbrium altissimum</i> L.	Tumblemustard
CACTACEAE	Cactus Family
<i>Opuntia polycantha</i> Haw.	Starvation cactus

Table 3-1. Vascular Plants Observed During 1995 (Continued)

<u>Scientific Name</u>	<u>Common Name</u>
CARYOPHYLLACEAE	Pink Family
<i>Arenaria franklinii</i> Dougl. var <i>franklinii</i>	Franklin's sandwort
<i>Holosteum umbellatum</i> L.	Jagged chickweed
CHENOPODIACEAE	Chenopod Family
<i>Chenopodium leptophyllum</i> (MOQ.) Wats.	Slimleaf goosefoot
<i>Grayia spinosa</i> (Hook.) MOQ.	Russian thistle
<i>Salsola kali</i> L.	
FABACEAE	Pea Family
<i>Astragalus purshii</i> Dougl.	Wooly-pod milk-vetch
<i>Astragalus sclerocarpus</i> Gray	Stalked-pod milk-vetch
<i>Psoralea lanceolata</i> Pursh	Lance-leaf scurf-pea
GERANIACEAE	Geranium Family
<i>Erodium cicutarium</i> (L.) L'Her.	Filaree, storks-bill
HYDROPHYLLACEAE	Waterleaf Family
<i>Phacelia hastata</i> Dougl.	Whiteleaf phacelia
<i>Phacelia linearis</i> (Pursh) Holz.	Threadleaf phacelia
LILIACEAE	Lily Family
<i>Brodiaea douglasii</i> Wats.	Douglas' brodiaea
<i>Calochortus macrocarpus</i> Dougl.	Sego lily
<i>Fritillaria pudica</i> (Pursh) Spreng.	Chocolate lily
LOASACEAE	Blasing-star Family
<i>Mentzelia albicaulis</i> Dougl. Ex Hook.	White-stemmed mentzelia
MALVACEAE	Mallow Family
<i>Sphaeralcea munroana</i> (Dougl.) Spach Ex Gray	White-stemmed globe-mallow
ONAGRACEAE	Evening-primrose Family
<i>Oenothera pallida</i> Lindl. var. <i>pallida</i>	White-stemmed evening-primrose
PLANTAGINACEAE	Plantain Family
<i>Plantago patagonica</i> Jacq.	Indian-wheat



Table 3-1. Vascular Plants Observed During 1995 (Continued)

<u>Scientific Name</u>	<u>Common Name</u>
POACEAE	Grass Family
<i>Agropyron cristatum</i> (L.) Gaertn.	Crested wheatgrass
<i>Agropyron dasystachyum</i> (Hook.) Scribn.	Thick-spiked wheatgrass
<i>Agropyron spicatum</i> (Pursh) Scribn. & Smith	Bluebunch wheatgrass
<i>Bromus tectorum</i> L.	Cheatgrass
<i>Festuca octoflora</i> Walt.	Six-weeks fescue
<i>Koeleria cristata</i> Pers.	Prairie Junegrass
<i>Oryzopsis hymenoides</i> (R&S) Ricker	Indian ricegrass
<i>Poa sandbergii</i> Vasey	Sandberg's bluegrass
<i>Sitanion hystrix</i> (Nutt.) Smith	Bottlebrush squirreltail
<i>Stipa comata</i> Trin. & Rupr.	Needle-and-thread
POLEMONIACEAE	Phlox Family
<i>Gilia minutiflora</i> Benth.	Gilia
<i>Gilia sinuata</i> Dougl.	Shy gilia
<i>Leptodactylon pungens</i> (Torr.) Nutt.	Granite gilia
<i>Microsteris gracilis</i> (Hook.) Greene var. <i>humilior</i> (Hook.) Cronq.	Pink microsteris
<i>Phlox longifolia</i> Nutt.	Long-leaf phlox
POLYGONACEAE	Buckwheat Family
<i>Eriogonuum niveum</i> Dougl.	Snow buckwheat
<i>Rumex venosus</i> Pursh	Wild begonia
RANUNCULACEAE	Buttercup Family
<i>Delphinium nuttallianum</i> Pritz. ex Walpers	Larkspur
ROSACEAE	Rose Family
<i>Purshia tridentata</i> (Pursh) DC	Antelope Bitterbrush
SANTALACEAE	Sandalwood Family
<i>Comandra umbellata</i> (L.) Nutt.	Bastard toad-flax
SAXIFRAGACEAE	Saxifrage Family
<i>Ribes aureum</i> Pursh	Golden current
SCROPHULARIACEAE	Figwort Family
<i>Penstemon acuminatus</i> Dougl.	Sand-dune penstemon

Table 3-1. Vascular Plants Observed During 1995 (Continued)

<u>Scientific Name</u>	<u>Common Name</u>
VALERIANACEAE	Valerian Family
<i>Plectritis macrocera</i> T&G	Longhorn plectritis

3.3.1 Herbaceous Cover

Total herbaceous cover averaged 92.39% in 1995 which represents an increase of 105% from 1994 (45.05). With the exception of station GO6, all other stations showed an increase of 45% or greater in total herbaceous cover. *Bromus tectorum* continues to be the dominant annual grass with an average cover of 34.42%, an increase of 35.25%. Total perennial grass cover was 27.60%, an increase of 212%. As in previous years, the dominant perennial grass was *Poa sandbergii* with an average cover of 19.69%. The most significant change in cover occurred in the annual forbs. Total annual forb cover increased 395% from last year. *Sisymbrium altissimum* had an average cover of 7.44% compared to 0.54 % last year. *Draba verna* with a previous cover of 0.88% increased 509% with an average cover of 5.36% for 1995. The total perennial forb cover was 3.5%.

Frequency values (%) increased at eight of the fifteen stations. The most significant increase in frequency values was seen in annual forbs. Only two stations (SO2 and SO4) showed a decrease in the number of annual forb species per site. Station GO5 had an increase of six annual forb species. The total species per site for station GO5 increased from 14 to 21. Station SO6 increased from 7 to 14 in total species per site. Table 3-3 shows mean frequency values (%) by species for each sampling station.

Table 3-2. Herbaceous Cover for Fifteen Sampling Stations (%)

	G01	G02	G03	G04	G05	G06	G07	G08	S01	S02	S03	S04	S05	S06	S07	AVG. G01-S07
Annual Grasses																
<i>Bromus tectorum</i>	78.65	70.60	55.50	4.80	10.50	27.60	53.65	60.20	31.80	9.25	25.80	36.15	46.75	4.00	1.10	34.42
<i>Festuca octoflora</i>	0.00	0.00	0.00	0.00	0.60	0.05	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.05
Total Annual Grass Cover	78.65	70.60	55.50	4.80	11.10	27.65	53.65	60.20	31.80	9.25	25.80	36.15	46.25	4.00	1.10	34.47
Perennial Grasses																
<i>Agropyron spicatum</i>	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.30	0.00	0.00	0.02
<i>Oryzopsis hymenoides</i>	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	2.60	0.00	0.00	0.00	2.30	0.00	0.33
<i>Poa sandbergii</i>	10.15	18.00	4.00	38.25	47.45	1.95	16.75	16.25	17.60	21.80	30.00	8.64	0.00	51.85	12.65	19.69
<i>Stipa comata</i>	0.00	0.00	0.00	26.60	0.00	2.45	0.00	6.65	0.00	14.40	0.00	0.00	2.00	0.40	0.00	7.50
Total Perennial Grass Cover	10.15	18.00	4.00	64.85	47.45	4.40	16.75	22.90	17.60	98.80	30.00	8.64	2.30	55.55	12.65	27.60
Annual Forbs																
<i>Amsinckia lycopsoides</i>	0.05	0.10	9.50	0.05	0.50	0.05	11.20	0.05	4.15	0.35	0.00	0.00	3.30	0.10	2.35	2.12
<i>Brodiaea douglasii</i>	0.00	0.00	0.00	0.00	0.05	0.00	0.00	0.05	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01
<i>Cryptantha cirmumscissa</i>	0.00	0.00	0.00	0.05	0.05	0.15	0.00	0.00	0.15	0.00	0.00	0.00	0.15	0.00	0.00	0.04
<i>Descurainia pinnata</i>	0.00	0.10	0.10	0.00	0.05	0.00	1.70	0.00	0.55	0.00	0.00	0.00	0.00	1.85	57.80	4.14
<i>Draba verna</i>	16.55	15.15	3.65	3.30	14.00	0.05	5.55	8.15	4.45	0.00	4.60	2.00	2.00	0.90	0.10	5.36
<i>Erodium cicutarium</i>	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.85	0.00	0.00	0.06
<i>Franseria acanthocarpa</i>	0.00	0.00	3.15	0.10	0.10	0.40	1.40	0.05	0.95	0.00	0.00	0.00	0.30	0.05	0.00	0.43
<i>Gilia minusflora</i>	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.20	0.00	0.00
<i>Holosteum umbellatum</i>	2.30	8.65	2.75	3.70	9.30	0.35	4.60	9.10	7.10	5.70	6.55	2.60	2.90	0.00	0.60	4.41
<i>Mentzelia albicaulis</i>	0.00	0.00	0.00	0.00	0.00	0.00	0.05	0.00	0.00	0.05	0.00	0.10	0.00	0.00	0.00	0.01
<i>Microrsteris gracilis</i>	0.15	0.00	3.80	0.15	0.60	2.30	2.45	0.20	7.60	0.00	9.50	0.00	1.20	0.00	0.00	1.29
<i>Phacelia linearis</i>	0.00	0.00	0.05	0.05	0.10	0.10	0.00	0.00	0.10	0.00	0.00	0.00	0.20	0.00	0.00	0.04
<i>Pyrago patagonica</i>	0.40	1.15	0.00	1.70	0.00	0.00	0.00	0.40	0.00	0.00	5.00	0.00	1.45	0.00	0.50	0.71
<i>Thymus kalli</i>	1.35	0.15	0.15	0.10	1.45	0.20	0.05	0.05	5.40	0.45	0.00	0.45	0.65	0.05	0.00	0.70
<i>Thymus altissimum</i>	2.95	1.90	0.30	0.00	0.10	0.15	3.85	2.60	1.00	1.10	0.65	44.47	1.70	11.15	40.75	7.44
Total Annual Forb Cover	23.75	27.50	23.35	9.65	26.25	3.75	30.85	20.65	31.45	6.65	17.75	49.62	14.75	14.30	102.1	26.82
Perennial Forbs																
<i>Achillea millefolium</i>	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.05	0.00	0.00	0.00	1.30	0.00	0.10
<i>Aster canescens</i>	0.00	0.00	0.10	0.00	0.00	0.00	0.05	0.00	1.75	0.00	0.00	0.30	0.00	0.00	0.00	0.88
<i>Astragalus sclerocarpus</i>	0.00	0.00	0.00	0.00	0.00	1.30	0.00	0.00	0.00	0.00	0.00	1.05	0.30	0.00	0.00	0.09
<i>Balsamorhiza careyana</i>	0.00	0.00	0.00	0.00	3.40	0.05	0.00	0.00	0.00	0.00	0.00	2.30	1.70	0.00	0.00	0.43
<i>Comandra umbellata</i>	0.00	0.00	0.00	0.00	0.60	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.05
<i>Crepis atrabarba</i>	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	2.40	0.00	0.00	0.00	0.00	0.16
<i>Cymopterus terebinthinus</i>	0.00	0.60	0.00	0.00	0.00	0.00	4.15	0.05	0.00	0.00	6.95	0.00	0.00	0.00	6.45	1.51
<i>Eriogonum niveum</i>	0.00	0.00	0.00	0.00	1.40	0.00	0.00	0.00	0.05	0.00	0.00	0.00	6.75	0.00	0.00	0.55
<i>Genothera pallida</i>	0.00	0.00	0.20	0.20	0.85	0.15	0.00	0.00	0.05	0.05	0.00	0.00	0.00	0.00	0.00	0.12
<i>Phlox longifolia</i>	0.10	0.05	0.05	0.50	0.90	0.00	0.05	2.80	0.30	0.35	0.00	0.00	0.00	3.00	0.00	0.42
<i>Rumex venosus</i>	0.05	0.00	0.10	0.00	0.00	0.00	0.00	0.00	0.10	0.00	0.00	0.00	0.00	0.00	0.00	0.01
Total Perennial Forb Cover	0.15	0.65	0.45	0.25	7.15	5.65	0.15	2.80	2.25	7.40	2.40	3.65	8.48	4.60	6.45	3.50
Total Herbaceous Cover	112.7	116.8	85.30	79.55	91.95	41.45	101.4	106.6	83.10	122.1	75.95	98.06	72.28	78.45	122.3	92.39

Table 3-3. Mean Frequency Values (%) by Species for Each Sampling Station

	GO1	GO2	GO3	GO4	GO5	GO6	GO7	GO8	SO1	SO2	SO3	SO4	SO5	SO6	SO7
Annual Grasses															
<i>Bromus tectorum</i>	100	100	100	76	96	98	96	96	100	52	90	100	100	34	34
<i>Festuca octoflora</i>					14	2						4			
Perennial Grasses															
<i>Agropyron spicatum</i>													2		
<i>Oryzopsis hymenoides</i>										10				14	
<i>Poa sandbergii</i>	86	82	8	96	94	10	48	56	38	52	100	30		86	80
<i>Stipa comata</i>				74		4		12		50			4	6	
Annual Forbs															
<i>Amsinckia lycopsoides</i>	2	4	44	2	10	2	56	2	40	4			26	4	16
<i>Chenopodium leptophyllum</i>													6		
<i>Cryptantha circumscissa</i>				2	2	4			6						
<i>Cryptantha pterocarya</i>									6						
<i>Descurainia pinnata</i>		4	4		2		2		12					36	96
<i>Draba verna</i>	94	98	98	82	98	2	92	98	88		94	60	70	26	4
<i>Erodium cicutarium</i>													2		
<i>Franseria acanthacarpa</i>			66	4	4	16	36	2	18				12	2	
<i>Gilia sinuata</i>					2					2					
<i>Holosteum umbellatum</i>	62	98	90	52	98	4	74	96	94	58	94	54	76		14
<i>Layia glandulosa</i>					2										
<i>Mentzelia albicaulis</i>							2			2		4			
<i>Microsteris gracilis</i>	6		72	6	14	62	20	8	68		28		28		
<i>Phacelia linearis</i>			2	2	4				4	4	22		8		
<i>Plantago patagonica</i>	16	26		48				6			74		28		10
<i>Salsola kali</i>	54	6	6	4	38	8	2	2	62	18		18	26	2	
<i>Sisymbrium altissimum</i>	30	26	12		4	6	20	16	20	4	26	86	20	52	78
<i>Tragopogon dubius</i>												2			
Perennial Forbs															
<i>Achillea millifolium</i>															4
<i>Aster canescens</i>			4				2						2		
<i>Astragalus purshii</i>															
<i>Astragalus sclerocarpus</i>						4					4			2	
<i>Balsamorhiza careyana</i>				4	2						6	2			
<i>Brodiaea douglasii</i>				2				2	2						
<i>Comandra umbellata</i>				4											
<i>Crepis atrabarba</i>															
<i>Cymopterus terebinthinus</i>		4				22									
<i>Eriogonum niveum</i>					6				8				20		
<i>Oenothera pallida</i>			8	8	14	6			2	2		4	40		
<i>Phlox longifolia</i>	4	2	2	10	8		2	32	2	4			6	6	
<i>Rumex venosus</i>	2		4						4						
Total Species per Site	11	11	15	14	21	16	13	13	18	11	9	10	19	14	8



Table 3-3. Mean Herbaceous Cover for 1975 through 1995

CLASS	YEAR	S01	S02	S03	S04	S05	X S01-5	S06	S07	X5	G01	G02	G03	G04	X G01-4	G05	G06	G07	G08	XG	X5G	G01-4, S01-5
AG	1975	49.90	35.30	43.80						43.00	43.90	43.00								43.45	43.18	43.18
PG	1975	0.60	2.00	4.50						2.37	3.70	5.50								4.60	3.26	3.26
AF	1975	14.60	11.70	11.70						12.67	29.50	13.00								21.25	16.10	16.10
PF	1975	4.30	0.90	1.80						2.33	1.50	2.10								1.80	2.12	2.12
ALL	1975	69.40	49.90	61.80						60.37	78.60	63.60								71.10	64.66	64.66
AG	1976	50.70	40.90	34.30						41.97	71.20	51.60								61.40	49.74	49.74
PG	1976	0.40	10.50	10.30						7.07	4.40	3.10								3.75	5.74	5.74
AF	1976	5.50	5.30	7.20						6.00	11.90	8.50								10.20	7.68	7.68
PF	1976	0.00	0.50	0.20						0.23	0.00	0.20								0.10	0.18	0.18
ALL	1976	56.60	57.20	52.00						55.27	87.50	63.40								75.45	63.34	63.34
AG	1977	1.35	0.65	1.90						1.30	5.20	1.45								3.33	2.11	2.11
PG	1977	0.35	11.30	8.28						6.64	3.25	2.90								3.08	5.22	5.22
AF	1977	0.25	0.05	0.90						0.40	2.40	9.35								5.88	2.59	2.59
PF	1977	0.55	0.60	1.42						0.86	0.05	6.30								3.18	1.78	1.78
ALL	1977	2.50	12.60	12.50						9.20	10.90	20.00								15.45	11.70	11.70
AG	1978	51.00	67.00	51.00						56.33	68.00	42.00								55.00	55.80	55.80
PG	1978	3.00	18.00	11.00						10.67	8.00	7.00								7.50	9.40	9.40
AF	1978	34.00	10.00	33.00						27.00	23.00	25.00								24.00	25.80	25.80
PF	1978	8.00	0.00	5.00						4.33	2.00	3.00								2.50	3.60	3.60
ALL	1978	100.00	95.00	100.00						98.33	101.00	77.00								89.00	97.60	94.60
AG	1979	25.00	29.00	9.00						21.00	31.00	10.00								20.50	20.80	20.80
PG	1979	1.00	18.00	11.00						10.00	7.00	5.00								6.00	8.40	8.40
AF	1979	2.00	4.00	10.00						5.33	43.00	33.00								38.00	18.40	18.40
PF	1979	11.00	0.00	3.00						4.67	0.00	7.00								3.50	4.20	4.20
ALL	1979	39.00	51.00	33.00						41.00	81.00	55.00								68.00	51.80	51.80
AG	1980	50.40	51.80	24.30	56.20	56.40	47.82			47.82	64.30	77.80	73.80	12.30	57.05					57.05	51.92	51.92
PG	1980	1.00	7.20	23.30	10.90	0.10	8.50			8.50	28.30	64.00	0.10	26.60	29.75					29.75	17.94	17.94
AF	1980	7.60	4.20	22.50	3.40	14.10	10.36			10.36	7.30	5.00	28.70	4.90	11.48					11.48	10.86	10.86
PF	1980	2.20	2.20	4.70	4.60	1.80	3.10			3.10	0.40	0.00	0.00	4.60	1.25					1.25	2.28	2.28
ALL	1980	61.20	65.40	74.80	75.10	72.40	69.78			69.78	100.30	146.80	102.60	48.40	99.53					99.53	83.00	83.00
AG	1981	74.80	54.60	66.50	49.80	76.20	64.38			64.38	77.40	84.00	88.40	48.90	74.68					74.68	68.96	68.96
PG	1981	0.10	4.70	14.30	5.80	0.00	4.98			4.98	19.60	25.90	0.00	36.70	20.55					20.55	11.90	11.90
AF	1981	5.30	3.50	18.20	1.20	12.50	8.14			8.14	15.90	11.90	17.50	5.90	12.80					12.80	10.21	10.21
PF	1981	0.00	3.20	0.70	4.90	0.50	1.86			1.86	0.20	0.00	0.00	1.90	0.53					0.53	1.27	1.27
ALL	1981	80.20	66.00	99.70	61.70	89.20	79.36			79.36	113.10	121.80	105.90	93.40	108.55					108.55	92.33	92.33
AG	1982	51.50	25.80	36.60	32.70	20.00	33.32			33.32	42.20	45.50	51.00	22.90	40.40					40.40	36.47	36.47
PG	1982	0.40	6.40	17.90	4.30	0.80	5.96			5.96	11.20	11.60	0.10	31.30	13.55					13.55	9.33	9.33
AF	1982	4.60	4.20	7.50	1.60	17.30	7.04			7.04	9.70	4.60	4.60	4.10	5.75					5.75	6.47	6.47
PF	1982	0.20	4.30	0.70	6.20	1.00	2.48			2.48	0.30	0.00	1.30	3.80	1.35					1.35	1.98	1.98
ALL	1982	56.70	40.70	62.70	44.80	39.10	48.80			48.80	63.40	61.70	57.00	62.10	61.05					61.05	54.24	54.24



Table 3-3. Mean Herbaceous Cover for 1975 through 1995 (continued)

CLASS	YEAR	S01	S02	S03	S04	S05	X S01-5	S06	S07	XS	G01	G02	G03	G04	X G01-4	G05	G06	G07	G08	XG	XSG	G01-4, S01-5
AG	1983	53.80	37.60	33.65	36.75	31.85	38.73			38.73	49.50	39.55	62.75	17.55	42.35					42.34	40.33	40.33
PG	1983	2.15	7.70	14.45	6.40	1.29	6.40			6.40	2.10	15.75	0.00	25.50	10.84					10.84	8.37	8.37
AF	1983	8.20	7.85	12.55	3.45	22.35	10.88			10.88	18.70	8.85	8.65	6.65	10.71					10.71	10.81	10.81
PF	1983	0.70	3.10	1.05	4.40	1.95	2.24			2.24	0.65	0.05	2.10	4.00	1.70					1.70	2.00	2.00
ALL	1983	64.85	56.25	61.70	51.00	57.44	58.25			58.25	70.95	64.20	73.50	53.70	65.59					65.59	61.51	61.51
AG	1984	41.50	32.75	39.35	36.30	36.50	37.28			37.28	60.85	71.30	60.85	9.60	50.65					50.65	43.22	43.22
PG	1984	1.85	8.80	11.55	8.55	0.40	6.23			6.23	1.20	4.45		25.00	10.22					10.22	6.87	7.73
AF	1984	12.35	8.10	11.10	4.00	13.40	9.79			9.79	20.65	9.70	19.45	7.95	14.44					14.44	11.86	11.86
PF	1984	0.30	4.00	0.75	6.55	0.65	2.45			2.45	0.70	0.20	1.10	1.25	0.81					0.81	1.72	1.72
ALL	1984	56.00	53.65	62.75	55.40	50.95	55.75			55.75	83.40	85.65	81.40	43.80	73.56					73.56	63.67	63.67
AG	1985	2.10	2.15	14.60	4.95	27.05	10.17			10.17	8.00	8.10	18.30	7.25	10.41					10.41	10.28	10.28
PG	1985	1.05	4.70	17.85	2.40	1.85	5.57			5.57	9.20	17.95	0.00	13.90	10.26					10.26	7.66	7.66
AF	1985	0.70	1.35	9.40	2.30	4.75	3.70			3.70	18.20	8.15	7.55	3.05	9.24					9.24	6.16	6.16
PF	1985	0.00	1.35	1.15	3.00	0.25	1.15			1.15	0.80	0.10	2.35	0.90	1.04					1.04	1.10	1.10
ALL	1985	3.85	9.55	43.00	12.65	33.90	20.59			20.59	36.20	34.30	28.20	25.10	30.95					30.95	25.19	25.19
AG	1986	17.45	1.95	7.20	11.45	13.05	10.22			10.22	9.40	4.65	13.25	7.35	8.66					8.66	9.53	9.53
PG	1986	2.20	10.75	17.25	9.85	1.30	8.27			8.27	19.85	38.65	0.00	26.00	21.13					21.13	13.98	13.98
AF	1986	25.40	16.65	38.10	10.25	16.70	21.42			21.42	27.65	34.15	25.45	8.70	23.99					23.99	22.56	22.56
PF	1986	1.15	5.35	2.30	9.15	1.25	3.84			3.84	1.80	1.95	0.05	2.55	1.59					1.59	2.84	2.84
ALL	1986	46.20	34.70	64.85	40.70	32.30	43.75			43.75	58.70	79.40	38.75	44.60	55.36					55.36	48.91	48.91
AG	1987	28.90	9.95	7.80	19.05	33.40	19.82			19.82	23.85	9.45	51.65	4.65	22.40					22.40	20.97	20.97
PG	1987	3.60	21.90	42.65	19.55	2.30	18.00			18.00	32.45	58.79	0.05	45.95	34.31					34.31	25.25	25.25
AF	1987	12.56	8.50	10.80	6.55	11.40	9.96			9.96	10.30	11.32	14.00	3.25	9.72					9.72	9.85	9.85
PF	1987	5.00	6.00	2.00	10.40	1.75	5.03			5.03	0.90	1.90	0.15	1.55	1.13					1.13	3.29	3.29
ALL	1987	50.06	46.35	63.25	55.35	48.85	52.81			52.81	67.50	81.46	65.85	55.40	67.53					67.53	59.36	59.36
AG	1988	13.80	5.05	8.10	13.80	10.15	10.18	10.40	12.24	10.51	22.95	10.10	16.75	4.80	13.65	11.95	19.20	15.85	10.40	14.00	12.32	11.72
PG	1988	1.75	8.40	11.95	9.40	3.35	6.97	16.85	17.50	9.89	17.85	21.70	0.05	30.20	17.45	9.50	12.05	10.45	14.30	14.51	12.34	11.63
AF	1988	6.08	5.25	3.60	3.10	4.00	4.41	0.00	0.35	3.20	6.30	16.15	7.55	1.80	7.95	1.20	1.45	12.35	6.12	6.61	5.16	5.98
PF	1988	11.55	15.75	2.10	4.85	3.25	7.50	0.10	0.00	5.37	0.20	2.00	0.00	4.40	1.65	15.25	8.70	2.45	4.34	4.34	4.79	4.90
ALL	1988	33.18	34.45	25.75	31.15	20.75	29.06	27.35	30.09	28.96	47.30	49.95	24.35	41.20	40.70	37.90	41.40	41.10	32.52	39.47	34.60	34.23
AG	1989	21.85	12.50	12.45	10.25	32.90	17.99	15.00	47.65	21.80	22.50	13.20	65.85	3.05	26.15	22.35	35.10	38.05	12.05	26.52	24.05	21.62
PG	1989	8.30	29.55	64.00	13.00	1.25	23.22	30.35	37.50	26.28	60.40	59.60	0.05	49.55	42.40	36.75	16.20	32.05	48.95	37.94	32.54	31.74
AF	1989	12.50	6.95	13.05	6.45	11.10	10.01	0.85	5.15	8.01	12.85	5.90	42.20	2.85	15.95	8.85	13.55	13.05	13.95	14.15	11.48	12.65
PF	1989	4.45	14.50	4.40	8.20	0.55	6.42	0.10	0.00	4.60	3.85	1.10	0.05	3.00	2.00	6.45	10.40	12.90	10.60	6.04	5.23	4.46
ALL	1989	47.10	63.50	93.90	37.90	45.80	57.64	46.30	90.30	60.69	99.60	79.80	108.15	58.45	86.50	74.40	75.25	96.05	85.55	84.66	73.31	70.47
AG	1990	36.80	16.80	17.50	32.40	53.35	31.37	12.90	5.45	25.03	18.60	7.75	61.55	13.65	25.39	23.80	35.45	36.55	19.75	27.01	26.06	28.71
PG	1990	3.30	12.85	18.35	12.70	0.05	9.45	18.40	17.55	11.89	18.70	0.00	0.00	30.00	12.18	11.50	10.70	9.30	12.10	11.59	11.73	10.66
AF	1990	7.95	2.60	3.15	4.55	8.90	6.43	0.10	0.00	4.61	7.75	2.35	15.70	3.35	7.90	2.75	6.90	8.95	7.00	6.84	5.80	6.81
PF	1990	0.40	9.55	1.75	3.90	0.05	3.313	0.00	0.00	2.24	0.00	0.05	0.05	1.20	3	3.95	8.55	0.05	0.20	1.76	1.98	1.88
ALL	1990	48.45	41.80	45.20	53.35	62.35	50.29	31.40	23.00	43.69	45.05	10.15	77.30	48.20	45.18	42.40	61.60	53.85	39.05	47.20	45.56	48.02
AG	1991	40.25	15.25	40.05	38.55	48.15	35.85	17.85	5.90	25.14	26.15	20.80	65.55	18.90	32.85	36.95	37.25	48.30	38.25	36.52	33.81	34.52
PG	1991	7.60	32.05	26.35	14.45	2.30	11.14	38.40	60.60	25.96	41.75	50.55	1.35	38.70	29.09	23.55	12.80	0.00	22.85	23.94	26.14	11.12
AF	1991	36.25	15.00	16.75	37.30	21.60	24.29	4.85	7.30	19.86	0.25	4.20	13.35	1.85	4.92	4.75	6.30	35.13	16.65	10.31	14.77	16.26
PF	1991	4.45	6.35	1.95	2.35	0.30	3.08	0.30	0.00	2.24	0.00	0.10	0.00	0.90	0.25	3.35	12.20	0.05	1.70	2.29	2.25	1.88
ALL	1991	88.55	63.65	85.10	89.65	72.35	74.36	61.40	73.80	76.36	61.15	75.65	80.25	60.35	66.11	68.60	68.55	83.48	79.45	72.19	76.97	63.78
AG	1992	30.30	30.20	42.60	55.95	51.60	42.13	23.90	15.20	35.67	48.70	64.25	53.15	34.24	30.09	46.00	41.80	66.15	55.15	51.18	43.95	45.67
PG	1992	3.25	15.65	11.40	5.40	2.39	7.62	31.30	33.80	14.74	25.60	20.00	0.00	32.20	19.45	18.60	10.20	5.95	8.80	15.17	14.97	12.88
AF	1992	9.85	5.55	11.95	16.40	8.95	10.54	4.65	23.05	11.48	13.15	8.15	15.05	7.15	10.87	7.45	10.20	8.80	17.25	10.93	11.19	10.693
PF	1992	9.15	10.70	2.25	4.25	1.05	5.48	0.65	6.00	4.01	0.10	0.25	0.30	0.75	0.35	1.95	12.55	1.35	3.85	2.64	3.28	2.00
ALL	1992	52.55	62.10	68.20	82.00	63.99	65.77	60.50	72.70	65.90	87.55	92.65	68.50	74.34	80.76	74.20	74.75	82.25	85.05	79.92	73.39	72.44




Table 3-3. Mean Herbaceous Cover for 1975 through 1995 (continued)

CLASS	YEAR	S01	S02	S03	S04	S05	X S01-5	S06	S07	XS	O01	O02	O03	O04	X O01-4	O05	O06	O07	O08	XG	XSG	O01-4, S01-5
AG	1993	27.70	34.65	53.45	58.25	48.20	44.45	23.65	57.95	43.41	46.90	68.65	43.40	29.20	47.04	38.35	28.90	68.85	59.60	47.98	45.85	45.74
PG	1993	7.15	22.14	16.25	12.85	4.00	12.48	46.10	23.15	18.81	48.25	23.35	2.00	46.10	29.93	31.40	15.40	12.25	16.35	24.39	21.78	21.2
AF	1993	12.95	8.70	12.90	14.80	13.25	12.52	2.15	9.85	10.66	13.45	5.95	22.60	10.20	13.05	10.90	16.45	13.45	9.02	12.75	11.77	12.79
PF	1993	13.70	12.70	2.690	8.65	7.40	9.01	0.05	0.05	6.45	1.45	0.00	0.75	2.15	1.09	5.85	8.85	1.55	4.50	3.14	4.68	5.05
ALL	1993	58.10	77.49	85.20	94.55	71.35	77.34	71.95	88.90	78.22	110.00	97.95	66.25	87.55	90.44	86.00	69.60	94.40	89.27	87.63	84.08	83.89
AG	1994	23.30	11.00	28.40	21.20	35.50	23.88	5.35	9.70	19.21	47.50	61.85	27.50	9.05	36.48	6.35	16.70	42.40	51.20	32.88	26.50	30.18
PG	1994	2.65	16.95	5.70	3.75	2.15	6.24	11.20	9.55	7.42	5.50	4.30	0.60	34.30	11.18	9.55	16.65	2.20	7.20	10.04	8.82	8.71
AF	1994	8.10	2.70	7.40	3.30	7.65	5.83	0.15	2.90	4.60	4.10	1.90	18.15	2.50	6.66	2.90	6.65	11.35	2.25	6.14	5.42	6.25
PF	1994	2.25	7.15	3.60	4.50	7.50	5.00	1.50	6.45	4.68	0.15	0.05	11.25	0.60	3.01	5.30	11.10	0.75	2.80	4.00	4.32	4.01
ALL	1994	36.30	37.80	45.10	32.75	52.80	40.95	18.00	28.60	35.91	57.25	68.10	57.50	46.45	57.33	23.40	51.60	56.70	63.45	53.06	45.05	49.14
AG	1995	31.80	9.25	25.8	36.15	46.75	29.95	4.0	1.1	22.12	78.65	70.60	55.50	4.80	52.39	11.10	27.65	53.65	60.20	45.21	34.47	39.92
PG	1995	17.60	98.80	30.0	8.64	2.3	31.47	55.55	12.65	32.22	10.15	18.0	4.00	64.85	24.25	47.45	4.40	16.75	22.90	26.59	27.60	28.26
AF	1995	31.45	6.65	17.75	49.62	14.75	24.04	14.30	102.1	33.81	23.75	27.5	23.35	9.65	21.06	26.25	3.75	30.85	20.65	20.72	26.96	22.72
PF	1995	2.25	7.4	2.4	3.65	8.48	4.84	4.6	6.45	5.03	0.15	0.65	0.45	0.38	7.15	5.65	0.15	2.80	2.16	3.50	2.86	
ALL	1995	83.10	122.1	75.95	98.06	72.28	90.30	78.45	122.30	93.18	112.70	116.75	83.30	79.55	98.08	91.95	41.45	101.40	106.55	91.71	92.39	93.76

Figure 3-3 shows a comparison of the current data with previous data. Growing season (October 94 - April 95) precipitation (21.06 cm) increased 465% from the previous season (3.73 cm). According to Battelle Northwest Laboratories, the months of January through April were recorded as the wettest four months of the year on record. The mean temperature during the growing season was 6.43°C compared with 6.161°C for 1994. A comparison of mean cover and precipitation for 1982 through 1995 can be seen in Figure 3-4.

Fig. 3-3. Mean Herbaceous Cover for 1975-1995



3.3.2 Herbaceous Phytomass

The increase (260%) in herbaceous phytomass is in direct correlation to the increase in herbaceous cover. At grassland and shrub stations, the herbaceous phytomass production averaged 107g/m² and 123g/m² respectively. Mean herbaceous phytomass production at grassland and shrub stations is shown graphically in Figure 3-5 and summarized in Table 3-6.

Fig. 3-4. Mean Herbaceous Cover and Total Precipitation




Table 3-5. Herbaceous Phytomass for 1995

DATE	SITE	PLOT	WT.(g)	WT./m ²	DATE	SITE	PLOT	WT.(g)	WT./m ²
05/25	GO1	35-3	32.0	319.8	05/25	GO2	45-9	29.3	293.3
05/25	GO1	2-6	17.5	175.2	05/25	GO2	29-4	15.1	151.4
05/25	GO1	31-7	16.7	167.1	05/25	GO2	2-6	17.2	171.6
05/25	GO1	45-7	29.1	291.4	05/25	GO2	31-7	12.9	129.1
05/25	GO1	29-4	8.9	89.1	05/25	GO2	35-3	12.7	127.0
		AVG	20.9	208.5			AVG	17.4*	174.5
		STD	8.5	85.3			STD	6.2	61.6

DATE	SITE	PLOT	WT.(g)	WT./m ²	DATE	SITE	PLOT	WT.(g)	WT./m ²
05/18	GO3	31-7	0.4	3.9	05/15	GO4	29-4	2.6	26.0
05/18	GO3	2-6	1.1	11.4	05/15	GO4	2-6	8.7	86.6
05/18	GO3	29-4	2.5	24.9	05/15	GO4	25-3	7.9	78.8
05/18	GO3	45-9	4.7	46.9	05/15	GO4	31-7	11.1	111.2
05/18	GO3	35-3	8.0	80.2	05/15	GO4	45-9	2.5	24.9
		AVG	3.3	33.5			AVG	6.6	65.5
		STD	2.8	27.6			STD	3.4	34.4

DATE	SITE	PLOT	WT.(g)	WT./m ²	DATE	SITE	PLOT	WT.(g)	WT./m ²
05/16	GO5	29-4	1.4	13.6	05/17	GO6	31-7	7.2	71.5
05/16	GO5	45-9	2.4	23.9	05/17	GO6	29-4	13.4	134.0
05/16	GO5	2-6	7.1	70.9	05/17	GO6	35-3	8.6	86.1
05/16	GO5	35-6	4.7	47.3	05/17	GO6	45-9	3.1	30.6
05/16	GO5	31-7	2.2	21.9	05/17	GO6	2-6	14.0	139.7
		AVG	3.6	35.5			AVG	9.2	92.4
		STD	2.1	20.9			STD	4.1	40.7

DATE	SITE	PLOT	WT.(g)	WT./m ²	DATE	SITE	PLOT	WT.(g)	WT./m ²
05/18	GO7	35-3	9.3	16.3	05/25	GO8	45-9	22.4	224.4
05/18	GO7	31-7	5.7	27.9	05/25	GO8	29-4	10.2	102.2
05/18	GO7	29-4	2.4	50.9	05/25	GO8	35-3	16.8	168.2
05/18	GO7	45-9	26.0	46.1	05/25	GO8	2-6	8.6	85.9
05/18	GO7	2-6	10.1	66.4	05/25	GO8	31-7	11.6	115.6
		AVG	10.7	41.5			AVG	13.9	139.3
		STD	.1	17.6			STD	5.1	50.7

DATE	SITE	PLOT	WT.(g)	WT./m ²	DATE	SITE	PLOT	WT.(g)	WT./m ²
05/25	SO1	2-6	12.3	123.0	05/26	SO2	2-6	6.5	65.2
05/25	SO1	35-3	5.2	51.9	05/26	SO2	29-4	6.4	63.8
05/25	SO1	45-9	2.6	26.0	05/26	SO2	31-7	14.9	149.3
05/25	SO1	31-7	5.0	49.7	05/26	SO2	35-3	6.8	68.4
05/25	SO1	29-4	13.2	131.9	05/26	SO2	45-9	7.7	76.5
		AVG	7.7	76.5			AVG	8.5	84.6
		STD	4.3	42.7			STD	3.3	32.6

DATE	SITE	PLOT	WT.(g)	WT./m ²	DATE	SITE	PLOT	WT.(g)	WT./m ²
05/12	SO3	2-6	4.8	47.5	05/26	SO4	35-3	59.9	599.4
05/12	SO3	31-3	4.7	47.0	05/26	SO4	31-7	15.3	153.1
05/12	SO3	47-4	2.4	24.4	05/26	SO4	29-2	22.4	224.0
05/12	SO3	41-5	2.0	20.3	05/26	SO5	45-9	13.2	132.1
05/12	SO3	19-7	13.0	130.0	05/26	SO4	2-6	11.2	111.8
		AVG	5.4	53.8			AVG	24.4	244.1
		STD	4.0	39.7			STD	18.2	181.6

DATE	SITE	PLOT	WT.(g)	WT./m ²	DATE	SITE	PLOT	WT.(g)	WT./m ²
05/16	SO5	2-6	8.7	87.1	05/17	SO6	31-7	7.7	77.0
05/16	SO5	29-4	6.1	61.3	05/17	SO6	29-4	6.9	69.2
05/16	SO5	35-3	2.7	27.0	05/17	SO6	35-3	5.6	56.4
05/16	SO5	31-7	31.3	312.6	05/17	SO6	45-9	6.3	63.2
05/16	SO5	45-9	8.2	81.5	05/17	SO6	2-6	10.2	101.7
		AVG	11.4	113.9			AVG	7.4	73.5
		STD	10.4	101.6			STD	1.6	15.6

DATE	SITE	PLOT	WT.(g)	WT./m ²
05/08	SO7	45-9	18.0	180.1
05/08	SO7	35-3	2.5	25.2
05/08	SO7	2-6	24.5	245.1
05/08	SO7	29-4	29.4	294.3
05/08	SO7	31-7	32.8	327.9
		AVG	21.5	214.5
		STD	10.7	106.9

Phytomass Summary

MEAN GO1-GO8	107.0 Grams/sq. meter
MEAN SO1-SO7	123.0 Grams/sq. meter




Table 3-6 presents mean phytomass values for each station in each year since 1975.

Table 3-6. Comparison of Herbaceous Phytomass (g/m²) for 1975 through 1995

YEAR	GO1	GO2	GO3	GO4	GO5	GO6	GO7	GO8	SO1	SO2	SO3	SO4	SO5	SO6	SO7
1975	359	302	--	--	--	--	--	--	126	144	88	--	--	--	--
1976	108	258	--	--	--	--	--	--	137	98	177	--	--	--	--
1977	21	11	--	--	--	--	--	--	4	7	7	--	--	--	--
1978	166	162	--	--	--	--	--	--	173	128	115	--	--	--	--
1979	64	37	--	--	--	--	--	--	21	28	16	--	--	--	--
1980	160	68	53	79	--	--	--	--	36	63	43	78	71	--	--
1981	200	255	261	159	--	--	--	--	180	115	31	52	81	--	--
1982	90	60	62	113	--	--	--	--	98	24	22	39	184	--	--
1983	77	137	64	82	--	--	--	--	171	232	54	68	136	--	--
1984	94	116	133	67	--	--	--	--	104	57	95	93	43	--	--
1985	70	27	12	37	--	--	--	--	5	1	27	11	61	--	--
1986	50	61	32	35	--	--	--	--	35	112	25	176	42	--	--
1987	83	77	134	90	--	--	--	--	62	144	48	108	145	--	--
1988	34	14	16	61	--	--	--	--	59	73	15	24	19	--	--
1989	173.3	65.7	105.1	49.5	43.2	61	113.1	112.3	53.9	72.8	67	39.8	103.7	72.7	149.5
1990	13.6	4.1	64	73.2	36.8	39.8	29.1	10	32.8	78.3	28.2	30.9	43.4	34	6.1
1991	87.7	97.2	161.6	67.6	171.8	101.4	168.4	137.3	225.1	58.2	87.6	185.2	111.3	225.1	226
1992	142.4	109.4	82.7	60	54.4	49.4	181.4	74.3	49.2	147.5	90.7	80.3	110.3	101.3	187.3
1993	146	156.6	70.3	109.8	75.3	162	150.7	100.3	80.2	84.1	91.7	261.4	173.1	93.5	330.3
1994	45.7	48.8	49.2	15.6	13.2	60.1	41.5	44.1	27.3	13.4	20.5	19.8	60.8	7	19.1
1995	208.5	174.5	33.5	65.5	35.5	92.4	106.9	139.3	76.5	84.6	53.8	244.1	113.9	73.5	214.5

Fig. 3-5. Phytomass at Grassland and Shrub Stations for 1975 through 1995






3.3.3 Soil Chemistry

In comparison to previous years data, there has been no significant change in soil chemistry for the fifteen sampling stations. The following Table (3-8) is a summation of soil chemistry for 1995.

Table 3-8 Summary of Soil Chemistry for 1995

	pH	Conductivity μS/cm	Sulfate μg/gm	Chloride μg/gm	Copper μg/gm	Zinc μg/gm	Sodium μg/gm	Bicarbonate (meq/HCO ₃ /gm)
GO1	6.91	29.9	1.01	0.205	8.9	46.7	0.047	0.0011
GO2	7.21	36.3	0.94	0.330	10.9	53.7	0.053	0.0010
GO3	7.13	71.8	6.43	0.606	9.0	46.4	0.046	0.0018
GO4	6.84	15.4	1.34	0.160	8.1	46.4	0.053	0.0005
GO5	6.94	19.2	0.86	0.215	7.8	38.3	0.043	0.0007
GO6	6.85	16.6	0.63	0.163	8.1	43.1	0.054	0.0005
GO7	7.08	61.1	0.83	0.205	12.7	47.6	0.056	0.0014
GO8	7.11	28.1	0.85	0.247	12.8	42.8	0.047	0.0009
SO1	7.14	33.7	1.15	0.277	9.9	33.5	0.049	0.0010
SO2	7.62	19.1	0.33	0.196	3.1	15.2	0.032	0.0010
SO3	6.22	131	2.01	0.528	10.0	53.9	0.057	0.0006
SO4	7.166	18.6	0.32	0.209	9.0	43.4	0.051	0.0008
SO5	7.03	23.9	0.37	0.143	7.8	42.6	0.051	0.0007
SO6	7.91	43.7	0.39	0.205	8.7	41.3	0.042	0.0030
SO7	8.50	212	5.85	2.51	13.1	59.2	0.056	0.0054



4.0 REFERENCES

Battelle Pacific Northwest Laboratories. 1976. Aquatic ecological studies conducted near WNP-1, 2, and 4, September 1974 through September 1975. Supply System Columbia River ecology studies Vol. 2. Richland, WA.

Battelle Pacific Northwest Laboratories. 1977. Aquatic ecological studies near WNP-1, 2, and 4, October 1975 through February 1976. Supply System Columbia River ecology studies Vol. 3. Richland, WA.

Battelle Pacific Northwest Laboratories. 1978. Aquatic ecological studies near WNP-1, 2, and 4, March through December 1976. Supply System Columbia River ecology studies Vol. 4. Richland, WA.

Battelle Pacific Northwest Laboratories. 1979a. Aquatic ecological studies near WNP-1, 2, and 4, March through December 1977. Supply System Columbia River ecology studies Vol. 5. Richland, WA.

Battelle Pacific Northwest Laboratories. 1979b. Aquatic ecological studies near WNP-1, 2, and 4, January through August 1978. Supply System Columbia River ecology studies Vol. 6. Richland, WA.

Beak Consultants, Inc. 1980. Aquatic ecological studies near WNP-1, 2, and 4, August 1978 through March 1980. Supply System Columbia River ecology studies Vol. 7. Portland, OR.

Beak Consultants, Inc. 1981. Terrestrial monitoring studies near WNP-1, 2, and 4, May through December 1980. Portland, OR.

Beak Consultants, Inc. 1982a. Terrestrial monitoring studies near WNP-1, 2, and 4, May through December 1981. Portland, OR.

Beak Consultants, Inc. 1982b. Terrestrial monitoring studies near WNP-1, 2, and 4, May through August 1982. Portland, OR.

Black, C.A. et al., 1965. Methods of Soil Analysis. Academic Press, Inc. New York, New York.

Daubenmire, R. 1968. Plant Communities. Harper and Row, New York, New York.

Davis, W. III and T.E. Northstrom. 1987. Review of the environmental monitoring program for WNP-2 with recommendations for design of continuing studies. Washington Public Power Supply System. Richland, WA.

4.0 REFERENCES (Continued)

Droppo, J.G., C.E. Hane and R.K. Woodruff. November 1976. Atmospheric Effects of Circular Mechanical Draft Cooling Towers at Washington Public Power Supply System Nuclear Power Plant Number Two.

Environmental Protection Agency. August 1978. Quality Assurance Guidelines for Biological Testing, EPA/600/4-78/043.

Environmental Protection Agency. 1991. Methods for the Determination of Metals in Environmental Samples. EPA/600/4-91-010.

Environmental Protection Agency. September 1991. Methods for Measuring the Acute Toxicity of Effluents to Freshwater and Marine Organisms. EPA/600/4-85/013.

Gilman, Lee B. 1989. Microwave Sample Preparation. CEM Corporation.

Hamilton, M.A., R.C. Russo, and R.V. Thurston. 1977. Trimmed Spearman-Kärber Method for Estimating Median Lethal Concentrations in Toxicity Bioassays. Environ. Sci. Technol. 11(7) : 714-719; Correction 12(4) : 417 (1978).

Klemmedson, J.O. and J.G. Smith, 1964. Cheat Grass (*Bromus tectorum* L.) Bot. Rev. 30; 226-262.

Mudge, J.E., T.B. Stables and W. Davis III. 1982. Technical review of the aquatic monitoring program of WNP-2. Washington Public Power Supply System. Richland, WA.

Northstrom, T.E., J.L. Hickam and T.B. Stables. 1984. Terrestrial monitoring studies for 1983. Washington Public Power Supply System. Richland, WA.

Rickard, W.H. and K.A. Gano. 1976. Terrestrial ecology studies in the vicinity of Washington Public Power Supply system Nuclear Power Projects 1 and 4. Progress report for the period July 1974 to June 1975. Battelle Pacific Northwest Laboratories. Richland, WA.

Rickard, W.H. and K.A. Gano. 1977. Terrestrial ecology studies in the vicinity of Washington Public Power Supply System Nuclear Power Projects 1 and 4. Progress report for 1976. Battelle Pacific Northwest Laboratories. Richland, WA.

Rickard, W.H. and K.A. Gano. 1979a. Terrestrial ecology studies in the vicinity of Washington Public Power Supply System Nuclear Power Projects 1 and 4. Progress report for 1977. Battelle Pacific Northwest Laboratories, Richland, WA.

4.0 REFERENCES (Continued)

Rickard, W.H. and K.A. Gano. 1979b. Terrestrial ecology studies in the vicinity of Washington Public Power Supply System Nuclear Power Projects 1 and 4. Progress report for 1978. Battelle Pacific Northwest Laboratories, Richland, WA.

Schleder, L.S. 1982. Preoperational animal studies near WNP-1, 2, and 4. Annual report for 1981. Washington Public Power Supply System, Richland, WA.

Schleder, L.S. 1983. Preoperational animal studies near WNP-1, 2, and 4. Annual report for 1982. Washington Public Power Supply System, Richland, WA.

Schleder, L.S. 1984. Preoperational animal studies near WNP-1, 2, and 4. Annual report for 1983. Washington Public Power Supply System, Richland, WA.

Shipley, B.L. et al., 1980. NUREG/CR1231. Remote sensing for detection and monitoring of salt stress on vegetation: Evaluation and guidelines. Final report. September 1976-March 1979. Nuclear Regulatory Commission. Washington, DC.

Standard Methods for the Examination of Water and Wastewater. 1989. 18th Edition, APHA, AWWA, WPCF, Washington, D.C., 1989.

Standard Methods for the Examination of Water and Wastewater. 1995. 19th Edition, APHA, AWWA, WEF, Washington, D.C., 1995.

Washington Department of Ecology. 1992. Water Quality Standards for Surface Waters of the State of Washington (WAC 173-201A). Water Quality Planning Office of Water Programs. Olympia, WA.

Washington Public Power Supply System. Operational Ecological Monitoring Program for Nuclear Plant 2. Annual Reports for 1985-1994. Richland, WA.

