

PUBLIC SUBMISSION

As of: 12/7/17 10:44 AM Received: December 01, 2017 Status: Pending_Post Tracking No. 1k1-903s-zz8v Comments Due: January 08, 2018 Submission Type: Web
--

Docket: NRC-2017-0215

Yttrium-90 Microsphere Brachytherapy Sources and Devices Therasphere and SIR-Spheres

Comment On: NRC-2017-0215-0001Yttrium-90 Microsphere Brachytherapy Sources and Devices TheraSphere and SIR-Spheres;
Draft Guidance for Comment**Document:** NRC-2017-0215-DRAFT-0086

Comment on FR Doc # 2017-24129

Submitter Information

Name: Laurence Donahue**Address:**Aurora Health Care Medical Group, 2900 W. Oklahoma Avenue
Milwaukee, WI, 53215**Email:** laurence.donahue@aurora.org

11/7/2017

82 FR 51655

85

General Comment

I am writing as a current interventional radiologist and a member of The Society of Interventional Radiology (SIR), a physician association comprised of over 6,100 members representing the majority of practicing interventional radiologists in the United States. I am corresponding to voice my strong opposition to the proposed changes to the "Yttrium-90 Microsphere Brachytherapy Sources and Devices TheraSphere and SIR-Spheres Licensing Guidance" that would eliminate vendor involvement from the Interventional Radiology pathway to Authorized User (AU) status.

The current process, known as Pathway 2, has been in place since the concept evolved that Interventional Radiologists are the natural AU's of these devices. For these procedure, Interventional Radiologists:

Perform required dosimetry to deliver appropriate activity to the patient

SUNSI Review Complete

Template = ADM - 013

E-RIDS= ADM-03

Add= L. Dimerick (2017) 12/07/2017

Oversee and ensure appropriate Yttrium-90 (Y90) device handling and preparation
Directly administer the therapy to the patient
Coordinate the longitudinal care of patients following these procedures

Interventional Radiologists are the only Authorized Users capable of performing all four steps above. The existing collaboration between physicians and industry helps ensure safe and comprehensive training in the use of Y90 devices.

The existing guidelines have been tremendously successful while maintaining impeccable safety. Manufacturer And User facility Device Experience (MAUDE) reports have remained 10/year for both devices since 2013. The majority of the MAUDE reports focus on procedural complication and treatment toxicities seen with all types of hepatic embolization, not specific to the Y90 devices. It seems statistically implausible that reduced vendor involvement will result in a measurable improvement in safety.

Proposed changes to the current arrangement, in which physicians and industry work closely together to ensure the appropriate training of interventional radiologists in the safe use of these devices will make it exceedingly difficult for Interventional Radiologists developing a clinical practice in radioembolization. Without the current direct training provided offsite by the device vendors, physician training will have to be solely performed by direct proctoring. Securing physician proctors is a challenge and can result in the delay in care, impacting cancer outcomes; physicians have limited time and availability away from their own clinical practices. Placing additional responsibilities on physician proctors may also have the untoward effect of limiting access to care, particularly for programs in underserved areas. The unanticipated consequence of the proposed changes is that training Interventional Radiologists in the safe and effective use of these devices will suffer greatly and patient access to care will diminish.

In summary, Interventional Radiologists deliver high quality minimally invasive care via imaging guidance, employing a variety of technologies. Training with other devices, such as aortic stent grafts, spinal augmentation devices, and atherectomy tools frequently involves a combination of vendor and physician collaboration. These relationships are a supplement to core training in hepatic embolization that is accomplished in fellowship. However, fine details regarding all devices may not be included in all programs. The existing NRC guidelines have facilitated training Interventional Radiologists in the safe and effective use of the Y90 devices, benefiting patients, physicians, and the government. There is no evidence of a need for change to the current NRC guidelines.

Regards,

Laurence Donahue, M.D.
Interventional Radiology

Aurora Health Care Medical Group
2900 W. Oklahoma Avenue
Milwaukee, WI 53215
laurence.donahue@aurora.org