

ENVIRONMENTAL PROTECTION PLAN

FOR

SALEM GENERATING STATION

UNIT NOS. 1 AND 2

PUBLIC SERVICE ELECTRIC AND GAS COMPANY

DOCKET NOS. 50-272 AND 50-311

8511250343 851119
PDR ADOCK 05000272
P PDR

ENVIRONMENTAL PROTECTION PLAN

TABLE OF CONTENTS

| | | Page |
|-----|--|-------|
| 1.0 | OBJECTIVES | 1 - 1 |
| 2.0 | ENVIRONMENTAL PROTECTION ISSUES | 2 - 1 |
| 3.0 | CONSISTENCY REQUIREMENTS | 3 - 1 |
| 3.1 | PLANT DESIGN AND OPERATION | 3 - 1 |
| 3.2 | REPORTING RELATED TO THE NJPDES PERMIT AND STATE CERTIFICATION (PURSUANT TO SECTION 401 OF THE CLEAN WATER ACT) | 3 - 3 |
| 3.3 | CHANGES REQUIRED FOR COMPLIANCE WITH OTHER ENVIRONMENTAL REGULATIONS | 3 - 4 |
| 3.4 | CHANGES IN ENVIRONMENTAL PROTECTION PLAN | 3 - 5 |
| 4.0 | UNUSUAL OR IMPORTANT ENVIRONMENTAL EVENTS | 4 - 1 |
| 5.0 | REFERENCES | 5 - 1 |

1.0 OBJECTIVES

This Environmental Protection Plan (EPP) provides for protection of the environment during operation of the Salem Generating Station. The principal objectives of this EPP are to:

1. Verify that the plant is operated in an environmentally acceptable manner, as established by the Final Environmental Statement (FES) and other NRC environmental impact assessments.
2. Coordinate NRC requirements and maintain consistency with other Federal, State and local requirements for environmental protection.
3. Keep NRC informed of the environmental effects of facility operation and of actions taken to control those effects.

Environmental concerns identified in the FES which relate to water quality matters are regulated by way of the licensee's NJPDES permit.

2.0 ENVIRONMENTAL PROTECTION ISSUES

In the FES, dated April 1973, the staff considered the environmental impacts associated with the operation of the Salem Generating Station. Certain environmental issues were identified which required study or license conditions to resolve environmental concerns and to assure adequate protection of the environment. The Appendix B Environmental Technical Specifications (ETS) issued with the operating license included discharge restrictions and monitoring programs related to:

1. Protection of the aquatic environment by limiting the thermal characteristics of the discharge
2. Protection of the aquatic environment from biocide used in plant operations
3. Protection of the aquatic environment from suspended solids and changes in pH in releases from the non-radioactive liquid waste disposal system
4. Surveillance programs for dissolved gases, suspended solids, chemical releases, and the general aquatic ecological surveys to establish impact of plant operation on the biotic environment

5. Special studies of experimental entrainment and to document the thermal plume and intake velocity
6. Surveillance programs to determine the impact of plant operation on the nesting of diamondback terrapin, osprey and southern bald eagle in the plant vicinity.

Aquatic requirements and programs 1 through 5 above were deleted by License Amendments 51 (Unit 1) and 18 (Unit 2), dated March 14, 1983 and March 11, 1983, respectively. These issues are now addressed by the effluent limitations and monitoring requirements contained in the effective NJPDES Permit No. NJ0005622 under the jurisdiction of the State of New Jersey. The NRC will rely on this agency for regulation of matters involving water quality and aquatic biota. The NRC requirements for the terrestrial issue, 6 above, have been completed and are terminated by the submittal of Reference 1 in conjunction with this EPP.

3.0 CONSISTENCY REQUIREMENTS

3.1 PLANT DESIGN AND OPERATION

The licensee may make changes in station design or operation or perform tests or experiments affecting the environment provided such changes, tests or experiments do not involve an unreviewed environmental question and do not involve a change in this EPP*. Changes in plant design or operation or performance of tests or experiments which do not affect the environment are not subject to the requirements of this EPP. Activities governed by Section 3.3 are not subject to the requirements of this section.

Before engaging in activities which may significantly affect the environment, the licensee shall prepare and record an environmental evaluation of such activity. Activities are excluded from this requirement if all measurable non-radiological effects are confined to the on-site areas previously disturbed during site preparation and plant construction. When the evaluation indicates that such activity

* This provision does not relieve the licensee of the requirements of 10 CFR 50.59.

involves an unreviewed environmental question, the licensee shall provide a written evaluation of such activities and obtain prior approval from the NRC. When such activity involves a change in the EPP, such activity and change to the EPP may be implemented only after NRC approval as set forth in Section 3.4 of this EPP.

A proposed change, test or experiment shall be deemed to involve an unreviewed environmental question if it concerns (a) a matter which may result in a significant increase in any adverse environmental impact previously evaluated in the FES, supplements to the FES, environmental impact appraisals, or in any decisions of the Atomic Safety and Licensing Board; or (b) a significant change in effluents or power level; or (c) a matter not previously reviewed and evaluated in the documents specified in (a) of this Subsection, which may have a significant adverse environmental impact.

The licensee shall maintain records of changes in facility design or operation and of tests and experiments carried out pursuant to this Subsection. These records shall include a written evaluation which provides bases for the determination that the change, test, or experiment does not involve an unreviewed environmental question or constitute a decrease in the effectiveness of this EPP to meet the objectives specified in Section 1.0.

3.2 REPORTING RELATED TO THE NJPDES PERMIT AND STATE CERTIFICATION (PURSUANT TO SECTION 401 OF THE CLEAN WATER ACT).

Violations of the NJPDES Permit Conditions shall be reported to the NRC by submittal of copies of the reports required by the NJPDES Permit.

The licensee shall provide the NRC with a copy of any 316(a) or (b) studies and/or related documentation at the same time it is submitted to the permitting agency.

Changes to, or renewals of, the NJPDES Permit or the State certification shall be reported to the NRC within 30 days following the date the change or renewal is approved. If a permit or certification, in part or in its entirety, is appealed and stayed, the NRC shall be notified within 30 days following the date the stay is granted.

The NRC shall be notified of changes to the effective NJPDES Permit proposed by the licensee by submittal of a copy of the proposed change at the same time it is submitted to the permitting agency. The licensee shall provide the NRC with a copy of the application for renewal of the NJPDES Permit at the same time the application is submitted to the permitting agency.

3.3 CHANGES REQUIRED FOR COMPLIANCE WITH OTHER ENVIRONMENTAL REGULATIONS.

Changes in plant design or operation and performance of tests or experiments which are required to achieve compliance with other Federal, State or local environmental regulations are not subject to the requirements of this Section.

3.4 CHANGES IN ENVIRONMENTAL PROTECTION PLAN

Requests for changes in the EPP shall include an assessment of the environmental impact of the proposed change and a supporting justification. Implementation of such changes in the EPP shall not commence prior to NRC written approval of the proposed changes.

4.0 UNUSUAL OR IMPORTANT ENVIRONMENTAL EVENTS

Any occurrence of an unusual or important event that indicates, or could result in, significant environmental impact causally related to station operation shall be recorded and promptly reported to the NRC within 24 hours followed by a written report within 30 days. No routine monitoring programs are required to implement this condition.

The written report shall (a) describe, analyze and evaluate the event, including the extent and magnitude of the impact and plant operating characteristics, (b) describe the probable cause of the event, (c) indicate the action taken to correct the reported event, (d) indicate the corrective action taken to preclude repetition of the event and to prevent similar occurrences involving similar components or systems, and (e) indicate the agencies notified and their preliminary responses.

Events reportable under this subsection which also require reports to other Federal, State or local agencies shall be reported in accordance with those reporting requirements in lieu of the requirements of this subsection. The NRC shall be provided a copy of such report at the same time it is submitted to the other agency.

The following are examples of potentially significant environmental events: excessive bird impaction events; onsite plant or animal disease outbreaks; mortality or unusual occurrence of any species protected by the Endangered Species Act of 1973; unusual fish kills; and an increase in nuisance organisms or conditions.

5.0 REFERENCES

1. An Environmental Monitoring Program 1974 - 1984 on
Diamondback Terrapin Nesting and Osprey Nesting/Bald Eagle
Occurrence in the Vicinity of Artificial Island,
V. J. Schuler Associates, Inc., March 28, 1985.

An Environmental Monitoring Program
1974-1984

on
Diamondback Terrapin Nesting
and
Osprey Nesting/Bald Eagle Occurrence
In the Vicinity of Artificial Island


a Summary

for

Public Service Electric and Gas Co.
Nuclear Department
Hancocks Bridge, NJ 08038

by

V.J. Schuler Associates, Inc.
100 South Cass Street
Middletown, Delaware 19709


Victor J. Schuler, President

8511250347 851119
PDR ADCK 05000272
P PDR

March 28, 1985

TABLE OF CONTENTS

| | Page |
|---|------|
| 1.0 INTRODUCTION..... | 1 |
| 2.0 DIAMONDBACK TERRAPIN NESTING..... | 2 |
| 2.1 STUDY SCOPE..... | 2 |
| 2.2 STUDY AREA..... | 3 |
| 2.3 MATERIALS AND METHODS..... | 4 |
| 2.4 DATA ANALYSIS..... | 5 |
| 2.5 DISCUSSION..... | 5 |
| 2.5.1 Nesting Period..... | 5 |
| 2.5.2 Nesting Activity..... | 6 |
| 2.5.3 Predators..... | 7 |
| 2.5.4 Age and Size of Nesting Females..... | 8 |
| 2.6 OVERVIEW..... | 8 |
| 3.0 OSPREY NESTING/BALD EAGLE OCCURRENCE..... | 10 |
| 3.1 STUDY SCOPE..... | 10 |
| 3.2 STUDY AREA..... | 11 |
| 3.3 MATERIALS AND METHODS..... | 11 |
| 3.4 DISCUSSION..... | 11 |
| 3.4.1 Osprey..... | 11 |
| 3.4.2 Bald Eagle..... | 13 |
| 3.5 OVERVIEW..... | 14 |
| 4.0 LITERATURE CITED..... | 16 |

LIST OF TABLES

| | |
|---|----|
| Table 1. Annual summary data on observed nesting, nest depredation, and hatchlings of diamondback terrapin observed at a beach north of Liston Point, DE, 1975-1984..... | 19 |
| Table 2. Annual summary data on observed nesting, nest depredation, and hatchlings of diamondback terrapin observed at a beach north of Sunken Ship Cove, NJ, 1975-1984..... | 20 |
| Table 3. Annual summary data on observed nesting, nest depredation, and hatchlings of diamondback terrapin observed at a beach north of Hope Creek, NJ, 1975-1984..... | 21 |
| Table 4. Summary of osprey nesting activity within 16 km of southern Artificial Island: 1974-1984..... | 22 |

TABLE OF CONTENTS
CONTINUED

LIST OF FIGURES

| | Page |
|--|------|
| Figure 1. Diamondback terrapin study sites, Salem Generating Station Tech Spec Study, 1975-1984..... | 24 |
| Figure 2. Number of adult diamondback terrapin and tracks observed per sampling date during 1979 through 1984 at a beach on the Delaware River near Liston Point, DE..... | 25 |
| Figure 3. Number of adult diamondback terrapin and tracks observed per sampling date during 1979 through 1984 at a beach on the Delaware River near Sunken Ship Cove, NJ..... | 26 |
| Figure 4. Number of adult diamondback terrapin and tracks observed per sampling date during 1979 through 1984 at a beach on the Delaware River near Hope Creek, NJ..... | 27 |
| Figure 5. Annual mean number of observed adult diamondback terrapin and tracks at Liston Point, DE, and Hope Creek and Sunken Ship Cove, NJ, 1975-1984..... | 28 |
| Figure 6. Nesting sites (O), historical and present, observed during SNGS Osprey Study, 1974-1984. Dashed (--) line is new transmission line nearing completion in 1984..... | 29 |
| Figure 7. Tower locations on transmission lines from SNGS; Osprey study, 1974-1984. O indicates tower used by nesting osprey..... | 30 |
| Figure 8. Northern extension of Salem-Keeney line, Osprey study, 1974-1984. O indicates tower used by nesting osprey..... | 31 |

SECTION 1.0
INTRODUCTION

Annually from 1972 through 1984, various aspects of the terrestrial ecology of Artificial Island and vicinity have been studied as part of the Salem Nuclear Generating Station (SNGS) Environmental Technical Specifications (Tech Specs), as required by the U.S. Nuclear Regulatory Commission (NRC). Findings have been presented in Annual Progress Reports (Schuler, 1974-1977; PSE&G, 1978-1984), and summary reports by Hardin (1980) covering 1972-1978 and PSE&G (1983) covering 1974-1981.

Early studies generated a perception of the pre SNGS-operation local terrestrial community and representative elements. Later studies traced variously-selected elements in the post SNGS-operation environment. Vegetative associations were studied during 1972-1974, small-mammal populations during 1972-1973, and bird seasonal/migratory occurrence during 1972-1979.

The two topic studies of this summary report, osprey nesting/bald eagle occurrence and diamondback terrapin nesting, were begun in 1974 and 1975, respectively, and continued through 1984. This report integrates the annual information collected over the decade of observations. It presents general ranges, means, and trends reflected in the data and attempts to describe the ecological impact of SNGS on these species.

SECTION 2.0 DIAMONDBACK TERRAPIN NESTING

Perhaps the most common reptile in the river and marshes of the Delaware Bay is the northern diamondback terrapin, Malaclemys terrapin terrapin. The literature generally describes the environment of this subspecies as the salt and brackish coastal waters from Cape Cod to Cape Hatteras. In the early part of this century, market-for-table demand was high and populations were heavily exploited, some to the point of serious regulatory concern. As a result, capture seasons were closed, and possession was prohibited. This protection allowed a population recovery and a coincident drying-up of market interest, and populations have again become well established.

Hurd et al. (1979), in a two-year study of a population in a Delaware salt-marsh designed to reflect on seasonal population phenomena related to ecology, described a terrapin density of 1.8 individuals per linear meter of tidal creek, which they described as a large population. They also commented on the paucity of information concerning population dynamics and ecological relationships of diamondback terrapin in nature. This present study provides some useful insight into the biology of the local terrapin population and its utilization of local nesting areas.

2.1 STUDY SCOPE

Study of diamondback terrapin focused on reproduction-related parameters which could be monitored at local nesting-beaches. Observations of nesting activity and effort, age of nesting females, nest activity and hatching success, and predation were made, typically from June-November, in all years. From 1975-1982, three local beaches which had been determined to support nesting were monitored. Two were on the New Jersey shore and proximal to Salem; the other was established in Delaware to possibly reflect behavioral differences at east- versus west-shore beaches. Data through 1982 failed to evidence such differences, and in 1983 effort was restricted to the one beach in Delaware and the up-river site in New Jersey.

2.2 STUDY AREA

Observations through 1984 were made just north of Liston Point, Delaware and at Sunken Ship Cove and, through 1982, at Hope Creek in New Jersey (Figure 1).

Sunken Ship Cove is at the southeastern end of Artificial Island. The beach is partially bounded by a breakwater and lies half within the cove and half east of the cove. The area monitored is 213 m (700 ft) long and from 15 to 38 m (49-125 ft) wide. Primary vegetation consists of a dense stand of saltmeadow cordgrass (Spartina patens) with reed canary grass, Phalaris arundinacea; sea rocket, Cakile edentula; and wild radish, Raphanus raphanistrum, occurring in clumps.

Hope Creek beach is some 209 m (680 ft) southeast of the mouth of Hope Creek. The site is 213 m (700 ft) long and 3 to 5 m (10-16 ft) wide. Vegetation originally included a 3 m (10 ft) wide stand of saltmarsh cordgrass in the intertidal zone, with common reed, Phragmites australis (formerly P. communis); saltmeadow cordgrass, and groundsel bush, Baccaris halimifolia, occurring above mean high tide. Behind this strip of beach is an intermediate type tidal marsh. The site has become almost entirely covered with common reed during the period of study.

The Liston Point site is 397 m (1,300 ft) long and from 20 to 24 m (60-80 ft) wide. Primary vegetation includes saltmeadow cordgrass and American beachgrass, Ammophila brenligulata, in sparse to dense stands, with marsh elder, Iva frutescens, and sedge, Cyperus sp., occurring in clumps. This is located behind a 7 to 12 m (20-40 ft) wide shoreline strip of sand. An intertidal stand from 3 to 5 m (10-16 ft) wide of mostly saltmarsh cordgrass occurs on the southern half of the site.

Liston Point has the highest elevation and Hope Creek beach the lowest of the three sites. Sections of the Hope Creek site are occasionally inundated during storms.

The amount of human disturbance varies greatly at the three sites. Sunken Ship Cove is used for fishing, swimming, and picnicing. Hope Creek is rather isolated and probably visited only by present-study personnel during sampling. Liston Point is used occasionally for recreation by local inhabitants.

2.3 MATERIALS AND METHODS

The study sites were searched during daylight, typically from June through November. Weekly searches for evidence of nesting were made in early June. After first evidence of nesting, beaches were monitored several times (2-5) a week through July. Searches for depredated nests and emerging hatchlings were made several times a week from August through September and occasionally (if weather stayed warm) into November. Weather and tide occasionally prevented visiting all beaches on the same day. Each visit consisted of walking the beach and counting turtles, crawl tracks (incoming only), depredated nests, and eggs.

Effort was taken to minimize disturbance of nesting terrapins. Wherever possible they were not disturbed until after nesting; females typically attempted to leave the area when they sensed the observers. It is probable that disturbance from beaching the boat and subsequent monitoring activities interrupted turtles at the various stages of nesting.

Females were caught by hand and the length and width (mm) of the carapace and plastron were measured. A numbered spaghetti tag was placed in a hole drilled in either the eleventh marginal or one of the postcentral laminae after Porter (1972), with the hole location being part of a binary code which keyed to the tagging event. This location enables easy drilling and placement of the tag and offers minimal interference to the activities of the terrapin.

Hatchlings were enumerated from crawl tracks or by digging them out of nests. Young were returned to the point of capture.

The number(s) of the tag(s) attached, length and width measurements, general location of the nest, time, date, tidal stage, weather, number of turtles observed but not tagged, and number of terrapin tracks observed on the beaches were recorded. Tide data were taken from National Oceanic and Atmospheric Administration (NOAA) (1977) and measurements of cloud cover from NOAA (1972).

In all years except 1976 and 1977, nests that contained unbroken eggs were marked with a stake, and the number of eggs was recorded. These nests were located by following tracks, finding females on the nests, and by random search. Depredated nests were counted and the number of eggs destroyed at each nest estimated by counting egg shells in and near the nest. Scattered individual egg shells were not counted. All shell fragments were buried or removed from the study area after counting. In 1976 and 1977, nests were

covered with a wire enclosure to prevent depredation, and weekly measurements of temperature and soil moisture (g/100c) were taken.

2.4 DATA ANALYSIS

Intensity of nesting activity for each study site was annually estimated for the period from the first observed occurrence of nesting turtles or adult tracks to the last observed occurrence. A log (x+1) transformation was employed to allow for the occurrence of zero observed turtles and tracks in the data. Plots of the annual mean log (x+1) number of turtles and adult tracks for each year are presented for comparison.

2.5 DISCUSSION

Findings on the discrete behavioral, biologic and ecologic parameters monitored during the study are summarized categorically. Collectively, they characterize the schedules and relationships exhibited by local diamondback terrapin during their usage of the studied beaches.

2.5.1 Nesting Period

Nesting has regularly begun during early- to mid-June and continued through mid- to late July (Figures 2, 3, 4). The earliest date of observed nesting evidence was June 4, 1981 at Liston Point beach. Nesting at the different study sites has always commenced within a few days of each other, suggesting a synchronization in the local population. Nesting also appears cyclic in that there are typically two major peaks and perhaps a third to several lesser peaks during a season. This implies a hormonal synchrony, the existence of which is supported by a general correlation of nesting pattern with photoperiod, temperature, and even lunar stage. Photoperiod is suggested in first nesting evidence being annually observed about June 10, the time of earliest sunrise in the year. Burger and Montevecchi (1975) also observed first nesting on about June 10 at a site on the Atlantic coast of southern New Jersey. Earlier, Burger (1937) had stated that sexual behavior of turtles might be controlled by light. Temperature is suggested as a factor since in years when the period April through mid-June has

been relatively cool nesting started in mid-June but did not peak until warming had occurred. The actual temperature probably affects date of emergence from hibernation, subsequent mating, and timing as well as number of individuals involved in movement into the rookery. Lunar stage correlation might be evidenced by the apparent 14-day cycles within the nesting data curves in Figures 2, 3, and 4.

2.5.2 Nesting Activity

Nesting at the three study sites has followed a general pattern in which the Liston Point site has annually, based on the mean number of observed turtles and tracks per visit, been the most intensively used, and Sunken Ship Cove the least (Figure 5). Further, each site has evidenced a relatively similar increase or decrease in annual usage, i.e., population activity trends could be reasonably inferred from any one of the studied beaches. Liston Point, being the largest beach and relatively isolated, is explainably the most used. The Hope Creek beach has been physically diminished by erosion and encroaching marsh vegetation. The Sunken Ship Cove site, based on its size and appearance, might be expected to support more terrapin utilization were it not for the heavy usage by fishermen throughout the summer.

This pattern of usage is also reflected in numbers of nests, eggs, and hatchlings (Tables 1, 2, 3). Of course, these counts are conservative (low) and should be used only as relative indices for inter-beach comparisons and not for actual production or population-size inference. During 1975-1984, 3,741 nests were identified at Liston Point, 99 at Sunken Ship Cove, and 1,415 at Hope Creek. Observed nest depredation was greatest at Liston Point; of the 3,741 nests seen, only 247 had been undisturbed. Sunken Ship Cove evidenced the least; of 99 total nests, 43 had been undisturbed. The range of mean eggs per nest at the three sites was reasonably close; 8.25 at Liston Point, 7.15 at Sunken Ship Cove, and 6.12 at Hope Creek. Similar to the mean number eggs-per-nest patterns, the numbers of hatchlings represented in Column III in Tables 1, 2 and 3 are not as dissimilar as numbers of nests and eggs might suggest. Liston Point beach evidenced 37.8 x the number of total nests at Sunken Ship Cove, and 5.7 x the number of non-depredated nests and 6.12 x the number of non-depredated eggs. Yet, the ratio of hatchlings at the two sites was 1.83:1. Compared with Hope Creek beach, the ratio was somewhat higher at 2.92:1.

During monitoring of non-depredated nests, incubation took from as few as 49 days (in 1977) to as many as 100 days (in 1982). Typically, hatching occurred 65-75 days after eggs were laid. Carr (1952) reported that incubation periods for turtles normally range from 60-90 days, but are so strongly affected by temperature and humidity that no given species adheres very closely to a definite schedule. Spearman's coefficient of rank correlation on several year's data verified that length of incubation was negatively correlated with mean nest temperatures (i.e., as temperature decreases incubation time increases). In general, nests laid under vegetation had longer incubation periods, probably reflecting cooler temperatures due to shading. In the present study, soil moisture was not found to affect length of incubation of successful eggs. However, it did affect hatching success of fertile eggs. Unsuccessful embryo development (percent of embryos that died) was greater at the wetter nest sites. This mortality could reflect lower temperatures, decreased air availability, and perhaps fungal occurrence, that could accompany higher moisture levels.

Hatching typically began during the latter half of August, peaked during the next two weeks, decreased sharply during late September and occasionally continued at a low level into early October. Burger (1976) observed that young terrapin spent several days in the nest before emerging, perhaps as many as 11 days. In the present study in 1977, turtle nests were excavated and many fully-formed hatchlings were found, accompanying the unhatched eggs. We did not observe synchrony in either egg hatching or hatchling emergence. We did observe, as did Burger (1976), that most hatchlings emerge between 1200 and 1700 hrs, normally the warmest part of the day.

2.5.3 Predators

As described above, depredation of nests and predation on hatchlings was a significant statistic in this study (Tables 1, 2, 3). Most local common predators and scavengers likely exploit these early life stages. At Sunken Ship Cove, tracks of the Norway rat, Rattus norvegicus, and striped skunk, Mephitis mephitis, were occasionally observed. Tracks of mink, Mustela vison; Norway rat; common crow, Corvus brachyrhynchos; and, occasionally, muskrat, Ondatra zibethicus, were observed at Hope Creek beach. Mink; Norway rat; and raccoon, Procyon lotor, tracks were commonly observed at Liston Point. Track evidence indicated that mink; raccoon; Norway rat; crow; great black-backed gull, Larus marinus; and occasionally, great blue heron, Ardea herodias, and turkey vulture, Cathartes aura, also preyed on

hatchlings both in and out of the nests. Fox (whether red or gray is unknown) tracks were also seen at all locations.

2.5.4 Age and Size of Nesting Females

From 1975-1984, 380 nesting females were captured, examined and tagged. These were distributed as 202 at Liston Point, 175 at Hope Creek, and three at Sunken Ship Cove.

Mean plastron length was annually similar, typically 17.5-18 cm, as was plastron width at 12-14 cm. Carapace length was annually about 16.5 cm and width was 9-13 cm.

Age of captured specimens ranged from 5 to 20+ years. More than half had smooth shells, which Hildebrand (1932) stated may indicate age to perhaps 40+ years.

The turtle recapture pattern evidenced several factors. Only 25 specimens of the 380 tagged were recaptured, a recapture rate of only six percent. However, although population estimates were neither planned nor possible, the results do reflect on the parameters at which tagging was directed, namely, beach fidelity and, perhaps, growth information. All recaptures were at the beach of initial capture and tagging. Time to recapture ranged from 2 days to 5 years. Many recaptures were within days or weeks of tagging, showing a persistence to nesting or the act of re-nesting during the same season. Re-nesting was observed in cultured terrapin, from one to five nests per year, by Hildebrand (1932). The long-term recaptures are evidence of beach fidelity, a feature reported by Carr and Ogren (1960) and Carr and Carr (1972). Perceived change in physical size was slight; during a five-year interval one capture had increased in carapace length by only two percent, from 16.4 cm to 16.7 cm.

2.6 Overview

All observations on local diamondback terrapin suggest behavior, and response to environmental conditions, typical of the species and of a healthy biological population. During the near-decade of study, construction of SNGS Units 1 and 2 was completed and both units underwent power-level staging and reached 100 percent, or commercial, operation (Unit 1 on June, 1977 and Unit 2 on October 18, 1981). There is no evidence that operational levels or characteristics of SNGS have affected, in any way, the

activities or success of local diamondback terrapin. It is probable that the Artificial Island access road has indirectly had a negative effect on the degree of utilization of the Sunken Ship Cove beach as a nesting site. The road provides ready, and literally the only, land access to Sunken Ship Cove and the local Delaware River, and there is an established use pattern by fishermen, boaters and picnickers. This human recreational activity during the nesting period probably discourages or disrupts nesting behavior. However, it is unlikely that this very localized action has any substantive effect on the regional diamondback terrapin population.

SECTION 3.0

OSPREY NESTING/BALD EAGLE OCCURRENCE

The osprey, Pandion haliaetus, is a common summer-resident raptor in the study area, occurring annually between March and August during which time it breeds, nests, and rears its young. Nests built of sticks, reeds, and debris are constructed in natural and man-made structures including dead or dying trees, channel markers, and, increasingly in this area, electric transmission towers.

The species is listed as "endangered" in New Jersey (NJDEP, 1984). It had been federally classified as "status undetermined" (USDI, 1973) but has since been deleted from the Federal list. In New Jersey, prior to 1950 there were some 500 nesting pairs of osprey (Frier, 1982). However, chemical contamination (primarily DDT) of the environment and coincident loss of nesting sites caused a severe population reduction, and in 1974 there were only 50 known nesting pairs in the State (Frier, 1982). The cessation of DDT usage (in 1966) and, to a degree, the increase in nesting sites have encouraged a population recovery. This trend can be perceived in the local population discussed in this report.

The bald eagle, Haliaeetus leucacephalus, has historically wintered along major rivers and bays in New Jersey. Throughout much of its range the species has evidenced reduced reproductive success as a result of infertile or thin-shelled eggs, these being attributed to use of DDT and its occurrence in the food chain. The species is federally classified as "endangered". Frier (1982) listed one breeding pair remaining in New Jersey. Annually during 1974-1984, special note was taken of eagle sitings and reports in the course of all terrestrial studies. However, due to the absence of nesting activity in the area, no special study program or area was established.

3.1 STUDY SCOPE

Monitoring of osprey focused on nesting-related activities, behavior, and reproductive success. Numbers and locations of nests, vacant and occupied, and counts/estimates of eggs and number young fledged were the standard parameters. Monitoring was done each year, 1974-1984. Records were kept of bald eagle sightings and awareness of nesting activity was maintained.

3.2 STUDY AREA

Observations were made at historical and regularly-used nesting locations contained within the area shown in Figure 6. The most striking and dominant physical feature of the 232 km² region is the array of electric transmission lines, and the associated towers which support most of the local nesting. The region features a variety of habitats, e.g., bay, riverine, marsh, upland field, and wooded, and with availability of suitable nesting locations appears capable of supporting a local seasonal osprey population.

3.3 MATERIALS AND METHODS

Observations were made by boat and foot travel during 1974-1976, and from a PSE&G helicopter from 1975-1984. In 1974 and 1976, nests were closely inspected and exact counts of eggs, nestlings, and fledglings were recorded. These counts were possible when climbing-visits coincided with the feeding or other absence of the sitting female. In the mid-1970's, when helicopter became the prime observation mode, a reasonable distance (ca. 50 yards) from the nest was maintained to avoid frightening or otherwise disturbing the sitting birds. Observations were made with binoculars and data should be considered as semi-quantitative.

3.4 DISCUSSION

3.4.1 Osprey

During the decade of study, adult osprey have been annually recorded in the study area, as early as March 15 and as late as October 15. Soon after first sightings, activity at nesting sites was observed as the birds began nest building or refurbishment. Eggs were usually laid, and clutches were complete, by mid-April. Incubation takes about 28 days, and eggs hatched typically during mid- to late May. The young birds fledged by mid-July, and by mid-August most were independent of the nest. By mid-September, young and adults were leaving or had already left the study area for overwintering grounds in the West Indies and South America (Henny and Van Velzen, 1972).

Table 4 presents all data on nesting collected during the period of study. It shows the temporal and spatial expansion of site utilization and presents statistics and notations on osprey activity and success over the years.

Figure 6 shows nesting locations (historical and present) in non-PSE&G sites (e.g., nesting platforms, snag trees, transmission-line towers), and in the PSE&G transmission lines that emanate from SNGS. The locations of towers in PSE&G lines, with notations to those used by osprey, are shown in Figures 7 and 8. Most local nesting activity occurs in the PSE&G towers.

Table 4 does not list data collected in 1974 and 1976 during close-up inspection of nests and clutches. During these years, annual mean clutch size was 2.4 and 3.0 eggs, and mean hatching success was 33.3 and 44.4 percent. Mean success from nestling, or young, to fledgling stage was 75.0 and 87.5 percent. These levels approximate those described as generally good for a healthy population by Parnell and Walton (1972) in a discussion of osprey reproductive success in North Carolina. We have no local information on these parameters since 1976.

With completion of the Salem transmission towers in 1971, the number of local potential osprey nesting sites greatly increased. Commencing ca. 1975-1977 there has been an apparent shift in nesting activity from the old natural structures and man-made platforms to the transmission towers. In 1980, utilization of towers in the Salem-Keeney line was pronounced, and since 1981 utilization of the Salem-New Freedom North line has increased. The DP&L towers have been regularly used since 1981. Several towers, most notably tower 6/1 of the Salem-New Freedom North line, have accommodated coincident nests, although not all were active (breeding) nests. It would appear that the tower sites may offer some subtle attraction over the natural sites such as at Reedy Island or off the Smyrna River. The shift may also reflect the increasing human activity on the river vis a vis boating and fishing. The continued use of the Raccoon Ditch location invites speculation; it could reflect the site's relative isolation, or perhaps territorial partitioning by osprey pairs.

As one traces the nest locations and descriptors in Table 4, the distinction between "nest" and "active nest", and the concept of territory should be kept in mind. The territory occupied by one pair of breeding ospreys contains one or more nest structures. Pairs often have more than one nest, and what might be inferred by the observer as inactive nests may in fact be second or third nests of extant pairs rather than abandoned nests with no birds. One nest is used for brooding; the others are used for resting or other behavior by the adults. This is probably the situation at tower 6/1, where although multiple nests are listed only one is "active". Although the actual number of total nests is useful, it is the number of active nests, which equate to breeding territories, that is the more useful statistic in a population sense.

The summary statistics in Table 4 evidence a plateau in number of both nests and active nests from 1975-1979 and another general stabilization, but at a somewhat higher level, from 1980-1984. The dramatic increase in both categories from 1974 to 1975, the greatest change seen during the study, may reflect over-conservatism or inexperience by the earliest observer in 1974, but there seems little reason to question the 1975 data. There is an obvious and puzzling disparity between our local observations of active nests in 1975 and reference to local 1975 osprey success in the literature. Henny et al. (1977) state, relative to 1975: "Few ospreys now nest on the Delaware Bay side of New Jersey (Cape May Point to Wilmington Bridge). One nesting pair was seen from the air and by ground investigators and a second was located from the ground. We doubt that many other nests were in the area".

Breeding success, the realistic measure of population status, is assessed from the proportion: no. fledglings produced per no. active nests. To be meaningful and statistically testable, the area covered and the number of nests should be larger than available in the present study. However, the study area "population" by itself and as part of the much larger New Jersey population can be characterized to some degree. Summary data on no. of fledglings follow the two-plateau pattern mentioned earlier, i.e., the two periods 1975-1979 and 1980-1984, as does the index, no. fledglings/no. active nests. In five of the six years 1974-1979 the proportion is at least 1.0; during 1980-1984 it is less than 1.0 except in 1983. These levels can be reflected against the oft-cited reference by Henny and Wight (1969) that "0.95-1.30 young per active nest are required for population stability in ospreys". The 1974-1979 levels are within range and appear reasonable. However, the quantitative and qualitative nature of the inputs, and their potential effect on the index, should be considered. The lower number of fledgling to active nest ratios for 1980 to 1984 may be a result of overestimating the number of active nests. Nest-presence by two-year immature birds or three-year old non-breeders was included in the "active count." Whatever the reasons for observed levels during 1980-1984, the increase during 1983 and 1984 is a positive sign.

3.4.2 Bald Eagle

A total of 22 sightings of bald eagle in the general region were reported by Project observers during 1971 through 1984; 15 of these were through 1978. In 1979 there was one; in 1980, none; in 1981, four; in 1982, two; in 1983, none; and

in 1984, none. Of the 15 sightings prior to 1979, six were in the New Jersey portion of the region; five near northern Artificial Island and one near Hope Creek. Hardin (1978) stated in the Project Annual Report that bald eagle did not currently nest in the study area, but listed historical reference to a nest on Blackbird Creek in Delaware.

Frier (1982) listed specifics on the bald eagle as it occurs in New Jersey. She described a small wintering population throughout New Jersey which concentrates in the Dingman Ferry area of the Delaware River, the Brigantine National Wildlife Refuge area, and the Dividing Creek area of Cumberland County along the Delaware River. She reported one breeding pair remaining in New Jersey, in Cumberland County, and stated that "during winter of 1980 there were 16 wintering eagles observed."

Records of New Jersey Birds (NJ Audubon Society, 1984) references a pair of bald eagles constructing a nest in the spring of 1984 at a Salem County location and the observance of an immature bird at the site in April of that year. This location was not on the Project survey route which emphasized the transmission corridors; however, NJDEP personnel have corroborated the reported sightings. Apparently, the nest was not actually used in 1984. The Records issue also describes a sighting of an adult bald eagle over Audubon, NJ on March 8, 1984.

3.5 OVERVIEW

Appraisal of the local status of these two raptors requires speculation in addition to analysis of the limited data. Fortunately in the case of the osprey, in New Jersey there is an organized statewide research/reestablishment program underway by the New Jersey Non-Game and Endangered Species Program. This program has produced a body of information on osprey in New Jersey, and the local breeding population can be considered on the basis of the statewide database. That perspective was developed in conversation with the New Jersey Non-Game and Endangered Species Program office (J. Frier-Murza, Prog. Mgr., pers. comm.) on January 31, 1985. Perhaps the most singularly important and suggestive element in that conversation is that the Non-Game and Endangered Species Group is recommending to the State, i.e., the New Jersey Department of Environmental Protection, that the osprey be de-classified. Notice of intent to recommend this de-classification was published in the State (NJ) Register on February 19, 1985. Among the evidence supporting this recommendation is the trend in number of breeding osprey pairs in New Jersey. In 1973 there were 50; in 1981, 97;

and in 1984 there were 108. The statewide population productivity index has been within the Henny and Wight (1969) range of 0.95 to 1.3. Again in the conversation, there was mutual inclination to the position that the study-area "population" is behaving similarly to the NJ population, and that the available local statistics, being based on limited sample size, should not be rigorously considered on their own. It is accepted that the PSE&G transmission towers have contributed positively to the re-establishment of the osprey breeding population in New Jersey.

Further, with the completion of the new transmission line, which parallels the Salem-New Freedom North line, and the concomitant decrease in regular human activity near the towers of both lines, the number and availability of potential nesting sites will be increased. This can only advantage the local breeding osprey population.

Definition of status and speculation on potential of bald eagle in this area is difficult. The species has experienced a variety of negative factors including shooting, egg collection by oologists and starting in the 1940's, and of perhaps the most consequence, chemical contamination. The last involves primarily pesticides, most notably the long-lived DDT, which pass through the food chain to fish and other aquatic prey of the eagle and to the eagle itself. The most notable effect was a significantly reduced hatch rate. Abbott (1982) reports that in 1962, only 5 of 37 rechecked active nests produced young, compared to 31 successful of 35 rechecked nests in 1936, before chemical pesticides were commonly used. The use of DDT was formally banned in the mid 1960's, and there is evidence that populations may be increasing.

Abbott (1982) discussed the status of the bald eagle in Delaware, Maryland and Virginia. He stated that Delaware had four active bald eagle nests in 1981 for the first time on the survey since 1939. The nest at Bombay Hook National Wildlife refuge was abandoned in 1982, and he stated it had produced young in only three of the past twenty years (1970, 1976, and 1979). He listed several successful nests in lower Delaware. For the three states, he listed 87 active nests in 1978 and 94 in 1981, and cited numbers of eagles hatched as 59 in 1978, 67 in 1979, 74 in 1980 and 97 in 1981. This trend was taken as an encouraging indication that former problems with chemical pollutants, which affected egg hatching, are being worked out. Perhaps, if this perceived trend is real and limiting factors are easing, successful eagle nesting may again be observed in the study area.

SECTION 4.0
LITERATURE CITED

- Abbott, J. M. 1982. Our National Heritage. Delaware Conservationist. Vol. XXV(1):6-11.
- Burger, J. 1976. Behavior of hatchling diamondback terrapin (Malaclemys terrapin) in the field. Copeia 1976(4):742-748.
- Burger, J. and W. A. Montevechi. 1975. Nest site selection in the terrapin, Malaclemys terrapin. Copeia 1975(1):113-119.
- Burger, W. 1937. Experimental photoperiodicity in the male turtle, Pseudemys elegans (Wiel). Amer. Nat. 71:481-487.
- Carr, A. 1952. Handbook of turtles. Comstock Publishing Assoc., Cornell Univ. Press, Ithaca, NY. 542 p.
- Carr, A. and M. H. Carr. 1972. Site fixity in the Caribbean green turtle. Ecology 53:425-429.
- Carr, A. and L. Ogren. 1960. The ecology and migration of sea turtles. IV. The green turtle in the Caribbean Sea. Bull. Am. Mus. Nat. Hist. 121:1-48.
- Frier, J. 1982. Pandion haliaetus, osprey, fish hawk. Pages 196-198 in New Jersey's Endangered and Threatened Plants and Animals. Stockton State College, Center for Environmental Research, Pomona, NJ. 385 p.
- Hardin, D. L. 1980. The terrestrial ecology of Artificial Island and vicinity 1972-1978: A summary. For Public Service Electric and Gas Co., Newark, NJ. Ichthyological Associates, Inc., Middletown, DE. 70 p.
- Henny, C. J. et al. 1977. Mid-atlantic coast osprey population: present numbers, productivity, pollutant contamination, and status. J. Wildl. Manage. 41(2):1977.
- Henny, C. J. and W. T. Van Velzen. 1972. Migration patterns and wintering localities of American ospreys. J. Wildl. Man. 36(4):1133-1141.
- Henny, C. J. and H. M. Wight. 1969. An endangered osprey population: estimates of mortality and production. Auk 86(2):188-198.

Hildebrand, S. F. 1932. Growth of diamondback terrapins, size attained, sex ratios and longevity. *Zoologica* 9(15):551-563.

Hurd, L. E. et al. 1979. An ecological study of a natural population of diamondback terrapins (Malaclemys t. terrapin) in a Delaware Salt Marsh. *Estuaries* 2(1):28-33.

NJ Audubon Society. 1984. Records of New Jersey birds. Autumn 1984, Vol. X; No. 3.

NJDEP (New Jersey Department of Environmental Protection). 1974. New Jersey Administrative Code; 7;25-4.13. List of endangered species. NJ Dept. Environmental Protection, Division of Fish, Game & Wildlife.

NJ Audubon Society. 1984. Records of New Jersey birds. Autumn 1984, Vol. X; No. 3.

NOAA (National Oceanic and Atmospheric Administration). 1972. C & G S Chart 294, Delaware River, Smyrna River to Wilmington. National Ocean Survey, Washington, DC.

_____. 1977. Local climatological data - Wilmington, DE. National Climatic Center, Asheville, NC.

Parnell, J. F. and R. Walton. 1972. Osprey reproductive success in southeastern North Carolina. Pages 139-142 in Transactions of the North American Osprey Research Conference, College of William and Mary, Williamsburg, VA, 10-12 February 1972. 258 p.

Porter, K. 1972. Herpetology. W. B. Saunders & Co., Phila., PA. 524 p.

PSE&G (Public Service Electric and Gas Co.). 1978. Biotic environmental surveillance. Vol. 2 of 3, Annual environmental operating report (nonradiological). SNGS - Unit No. 1, January 1 through December 31, 1977. 679 p.

_____. 1979. Annual environmental operating report (nonradiological). SNGS - Unit No. 1, January 1 through December 31, 1978.

_____. 1980. Annual environmental operating report (nonradiological). SNGS - Units 1 and 2, January 1 through December 31, 1979.

_____. 1981. Annual environmental operating report (nonradiological). SNGS - Units 1 and 2, January 1 through December 31, 1980.

- _____. 1982. Annual environmental operating report (nonradiological). SNGS - Units 1 and 2, January 1 through December 31, 1981.
- _____. 1983. Annual environmental operating report (nonradiological). SNGS - Units 1 and 2, January 1 through December 31, 1982.
- _____. 1983. (unpublished). Environmental monitoring program diamondback terrapin, osprey, and eagle monitoring studies 1974-1981: A summary. 19 p.
- _____. 1984. Annual environmental operating report (nonradiological). SNGS - Units 1 and 2, January 1 through December 31, 1983.
- Schuler, V. J. 1974. An ecological study of the Delaware River in the vicinity of Artificial Island. Progress report for the period January through December 1972. Ichthyological Associates, Inc. 746 p.
- _____. 1975. An ecological study of the Delaware River in the vicinity of Artificial Island. Progress report for the period January through December 1973. Ichthyological Associates, Inc. 579 p.
- _____. 1976. An ecological study of the Delaware River in the vicinity of Artificial Island. Progress report for the period January through December 1974. Ichthyological Associates, Inc. 517 p.
- _____. 1977. An ecological study of the Delaware River in the vicinity of Artificial Island. Progress report for the period January through December 1975.
- USDI. 1973. Threatened wildlife of the U.S. Bur. Sport Fish and Wildlife Resource Publ. 114. 289 p.

Table 1. Annual summary data on observed nesting, nest depredation, and hatchlings of diamondback terrapin observed at a beach north of Liston Point, DE, 1975-1984.

| Year (# visits) | <u>I</u> Nests | | <u>II</u> Eggs | | <u>III</u> Total observed Turtles Tracks | | Hatchlings (Actual or Tracks) included in Column III |
|-----------------|-------------------|------|-------------------|-------|--|-------|---|
| | Non-Dep. | Dep. | Non-Dep. | Dep. | | | |
| 1975 (21) | 6 | 498 | 52 | 2,443 | 34 | 189 | 146 |
| 1976 (32) | 15 | 393 | 170 | 3,425 | 30 | 470 | 215 |
| 1977 (42) | 25 | 259 | 237 | 4,192 | 44 | 1,544 | 212 (32)* |
| 1978 (46) | 61 | 444 | 616 | 3,455 | 111 | 1,093 | 54 |
| 1979 (40) | 45 | 267 | 483 | 2,276 | 43 | 618 | 12 |
| 1980 (33) | 19 | 429 | 122 | 3,405 | 45 | 712 | 49 |
| 1981 (40) | 18 | 337 | 132 | 2,656 | 29 | 514 | 15 |
| 1982 (41) | 28 | 344 | 220 | 2,830 | 20 | 514 | 57 |
| 1983 (18) | 18 | 238 | 111 | 1,776 | 10 | 132 | 72 |
| 1984 (17) | 12 | 285 | 99 | 2,193 | 47 | 156 | 0 |

* - hatchlings observed in nests upon excavation.

Table 2. Annual summary data on observed nesting, nest depredation, and hatchlings of diamondback terrapin observed at a beach north of Sunken Ship Cove, NJ, 1975-1984.

| | | <u>I</u> | | <u>II</u> | | <u>III</u> | | Hatchlings (Actual or Tracks) |
|-----------------|------|--------------|------|-------------|------|----------------|--------|----------------------------------|
| | | <u>Nests</u> | | <u>Eggs</u> | | Total observed | | included in |
| Year (# visits) | | Non-Dep. | Dep. | Non-Dep. | Dep. | Turtles | Tracks | Column III |
| 1975 | (19) | 1 | 44 | 3 | 191 | 6 | 53 | 25 |
| 1976 | (32) | 8 | 0 | 57 | 0 | 7 | 112 | 79 |
| 1977 | (39) | 3 | 0 | 25 | 0 | 15 | 195 | 195 (15)* |
| 1978 | (42) | 2 | 3 | 20 | 16 | 12 | 71 | 33 |
| 1979 | (27) | 10 | 4 | 97 | 28 | 0 | 92 | 16 |
| 1980 | (32) | 6 | 3 | 52 | 13 | 3 | 129 | 84 |
| 1981 | (40) | 3 | 1 | 17 | 4 | 0 | 39 | 8 |
| 1982 | (42) | 6 | 0 | 62 | 0 | 0 | 38 | 6 |
| 1983 | (18) | 2 | 0 | 14 | 0 | 0 | 4 | 0 |
| 1984 | (17) | 2 | 1 | 19 | 90 | 0 | 40 | 34 |

* - hatchlings observed in nests upon excavation.

Table 3. Annual summary data on observed nesting, nest depredation, and hatchlings of diamondback terrapin observed at a beach north of Hope Creek, NJ, 1975-1984.

| Year (# visits) | <u>I</u> <u>Nests</u> | | <u>II</u> <u>Eggs</u> | | <u>III</u> Total observed | | Hatchlings (Actual or Tracks) included in Column III |
|-----------------|--------------------------|------|--------------------------|-------|------------------------------|--------|---|
| | Non-Dep. | Dep. | Non-Dep. | Dep. | Turtles | Tracks | |
| 1975 (21) | 8 | 518 | 31 | 2,814 | 34 | 23 | 17 |
| 1976 (32) | 11 | 132 | 123 | 915 | 74 | 68 | 80 |
| 1977 (45) | 35 | 170 | 298 | 1,124 | 108 | 266 | 177 (62)* |
| 1978 (44) | 31 | 235 | 192 | 1,354 | 89 | 281 | 1 |
| 1979 (40) | 3 | 64 | 26 | 473 | 13 | 132 | 0 |
| 1980 (26) | 9 | 97 | 48 | 699 | 17 | 117 | 13 |
| 1981 (39) | 5 | 49 | 20 | 234 | 8 | 54 | 12 |
| 1982 (39) | 19 | 29 | 119 | 191 | 31 | 126 | 1 |

* = hatchlings observed in nests upon excavation.


Table 4. Summary of osprey nesting activity within 16 km of southern Artificial Island: 1974-1984.


Key: Numbers indicate young fledged; *=Data collected by PSE&G Transmission & Development Dept.; N=Nest present, may have been active or constructed as housekeeping nest; A=Active nest, eggs observed or adults appeared to be incubating eggs and defending nest; Owl=great horned owl nested in former osprey nest; **=assumed number active. Helicopter observation began too late in the season to assess egg production.

| NEST LOCATION | 1974 | 1975 | 1976 | 1977 | 1978 | 1979 * | 1980 | 1981 | 1982 | 1983 | 1984 |
|---------------------------|------|------|------|------|------|--------|------|------|------|------|------|
| NEW JERSEY | | | | | | | | | | | |
| Transmission Line Towers | | | | | | | | | | | |
| <u>Salem-Keeney</u> | | | | | | | | | | | |
| Tower #12/1 | | | | | | | | | | N,N | 2 |
| 11/3 | | | | | | | | | 1 | 2 | A |
| 10/1 | | | | | | | | | | N | |
| 9/3 | | | | | | | | | | | A |
| 8/4 | | | A | N | N | A | N | A | N | | |
| 8/3 | | | | | | | N | A | | | |
| 5/1 | | | | | | N | A | | | | |
| 4/3 | | | | | 1 | N | N | A | N | A | A |
| 4/2 | | | | N | | N | N | N | | | |
| 4/1 | | 2 | A | | | | N,N | Owl | | | |
| 3/4 | | | A | | | A | N | | | | |
| 3/3 | | | | | | | | | | | A |
| 3/2 | A | | | | | A | N | A | | A | |
| 3/1 | | | 1 | N | N | | N | | | N | |
| <u>New Freedom: South</u> | | | | | | | | | | | |
| Tower # 5/3 | | | | | | | A | 1 | A | A | |
| 5/2 | | | | | | | | N | | | |
| 5/1 | | 2 | 1 | 1 | 3 | A | 2 | 2 | 2 | A | 3 |
| 4/1 | | N | N | N | | | | | | | |
| 3/3 | | N | N | N | | | A | 1 | 1 | 3 | 2 |
| 2/4 | | | | | 1 | A | | | | | |
| 2/3 | | | | | | N | | | | | |

Table 4.
Continued.

| NEST LOCATION | 1974 | 1975 | 1976 | 1977 | 1978 | 1979* | 1980 | 1981 | 1982 | 1983 | 1984 |
|---|------|------|------|------|------|-------|------|-------|------|------|-------|
| <u>New Freedom: North</u> | | | | | | | | | | | |
| Tower # 6/1 | | | 1 | 2 | A | A | 2 | A,N,N | 1 | A | A,N,N |
| 4/1 | | N | 1 | 1 | 1 | A | 1 | A | A | A | 1 |
| 3/4 | | N | | 2 | 3 | A | A | 2 | A | 2 | A |
| 3/3 | | | | | | | | | | | A |
| 2/3 | | | | | 1 | A | 1 | A | A | A | |
| Raccoon Ditch | | | | | | | | | | | |
| Old cedar tree | | | | | | | | N | A | 1 | A |
| Nest platform | | | | | | | | N | N | N | N |
| <hr/> Area not monitored <hr/> | | | | | | | | | | | |
| <u>DELAWARE</u> | | | | | | | | | | | |
| Delaware River | | | | | | | | | | | |
| Getty-Range Tower | A | 2 | 2 | 2 | N | A | 1 | A | A | 2 | 2 |
| Reedy Island | | | | | | | | | | | |
| East | A | | | | | | | | | | |
| West | N | 2 | 1 | | | | | | | | |
| Jetty | N | N | N | N | A | | | | | | |
| Transmission Line Towers | | | | | | | | | | | |
| DP&L #5015/47 | | | | | | | | A | A | Owl | N |
| DP&L #5015/46 | | | | | | | | | | 3 | 2 |
| <hr/> Under construction <hr/> Area not monitored <hr/> | | | | | | | | | | | |
| Smyrna River | | | | | | | | | | | |
| Range Tower | N | N | A | | | | | | | | |
| <u>SUMMARY</u> | | | | | | | | | | | |
| Nests | 6 | 10 | 13 | 11 | 11 | 14 | 18 | 18 | 14 | 18 | 18 |
| Active nests | 3 | 7** | 10 | 7 | 8 | 11 | 10 | 12 | 11 | 13 | 14 |
| Successful (i.e., young fledged) nests | 2 | 4 | 6 | 5 | 6 | 10 | 5 | 4 | 4 | 6 | 6 |
| Fledglings | 4 | 8 | 7 | 8 | 10 | 16 | 7 | 6 | 5 | 13 | 12 |
| Fledglings/active nest | 1.33 | 1.14 | 0.70 | 1.14 | 1.25 | 1.45 | 0.70 | 0.50 | 0.45 | 1.00 | 0.85 |
| Successful-/active nest | 0.67 | 0.57 | 0.60 | 0.71 | 0.75 | 0.81 | 0.50 | 0.33 | 0.36 | 0.46 | 0.42 |






PUBLIC SERVICE ELECTRIC AND GAS COMPANY

Number of adult diamondback terrapin and tracks observed per sampling date during 1979 through 1984 at a beach on the Delaware River near Liston Point, DE.


Figure 2



PUBLIC SERVICE ELECTRIC AND GAS COMPANY

Number of adult diamondback terrapin and tracks observed per sampling date during 1979 through 1984 at a beach on the Delaware River near Sunken Ship Cove, NJ.


Figure 3



PUBLIC SERVICE ELECTRIC AND GAS COMPANY

Number of adult diamondback terrapin and tracks observed per sampling date during 1979 through 1984 at a beach on the Delaware River near Hope Creek, NJ.


Figure 4



PUBLIC SERVICE ELECTRIC AND GAS COMPANY

Annual mean number of observed adult diamondback terrapin and tracks at Liston Point, DE, and Hope Creek and Sunken Ship Cove, NJ, 1975-1984.


Figure 5



PUBLIC SERVICE ELECTRIC AND GAS COMPANY

Nesting sites (O), historical and present, observed during SNGS Osprey Study, 1974-1984. Dashed (--) line is new transmission line nearing completion in 1984.


Figure 6



PUBLIC SERVICE ELECTRIC AND GAS COMPANY

Tower locations on transmission lines from
SNGS; Osprey study, 1974-1984.
○ indicates tower used by nesting osprey.

Figure 7



PUBLIC SERVICE ELECTRIC AND GAS COMPANY

Northern extension of Salem-Keeney line,
Osprey study, 1974-1984.

○ indicates tower used by nesting osprey.

Figure 8