IMPINGEMENT MONITORING PROGRAM

REPORT NO. 11 - FEBRUARY 1978

SALEM NUCLEAR GENERATING STATION UNIT NO. 1

DOCKET NO. 50-272
OPERATING LICENSE NO. DPR-70

Docket # Control # Of Documents
REGULATORY DOCKET FILE

PUBLIC SERVICE ELECTRIC AND GAS COMPANY NEWARK, NEW JERSEY MARCH 20, 1978

P77 54 19

750ff0029

SUMMARY

This is the eleventh monthly impingement monitoring program report required by Section 3.1.2.2 of the Environmental Technical Specifications (ETS). The ETS are Appendix B of the Salem Nuclear Generating Station (SNGS) Operating License No. DPR-70. The tabular summaries attached include daily estimates of the number and weight of fish impinged on the Circulating Water System (CWS) traveling screens (Table 2) and the Service Water System (SWS) traveling screens (Table 3). CWS operation is also summarized (Table 1).

The monitoring program at both the CWS and SWS consisted of three 24-hour sampling periods per week. The CWS screens were sampled four times per 24-hour period at approximately six-hour intervals. Samples were collected by diverting approximately three minutes of screen wash water to the counting pool located on the north end of the CWS intake structure. Survival rates were calculated for CWS samples only. SWS screen samples consisted of either one 24-hour or two 12-hour samples per day. These samples consisted of all fish impinged and found in the trash baskets located on the intake structure proper.

MDL:aec

TABLE 1. - START TIME AND DURATION OF IMPINGEMENT SAMPLES AT THE CIRCULATING MATER INTAKE, S.N. 6.S. UNIT 1

1 FEBRUARY THROUGH 28 FEBRUARY 1978. NUMBER OF AND SCREENS IN OPERATION, SCREEN OPERATING MODE INTERMITTENT, C= CONTINUOUS), WEIGHT (G/3 MIN. SAME OF DETRITUS, AND WATER TEMPERATURE (C)

HE INDICATED.

,	

DATE	SAMPLE START Time					DETRITUS WEIGHT (G/3 MIN SAMPLE)	WATER TEMPERATURE (C)
J2/g1/7B	1200	3	556655665666666666666666666666666666666	5	С	5,100	0 • 4
32/31/78	1800	3	5	5	C	17,600	0.9
J2/J2/78	0000	3	6	ć	C	6,100	0.9
32/02/7B	0600	3	6	ć	C	600	0.9
J2/02/7B	1200	3	5	5	C	23,800	1.0
32/02/78	1800	3	5	5	C	1,400	1.2
32/33/79	0000	3	6	6	C	6.000	0.8
32/03/7 8	0600	3	6	6	Ċ	400	0.9
22/06/7B	1200	3	5	5	Č	3,800	1.9
32/06/78	1800	3	6	6	Č	5,000	1.0
32/07/78	0000	3	5	6	C -	9,600	1.0
32/07/78	0600	3	6	6	C C	2,100	1.0
32/08/75	1200	3	5	5	Ċ	1,200	1.2
32/08/78	1600	3	6	6	t	4,600	0.0
32/09/7B	0000	3	6	0	C C	7,200	0.2
32/39/78	0600	3	0	D 4	Ċ	45,500	0.3 0.4
22/09/78	1210	3	0	0	C	3,800	0.0
32/09/7B	1600	3	D E	D 4	C	35,000 4,000	0.0
32/10/78	0000	3	2	0	C	60,000	0.0
32/13/7B	D6DD	3	0 .	D 4	Ċ	1,200	1.3
32/13/78	1200	3	0	4	Č	3,800	1.0
32/13/78	1800	3	0 4	6	C	3,000	0.3
32/14/78	0000	7	4	, , , , , , , , , , , , , , , , , , ,	Č	3,000	0.4
32/14/7B	0600 1200	3	6	6	ř	4,100	0.8
02/15/78 02/15/78	1800	3	6	6	Č	700	0.4
32/16/78	0000	₹	6	6	č	7,000	0.1
32/16/78	0600	3	6	ķ	č	500	0.0
32/16/78	1200	₹	6	6	č	10,900	0.9
32/16/78	1800	₹	6	6	č	100	1.2
32/17/78	0000	~	6	6	č	50,400	1.6
32/17/78	0600	ž	6	6	č	1,400	. 0.8
32/21/78	1200	3	6	6	č ·	3,900	0.1
32/21/7B	1800	3	6	6	C	8,200	0.4
32/22/75	0000	3	6	6	C	39,500	0.9
J2/22/7B	0600	3	6	6	С	17,000	0.1
32/22/78	1210	3	6	6	С	15,000	0.1
02/22/78	1810	3	6	6	C	35,000	0.1
32/23/78	0000	3	6	6	C	2,600	- 1.2
22/23/78	0600	3	6	6	C	41,000	- 1.2
32/23/78	1205	3	5	5	С	700	0.2
32/23/78	1800-	3	5	5	C	10,000	- 0.7
32/24/78	0000	3	5	5	C	2,800	2.0
32/24/78	0600	3	5	5	C	19,100	0.5
J2/27/78	1200	3	6	6	C	19,000	2.6
32/27/78	1800	3	5	5	č	72,000	2.9
32/28/78	0000	3	6	6	C	5,000	2.2
32/28/78	0600	3	6	6	C	11,000	3.0

F# MO+ 04-

TABLE 2. - IMPINGEMENT - S.N.G.S. CIRCULATING WATER SYSTEM INTAKE, 1 FEBRUARY 1978 TO 28 FEBRUARY 1978.

SAMPLING PERIOD - 02/01/78 1200 THROUGH 02/02/78 0600

			2 S	URV	IVAL	TOTAL Number	NEAN ESTIMATED NUMBER/		STANDARD	MEAN ESTIMATED bt/26 Hrs	STANDARD	(6	GHT M)	CH	IGTH IM)	
SPECIES	N	C F	L	٥	₽ •	COLLECTED	24 HOURS		DEVIATION	(K6)	DEVIATION	M1W.		MIN.	MAX.	
H. NUCHALIS	4	4	35	1	62	. 76	9360		8309.2	95•31	79.7	3.3	24.7	67.5	127.5	
M. AMERICANA	4	4	6	11	61	61	7320	,	8102.1	53.39	57.1	2.2	53.2	57.5	137.5	
M. SANATILIS	. 4	2	16	٥	83	12	1440		1838.2	12.27	15.3	4.3	17.0	72.5	107.5	
D. CEPEDIANUM	4	1	٥	Ō	100	1	120		240.0	5.07	10.1	42.3	42.3	152.5	152.5	
1. CATUS	4	2	50	0	50	2	240		277.1	106.05	128.0	346.2	537.6	292.5	342.5	,
TOTAL						154	18480			272.09						

N = NUMBER OF 3-MINUTE SAMPLES.

MEAN ESTIMATED WEIGHT PER 24 HOURS FOR EACH SPECIES WAS CALCULATED BY THE SAME METHOD AS MEAN ESTIMATED NUMBER.

CF = CAICH FREQUENCY (NUMBER OF SAMPLES IN WHICH THE SPECIES APPEARED).

X SURVIVAL: L = LIVE; D = DEAD; D* = DAMAGED.

MEAN ESTIMATED NUMBER PER 24 HOURS FOR EACH SPECIES WAS CALCULATED BY MULTIPLYING THE NUMBER OF SPECIMENS IN EACH 3-MINUTE SAMPLE BY 480 TO OBTAIN A 24-HOUR ESTIMATE. THE ESTIMATED NUMBER FOR EACH SAMPLE WAS SUMMED AND DIVIDED BY THE NUMBER OF SAMPLES TO OBTAIN THE MEAN.



SAMPLING PERIOD - 02/02/78 1200 THROUGH 02/03/78 0600

			1	S UR V I	VAL	TOTAL Number	MEAN ESTIMATED NUMBER/	STANDARD	MEAN ESTIMATED WT/26 HRS	STANDARD	₩E I	GHT M)		GTH M)
SPECIES	N	C F	Ĺ	D	D *	COLLECTED	24 HOURS	DEVIATION	(KB)	DEVIATION	MIN.	MAX.	MIN.	. XAM
H. NUCHALIS	4	4	45	13	40	22	2640	831.3	29.97	13.2	2.3	31.3	67.5	142.5
M. AMERICANA	4	4	0	6	93	16	1920	1239.3	38.14	51.4	3.0	213.4	62.5	217.5
D. CEPEDIANUM	4	1	0	100	۵	1	120	240.0	6 - 9 6	13.9	58.0	58.0	162.5	162.5
TOTAL						39	4680		75.09					

N = NUMBER OF 3-MINUTE SAMPLES.

CF = CATCH FREQUENCY (NUMBER OF SAMPLES IN WHICH THE SPECIES APPEARED).

I SURVIVAL: L = LIVE; D = DEAD; D+ = DAMAGED.

MEAN ESTIMATED NUMBER PER 24 HOURS FOR EACH SPECIES WAS CALCULATED BY MULTIPLYING THE NUMBER OF SPECIMENS IN EACH 3-MINUTE SAMPLE BY 48D TO DOTAIN A 24-HOUR ESTIMATE. THE ESTIMATED NUMBER FOR EACH SAMPLE WAS SUMMED AND DIVIDED BY THE NUMBER OF SAMPLES TO DOTAIN THE MEAN.

MEAN ESTINATED WEIGHT PER 24 HOURS FOR EACH SPECIES WAS CALCULATED BY THE SAME METHOD AS MEAN ESTIMATED NUMBER.

TABLE 2. - (CONTINUED).

SAMPLING PERIOD - 02/06/78 1200 THROUGH 02/07/78 0600

SPECIES	N	C F	l s L	urv D	IVAL D=	TOTAL NUMBER COLLECTED	MEAN ESTIMATED NUMBER/ 24 HOURS	STANDARD DEVIATION	MEAN ESTIMATED WT/24 HRS (KG)	STANDARD DEVIATION	WEIG (G		_	HAX.
N. AMERICANA N. SAMATILIS H. NUCHALIS D. CEPEDIANUM U. CHUSS	4 4 4	4 2 3 1	5 0 42 0	_	66 50 100	133 3 28 1	15960 360 3360 120 120	8676.8 459.5 2569.9 240.0 240.0	135.84 3.31 42.74 5.16 .81	95.6 4.2 37.3 10.3	2.0 3.8 3.6 43.2 6.8	86.1 18.5 19.6 43.2 6.8		177.5 117.5 127.5 147.5 97.5
TOTAL						166	19920		167.88		**************************************			

N = NUMBER OF 3-MINUTE SAMPLES.

CF = CATCH FREQUENCY (NUMBER OF SAMPLES IN WHICH THE SPECIES APPEARED).

K SURVIVAL: L = LIVE; D = DEAD; D= = DAMAGED.

MEAN ESTIMATED NUMBER PER 24 HOURS FOR EACH SPECIES WAS CALCULATED BY MULTIPLYING THE NUMBER OF SPECIMENS IN EACH 3-MINUTE SAMPLE BY 483 TO OBTAIN A 24-HOUR ESTIMATE. THE ESTIMATED NUMBER FOR EACH SAMPLE WAS SUMMED AND DIVIDED BY THE NUMBER OF SAMPLES TO OBTAIN THE MEAN.

MEAN ESTIMATED WEIGHT PER 24 HOURS FOR EACH SPECIES WAS CALCULATED BY THE SAME METHOD AS MEAN ESTIMATED NUMBER.

TABLE 2. - (CONTINUED).

SAMPLING PERIOD - 02/08/78 1200 THROUGH 02/09/78 0600

			x :	S UR V	LVAL	TOTAL Number	MEAN ESTIMATED NUMBER/	STANDARD	MEAN ESTIMATED WT/24 HRS	STANDARD	PE]	GHT M)	_	ETH M)
SPECIES	N	C F	L	D	D *	COLLECTED	24 HOURS	DEVIATION	(K5)	DEVIATION	MIN.	MAX.	MIN.	MAX.
M. AMERICANA	4	4	2	25	71	101	12120	7069.4	82-82	56.7	2.2	89.4	52.5	172.5
H. MUCHALIS	4	4	37	25 37	71 25	В	960	391.9	12.04	6.9	4.0	23.0	72.5	127.5
M. SAXATILIS	4	2	۵	100	0	2	240	277.1	1.12	1.3	3.7	5.7	72.5	72.5
I. CATUS	4	2	D	D	100	3	360	459.5	140.70	166.8	269.8	489.0	267.5	317.5
,A. AESTIVALIS	4	1	D	100	٥	1	120	240.0	•32	• 6	2.7	2.7	62.5	62.5
TOTAL						11'5	13800		237.00		****			

N = NJM3ER OF 3-MINUTE SAMPLES.

CF = CATCH FREQUENCY (NUMBER OF SAMPLES IN WHICH THE SPECIES APPEARED).

X SURVIVAL: L = LIVE; D = DEAD; D* = DAMAGED.

MEAN ESTIMATED NUMBER PER 24 HOURS FOR EACH SPECIES WAS CALCULATED BY MULTIPLYING THE NUMBER OF SPECIMENS IN EACH 3-MINUTE SAMPLE BY 48) TO OBTAIN A 24-HOUR ESTIMATE. THE ESTIMATED NUMBER FOR EACH SAMPLE WAS SUMMED AND DIVIDED BY THE NUMBER OF SAMPLES TO OBTAIN THE MEAN.

MEAN ESTIMATED WEIGHT PER 24 HOURS FOR EACH SPECIES WAS CALCULATED BY THE SAME METHOD AS MEAN ESTIMATED NUMBER.

TABLE 2. - (CONTINUED).

SAMPLING PERIOD - 02/09/78 1210 THROUGH 02/10/78 0600

			1 :	SURV	IVAL	TOTAL Number	MEAN ESTIMATED NUMBER/	STANDARD	MEAN ESTIMATED UT/26 HRS	STANDARD	⊌E I			G TH	
\$PECIES		C F	L 	۵	D •	COLLECTED	24 HOURS	DEVIATION	(K&)	DEVIATION	MIN.	. XAM	MIN.	MAX.	
M. AMERICANA	4	4		24	71	45	5400	3578.6	105.83	77.6	1.6	427.6	52.5	267.5	
H. MUCHALIS	7	3	50	_		7,	480	391.9	3.45	3.8	4.2	11.7	77.5	107.5	
M. SAKATILIS	4	1	0	_	100	ĩ	120	240.0	1.44	2.8	12.0	12.0	102.5	102.5	
I. NEBULOSUS	4	Ž	50	_		ž	240	277.1	19,27	23.6	60.0	100.6	172.5	197.5	
P. FLAVESCENS	4	1	Ō	Ō	100	1	120	240.0	1.09	2.1	9.1	9.1	97.5	97.5	
M. BILINEARIS	4	1	0	100	0	1	120	240.0	1.40	2.9	12.2	12.2	117.5	117.5	•
TOTAL						54	6480		132.54						

N = NUMBER OF 3-MINUTE SAMPLES.

CF = CATCH FREQUENCY (NUMBER OF SAMPLES IN WHICH THE SPECIES APPEARED).

X SURVIVAL: L = LIVE; D = DEAD; D+ = DAMAGED.

MEAN ESTIMATED NUMBER PER 24 HOURS FOR EACH SPECIES WAS CALCULATED BY MULTIPLYING THE NUMBER OF SPECIMENS IN EACH 3-MINUTE SAMPLE BY 480 TO OBTAIN A 24-HOUR ESTIMATE. THE ESTIMATED NUMBER FOR EACH SAMPLE WAS SUMMED AND DIVIDED BY THE NUMBER OF SAMPLES TO OBTAIN THE MEAN.

MEAN ESTIMATED WEIGHT PER 24 HOURS FOR EACH SPECIES WAS CALCULATED BY THE SAME METHOD AS MEAN ESTIMATED NUMBER.

TABLE 2. - (CONTINUED).

SAMPLING PERIOD - 02/13/78 1200 THROUGH C2/14/78 0600

			1 S	UŖV	IVAL	TOTAL Number	MEAN ESTIMATED NUMBER/	STANDARD	MEAN ESTIMATED ht/24 hrs	STANDARD		GHT M)	L EN	GTH M)
SPECIES	N	C F	L 	D	D 4	COLLECTED	24 HOURS	DEVIATION	(KB)	DEVIATION	MIN.	MAX.	MIN.	MAX.
1. AMERICANA	4	4	7	3	88	26	3120	3427.8	88.04	161.7	2.1	272.5	57.5	232.5
I- NUCHALIS	4	3	0	20	80	5	600	459.5	8.36	7.5	8.6	18.1	87.5	122.5
P. FLAVESCENS	4	1	100	0	C	1	120	240.0	1. 26	2.5	10.5	10.5	102.5	102.5
L. CATUS	4	1	٥	0	100	1	120	240.D	50.D6	100.1	417.2	417.2	302.5	302.5
. PSEUDOHARENGUS	4	1	D	0	D	1	120	240.0	.51	1.0	4.3	4.3	72.5	72.5
M. SANATILIS	4	1	0	0	100	2	240	486.0	2.86	5.7	11.5	12.4	102.5	102.5
TOTAL						36	4320		151.09					

N = NUMBER OF 3-MINUTE SAMPLES.

CF = CATCH FREQUENCY (NUMBER OF SAMPLES IN WHICH THE SPECIES APPEARED).

2 SURVIVAL: L = LIVE; D = DEAD; D+ = DAMAGED.

MEAN ESTIMATED WEIGHT PER 24 HOURS FOR EACH SPECIES WAS CALCULATED BY THE SAME METHOD AS MEAN ESTIMATED NUMBER.

MEAN ESTIMATED NUMBER PER 24 HOURS FOR EACH SPECIES WAS CALCULATED BY MULTIPLYING THE NUMBER OF SPECIMENS IN EACH 3-MINUTE SAMPLE BY 480 TO OBTAIN A 24-HOUR ESTIMATE. THE ESTIMATED NUMBER FOR EACH SAMPLE WAS SUMMED AND DIVIDED BY THE NUMBER OF SAMPLES TO OBTAIN THE MEAN.



SAMPLING PERIOD - 02/15/78 1200 THROUGH 02/16/78 0600

			1 S	URVI	V AL	TOTAL NUMBER	MEAN ESTIMATED NUMBER/	STANDARD	MEAN ESTIMATED WT/24 HRS	STANDARD		6НТ м)		IGTH
SPECIES		C F	L 	D 	D #	COLLECTED	24 HOURS	NOITAIV30	(KS)	DEVIATION	MIN.	. XAM	MIN.	MAX.
H. NUCHALIS	4	2	25	0	75	8	960	1615.9	13.79	25.8	4.3	22.9	82.5	127.5
M. SAXATILIS	4	ž	٥			4	480	554.2	7.60	8.7	10.1	22.1	97.5	122.5
N. AMERICANA	4	4	٥	29	70	17	2040	1536.7	22.15	23.6	1.9	65.5	57.5	162.5
P. FLAWESCENS	4	1	100	D	0	1	120	240.0	24-06	48.1	200.5	200.5	237.5	237.5
TOTAL						30	3600		67.60					

N = NUMBER OF 3-MINUTE SAMPLES.

CF = CATCH FREQUENCY (NUMBER OF SAMPLES IN WHICH THE SPECIES APPEARED).

Z SURVIVAL: L = LIVE; D = DEAD; D* = DAMAGED.

MEAN ESTIMATED NUMBER PER 24 HOURS FOR EACH SPECIES WAS CALCULATED BY MULTIPLYING THE NUMBER OF SPECIMENS IN EACH 3-MINUTE SAMPLE BY 48D TO OBTAIN A 24-HOUR ESTIMATE. THE ESTIMATED NUMBER FOR EACH SAMPLE WAS SUMMED AND DIVIDED BY THE NUMBER OF SAMPLES TO OBTAIN THE MEAN.

MEAN ESTIMATED WEIGHT PER 24 HOURS FOR EACH SPECIES WAS CALCULATED BY THE SAME METHOD AS MEAN ESTIMATED NUMBER.

TABLE 2. - (CONTINUED).

SAMPLING PERIOD - 02/16/78 1.00 THROUGH 02/17/78 0600

			1 5	urv	JVÁL	TOTAL NUMBER	MEAN ESTIMATED NUMBER/	STANDARD	MEAN ESTIMATED WT/24 HRS	STANDARD		6HT	_	eth Eth
SPECIES		C F	L 	D	D *	COLLECTED	24 HOURS	MOLTALVED	(K6)	DEVIATION	MIN.	. XAM	MIN.	MAX.
H. NUCHALIS	4	3	33	22	44	16	2160	2524.7	26.98	29•7	4.6	22.1	77 5	127.5
N. AMERICANA	- 4	3	٥	45		40	4800	5852.5	56.02	64.8	2.9	114.4	57.5	162.5
A. ROSTRATA	4	1	100	Ō		1	120	240.0	3.91	7.8	32.6	32.6	262.5	262.5
M. SAXATILIS	. 4	1	D	۵	100	1	120	240.0	3.64	7.2	30.4	30.4	132.5	132.5
												~		
TOTAL						60	7200		90.55					

N = NUMBER OF 3-MINUTE SAMPLES.

CF = CATCH FREQUENCY (NUMBER OF SAMPLES IN WHICH THE SPECIES APPEARED).

X SURVIVAL: L = LIVE; D = DEAD; D+ = DAMAGED.

MEAN ESTINATED NUMBER PER 24 HOURS FOR EACH SPECIES WAS CALCULATED BY MULTIPLYING THE NUMBER OF SPECIMENS IN EACH 3-MINUTE SAMPLE BY 483 TO OBTAIN A 24-HOUR ESTIMATE. THE ESTIMATED NUMBER FOR EACH SAMPLE WAS SUMMED AND DIVIDED BY THE NUMBER OF SAMPLES TO OBTAIN THE MEAN.

MEAN ESTIMATED WEIGHT PER 24 HOURS FOR EACH SPECIES WAS CALCULATED BY THE SAME METHOD AS MEAN ESTIMATED NUMBER.

TABLE 2. - (CONTINUED).

SAMPLING PERIOD - 02/21/78 1200 THROUGH 02/22/78 0600

			r s		IVAL	TOTAL NUMBER	HEAN ESTIMATED NUMBER/	STANDARD	MEAN ESTIMATED WT/24 HRS	STANDARD		M)	()	iGTH M)
SPECIES		C F	L 	D	D,*	COLLECTED	24 HOURS	DEVIATION	(K6)	DEVIATION	MIN.	MAX.	MIN.	MAX.
H. NUCHALIS		7	25	0	75		480	391.9	7.31	5.6	9.9	18.4	92.5	117.5
A. ROSTRATA	2	2	50	50		2	240	277.1	4.41	8.7	.1	36.7	62.5	282.5
	7				-	7.								
M. AMERICANA	•	•	11	20		34	4080	2431.7	63.67	85.8	2.2	136.7	57.5	197.5
C- CARPID	•	1	0	Ð	100	1	120	240.0	88.52	177.0	737.7	737.7	337.5	337.5
M. SAXATILIS	4	1	0	0	100	. 1	120	240.0	- 6 D	1.2	5.0	5.0	77.5	77.5
FOTAL						42	5040		164.51					

N = NUABER OF 3-MINUTE SAMPLES.

CF = CATCH FREQUENCY (NUMBER OF SAMPLES IN WHICH THE SPECIES APPEARED).

I SURVIVAL: L = LIVE; D = DEAD; DA = DAMAGED.

MEAN ESTIMATED NUMBER PER 24 HOURS FOR EACH SPECIES WAS CALCULATED BY MULTIPLYING THE NUMBER OF SPECIMENS IN EACH 3-MINUTE SAMPLE BY 4BJ TO OPTAIN A 24-HOUR ESTIMATE. THE ESTIMATED NUMBER FOR EACH SAMPLE WAS SUMMED AND DIVIDED BY THE NUMBER OF SAMPLES TO OBTAIN THE MEAN.

MEAN ESTIMATED WEIGHT PER 24 HOURS FOR EACH SPECIES WAS CALCULATED BY THE SAME METHOD AS MEAN ESTIMATED NUMBER.



SAMPLING PERIOD - 02/22/78 1210 THROUGH 02/23/78 0600

	 		z	URVI	VAL	TOTAL NUMBER	MEAN ESTIMATED NUMBER/	STANDARD	MEAN ESTIMATED UT/24 HRS	STANDARD	₩E1	-		IETH
SPECIES	N	CF	L	D	D A	COLLECTED	24 HOURS	DEVIATION	(K&)	DEVIATION	MIN.	MAX.	MIN.	MAX.
M. AMERICANA H. MUCHALIS	4	4 2		32 58		25 12 _,	3000 1440	1725 • 1 2036 • 4	30.56 11.92	30.7 17.7	2.6 2.6	95.5 20.5	_	182.5 112.5
FOTAL	 					37	4440		42.48					

N = NUMBER OF 3-MINUTE SAMPLES.

CF = CATCH FREQUENCY (NUMBER OF SAMPLES IN WHICH THE SPECIES APPEARED).

I SURVIVAL: L = LIVE; D = DEAD; D* = DAMAGED.

MEAN ESTIMATED NUMBER PER 24 HOURS FOR EACH SPECIES WAS CALCULATED BY MULTIPLYING THE NUMBER OF SPECIMENS IN EACH 3-MINUTE SAMPLE BY 480 TO OBTAIN A 24-HOUR ESTIMATE. THE ESTIMATED NUMBER FOR EACH SAMPLE WAS SUMMED AND DIVIDED BY THE NUMBER OF SAMPLES TO OBTAIN THE MEAN.

MEAN ESTIMATED WEIGHT PER 24 HOURS FOR EACH SPECIES WAS CALCULATED BY THE SAME METHOD AS MEAN ESTIMATED NUMBER.

TABLE 2. - (CONTINUED).

SAMPLING PERIOD - 02/23/78 1205 THROUGH 02/24/78 0600

SPECIES	N	C F	l S	URVIVAL D D4		MEAN ESTIMATED NUMBER/ 24 HOURS	STANDARD DEVIATION	MEAN ESTIMATED WT/24 HRS (KG)	STANDARD DEVIATION	HEI (G			MAX.
M. AMERICANA H. MUCHALIS M. SAKATILIS	4	4 1 1	0 60 0	5 94 0 40 D 1 00	5	2040 600 120	603.9 1200.0 240.0	51.28 3.44 .72	48.7 6.8 1.4	4.5 4.7 6.0	69.1 7.1 6.0	62.5 77.5 82.5	182.5 87.5 82.5
FOTAL					23	2760		55.44					

N = NUMBER OF 3-MINUTE SAMPLES.

CF = CATCH FREQUENCY (NUMBER OF SAMPLES IN WHICH THE SPECIES APPEARED).

X SURVIVAL: L = LIVE; D = DEAD; D* = DAMAGED.

MEAN ESTIMATED NUMBER PER 24 HOURS FOR EACH SPECIES WAS CALCULATED BY MULTIPLYING THE NUMBER OF SPECIMENS IN EACH 3-MINUTE SAMPLE BY 48D TO OBTAIN A 24-HOUR ESTIMATE. THE ESTIMATED NUMBER FOR EACH SAMPLE WAS SUMMED AND DIVIDED BY THE NUMBER OF SAMPLES TO OBTAIN THE MEAN.

MEAN ESTIMATED WEIGHT PER 24 HOURS FOR EACH SPECIES WAS CALCULATED BY THE SAME METHOD AS MEAN ESTIMATED NUMBER.

TABLE 2. - (CONTINUED).

SAMPLING PERIOD - 02/27/78 1200 THROUGH 02/28/78 0600

SPECIES	N	C F	X S	URV.	D+	TOTAL NUMBER COLLECTED	MEAN ESTIMATED NUMBER/ 24 HOURS	STANDARD DEVIATION	MEAN ESTIMATED WT/24 HRS (KG)	STANDARD DEVIATION	WEI (6) Min.	-		MAX.
G- ACULEATUS	4	2	100	0	3	2	240	277.1	• 5 1	.5	2.0	2.3	52.5	57.5
H. MUCHALIS	4	2	66	33	0	3	360	459.5	2.79	3.4	6.9	8.3	92.5	97.5
M. AMERICANA	4	4	0	٥	100	7	840	459.5	16.54	16.9	2.1	80.5	47.5	172.5
M. SAXATILIS	4	1	Đ	0	100	1	120	240.0	3.96	7.9	33.0	33.0	147.5	147.5
TOTAL						13	1560		23.80					

N = NJMBER OF 3-MINUTE SAMPLES.

CF = CATCH FREQUENCY (NUMBER OF SAMPLES IN WHICH THE SPECIES APPEARED).

I SURVIVAL: L = LIVE; D = DEAD; D+ = DAMAGED.

MEAN ESTIMATED NUMBER PER 24 HOURS FOR EACH SPECIES WAS CALCULATED BY MULTIPLYING THE NUMBER OF SPECIMENS IN EACH 3-MINUTE SAMPLE BY 483 TO OBTAIN A 24-HOUR ESTIMATE. THE ESTIMATED NUMBER FOR EACH SAMPLE WAS SUMMED AND DIVIDED BY THE NUMBER OF SAMPLES TO OBTAIN THE MEAN.

MEAN ESTIMATED WEIGHT PER 24 HOURS FOR EACH SPECIES WAS CALCULATED BY THE SAME METHOD AS MEAN ESTIMATED NUMBER.

IMPINGEMENT MONITORING PROGRAM
SALEM NUCLEAR GENERATING STATION UNIT 1
PSE&G, NEWARK, N.J.

FIN

BASS, A 760311MP.
FACILITY WARNING 10000000000

FACILITY MARNING 10000000000

BUSE SPECIES-FILE., 76031SP. READY

BEAT 76031SYM.402RPTIM

TABLE 3. - IMPINGEMENT - S.N.G.S. SERVICE WATER SYSTEM INTAKE, 1 FEBRUARY 1978 TO 28 FEBRUARY 1978.

SAMPLING PERIOD - 02/01/78

0730

10

02/02/78

0730

NUMBER OF PUMPS 3 - 3

			W£3	GHT	LEN	e1H
SPECIES	NUMBER	.T.(M)	MIN.	MAX.	MIN.	MAX.
H. NUCHALIS	25	280.30	3.6	21.1	72.5	127.5
M. AMERICANA	13	81.10	2.2	9.8	57.5	92.5
I - PUNCTATUS	1	3.50	3.5	3.5	67.5	67.5
M. SANATILIS	2	20.50	6 • D	14.5	67.5	107.5
TOTAL	41	385.40				

TABLE 3. - (CONTINUED).

SAMPLING PERIOD - 02/02/78 0730 TO 02/03/78 0730

NUMBER OF PUMPS 3 - 4


			WEI	6HT	LEN	 бтн
SPECIES	NUMBER	WT.(6M)	MIN.		MIN.	_ K A M
M. AMERICANA	6	46.40	1.1	18.1	57.5	102.5
H. NUCHALIS	5	32.90	1.8	15.8	62.5	117.5
D. CEPEDIANUM	1	46.10	46.1	46.1	157.5	157.5
TOTAL	12	125.40				

TABLE 3. - (CONTINUED).

SAMPLING PERIOD - 02/06/78 0730 TO 02/07/78 0730

NUMBER OF PUMPS 3 - 4

			WE	1 6 H T	LEN	6 T H
SPECIES	NUMBER	HT.(6M)	MIN.	M A X .	M1N.	MAX.
M. AMERICANA	16	214.80	3.9	121.8	57.5	197.5
H. NUCHALIS	4	43.10	5.9	24.4	82.5	117.5
P. FLAVESCENS	1	36.DO	36.0	36.0	147.5	147.5
M. SARATILIS	1	7.30	7.3	7.3	87.5	87.5
TOTAL	22	301.20				



SAMPLING PERIOD - 02/07/78 1500 TO 02/08/78

1500

NUMBER OF PUMPS 3 - 3

			WEIG	нт	LENG	TH	
SPECIES	NUMBER	WT. (GM)	MIN.	MAX	MIN.	MAX	
M. AMERICANA	190	1261.80	2.0	32.4	57.5	132.5	
C. CARPIO	1	33.50	33.5	33.5	112.5	112.5	
P. FLAVESCENS	1	10.20	10.2	10.2	97.5	97.5	
P. HETEROCLITUS	1	6.50	6.2	6.5	77.5	77.5	4
M. SAXATILIS	11	83.60	4.0	12.7	67.5	102.5	
H. NUCHALIS	18	168.30	4.1	24.0		162.5	
TOTAL	222	1563.90					

TABLE 3. - (CONTINUED).

SAMPLING PERIOD - 02/09/78 0730 TO 02/10/78 0730

NUMBER OF PUMPS 3 - 3

			WEI	GHT	LEN	GTH
SPECIES	NUMBER	(Ma).TH	MIN.	MAX.	MIN-	MAX.
M- AMERICANA	96	555.50	2.9	90.0	57.5	87.5
H. NUCHALIS	2	22.10	8 - 8	13.3	92.5	112.5
M- SAXATILIS	1	30-20	30.2	30.2	132.5	132.5
TOTAL	99	607.80				

TABLE 3. - (CONTINUED).

SAMPLING PERIOD - 02/13/78 0730 TO 02/14/76 0730

NUMBER OF PUMPS 3 - 3

			 Kej	6HT	LEN	 61H	
SPECIES	NUMBER	MT.(6M)	MIN.	MAX.	MIN.	MAX.	
H. NUCHALIS F. HETEROCLITUS M. AMERICANA	4 1 1	57.80 2.00 3.20	4.2 2.0 3.2	20.4 2.0 3.2	77.5 42.5 67.5	122.5 42.5 67.5	
TOTAL	6	63.00					

TABLE 3. - (CONTINUED).

SAMPLING PERIOD - 02/15/78 0730 TO 02/16/78 0730

NUMBER OF PUMPS 3 - 3

		AFT	6HT	LEN	6TH	
NUMBER	ut.(GM)	MIN.	MAX.	MIN.	MAX.	
4	26.50	26.5	26.5	107.5	127 5	
3	15.70	3.5	6.3	62.5	77.5	
1	17.50	17.5	17.5	112.5	112.5	
1	5 • 5 0	5.5	5.5	82.5	82 • 5	
i 						
6	65.20					
	1 3 1	1 26.50 3 15.70 1 17.50 1 5.50	1 26.50 26.5 3 15.70 3.5 1 17.50 17.5 1 5.50 5.5	1 26.50 26.5 26.5 3 15.70 3.5 6.3 1 17.50 17.5 17.5 1 5.50 5.5 5.5	1 26.50 26.5 26.5 127.5 3 15.70 3.5 6.3 62.5 1 17.50 17.5 17.5 112.5 1 5.50 5.5 5.5 82.5	1 26.50 26.5 26.5 127.5 127.5 3 15.70 3.5 6.3 62.5 77.5 1 17.50 17.5 17.5 112.5 112.5 1 5.50 5.5 5.5 82.5 82.5

TABLE 3. - (CONTINUED).

SAMPLING PERIOD - 02/16/78 0730 TO 02/17/78 0730

NUMBER OF PUMPS 3 - 3

				 GHT	LEN	 6TH
SPECIES	NUMBER	HT-(6M)	MIN.	MAX.	MIN.	MAX.
H. NUCHALIS	4	40.40	6.5	16.9	77.5	117.5
M. MENIDIA	1	2.50	2.5	2.5	62.5	62.5
M. AMERICANA	19	120.10	3.0	11.9	57.5	92.5
M. SAKATILIS	4	28.50	5.3	14.4	72.5	107.5
F. HETEROCLITUS	2	6.70	2.2	4.5	47.5	72.5
C. CARPID	1	61.20	61.2	61.2	137.5	137.5
P. FLAVESCENS	1	5 •00	5.0	5.0	77.5	77.5
TOTAL	32	264.40				

TABLE 3. - (CONTINUED).

SAMPLING PERIOD - 02/21/78 0730 TO 02/22/78 0730

NUMBER OF PUMPS 3 - 3

			WE I	6HT	LEN	 6тн
SPECIES	NUMBER	⊌T.(6M)	MIN.	MAX.	MIN.	. X A M
N. AMERICANA	9	46.20	2.5	8.6	52.5	77.5
H. NUCHALIS	3	39.20	7.6	20.0	92.5	127.5
6. ACULEATUS	1	1.70	1.7	1.7	62.5	62.5
TOTAL	13	87.10	•			

TABLE 3. - (CONTINUED).

SAMPLING PERIOD - 02/22/78 0730 TO 02/23/78 0730

NUMBER OF PUMPS 3 - 3

~~~~~~~			WE	16HT	LEN	 6 T H
SPECIES	NUMBER	ut.(Ma)	MIN.	MAX.	MIN.	MÁX.
P. FLAVESCENS	2	166.20	10.8	155.4	102.5	222.5
M. AMERICANA	11	115.30	2.5	45.9	62.5	152.5
H. NUCHALIS	5	82.50	3.2	20.0	72.5	147.5
I. NEBULOSUS	1	11.10	11.1	11.1	92.5	92.5
M. SALATILIS	2	20.00	8.5	11.5	72.5	102.5
***************************************						
TOTAL	21	395.10				

TABLE 3. - (CONTINUED).

SAMPLING PERIOD - 02/23/78 0730 TO 02/24/78 0730

NUMBER OF PUMPS 3 - 3

PACALCO				6HT	LEN		 	
SPECIES	NUMBER	AT.(6M)		MAX.	HIN.	. KAM	 	
M. AMERICANA	7	44.80	3.1	8.8	67.5	82.5		
H. NUCHALIS	5	22.50	3.1 3.6		72.5			
***************************************							 	
TOTAL	9	67.30						

TABLE 3. - (CONTINUED).

SAMPLING PERIOD - 02/27/78 0730 TO 02/28/78 0730

NUMBER OF PUMPS 3 - 3

			WEIGHT		LENGTH	
SPECIES	NUMBER	HT.(6M)	MIN.	MAX.	MIN.	MAX.
H. NUCHALIS	1	9.00	9.0	9.0	102.5	102.5
P. FLAVESCENS	1	8.10	8.1	8.1	87.5	87.5
M. AMERICANA	1	4.90	4.9	4.9	62.5	62.5
TOTAL .	3	22.00				

IMPINGEMENT MONITORING PROGRAM SALEM NUCLEAR GENERATING STATION UNIT 1 PSE&G, NEWARK, N.J.

SBRKPT PRINTS

,