


# Regional Tectonics and Seismicity of Eastern Nebraska

Annual Report  
July 1980 - June 1981



---

R. H. Burchett

Institute of Agriculture and Natural Resources  
University of Nebraska-Lincoln

Prepared for  
U.S. Nuclear Regulatory  
Commission

NOTICE

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, or any of their employees, makes any warranty, expressed or implied, or assumes any legal liability or responsibility for any third party's use, or the results of such use, of any information, apparatus product or process disclosed in this report, or represents that its use by such third party would not infringe privately owned rights.

Available from

GPO Sales Program  
Division of Technical Information and Document Control  
U. S. Nuclear Regulatory Commission  
Washington, D. C. 20555

Printed copy price: \$5.50

and

National Technical Information Service  
Springfield, Virginia 22161

# Regional Tectonics and Seismicity of Eastern Nebraska

Annual Report  
July 1980 - June 1981

---

Manuscript Completed: October 1981

Date Published: November 1981

---

Prepared by  
R. R. Burnett

Nebraska Geological Survey  
Institute of Agriculture and Natural Resources  
University of Nebraska-Lincoln  
Lincoln, NE 68588-0517

Prepared for  
Division of Health, Siting and Waste Management  
Office of Nuclear Regulatory Research  
U.S. Nuclear Regulatory Commission  
Washington, D.C. 20555  
NRC FIN B5774

#### Availability of Reference Materials Cited in NRC Publications

Most documents cited in NRC publications will be available from one of the following sources:

1. The NRC Public Document Room, 1717 H Street., N.W.  
Washington, DC 20555
2. The NRC/GPO Sales Program, U.S. Nuclear Regulatory Commission,  
Washington, DC 20555
3. The National Technical Information Service, Springfield, VA 22161

Although the listing that follows represents the majority of documents cited in NRC publications, it is not intended to be exhaustive.

Referenced documents available for inspection and copying for a fee from the NRC Public Document Room include NRC correspondence and internal NRC memoranda; NRC Office of Inspection and Enforcement bulletins, circulars, information notices, inspection and investigation notices; Licensee Event Reports; vendor reports and correspondence; Commission papers; and applicant and licensee documents and correspondence.

The following documents in the NUREG series are available for purchase from the NRC/GPO Sales Program: formal NRC staff and contractor reports, NRC-sponsored conference proceedings, and NRC booklets and brochures. Also available are Regulatory Guides, NRC regulations in the *Code of Federal Regulations*, and *Nuclear Regulatory Commission Issuances*.

Documents available from the National Technical Information Service include NUREG series reports and technical reports prepared by other federal agencies and reports prepared by the Atomic Energy Commission, forerunner agency to the Nuclear Regulatory Commission.

Documents available from public and special technical libraries include all open literature items, such as books, journal and periodical articles, transactions, and codes and standards. *Federal Register* notices, federal and state legislation, and congressional reports can usually be obtained from these libraries.

Documents such as theses, dissertations, foreign reports and translations, and non-NRC conference proceedings are available for purchase from the organization sponsoring the publication cited.

Single copies of NRC draft reports are available free upon written request to the Division of Technical Information and Document Control, U.S. Nuclear Regulatory Commission, Washington, DC 20555.

TABLE OF CONTENTS

	Page
ASTRACT . . . . .	1
STRUCTURE OF THE ELMONT LIMESTONE IN CENTRAL OTOE COUNTY, NEBRASKA, R. R. Burchett and M. R. Bolitho .	2
Introduction . . . . .	2
Scope of Investigation . . . . .	5
Results of Investigation . . . . .	10
References . . . . .	14
EARTHQUAKE MONITORING NETWORK IN NEBRASKA, R. R. Burchett and Russell Smith . . . . .	15
GRAVITY AND GROUND MAGNETIC INVESTIGATIONS IN EASTERN NEBRASKA, R. F. Diffendal Jr. and R. R. Burchett . . . . .	21
Introduction . . . . .	21
Gravity Study . . . . .	21
Ground Magnetic Study . . . . .	25
References . . . . .	44
SOME PRECAMBRIAN GABBROIC ROCKS FROM SOUTHEAST NEBRASKA, S. B. Treves . . . . .	45
Introduction . . . . .	45
Location . . . . .	45
Summary and Conclusions . . . . .	53
References . . . . .	54
APPENDIX A . . . . .	55

LIST OF ILLUSTRATIONS

Figure	Page
1 Principal Structural Features of Nebraska . . . . .	3
2 Structural Contour - Top of the Elmont Limestone (Pennsylvanian) in Central Otoe County. . . . .	4
3 Location of Test Holes Drilled During 1981 in Central Otoe County . . . . .	6
4 Composite Section of Pennsylvanian Rocks (Virgil Series) Drilled in Central Otoe County. . . . .	7
5 Composite Section of Permian Rocks (Big Blue Series) Drilled in Central Otoe County . . . . .	8
6 Geologic Bedrock of Central Otoe County . . . . .	9
7 Elevation of Precambrian Surface. . . . .	12
8 Location of Earthquake Monitoring Stations in Nebraska as of July 1, 1981 . . . . .	16
9 Microearthquakes in Nebraska. . . . .	20
10 Location of Gravity and Magnetic Studies in Eastern Nebraska. . . . .	22
11 Location of Bouguer Gravity Stations in Eastern Nebraska. . . . .	24
12 Bouguer Gravity of Central Otoe County. . . . .	26
13 Location of Bouguer Gravity Stations in Clay County. . . . .	27
14 Location of Bouguer Gravity Stations in Fillmore County. . . . .	28
15 Location of Bouguer Gravity Station in Hamilton County. . . . .	29

Figure	Page
16 Location of Bouguer Gravity Stations in Nuckolls County. . . . .	30
17 Location of Bouguer Gravity Stations in Polk County. . . . .	31
18 Location of Bouguer Gravity Stations in Thayer County. . . . .	32
19 Location of Bouguer Gravity Stations in York County. . . . .	33
20 Bouguer Gravity of Clay County. . . . .	34
21 Bouguer Gravity of Fillmore County. . . . .	35
22 Bouguer Gravity of Hamilton County. . . . .	36
23 Bouguer Gravity of Nuckolls County. . . . .	37
24 Bouguer Gravity of Polk County. . . . .	38
25 Bouguer Gravity of Thayer County. . . . .	39
26 Bouguer Gravity of York County. . . . .	40
27 Bouguer Gravity Anomaly Map in Eastern Nebraska . .	41
28 Total Magnetic Intensity of Northeastern Cass and Eastern Sarpy Counties. . . . .	43
29 Index Map of Southeast Nebraska and Northeast Kansas . . . . .	46
30 $^{40}\text{Ar}/^{39}\text{Ar}$ Age Spectrum Diagram for St-1 and St-2 . .	52

LIST OF TABLES

Table		Page
1	Microearthquakes in Nebraska . . . . .	19
2	Modal Analyses of Specimens from St-1 and St-2 .	43
3	Chemical Compositions . . . . .	50
4	Age Spectrum Analysis Analytical Data . . . . .	51

## ABSTRACT

This annual report presents and interprets the information obtained by the Conservation and Survey Division (Nebraska Geological Survey) during contract year July 1, 1980, to June 30, 1981, under contract NRC-04-76-315 with the U. S. Nuclear Regulatory Commission. The information pertains to the geology, structure, tectonics, and seismicity of eastern Nebraska with emphasis on central Otoe County. Some of the information presented here results from a combination of studies begun in earlier years but the greater part results from studies begun during the contract year.

The scope of the studies is summarized as follows:

1. Rock outcrops in Otoe County were reexamined and reevaluated, and 22 test holes were drilled to determine the altitude of the upper surface of the Elmont Limestone of Pennsylvanian age;
2. One new seismograph was installed in eastern Nebraska;
3. Gravity surveys in eastern Nebraska were extended;
4. Ground magnetic surveys in Otoe County were made and evaluated.
5. Age dating of two Precambrian cores.

Discussion of the results of these studies constitute the remainder of this report.


STRUCTURE OF THE ELMONT LIMESTONE IN CENTRAL OTOE COUNTY, NEBRASKA

R. R. Burchett and M. F. Bolitho

Introduction


A specific site area in central Otoe County, Nebraska was chosen for study in 1980-81 because it overlies the southwestern extension of the Union Fault (fig. 1). The Union Fault defines the southern margin of the northeast-southwest trending midcontinent gravity anomaly. Southward from it the rock strata either are downthrown or dip steeply into the Forest City Basin.

The primary purposes of this investigation were to acquire, by drilling test holes, subsurface data needed to correlate buried strata with strata that crop out and to gain a better understanding of buried structural features in eastern Nebraska. The top of the Elmont Limestone, a member of the Emporia Formation of the Wabaunsee Group (Virgil Series) of the Pennsylvanian System, was chosen as a datum plane for a structure map because the Elmont underlies most of the study area at a shallow depth and is easily identified in rock cuttings obtained by drilling. All available data on the altitude of this surface in outcrops and in test holes were used as control points for contour lines depicting the configuration of the Elmont's upper surface (fig. 2).



PRINCIPAL STRUCTURAL FEATURES OF NEBRASKA (Carlson, 1970)

Figure 1



Compilation October 1981  
Principal Investigator: R. R. Burchett

1/2 1 MILES  
1/2 1 KILOMETERS

Prepared by R. R. Burchett  
Nebraska Geological Survey  
Conservation and Survey Division  
Institute of Agriculture and Natural Resources  
University of Nebraska-Lincoln  
Supported by the United States Nuclear Regulatory Commission  
under contract NRC-O4-76-315



STRUCTURAL CONTOUR-TOP OF ELMONT LIMESTONE (PENNSYLVANIAN) IN CENTRAL OTOE COUNTY


Figure 2

### Scope of Investigation

Twenty-two shallow rotary test holes were drilled in central Otoe County (fig. 3) by the Rieschick Drilling Company of Falls City, Nebraska. The holes averaged about 182 feet (55.6 m) in depth, and each was logged electrically as well as thru visual examination of the rock cuttings. Some of the holes penetrated the Elmont Limestone; others were drilled to an identifiable horizon whose height above or depth below the Elmont is known, thus providing an Elmont datum. The holes were drilled in May and June of 1981 under a cooperative agreement between the Conservation and Survey Division (Nebraska Geological Survey) and the U.S. Nuclear Regulatory Commission (Contract NRC-04-76-315).


A binocular microscope was used for examination of the cuttings. Logs of the test holes, based on descriptions of the cuttings, are presented in Appendix A. Figures 4 and 5 show a composite section of the Lower Permian and Upper Pennsylvanian rocks drilled in the study area, together with a composite electric log of those rocks.

The geologic map illustrated in figure 6 shows the distribution of groups ranging in age from Late Pennsylvanian to Early Permian. Location of bedrock outcrops, mostly along valley sides, are shown in solid black. A careful reexamination of outcrops in the area provided additional data for mapping purposes. By determining the altitude of many outcrops, the investigators obtained additional vertical control points for the structure map.



LOCATION OF TEST HOLES DRILLED DURING 1981	CENTRAL OTOE COUNTY
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
11	
12	
13	
14	
15	
16	
17	
18	
19	
20	
21	
22	
23	
24	
25	
26	
27	
28	
29	
30	
31	
32	
33	
34	
35	
36	
37	
38	
39	
40	
41	
42	
43	
44	
45	
46	
47	
48	
49	
50	
51	
52	
53	
54	
55	
56	
57	
58	
59	
60	
61	
62	
63	
64	
65	
66	
67	
68	
69	
70	
71	
72	
73	
74	
75	
76	
77	
78	
79	
80	
81	
82	
83	
84	
85	
86	
87	
88	
89	
90	
91	
92	
93	
94	
95	
96	
97	
98	
99	
100	
101	
102	
103	
104	
105	
106	
107	
108	
109	
110	
111	
112	
113	
114	
115	
116	
117	
118	
119	
120	
121	
122	
123	
124	
125	
126	
127	
128	
129	
130	
131	
132	
133	
134	
135	
136	
137	
138	
139	
140	
141	
142	
143	
144	
145	
146	
147	
148	
149	
150	
151	
152	
153	
154	
155	
156	
157	
158	
159	
160	
161	
162	
163	
164	
165	
166	
167	
168	
169	
170	
171	
172	
173	
174	
175	
176	
177	
178	
179	
180	
181	
182	
183	
184	
185	
186	
187	
188	
189	
190	
191	
192	
193	
194	
195	
196	
197	
198	
199	
200	
201	
202	
203	
204	
205	
206	
207	
208	
209	
210	
211	
212	
213	
214	
215	
216	
217	
218	
219	
220	
221	
222	
223	
224	
225	
226	
227	
228	
229	
230	
231	
232	
233	
234	
235	
236	
237	
238	
239	
240	
241	
242	
243	
244	
245	
246	
247	
248	
249	
250	
251	
252	
253	
254	
255	
256	
257	
258	
259	
260	
261	
262	
263	
264	
265	
266	
267	
268	
269	
270	
271	
272	
273	
274	
275	
276	
277	
278	
279	
280	
281	
282	
283	
284	
285	
286	
287	
288	
289	
290	
291	
292	
293	
294	
295	
296	
297	
298	
299	
300	
301	
302	
303	
304	
305	
306	
307	
308	
309	
310	
311	
312	
313	
314	
315	
316	
317	
318	
319	
320	
321	
322	
323	
324	
325	
326	
327	
328	
329	
330	
331	
332	
333	
334	
335	
336	
337	
338	
339	
340	
341	
342	
343	
344	
345	
346	
347	
348	
349	
350	
351	
352	
353	
354	
355	
356	
357	
358	
359	
360	
361	
362	
363	
364	
365	
366	
367	
368	
369	
370	
371	
372	
373	
374	
375	
376	
377	
378	
379	
380	
381	
382	
383	
384	
385	
386	
387	
388	
389	
390	
391	
392	
393	
394	
395	
396	
397	
398	
399	
400	
401	
402	
403	
404	
405	
406	
407	
408	
409	
410	
411	
412	
413	
414	
415	
416	
417	
418	
419	
420	
421	
422	
423	
424	
425	
426	
427	
428	
429	
430	
431	
432	
433	
434	
435	
436	
437	
438	
439	
440	
441	
442	
443	
444	
445	
446	
447	
448	
449	
450	
451	
452	
453	
454	
455	
456	
457	
458	
459	
460	
461	
462	
463	
464	
465	
466	
467	
468	
469	
470	
471	
472	
473	
474	
475	
476	
477	
478	
479	
480	
481	
482	
483	
484	
485	
486	
487	
488	
489	
490	
491	
492	
493	
494	
495	
496	
497	
498	
499	
500	
501	
502	
503	
504	
505	
506	
507	
508	
509	
510	
511	
512	
513	
514	
515	
516	
517	
518	
519	
520	
521	
522	
523	
524	
525	
526	
527	
528	
529	
530	
531	
532	
533	
534	
535	
536	
537	
538	
539	
540	
541	
542	
543	
544	
545	
546	
547	
548	
549	
550	
551	
552	
553	
554	
555	
556	
557	
558	
559	
560	
561	
562	
563	
564	
565	
566	
567	
568	
569	
570	
571	
572	
573	
574	
575	
576	
577	
578	
579	
580	
581	
582	
583	
584	
585	
586	
587	
588	
589	
590	
591	
592	
593	
594	
595	
596	
597	
598	
599	
600	
601	
602	
603	
604	
605	
606	
607	
608	
609	
610	
611	
612	
613	
614	
615	
616	
617	
618	
619	
620	
621	
622	
623	
624	
625	
626	
627	
628	
629	
630	
631	
632	
633	
634	
635	
636	
637	
638	
639	
640	
641	
642	
643	
644	
645	
646	
647	
648	
649	
650	
651	
652	
653	
654	
655	
656	
657	
658	
659	
660	
661	
662	
663	
664	
665	
666	
667	
668	
669	
670	
671	
672	
673	
674	
675	
676	
677	
678	
679	
680	
681	
682	
683	
684	
685	
686	
687	
688	
689	
690	
691	
692	
693	
694	
695	
696	
697	
698	
699	
700	
701	
702	
703	
704	
705	
706	
707	
708	
709	
710	
711	
712	
713	
714	
715	
716	
717	
718	
719	
720	
721	
722	
723	
724	
725	
726	
727	
728	
729	
730	
731	
732	
733	
734	
735	
736	
737	
738	
739	
740	
741	
742	
743	
744	
745	
746	
747	
748	
749	
750	
751	
752	
753	
754	
755	
756	
757	
758	
759	
760	
761	
762	
763	</

Figure 3



Prepared by R. H. Johnson and W. H. Bell  
Geological Survey, Kansas  
for the State Geological Survey  
Division of Mineral Resources  
Department of Revenue, State of Kansas  
July 1952

COMPOSITE SECTION OF PERMIAN ROCKS (BIG BLUE SERIES) DRILLED IN CENTRAL OTOE COUNTY

Figure 4




Figure 5




Figure 6

Most upland and lowland areas are mantled by unconsolidated Quaternary deposits such as loess, till, and alluvium. The maximum aggregate thickness of these deposits is 400 feet (122 m).


#### Results of Investigation

Interpretations based on drilling results and field observations during this investigation are summarized as follows:

- (1) The Union Fault probably is not a single structure, as previously interpreted. Instead, it is a complex zone of faults and steep dips affecting rocks as young as Permian in age.
- (2) Pennsylvanian and Permian age rocks are faulted along the Union Fault. The direction and angle of dip exhibited by these rocks, on both sides of the Union Fault, differ markedly within very short distances, thus indicating that the structural pattern is highly complex. The greatest dip measured about 15°SE, just east of the area of study, is in the SW sec. 6, T. 9 N., R. 13 E., in Otoe County (Burchett, 1980).
- (3) Only two faults have actually been observed in outcrops near the study area, and both are located along the Union Fault. One is in the NW NW sec. 21, T. 10 N., R. 14 E., Cass County, and was observed by Condra and Reed (1938) many years ago before quarrying destroyed the outcrop. Throw along this fault was reported to be 8.8 feet (2.7 m) in the Deer Creek Formation of Pennsylvanian age, but no mention was made whether that or any other faults involved Quaternary deposits. The other fault is in the NE SW

sec. 6, T. 9 N., R. 13 E., Otoe County. Here the Lecompton Formation of Pennsylvanian age is displaced about 3 feet (0.9 m). Overlying Quaternary deposits do not appear to be faulted.

- (4) Displacements along the faults not observable in outcrops but indicated on figure 2 are interpretive for the following reasons: Thick glacial deposits mantle the bedrock throughout most of the study area; outcrops are of small areal extent and consist of strata so similar that their correlation is difficult; and the distance between outcrops and test holes available for interpretation ranges from 0.1 (0.161 km) to 1 mile (1.61 km). The greatest interpreted throw of the Elmont Limestone on the Union Fault, about 200 feet (61.0 m), is in the southwestern part of the study area.
- (5) The altitude of the top of the Elmont Limestone (fig. 2) ranges from about 800 feet (243.8 m) above mean sea level in the southwestern part of the study area to more than 1100 feet (335.3 m) north of the Union Fault.
- (6) Faults and/or steep dips in the area probably reflect faulting and steep dips at depth. Deeper structures are indicated by contours drawn on the base of the Hertha Limestone, a formation in the Kansas City Group of the Missouri Series of the Pennsylvanian System (Burchett, 1978) and on the surface of Precambrian rocks (Carlson, 1967).
- (7) Several deep drill holes have penetrated the Precambrian age rocks in Cass, Otoe, Lancaster, and adjacent counties to the south. Locations of these holes and the elevation of the Precambrian surface are shown in figure 7.



ELEVATION OF PRECAMBRIAN SURFACE

Figure 7

Interpretation of these elevations indicates that faulting probably occurred in the Precambrian rocks along the Union Fault.

#### REFERENCES

- Burchett, R. R. 1978. Regional tectonics and seismicity of eastern Nebraska, Annual Report, June 1, 1976--June 1, 1977. Available from National Technical Information Service, Springfield, VA 22161. (NUREG/CR-0053)
- Burchett, R. R. 1980. Regional tectonics and seismicity of eastern Nebraska, Annual Report, June 1978--May 1979, p. 10. Available from National Technical Information Service, Springfield, VA 22161. (NUREG/CR-1328)
- Carlson, M. P. 1967. Precambrian well data in Nebraska including rock type and surface configuration. Nebraska Geological Survey Bulletin 25. Lincoln, Nebraska: Conservation and Survey Division, University of Nebraska.
- Carlson, M. P. 1970. Distribution and subdivision of Precambrian and Lower and Middle Paleozoic rocks in the subsurface of Nebraska. Nebraska Geological Survey Report of Investigations 3. Lincoln, Nebraska: Conservation and Survey Division, University of Nebraska.
- Condra, G. E., and Reed, E. C. 1938. The Redfield anticline of Nebraska and Iowa. Nebraska Geological Survey Paper 12. Lincoln, Nebraska: Conservation and Survey Division, University of Nebraska.

## EARTHQUAKE MONITORING NETWORK IN NEBRASKA


R. R. Burchett & Russell Smith

The Conservation and Survey Division (Nebraska Geological Survey) and the Geology Department of the University of Nebraska-Lincoln collaborated in the installation of a portable micro-earthquake station in Nebraska during contract year 1980-81. This station, near Dwight, was a replacement for the station near Columbus which developed high background noise.

As of July 1, 1981, a total of seven earthquake monitoring stations were operating in Nebraska. Locations of these stations are shown in figure 8. Precise location and other pertinent information about these stations are given below:

Site designation LCN. Location: E/2 NE NE SE sec. 23, T. 10 N., R. 6 E., lat.  $40^{\circ} 48' 42''$  N., long.  $96^{\circ} 42' 07''$  W.; on University of Nebraska-Lincoln campus at Lincoln in Lancaster County, Nebraska. Instrument altitude: 1,165 ft. (355 m) msl datum. Operation begun September 1, 1977.

Site designation DTN: Location: Center SE NW SW sec. 6, T. 12 N., R. 4 E.; lat.  $41^{\circ} 01' 18''$  N., long.  $97^{\circ} 02' 09''$  W.; on James Tesina farm near Dwight, in Seward County, Nebraska. Instrument altitude: 1,523 ft. (464 m) msl datum. Operation begun August 25, 1980.



**LOCATION OF EARTHQUAKE MONITORING STATIONS IN NEBRASKA AS OF JULY 1, 1980**

Figure 8

Site designation JHN. Location: NW corner SW NE sec. 36, T. 6 N., R. 12 E.; lat.  $40^{\circ} 26' 49''$  N., long.  $96^{\circ} 01' 03''$  W.; on Douglas Boellstorff farm near Johnson in Nemaha County, Nebraska. Instrument altitude: 1,080 ft. (329 m) msl datum. Operation begun December 5, 1978.

Site designation HMN. Location. NW SW SE NW sec. 26, T. 4 N., R. 5 E.; lat.  $40^{\circ} 17' 11''$  N., long.  $96^{\circ} 50' 08''$  W.; on Homestead National Monument of America Site near Beatrice in Gage County, Nebraska. Instrument altitude: 1,207 ft. (368 m) msl datum. Operation begun June 20, 1979.

Site designation SGN. Location: SE NW NE SE sec. 30, T. 13 N., R. 11 E.; lat.  $41^{\circ} 3' 57''$  N., long.  $96^{\circ} 13' 7''$  W.; on Leroy Nielson farm near Springfield in Sarpy County, Nebraska. Instrument altitude: 1,140 ft. (348 m) msl datum. Operation begun June 20, 1980.

Site designation MKN. Location: NW SE NW sec. 25, T. 5 N., R. 26 W.; lat.  $40^{\circ} 22.44'$  N.,  $100^{\circ} 13.50'$  W.; near Medicine Creek Dam in Frontier County, Nebraska. Instrument altitude 2,395 ft. (730 m) msl datum. Operation begun March 11, 1979.

Site designation INN. Location. NW SW SW sec. 29, T. 3 N., R. 27 W.; lat.  $40^{\circ} 11.55'$  N.,  $100^{\circ} 24.01'$  W.; near Indianola in Red Willow County, Nebraska. Instrument altitude 2,500 ft. (762 m) msl datum. Operation begun September 15, 1979.

All stations are equipped with Geotech Portacorders model RV-320 and Geotech model S-13 seismometers except the HMN and MKN stations are equipped with Geotech model S-500 seismometers and INN with a GSC-20D seismometer.

Each station is operated by a volunteer who agreed to (1) permit installation of a seismometer vault on his property, (2) provide housing for the amplifier-filter-recorder-clock system and the WWV time-signal radio receiver, and (3) change the seismograph recorder charts, replenish the ink supply, and reset the clock whenever necessary on a 365-day-per-year basis. All seismographic records with the exception of Medicine Creek and Indianola are sent to the University of Nebraska-Lincoln for preliminary analysis and then are forwarded to the Kansas Survey for more detailed analysis. The seismographic records from Medicine Creek and Indianola are sent directly to the Kansas Geological Survey.

The Nebraska seismometers, together with seismometers in the adjacent part of Kansas, provide continuous coverage of seismic activity along the buried Nemaha Ridge.

Thirty microearthquakes centering in Nebraska were recorded between August 1977 and July 1981. Table 1 and figure 9 show the location of these microearthquakes.

TABLE I

## MICROEARTHQUAKES IN NEBRASKA


Map Ref.	Date	Origin Time (UTC) <sup>1</sup>	Latitude Degrees North	Longitude Degrees West	Depth <sup>2</sup> (Km)	Magnitude <sup>3</sup>
1	1977 Aug 18	10:34:25.21	41:24.90	98:28.07	5.00	2.5
2	1977 Dec 1	13:04:34.20	40:18.53	100:22.00	5.00	2.3
3	1977 Dec 1	13:22:38.57	40:12.52	100:17.87	5.00	2.4
4	1978 Jan 13	20:15:33.39	40:05.58	95:42.00	5.00	1.7
5	1978 Feb 3	0:25:47.62	40:01.92	100:20.00	5.00	2.4
6	1978 Sep 14	8:06:18.59	40:53.76	100:22.00	5.00	2.2
7	1979 Apr 8	22:46:10.41	40:58.12	98:33.83	0.67	2.4
8	1979 Jun 6	16:16:21.91	40:08.61	100:20.88	1.00	2.5
9	1979 Jun 12	11:13:11.88	40:24.30	90:03.26	2.07	1.8
10	1979 Jul 16	0:03:48.18	40:10.07	100:17.22	5.00	2.7
11	1979 Jul 16	1:34:20.32	40:11.59	100:20.70	5.00	2.5
12	1979 Jul 16	5:27:01.42	40:11.45	100:20.00	9.08	1.3
13	1979 Jul 15	6:08:09.89	40:11.32	100:20.73	11.09	1.5
14	1979 Jul 16	7:05:56.02	40:12.00	100:19.90	7.08	1.1
15	1979 Jul 24	4:16:45.09	40:12.47	100:26.00	0.88	2.2
16	1979 Jul 24	8:04:46.26	40:27.94	99:37.38	0.87	1.9
17	1979 Aug 2	4:16:21.66	40:10.34	100:21.44	0.84	2.5
18	1979 Aug 13	11:09:47.65	40:06.80	100:30.10	1.50	1.7
19	1979 Aug 14	23:59:31.37	40:10.39	100:20.58	1.76	1.5
20	1979 Aug 15	6:45:53.87	40:08.68	100:20.34	1.01	1.5
21	1979 Aug 15	16:07:07.14	40:08.49	100:26.43	1.23	1.3
22	1979 Aug 31	8:00:11.70	40:08.31	100:20.22	1.03	2.2
23	1979 Nov 19	4:58:43.40	40:14.86	100:02.17	13.57	1.5
24	1979 Nov 29	22:02:31.21	40:09.80	100:21.64	3.15	1.9
25	1980 Apr 26	14:21:48.50	40:43.99	99:43.91	5.00	2.3
26	1980 Aug 13	5:50:11.83	41:53.59	97:06.01	9.90	2.1
27	1981 Mar 13	12:42:16.68	40:53.44	99:41.67	5.00	2.4
28	1981 Mar 20	5:09:48.17	40:10.31	100:19.60	1.75	1.4
29	1981 Apr 20	18:18:15.28	41:01.02	97:49.49	20.80	2.4
30	1981 Jun 26	18:55:02.07	41:31.75	97:39.06	5.00	2.8

<sup>1</sup>(UTC) Coordinated Universal Time.--Subtract 6 hours for Central Standard Time.

<sup>2</sup>Depth calculated in kilometers or fixed at 5.00 km.

<sup>3</sup>Duration magnitude calculated from equation derived by Oklahoma Geological Observatory.

20



## MICROEARTHQUAKES IN NEBRASKA

Figure 9

GRAVITY AND GROUND MAGNETIC INVESTIGATIONS IN EASTERN NEBRASKA


R. F. Diffendal, Jr. and R. R. Burchett

Introduction

The purpose of this investigation was to extend the area covered by gravity and ground magnetic data related to the Nemaha Uplift and associated structures in eastern Nebraska (fig. 10). Gravity data were collected from Clay, Fillmore, Hamilton, Nuckolls, Polk, Thayer, and York counties. Additional ground magnetic data and gravity data were collected for central Otoe County.

Gravity Study

Gravimetric surveying is a geophysical technique that aids in interpretation of subsurface geologic structure. A Worden Master, a very sensitive weighing device which measures differences in gravity intensity (acceleration) at specific points on the earth's surface, is used in this type of surveying. The principal causes of gravity anomalies generally are either the dissimilar compositions of crystalline basement rocks and overlying sedimentary rocks or density contrasts produced by vertical or horizontal displacements of rocks. However, other phenomena also can contribute to differences in gravity intensity and in combination with the principal causes can make gravity data somewhat ambiguous.



LOCATIONS OF GRAVITY AND MAGNETIC STUDIES IN EASTERN NEBRASKA

Figure 10

All gravity base stations in the Nebraska Gravity Network are tied to the National Gravity Network base station located at the Lincoln Municipal Airport. Gravity stations established during this study were located at section corners and in some cases half-mile section lines (fig. 11). Land-surface altitudes at most stations could be determined directly from 7.5-minute topographic quadrangles. For other stations, land-surface altitudes were established from topographic contours and are accurate within  $\pm 2$  feet.

Field gravity data were corrected for latitude, meter drift and elevation; earth tides were not considered. The maximum error in station altitudes was enough to produce variations of  $\pm 0.2$  milligal. Slight inaccuracies in station locations--none greater than 0.1 minute of latitude--could have produced errors of no more than  $\pm 0.08$  milligal. A  $\pm 0.02$  milligal error was possible from either tidal or instrumental sources. Therefore, the maximum possible was  $\pm 0.3$  milligal but generally was much less because some errors were compensatory.

Bouguer values were calculated from assumed computational density values of 2.5, 2.6, and 2.67 gm/cm<sup>3</sup>. The average density value of 2.69 gm/cm<sup>3</sup> obtained by Muehlberger and others (1964) for two samples of Precambrian crystalline rocks indicated that 2.67 gm/cm<sup>3</sup> would be the most realistic Bouguer computational density value. These three densities were used to compute the efforts of observed gravity versus theoretical gravity for anomaly interpretation.

In central Otoe County 253 gravity stations, in addition to those previously established, were used to produce a

LOCATION OF BOUGUER GRAVITY STATIONS  
IN EASTERN NEBRASKA




Figure 11

hand-contoured Bouguer gravity map (fig. 12). In addition about 980 gravity stations were occupied in Clay (fig. 13), Fillmore (fig. 14), Hamilton (fig. 15), Nuckolls (fig. 16), Polk, (fig. 17), Thayer (fig. 18), and York (fig. 19). Data collected at these stations were used to produce Bouguer gravity maps for each of the counties (figs. 20, 21, 22, 23, 24, 25, & 26). These data were also integrated with previously collected data to generate a comprehensive regional Bouguer gravity map of eastern Nebraska (fig. 27).

#### Ground Magnetic Study

The instrument used in the ground magnetic study was a Geometrics® 826 Proton Magnetometer. This instrument measures the total magnetic field with an accuracy of  $\pm 1$  gamma. Magnetic anomalies are produced by differences in the degree of rock magnetization (polarization). Although, to a large extent, magnetic intensity is related directly to the percentage of the mineral magnetite in rocks, depth to the magnetic source rocks is another variable that affects the measurements.

During field operations about 300 magnetic stations were established at 0.5- and 1.0- mile spacings in central Otoe County. Sites selected as magnetic stations were well-removed from artificial magnetic sources such as fences, power lines, and pipelines. Corrections for diurnal magnetic drift were made by making observations at a base station at 3-hour intervals and then using those observations to correct, by linear interpolation, the readings made at field stations between base stations. No corrections






Figure 12



LOCATION OF BOUGUER GRAVITY STATIONS IN CLAY COUNTY


Figure 13




LOCATION OF BOUGUER GRAVITY STATIONS IN FILLMORE COUNTY

Figure 14

# LOCATION OF BCUGUER GRAVITY STATIONS IN HAMILTON COUNTY




Scale  
Figure 15




LOCATION OF BOUGUER GRAVITY STATIONS IN NUCKOLLS COUNTY

Figure 16




LOCATION OF BOUGUER GRAVITY STATIONS IN POLK COUNTY


Figure




LOCATION OF BOUGUER GRAVITY STATIONS IN THAYER COUNTY

Figure 18






Isogal-Bouguer gravity contours  
 Contour interval, 5 milligals. Hachures  
 indicate closed areas of lower gravity  
 values. Density used in Bouguer  
 calculations, 2.67 gm/cc. Gravity values  
 tied to National Gravity Network




### BOUGUER GRAVITY OF CLAY COUNTY

Figure 20



### [sogg]-Bouguer gravity contours

Contour interval, 5 milligals. Hachures indicate closed areas of lower gravity values. Density used in Bouguer calculations, 2.67 gm/cc. Gravity values tied to National Gravity Network



BOUGUER GRAVITY OF FILLMORE COUNTY

Figure 21

## BOUGUER GRAVITY OF HAMILTON COUNTY

Isogal-Bouguer gravity contours  
Contour interval, 5 milligals. Hachures  
indicate closed areas of lower gravity  
values. Density used in Bouguer  
calculations, 2.67 gm/cc. Gravity values  
tied to National Gravity Network






Figure 22




Isogal-Bouguer gravity contours  
 Contour interval, 5 milligals. Hachures  
 indicate closed areas of lower gravity  
 values. Density used in Bouguer  
 calculations, 2.67 gm/cc. Gravity values  
 tied to National Gravity Network




### BOUGUER GRAVITY OF NUCKOLLS COUNTY

Figure 23





F6  
 Isogal-Bouguer gravity contours  
 Contour interval, 5 milligals. Hachures  
 indicate closed areas of lower gravity  
 values. Density used in Bouguer  
 calculations, 2.67 gm/cc. Gravity values  
 tied to National Gravity Network



### BOUGUER GRAVITY OF POLK COUNTY

Figure 24





### BOUGUER GRAVITY OF YORK COUNTY

Figure 26




Figure 27

were made for latitude. The final magnetic intensity data were contoured by hand (fig. 28) and show very clearly the southern edge of the midcontinent anomaly.




Figure 28

References

Muehlberger, W. R., Denison, R. E., and Lidiak, E. G. 1964.  
Buried basements rocks of the United States of America  
and Canada: Final Report, Appendix to vol. II, contract  
AF49(638)-1115/ARPA Order No. 180-62. Univ. Texas, Austin.

## SOME PRECAMBRIAN GABBROIC ROCKS FROM SOUTHEAST NEBRASKA

S. B. Treves

### Introduction

Our papers on the Elk Creek carbonatite of southeast Nebraska (1972, 1972b, 1975) produced a flurry of leasing, geophysical surveys and, finally, drilling of the Elk Creek anomaly and some smaller, nearby anomalies. Recently two cores from boreholes that were drilled to basement at the site of the nearby Steinhauer anomaly were released to the Conservation and Survey Division for reference purposes and study. These cores are described in the sections that follow.


### Location

The cores are designated Steinhauer 1 and 2, and are referred to as St-1 and St-2 in the rest of the text. St-1 was drilled (fig. 29) at the center of the north boundary line of the NW 1/4 NE 1/4 NW 1/4 Sec. 19 T.3N. R.11E. St-2 was drilled at the center of the south boundary line of the SW 1/4 SE 1/4 SW 1/4 Sec. 14 T.3N. R.12E.

### Core Description

Analysis of the core of St-1 shows that basement rocks were encountered at a depth of 800.5 feet and that drilling was continued to a depth of 994 feet, producing 194 feet of core. Analysis of the core from St-2 shows that basement rocks were encountered at a depth of 971 feet. Additional drilling to a depth of 1072 feet produced 101 feet of core. The contact of the basement rocks with the overlying, younger sedimentary rocks is sharp and nonconformable. It is clearly not an intrusive contact, as fragments of the older rock occur as clasts in the basal unit of the overlying sedimentary unit.

The upper portions of both cores are deeply weathered and stained by iron oxides. The St-1 core is weathered to a depth of 67 feet; St-2 core is only



**INDEX MAP OF**  
**Southeast Nebraska and Northeast Kansas**

50 Miles

Figure 29

weathered to a depth of 5 feet. The rocks of both cores are extensively jointed and cut by veins of secondary calcite and quartz. The orientation of the joints ranges from horizontal to vertical, with most being inclined at angles that range from 30° to 60°. Slickensides are common. The rocks immediately adjacent to the veins and most of the joints are altered, sometimes extensively.

Below the weathered zone the rocks are strikingly fresh and unaltered and are black and dark grey. Hand specimens show abundant plagioclase and lesser amounts of pyroxene, olivine, and opaque minerals. The average grain size ranges from about 2 mm to about 10 mm. Specimens from both holes are identical in hand specimens.

Thin-section analysis of unaltered specimens of core indicate the rocks are gabbroic and that they consist primarily of olivine, plagioclase, pyroxene, opaque minerals, biotite and traces of apatite and alkali feldspar. The olivine is magnesium-rich and is generally about Fo<sub>80</sub>. Locally, it is altered to serpentine. The plagioclase is generally sodic labradorite. It is complexly twinned and generally unaltered. In plagioclase-rich rocks the fabric shown by the plagioclase suggests that it may be a cumulate phase. Some plagioclase shows slight alteration to sericite. The pyroxenes consist of orthopyroxene and clinopyroxene. The orthopyroxenes are inverted pigeonites and contain lamellae of augite that are parallel to the (001) of the original pigeonite. The clinopyroxene is augite, which contains exsolved blebs of pigeonite. Orthopyroxenes are much more abundant than clinopyroxenes in these rocks. The opaque minerals are primarily ilmenite, but some sulphides are present. In many sections the ilmenite and the mafic minerals are mantled with overgrowths of biotite. Apatite occurs as prismatic, interstitial crystals or as inclusions in the plagioclase. Alkali feldspar occurs in the interstices between plagioclase grains.

Modal analyses of the gabbroic rocks are presented in table 2. The analyses show that the upper reaches of St-1 consist primarily of anorthositic gabbro,

Elevation	Olivine	Pyroxene	Plagioclase*	Opaque Minerals	Biotite	Apatite	Alkali Feldspar
St-1	,						
847	9	6	75(57)	5	3	Tr	Tr
839.8	5	6	74(65)	11	2	Tr	Tr
842	9	7	75(52)	7	2	Tr	Tr
944.5	7	23	57(56)	11	2	Tr	Tr
989.5	11	28	52(57)	7	1	Tr	Tr
994.5 <sup>+</sup>	4	21	66(55)	7	1	Tr	Tr
St-2							
974	36	13	32(58)	18	Tr	Tr	Tr
976	39	7	35(50)	16	2	Tr	Tr
978	33	10	40(35)	16	1	Tr	Tr
1000	37	5	35(58)	22	3	Tr	Tr
1032.5	38	5	33(58)	20	3	Tr	Tr
1037	38	5	39(60)	16	2	Tr	Tr
1045	45	4	25(55)	17	2	Tr	Tr
1051	18	5	39(55)	17	2	Tr	Tr
1060	43	4	34(62)	17	1	Tr	Tr
1067.5 <sup>+</sup>	31	20	29(62)	18	2	Tr	Tr
1070.5	30	13	31(50)	22	3	Tr	Tr

<sup>†</sup>Chemical analysis

\*Figure in brackets is anorthite content

Table 2. Modal Analyses of Specimens from St-1 and St-2.

whereas the lower reaches consist of olivine gabbro. The upper and lower reaches of St - 2, in contrast, consist of ilmenite-rich olivine gabbros, whereas the middle reaches consist of troctolitic, ilmenite-rich gabbros. The cores seem to exhibit a coarse lithologic layering. It does not seem possible to correlate units between the holes.

Near the bottom of both holes the core shows several thin, horizontal layers of grey carbonate. Thickness ranges from a few centimeters to about 10 centimeters. These units consist primarily of calcite, phlogopite, apatite(?) and opaque minerals and may be carbonatites. Further work will be necessary before a final determination is possible.

Two chemical analyses of fresh specimens of the core were made. The results are presented in table 3. It is readily apparent that both rocks are abnormally rich in ilmenite. If allowances are made for this feature, the rocks are not too much unlike Nockolds' average gabbro. Analysis three is of a basalt obtained from a borehole, Radenslaben No. 1, drilled on the midcontinent gravity high near Omaha, Nebraska. More chemical analyses will be required to completely characterize these rocks.

Two  $^{39}\text{Ar}/^{40}\text{Ar}$  dates (Dalrymple and Lanphere, 1971) were determined on plagioclase separates from St - 1 and St - 2 by J. Sutter. The pertinent analytical data and  $^{39}\text{Ar}/^{40}\text{Ar}$  age spectrum diagrams are presented in table 4 and figure 30. Sutter reports that both spectra are indicative of the loss of radiogenic  $^{40}\text{Ar}$  since crystallization of the gabbros. Thus, none of the ages indicated by the diagram is the time of crystallization of the gabbros. The spectra further indicate that the gabbros are older than the oldest age measured, which was 1.18 b.y. on the fuse fraction of St - 1 plagioclase. Sutter suggests that both specimens are likely to be the same age and that the age is probably not much in excess of 1.2 b.y., which makes these rocks Keweenawan in age (Goldich *et al.*, 1960).

	(1)	(2)	(3) <sup>†</sup>	(4)	(5)
SiO <sub>2</sub>	49.60	34.30	48.50	50.83	48.36
TiO <sub>2</sub>	5.37	12.50	1.48	2.03	1.32
Al <sub>2</sub> O <sub>3</sub>	15.00	8.52	15.80	14.07	16.84
Fe <sub>2</sub> O <sub>3</sub>	14.40*	35.00*	2.99	2.88	2.55
FeO	--	--	8.74	9.00	7.92
MnO	0.14	0.22	0.19	0.18	0.18
MgO	3.77	5.14	7.44	6.34	8.06
CaO	5.99	2.98	10.20	10.42	11.07
Na <sub>2</sub> O	2.36	1.04	2.37	2.23	2.26
K <sub>2</sub> O	0.86	0.57	0.34	0.82	0.56
P <sub>2</sub> O <sub>5</sub>	0.05	0.05	0.10	0.23	0.24
H <sub>2</sub> O+	--	--	1.52	0.91	0.64

\*total iron

†composite sample

1. St-1; 994.5 ft.
2. St-2; 1067.5 ft.
3. Radenslaben No. 1, C SW 1/4 SE 1/4 NW 1/4 Sec. 28 T13N R8E
4. Average tholeiitic basalt (Nockolds, 1954)
5. Average gabbro (Nockolds, 1954)

Table 3. Chemical Compositions.

## St - 1 Plagioclase 944.5 ft.

Temp. °C.	$^{39}\text{Ar}$ % of Total	Radio- genic $^{40}\text{A}$ %	Age in m.y.*
500	1.69	8.65	> 1,000 m.y.
800	11.79	39.11	1053.883 ± 21.336
1000	15.59	40.50	978.892 ± 19.464
1050	32.05	67.75	1053.472 ± 11.847
1100	15.45	48.21	1134.997 ± 18.098
Fuse	23.41	62.02	1178.587 ± 13.789
Total Gas	100.00	49.85	1082.873

## St - 2 Plagioclase

400	0.82	55.97	575.340 ± 52.792
500	1.97	80.07	818.143 ± 17.583
600	5.78	92.00	964.203 ± 10.950
700	5.22	93.26	972.995 ± 10.738
800	8.43	95.62	979.916 ± 9.901
900	7.44	94.88	965.828 ± 9.906
1050	25.85	98.03	929.321 ± 8.847
Fuse	44.50	98.31	1115.876 ± 10.087
Total Gas	100.00	96.55	1021.017

\*Post-1977 constant

Table 4. Age Spectrum Analysis Analytical Data.





Figure 30.  $^{40}\text{Ar}/^{39}\text{Ar}$  age spectrum diagram for St-1 and St-2.

### Summary and Conclusions

The gabbroic rocks associated with the Steinhauer anomaly are intrusive into older granitic rocks. They constitute a plug or plugs that occur on the west flank of the Nemaha uplift to the south and northeast of the midcontinent gravity high. The coarse-grained Steinhauer gabbros do not resemble the ophitic basalts of the midcontinent gravity high in appearance or mineralogy. They also appear to be older than the basalts in Nebraska (Goldich *et al.*, 1966). They do, however, resemble the basalts in terms of their chemistry and, hence, may be products of the same igneous event. Further, the Steinhauer gabbros resemble the Keweenawan rocks of the Duluth gabbro as described by Weiblen and Morey (1980). In light of the above, it seems that the Steinhauer gabbros are probably an early intrusive phase of the igneous activity, in Nebraska, that gave rise to the Keweenawan basaltic rocks of the midcontinent gravity high and the Keweenawan basalts and gabbros of the Lake Superior region.

### References

- Brookins, D. G., Treves, S. B., and Bolivar, S. L. 1975. Elk Creek, Nebraska, carbonatite: strontium geochemistry. *Earth Planetary Science Letters*, v. 28, p. 79-82.
- Dalrymple, G. B. and Lanphere, M. A. 1971.  $^{40}\text{Ar}/^{39}\text{Ar}$  technique of K-Ar dating: a comparison with conventional technique. *Earth and Planetary Science Letters*, v. 12, pp. 300-308.
- Goldich, S. S., Lidiak, E. G., Hedge, C. E., and Walthall, F. G. 1966. Geochronology of the Midcontinent Region, United States. *Journal of Geophysical Research*, v. 71, pp. 5379-5407.
- Nockolds, S. R. 1954. Average chemical composition of some igneous rocks. *Geological Society of America Bulletin*, v. 65, pp. 1007-1032.
- Treves, S. B., Smith, R., Carlson, M. P., and Coleman, G. 1972. Elk Creek carbonatite, Johnson and Pawnee counties, Nebraska. *Geological Society of America Abstracts with Programs*, v. 4, p. 297.
- Treves, S. B., Smith, R., Rinehart, J., Coleman, G., and Carlson, M. P. 1972. Petrography and mineralogy of the Elk Creek carbonatite, Nebraska. *Geological Society of America Abstracts with Programs*, v. 4, p. 352.
- Weiblen, P. W. and Morey, G. B. 1980. A summary of the stratigraphy, petrology, and structure of the Duluth complex. *American Journal of Science*, v. 280-A, pp. 88-133.

## APPENDIX A

This appendix consists of descriptions of the twenty-two test holes drilled in central Otoe County. Locations of these test holes are shown in figure 3 of the text.

All measurements in this appendix are expressed in feet. To convert to the International System of Units, multiply feet by .3048 to obtain meters.

Test Hole 1-81

Location: Otoe County, SW SW NW sec. 19, T. 8 N., R. 10 E., approximately 1950 feet south of north section line and 20 feet east of west section line.

Ground-level elevation: 1182.5 feet above mean sea level.

Started: May 7, 1981. Completed: May 7, 1981.

<u>Description</u>	<u>Depth, in feet</u>	
	<u>From</u>	<u>To</u>
Quaternary System:		
Soil (no sample).....	0.0	- 5.0
Clay, light yellowish brown; moderately silty.....	5.0	- 10.0
Clay, light yellowish brown; contains gravel.....	10.0	- 23.0
Clay, dark gray, slightly silty.....	23.0	- 37.5
Clay, light greenish gray.....	37.5	- 48.0
Clay, olive, silty.....	48.0	- 52.5
Clay, medium to dark gray, slightly silty....	52.5	- 66.0
Clay, medium to dark gray, slightly silty; contains black carbonaceous material.....	66.0	- 69.0
Clay, medium gray, slightly silty.....	69.0	- 83.0
Clay, light gray, slightly silty.....	83.0	- 86.0
Clay, medium gray, slightly silty.....	86.0	- 88.0
Clay, light gray, slightly silty.....	88.0	- 92.0
Clay, light gray, sandy, silty.....	92.0	- 96.0

Pennsylvanian System - Virgil Series - Wabaunsee Group:

Willard Formation:

Shale, light greenish gray.....	96.0	- 97.2
Limestone, medium gray, very finely crystalline; contains fusulinids.....	97.2	- 97.8
Shale, pale olive.....	97.8	- 99.0
Shale, light reddish gray.....	99.0	- 100.1

Emporia Formation:

Elmont Member:

Limestone, medium gray to tan, finely crystalline, pseudo-oolitic; contains bryozoans, crinoids and <u>Osagia</u> .....	100.1	- 102.8
Shale, light reddish gray.....	102.8	- 103.7
Limestone, very light gray, finely crystalline; contains <u>Osagia</u> .....	103.7	- 105.4

	<u>Description</u>	<u>Depth, in feet</u>	
		<u>From</u>	<u>To</u>
Harveyville Member:			
Shale, light to medium gray.....	105.4	-	106.9
Reading Member:			
Limestone, light gray, finely crystalline; contains crinoids, brachiopods, and Osagia.....	106.9	-	110.0
Shale, light to medium gray.....	110.0	-	110.2
Limestone, medium gray, finely crystalline, pseudo-oolitic in part; contains brachiopods and Osagia.....	110.2	-	113.0
Auburn Formation:			
Shale, light greenish gray.....	113.0	-	114.0
Shale, red.....	114.0	-	123.2
Shale, red mottled tan.....	123.2	-	126.0
Shale, light greenish gray mottled olive.....	126.0	-	130.0
Shale, medium to dark gray.....	130.0	-	138.2
Wakarusa Formation:			
Limestone, medium to dark gray, finely crys- talline; contains fusulinids; interbedded with shale, medium gray.....	138.2	-	140.6
Limestone, light gray, finely crystalline....	140.6	-	142.7
Soldier Creek Formation:			
Shale, medium gray.....	142.7	-	145.2
Limestone, medium gray, finely crystalline; contains fusulinids, crinoids, Osagia, and "black inclusions"; interbedded with shale, medium gray.....	145.2	-	145.8
Shale, reddish gray.....	145.8	-	147.2
Limestone, light gray, very finely crys- talline; contains crinoids.....	147.2	-	150.4
Shale, dark gray.....	150.4	-	151.5
Limestone, light gray to tan, finely crys- talline; contains Osagia.....	151.5	-	152.1
Shale, medium gray.....	152.1	-	155.0
Shale, light gray.....	155.0	-	156.2
Shale, medium gray.....	156.2	-	160.1
Burlingame Formation:			
Limestone, medium gray, finely crystalline... 160.1	-	161.0	
Limestone, medium gray, variably crystal- line; contains fusulinids, crinoids, brachiopods, pyrite, and "black inclusions".....	161.0	-	161.6
Shale, light greenish gray.....	161.6	-	162.0
Shale, light to medium gray.....	162.0	-	163.5
Limestone, light gray, very finely crys- talline; contains brachiopods and glauconite.....	163.5	-	164.8
Scranton Formation:			
Silver Lake - White Cloud Members:			
Shale, pale olive.....	164.8	-	167.0

<u>Description</u>	<u>Depth, in feet</u>	
	<u>From</u>	<u>To</u>
Shale, red mottled green.....	167.0	- 167.2
Shale, light greenish gray.....	167.2	- 170.0
Shale, reddish gray.....	170.0	- 178.3
Shale, reddish gray; interbedded with lime-stone, light greenish gray.....	178.3	- 182.9
Shale, light greenish gray.....	182.9	- 183.0
Limestone, light greenish gray, finely crystalline.....	183.0	- 183.9
Shale, pale olive.....	183.9	- 184.5
Shale, light gray.....	184.5	- 194.0
Coal, black.....	194.0	- 194.5
Shale, light greenish gray.....	194.5	- 213.3
Shale, light greenish gray; interbedded with thin limestone streaks.....	213.3	- 213.5
Shale, medium gray.....	213.5	- 240.3
Shale, light gray, silty.....	240.3	- 242.0
Howard Formation:		
Limestone, medium to dark gray, finely crystalline; contains abundant crinoids...	242.0	- 243.3
Shale, dark gray.....	243.3	- 245.2
Limestone, light to medium gray, very finely crystalline; contains crinoids, fusulinids, pyrite, and "black inclusions".....	245.2	- 250.8
Severy Formation:		
Shale, medium gray.....	250.8	- 251.0
Coal, black.....	251.0	- 251.4
Shale, medium gray.....	251.4	- 257.0

#### Test Hole 2-81

Location: Otoe County, NE corner SE sec. 17, T. 3 N., R. 10 E., approximately 2578 feet north of south section line and 22 feet west of east section line.

Ground-level elevation: 1152.0 feet above mean sea level.

Started: May 7, 1981. Completed: May 7, 1981.

Total depth: 227.0 feet.

<u>Description</u>	<u>Depth, in feet</u>	
	<u>From</u>	<u>To</u>
Quaternary System: Soil (no sample).....	0.0	- 3.0

<u>Description</u>	<u>Depth, in feet</u>	
	<u>From</u>	<u>To</u>
Clay, yellowish brown; contains gravel.....	3.0	- 7.0
Clay, yellowish brown; contains sand and gravel.....	7.0	- 10.0
Clay, light brownish gray, slightly silty....	10.0	- 17.0
Sand, coarse; contains gravel.....	17.0	- 19.0
Clay, tan, slightly silty.....	19.0	- 23.5
Clay, light brown.....	23.5	- 24.0
Clay, light brown to light gray.....	24.0	- 31.0
Clay, medium gray to tan, silty.....	31.0	- 42.0
Clay, medium gray, silty.....	42.0	- 47.0
Clay, light to medium gray, silty.....	47.0	- 54.5
Clay, light greenish gray, silty.....	54.5	- 59.5
Clay, medium greenish gray, silty.....	59.5	- 64.3
Clay, medium to dark gray, silty.....	64.3	- 68.0
Clay, medium gray to greenish gray.....	68.0	- 79.5
Sand, fine to medium; contains gravel.....	79.5	- 89.0
Pennsylvanian System - Virgil Series - Wabaunsee Group:		
Auburn Formation:		
Shale, pale olive.....	89.0	- 93.0
Shale, light greenish gray.....	93.0	- 94.0
Shale, light to medium gray.....	94.0	- 103.5
Wakarusa Formation:		
Limestone, medium gray, finely crystalline; contains crinoids, fusulinids, brachio- pods, and <u>Osagia</u> .....	103.5	- 107.4
Soldier Creek Formation:		
Shale, medium gray.....	107.4	- 108.5
Limestone, medium gray to tan, finely crystalline.....	108.5	- 110.2
Limestone, light gray, finely crystalline, pseudo-oolitic; contains <u>Osagia</u> and glauconite.....	110.2	- 114.2
Shale, light gray.....	114.2	- 117.0
Shale, medium gray.....	117.0	- 120.1
Shale, light gray.....	120.1	- 126.2
Burlingame Formation:		
Limestone, medium gray to tan, finely crys- talline; contains fusulinids and <u>Osagia</u> ...	126.2	- 127.5
Shale, light gray.....	127.5	- 128.6
Limestone, tan, finely crystalline; contains fusulinids, <u>Osagia</u> , glauconite, and stylolites.....	128.6	- 131.0
Scranton Formation:		
Silver Lake - White Cloud Members:		
Shale, light greenish gray mottled olive....	131.0	- 132.8
Shale, light gray.....	132.8	- 150.5
Shale, light gray; contains brown sandstone streaks and pyrite.....	150.5	- 151.0

	<u>Description</u>	<u>Depth, in feet</u>	
		<u>From</u>	<u>To</u>
	Shale, light gray; contains brown sandstone streaks and black carbonaceous material....	151.0	- 163.4
	Shale, light gray; contains brown sandstone streaks.....	163.4	- 179.3
	Sandstone, medium to dark gray.....	179.?	- 180.1
	Shale, light gray, silty.....	180.1	- 192.0
	Shale, medium gray mottled tan; contains brown sandstone streaks.....	192.0	- 198.4
	Shale, light gray; contains brown sandstone streaks and trace of coal at 209.0.....	198.4	- 212.0
	Limestone, medium gray, very finely crystalline; contains brachiopods, crinoids, and pyrite.....	212.0	- 212.2
	Shale, light gray; contains brown sandstone streaks.....	212.2	- 214.3
Howard Formation:			
	Limestone, medium gray, very finely crystalline; contains crinoids and abundant fusulinids.....	214.3	- 216.2
	Limestone, light gray, very finely crystalline; contains crinoids, fusulinids, <u>Osagia</u> , and pyrite.....	216.2	- 222.4
Severy Formation:			
	Shale, black, carbonaceous.....	222.4	- 222.7
	Shale, light gray.....	222.7	- 227.0

#### Test Hole 3-81

Location: Otoe County, NW corner sec. 21, T. 8 N. R. 10 E., approximately 138 feet south of north section line and 18 feet east of west section line.

Ground-level elevation: 1170.0 feet above mean sea level.

Started: May 3, 1981. Completed: May 8, 1981.

Total Depth: 182.0 feet.

	<u>Description</u>	<u>Depth, in feet</u>	
		<u>From</u>	<u>To</u>
Quaternary System:			
	Soil (no sample).....	0.0	- 6.0
	Clay, brown.....	6.0	- 18.0
	Clay, light brown to tan, sandy.....	18.0	- 22.0
	Clay, gray to tan, sandy.....	22.0	- 25.0

<u>Description</u>	<u>Depth, in feet</u>	
	<u>From</u>	<u>To</u>
Clay, tan, very sandy.....	25.0	- 32.0
Clay, brown, very sandy.....	32.0	- 44.0
Clay, light brown, sandy.....	44.0	- 51.0
Sand, fine.....	51.0	- 54.5
Clay, light brown, very sandy.....	54.5	- 59.0
Clay, light gray, silty.....	59.0	- 62.5
Clay, medium gray, very sandy.....	62.5	- 71.0
Clay, dark gray.....	71.0	- 76.5
Clay, light gray, silty, sandy.....	76.5	- 88.0
Clay, greenish gray, sandy.....	88.0	- 95.5
Pennsylvanian System - Virgil Series - Wabaunsee Group:		
Auburn Formation:		
Shale, red.....	95.5	- 98.5
Shale, olive, mottled yellow and red.....	98.5	- 100.3
Shale, olive.....	100.3	- 103.0
Shale, medium gray.....	103.0	- 112.8
Wakarusa Formation:		
Limestone, medium gray, finely crystalline; contains crinoids, fusulinids, brachio- pods, and pyrite.....	112.8	- 115.0
Limestone, medium gray, finely crystalline; contains fusulinids, pyrite, and "black inclusions".....	115.0	- 117.2
Soldier Creek Formation:		
Shale, medium gray.....	117.2	- 118.3
Limestone, light gray to tan, very finely crystalline; interbedded with shale, light greenish gray.....	118.3	- 120.0
Shale, reddish gray.....	120.0	- 121.1
Shale, light greenish gray mottled red.....	121.1	- 121.8
Limestone, light gray, very finely crystal- line; contains <u>Osagia</u> and glauconite.....	121.8	- 123.8
Shale, light greenish gray.....	123.8	- 127.0
Shale, light gray.....	127.0	- 130.0
Shale, medium gray.....	130.0	- 135.1
Burlingame Formation:		
Limestone, medium gray, finely crystalline...	135.1	- 135.6
Shale, light gray.....	135.6	- 136.8
Limestone, light gray, very finely crystal- line; contains glauconite.....	136.8	- 137.5
Limestone, light greenish gray, finely crys- talline; contains abundant glauconite.....	137.5	- 139.3
Scranton Formation:		
Silver Lake - White Cloud Members:		
Sandstone, tan; interbedded with shale, greenish gray.....	139.3	- 141.8
Shale, black, carbonaceous.....	141.8	- 142.2
Shale, light greenish gray.....	142.2	- 145.0
Shale, red.....	145.0	- 149.5

	<u>Description</u>	<u>Depth, in feet</u>	
		<u>From</u>	<u>To</u>
	Shale, green mottled red.....	149.5	- 151.0
	Shale, red mottled green.....	151.0	- 155.3
	Shale, green mottled red.....	155.3	- 155.9
	Shale, red mottled green.....	155.9	- 158.5
	Shale, olive mottled red.....	158.5	- 159.5
	Shale, medium gray.....	159.5	- 167.2
	Coal, black.....	167.2	- 167.5
	Shale, medium gray.....	167.5	- 182.0

Test Hole 4-81

Location: Otoe County, SE corner NE NE SE sec. 5, T. 8 N., R. 10 E., approximately 2012 feet north of south section line and 30 feet west of east section line.

Ground-level Elevation: 1149.0 feet above mean sea level.

Started: May 8, 1981. Completed: May 8, 1981.

Total Depth: 137.0 feet.

	<u>Description</u>	<u>Depth, in feet</u>	
		<u>From</u>	<u>To</u>
Quaternary System:			
	Soil (no sample).....	0.0	- 3.0
	Clay, brown, silty.....	3.0	- 6.5
	Clay, light tan, very silty.....	6.5	- 20.0
	Clay, light tan, sandy.....	20.0	- 38.5
	Clay, tan, silty.....	38.5	- 43.0
	Clay, gray, sandy, silty.....	43.0	- 47.0
	Clay, medium gray, sandy.....	47.0	- 51.0
	Clay, light gray, sandy.....	51.0	- 62.0
	Clay, medium gray, sandy.....	62.0	- 69.0
	Clay, medium gray; contains gravel.....	69.0	- 74.5

Pennsylvanian Series - Virgil Series - Wabaunsee Group:

Auburn Formation:

Shale, olive..... 74.5 - 76.8

Wakarusa Formation:

Limestone, medium gray, finely crystalline;  
interbedded with shale, medium gray..... 76.8 - 79.6

Shale, black; contains carbonaceous  
material..... 79.6 - 80.0

<u>Description</u>	<u>Depth, in feet</u>	
	<u>From</u>	<u>To</u>
Limestone, medium gray, finely to very finely crystalline; contains brachiopods and pyrite.....	80.0	-
Soldier Creek Formation:	80.4	-
Shale, medium gray.....	80.4	-
Shale, reddish gray.....	82.0	-
Limestone, light gray, very finely crystalline.....	84.2	-
Shale, gray to tan.....	86.1	-
Shale, medium gray.....	87.8	-
Shale, dark gray.....	92.0	-
Burlingame Formation:	98.3	-
Limestone, medium gray, finely crystalline; contains gastropods and pyrite.....	98.3	-
Limestone, medium gray, variably crystal- line; contains <u>Osagia</u> .....	100.4	-
Scranton Formation:	100.4	-
Silver Lake - White Cloud Members:	101.4	-
Shale, medium greenish gray.....	101.4	-
Shale, medium olive gray.....	102.6	-
Shale, grayish red.....	104.3	-
Shale, light green.....	105.0	-
Shale, dark red.....	108.0	-
Shale, red.....	111.5	-
Shale, red; interbedded with limestone, gray.....	115.0	-
Shale, dark red.....	115.5	-
Shale, olive.....	118.0	-
Shale, medium gray.....	118.8	-
Shale, light gray.....	126.7	-
Coal, black.....	128.0	-
Shale, light gray.....	128.4	-
Shale, medium gray.....	130.0	-
	137.0	

Test Hole 5-81

Location: Otoe County, SE corner sec. 4, T. 8 N., R. 10 E.,  
approximately 94 feet north of south section line and  
15 feet west of east section line.

Ground-level elevation: 1159.5 feet above mean sea level.

Started: May 11, 1981. Completed: May 11, 1981.

Total Depth: 136.0 feet.

<u>Description</u>	<u>Depth, in feet</u>	
	<u>From</u>	<u>To</u>
Quaternary System:		
Soil (no sample).....	0.0	- 3.0
Clay, brown, silty, sandy.....	3.0	- 6.5
Clay, light brown, silty, sandy.....	6.5	- 11.0
Clay, medium gray to brown, silty, sandy.....	11.0	- 14.5
Clay, brown, sandy.....	14.5	- 17.0
Clay, brown, very sandy.....	17.0	- 24.5
Clay, medium gray to brown, sandy.....	24.5	- 38.0
Clay, medium gray, slightly sandy.....	38.0	- 43.4
Pennsylvanian System - Virgil Series - Wabaunsee Group:		
Emporia Formation:		
Reading Member:		
Limestone, light gray, finely crystalline....	43.4	- 44.9
Shale, medium gray.....	44.9	- 46.2
Limestone, tan, weathered, finely crystalline; contains fusulinids, crinoids, <u>Osagia</u> and manganese staining.....	46.2	- 48.1
Auburn Formation:		
Shale, olive.....	48.1	- 50.1
Shale, red.....	50.1	- 58.0
Shale, red mottled green.....	58.0	- 59.0
Shale, red.....	59.0	- 63.7
Shale, red; interbedded with limestone, medium gray.....	63.7	- 66.5
Shale, green mottled red.....	66.5	- 70.8
Shale, medium gray.....	70.8	- 79.3
Wakarusa Formation:		
Limestone, medium gray, finely crystalline; interbedded with shale, medium gray.....	79.3	- 80.3
Limestone, medium gray, finely crystalline; contains fusulinids and brachiopods.....	80.3	- 83.2
Limestone, light to medium gray, finely crystalline; contains fusulinids.....	83.2	- 84.0
Soldier Creek Formation:		
Shale, medium gray mottled red.....	84.0	- 85.3
Limestone, medium gray, very finely crystalline.....	85.3	- 86.2
Shale, medium reddish gray.....	86.2	- 87.2
Shale, medium gray.....	87.2	- 88.6
Limestone, light gray, finely crystalline; contains <u>Osagia</u> .....	88.6	- 89.6
Limestone, light gray, very finely crystalline; contains glauconite.....	89.6	- 91.8
Shale, medium gray.....	91.8	- 101.2
Burlingame Formation:		
Limestone, medium gray, finely crystalline; contains "black inclusions".....	101.2	- 103.1
Shale, medium greenish gray.....	103.1	- 104.2
Limestone, light gray, very finely crystalline; interbedded with shale, red.....	104.2	- 106.5

	<u>Description</u>	<u>Depth, in feet</u>		
		<u>From</u>	<u>To</u>	
<b>Scranton Formation:</b>				
Silver Lake - White Cloud Members:				
Shale, red.....	106.5	-	107.0	
Shale, red mottled gray.....	107.0	-	109.2	
Shale, green.....	109.2	-	110.2	
Shale, red.....	110.2	-	117.2	
Shale, red; interbedded with limestone, medium gray.....	117.2	-	120.1	
Shale, light gray.....	120.1	-	122.?	
Shale, medium gray.....	122.3	-	123.1	
Limestone, medium gray, very finely crys- talline; contains <u>Osagia</u> ; interbedded with shale, dark gray.....	123.1	-	123.3	
Shale, light gray.....	123.3	-	129.8	
Coal, black.....	129.8	-	130.2	
Shale, medium gray.....	130.2	-	136.0	

Test Hole 6-81

Location: Otoe County, SW corner sec. 1, T. 8 N., R. 10 E.,  
approximately 207 feet north of south section line  
and 103 feet east of west section line.

Ground-level elevation: 1133.5 feet above mean sea level.

Started: May 11, 1981. Completed: May 11, 1981.

Total Depth: 205.0 feet.

	<u>Description</u>	<u>Depth, in feet</u>		
		<u>From</u>	<u>To</u>	
<b>Quaternary System:</b>				
Soil (no sample).....				
Clay, gray to tan, sandy.....	0.0	-	4.5	
Clay, gray, silty, sandy.....	4.5	-	10.0	
Clay, gray to brown, silty, sandy.....	10.0	-	19.0	
Clay, gray, sandy.....	19.0	-	25.0	
Clay, brown to gray, sandy.....	25.0	-	32.0	
Clay, gray, sandy.....	32.0	-	38.5	
Clay, gray, very silty, sandy.....	38.5	-	47.0	
Clay, dark gray, very silty, sandy.....	47.0	-	105.5	
	105.5	-	108.9	

<u>Description</u>	<u>Depth, in feet</u>	
	<u>From</u>	<u>To</u>
Pennsylvanian System - Virgil Series - Wabaunsee Group:		
Burlingame Formation:		
Shale, gray to dark gray.....	108.9	- 109.3
Shale, olive.....	109.3	- 110.2
Shale, olive, interbedded with limestone, gray.....	110.2	- 111.4
Shale, medium gray.....	111.4	- 112.1
Limestone, gray to tan, variably crystal- line; contains glauconite; interbedded with shale, olive.....	112.1	- 113.0
Scranton Formation:		
Silver Lake - White Cloud Members:		
Shale, light green.....	113.0	- 114.8
Shale, black.....	114.8	- 115.4
Shale, medium greenish gray.....	115.4	- 123.5
Shale, red mottled green.....	123.5	- 134.2
Shale, light gray.....	134.2	- 139.5
Coal, black.....	139.5	- 139.9
Shale, light gray.....	139.9	- 167.0
Shale, medium gray.....	167.0	- 188.3
Howard Formation:		
Limestone, light gray, finely crystalline; contains brachiopods; interbedded with shale, medium gray.....	188.3	- 190.4
Limestone, medium gray, very finely crystal- line; interbedded with shale, medium gray.....	190.4	- 190.7
Shale, medium gray.....	190.7	- 193.4
Limestone medium gray, very finely crystal- line; contains fusulinids, <u>Osagia</u> , and "black inclusions".....	193.4	- 194.7
Limestone, light gray to tan, very finely crystalline; contains fusulinids, <u>Osagia</u> , and coarsely crystalline calcite streaks..	194.7	- 198.6
Severy Formation:		
Coal, black.....	198.6	- 199.4
Siltstone, light gray.....	199.4	- 205.0

#### Test Hole 7-81

Location: Otoe County, NE corner SE NE sec. 14, T. 8 N.,  
R. 10 E., approximately 1920 feet south of north  
section line and 19 feet west of east section line.

Ground-level elevation: 1124.5 feet above mean sea level.

Started: May 11, 1981. Completed: May 11, 1981.

Total Depth: 130.0 feet.

	<u>Description</u>	<u>Depth, in feet</u>	
		<u>From</u>	<u>To</u>
Quaternary System:			
Soil (no sample).....		0.0	- 6.0
Clay, medium gray, very sandy.....		6.0	- 10.0
Clay, tan to medium gray, sandy.....		10.0	- 15.0
Clay, tan, sandy.....		15.0	- 17.0
Pennsylvanian System - Virgil Series - Wabaunsee Group:			
Willard Formation:			
Shale, medium gray; contains sandstone streaks.....		17.0	- 23.0
Shale, light gray; contains sandstone streaks.....		23.0	- 32.0
Shale, medium gray.....		32.0	- 44.5
Shale, medium gray, sandy.....		44.5	- 48.5
Shale, medium greenish gray.....		48.5	- 51.0
Shale, green mottled red.....		51.0	- 52.0
Shale, red and green.....		52.0	- 53.4
Emporia Formation:			
Elmont - Reading Members:			
Limestone, medium gray, finely crystalline, pseudo-oolitic; contains glauconite and pyrite.....		53.4	- 54.2
Shale, red.....		54.2	- 56.4
Shale, medium gray; interbedded with limestone, medium gray.....		56.4	- 56.8
Limestone, light gray, very finely crystalline; contains <u>Osagia</u> and coarsely crystalline calcite streaks.....		56.8	- 58.5
Shale, medium gray.....		58.5	- 59.8
Limestone, light gray to tan, very finely crystalline; contains brachiopods, fusulinids, <u>Osagia</u> , and glauconite.....		59.8	- 62.2
Auburn Formation:			
Shale, light green.....		62.2	- 63.5
Shale, red.....		63.5	- 71.2
Shale, red mottled green.....		71.2	- 71.6
Shale, red.....		71.6	- 77.5
Shale, red, very sandy.....		77.5	- 81.0
Shale, olive.....		81.0	- 83.6
Shale, grayish green.....		83.6	- 85.2
Shale, medium gray.....		85.2	- 93.3
Wakarusa Formation:			
Limestone, medium gray, finely crystalline; contains brachiopods, fusulinids, and pyrite.....		93.3	- 93.9

<u>Description</u>	<u>Depth, in feet</u>	
	<u>From</u>	<u>To</u>
Limestone, medium gray, finely crystalline; contains brachiopods, <u>Osagia</u> , "black inclusions," and abundant fusulinids; interbedded with shale, green.....	93.9	- 98.2
Soldier Creek Formation:		
Shale, medium gray.....	98.2	- 100.7
Shale, light grayish green.....	100.7	- 103.2
Limestone, light gray to tan, finely crystalline; contains <u>Osagia</u> .....	103.2	- 105.1
Shale, medium gray.....	105.1	- 107.8
Shale, medium olive gray.....	107.8	- 110.5
Shale, medium gray.....	110.5	- 116.1
Burlingame Formation:		
Limestone, light greenish gray, finely crystalline; contains glauconite, pyrite, and "black inclusions".....	116.1	- 119.9
Scranton Formation:		
Silver Lake - White Cloud Members:		
Shale, medium greenish gray.....	119.9	- 121.2
Shale, black.....	121.2	- 122.7
Shale, grayish green.....	122.7	- 127.5
Shale, light green.....	127.5	- 130.0

#### Test Hole 8-81

Location: Otoe County, SW NW SW sec. 7, T. 8 N., R. 11 E., approximately 1700 feet north of south section line and 20 feet east of west section line.

Ground-level elevation: 1093.0 feet above mean sea level.

Started: May 13, 1981. Completed: May 13, 1981.

Total Depth: 122.0 feet.

<u>Description</u>	<u>Depth, in feet</u>	
	<u>From</u>	<u>To</u>
Quaternary System:		
Soil (no sample).....	0.0	- 4.0
Clay, light brown, sandy.....	4.0	- 11.5
Clay, light brown, very sandy.....	11.5	- 15.5
Clay, brown, sandy, silty.....	15.5	- 21.0
Clay, brown, sandy.....	21.0	- 26.0

<u>Description</u>	<u>Depth, in feet</u>	
	<u>From</u>	<u>To</u>
Pennsylvanian System - Virgil Series - Wabaunsee Group:		
Auburn Formation:		
Shale, red, very sandy.....	26.0	- 27.2
Shale, red, and siltstone, red.....	27.2	- 32.7
Shale, olive.....	32.7	- 35.0
Shale, greenish gray.....	35.0	- 36.8
Shale, medium gray.....	36.8	- 41.2
Shale, medium gray; contains abundant pyrite.....	41.2	- 47.2
Wakarusa Formation:		
Shale, medium gray; interbedded with limestone, gray.....	47.2	- 48.1
Limestone, medium gray, finely crystalline; contains brachiopods.....	48.1	- 49.4
Shale, black.....	49.4	- 49.5
Limestone, medium gray, finely crystalline; contains brachiopods.....	49.5	- 50.6
Shale, black.....	50.6	- 50.8
Limestone, medium gray, finely crystalline; contains fusulinids and brachiopods.....	50.8	- 51.9
Soldier Creek Formation:		
Shale, light gray.....	51.9	- 53.4
Limestone, light gray to tan, very finely crystalline; contains fusulinids.....	53.4	- 53.9
Shale, red.....	53.9	- 55.0
Shale, red mottled green.....	55.0	- 56.1
Limestone, light gray, very finely crystalline; contains <u>Osagia</u> .....	56.1	- 57.0
Limestone, light greenish gray, very finely crystalline; contains fusulinids, <u>Osagia</u> , and glauconite.....	57.0	- 59.6
Shale, light green.....	59.6	- 60.2
Shale, light gray.....	60.2	- 62.3
Shale, olive gray .....	62.3	- 63.0
Shale, medium gray.....	63.0	- 71.2
Shale, medium to very dark gray.....	71.2	- 71.4
Burlingame Formation:		
Limestone, medium to dark gray, finely crystalline; contains <u>Osagia</u> .....	71.4	- 71.6
Limestone, medium gray, very finely crystalline; contains "black inclusions" and pyrite.....	71.6	- 73.0
Limestone, medium gray, finely crystalline; contains fusulinids, <u>Osagia</u> , glauconite, pyrite, and "black inclusions".....	73.0	- 74.2
Limestone, light greenish gray, very finely crystalline; contains <u>Osagia</u> and glauconite.....	74.2	- 75.1

	<u>Description</u>	<u>Depth, in feet</u>		
		<u>From</u>	<u>To</u>	
<b>Scranton Formation:</b>				
<b>Silver Lake - White Cloud Members:</b>				
Shale, light grayish green.....	75.1	-	75.6	
Shale, dark gray.....	75.6	-	77.5	
Shale, medium gray.....	77.5	-	79.0	
Shale, light green.....	79.0	-	84.4	
Shale, red.....	84.4	-	87.8	
Shale, red; interbedded with limestone, gray.....	87.8	-	90.1	
Shale, red mottled green.....	90.1	-	90.3	
Shale, red.....	90.3	-	92.0	
Shale, red mottled green.....	92.0	-	93.5	
Shale, light gray.....	93.5	-	95.1	
Limestone, medium greenish gray, medium crystalline; contains abundant brachiopods.....	95.1	-	95.4	
Shale, medium gray.....	95.4	-	96.5	
Coal, black.....	96.5	-	97.0	
Shale, light greenish gray.....	97.0	-	102.4	
Shale, black.....	102.4	-	102.6	
Shale, light gray.....	102.6	-	122.0	

Test Hole 9-81

Location: Otoe County, NW NE NW sec. 19, T. 8 N., R. 11 E.,  
approximately 19 feet south of north section line  
and 1775 feet east of west section line.

Ground-level elevation: 1160.0 feet above mean sea level.

Started: May 13, 1981. Completed: May 13, 1981.

Total Depth: 120.0 feet.

	<u>Description</u>	<u>Depth, in feet</u>		
		<u>From</u>	<u>To</u>	
<b>Quaternary System:</b>				
Soil (no sample).....				
Clay, brown, sandy.....	0.0	-	3.0	
Clay, medium gray to brown, sandy.....	3.0	-	4.5	
Clay, light brown, very sandy.....	4.5	-	6.0	
Clay, olive, sandy.....	6.0	-	8.0	
Clay, tan, sandy.....	8.0	-	10.0	
	10.0	-	10.5	

	<u>Description</u>	<u>Depth, in feet</u>	
		<u>From</u>	<u>To</u>
Permian System - Big Blue Series - Council Grove Group:			
Red Eagle Formation:			
Bennett Member:			
(no sample).....		10.5	-
Glenrock Member:			
(no sample).....		15.5	-
Johnson Formation:			
(no sample).....		17.0	-
Limestone, olive to medium gray, very finely crystalline.....		20.0	-
Limestone, light gray, very finely crystalline.....		21.0	-
Shale, grayish green.....		23.0	-
Limestone, medium greenish gray, very finely crystalline.....		23.5	-
Limestone, gray to pale red, very finely crystalline.....		24.0	-
Shale, green.....		27.4	-
Shale, tan, limy.....		27.5	-
Foraker Formation:			
Long Creek Member:			
Limestone, tan, very finely crystalline.....		28.5	-
Limestone, medium gray, very finely crystalline.....		30.0	-
Limestone, tan, very finely crystalline.....		30.4	-
Limestone, medium gray, finely crystalline....		32.5	-
Hughes Creek Member:			
Shale, dark gray; interbedded with lime- stone, medium gray.....		34.0	-
Limestone, medium gray, medium to finely crystalline; contains fusulinids, brachiopods, crinoids, and <u>Osagia</u> .....		38.8	-
Shale, light gray.....		41.1	-
Shale, medium gray; interbedded with lime- stone, medium gray.....		41.8	-
Shale, medium to dark gray.....		45.0	-
Limestone, light gray to medium gray, finely crystalline; contains brachiopods, fusulnids, and pyrite.....		47.5	-
Shale, light gray.....		48.4	-
Limestone, light gray, finely crystalline; contains brachiopods and crinoids; interbedded with shale, light gray.....		50.0	-
Shale, light gray.....		51.6	-
Shale, light grayish green.....		52.0	-
Limestone, medium gray, finely crystalline...		53.9	-
Shale, very dark gray.....		55.1	-
Limestone, medium gray, finely crystalline; contains <u>Osagia</u> and "black inclusions"....		56.8	-
Shale, medium gray.....		58.4	-

	<u>Description</u>	<u>Depth, in feet</u>	
		<u>From</u>	<u>To</u>
Limestone, dark gray, finely crystalline; contains <u>Osagia</u> .....	59.0	-	59.2
Shale, medium gray.....	59.2	-	60.7
Limestone, medium gray, finely crystalline...	60.7	-	61.1
Shale, medium gray.....	61.1	-	61.6
Limestone, medium gray, medium crystalline; contains brachiopods, crinoids, fusulinids, and <u>Osagia</u> .....	61.6	-	63.3
Limestone, medium gray, finely crystalline; contains crinoids, fusulinids, brachio- pods, and "black inclusions".....	63.3	-	64.0
Limestone, light gray, finely crystalline; contains abundant crinoids; interbedded with shale, medium gray.....	64.0	-	64.5
Shale, medium to very dark gray.....	64.5	-	65.0
Limestone, medium gray, very finely crystalline.....	65.0	-	65.6
Shale, medium to dark gray.....	65.6	-	65.9
Limestone, medium gray, finely crystalline...	65.9	-	67.1
Shale, medium greenish gray.....	67.1	-	68.1
Shale, medium gray.....	68.1	-	69.0
Limestone, medium gray, finely crystalline; contains brachiopods, <u>Osagia</u> , and "black inclusions".....	69.0	-	71.4
Shale, medium gray.....	71.4	-	74.1
Shale, black.....	74.1	-	74.3
Americus Member:			
Limestone, medium gray, very finely crystal- line; contains brachiopods, bryozoans, and black carbonaceous material.....	74.3	-	75.8
Shale, dark gray.....	75.8	-	76.8
Limestone, light gray, finely crystalline; contains crinoids; interbedded with shale, dark gray.....	76.8	-	77.0
Limestone, medium gray, finely crystalline; contains brachiopods and fusulinids.....	77.0	-	77.3
Shale, medium to dark gray.....	77.3	-	77.7
Limestone, light gray, very finely crystal- line; contains brachiopods and crinoids...	77.7	-	78.1
Admire Group:			
West Branch - Hamlin Formations:			
Shale, light gray.....	78.1	-	82.3
Limestone, medium gray, very finely crystalline.....	82.3	-	82.6
Shale, medium gray.....	82.6	-	88.1
Limestone, medium gray, very finely crystal- line; contains pyrite.....	88.1	-	88.3
Shale, medium gray.....	88.3	-	88.9

<u>Description</u>	<u>Depth, in feet</u>	
	<u>From</u>	<u>To</u>
Limestone, medium gray, finely crystalline; contains brachiopods, <u>Osagia</u> , and pyrite..	88.9	- 90.1
Shale, green.....	90.1	- 92.0
Shale, reddish green.....	92.0	- 93.8
Shale, green.....	93.8	- 94.7
Limestone, cream, very finely crystalline....	94.7	- 95.8
Shale, green.....	95.8	- 96.0
Shale, red.....	96.0	- 97.0
Shale, medium gray.....	97.0	- 97.4
Limestone, light gray, very finely crystalline.....	97.4	- 98.0
Shale, grayish green.....	98.0	- 98.4
Shale, red.....	98.4	- 99.0
Shale, olive.....	99.0	- 99.5
Shale, red mottled green.....	99.5	- 100.0
Shale, olive.....	100.0	- 100.3
Limestone, very light gray, very finely crystalline; contains glauconite.....	100.3	- 102.5
Shale, red; interbedded with limestone, gray.....	102.5	- 104.0
Shale, olive.....	104.0	- 104.4
Shale, green; interbedded with limestone, gray.....	104.4	- 106.2
Shale, red.....	106.2	- 108.2
Shale, green.....	108.2	- 109.0
Shale, olive.....	109.0	- 109.8
Shale, medium gray.....	109.8	- 111.0
Shale, green.....	111.0	- 112.1
Shale, light gray.....	112.1	- 113.8
Shale, dark green.....	113.8	- 114.5
Shale, green; interbedded with limestone, gray.....	114.5	- 115.8
Shale, red mottled green.....	115.8	- 116.9
Limestone, light greenish gray, very finely crystalline; contains glauconite.....	116.9	- 120.0

Test Hole 1C-81

Location: Otoe County, NE corner SE sec. 30, T. 3 N., R. 11 E.,  
approximately 2 $\frac{1}{2}$ 63 feet north of south section line  
and 20 feet west of east section line.

Ground-level elevation: 1136.5 feet above mean sea level.

Started: May 14, 1981. Completed: May 14, 1981.

Total Depth: 130.0 feet.

<u>Description</u>	<u>Depth, in feet</u>	
	<u>From</u>	<u>To</u>
Quaternary System:		
Soil (no sample).....	0.0	- 3.0
Clay, tan, sandy.....	3.0	- 3.2
Sand, medium.....	8.2	- 8.5
Clay, tan, sandy.....	8.5	- 21.0
Clay, brown, sandy.....	21.0	- 26.0
Clay, tan.....	26.0	- 28.0
Clay, brown.....	28.0	- 32.0
Permian System - Big Blue Series - Council Grove Group:		
Foraker Formation:		
Hughes Creek Member:		
Shale, olive; interbedded with limestone, gray.....	32.0	- 34.3
Shale, olive gray; interbedded with lime- stone, gray.....	34.3	- 37.5
Limestone, medium gray, finely crystalline...	37.5	- 37.9
Shale, olive.....	37.9	- 38.6
Limestone, tan, finely crystalline; contains brachiopods and "black inclusions".....	38.6	- 39.0
Limestone, medium gray to tan, finely crys- talline; contains brachiopods and "black inclusions".....	39.0	- 40.1
Shale, olive; interbedded with limestone, gray.....	40.1	- 41.0
Shale, dark gray.....	41.0	- 42.4
Limestone, medium gray, finely crystalline, pseudo-oolitic; contains brachiopods.....	42.4	- 43.0
Shale, medium gray.....	43.0	- 43.1
Limestone, medium gray, finely crystalline; contains <u>Osagia</u> .....	43.1	- 44.8
Shale, medium gray.....	44.8	- 47.8
Shale, black.....	47.8	- 48.2
Americus Member:		
Limestone, medium to dark gray, very finely crystalline; contains fusulinids and brachiopods.....	48.2	- 49.2
Shale, dark gray.....	49.2	- 50.1
Limestone, medium gray, very finely crystal- line; contains fusulinids and brachio- pods.....	50.1	- 50.2
Shale, dark gray.....	50.2	- 50.7
Limestone, medium gray, very finely crystalline.....	50.7	- 51.3
Shale, dark gray.....	51.3	- 51.5
Limestone, medium gray, finely crystalline; contains brachiopods.....	51.5	- 52.0

	<u>Description</u>	<u>Depth, in feet</u>		
		<u>From</u>	<u>To</u>	
Admire Group:				
Hamlin Formation:				
Oaks Member:				
	Shale, medium gray.....	52.0	-	
	Limestone, medium gray, very finely crystalline; contains crinoids.....	57.1	-	
	Shale, medium gray.....	57.8	-	
	Limestone, medium gray, very finely crystalline; contains fusulinids.....	58.5	-	
	Shale, light green.....	59.6	-	
	Limestone, very light gray, lithographic; interbedded with shale, green.....	62.0	-	
	Shale, green to gray.....	63.5	-	
	Shale, red mottled green.....	65.0	-	
	Shale, green.....	66.8	-	
	Shale, dark gray.....	67.5	-	
Houchens Creek Member:				
	Limestone, light gray, finely crystalline....	69.3	-	
Stine Member:				
	Shale, olive gray.....	70.1	-	
	Shale, green.....	72.0	-	
	Limestone, medium gray, finely crystalline...	72.7	-	
	Shale, red mottled green.....	73.8	-	
	Shale, light gray.....	74.7	-	
	Limestone, light greenish gray, finely crystalline; contains <u>Osagia</u> and abundant glauconite.....	75.9	-	
	Shale, medium olive gray.....	77.2	-	
	Shale, olive.....	78.8	-	
	Shale, medium gray.....	79.8	-	
	Shale, dark gray.....	81.5	-	
Five Point Formation:				
	Limestone, light gray, finely crystalline; contains crinoids, brachiopods, <u>Osagia</u> , and glauconite.....	85.0	-	
West Branch Formation:				
	Shale, light gray.....	86.2	-	
	Shale, green.....	88.1	-	
	Shale, reddish gray.....	89.3	-	
	Limestone, light gray, very finely crystalline.....	91.0	-	
	Shale, light green.....	92.1	-	
	Shale, light gray.....	93.4	-	
	Shale, black.....	100.0	-	
	Shale, light gray.....	100.5	-	
Falls City Formation:				
Lehmer Member:				
	Limestone, light gray, very finely crystalline.....	102.1	-	
	line.....	104.0		

<u>Description</u>	<u>Depth, in feet</u>	
	<u>From</u>	<u>To</u>
Limestone, medium gray, very finely crystalline; contains pyrite and "black inclusions".....	104.0	- 107.0
Shale, light grayish green; interbedded with limestone, light gray.....	107.0	- 108.9
Reserve Member:		
Shale, medium gray.....	108.9	- 110.2
Miles Member:		
Limestone, medium gray, very finely crystalline.....	110.2	- 111.0
Onaga Formation:		
Shale, medium gray.....	111.0	- 111.4
Shale, green; interbedded with limestone, gray.....	111.4	- 112.0
Shale, light green.....	112.0	- 113.0
Shale, very light green.....	113.0	- 115.0
Shale, green.....	115.0	- 117.4
Shale, red.....	117.4	- 119.7
Shale, olive.....	119.7	- 119.9
Limestone, light greenish gray, very finely crystalline.....	119.9	- 120.7
Shale, red.....	120.7	- 124.5
Shale, light gray.....	124.5	- 127.7
Pennsylvanian System - Virgil Series - Wabaunsee Group:		
Wood Siding Formation:		
Brownville Member:		
Limestone, light gray, finely crystalline; contains <u>Osagia</u> .....	127.7	- 129.4
Shale, light gray.....	129.4	- 130.0

#### Test Hole 11-81

Location: Otoe County, SW corner sec. 25, T. 9 N., R. 11 E., approximately 205 feet north of south section line and 21 feet east of west section line.

Ground-level elevation: 1239.0 feet above mean sea level.

Started: May 14, 1981. Completed: May 14, 1981.

Total Depth: 302.0 feet.

<u>Description</u>	<u>Depth, in feet</u>	
	<u>From</u>	<u>To</u>
Quaternary System:		
Soil (no sample).....	0.0	- 3.0
Clay, tan, silty.....	3.0	- 5.0
Clay, tan, silty, sandy.....	5.0	- 12.0
Clay, red, sandy.....	12.0	- 17.5
Clay, tan, sandy.....	17.5	- 25.5
Clay, light greenish gray.....	25.5	- 32.0
Clay, medium gray, very sandy.....	32.0	- 37.0
Clay, gray to brown, sandy.....	37.0	- 45.0
Clay, light grayish green to brown, sandy.....	45.0	- 47.5
Clay, gray to olive brown, sandy.....	47.5	- 55.0
Clay, olive, sandy.....	55.0	- 70.4
Sand, very fine.....	70.4	- 77.0
Sand, fine.....	77.0	- 107.0
Sand, very fine.....	107.0	- 115.5
Sand, fine.....	115.5	- 131.0
Sand, fine, very silty.....	131.0	- 137.5
Clay, medium gray, sandy, silty.....	137.5	- 173.0
Clay, medium gray.....	173.0	- 182.0
Sand, coarse.....	182.0	- 185.0
Sand, coarse; contains gravel.....	185.0	- 195.2

Pennsylvanian System - Virgil Series - Wabaunsee Group:

Willard Formation:

Limestone, medium to dark gray, finely crystalline.....	195.2	- 197.0
Shale, light gray.....	197.0	- 200.6
Shale, medium gray.....	200.6	- 201.1
Shale, light greenish gray.....	201.1	- 203.0
Shale, dark red.....	203.0	- 204.4
Shale, grayish green.....	204.4	- 205.4

Emporia Formation:

Elmont - Reading Members:

Limestone, light gray to pale red, very finely crystalline; contains crinoids.....	205.4	- 208.3
Shale, olive.....	208.3	- 209.6
Limestone, light gray to tan, very finely crystalline; contains <u>Osagia</u> .....	209.6	- 211.0
Shale, light gray.....	211.0	- 211.8
Limestone, very light gray, very finely crystalline, pseudo-oolitic; contains brachiopods, crinoids, <u>Osagia</u> , and glaucnrite.....	211.8	- 214.4
Limestone, light gray, finely crystalline; contains crinoids, bryozoans, and <u>Osagia</u> .....	214.4	- 215.5

Auburn Formation:

Shale, light grayish green mottled red.....	215.5	- 218.0
Shale, red.....	218.0	- 234.0
Shale, olive.....	234.0	- 235.6

<u>Description</u>	<u>Depth, in feet</u>	
	<u>From</u>	<u>To</u>
Shale, medium gray.....	235.6	- 239.8
Sandstone, tan.....	239.8	- 240.0
Shale, medium gray.....	240.0	- 248.1
Wakarusa Formation:		
Limestone, medium gray, finely crystalline; contains crinoids.....	248.1	- 248.9
Shale, medium gray.....	248.9	- 252.1
Shale, black.....	252.1	- 252.2
Limestone, medium gray, finely crystalline; contains brachiopods, pyrite, and "black inclusions".....	252.2	- 253.0
Soldier Creek Formations:		
Shale, light gray.....	253.0	- 257.0
Shale, red.....	257.0	- 258.9
Shale, medium gray.....	258.9	- 259.3
Limestone, very light gray, finely crystal- line; contains <u>Osagia</u> and glauconite.....	259.3	- 261.6
Shale, light grayish green.....	261.6	- 262.4
Shale, red.....	262.4	- 266.0
Shale, light gray.....	266.0	- 275.3
Burlingame Formation:		
Limestone, medium to dark gray, finely crys- talline, pseudo-oolitic; contains brachiopods and <u>Osagia</u> .....	275.3	- 277.0
Shale, medium gray.....	277.0	- 278.2
Limestone, medium gray, finely crystalline, pseudo-oolitic; contains fusulinids and <u>Osagia</u> .....	278.2	- 279.3
Scranton Formation:		
Silver Lake - White Cloud Members:		
Shale, green.....	279.3	- 280.3
Shale, light gray.....	280.3	- 284.8
Shale, black.....	284.8	- 286.5
Shale, light greenish gray.....	286.5	- 287.0
Shale, green.....	287.0	- 295.5
Shale, red mottled green.....	295.5	- 299.0
Shale, red mottled green; interbedded with limestone, medium gray.....	299.0	- 302.0

Test Hole 12-81

Location: Otoe County, SW corner sec. 13, T. 9 N., R. 11 E.,  
approximately 212 feet north of south section line  
and 22 feet east of west section line.

Ground-level elevation: 1132.5 feet above mean sea level.

Started: May 14, 1981. Completed: May 14, 1981.

Total Depth: 92.0 feet.

	<u>Description</u>	<u>Depth, in feet</u>	
		<u>From</u>	<u>To</u>
Quaternary System:			
Soil (no sample).....		0.0	- 8.0
Clay, tan, very sandy.....		8.0	- 14.5
Sand, medium to coarse.....		14.5	- 15.0
Clay, tan, sandy.....		15.0	- 16.0
Clay, medium gray, sandy.....		16.0	- 17.0
Pennsylvanian System - Virgil Series - Shawnee Group:			
Topeka Formation:			
Curzon Member:			
(no sample).....		17.0	- 22.0
Iowa Point Member:			
Shale, light gray.....		22.0	- 22.6
Hartford Member:			
Limestone, tan to light gray, very finely crystalline; contains <u>Osagia</u> and manganese staining.....		22.6	- 23.0
Shale, black.....		23.0	- 23.1
Limestone, medium gray, finely crystalline...		23.1	- 24.2
Calhoun Formation:			
Shale, medium gray.....		24.2	- 25.0
Tecumseh - Deer Creek Formations:			
Ervine Creek Member:			
Limestone, light bluish gray, medium to coarsely crystalline; contains crinoids, brachiopods, and <u>Osagia</u> .....		25.0	- 28.0
Limestone, light bluish gray, finely crys- talline; contains <u>Osagia</u> and glauconite...		28.0	- 30.1
Limestone, light gray, very finely crystal- line; contains <u>Osagia</u> and fusulinids.....		30.1	- 32.2
Limestone, medium to dark gray, finely crystalline; contains pyrite, abundant fusulinids, and abundant black carbonaceous material.....		32.2	- 34.0
Limestone, dark gray, finely crystalline; contains pyrite, black carbonaceous material, and fusulinids.....		34.0	- 36.3
Shale, black.....		36.3	- 36.4
Limestone, medium gray, finely crystalline; contains <u>Osagia</u> and abundant fusulinids...		36.4	- 38.6
Shale, medium gray.....		38.6	- 39.5
Limestone, light gray, finely crystalline; contains crinoids, fusulinids, pyrite, and <u>Osagia</u> .....		39.5	- 40.4
Larsh Member:			
Shale, medium gray.....		40.4	- 41.0

	<u>Description</u>	<u>Depth, in feet</u>	
		<u>From</u>	<u>To</u>
	Shale, black.....	41.0	- 42.4
Rock Bluff Member:			
	Limestone, light gray to tan, very finely crystalline; contains <u>Osagia</u> and pyrite...	42.4	- 44.3
Oskaloosa - Rakes Creek Members:			
	Shale, medium gray.....	44.3	- 45.2
	Shale, medium greenish gray.....	45.2	- 47.6
	Siltstone, light greenish gray, sandy.....	47.6	- 55.0
	Shale, green.....	55.0	- 69.5
	Shale, red.....	69.5	- 72.9
Ost Member:			
	Limestone, light gray, very finely crystal- line, silty; contains <u>Osagia</u> .....	72.9	- 75.8
	Limestone, very light grayish green to white, very finely crystalline to litho- graphic; contains <u>Osagia</u> and glauconite...	75.8	- 79.1
Kenosha Member:			
	Shale, red mottled green.....	79.1	- 80.8
Lecompton Formation:			
Avoca Member:			
	Limestone, light gray, very finely crystal- line; contains <u>Osagia</u> and brachiopods.....	80.8	- 81.2
	Limestone, light to medium gray, finely crystalline; pseudo-oolitic in part; contains crinoids and <u>Osagia</u> .....	81.2	- 82.7
King Hill Member:			
	Shale, red.....	82.7	- 87.5
	Shale, medium greenish gray.....	87.5	- 89.6
Beil Member:			
	Limestone, very light gray to white, very finely crystalline; contains pyrite, glauconite, coral, and <u>Osagia</u> .....	89.6	- 92.0

#### Test Hole 13-81

Location: Otoe County, SW corner sec. 24, T. 9 N., R. 11 E.,  
approximately 85 feet north of south section line  
and 73 feet east of west section line.

Ground-level elevation: 1148.5 feet above mean sea level.

Started: May 20, 1981. Completed: May 20, 1981.

Total Depth: 197.0 feet.

<u>Description</u>	<u>Depth, in feet</u>		
	<u>From</u>	<u>To</u>	
Quaternary System:			
Soil (no sample).....	0.0	-	6.0
Clay, medium gray, sandy.....	6.0	-	14.0
Clay, tan, sandy.....	14.0	-	25.0
Clay, olive gray, sandy.....	25.0	-	32.0
(no sample).....	32.0	-	40.0
Clay, medium gray, sandy, silty.....	40.0	-	49.0
Clay, medium gray, silty, very sandy.....	49.0	-	55.0
Clay, medium gray, sandy, silty.....	55.0	--	100.5
Sand, coarse; contains gravel.....	100.5	-	107.0
Clay, medium gray, sandy.....	107.0	-	111.5
Pennsylvanian System - Virgil Series - Wabaunsee Group:			
Scranton Formation:			
Silver Lake - White Cloud Members:			
Shale, medium gray.....	111.5	-	120.0
Coal, black.....	120.0	-	120.2
Shale, medium gray.....	120.2	-	137.0
Shale, medium gray; contains brown sandstone streaks.....	137.0	-	162.9
Howard Formation:			
Limestone, dark gray, finely crystalline; contains crinoids.....	162.9	-	163.3
Shale, medium greenish gray.....	163.3	-	164.4
Limestone, dark gray, finely crystalline; contains brachiopods, pyrite, and "black inclusions".....	164.4	-	165.0
Shale, light greenish gray.....	165.0	-	168.1
Limestone, medium gray, finely crystalline; contains <u>Osagia</u> , coral, brachiopods, pyrite, "black inclusions", and abundant fusulinids.....	168.1	-	170.0
Limestone, light gray, very finely crystal- line; contains <u>Osagia</u> and pyrite.....	170.0	-	171.7
Severy Formation:			
Shale, light gray.....	171.7	-	172.2
Shale, black.....	172.2	-	173.0
Shale, medium grayish green.....	173.0	-	176.0
Coal, black.....	176.0	-	177.0
Shale, medium grayish green.....	177.0	-	183.0
Shawnee Group:			
Topeka Formation:			
Coal Creek Member:			
Limestone, dark gray, finely crystalline; contains pyrite.....	183.0	-	185.0

	<u>Description</u>	<u>Depth, in feet</u>	
		<u>From</u>	<u>To</u>
	Shale, medium gray.....	185.0	- 185.8
	Limestone, medium gray, very finely crystalline; contains brachiopods, coral, and pyrite.....	185.8	- 187.2
	Shale, medium gray.....	187.2	- 188.0
	Limestone, medium to dark gray, very finely crystalline; contains fusulinids, crinoids, and pyrite.....	188.0	- 189.6
	Limestone, light gray, finely crystalline; contains fusulinids, crinoids, pyrite, and abundant brachiopods.....	189.6	- 190.0
Holt Member:			
	Shale, medium gray.....	190.0	- 191.0
	Shale, black.....	191.0	- 192.1
DuBois Member:			
	Limestone, light to very dark gray, finely crystalline; contains brachiopods, pyrite, "black inclusions", and abundant crinoids.....	192.1	- 193.4
Turner Creek Member:			
	Shale, medium greenish gray.....	193.4	- 196.1
Sheldon Member:			
	Limestone, light to medium gray, very finely crystalline, pseudo-oolitic in part; contains crinoids, pyrite, and abundant fusulinids.....	196.1	- 197.0

#### Test Hole 14-81

Location: Otoe County, NW corner sec. 2, T. 8 N., R. 11 E., approximately 19 feet south of north section line and 177 feet east of west section line.

Ground-level elevation: 1190.0 feet above mean sea level.

Started: May 20, 1981. Completed: May 20, 1981.

Total Depth: 182.0 feet.

	<u>Description</u>	<u>Depth, in feet</u>	
		<u>From</u>	<u>To</u>
Quaternary System:			
	Soil (no sample).....	0.0	- 3.0
	Clay, tan, sandy.....	3.0	- 8.5
	Clay, medium olive gray, sandy.....	8.5	- 16.0

<u>Description</u>	<u>Depth, in feet</u>	
	<u>From</u>	<u>To</u>
Clay, tan to brown, sandy.....	16.0	- 27.0
Clay, tan, sandy.....	27.0	- 29.0
Clay, olive, sandy.....	29.0	- 39.0
Clay, dark gray, sandy.....	39.0	- 42.0
Clay, reddish gray, sandy.....	42.0	- 47.5
Clay, dark reddish gray, sandy.....	47.5	- 50.0
Clay, medium gray, sandy.....	50.0	- 53.0
Clay, light grayish green, sandy.....	53.0	- 70.0
Clay, light grayish green, very sandy.....	70.0	- 71.0
Clay, tan, sandy.....	71.0	- 77.0
Pennsylvanian System - Virgil Series - Wabaunsee Group:		
Willard Formation:		
Shale, olive.....	77.0	- 82.0
Shale, medium gray.....	82.0	- 84.7
Shale, medium gray, limy.....	84.7	- 89.0
Shale, medium gray.....	89.0	- 93.3
Emporia Formation:		
Elmont - Reading Members:		
Limestone, light gray, very finely crystalline.....	93.3	- 96.0
Shale, red.....	96.0	- 97.1
Limestone, light gray, finely crystalline; contains <u>Osagia</u> .....	97.1	- 99.2
Shale, medium gray.....	99.2	- 100.0
Limestone, light gray to tan, very finely crystalline, weathered; contains <u>crinoids, Osagia, and glauconite</u> .....	100.0	- 102.8
Auburn Formation:		
Shale, dark gray.....	102.8	- 103.8
Shale, medium grayish green.....	103.0	- 104.8
Shale, red.....	104.8	- 118.6
Shale, red mottled olive.....	118.6	- 120.0
Shale, medium gray.....	120.0	- 124.5
Coal, black.....	124.5	- 124.6
Shale, medium gray.....	124.6	- 133.6
Wakarusa Formation:		
Limestone, medium gray, finely crystalline; contains brachiopods and black carbonaceous material.....	133.6	- 134.8
Shale, medium gray.....	134.8	- 135.0
Limestone, medium to dark gray, finely crys- talline; contains brachiopods and "black inclusions".....	135.0	- 138.0
Soldier Creek Formation:		
Shale, medium gray.....	138.0	- 140.8
Limestone, light gray, very finely crystal- line, pseudo-oolitic; contains brachiopods.....	140.8	- 141.0
Shale, red.....	141.0	- 144.0

	<u>Description</u>	<u>Depth, in feet</u>	
		<u>From</u>	<u>To</u>
	Shale, gray mottled red.....	144.0	- 145.0
	Limestone, light gray, finely crystalline, pseudo-oolitic in part; contains glauconite and abundant <u>Osagia</u> .....	145.0	- 147.0
	Limestone, light gray, very finely crystal- line; contains <u>Osagia</u> .....	147.0	- 147.6
	Shale, green.....	147.6	- 148.1
	Shale, red.....	148.1	- 150.8
	Shale, medium gray.....	150.8	- 160.1
Burlingame Formation:			
	Limestone, medium gray, medium crystalline; contains coral, brachiopods, bryozoans, gastropods, and pyrite.....	160.1	- 161.4
	Shale, medium gray.....	161.4	- 162.8
	Limestone, medium to dark gray, medium crystalline, pseudo-oolitic; contains crinoids and <u>Osagia</u> .....	162.8	- 163.0
Scranton Formation:			
	Silver Lake - White Cloud Members:		
	Shale, light green.....	163.0	- 163.8
	Shale, light gray.....	163.8	- 164.8
	Shale, light gray; interbedded with lime- stone, light gray.....	164.8	- 167.5
	Coal, black.....	167.5	- 169.0
	Shale, light gray.....	169.0	- 170.5
	Shale, light gray; interbedded with lime- stone, medium gray.....	170.5	- 171.5
	Shale, light greenish gray.....	171.5	- 175.4
	Shale, red.....	175.4	- 180.7
	Limestone, light tan to gray, lithographic; interbedded with shale, red.....	180.7	- 182.0

#### Test Hole 15-81

Location: Otoe Conty, NE NW NW sec. 10, T. 8 N., R. 11 E.,  
approximately 28 feet south of north section line  
and 1036 feet east of west section line.

Ground-level elevation: 1109.0 feet above mean sea level.

Started: May 21, 1981. Completed: May 21, 1981.

Total Depth: 212.0 feet.

	<u>Description</u>	<u>Depth, in feet</u>	
		<u>From</u>	<u>To</u>
Quaternary System:			
Soil (no sample).....	0.0	-	5.0
Clay, black, sandy.....	5.0	-	13.0
Clay, dark greenish gray, sandy.....	13.0	-	22.0
Clay, green, sandy.....	22.0	-	32.5
Clay, medium gray, sandy.....	32.5	-	37.0
Clay, dark gray.....	37.0	-	48.0
Clay, light gray.....	48.0	-	68.5
Clay, light gray, sandy.....	68.5	-	96.0
Clay, light gray, contains sand, coarse, and gravel.....	96.0	-	106.0
Pennsylvanian System - Virgil Series - Wabaunsee Group:			
Stotler Formation:			
Dry Member:			
Shale, light greenish gray.....	106.0	-	107.3
Dover Member:			
Limestone, dark gray, finely crystalline, pseudo-oolitic in part; contains crinoids and "black inclusions".....	107.3	-	108.0
Pillsbury Formation:			
Shale, light gray.....	108.0	-	129.2
Zeandale Formation:			
Maple Hill Member:			
Limestone, dark gray, finely crystalline; contains crinoids, brachiopods, black carbonaceous material, and pyrite.....	129.2	-	130.8
Wamego Member:			
Shale, medium gray.....	130.8	-	134.9
Tarkio Member:			
Limestone, tan, lithographic.....	134.9	-	135.1
Shale, medium gray.....	135.1	-	136.1
Limestone, light gray, finely crystalline; contains <u>Osagia</u> .....	136.1	-	138.6
Willard Formation:			
Siltstone, green, very sandy.....	138.6	-	146.4
Sandstone, grayish green.....	146.4	-	147.5
Shale, medium gray.....	147.5	-	169.2
Limestone, medium gray, finely crystalline; contains brachiopods and coral.....	169.2	-	174.8
Coal, black.....	174.8	-	175.0
Limestone, medium gray, finely crystalline; contains pyrite.....	175.0	-	175.2
Shale, light grayish green.....	175.2	-	179.0
Emporia Formation:			
Elmont - Reading Members:			
Limestone, light gray to tan, very finely crystalline; contains <u>Osagia</u> and abundant pyrite.....	179.0	-	180.0

<u>Description</u>	<u>Depth, in feet</u>	
	<u>From</u>	<u>To</u>
Shale, medium gray; interbedded with limestone, grayish green.....	180.0	- 181.0
Shale, reddish gray.....	181.0	- 181.6
Shale, medium gray.....	181.6	- 182.0
Limestone, light gray to tan, very finely crystalline; contains <u>Osagia</u> .....	182.0	- 183.7
Shale, medium gray.....	183.7	- 185.1
Limestone, light gray, very finely crystalline; contains brachiopods, <u>Osagia</u> , and glauconite.....	185.1	- 187.7
Auburn Formation:		
Shale, green mottled red.....	187.7	- 188.0
Shale, red mottled green.....	188.0	- 192.0
Shale, red.....	192.0	- 197.0
Shale, red mottled olive.....	197.0	- 204.3
Shale, olive.....	204.3	- 205.0
Shale, light gray.....	205.0	- 206.0
Shale, medium gray.....	206.0	- 212.0

#### Test Hole 16-81

Location: Otoe County, NW corner SW sec. 18, T. 8 N., R. 11 E., approximately 2551 feet north of south section line and 20 feet east of west section line.

Ground-level elevation: 1087.0 feet above mean sea level.

Started: May 21, 1981. Completed: May 21, 1981.

Total Depth: 317.0 feet.

<u>Description</u>	<u>Depth, in feet</u>	
	<u>From</u>	<u>To</u>
Quaternary System:		
Soil (no sample).....	0.0	- 3.0
Clay, brown, sandy.....	3.0	- 9.5
Clay, dark brown, very sandy.....	9.5	- 14.5
Clay, tan, sandy.....	14.5	- 33.0
Sand, medium to coarse; contains gravel.....	33.0	- 37.0
Clay, olive gray.....	37.0	- 48.5
Clay, olive gray, very sandy.....	48.5	- 58.0
Sand, coarse; contains gravel.....	58.0	- 67.7

<u>Description</u>	<u>Depth, in feet</u>	
	<u>From</u>	<u>To</u>
Permian System - Big Blue Series - Admire Group:		
Onaga Formation:		
Shale, tan, weathered.....	67.7	-
Shale, light bluish green.....	68.2	-
Shale, dark red.....	69.5	-
Limestone, light gray, very finely crystalline.....	69.7	-
Shale, red.....	71.2	-
Shale, red mottled olive.....	73.0	-
Shale, medium gray.....	77.5	-
Pennsylvanian System - Virgil Series - Wabaunsee Group:		
Wood Siding Formation:		
Brownville Member:		
Limestone, light gray, finely crystalline....	80.3	-
Pony Creek - Plumb Members:		
Shale, light gray.....	83.5	-
Shale, light green.....	84.0	-
Shale, red mottled green.....	85.0	-
Shale, light green mottled red.....	86.7	-
Sandstone, medium greenish gray.....	88.4	-
Shale, red.....	90.0	-
Shale, green.....	93.4	-
Shale, medium gray.....	96.5	-
Nebraska City Member:		
Limestone, medium gray, medium crystalline; contains brachiopods.....	100.4	-
Pillsbury - Root Formations:		
Coal, black.....	102.6	-
Shale, green.....	103.4	-
Shale, red mottled green.....	112.1	-
Limestone, medium gray, medium crystalline; contains <u>Osacia</u> .....	113.1	-
Shale, green.....	113.6	-
Shale, red.....	114.6	-
Shale, olive.....	121.3	-
Shale, medium greenish gray.....	124.0	-
Shale, medium gray.....	145.0	-
Zeandale Formation:		
Maple Hill Member:		
Limestone, dark gray, finely crystalline; contains brachiopods and fusulinids.....	173.2	-
Willard Formation:		
Shale, medium gray.....	174.4	-
Sandstone, medium gray, very limy.....	176.3	-
Shale, light greenish gray.....	177.6	-
Sandstone, light gray; contains black carbonaceous material.....	191.0	-
Shale, medium gray.....	202.3	-
		218.8

<u>Description</u>	<u>Depth, in feet</u>	
	<u>From</u>	<u>To</u>
Shale, medium gray; contains limy sandstone streaks throughout and limestone streaks near bottom.....	218.8	- 223.3
Shale, medium gray; interbedded with limestone, medium gray.....	223.3	- 225.0
Shale, olive.....	225.0	- 227.0
Shale, red mottled gray.....	227.0	- 231.4
Emporia Formation:		
Elmont - Reading Members:		
Limestone, light grayish green, very finely crystalline; contains crinoids and glauconite.....	231.4	- 232.0
Limestone, very light gray, very finely crystalline, pseudo-oolitic; contains <u>Osagia</u> and pyrite.....	232.0	- 234.6
Shale, light greenish gray.....	234.6	- 235.1
Limestone, very light gray, very finely crystalline, pseudo-oolitic; contains brachiopods, crinoids, <u>Osagia</u> , and glauconite.....	235.1	- 240.3
Auburn Formation:		
Shale, green.....	240.3	- 240.8
Shale, red.....	240.8	- 251.3
Sandstone, light grayish green.....	251.3	- 258.5
Shale, medium gray.....	258.5	- 268.9
Wakarusa Formation:		
Limestone, medium gray, finely crystalline; contains brachiopods and pyrite.....	268.9	- 272.3
Soldier Creek Formation:		
Shale, medium gray.....	272.3	- 273.0
Limestone, light greenish gray, lithographic; interbedded with shale, medium gray.....	273.0	- 275.0
Shale, green mottled red.....	275.0	- 277.4
Limestone, very light gray, very finely crystalline; contains <u>Osagia</u> .....	277.4	- 280.4
Shale, olive.....	280.4	- 281.4
Shale, medium gray.....	281.4	- 290.9
Burlingame Formation:		
Limestone, light gray, finely crystalline; contains black carbonaceous material.....	290.9	- 295.7
Scranton Formation:		
Silver Lake - White Cloud Members:		
Shale, medium gray.....	295.7	- 296.7
Shale, black.....	296.7	- 298.7
Shale, medium gray.....	298.7	- 301.6
Limestone, medium gray, finely crystalline; contains fusulinids and pyrite; interbedded with shale, medium gray.....	301.6	- 303.0
Shale, black.....	303.0	- 303.2

	<u>Description</u>	<u>Depth, in feet</u>	
		<u>From</u>	<u>To</u>
Limestone, light grayish green, very finely crystalline; interbedded with shale, green.....		303.2	- 312.0
Shale, red.....		312.0	- 313.0
Shale, red; contains limestone streaks.....		313.0	- 317.0

Test Hole 17-81

Location: Otoe County, SE SW SW sec. 8, T. 8 N., R. 11 E., approximately 26 feet north of south section line and 1150 feet east of west section line.

Ground-level elevation: 1063.0 feet above mean sea level.

Started: May 22, 1981. Completed: May 22, 1981.

Total Depth: 137.0 feet.

	<u>Description</u>	<u>Depth, in feet</u>	
		<u>From</u>	<u>To</u>
Quaternary System:			
Soil (no sample).....		0.0	- 6.0
Clay, brown, sandy.....		6.0	- 15.0
Clay, olive gray, sandy.....		15.0	- 17.5
Clay, tan to gray, sandy.....		17.5	- 23.0
Clay, olive, very sandy.....		23.0	- 26.0
Clay, medium gray, sandy.....		26.0	- 26.5
Sand, medium to coarse.....		26.5	- 29.0
Clay, medium gray, very sandy.....		29.0	- 34.5
Sand, medium to coarse.....		34.5	- 35.0

Pennsylvanian System - Virgil Series - Wabaunsee Group:

Soldier Creek Formation:

Shale, red.....	35.0	-	37.2
Limestone, yellow to tan, finely crystalline, weathered; contains manganese staining.....	37.2	-	37.4
Shale, medium gray to tan.....	37.4	-	39.0
Shale, red.....	39.0	-	44.5
Shale, medium gray.....	44.5	-	47.4

Burlingame Formation:

Limestone, light gray, finely crystalline, pseudo-oolitic; contains bryozoans and Osagia.....	47.4	-	50.2
---	------	---	------

	<u>Description</u>	<u>Depth, in feet</u>		
		<u>From</u>	<u>To</u>	
Scranton Formation:				
Silver Lake Member:				
Shale, medium gray.....	50.2	-	51.0	
Shale, green.....	51.0	-	51.6	
Shale, red.....	51.6	-	60.5	
Shale, green mottled red.....	60.5	-	61.5	
Shale, medium gray.....	61.5	-	65.4	
Rulo Member:				
Limestone, medium gray, finely crystalline...	65.4	-	66.3	
Cedarvale Member:				
Shale, medium gray.....	66.3	-	67.0	
Shale, black.....	67.0	-	68.1	
Shale, medium gray.....	68.1	-	73.3	
Shale, green.....	73.3	-	74.3	
Shale, red.....	74.3	-	75.1	
Shale, green mottled red.....	75.1	-	77.1	
Happy Hollow Member:				
Limestone, light greenish gray, finely crystalline, silty; contains fusulines, bryozoans, <u>Osagia</u> and glauconite.....	77.1	-	79.3	
Limestone, tan to light greenish gray, finely crystalline; contains crinoids and <u>Osagia</u> .....	79.3	-	80.4	
White Cloud Member:				
Shale, red mottled olive.....	80.4	-	82.5	
Shale, olive.....	82.5	-	83.5	
Shale, light greenish gray.....	83.5	-	86.5	
Shale, medium greenish gray.....	86.5	-	88.0	
Shale, medium gray, sandy.....	88.0	-	102.0	
Shale, medium gray.....	102.0	-	122.0	
Shale, medium gray; contains brown sandstone streaks.....	122.0	-	137.0	

#### Test Hole 18-81

Location: Otoe County, SW corner sec. 13, T. 8 N., R. 10 E.,  
approximately 93 feet north of south section line  
and 19 feet east of west section line.

Ground-level elevation: 1165.0 feet above mean sea level.

Started: May 22, 1981. Completed: May 22, 1981.

Total Depth: 122.0 feet.

<u>Description</u>	<u>Depth, in feet</u>	
	<u>From</u>	<u>To</u>
Quaternary System:		
Soil (no sample).....	0.0	- 4.5
Clay, tan, sandy.....	4.5	- 9.0
Clay, dark brown, sandy.....	9.0	- 13.5
Clay, tan, sandy.....	13.5	- 14.5
Permian System - Big Blue Series - Council Grove Group:		
Red Eagle Formation:		
Bennett Member:		
Shale, very light green to tan, weathered....	14.5	- 19.4
Glenrock Member:		
Limestone, tan to light gray, finely crystalline, weathered; contains fusulinids.....	19.4	- 20.5
Limestone, tan, very finely crystalline, weathered; contains abundant <u>Osagia</u> and manganese staining.....	20.5	- 21.2
Limestone, tan to medium gray, finely crystalline; contains manganese staining.....	21.2	- 22.5
Johnson Formation:		
Shale, green.....	22.5	- 24.0
Shale, medium gray.....	24.0	- 28.4
Shale, dark green.....	28.4	- 29.0
Shale, red mottled gray.....	29.0	- 29.7
Shale, medium gray.....	29.7	- 30.6
Foraker Formation:		
Long Creek Member:		
Limestone, light greenish gray, very finely crystalline, argillaceous.....	30.6	- 33.0
Limestone, light greenish gray to medium gray, very finely crystalline.....	33.0	- 35.0
Shale, green.....	35.0	- 35.2
Limestone, medium gray, very finely crystalline to lithographic.....	35.2	- 37.1
Hughes Creek Member:		
Shale, medium gray.....	37.1	- 38.1
Shale, light gray.....	38.1	- 39.0
Shale, dark gray.....	39.0	- 43.2
Limestone, very dark gray, medium crystalline, pseudo-oolitic; contains abundant <u>Osagia</u> .....	43.2	- 45.1
Limestone, medium gray, finely crystalline; contains brachiopods, <u>Osagia</u> , and "black inclusions".....	45.1	- 46.3
Shale, medium gray.....	46.3	- 52.8
Limestone, medium gray, finely crystalline; contains abundant brachiopods; interbedded with shale, medium gray.....	52.8	- 53.0
Shale, medium gray.....	53.0	- 53.5

<u>Description</u>	<u>Depth, in feet</u>	
	<u>From</u>	<u>To</u>
Limestone, light gray, finely crystalline; contains abundant brachiopods; interbedded with shale, medium gray.....	53.5	- 54.5
Shale, light greenish gray.....	54.5	- 59.2
Limestone, dark gray, finely crystalline; contains brachiopods and pyrite.....	59.2	- 60.0
Shale, very dark gray.....	60.0	- 61.7
Limestone, medium gray, variably crystal- line; contains <u>Osagia</u> , coral, "black inclusions", and pyrite.....	61.7	- 63.2
Shale, medium gray.....	63.2	- 63.6
Limestone, medium to dark gray, finely crystalline; contains fusulinids, crinoids, <u>Osagia</u> , and pyrite.....	63.6	- 63.8
Shale, medium gray.....	63.8	- 66.5
Limestone, medium to dark gray, finely crystalline; contains brachiopods, crinoids, and <u>Osagia</u> ; interbedded with with shale, medium gray.....	66.5	- 70.2
Shale, black.....	70.2	- 71.0
Limestone, medium gray, very finely crystal- line; contains coral, crinoids, and <u>Osagia</u> .....	71.0	- 72.2
Shale, medium gray.....	72.2	- 74.0
Limestone, medium to dark gray, finely crystalline, pseudo-oolitic; contains crinoids and <u>Osagia</u> .....	74.0	- 76.0
Shale, medium gray.....	76.0	- 76.5
Limestone, medium gray, medium crystalline; contains abundant <u>Osagia</u> .....	76.5	- 76.7
Shale, medium gray.....	76.7	- 78.2
Shale, black.....	78.2	- 78.6
Americus Member:		
Limestone, medium gray, finely crystalline; contains crinoids and pyrite.....	78.6	- 79.8
Shale, medium gray.....	79.8	- 81.5
Limestone, light gray, very finely crystal- line; contains crinoids.....	81.5	- 82.0
Limestone, medium gray, very finely crystalline.....	82.0	- 82.9
Admire Group:		
West Branch - Hamlin Formations:		
Shale, medium gray.....	82.9	- 86.4
Limestone, medium gray, very finely crystalline.....	86.4	- 86.7
Shale, medium gray.....	86.7	- 88.3
Limestone, light gray, lithographic; interbedded with shale, green.....	88.3	- 89.0

<u>Description</u>	<u>Depth, in feet</u>		
	<u>From</u>	<u>To</u>	
Limestone, very light grayish green, very finely crystalline; contains glauconite; interbedded with shale, green, at 91.0-92.4.....	89.0	-	92.4
Shale, red.....	92.4	-	95.0
Shale, red; interbedded with limestone, light gray.....	95.0	-	98.3
Limestone, very light grayish green, very finely crystalline; contains glauconite; interbedded with shale, green.....	98.3	-	101.0
Shale, red.....	101.0	-	104.2
Limestone, medium gray, very finely crystalline; contains brachiopods and crinoids...	104.2	-	104.8
Shale, medium gray.....	104.8	-	107.0
Shale, red.....	107.0	-	109.5
Shale, grayish green.....	109.5	-	110.5
Shale, pale red; interbedded with limestone, light gray.....	110.5	-	114.0
Shale, medium gray; interbedded with limestone, light gray.....	114.0	-	115.0
Shale, olive.....	115.0	-	117.0
Shale, green.....	117.0	-	118.0
Shale, olive.....	118.0	-	119.5
Shale, gray mottled olive.....	119.5	-	120.5
Limestone, dark gray, finely crystalline; contains <u>Osagia</u> , pyrite, and black carbonaceous material.....	120.5	-	120.8
Shale, medium gra.....	120.8	-	122.0

Test Hole 19-81

Location: Otoe County, SW corner sec. 11, T. 10 N., R. 8 E., approximately 23 feet north of south section line and 103 feet east of west section line.

Ground-level elevation: 1143.0 feet above mean sea level.

Started: May 22, 1981. Completed: May 22, 1981.

Total Depth: 152.0 feet.

	<u>Description</u>	<u>Depth, in feet</u>	
		<u>From</u>	<u>To</u>
Quaternary System:			
Soil (no sample).....		0.0	- 4.0
Silt, brown, sandy.....		4.0	- 9.0
Clay, brown, very sandy.....		9.0	- 14.0
Clay, brown, sandy.....		14.0	- 22.0
Clay, light brown, sandy.....		22.0	- 23.5
Clay, medium gray.....		28.5	- 29.0
Clay, dark gray.....		29.0	- 35.0
Clay, black.....		35.0	- 38.0
Clay, dark gray.....		38.0	- 39.0
Clay, medium gray.....		39.0	- 41.5
Clay, tan.....		41.5	- 48.0
Pennsylvanian System - Virgil Series - Wabaunsee Group:			
Willard Formation:			
Shale, olive.....		48.0	- 50.0
Limestone, medium gray, finely crystalline, pseudo-oolitic; contains brachiopods, crinoids, and <u>Osagia</u> .....		50.0	- 50.6
Shale, light gray.....		50.6	- 55.0
Shale, red.....		55.0	- 55.7
Shale, medium gray.....		55.7	- 56.5
Emporia Formation:			
Elmont - Reading Members:			
Limestone, light grayish green, very finely crystalline, pseudo-oolitic in part; contains crinoids and <u>Osagia</u> .....		56.5	- 57.1
Shale, medium gray; interbedded with lime- stone, gray.....		57.1	- 59.0
Shale, medium gray.....		59.0	- 59.6
Limestone, light gray, very finely crystal- line; contains <u>Osagia</u> .....		59.6	- 61.2
Shale, medium gray.....		61.2	- 62.2
Limestone, light gray, very finely crystal- line; contains <u>Osagia</u> .....		62.2	- 65.5
Auburn Formation:			
Shale, medium greenish gray.....		65.5	- 67.0
Shale, red.....		67.0	- 80.0
Shale, olive.....		80.0	- 82.0
Shale, light gray.....		82.0	- 84.5
Shale, medium gray.....		84.5	- 92.0
Shale, medium gray; contains thin sandstone streaks.....		92.0	- 94.5
Wakarusa Formation:			
Limestone, light gray, finely crystalline; interbedded with shale, medium gray.....		94.5	- 98.5
Soldier Creek Formation:			
Shale, medium gray.....		98.5	- 100.5
Shale, medium gray; interbedded with lime- stone, light gray.....		100.5	- 103.6

<u>Description</u>	<u>Depth, in feet</u>		
	<u>From</u>	<u>To</u>	
Limestone, light gray, irregularly crystalline; contains <u>Osagia</u> and glauconite.....	103.6	-	106.5
Shale, medium gray.....	106.5	-	107.0
Shale, medium gray; contains brown sandstone streaks.....	107.0	-	117.1
Burlingame Formation:			
Limestone, light to medium gray, finely crystalline; contains <u>Osagia</u> .....	117.1	-	119.0
Limestone, light gray, finely crystalline; contains glauconite.....	119.0	-	120.2
Limestone, light gray, finely crystalline; contains crinoids; interbedded with shale, medium gray.....	120.2	-	120.6
Scranton Formation:			
Silver Lake - White Cloud Members:			
Shale, black.....	120.6	-	121.0
Shale, green.....	121.0	-	127.5
Shale, red.....	127.5	-	139.0
Shale, medium gray.....	139.0	-	142.3
Coal, black.....	142.3	-	142.6
Shale, medium gray.....	142.6	-	152.0

#### Test Hole 20-81

Location: Otoe County, NW NE NW sec. 21, T. 8 N., R. 10 E., approximately 12 feet south of north section line and 1362 feet east of west section line.

Ground-level elevation: 1213.0 feet above mean sea level.

Started: May 26, 1981. Completed: May 26, 1981.

Total Depth: 197.0 feet.

<u>Description</u>	<u>Depth, in feet</u>		
	<u>From</u>	<u>To</u>	
Quaternary System:			
Soil (no sample).....	0.0	-	4.0
Clay, tan, sandy.....	4.0	-	8.5
Clay, olive, sandy.....	8.5	-	19.0
Clay, medium gray, sandy.....	19.0	-	22.0
Clay, brown, sandy.....	22.0	-	34.0
Sand, medium, clayey.....	34.0	-	41.0
Sand, coarse, and gravel.....	41.0	-	43.0
Gravel, clayey.....	43.0	-	47.0

<u>Description</u>	<u>Depth, in feet</u>	
	<u>From</u>	<u>To</u>
Clay, medium gray, sandy, very silty.....	47.0	- 62.0
Clay, light gray, sandy.....	62.0	- 71.0
Clay, reddish gray, sandy.....	71.0	- 77.0
Clay, reddish gray, very sandy.....	77.0	- 85.8
Permian System - Big Blue Series - Council Grove Group:		
Grenola Formation:		
Burr Member:		
Limestone, light gray, very finely crystalline, pseudo-oolitic; contains brachipods.....	85.8	- 88.7
Roca Formation:		
Shale, light grayish green.....	88.7	- 92.5
Shale, very light gray.....	92.5	- 94.0
Shale, medium grayish green.....	94.0	- 97.6
Shale, red mottled green.....	97.6	- 101.0
Shale, red mottley gray.....	101.0	- 101.8
Shale, medium gray; contains thin limestone steaks.....	101.8	- 104.0
Shale, light gray.....	104.0	- 106.5
Shale, red.....	106.5	- 107.4
Shale, medium gray.....	107.4	- 108.8
Limestone, very light gray, lithographic.....	108.8	- 109.0
Shale, reddish gray.....	109.0	- 110.5
Shale, grayish green.....	110.5	- 112.7
Red Eagle Formation:		
Howe Member:		
Shale, medium gray, limy.....	112.7	- 113.3
Shale, grayish tan.....	113.3	- 115.2
Limestone, medium gray to tan, variably crystalline, pseudo-oolitic; contains <u>Osagia</u> .....	115.2	- 118.4
Bennett Member:		
Shale, light gray.....	118.4	- 121.0
Shale, gray mottled olive.....	121.0	- 124.0
Shale, black.....	124.0	- 124.7
Shale, medium gray.....	124.7	- 125.5
Shale, black.....	125.5	- 126.5
Glenrock Member:		
Limestone, medium to dark gray, very finely crystalline; contains abundant black carbonaceus material.....	126.5	- 128.4
Johnson Formation:		
Shale, light grayish green.....	128.4	- 129.4
Shale, medium gray.....	129.4	- 130.1
Limestone, medium gray, finely crystalline; interbedded with shale, medium gray.....	130.1	- 131.1
Shale, medium greenish gray.....	131.1	- 137.2
Shale, red.....	137.2	- 138.2
Shale, medium gray.....	138.2	- 139.0

<u>Description</u>	<u>Depth, in feet</u>	
	<u>From</u>	<u>To</u>
Foraker Formation:		
Long Creek Member:		
Limestone, light gray, very finely crystalline.....	139.0	- 141.5
Limestone, medium gray, very finely crystal- line; contains glauconite.....	141.5	- 142.5
Hughes Creek Member:		
Shale, light gray.....	142.5	- 145.0
Shale, medium gray.....	145.0	- 152.0
Limestone, medium to dark gray, variably crystalline, pseudo-oolitic; contains brachiopods, <u>Osagia</u> , and "black inclusions".....	152.0	- 154.5
Shale, medium gray.....	154.5	- 162.0
Shale, light gray.....	162.0	- 167.8
Limestone, medium gray, finely crystalline...	167.8	- 168.5
Shale, black.....	168.5	- 170.1
Limestone, medium gray, finely crystalline; contains fusulinids, brachiopods, <u>Osagia</u> , and "black inclusions".....	170.1	- 171.9
Shale, dark gray.....	171.9	- 176.5
Shale, medium gray.....	176.5	- 178.8
Shale, medium gray; interbedded with lime- stone, light gray.....	178.8	- 180.0
Shale, dark gray.....	180.0	- 180.4
Limestone, medium gray, finely crystalline; contains brachiopods, <u>Osagia</u> , and "black inclusions".....	180.4	- 181.5
Shale, medium gray.....	181.5	- 183.0
Limestone, medium gray, finely crystalline, pseudo-oolitic; contains brachiopods, bryozoans, crinoids, fusulinids, and <u>Osagia</u> .....	183.0	- 185.0
Shale, medium gray.....	185.0	- 188.0
Shale, black.....	188.0	- 188.4
Americus Member:		
Limestone, dark gray, very finely crystalline.....	188.4	- 189.7
Shale, dark gray.....	189.7	- 191.3
Limestone, medium gray, finely crystalline; contains coral, crinoids, and brachiopods.....	191.3	- 192.4
Admire Group:		
Hamlin Formation:		
Shale, medium gray.....	192.4	- 196.0
Limestone, medium gray, finely crystalline; pseudo-oolitic; contains brachiopods and glauconite.....	196.0	- 197.0

Test Hole 21-81

Location: Otoe County, SW corner NW sec. 21, T. 8 N., R. 10 E., approximately 2604 feet south of north section line and 30 feet east of west section line.

Ground-level elevation: 1236.0 feet above mean sea level.

Started: May 26, 1981. Completed: May 26, 1981.

Total Depth: 287.0 feet.

<u>Description</u>	<u>Depth, in feet</u>	
	<u>From</u>	<u>To</u>
Quaternary System:		
Soil (no sample).....	0.0	-
Clay, red, silty.....	5.0	9.5
Clay, tan, sandy.....	9.5	-
Clay, tan, very sandy.....	13.0	-
Clay, olive gray, sandy.....	22.0	-
Clay, brown to gray, sandy.....	29.5	-
Clay, olive gray, sandy.....	41.0	-
Sand, fine, silty.....	49.0	-
Clay, gray, very sandy; contains gravel.....	61.5	-
Clay, dark gray, very sandy; contains gravel.....	72.0	-
Clay, light gray, very sandy.....	85.0	-
Clay, light brown to gray, sandy.....	92.0	-
Clay, light gray, sandy.....	108.0	-
Clay, medium gray, sandy.....	112.0	-
Clay, dark gray, sandy.....	128.0	-
Clay, medium gray.....	143.0	-
Clay, dark gray.....	148.0	-
Clay, medium gray, sandy.....	152.5	-
Clay, medium gray.....	158.0	-
Clay, dark gray.....	169.0	-
Clay, light gray, sandy.....	175.0	-
Sand, coarse; contains gravel.....	181.4	-
Clay, greenish gray.....	183.0	-
	190.0	190.0
Permian System - Big Blue Series - Admire Group:		
West Branch Formation:		
Shale, light green.....	190.0	-
Limestone, light gray to green, very finely crystalline; contains bryozoans, brachiopods, and crinoids.....	191.6	-
Shale, light gray.....	192.1	-
Shale, medium gray.....	199.0	-
Shale, dark gray.....	200.0	-
Falls City Formation:		
Limestone, light gray, finely crystalline....	202.6	-
	203.5	

	<u>Description</u>	<u>Depth, in feet</u>	
		<u>From</u>	<u>To</u>
	Shale, medium gray.....	203.5	- 203.7
	Limestone, light gray, very finely crystalline.....	203.7	- 207.2
	Shale, medium gray.....	207.2	- 210.4
	Limestone, medium to dark gray, finely crystalline.....	210.4	- 212.0
	Limestone, light gray, very finely crystal- line; contains <u>Osagia</u> and "black inclusions".....	212.0	- 213.5
Onaga Formation:			
	Hauxby - Towle Members:		
	Shale, medium greenish gray.....	213.5	- 215.5
	Shale, light grayish green.....	215.5	- 218.5
	Shale, medium gray.....	218.5	- 220.5
	Shale, red mottled green.....	220.5	- 223.0
	Limestone, bluish gray to very light gray, very finely crystalline; interbedded with shale, red.....	223.0	- 224.0
	Shale, red.....	224.0	- 229.0
	Shale, olive.....	229.0	- 231.5
Pennsylvanian System - Virgil Series - Wabaunsee Group:			
Wood Siding Formation:			
	Brownville Member:		
	Limestone, very light gray, variably crys- talline; contains fusulinids.....	231.5	- 233.6
	Pony Creek - Plumb Members:		
	Shale, medium gray.....	233.6	- 236.2
	Shale, red.....	236.2	- 241.2
	Limestone, light gray, finely crystalline; interbedded with shale, red.....	241.2	- 241.5
	Shale, red.....	241.5	- 244.5
	Shale, light grayish green.....	244.5	- 245.5
	Shale, dark gray.....	245.5	- 250.4
Nebraska City Member:			
	Limestone, dark gray, very finely crystal- line; contains brachiopods, fusulinids, black carbonaceous material, and pyrite...	250.4	- 251.0
Stoler - Root Formations:			
	Shale, black.....	251.0	- 252.5
	Coal, black.....	252.5	- 252.6
	Shale, medium gray.....	252.6	- 259.0
	Shale, red mottled green.....	259.0	- 261.5
	Limestone, very light gray to white, very finely crystalline to lithographic; contains fusulinids and crinoids.....	261.5	- 265.3
	Shale, red.....	265.8	- 267.0
	Shale, olive.....	267.0	- 272.5
	Shale, medium gray.....	272.5	- 287.0

Test Hole 22-81

Location: Otoe County, SE corner sec. 8, T. 8 N., R. 11 E., approximately 136 feet north of south section line and 20 feet west of east section line.

Ground-level elevation: 1071.5 feet above mean sea level.

Started: May 27, 1981. Completed: May 27, 1981.

Total Depth: 171.0 feet.

	<u>Description</u>	<u>Depth, in feet</u>	
		<u>From</u>	<u>To</u>
Quaternary System:			
Soil (no sample).....		0.0	- 5.0
Silt, black.....		5.0	- 8.0
Clay, medium gray.....		8.0	- 13.0
Clay, green.....		13.0	- 23.0
Clay, medium gray.....		23.0	- 30.5
Clay, greenish gray, silty.....		30.5	- 73.5
Sand, medium to coarse, and gravel, medium....		73.5	- 78.6
Pennsylvanian System - Virgil Series - Wabaunsee Group:			
Wood Siding Formation:			
Nebraska City Member:			
Limestone, medium greenish gray, finely crystalline; contains brachiopods and glauconite.....		78.6	- 79.5
Pillsbury - Root Formations:			
Shale, black.....		79.5	- 81.2
Shale, medium greenish gray.....		81.2	- 87.4
Shale, red.....		87.4	- 90.3
Shale, red; interbedded with limestone, gray.....		90.3	- 93.7
Limestone, light gray, very finely crystalline; contains fusulinids.....		93.7	- 94.2
Shale, light greenish gray.....		94.2	- 96.4
Limestone, medium gray, finely crystalline; interbedded with shale, medium gray.....		96.4	- 98.5
Shale, grayish olive.....		98.5	- 100.7
Shale, medium gray.....		100.7	- 128.9
Sandstone, gray; contains black carbonaceous material.....		128.9	- 129.9
Shale, medium gray.....		129.9	- 147.0
Shale, medium gray; contains pyrite.....		147.0	- 152.0
Shale, medium gray.....		152.0	- 158.6
Zeandale Formation:			
Maple Hill Member:			
Limestone, medium gray, very finely crystalline; contains brachiopods.....		158.6	- 160.2

<u>Description</u>	<u>Depth, in feet</u>	
	<u>From</u>	<u>To</u>
Wamego Member:		
Shale, medium gray.....	160.2	- 163.8
Tarkio Member:		
Limestone, medium gray, very finely crystal-line; contains brachiopods.....	163.8	- 164.5
Shale, medium greenish gray.....	164.5	- 167.0
Shale, dark gray.....	167.0	- 167.2
Limestone, medium gray, very finely crystalline.....	167.2	- 168.6
Limestone, light gray, finely crystalline; contains abundant brachiopods; interbedded with shale, green.....	168.6	- 168.9
Willard Formation:		
Shale, very dark gray.....	168.9	- 170.1
Shale, medium gray.....	170.1	- 171.0

NRC FORM 335 (7-77) U.S. NUCLEAR REGULATORY COMMISSION BIBLIOGRAPHIC DATA SHEET		1. REPORT NUMBER (Assigned by DDCI) NUREG/CR-2411
4. TITLE AND SUBTITLE (Add Volume No., if appropriate) Regional Tectonics and Seismicity of Eastern Nebraska Annual Report July 1980 - June 1981		2. (Leave blank)
7. AUTHOR(S) R. R. Burchett		3. RECIPIENT'S ACCESSION NO.
9. PERFORMING ORGANIZATION NAME AND MAILING ADDRESS (Include Zip Code) Nebraska Geological Survey Institute of Agriculture and Natural Resources University of Nebraska-Lincoln Lincoln, NE 68588-0517		5. DATE REPORT COMPLETED MONTH October YEAR 1981
12. SPONSORING ORGANIZATION NAME AND MAILING ADDRESS (Include Zip Code) Division of Health, Siting and Waste Management Office of Nuclear Regulatory Research U.S. Nuclear Regulatory Commission Washington, DC 20555		6. (Leave blank)
13. TYPE OF REPORT Technical		7. DATE REPORT ISSUED MONTH November YEAR 1981
15. SUPPLEMENTARY NOTES		8. (Leave blank)
16. ABSTRACT (200 words or less) <p>This annual report presents and interprets the information obtained by the Conservation Survey Division (Nebraska Geological Survey) during contract year July 1, 1980 to June 30, 1981, under contract NRC 04 76 315 with the USNRC. The information pertains to the geology, structure, tectonics, and seismicity of eastern Nebraska with emphasis on central Otoe County. Some of the information presented here result from a combination of studies begun in earlier years but the greater part results from studies begun during the contract year.</p> <p>The scope of the studies is summarized as follows:</p> <ol style="list-style-type: none"> <li>1. Rock outcrops in Otoe County were reexamined and reevaluated, and 22 test holes were drilled to determine the altitude of the upper surface of the Elmont Limestone of Pennsylvanian age;</li> <li>2. One new seismograph was installed in eastern Nebraska;</li> <li>3. Gravity surveys in eastern Nebraska were extended;</li> <li>4. Ground magnetic surveys in Otoe County were made and evaluated.</li> <li>5. Age dating of two Precambrian cores.</li> </ol> <p>Discussion of the results of these studies constitute the remainder of this report.</p>		9. PERIOD COVERED (Inclusive dates) July 1, 1980 - June 30, 1981
17. KEY WORDS AND DOCUMENT ANALYSIS Nebraska, Tectonics, Seismic Network		10. PROJECT/TASK/WORK UNIT NO. FIN B5774
17a. DESCRIPTORS		
17b. IDENTIFIERS/OPEN-ENDED TERMS		
18. AVAILABILITY STATEMENT Unlimited		19. SECURITY CLASS (This report) Unclassified
		20. SECURITY CLASS (This page) Unclassified
		21. NO. OF PAGES
		22. PRICE \$

UNITED STATES  
NUCLEAR REGULATORY COMMISSION  
WASHINGTON, D. C. 20585

OFFICIAL BUSINESS  
PENALTY FOR PRIVATE USE, \$300

POSTAGE AND FEES PAID  
U.S. NUCLEAR REGULATORY  
COMMISSION



120555064215 2 ANRAK6  
US NRC  
ADM DOCUMENT CONTROL DESK  
PDR  
016 DC 20555  
WASHINGTON