TELEDYNE ISOTOPES

MIDWEST LABORATORY

700 LANDWEHR ROAD

NORTHBROOK, ILLINOIS 60062-2310

(312) 564-0700 FAX (312) 56: 9.17

NORTHERN STATES POWER COMPANY MINNEAPOLIS, MINNESOTA

PRAIRIE ISLAND NUCLEAR GENERATING PLANT Docket No. 50-282 License No. DPR-42 50-306 DPR-60

ANNUAL REPORT to the UNITED STATES NUCLEAR REGULATORY COMMISSION

Radiation Environmental Monitoring Program January 1, 1988 to December 31, 1988

Prepared Under Contract
by
TELEDYNE ISOTOPES MIDWEST LABORATORY
Project No. 8010

Approved by:

L. G. Huebner General Manager

PREFACE

The staff of Teledyne Isotopes Midwest Laboratory was responsible for the acquisition of data presented in this report. Samples were collected by members of the staff of the Environmental & Regulatory Activities Department.

The report was prepared by L. G. Huebner, General Manager, Teledyne Isotopes Midwest Laboratory. He was assisted in the report preparation by other staff members of this laboratory.

TABLE OF CONTENTS

Section		Page
	Preface	ii
	List of Tables	iv
1.0	INTRODUCTION	1
2.0	SUMMARY	2
3.0	RADIATION ENVIRONMENTAL MONITORING PROGRAM (REMP)	3
4.0 5.0 6.0	3.1 Program Design and Data Interpretation 3.2 Program Description	3 4 5 6 6 6 8 8 8 12 24
APPENDICE		
Α	Interlaboratory Comparison Program Results	A-1
В	Data Reporting Conventions	B-1
С	Maximum Permissible Concentrations of Radio- activity in Air and Water Above Natural Background in Unrestricted Areas	C-1

LIST OF TABLES

No.	<u>Title</u>			Page
5.1	Sample Collection and Analysis Program, 1988			13
5.2	Sampling Locations			15
5.3	Missed Collections and Analyses, 1988			17
5.4	Environmental Radiological Monitoring Program Summary			18
In addition	on, the following tables are in the Appendices:			
Appendix A				
A-1	Interlaboratory Comparison Program Results, 1984-1988			A-3
A-2	Interlaboratory Comparison Program Results (TLDs)			A-17
A-3	In-house Spiked Samples			A-20
A-3	In-house "Blank" Samples			A-24
	Attachment B: Acceptance criteria for spiked samples			A-28
	Addendum to Appendix A: Explanation of the Results Outside of Control Limits			A-29
Appendix (
C-1	Maximum Permissible Concentrations of Radioactivity in Air and Water Above Natural Background in Unrestricted Areas			C-2

1.0 INTRODUCTION

This report summarizes and interprets results of the Radiation Environmental Monitoring Program (REMP) conducted by Teledyne Isotopes Midwest Laboratory at the Prairie Island Nuclear Generating Plant, Red Wing, Minnesota, during the period January - December, 1988. This program monitors the levels of radioactivity in the air, terrestrial, and aquatic environments in order to assess the impact of the plant on its surroundings.

Tabulations of the individual analyses made during the year are not included in this report. These data are included in a reference document (Teledyne Isotopes Midwest Laboratory, 1988) available at Northern States Power Company, Nuclear Generation Department.

Prairie Island Nuclear Generating Plant is located on the Mississippi River in Goodhue County, Minnesota, and operated by Northern States Power Company. The plant has two 550 MWe pressurized water reactors. Unit 1 achieved initial criticality on 1 December 1973. Commercial operation at full power began on 16 December 1973. Unit 2 achieved initial criticality on 17 December 1974. Commercial operation at full power began on 21 December 1974.

2.0 SUMMARY

The Radiation Environmental Monitoring Program (REMP) required by the U.S. Nuclear Regulatory Commission (NRC) Technical Specifications for the Prairie Island Nuclear Generating Plant is described. Results for 1988 are summarized and discussed.

Program findings show background levels of radioactivity in the environmental samples collected in the vicinity of the Prairie Island Nuclear Generating Plant. No effect on the environment due to the operation of the plant is indicated.

3.0 RADIATION ENVIRONMENTAL MONITORING PROGRAM (REMP)

3.1 Program Design and Data Interpretation

The purpose of the Radiation Environmental Monitoring Program (REMP) at the Prairie Island Nuclear Generating Plant is to assess the impact of the plant on its environment. For this purpose, samples are collected from the air, terrestrial, and aquatic environments and analyzed for radioactive content. In addition, ambient gamma radiation levels are monitored by thermoluminescent dosimeters (TLDs).

Sources of environmental radiation include the following:

- (1) Natural background radiation arising from cosmic rays and primordial radionuclides;
- (2) Fallout from atmospheric nuclear detonations;
- (3) Releases from nuclear power plants;
- (4) Industrial and medical radioactive waste; and
- (5) Fallout from nuclear accidents.

In interpreting the data, effects due to the plant must be distinguished from those due to other sources.

A major interpretive aid in assessment of these effects is the design of the monitoring program at the Prairie Island Plant which is based on the indicator-control concept. Most types of samples are collected both at indicator locations (nearby, downwind, or downstream) and at control locations (distant, upwind, or upstream). A plant effect would be indicated if the radiation level at an indicator location was significantly larger than that at the control location. The difference would have to be greater than could be accounted for by typical fluctuations in radiation levels arising from other sources.

An additional interpretive technique involves analyses for specific radionuclides present in the environmental samples collected from the plant site. The plant's monitoring program includes analyses for tritium and iodine-131. Most samples are also analyzed for gamma-emitting isotopes with results for the following groups quantified: zirconium-95,

cesium-137, cerium-144, beryllium-7, and potassium-40. The first three gamma-emitting isotopes were selected as radiological impact indicators because of the different characteristic proportions in which they appear in the fission product mix produced by a nuclear reactor and that produced by a nuclear detonation. Each of the three isotopes is produced in roughly equivalent amounts by a reactor: each constitutes about 10% of the total activity of fission products 10 days after reactor shutdown. On the other hand, 10 days after a nuclear explosion, the contributions of zirconium-95, cerium-144, and cesium-137 to the activity of the resulting debris are in the approximate ratio 4:1:0.03 (Eisenbud, 1963). Beryllium-7 is of cosmogenic origin and potassium-40 is a naturally-occurring isotope. They were chosen as calibration monitors and should not be considered radiological impact indicators.

The other group quantified consists of niobium-95, ruthenium-103, and -106, cesium-134, barium-lanthanum-140, and cerium-141. These isotopes are released in small quantities by nuclear power plants, but to date their major source of injection into the general environment has been atmospheric nuclear testing. Nuclides of the final group, manganese-54, iron-59, cobalt-58, and -60, and zinc-65, are activation products and arise from activation of corrosion products. They are typical components of a nuclear power plant's effluents, but are not produced in significant quantities by nuclear detonations.

Other means of distinguishing sources of environmental radiation can be employed in interpreting the data. Current radiation levels can be compared with previous levels, including those measured before the Plant became operational. Results of the plant's monitoring program can be related to those obtained in other parts of the world. Finally, results can be related to events known to cause elevated levels of radiation in the environment, e.g., atmospheric nuclear detonations.

3.2 Program Description

The sampling and analysis schedule for the environmental radiation monitoring program at Prairie Island is summarized in Table 5.1 and briefly reviewed below. Table 5.2 defines the sampling location codes used in Table 5.1 and specifies for each location its type (indicator or control) and its distance, direction, and sector relative to the reactor site. To assure that sampling is carried out in a reproducible manner, detailed sampling procedures have been prescribed (Teledyne Isotopes Midwest Laboratory, 1987).

To monitor the air environment, airborne particulates are collected on membrane filters by continuous pumping at five locations. Also, airborne iodine is collected by continuous pumping through charcoal filters at all of these locations. Filters are changed and counted weekly. Particulate filters are analyzed for gross beta activity and charcoal filters for

iodine-131. A monthly composite of all particulate filters is gammascanned on an HP Ge or Ge(Li) detector. One of the five locations is a control (P-1), and four are indicator (P-2, P-3, P-4, and P-6). One of the indicators (P-3) is located near the residence expected to be most susceptible to any atmospheric emissions from the plant (highest D/Q residence).

As a "Lessons Learned" commitment, ambient gamma radiation is monitored at thirty-two (32) locations, using CaSo4:Dy dosimeter with four sensitive areas at each location: ten (10) in an inner ring in the general area of the site boundary, fifteen (15) in the outer ring within 4-5 mile radius, six (6) at special interest locations and one control location, 11.1 miles distant from the plant. They are replaced and measured quarterly. Also, a complete emergency set of TLDs for all locations is placed in the field at the same time as regular sets. The emergency set is returned to TIML quarterly for annealing and repackaging.

Milk samples are collected monthly from five farms (four indicator and one control). If the milch animals are on pasture; the milk is collected biweekly during the growing season (May - November). All samples are analyzed for iodine-131 and gamma-emitting isotopes.

For additional monitoring of the terrestrial environment, leafy green vegetables (cabbage) are collected annually from the highest D/Q garden and a control location (P-25) and analyzed for iodine-131. Corn is collected annually only from fields irrigated with river water and a control location (P-25) and analyzed for gamma-emitting isotopes. Also, well water is collected quarterly from four locations and analyzed for tritium and gamma-emitting isotopes.

River water is collected weekly at two locations, one upstream of the plant (P-5) and one downstream (P-6, Lock and Dam No. 3). Monthly composites are analyzed for gamma-emitting isotopes. Quarterly composites are analyzed for tritium.

Drinking water is collected weekly from the City of Red Wing well. Monthly composites are analyzed for gross beta, iodine-131, and gamma-emitting isotopes. Quarterly composites are analyzed for tritium.

The aquatic environment is also monitored by semi-annual upstream and downstream collections of fish, periphyton or invertebrates, and bottom sediments. Shoreline sediment is collected semi-annually from one location. All samples are analyzed for gamma-emitting isotopes.

3.3 Program Execution

The Program was executed as described in the preceding section with the following exceptions:

1. The TLD data for the first quarter of 1988 for Location P-05S was not available because TLD was lost in the field.

Deviations from the program are summarized in Table 5.3.

3.4 Laboratory Procedures

All iodine-131 analyses in milk and drinking water were made by using a sensitive radiochemical procedure which involves separation of the element by use of an ion-exchange resin and subsequent beta counting.

All gamma-spectroscopic analyses were performed with an HP Ge or Ge(Li) detector. Levels of iodine-131 in cabbage were determined by HP Ge or Ge(Li) spectrometry. Levels of airborne iodine-131 in charcoal samples were measured by HP Ge or Ge(Li) spectrometry.

Tritium levels were determined by liquid scintillation technique.

Analytical procedures used by the Teledyne Isotopes Midwest Laboratory are specified in detail elsewhere (Teledyne Isotopes Midwest Laboratory, 1985). Procedures are based on those prescribed by the National Center for Radiological Health of the U. S. Public Health Service (U. S. Public Health Service, 1967) and by the Health and Safety Laboratory of the U. S. Atomic Energy Commission, 1972).

Teledyne Isotopes Midwest Laboratory has a comprehensive quality control/quality assurance program designed to assure the reliability of data obtained. Details of TIML's Quality Assurance Program are presented elsewhere (Teledyne Isotopes Midwest Laboratory, 1988). The TIML Quality Assurance Program includes participation in Interlaboratory Comparison (Crosscheck) Programs. Results obtained in crosscheck programs are presented in Appendix A.

3.5 Program Modifications

During the growing season, milk producers were monitored to determine when the dairy animals were on pasture or fresh cut feed. The frequency of milk samples was increased to semimonthly at one indicator location (Gustafson) and one control location (Kinneman) during the period because animals were on pasture or fresh cut feed.

3.6 Land Use Census

In accordance with Technical Specification 4.10, paragraph B1, a land use census is conducted in order to identify the location of the nearest milk

animal, the nearest residence, and the nearest garden of greater than 500 ft² producing fresh leafy vegetables in each of the 16 meteorological sectors within a distance of 5 miles. This census is conducted at least once per 12 months between the dates of May 1 and October 31. New locations are added to the radiological environmental monitoring program within 30 days, and sampling locations having lower calculated doses or a lower dose commitment may be deleted from this monitoring program after October 31 of the year in which the land use census was conducted.

This land use census insures the updating of the radiation environmental monitoring program should sampling locations change within the 5 mile radius from the plant.

The 1988 Land Use Census was completed on July 8, 1988. This census did not identify any locations of exposure pathways different from those used in the program during the first six months of the year. Milk and garden sample locations did not change due to the requirements of the land use census.

4.0 RESULTS AND DISCUSSION

All of the scheduled collections and analyses were made except those listed in Table 5.3.

All results are summarized in Table 5.4 in a format recommended by the Nuclear Regulatory Commission in Regulatory Guide 4.8. For each type of analysis of each sampled medium, this table lists the mean and range for all indicator locations and for all control locations. The locations with the highest mean and range are also shown.

4.1 Atmospheric Nuclear Detonations and Nuclear Accidents

There were no reported atmospheric nuclear tests in 1988. The last reported test was conducted on October 16, 1980 by the People's Republic of China. The reported yield was in the 200 kiloton to 1 megaton range.

There were no reported accidents at nuclear reactor facilities in 1988.

4.2 Program Findings

Results obtained show background levels of radioactivity in the environmental samples collected in the vicinity of the Prairie Island Nuclear Generating Plant in 1988.

Ambient Radiation (TLDs)

Ambient radiation was measured in the general area of site boundary, at outer ring 4 - 5 mi distant from the Plant, at special interest areas, and at one control location. The means ranged from 14.9 mR/91 days at inner ring locations to 15.8 mR/91 days at outer ring locations. The mean at special locations was 14.4 mR/91 days and 16.0 mR/91 days at the control location. The differences are not statistically significant. The dose rates measured at all indicator and control locations were similar to those observed in 1978 (12.1 and 15.1 mR/91 days, respectively); in 1979 (12.6 and 15.3 mR/91 days, respectively); in 1980 (11.2 and 13.5 mR/91 days, respectively); in 1982 (12.0 and 13.0 mR/91 days, respectively); in

1983 (13.0 and 14.9 mR/91 days, respectively); in 1984 (13.9 and 15.3 mR/91 days, respectively); in 1985 (13.9 and 15.3 mR/91 days, respectively); in 1986 (16.6 and 17.0 mR/91 days, respectively) and in 1987 (15.4 and 16.0 mR/91 days, respectively). No plant effect on ambient gamma radiation was indicated.

Airborne Particulates

The average annual gross beta concentration in airborne particulates was identical at both indicator and control locations (0.030 pCi/m³) and was slightly higher than the levels observed in 1982 (0.026 pCi/m³), 1983 (0.023 pCi/m³), 1984 (0.024 pCi/m³), 1985 (0.025 pCi/m³), 1986 (0.025 pCi/m³), and 1987 (0.024 pCi/m³). The average of 0.025 pCi/m³ for 1986 does not include the results from May 19 to June 9, 1986, which were influenced by the accident at Chernobyl.

A spring peak in beta activity had been observed almost annually for many years (Wilson et al., 1969). It had been attributed to fallout of nuclides from the stratosphere (Gold et al., 1964). It was pronounced in 1981, occurred to a lesser degree in 1982, and did not occur in 1983, 1984, 1985, 1987 or 1988. In 1986, the spring peak could not be identified because it was overshadowed by the releases of radioactivity from Chernobyl. The highest averages for gross beta were for the month of December and the fourth quarter, as in 1983, 1984, 1985, 1986 (exclusive of the period between May 19, 1986 and June 9, 1986) and 1987.

Two pieces of evidence indicate conclusively that the elevated activity observed during the fourth quarter was not attributable to the Plant operation. In the first place, elevated activity of similar size occurred simultaneously at both indicator and control locations. Secondly, an identical pattern was observed at the Monticello Nuclear Generating Plant, about 100 miles distant from the Prairie Island Nuclear Generating Plant (Northern States Power Company, 1988).

Gamma spectroscopic analysis of quarterly composites of air particulate filters yielded similar results for inc. ator and control locations. Beryllium-7, which is produced continously in the upper atmosphere by cosmic radiation (Arnold and Al-Salih, 1955), was detected in all samples. All other gamma-emitting isotopes were below their respective LLD limits.

Airborne Iodine

Weekly levels of airborne iodine-131 were below the lower limit of detection (LLD) of 0.07 pCi/ m^3 in all samples.

Milk

Iodine-131 results were below the detection limit of 1.0 pCi/l in all samples.

Cs-137 results were below the LLD level of 15 pCi/l in all samples. No other gamma-emitting isotopes, except potassium-40, were detected in any milk samples. This is consistent with the finding of the National Center for Radiological Health that most radiocontaminants in feed do not find their way into milk due to the selective metabolism of the cow. The common exceptions are radioisotopes of potassium, cesium, strontium, barium, and iodine (National Center for Radiological Health, 1968).

In summary, the milk data for the 1988 show no radiological effects of the plant operation.

Drinking Water

In drinking water from the City of Red Wing well, tritium activity was below the LLD level of 330 pCi/l in all samples. Iodine-131 activity was also below the LLD level at 1.0 pCi/l in all samples. As with the other well water samples, all analyses for gamma-emitting isotopes yielded results below detection limits. Gross beta averaged 8.0 pCi/l and was similar to the levels observed in 1979 (10.5 pCi/l), 1980 (11.8 pCi/l), 1981 (10.7 pCi/l), 1982 (8.9 pCi/l), 1983 (8.0 pCi/l), 1984 (7.9 pCi/l), 1985 (7.1 pCi/l), 1986 (6.8 pCi/l) and 1987 (7.9 pCi/l).

River Water

At the upstream and downstream collection sites, quarterly composite tritium levels were below the LLD level of 330 pCi/l in all samples.

River water was also analyzed for gamma-emitting isotopes. All gamma-emitting isotopes were below their respective detection limits. There was no indication of a plant effect.

Well Water

At the control well P-25, Kinneman Farm and three indicator wells (P-8, Community Center; P-10, Lock and Dam No. 3; and P-9, Plant Well No. 2) no tritium was detected above LLD level of 330 pCi/l in all samples.

Gamma-emitting isotopes were below the detection limits in all samples.

Crops

Two samples of cabbage were collected in August and analyzed for I-131. The I-131 level was below $0.034~\rm pCi/g$ wet weight in both samples. There was no indication of a plant effect.

Fish

Fish samples were collected in June, September, and October, 1988. The only isotope detected was naturally-occuring potassium-40 and there was no significant difference between upstream and downstream results. There was no indication of a plant effect.

Aquatic Insects or Periphyton

Aquatic insects (invertebrates) or periphyton were collected in June, May, and September, 1986. The samples were analyzed for gamma-emitting isotopes. All gamma-emitting isotopes were below their respective LLDs. No plant effect was indicated.

Bottom and Shoreline Sediments

Sediment collections were made in April and September, 1988. The samples were analyzed for gamma-emitting isotopes.

Cs-137 was detected in two bottom sediment upstream samples and averaged 0.083 pCi/g dry weight.

All other gamma-emitting isotopes, except naturally-occurring potassium-40, were below their respective LLDs. No plant effect was indicated. 5.0 TABLES

Table 5.1 Sample collection and analysis program, 1988.

Prairie Island

		Locations	Collection Type and	Analysis Type and
Medium	No.	Codes (and Type)a	Frequencyb	Frequency ^C
Ambient radiation (TLDs)	35	P-01A - P-10A P-01B - P-15B P-01S - P-06S P-01C	0/0	Ambient gamma
Airborne particulates	ro.	P-1(C), P-2, P-3, P-4, P-6	C/W	CS, GS (QC of each location)
Airborne iodine	r.	P-1(C), P-2, P-3 P-4, P-6	C/W	1-131
Milk	ro.	P-16 to P-18, P-25(C), P-14	p'4/9	I-131, GS
River water	2	P-5(C), P-6	M/9	GS(MC), H-3(QC)
Drinking water	1	p-11	M/9	GB(MC), I-131(MC) GS(MC), H-3(QC)
Well water	4	P-25(C), P-6, P-8, P-9	0/9	Н-3, 65
Edible cultivated crops - leafy green vegetables	2	P-25(C), P-24	G/A	I-131

Sample collection and analysis program, 1988 (continued) Table 5.1.

מש	3
C	5
U	5
a	
7	
23	5
à	

Type)a Frequencyb -20			Locations	Collection Type and	Analysis Type and
2 P-25(C), P-20 G/A 2 P-5(C), P-6 G/SA 2 P-5(C), P-6 G/SA 2 P-5(C), P-6 G/SA 4 1 P-12 G/SA	Medium	No.	Codes (and Type)a	Frequencyb	Frequency ^C
2 P-5(C), P-6 G/SA 2 P-5(C), P-6 G/SA 2 P-5(C), P-6 G/SA t 1 P-12 G/SA	Edible cultivated crops - corn	2	P-25(C), P-20	G/A	99
2 P-5(C), P-6 G/SA 2 P-5(C), P-6 G/SA ent 1 P-12 G/SA	Fish (one species edible portion)	2	P-5(C), P-6	6/SA	98
2 P-5(C), P-6 G/SA ent 1 P-12 G/SA	Periphyton or invertebrates	2	P-5(C), P-6	6/SA	9
1 P-12 G/SA	Bottom sediment	2	P-5(C), P-6	G/SA	99
	Shoreline sediment	-	P-12	6/SA	65

a Location codes are defined in Table 5.2. Control stations are indicated by (C). All other stations are indicators. 0

as follows: W = weekly, M = monthly, Q = quarterly, SA = semi-annually, A = annually. Analysis type is coded as follows: GB = gross beta, GS = gamma spectroscopy, H-3 = tritium, I-131 Collection type is coded as follows: C/ = continuous, G/ = grab. Collection frequency is coded

iodine 131. Analysis frequency is coded as follows: MC = monthly composite, QC = quarterly composite. Milk is collected biweekly during the grazing season (May - November) if milch animals are on pasture. 0

Table 5.2. Sampling locations.

Code	Typea	Name	Loc	cation
P-1	С	Air Station P-1	16.5 mi	@ 348°/NNW
P-2		Air Station P-2		@ 294°/WNW
P-3		Air Station P-3	0.8 mi	@ 313°/NW
P-4		Air Station P-4		@ 359°/N
P-5	C	Upstream of Plant		@ 60°/ENE
P-6		Lock & Dam #3 & Air Station P-6		@ 129°/SE
P-8		Community Center		@ 304°/NW
P-9		Plant Well #2	0.3 mi	@ 306°/NW
P-11		City of Red Wing		@ 135°/SE
P-12		Recreational Area	3.4 mi	@ 116°/ESE
P-14		Gustafson Farm		@ 168°/SSE
P-16		Johnson Farm		@ 60°/ENE
P-17		Place Farm		@ 25°/NNE
P-18		Christensen Farm	3.7 mi	@ 88°/E
P-20		River Irrigated Corn Field*		
P-24		Highest D/Q Garden**		
P-25	C	Kinneman Farm	11.1 mi	@ 331°/NNW
P-01A		Property Line		@ 359°/N
P-02A		Property Line	0.3 mi	@ 19°/NNE
P-03A		Property Line	0.5 mi	@ 183°/S
P-04A		Property Line		@ 204°/SSW
P-05A		Property Line		@ 225°/SW
P-06A		Property Line	0.4 mi	@ 249°/WSW
P-07A		Property Line	0.4 mi	@ 268°/W
P-08A		Property Line	0.4 mi	@ 291°/NNV
P-09A		Property Line		@ 317°/NW
P-10A		Property Line		@ 333°/NNW
P-01B		Thomas Killian Residence		@ 355°/N
P-02B		Roy Kinneman Farm		@ 17°/NNE
P-03B		Wayne Anderson Farm		@ 46°/NE
P-04B		Nelson Drive (Road)	4.2 mi	@ 61°/ENE
P-05B		County Road E and Coulee		@ 97°/E
P-06B		William Houschildt Residence		@ 112°/ESE
P-07B		Red Wing Service Center		@ 140°/SE
P-08B		David Wnuk Residence		@ 165°/SSE
P-09B		Highway 19 South		@ 187°/S
P-10B		Cannondale Farm	4 9 mi	@ 200°/SSW

a "C" denotes control location. All other locations are indicators.

^{*} Collected only if river water is used to irrigate the cornfields (Technical Specification Revision No. 80, effective 11-14-86).


^{**} This location is not determined until after the Land Use Census is completed.


Table 5.2. Sampling locations (continued)

Prairie Island

Code Type	Name	Location
P-11B	Wallace Weberg Farm	4.5 mi @ 221°/SW
P-12B	Roy Gergen Farm	4.5 mi @ 247°/WSW
P-13B	Thomas O'Rourke Farm	4.4 mi @ 270°/W
P-14B	David J. Anderson Farm	4.9 mi @ 306°/NW
P-15B	Holst Farms	4.2 mi @ 347°/NNW
P-01S	Federal Lock & Dam #3	1.6 mi @ 129°/SE
P-02S	Charles Suter Residence	0.6 mi @ 158°/SSE
P-03S	Carl Gustafson Farm	2.2 mi @ 168°/SSE
P-04S	Richard Burt Residence	2.0 mi @ 228°/SW
P-05S	Kenney Store	2.0 mi @ 270°/W
P-06S	Earl Flynn Farm	2.5 mi @ 299°/WNW
P-01C	Robert Kinnemen Farm	11.1 mi @ 331°/NNW

a "C" denotes control location. All other locations are indicators.





Mote: Map numbers correspond to code numbers without the prefix P -




Table 5.3. Missed collections and analyses, 1988. Prairie Island Nuclear Generating Plant. All required samples were collected and analyzed as scheduled except the following.

Sample	Analysis	Location	Collection Date or Period	Comments
Thermoluminescent Dosimeters (TLDs)	Ambient Radiation	P-05S	1st Qtr. 1988	Lost in the field.

Table 5.4. Environmental Radiological Monitoring Program Summary.

Goodhue, Minnesota Reporting Period January - Decembe

Number of	Non-routine Resultse	0	9	5	0	0			0	0	0	0	0	0	0	0	0
Control	Mean (F) Range	(See control below)	(See control below)	(See control below)	16.0 (4/4)	0.030 (53/53) (0.014-0.058)			0.098 (4/4)	4LD	4LD	dLD.	4LLD	4FFD	dL0	4LD	44.0
Highest	Mean (F) Range	17.0 (4/4) (16.0-18.3)	(16.0-19.9)	16.5 (4/4)	16.0 (4/4)	0.031 (53/53) (0.014-0.085) 0.031 (52/52)	(0.008-0.085)		0.098 (4/4)	,			1	1	,	1	1
Location with Highest Annual Mean	Locationd	P-03A, Property Line 0.5 mi@ 183°/S	P-02B, Roy Kinneman Farm, 4.8 mi 0 17°/NNE	P-03S, C. Gustafson Farm, 2.2 mi @ 168°/SSE	P-01C, R. Kinnenan Farm, 11.1 mi 6 331°/NNM	P-2, Station P-2 P-6, Station P-6	1.6 mt @ 129°/NNW		P-1, Station P-1 0.5 @ 294°/WNW	,	,	,	1	,		,	1
Indicator	Mean (F)C Range	14.9 (40/45) (11.3-19.1)	15.8 (60/60) (12.1-19.9)	14.4 (23/23) (10.8-19.4)	None	0.030 (211/212)			0.086 (16/16)	<pre>cffp</pre>	410	<pre></pre>	<lld< td=""><td>4LD</td><td>4FFD</td><td>dll></td><td>4LD</td></lld<>	4LD	4FFD	dll>	4LD
	ęáni	3.0	3.0	3.0	3.0	0.002			0.022	0.0022	0.0028	0.0017	0.0038	0.0035	0.0025	0.016	0.0017
Type and	Number of Analyses ^a	Gamma 40	Сатта 60	Gamma 23	балта 4	68 265		65 20	Be-7	Mn-54	Co-58	09-03	Zn-65	Zr-Nb-95	Ru-103	Ru-106	Cs-134
Sample	Type (Units)	TLD (mR/91 days) (Inner Ring, General Area at Site Roundary)	TLD (mR/91 days) (Outer Ring, 4-5 miles distant)	TLD (mR/91 days) (Special Interest Areas)	TLD (mR/91 days) (control)	Airborne Particulates (pCi/m3)											

Table 5.4. Environmental Radiological Monitoring Program Summary (continued)

50-282, 50-306	lanuary - Decembe	
Docket No.	Reporting Feriod	
Name of Facility Prairie Island Nuclear Generating Plant Docket	Location of Facility Goodhue, Minnesota	(County, State)

Sample	Type and		Indicator	Location with Highest Annual Mean	Highest	Control	Number of
Type (Units)	Number of Analyses ^a	11.0b	Mean (F)C Range ^C	Locationd	Nean (F) Range	Mean (F) Range	Non-routine Resultse
Airborne	Cs-137	0.0019	<pre><pre></pre></pre>	,		4LD	0
(pCi/m³)	Ba-La-140	0.0079	orro	,	,	حرر ق	0
(continued)	Ce-141	0.0047	of LD	,	1	dL10	0
	Ce-144	0.011	dlp.		1	dLL0	0
Airborne Iodine (pCi/m3)	1-131 265	0.37	410		,	4TD	0
Milk	1-131 74	1.0f	41.0	,		4LL0	0
there is	65 74						
	K-40	100	1380 (5?/57) (1120-1570)	P-14, Gustafson Farm 2.2 mi @ 168º/SSE	1450 (17/17) (1350-1570)	1360 (17/17) (1210-1520)	0
	Cs-134	15	977>			440	0
	Cs-137	15	410			<pre><fre></fre></pre>	0
	Ba-La-140	15	0715		,	4LD	0
Drinking Water (pci/1)	68 12	1.0	8.0 (12/12) (6.4-9.9)	P-11, City of Red Wing, 7.1 mi @ 135°/SE	8.0 (12/12)	None	0
	1-131 12	1.0	QTD .		,	None	0
	H-3 4	330	dl.b	,	1	None	0
	65 12						
	Mn-54	15	41.0	,	,	None	0
	Fe-59	30	4LD		,	None	0
	60-58	15	4LD	,	,	None	0
	09-00	15	410	,	,	None	0
	Zn-65	30	dl.b	,	1	None	0
	Zr-Nb-95	15	dLLD.	,	,	None	0

Table 5.4. Environmental Radiological Monitoring Program Susmary (continued)

Locat	Location of Facility	905	y Goodhue, Minnesota	Reporting Period	January	anuary - December 1903	ı
			(County, State)				1
Sample	Type and		Indicator	Location with Highest Annual Mean	Highest	Control	Number of
Type (Units)	Number of Analyses ³	q077	Mean (F)C RangeC	Locationd	Mean (F) Range	Mean (F) Range	Non-routine Resultse
Drinking Water	Cs-13%	10	41.0	3		None	0
(continued)	Cs-137	10	dLD.		,	None	0
	Ba-La-140	1.5	44.0	,	,	None	0
	Ce-144	46	<pre></pre>	,	,	None	ō
River Water	H-3 8	330	410	ı		<pre></pre>	0
(pt./1)	65 24						
	Mn-54	15	410	1		<pre><fre></fre></pre>	0
	Fe-59	30	<pre><pre></pre></pre>		í	<pre><pre></pre></pre>	0
	Co-58	15	4LLD		4	<pre></pre>	0
	09-03	15	dLD.		,	<pre></pre>	0
	Zn-65	30	<pre></pre>		,	d110	0
	Cs-134	15	4110	,	,	<pre></pre>	0
	Cs-137	18	410	,	,	4770	0
	Ba-La-140	15	410	,	,	CLL5	0
	Ce-144	79	<lld< td=""><td>,</td><td>,</td><td>dLD</td><td>0</td></lld<>	,	,	dLD	0
Well Water	н-3 16	330	410	1		dll)	0
(pc1/1)	65 16						
	Mn-54	15	<lld< td=""><td>,</td><td></td><td>41.0</td><td>0</td></lld<>	,		41.0	0
	Fe-59	30	410		,	410	0
	35-03	15	<pre><pre></pre></pre>		,	QTD	0
	09-03	1,5	4LD	1	1	410	0
	Zn-65	30	410		1	410	0
	Zr-Nb-95	15	410	,	,	dL0	0
	Cs-134	10	410	1	1	4LLB	0

Table 5.4. Environmental Radiological Monitoring Program Summary (continued)

Local	Location of Facility	500	Goodhue, Minnesota	Reporting Period		January - December 1988	
			(County, State)				
Sample Type (Units)	Type and Number of Analysesa	q077	Indicator Locations Mean (F)C RangeC	Location with Highest Annual Mean Locationd	Highest ean Mean (F) Range	Control Locations Mean (F) Range	Number of Non-routin Results
Well Water	09-03	15	dLL0		1	QTT>	0
(continued)	Zn-65	30	<pre><fre></fre></pre>	,	,	410	0
	Zr-Nb-95	15	«TED	,	,	<pre></pre>	0
	Cs-134	10	on.		,	<pre><pre></pre></pre>	0
	Cs-137	10	crrp	,	,	4110	0
	Ba-La-140	15	«TTD		,	dLL0	0
	Ce-144	81	4110		,	<pre></pre>	0
Crops-Cabbage (pCi/g wet)	1-131 2	0.034	410	,	,	4110	0
Fish - Flesh	65 4						
(pCirg wet)	K-40	5,7	2.82 (2/2)	P-5(C), Upstream of Plant, 0.6 m ² 8	3.03 (2/2) (2.68-3.36)	3.03 (2/2) (2.68-3.38)	0
	Mn-54	0.040	4110	,	,	4LLD	0
	Fe-59	0.076	du.		1	410	0
	6-58	0.084	410	1		41.0	0
	09-03	0.037	410	7		quo	0
	Zn-65	0.11	dLD	,	,	4LLD	0
	Zr-Nb-95	0.074	410	,	1	410	0
	Cs-134	0.033	dLD	1	1	<pre></pre>	0
	Cs-137	0.042	413	1	1	4LLD	9
	Ba-La-140	0.11	4.00			4770	0

Table 5.4. Environmental Radiological Monitoring Program Summary (continued)

Loca	Location of Facility	60	Goodhue, Minnesota	Reporting Period	7	anuary - December 1988	
			(County, State)				1
Sample	Type and		Indicator	Location with Highest Annual Mean	Highest	Control	Number o
Type (Units)	Number of Analyses ^a	11.0b	Mean (F)C RangeC	Locationd	Mean (F) Range	Mean (F) Range	Non-routi Results
Invertebrates	53						
(pt1/g wet)	Be-7	3.03	4LD	,	,	4LLD	0
	K-40	7.46	d1.D		,	4LLD	0
	Mn-54	0.29	CLID	,	,	41.0	0
	69-03	0.37	d'us	,	•	4LD	0
	09-03	9.26	077>		,	4LD	0
	Zu-65	0.71	41.0		,	<	0
	Zr-Mb-95	0.54	CLLD.	,	1	4LLD	0
	Ru-103	0.43	410	,	,	4LL	0
	Ru-106	2.56	CLLD.	,		QTT>	0
	Cs-134	0.29	410	,	,	<pre><pre></pre></pre>	0
	Cs-137	0.29	d1.0		ı	<pre></pre>	0
	Ba-La-140	1.52	410			<pre></pre>	0
	Ce-141	69.0	4110	,	,	4LLD	0
	Ce-144	1.58	QLD.			4LLD	0
Sottom and	9 59						
Sediments (pCi/g dry)	Be-7	0.46	4110	P-5(C), Upstream of Plant, 0.6 mi @ 60°/ENE	0.49 (1/2)	0.49 (1/2)	0
	K-40	1.0	8.52 (4/4) (7.55-10.10)	P-5(C), Upstream of Plant, 0.6 mi 0 60°/ENE	9.26 (2/2) (9.24-9.27)	9.26 (2/2)	0
	Mn-54	0.036	4LLD		,	<pre></pre>	0
	Co-58	0.051	4110	,	•	<lld< td=""><td>0</td></lld<>	0

Table 5.4. Environmental Radiological Monitoring Program Summary (continued)

		Number of	Non-routine Resultse	0	0	0	0	0	0	0	0
50-282, 50-306	January - December 1988	Control	Mean (F) Range	<pre></pre>	4LD	<pre></pre>	dTD <	לרום	dL0	0.083 (2/2)	4LLD
50-282,		Highest	Mean (F) Range	1	,	,		1		0.083 (2/2)	
Plant Docket No.	Reporting Period	Location with Highest Annual Mean	Locationd	ě	,	1	,	1	,	P-5(C), Upstream of Plant, 0.6 mi @ 60°/ENE	,
Prairie Island Muclear Jenerating Plant	Location of Facility Goodhue, Minnesota (County, State)	Indicator	Mean (F)C RangeC	<ll0< td=""><td>CLLD.</td><td>4LLD</td><td>4110</td><td><lld< td=""><td><pre></pre></td><td>- LLD</td><td>- CLED</td></lld<></td></ll0<>	CLLD.	4LLD	4110	<lld< td=""><td><pre></pre></td><td>- LLD</td><td>- CLED</td></lld<>	<pre></pre>	- LLD	- CLED
Name of Facility Prairie Island		quTT		0.043	0.087	0.059	690.0	0.25	0.031	0.031	0,73
		Type and	Type and Number of Analysesa		.u-65	Zr-Nb-95	Ru-103	Ru-106	Cs-134	Cs-137	Ba-La-140
Name o		Sample	Type (Units)	Bottom and	Sediments	(continued)					

GB = Gross beta; GS = gamma scan.
LLD = Hominal lower limit of detection based on 4.66 sigma error for background sample.
Mean and range based upon detectable measurements only. Fraction of detectable measurements at specified location is indicated in parentheses (F).

0 0

410

1

4LLD

0,16 0.18

Ce-141 Ce-144

Locations are specified (1) by name and code (Table 2) and (2) distance, direction, and sector relative to reactor site.

Non-routine results are those which exceed ten times the control station value. If no control station value is available, the result is considered non-routine if it exceeds ten times the preoperational value for the location.

Iwo results (<1.1 and <1.2) were excluded from the determination of LLD for I-13% in milk. Elevated LLDs resulted from delay in analyses due to relocation of laboratory.

6.0 REFERENCES CITED

- Arnold, J. R. and H. A. Al-Salih. 1955. Beryllium-7 Produced by Cosmic Rays. Science 121: 451-453.
- Eisenbud, M. 1963. Environmental Radioactivity, McGraw-Hill, New York, New York, pp. 213, 275 and 276.
- Gold, S., H. W. Barkhau, B. Shlein, and B. Kahn, 1964. Measurement of Naturally Occuring Radionuclides in Air, in the Natural Radiation Environment, University of Chicago Press, Chicago, Illinois, 369-382.
- Hazleton Environmental Sciences Corporation. 1979a. Radiation Environmental Monitoring for Prairie Island Nuclear Generating Plant, Complete Analysis Data Tables, January December 1978.
- Nuclear Generating Plant, Complete Analysis Data Tables, January December 1978.
- Nuclear Generating Plant, Complete Analysis Data Tables, January December 1979.
- . 1980b. Radiation Environmental Monitoring for Prairie Island Nuclear Generating Plant, Complete Analysis Data Tables, January December 1979.
- Nuclear Generating Plant, Complete Analysis Data Tables, January December 1980.
- Nuclear Generating Plant, Complete Analysis Data Tables, January December 1980.
- Nuclear Generating Plant, Complete Analysis Data Tables, January December 1981.
- . 1982b. Radiation Environmental Monitoring for Prairie Island Nuclear Generating Plant, Complete Analysis Data Tables, January -December 1981.
- Nuclear Generating Plant, Complete Analysis Data Tables, January December 1982.

- Hohenemser, C.M. Deicher, A. Ernst, H. Hofsass, G. Lindner, E. Racknagel. 1986. "Chernobyl," Chemtech, October 1986, pp. 596-605.
- National Center for Radiological Health, 1968. Radiological Health and Data Reports, Vol. 9, Number 12, 730-746.
- Northern States Power Company. 1979. Prairie Island Nuclear Generating Plant, Annual Radiation Environmental Monitoring Report to the U.S. Nuclear Regulatory Commission, January 1, 1978 to December 31, 1978 (prepared by Hazleton Environmental Sciences). Minneapolis, Minnesota.
- Radiation Environmental Monitoring Report to the U.S. Nuclear Regulatory Commission, January 1, 1979 to December 31, 1979 (prepared by Hazleton Environmental Sciences). Minneapolis, Minnesota.
- Radiation Environmental Monitoring Report to the U.S. Nuclear Regulatory Commission, January 1, 1980 to December 31, 1980 (prepared by Hazleton Environmental Sciences). Minneapolis, Minnesota.
- Radiation Environmental Monitoring Report to the U.S. Nuclear Regulatory Commission, January 1, 1981 to December 31, 1981 (prepared by Hazleton Environmental Sciences). Minneapolis, Minnesota.
- Radiation Environmental Monitoring Report to the U.S. Muclear Regulatory Commission, January 1, 1982 to December 31, 1982 (prepared by Hazleton Environmental Sciences). Minneapolis, Minnesota.
- Radiation Environmental Monitoring Report to the U.S. Nuclear Regulatory Commission, January 1, 1983 to December 31, 1983 (prepared by Teledyne Isotopes Midwest Laboratory). Minneapolis, Minnesota.
- Radiation Environmental Monitoring Report to the U.S. Nuclear Regulatory Commission, January 1, 1984 to December 31, 1984 (prepared by Teledyne Isotopes Midwest Laboratory). Minneapolis, Minnesota.
- Radiation Environmental Monitoring Report to the U.S. Nuclear Regulatory Commission, January 1, 1985 to December 31, 1985 (prepared by Teledyne Isotopes Midwest Laboratory). Minneapolis, Minnesota.
- Radiation Environmental Monitoring Report to the U.S. Nuclear Regulatory Commission, January 1, 1986 to December 31, 1986 (prepared by Teledyne Isotopes Midwest Laboratory). Minneapolis, Minnesota.

1988. Prairie Island Nuclear Generating Plant, Annual Radiation Environmental Monitoring Report to the U.S. Nuclear Regulatory Commission, January 1, 1987 to December 31, 1987 (prepared by Teledyne Isotopes Midwest Laboratory). Minneapolis, Minnesota. Teledyne Isotopes Midwest Laboratory. 1971a. Quality Control Program, Revision 7. 20, November 1984. . 1985. Quality Control Procedures Manual, Revision 6. 09 August 1985. 1985. Quality Assurance Program Manual, Revision 1, 15 April 1985 . 1985. Analytical Procedures Manual, Revision 5, 29 July 1985. 1984a. Radiation Environmental Monitoring for Monticello Nuclear Generating Plant, Complete Analysis Data Tables, January -December 1983. . 1984b. Radiation Environmental Monitoring for Prairie Island Nuclear Generating Plant. Complete Analysis Data Tables, January -December 1983. 1985a. Radiation Environmental Monitoring for Monticello Nuclear Generating Plant, Complete Analyses Data Tables, January -December 1984. . 1985b. Radiation Environmental Monitoring for Prairie Island Nuclear Generating Plant, Complete Analyses Data Tables, January -December 1984. 1986a. Radiation Environmental Monitoring for Monticello Nuclear Generating Plant, Complete Analyses Data Tables, January -December 1985. . 1986b. Radiation Environmental Monitoring for Prairie Island Nuclear Generating Plant, Complete Analyses Data Tables, January -December 1985. . 1987. Sampling Procedures, Prairie Island Nuclear Generating Plant, Revision 16, 18 December 1987. 1987a. Radiation Environmental Monitoring for Monticello Nuclear Generating Plant, Complete Analyses Data Tables, January -December 1986. . 1987b. Radiation Environmental Monitoring for rairie Island Nuclear Generating Plant, Complete Analyses Data Tables, January -December 1986.

- Nuclear Generating Plant, Complete Analyses Data Tables, January December 1987.
- Nuclear Generating Plant, Complete Analyses Data Tables, January December 1987.
- Nuclear Generating Plant, Complete Analyses Data Tables, January December 1988.
- Nuclear Generating Plant, Complete Analyses Data Tables, January December 1988.
- U. S. Atomic Energy Commission. 1972. HASL Procedures Manual, Health and Safety Laboratory, New York, NY., 10014.
- U. S. Public Health Service. 1967. Radioassay Procedures for Environmental Samples, National Center for Radiological Health, Rockville, Maryland (Public Health Service Publication No. 999-RH-27).
- Wilson, D. W., G. M. Ward and J. E. Johnson. 1969. In Environmental Contamination by Radioactive Materials, International Atomic Energy Agency. p. 125.

Appendix A

Interlaboratory Comparison Program Results

NOTE: TIML participates in intercomparison studies administered by U.S. EPA Environmental Monitoring Systems Laboratory, Las Vegas, Nevada. The results are reported in Appendix A. Also reported are results of inhouse spikes and blanks. Appendix A is updated twice a year and the complete Appendix is included in January and July monthly reports only. Please refer to January and July Reports for information.

January, 1989

Appendix A

Interlaboratory Comparison Program Results

Teledyne Isotopes Midwest Laboratory (formerly Hazleton Environmental Sciences) has participated in interlaboratory comparison (crosscheck) programs since the formulation of its quality control program in December 1971. These programs are operated by agencies which supply environmental-type samples (e.g., milk or water) containing concentrations of radionuclides known to the issuing agency but not to participant laboratories. The purpose of such a program is to provide an independent check on the laboratory's analytical procedures and to alert it to any possible problems.

Participant laboratories measure the concentrations of specified radionuclides and report them to the issuing agency. Several months later, the agency reports the known values to the participant laboratories and specifies control limits. Results consistently higher or lower than the known values or outside the control limits indicate a need to check the instruments or procedures used.

The results in Table A-1 were obtained through participation in the environmental sample crosscheck program for milk, water, air filters, and food samples during the period May 1984 through November, 1988. This program has been conducted by the U.S. Environmental Protection Agency Intercomparison and Calibration Section, Quality Assurance Branch, Environmental Monitoring and Support Laboratory, Las Vegas, Nevada.

The results in Table A-2 were obtained for thermoluminescent dosimeters (TLD's) during the period 1976, 1977, 1979, 1980, 1984, and 1985-1986 through participation in the Second, Third, Fourth, Fifth, Seventh, and Eighth International Intercomparison of Environmental Dosimeters under the sponsorships listed in Table A-2.

Table A-3 lists results of the analyses on in-house spiked samples.

Table A-4 lists results of the analyses on in-house "blank" samples.

Attachment B lists acceptance criteria for "spiked" samples.

Addendum to Appendix A provides explanation for out of limit results.

Table A-1. U.S. Environmental Protection Agency's crosscheck program, comparison of EPA and Teledyne Isotopes Midwest Laboratory results for milk, water, air filters, and food samples, 1984 through 1988.

					Concentration in pCi/lb TIML Result EPA Resultd				
	Sample	Date			Military and a partners with the partners of t	North Control of the			
Code	Туре	Collected	Analysis	±2σ ^C	1s, N=1	Control Limits			
STW-358	Water	May 1984	Gr. alpha Gr. beta	3.0±0.6 6.7±1.2	3±5.0 6±5.0	0.0-11.7			
			di. Deta	0.711.2	0.010	0.0-14.7			
STM-366	Milk	June 1984	Sr-89	21±3.1	25±5.0	16.3-33.7			
			Sr-90	13±2.0	17±1.5	14.4-19.6			
			I-131		43±6.0	32.6-53.4			
			Cs-137	38±4.0	35±5.0	26.3-43.7			
			K	1577±172	1496±75	1336-1626			
STW-368	Water	July 1984	Gr. alpha	5.1±1.1	6±5.0	0.0-14.7			
			Gr. beta	11.9±2.4	13±5.0	4.3-21.7			
STW-369	Water	August 1984	I-131	34.3±5.0	34.0±6.0	23.6-44.4			
STW-370	Water	August 1984	H-3	3003±253	2817±356	2200-3434			
STF-371	Food	July 1984	Sr-89	22.0±5.3	25.0±5.0	14.3-33.7			
			Sr-50	14.7±3.1	20.0±1.5	17.4-22.6			
			I-131	<172	39.0±6.0	28.6-49.4			
			Cs-137	24.0±5.3	25.0±5.0	14.3-33.7			
			K	2503±132	2605±130	2379-2831			
TAF-372	Air	August 1984		15.3±1.2	17±5.0	8.3-25.7			
	Filter		Gr. beta	56.0±0.0	51±5.0	42.3-59.7			
			Sr-90	14.3±1.2	18±1.5	15.6-20.4			
			Cs-137	21.0±2.0	15±5.0	6.3-23.7			
STW-375	Water	Sept 1984	Ra-226	5.1±0.4	4.9±0.7	3.6-6.2			
			Ra-228	2.2±0.1	2.3±0.4	1.7-2.9			
STW-377	Water	Sept 1984	Gr. alpha	3.3±1.2	5.0±5.0	0.0-13.7			
			Gr. beta	12.7±2.3	16.0±5.0	7.3-24.7			
STW-379	Water	Oct 1984	H-3	2860±312	2810±205	2454-3166			
STW-380	Water	Oct 1984	Cr-51	<36	40±5.0	31.3-48.7			
			Co-60	20.3±1.2	20±5.0	11.3-28.7			
			Zn-65	150±8.1	147±5.0	138.3-155.7			
			Ru-106	<30	47±5.0	36.3-55.7			
			Cs-134	31.3±7.0	31±5.0	22.3-39.7			
			Cs-137	26.7±1.2	24±5.0	15.3-32.7			

Table A-1. (continued)

				PENERTY SACREMENTAL ARTEST CONTRACTOR TO A MANAGEMENT AND A STATE OF THE PENERTY	entration i	THE RESIDENCE OF THE PARTY OF T
Lab	Sample	Date		TIML Result	ATTENDED THE REAL PROPERTY OF THE PARTY OF T	
Code	Туре	Collected	Analysis	±2σC	1s, N=1	Control Limits
STM-382	Milk	Oct 1984	Sr-89	15.7±4.2	22±5.0	13.3-30.7
			Sr-90	12.7±1.2	16±1.5	13.4-18.6
			I-131	41.7±3.1	42±6.0	31.6-42.4
			Cs-137		32±5.0	23.3-40.7
			K	1447±66	1517±76	1386-1648
STW-384		Oct 1984	Gr. alpha			5.3-22.7
	(Blind)	Sample A				2.2-3.8
					2.1±0.3	1.6-2.6
			Uranium	NAe	5.0±6.0	0.0-15.4
		Sample B			64±5.0	55.3-72.7
			Sr-89	10.7±4.6	11±5.0	2.3-19.7
			Sr-90	7.3±1.2	12±1.5	9.4-14.6
			Co-60	16.3±1.2	14±5.0	5.3-22.7
			Cs-134	<2		0.0-10.7
			Cs-137	16.7±1.2	14±5.0	5.3-22.7
TAF-387		Nov 1984	Gr. alpha	18.7±1.2	15±5.0	6.3-23.7
	Filter		Gr. beta	59.0±5.3	52±5.0	43.3-60.7
			Sr-90	18.3±1.2		18.4-23.6
			Cs-137	10.3±1.2	10±5.0	1.3-18.7
STW-388	Water	Dec 1984	I-131	28.0±2.0	36±6.0	25.6-36.4
STW-389	Water	Dec 1984	H-3	3583±110	3182±360	2558-3806
STW-391	Water	Dec 1984	Ra-226	8.4±1.7	8.6±1.3	6.4-10.8
			Ra-228	3.1±0.2	4.1±0.6	3.0-5.2
STW-392	Water	Jan 1985	Sr-89			0.0-11.7
			Sr-90	27.3±5.2	30.0±1.5	27.4-32.6
STW-393	Water	Jan 1985	Gr. alpha	3.3±1.2	5±5.0	0.0-13.7
			Gr. beta	17.3±3.0	15±5.0	6.3-23.7
STF-395	Food	Jan 1985	Gr. alpha	4.7±2.3	6.0±5.0	0.0-14.7
			Gr. beta	11.3±1.2	15.0±5.0	6.3-23.7
			Sr-89	25.3±6.4	34.0±5.0	25.3-42.8
			Sr-90	11.3±1.2 25.3±6.4 27.0±8.8	26.0±1.5	23.4-28.6
			1-131	38. UIZ. U	35.010.0	24.0-45.4
			Cs-137	32.7±2.4		
			K	1410±212	1382±120	1174-1590

Table A-1. (continued)

				Concentration in pCi/1D			
Lab	Sample	Date		TIML Result	EPA Result ^d		
Code	Туре	Collected	Analysis	±2σ ^C	1s, N=1	Control Limits	
STW-397	Water	Feb 1985	Cr-51	<29	48±5.0	39.3-56.7	
			Co-60	21.3±3.0	20±5.0	11.3-28.7	
			Zn-65	53.7±5.0	55±5.0	46.3-63.7	
			Ru-106	<23	25±5.0	16.3-33.7	
			Cs-134	32.3±1.2	35±5.0	26.3-43.7	
			Cs-137	25.3±3.0	25±5.0	16.3-33.7	
STW-398	Water	Feb 1985	H-3	3869±319	3796±634	3162-4430	
STM-400	Milk	March 1985	I-131	7.3±2.4	9.0±0.9	7.4-10.6	
STW-402	Water	March 1985	Ra-226	4.6±0.6	5.0±0.8	3.7-6.3	
			Ra-228	<0.8	9.0±1.4	6.7-11.3	
		Reanalysis	Ra-228	9.0±0.4			
STW-404	Water	March 1985	Gr. alpha	4.7±2.3	6±5.0	0.0-14.7	
			Gr. beta	11.3±1.2	15±5.0	6.3-23.7	
STAF-405	Air	March 1985	Gr. alpha	9.3±1.0	10.0±5.0	1.3-18.7	
	Filter		Gr. beta	42.0±1.1	36.0±5.0	27.3-44.7	
			Sr-90	13.3±1.0	15.0±1.5	12.4.17.6	
			Cs-137	6.3±1.0	6.0±5.0	0.0-14.7	
STW-407	Water	April 1985	I-131	8.0±0.0	7.520.8	6.2-8.8	
STW-408	Water	April 1985	H-3	3399±150	3559±630	2929-4189	
STW-409	Water	April 1985					
	(Blind)		Gr. alpha	29.7±1.8	32.0±5.0	23.3-40.7	
	Sample A		Ra-226	4.4±0.2	4.1±0.6	3.1-5.1	
			Ra-228	NAe	15.2±0.9	4.6-7.8	
			Uranium	NAe	7.0±6.0	0.0-17.4	
	Sample B		Gr. beta	74.3±11.8	72.0±5.0	63.3-80.7	
			Sr-89	12.3±7.6	10.0±5.0	1.3-18.7	
			Sr-90			12.4-17.6	
			Co-60	14.7±2.4			
			Cs-134	12.0±2.0		6.3-23.7	
			Cs-137	14.0±2.0	12.0±5.0	3.3-20.7	

Table A-1. (continued)

				Concentration in pCi/lb			
Lab	Sample	Date		TIML Result	EPA Result ^d		
Code	Туре	Collected	Analysis	±2σ ^C	1s, N=1 C	ontrol Limit	
STW-413	Water	May 1985	Sr-89 Sr-90	36.0±12.4 14.3±4.2	39.0±5.0 15.0±1.5	30.3-47.7 12.4-17.6	
STW-414	Water	May 1985	Gr. alpha Gr. beta	8.3±4.1 8.7±1.2	12.0±5.0 11.0±5.0	3.3-20.7 2.3-19.7	
STW-416	Water	June 1985	Cr-51 Co-60 Zn-65 Ru-106 Cs-134 Cs-137	44.7±6.0 14.3±1.2 50.3±7.0 55.3±5.8 32.7±1.2 22.7±2.4	44.0±5.0 14.0±5.0 47.0±5.0 62.0±5.0 35.0±5.0 20.0±5.0	45.3-52.7 5.3-22.7 38.3-55.7 53.3-70.7 26.3-43.7 11.3-28.7	
STW-418	Water	June 1985	H-3	2446±132	2416±351	1807-3025	
STM-421	Milk	June 1985	Sr-89 Sr-90 I-131 Cs-137 K	10.3±4.6 9.0±2.0 11.7±1.2 12.7±1.2 1512±62	11.0±5.0 11.0±1.5 11.0±6.0 11.0±5.0 1525±132	2.3-19.7 8.4-13.6 0.6-21.4 2.3-19.7 1393-1657	
STW-423	Water	July 1985	Gr. alpha Gr. beta	5.0±0.0 5.0±2.0	11.0±5.0 8.0±5.0	2.3-19.7 0.0-16.7	
STW-425	Water	August 1985	I-131	25.7±3.0	33.0±6.0	22.6-43.4	
STW-426	Water	August 1985	H-3	4363±83	4480±447	3704-5256	
STAF-427	Aim Filter	August 1985	Gr. alpha Gr. beta Sr-90 Cs-137	11.3±0.6 46.0±1.0 17.7±0.6 10.3±0.6	44.0±5.0 18.0±1.5	4.3-21.7 35.3-52.7 15.4-20.6 0.0 16.7	
STW-429	Water	Sept 1985	Sr-89 Sr-90	15.7±0.6 7.0±0.0		11 3-28.7	
STW-430	Water	Sept 1985	Ra-226 Ra-228	8.2±0.3 4.1±0.3	8.9±1.3 4.6±0.7	6.6-11.1 3.4-5.8	
STW-431	Water	Sept 1985	Gr. alpha Gr. beta		8.0±5.0 8.0±5.0		

Table A-1. (continued)

					Concentration in pCi/lb			
Lab	Sample	D	ate	Analysis	TIML Result	EPA Resultd		
Code	Туре	Collected	lected		±2σ ^C	1s, N=1	Control Limits	
STW-433	Water	Oct	1985	Cr-51 Co-60 Zn-65 Ru-106 Cs-134 Cs-137	<13 19.3±0.6 19.7±0.6 <19 17.0±1.0 19.3±1.2	21.0±5.0 20.0±5.0 19.0±5.0 20.0±5.0 20.0±5.0	11.3-28.7 10.3-27.7 11.3-28.7 11.3-28.7	
STW-435	Water	Oct	1985	H-3	1957±50	1974±345	1376-2572	
STW-436 437	Water (31ind)	Oct	1985					
	Sample A			Gr. alpha Ra-226 Ra-228 Uranium	53.0±1.0 5.9±0.1 8.2±0.1 NA ^e		4.1-7.9 7.5-12.7	
	Sample B			Gr. beta Sr-89 Sr-90 Co-60 Cs-134 Cs-137	85.7±2.5 21.3±1.5 10.3±0.6 18.3±1.2 16.3±1.2 19.0±1.0	75.0±5.0 27.0±5.0 9.0±1.5 18.0±5.0 18.0±5.0	18.3-35.7 6.4-11.6 9.3-26.7 9.3-26.7	
STM-439	Milk	Oct	1985	Sr-89 Sr-90 I-131 Cs-137	50.3±0.6 23.3±0.6 45.7±3.2 60.7±0.6 1547±29	48.0±5.0 26.0±1.5 42.0±6.0 56.0±5.0 1540±77	23.4-28.6 31.6-52.4 47.3-64.7	
STW-441	Water	Nov	1985	Gr. alpha Gr. beta		10.0±5.0 13.0±5.0	1.3-18.7 4.3-21.7	
STW-443	Water	Dec	1985	I-131	46.7±2.1	45.0±6.0	34.6-55.4	
STW-444	Water	Dec	1985	Ra-226 Ra-228	6.5±0.1 6.1±0.1	7.1±1.1 7.3±1.1	5.2-9.0 5.4-9.2	
STW-445	Water	Jan	1986	Sr-89 Sr-90	29.7±2.5 13.7±0.6	31.0±5.0 15.0±1.5		
STW-446	Water	Jan	1986		3.0±0.0 5.3±0.6		0.0-11.7 0.0-15.7	

Table A-1. (continued)

				Concentration in pCi/lb			
Lab	Sample	Date		TIML Result	EPA	Resultd	
Code	Туре	Collected	Analysis	±20°	1s, N=1	Control Limits	
STF-447	Food	Jan 1986	Sr-89	24.3±2.5	25.0±5.0	16.3-33.7	
			Sr-90	17.3±0.6	10.0±1.5		
			I-131	22.7±2.3	20.0±6.0		
			Cs-137	16.3±0.6	15.0±5.0		
			K	927±46	950±144	701-1199	
STW-448	Water	Feb 1986	Cr-51	45.0±3.6	38.0±5.0		
			Co-60	19.7±1.5	18.0±5.0		
			Zn-65	44.0±3.5	40.0±5.0		
			Ru-106	<9.0		0.0-8.7	
			Cs-134			21.3-38.7	
			Cs-137	23.7±0.6	22.0±5.0	13.3-30.7	
STW-449	Water	Feb 1986	H-3	5176±48	5227±525	4317-6137	
STW-450	Water	Feb 1986	U total	8.0±0.0	9.0±6.0	0.0-19.4	
STM-451	Milk	Feb 1986	I-131	7.0±0.0	9.0±6.0	0.0-19.4	
STW-452	Water	March 1986	Ra-226	3.8±0.1	4.1±0.6	3.0-5.2	
			Ra-228	11.0±0.5	12.4±1.8	9.2-15.5	
STW-453	Water	March 1986	Gr. alpha	6.7±0.6	15.0±5.0	6.3-23.7	
			Gr. beta	7.3±0.6	8.0±5.0	0.0-16.7	
STW-454	Water	April 1986	I-131	7.0±0.0	9.0±6.0	0.0-19.4	
STW-455 456	Water (Blind)	April 1986					
	Sample A	4	Gr. alpha	15.0±1.0	17.0±5.0	8.3-25.7	
			Ra-226	3.1±0.1		2.1-3.7	
						1.5-2.5	
			Uranium	4.7±0.6	5.0±6.0	0.0-15.4	
	Sample B	3	Gr. beta	28.7±1.2	35.0±5.0		
			Sr-89	5.7±0.6	7.0±5.0	0.0-15.7	
			Sr-90	7.0±0.0	7.0±1.5	4.4-9.6 1.3-18.7	
			Co-60		10.0±5.0	1.3-18.7	
			Cs-134 Cs-137	4.0±1.7 5.3±0.6	5.0±5.0 5.0±5.0	0.0-13.7	
			US-13/	5.310.0	5.015.0	0.0-13.7	

Table A-1. (continued)

				A COLOR SERVICIO DE LA CONTRA PRESENTA DE LA COLOR DE	tration in	NAME OF THE PARTY
Lab	Sample	Date		TIML Result	AND DESCRIPTION OF THE PERSON	A Result ^d
Code	Туре	Collected	Analysis	±2σ ^C	1s, N=1	Control Limits
STAF-457	Air	April 1986	Gr. alpha	13.7±0.6	15.0±5.	0 6.3-23.7
	Filter		Gr. beta	46.3±0.6	47.0±5.	0 38.3-55.7
			Sr-90	14.7±0.6	18.0±1.	5 15.4-20.6
			Cs-137	10.7±0.6	10.0±5.	0 1.3-18.7
CTU-458	Urine	April 1986	Tritium	4313±70	4423±189	9 4096-4750
STW-459	Water	May 1986	Sr-89	4.3±0.6	5.0±5.0	0.0-13.7
			Sr-90	5.0±0.0	5.0±1.	5 2.4-7.6
STW-460	Water	May 1986	Gr. alpha	5.3±0.6	8.0±5.0	0.0-16.7
			Gr. beta	11.3±1.2	15.0±5.	0 6.3-23.7
STW-461	Water	June 1986	Cr-51	<9.0	0.0±5.0	0.0-8.7
			Co-60	66.0±1.0	66.0±5.0	57.3-74.7
			Zn-65	87.3±1.5	86.0±5.0	77.3-94.7
			Ru-106	39.7±2.5	50.0±5.0	
			Cs-134	49.3±2.5	49.0±5.0	
			Cs-137	10.3±1.5	10.0±5.	
STW-462	Water	June 1986	Tritium	3427±25	3125±36	2499-3751
STM-464	Milk	June 1986	Sr-89	<1.0	0.0±5.0	
			Sr-90	15.3±0.6	16.0±1.5	13.4-18.6
			I-131	48.3±2.3	41.0±6.0	30.6-51.4
			Cs-137	43.7±1.5	31.0±5.0	22.3-33.7
			K	1567±114	1600±80	1461-1739
STW-465	Water	July 1986	Gr. alpha	4.7±0.6	6.0±5.0	
			Gr. beta	18.7±1.2	18.0±5.0	9.3-26.7
STW-467	Water	August 1986	I-131	30.3±0.6	45.0±6.0	34.4-55.4
STW-468	Water	August 1986	Pu-239	11.3±0.6	10.1±1.0	8.3-11.9
STW-469	Water	August 1986	Uranium	4.0±0.0	4.0±6.0	0.0-14.4
STAF-470	Air	Sept 1986	Gr. alpha	19.3±1.5	22.0±5.0	13.3-30.7
471	Filter			64.0±2.6		57.3-74.7
472			3r-90	22.0±1.0		19.4-24.6
			Cs-137	25.7±1.5	22.0±5.0	
STW-473	Water	Sept 1986	Ra-226	6.0±0.1	6.1±0.9	4.5-7.7
			Ra-228	8.7±1.1	9.1±1.4	

Table A-1. (continued).

				Concentration in pCi/lb			
Lab	Sample	Date		TIML Result	EPA	Resultd	
Code	Туре	Collected	Analysis	±2σ ^C	1s, N=1	Control Limits	
STW-474	Water	Sept 1986	Gr. alpha Gr. beta	16.3±3.2 9.0±1.0	15.0±5.0 8.0±5.0		
STW-475	Water	Oct 1986	Cr-51 Co-60 Zn-65 Ru-106 Cs-134 Cs-137	63.3±5.5 31.0±2.0 87.3±5.9 74.7±7.4 25.7±0.6 46.3±1.5	59.0±5.0 31.0±5.0 85.0±5.0 74.0±5.0 28.0±5.0	22.3-39.7 76.3-93.7 65.3-82.7 19.3-36.7	
STW-476	later	Oct 1986	H-3	5918±60	5973±597	4938-7008	
SPW-477 478	Water (Blind)	Oct 1986					
	Sample A	4	Gr. alpha Ra-226 Ra-228 Uranium	34.0±6.0 5.8±0.2 2.7±1.0 11.0±0.0	40.0±5.0 6.0±0.9 5.0±0.8 10.0±6.0	4.4-7.6 3.7-6.3	
	Sample E	3	Gr. beta Sr-89 Sr-90 Co-60 Cs-134 Cs-137	38.7±1.2 5.0±0.0 3.0±0.0 24.7±1.2 11.0±2.0 9.3±1.2	51.0±5.0 10.0±5.0 4.0±1.5 24.0±5.0 12.0±5.0 8.0±5.0	1.3-18.7 1.4-6.6 15.3-32.7 3.3-20.7	
STM-479	Milk	Nov 1986	Sr-89 Sr-90 I-131 Cs-137	7.7±1.2 1.0±0.0 52.3±3.1 45.7±3.1 1489±104	9.0±5.0 0.0±1.5 49.0±6.0 39.0±5.0 1565±78	0.0-2.6 38.6-59.4	
STU-480	Urine	Nov 1986	H-3	5540±26	5257±912	4345-6169	
STW-481	Water	Nov 1986	Gr. alpha Gr. beta	12.0±4.0 20.0±3.5	20.0±5.0 20.0±5.0		
STW-482	Water	Dec 1986	Ra-226 Ra-228	6.7±0.2 5.2±0.2	6.8±1.0 11.1±1.7	5.0-8.6 8.2-14.0	
STW-483	Water	Jan 1987	Sr-89 Sr-90	19.7±5.0 21.0±2.0	25.0±5.0 25.0±1.5		

Table A-1. (continued).

					Printed Strategy region and an extra strategy of the strategy	tration in	AND
Lab		D			TIML Result	emission compared and control or provide the	Resultd
Code	Туре	Col	lected	Analysis	±2oc	1s, N=1	Control Limits
STW-484	Water	Jan	1987	Pu-239	17.0±2.3	16.7±1.7	13.8-19.6
S (F-486	Food	lan	1987	Sr-90 I-131 Cs-137 K	36.0±4.0 78.0±3.4 89.7±3.0 942±56	49.0±10.0 78.0±8.0 84.0±5.0 980±49	64.1-91.9
STF-487	Food (Blank)		1987	SR-90 I-131 Cs-137 K	2.0±0.0 <3 <2 993±102		
STW-488	Water	Feb	1987	Co-60 Zn-65 Ru-106 Cs-134 Cs-137	96.0±7.2		41.3-58.7 82.3-99.7 91.3-108.7 50.3-67.7 78.3-95.7
STW-489	Water	Feb	1987	H-3	4130±140	4209±420	3479-4939
STW-490	Water	Feb	1987	Uranium	8.3±1.2	8.0±6.0	0.0-18.4
STM-491	Milk	Feb	1987	I-131	10.0±0.0	9.0±0.9	7.4-10.6
STW-492	Water	Mar	1987	Gr. alpha Gr. beta	3.7±1.2 11.3±1.2	3.0±5.0 13.0±5.0	0.0-11.7 4.3-21.7
STW-493	Water	Mar	1987	Ra-226 Ra-228	7.0±0.1 7.1±2.3	7.3±1.1 7.5±1.1	5.4-9.2 5.5-9.5
STW-4.14	Water	Apr	1987	I-131	8.0±0.0	7.0±0.7	5.8-8.2
STAF-495	Air Filter	Apr	1987	Sr-90	15.0±0.0 41.0±2.0 16.3±1.2 7.0±0.0	17.0±1.5	34.3-51.7 14.4-19.6
	Water (Blind)	Apr	1987				
	Sample A				30.7±1.2 3.9±0.2 4.9±0.9 5.0±0.0	3.9±0.6 4.0±0.6	16.1-43.9 2.9-4.9 3.0-5.0 0.0-15.4

Table A-1. (continued)

				Concen	tration in	pCi/1b	
Lab	Sample	Date		TIML Result			
Code	Туре	Collected	Analysis	±2σ ^C	1s, N=1	Control Limits	
STW-496 497	Water (Blind)	Apr 1987				According to the second	
	Sample E	3	Gr. Beta Sr-89 Sr-90 Co-60 Cs-134 Cs-137	69.3±9.4 16.3±3.0 10.0±0.0 8.3±3.0 19.0±2.0 14.7±1.2	66.0±5.0 19.0±5.0 10.0±1.5 8.0±5.0 20.0±5.0	10.3-27.7 7.4-12.6 0.0-16.7 11.3-28.7	
STU-498	Urine	Apr 1987	H-3	6017±494	5620±795	4647-6593	
STW-499	Water	May 1987	Sr-89 Sr-90	38.0±6.0 21.0±2.0	41.0±5.0 20.0±1.5	The second secon	
STW-500	Water	May 1987	Gr. alpha Gr. beta	9.0±3.4 10.3±1.2	11.0±5.0 7.0±5.0		
STW-501	Water	June 1987	Cr-51 Co-60 Zn-65 Ru-106 Cs-134 Cs-137	40.0±8.0 60.3±3.0 11.3±5.0 78.3±6.4 36.7±3.0 80.3±4.2	41.0±5.0 64.0±5.0 10.0±5.0 75.0±5.0 40.0±5.0 80.0±5.0	55.3-72.7 1.3-18.7 66.3-83.7 31.3-48.7	
STW-502	Water	June 1987	H-3	2906±86	2895±357	2277-3513	
STW-503	Water	June 1987	Ra-226 Ra-228	6.9±0.1 13.3±1.0	7.3±1.1 15.2±2.3	5.4-9.2 11.2-19.2	
STM-504	Milk	June 1987	Sr-89 Sr-90 I-131 Cs-137	57.0±4.3 32.0±1.0 64.0±2.0 77.7±0.6 1383±17	69.0±5.0 35.0±1.5 59.0±6.0 74.0±5.0 1525±76	48.6-69.4 65.3-82.7	
STW-505	Water	July 1987	Gr. alpha Gr. beta	2.3±0.7 4.0±1.0	5.0±5.0 5.0±5.0		
STF-506	Food	July 1987	I-131 Cs-137 K	82.7±4.6 53.7±3.0 1548±57	80.0±8.0 50.0±5.0 1680±84	41.3-58.7	
STW-507	Water	Aug 1987	I-131	45.7±4.2	48.0±6.0	37.6-58.4	
STW-508	Water	Aug 1987	Pu-239	5.8±0.2	5.3±0.5	4.4-6.2	

Table A-1. (continued)

			- Analysis	Concentration in pCi/lb			
Lab	Sample	Date		TIML Result	EPA Resultd		
Code	Туре	Collected		£σc	1s, N=1	Control Limits	
STW-509	Water	Aug 1987	Uranium	13.3±0.3	13.0±6.0	2.6-23.4	
STAF-510	Air Filter	Aug 1987	Gr. alpha Gr. beta Sr-90 Cs-137	9.7±0.4 28.3±0.6 10.0±0.9 10.0±1.0	10.0±5.0 30.0±5.0 10.0±1.5 10.0±5.0	21.3-38.7 7.4-12.5	
STW-511	Water	Sept 1987	R a-226 R a-228	9.9±0.1 8.1±1.4	9.7±1.5 6.3±1.0		
STW-512	Water	Sept 1987	Gr. alpha Gr. beta	2.0±0.6 11.3±1.3	4.0±.0 12.0±5.0		
STW-513	Water	Oct 1987	H-3	4473±100	4492 ±449	3714-5270	
STW-514	Water (Blind)	Oct 1987					
	Sample A		Gr. alpha Ra-226 Ra-228 Uranium	29.3 ±2.6 4.9 ±0.1 4.2 ±1.0 3.0 ±0.1	28.0±7.0 4.8±0.7 3.6±0.5 3.0±6.0	3.6-6.1	
	Sample B		S r-89 S r-90 C o-60 C s-134 C s-137	14 .3 ±1 .3 9 .7 ±0 .4 16 .7 ±3 .0 16 .7 ±2 .3 24 .3 ±3 .3	16.0±5.0 10.0±1.5 16.0±5.0 16.0±5.0 24.0±5.0	7.3-24.7	
STW-516	Water	Oct 1987	C r-51 C o-60 Z n-65 R u-106 C s-134 C s-137	80 .3 ±17 .5 16 .0 ±2 .3 46 .3 ±5 .6 57 .3 ±15 .4 23 .7 ±2 .5 51 .7 ±3 .2	61.0±5.0	6.3-23.7 37.3-54.7 52.3-69.7 16.3-33.7	
STU-517	Urine	Nov 1987	H-3	7267 ±100	7432 ±743	6145-8719	
TW-518	Water	Nov 1987		3.0 ±2.0 15.7 ±2.3	7.0±5.0 19.0±5.0		
TW-519	Water	Dec 1987	I-131	26.0±3.0	26.0±6.0	15.6-36.4	

Table A-1. (continued)

				Concentration in pCi/lb			
Lab	Sample	Date		TIML Result	EPA	Resultd	
Code	Туре	Collected	Analysis	±2σ C	1s, N=1 (Control Limits	
STW-520	Water	Dec 1987	R a-226 R a-228	5.1 ±0.8 3.4 ±0.1	4.8±0.7 5.3±0.8	3.6-6.0 3.9-6.7	
STW-521	Water	Jan 1988	Sr-89 Sr-90	27 .3 ±5 .0 15 .3 ±1 .2	30.0±5.0 15.0±1.5	21 .3-38 .7 12 .4-17 .6	
STW-523	Water	Jan 1988	Gr. alpha Gr. beta	2.3±1.2 7.7±1.2	4.0±5.0 8.0±5.0	0.0-12.7	
STF-524	Food	Jan 1988	Sr-90	44.0±4.0 53.0±2.0 102.3±4.2 95.7±6.4 1010.7±158.5	102.0±10.2 91.0±5.0	50.2-59.8	
STW-525	Water	Feb 1988	C 0-60 Z n-65 R u-106 C s-134 C s-137	69.3±2.3 99.0±3.4 92.7±14.4 61.7±8.0 99.7±3.0	69.0±5.0 94.0±9.4 105.0±10.5 64.0±5.0 94.0±5.0	77.7-110.3 86.8-123.2	
STW-526	Water	Feb 1988	H-3	3453±103	3327 ±362	2700-3954	
STW-527	Water	Feb 1988	Uranium	3.0±0.0	3.0±6.0	0.0-13.4	
TM-538	Milk	Feb 1988	I-131	4.7±1.2	4.0±0.4	3.3-4.7	
STW-529	Water	Mar 1988	R a-226 R a-228	7.1±0.6 NA e	7.6±1.1 7.7±1.2		
STW-530	Water	Mar 1988	Gr. alpha Gr. beta	4.3±1.2 13.3±1.3		0.0-14.7	
TAF-531	Air Filter	Mar 1988	Gr. beta Sr-90	21 .0 ±2 .0 48 .0 ±0 .0 16 .7 ±1 .2 18 .7 ±1 .3	50.0±5.0 17.0±1.5	41.3-58.7 14.4-19.6	
TW-532	Water	Apr 1988	I-131	9.0±2.0	7.5±0.8	6.2-8.8	

Table A-1. (continued)

, ,,

				Concer	itration in p	C1/1b
Lab	Sample	Date		TIML Result	EPA	Resultd
Code	Туре	Collected	Analysis	±2 ₀ C	1s, N=1 (Control Limits
STW-533 534	Water (Blind)	Apr 1988				
	Sample A		Gr. alpha Ra-226 Ra-228 Uranium	ND f ND ND 6.0±0.0	46.0±11.0 6.4±1.0 5.6±0.8 6.0±6.0	27.0-65.0 4.7-8.1 4.2-7.0 0.0-16.4
	Sample B		Gr. beta Sr-89 Sr-90 Co-60 Cs-134 Cs-137	ND 3.3±1.2 5.3±1.2 63.3±1.3 7.7±1.2 8.3±1.2	57.0±5.0 5.0±5.0 5.0±1.5 50.0±5.0 7.0±5.0 7.0±5.0	48.3-65.7 0.0-13.7 2.4-7.6 41.3-58.7 0.0-15.7
STU-535	Urine	Apr 1988	H-3	6483±155	6202±620	5128-7276
STW-536	Water	Apr 1988	Sr-89 Sr-90	14.7±1.3 20.0±2.0	20.0±5.0 20.0±1.5	11.3-28.7 17.4-22.6
STW-538	Water	June 1988	Cr-51 Co-60 Zn-65 Ru-106 Cs-134 Cs-137	331.7±13.0 16.0±2.0 107.7±11.4 191.3±11.0 18.3±4.6 26.3±1.2	302.0±30.0 15.0±5.0 101.0±10.0 195.0±20.0 20.0±5.0 25.0±5.0	6.3-23.7 83.7-118.3 160.4-229.6
STW-539	Water	June 1988	H-3	5586±92	5565±557	4600-6530
STM-541	Milk	June 1988		52.7±3.1	51.0±5.0	31.3-48.7 54.8-65.2 78.4-109.6 42.3-59.7 1461.4-1738.6
STW-542	Water	July 1988		8.7±4.2 5.3±1.2	15.0±5.0 4.0±5.0	
STF-543	Food	July 1988	Sr-89 Sr-90 I-131 Cs-137 K	ND ND 115.0±5.3 52.7±6.4 1190.0±66.1	49.0±5.0	30.5-37.5 88.0-126.0

Table A-1. (continued)

				Concer	ntration in p(1/16
Lab	Sample	Date		TIML Result	EPA F	esultd
Code	Туре	Collected	Analysis	±20°C	1s, N=1 Co	entrol Limits
STW-544	Water	Aug 1988	I-131	80.0±0.0	76.0±8.0	62.1-89.9
STW-545	Water	Aug 1988	Pu-239	11.0±0.2	10.2±1.0	8.5-11.9
STW-546	Water	Aug 1988	Uranium	6.0±0.0	6.0±6.0	0.0-16.4
STAF-547	Air Filter	Aug 1988	Gr. alpha Gr. beta Sr-90 Cs-137	8.0±0.0 26.3±1.2 8.0±2.0 13.0±2.0	8.0±5.0 29.0±5.0 8.0±1.5 12.0±5.0	0.0-16.7 20.3-37.7 5.4-10.6 3.3-20.7
STW-548	Water	Sep 1988	Ra-226 Ra-228	9.3±0.5 5.8±0.4	8.4±2.6 5.4±1.6	6.2-10.6 4.0-6.8
STW-549	Water	Sep 1988	Gr. alpha Gr. beta	7.0±2.0 11.3±1.2	8.0±5.0 10.0±5.0	0.0-16.7 1.3-18.7
STW-550	Water	Oct 1988	Cr-51 Co-60 Zn-65 Ru-106 Cs-134 Cs-137	252.0±14.0 26.0±2.0 158.3±10.2 153.0±9.2 28.7±5.0 16.3±1.2	251.0±25.0 25.0±5.0 151.0±15.0 152.0±15.0 25.0±5.0 15.0±5.0	207.7-294.3 16.3-33.7 125.0-177.0 126.0-178.0 16.3-33.7 6.3-23.7
STW-551	Water	Oct 1988	H-3	2333.3±127.0	2316.0±350.0	1709.8-2927.2
STU-555	Urine	Nov 1988	H-3	3030.0±208.8	3025.0±359.0	2403.2-3646.8
STW-556	Water	Nov 1988	Gr. alpha Gr. beta	9.0±3.5 9.7±1.2	9.0±5.0 9.0±5.0	0.3-17.7 0.3-17.7

Results obtained by Teledyne Isotopes Midwest Laboratory as a participant in the environmental sample crosscheck program operated by the Intercomparison and Calibration Section, Quality Assurance Branch, Environmental Monitoring and Support Laboratory, U. S. Environmental Protection Agency (EPA), Las Vegas, Nevada.
b All results are in the pCi/l, except for elemental potassium (K) data in milk,

which are in mg/l; air filter samples, which are in pCi/filter; and food, which is in mg/kg.

C Unless otherwise indicated, the TIML results are given as the mean ±2 standard deviations for three determinations.

d USEPA results are presented as the known values and expected laboratory precision (1s, 1 determination) and control limits as defined by EPA.

e NA = Not analyzed.

f ND = No data. Not analyzed due to relocation of the lab.

Table A-2. Crosscheck program results, thermoluminescent dosimeters (TLDs).

		The same and the s	THE PERSON NAMED IN COLUMN TWO IS NOT THE OWNER, THE PERSON NAMED IN COLUMN TWO IS NOT THE OWNER, THE PERSON NAMED IN COLUMN TWO IS NOT THE OWNER, THE PERSON NAMED IN COLUMN TWO IS NOT THE OWNER, THE PERSON NAMED IN COLUMN TWO IS NOT THE OWNER, THE PERSON NAMED IN COLUMN TWO IS NOT THE OWNER, THE PERSON NAMED IN COLUMN TWO IS NOT THE OWNER, THE PERSON NAMED IN COLUMN TWO IS NOT THE OWNER, THE PERSON NAMED IN COLUMN TWO IS NOT THE OWNER, THE PERSON NAMED IN COLUMN TWO IS NOT THE OWNER, THE PERSON NAMED IN COLUMN TWO IS NOT THE OWNER, THE PERSON NAMED IN COLUMN TWO IS NOT THE OWNER, THE PERSON NAMED IN COLUMN TWO IS NOT THE OWNER, T	the second secon
Lab TLD Code Type	Measurement	Teledyne Result ±20ª	Known Value ^C	Average ±2 od (all participants)
2nd International Intercomparison ^b	ercomparison ^b			
115-2 CaF2:Mn	Field	17.0±1.9	17.1	16.4±7.7
a included in the control of the con	Lab	20.8±4.1	21.3	18.817.6
3rd International Intercomparisone	ercomparisone			
115-3 CaF2:Mn	Field	30.7±3.2	34.514.8	31.5±3.0
aina	Lab	89.6±6.4	91.7±14.6	86.2±24.0
4th International Intercomparisonf	ercomparisonf			
115-4 CaF2:Mn	Field	14.1±1.1	14.1±1.4	16.049.0
anna	Lab (LGW)	9.3±1.3	12.2±2.4	12.0±7.6
	Lab (High)	40.411.4	45.8±9.2	43.9±13.2
5th International Intercomparison9	ercomparison9			
115-5A CaF2:Mn	Field	31.4±1.8	30.046.0	30.2±14.6
0.00	Lab at	77.4±5.8	75.2±7.6	75.8±40.4
	Lab at	96.6±5.8	88.4±8.8	90.7±31.2

Table A-2. (Continued)

Teledyne Result Rown Field Field Rown Field Result Field ValueC ValueC Lif-loo Field Bulbs Lab at the end Third Thir					Æ	
Lif-100 Field 30.3±4.8 Chips Lab at beginning Lab at the end the end the end the comparisonh Lif-100 Field 75.4±2.6 Chips Lab (Co-60) 80.0±3.5 Lab (Co-60) 66.6±2.5 CaF2:Mn Field 71.5±2.6 Bulbs Lab (Co-60) 84.8±6.4 Lab (Co-60) 84.8±6.4 Lab (Co-60) 82.5±3.7 CaSO4:Dy Field 76.8±2.7 Caso4:Dy Field 76.8±2.7 Caso4:Dy Field 76.8±2.7 Caso4:Dy Field 76.9±3.7	Lab	TLD Type	Measurement	Teledyne Result ±2σ ^a	Known Value ^C	Average ±20 d (all participants)
Lab at B1.117.4 beginning Lab at the end the end the end the end the end the end Lif-100 Field T5.4±2.6 Chips Lab (Co-60) 80.0±3.5 Lab (Co-60) 84.8±6.4 Lab (Co-60) 84.8±6.4 Lab (Co-60) 78.8±1.6 Cards Lab (Co-60) 82.5±3.7 Lab (Co-60) 82.5±3.	115-58	LiF-100	Field	30.3±4.8	30.0±6.℃	30.2±14.6
Lab at the end the end the end the end ternational Intercomparisonh Lif-l00 Field Chips Lab (Co-60) CaF2:Mn Field Bulbs Lab (Co-60) CaF2:Mn Field CaF2:Mn Field CaS04:Dy Fi		Squa	Lab at beginning	81.1±7.4	75.2±7.6	75.8±40.4
LiF-100 Field 75.4±2.6 Chips Lab (Co-60) 80.0±3.5 Lab (Co-60) 66.6±2.5 CaF2:Mn Field 71.5±2.6 Bulbs Lab (Co-60) 84.8±6.4 Lab (Co-60) 78.8±1.6 CaSOq:Dy Field 76.8±2.7 Cards Lab (Co-60) 82.5±3.7 Lab (Cs-137) 79.0±3.2			Lab at the end	85.4±11.7	88.4±8.8	90.7±31.2
Lif-100 Field 75.4±2.6 Chips Lab (Co-60) 80.0±3.5 Lab (Cs-137) 66.6±2.5 CaF2:Mn Field 71.5±2.6 Bulbs Lab (Co-60) 84.8±6.4 Lab (Cs-137) 78.8±1.6 CaSO4:Dy Field 76.8±2.7 Cards Lab (Co-60) 82.5±3.7 Lab (Cs-137) 79.0±3.2	7th Inter	rnational Inte	ercomparisonh			
CaF ₂ :Mn Field (Co-60) 80.0±3.5 66.6±2.5 66.6±2.5 84.8±6.4 Lab (Co-60) 84.8±6.4 Lab (Cs-137) 78.8±1.6 CaSO ₄ :Dy Field 76.8±2.7 Cards Lab (Co-60) 82.5±3.7 Lab (Cs-137) 79.0±3.2	115-7A	LiF-100	Field	75.4±2.6	75.8±6.0	75.1±29.8
CaF ₂ :Mn Field 71.5±2.6 84.8±6.4 Lab (Co-60) 84.8±6.4 Lab (Co-137) 78.8±1.6 CaSO ₄ :Dy Field 76.8±2.7 Cards Lab (Co-60) 82.5±3.7 Lab (Co-60) 79.0±3.2		CHIPS	Lab (Co-60)	80.0±3.5	79.9±4.0	77.9±27.6
CaF2:Mn Field Bulbs Lab (Co-60) 84.8±6.4 Lab (Cs-137) 78.8±1.6 CaSO4:Dy Field Cards Lab (Co-60) 82.5±3.7 Lab (Cs-137) 79.0±3.2			Lab (Cs-137)	66.6±2.5	75.0±3.8	73.0±22.2
CaSO ₄ :Dy Field Co-60) 84.8±6.4 CaSO ₄ :Dy Field 76.8±2.7 Cards Lab (Co-60) 82.5±3.7 Lab (Cs-137) 79.0±3.2	115-78	CaF2:Mn	Field	71.5±2.6	75.8±6.0	75.1±29.8
CaSO4:Dy Field Caso4:Dy Field Cards Lab (Co-60) Lab (Cs-137) 78.8±1.6 76.8±2.7 79.5±3.7 79.0±3.2		Sqips	Lab (Co-60)	84.8±6.4	79.944.0	77.9±27.6
CaSO4:Dy Field 76.8±2.7 Sards Lab (Co-60) 82.5±3.7 Lab (Cs-137 79.0±3.2			Lab (Cs-137)	78.811.6	75.0±3.8	73.0±22.2
Lab (Co-60) 82.5±3.7 Lab (Cs-137 79.0±3.2	115-70	CaS04:Dy	Field	76.8±2.7	75.8±6.0	75.1±29.8
79.0±3.2		00000	Lab (Co-60)	82.5±3.7	79.9±4.0	77.9±27.6
			Lab (Cs-137	79.0±3.2	75.0±3.8	73.0±22.2

(Continued)

				壬	
Lab	TLD	Measurement	Teledyne Result ±20ª	Known Value ^C	Average 12 o d (all participants)
8th Inter	8th International Intercomparison	rcomparisoni			
115-8A	LiF-100	Field Site 1	29.5±1.4	29.7±1.5	28.9112.4
	Chips	Field Site 2	11.3±0.8	10.4±0.5	10.1±9.06
		Lab (Cs-137)	13.7±0.9	17.2±0.9	16.216.8
115-88	CaF2:Mn	Field Site 1	32.3±1.2	29.7±1.5	28.9±12.4
	Bulbs	Field Site 2	9.0±1.0	10.4±0.5	10.149.0
		Lab (Cs-137)	15.8±0.9	17.2±0.9	16.216.8
115-80	CaSO4: Dy	Field Site 1	32.3±0.7	29.7±1.5	28.9412.4
	Cards	Field Site 2	10.6±0.6	10.4±0.5	10.1±9.0
		Lab (Cs-137	18.1±0.8	17.2±0.9	16.2±6.8

Lab result given is the mean ±2 standard deviations of three determinations.

Second International Intercomparison of Environmental Dosimeters conducted in April of 1976 by the Health and Safety Laboratory (GASL), New York, New York, and the School of Public Health of the University of Texas, Houston, Texas.

Value determined by sponsor of the intercomparison using continuously operated pressurized ion chamber

Ridge Third International Intercomparison of Environmental Dosimeters conducted in summer of 1977 by Oak Mean ±2 standard deviations of results obtained by all laboratories participating in the program. 9

National Laboratory and the School of Public Health of the University of Texas, Houston, Texas.

Fourth International Intercomparison of Environmental Dosimeters conducted in summer of 1979 by the School of Public Health of the University of Texas, Houston, Texas.

Fifth International Intercomparison of Environmental Dosimeter conducted in fall of 1980 at Idaho Falls, Idaho and sponsored by the School of Public Health of the University of Texas, Houston, Texas and Environmental Measurements Laboratory, New York, New York, U.S. Department of Energy.

Seventh International Intercomparison of Environmental Dosimeters conducted in the spring and summer of 1984 at Las Vegas, Nevada, and sponsored by the U.S. Department of Energy, the U.S. Nuclear Regulatory Commission, and the U.S. Environmental Protection Agency. 5

Eighth International Intercomparison of Environmental Dosimeters conducted in the fall and winter 1985-1986 at New York, New York, and sponsored by the U.S. Department of Energy.

Table A-3. In-house spiked samples.

				Concentr	ation in p	Ci/1
Lab Code	Sample Type	Date Collected	Analysis	TIML Resuit n≈3	Known Activity	The state of the s
QC-MI-6	Milk	Feb. 1986	Sr-89 Sr-90 I-131 Cs-134 Cs-137	6.0±1.9 14.2±1.7 34.2±3.8 32.0±1.8 35.8±2.1	6.4±3.0 12.9±2.0 35.2±3.5 27.3±5.0 35.0±5.0	8.7 5.2 10.4 8.7 8.7
QC-W-14	Water	Mar. 1986	Sr-89 Sr-90	1.6±0.4 2.4±0.2	1.6±1.0 2.4±2.0	7.1 4.2
QC-W-15	Water	Apr. 1986	I-131 Co-60 Cs-134 Cs-137	44.9±2.4 10.6±1.7 30.2±2.4 21.9±1.9	41.5±7.0 12.1±5.0 25.8±8.0 19.9±5.0	10.6 7.1b 7.1b 7.1b
QC-MI-7	Milk	Apr. 1986	I-131 Cs-134 Cs-137	39.7±3.3 28.7±2.8 21.2±2.8	41.5±7.0 25.8±8.0 19.9±5.0	10.4 8.7 8.7
SPW-1	Water	May 1986	Gross alpha	15.8±1.8	18.0±5.0	5C
QC-W-16	Water	June 1986	Gross alpha Gross beta	16.2±0.7 38.4±3.5	16.9±2.5 30.2±5.0	8.7 8.7
QC-MI-9	Milk	June 1986	Sr-89 Sr-90 I-131 Cs-134 Cs-137	<1.0 12.6±1.8 38.9±7.0 33.0±3.4 38.5±2.8	0.0 13.3±3.0 34.8±7.0 36.1±5.0 39.0±5.0	7.1b 4.2b 10.4 8.7 8.7
SPW-2	Water	June 1986	Gross alpha	16.8±1.8	18.0±5.0	5C
SPW-3	Water	June 1986	Gross alpha	17.7±0.8	18.0±5.0	5C
QC-W-18	Water	Sep. 1986		34.7±5.6 51.1±7.0	31.3±5.0 43.3±8.0	8.7 8.7
QC-W-19	Water	Sep. 1986	Sr-89 Sr-90		15.6±3.5 6.2±2.0	7.1b 4.2b

Table A-3. In-house spiked samples (continued)

				Concentr	ation in p	Ci/1
Lab Code	Sample Type	Date Collected	Analysis	TIML Result n=3	Known Activity	Precision 1s, n=3
QC-W-21	Water	Oct. 1986	Co-60 Cs-134 Cs-137	19.2±2.2 31.7±5.2 23.8±1.0	18.5±3.0 25.6±8.0 21.6±5.0	8.7 8.7 8.7
QC-MI-11	Milk	Oct 1986	Sr-89	12.3±1.8	14.3±3.0	8.7
QC-W-20	Water	Nov. 1986	H-3	3855±180	3960±350	520b
QC-W-22	Water	Dec. 1986	Gross alpha Gross beta	9.8±1.4 21.7±2.0	11.2±4.0 23.8±5.0	8.7 8.7
QC-W-23	Water	Jan. 1987	I-131	29.8±2.5	27.9±3.0	10.4
QC-MI-12	Milk	Jan. 1987	I-131 Cs-137	36.5±1.3 32.6±4.2	32.6±5.0 27.4±8.0	10.4
SPM-13	Milk	Jan 1987	Sr-89 Sr-90 I-131 Cs-134 Cs-137	10.4±2.1 14.6±1.6 49.5±1.2 <1.6 33.3±0.6	12.2±4.0 12.6±3.0 54.9±8.0 0.0 27.4±8.0	8.7 5.2 10.4 8.7 8.7
SPW-24	Water	Mar 1987	Sr-89 Sr-90	24.7±3.6 23.9±3.8	25.9±5.0 22.8±8.0	8.7 5.2
SPW-25	Water	Apr 1987	I-131	28.0±1.9	29.3±5.0	10.6
SPM-14	Milk	Apr 1987	I-131 Cs-134 Cs-137	25.0±2.2 <2.1 34.2±2.0	23.9±5.0 0.0 27.2±7.0	10.4 8.7 8.7
SPW-26	Water	Jun 1987	H-3 Co-60 Cs-134 Cs-137	3422±100 24.8±1.4 <2.0 21.2±0.5	3362±300 26.5±7.0 0.0 21.6±7.0	520 8.7 8.7 8.7
SPW-27	Water	Jun 1987	Gr. alpha Gr. beta	8.5±1.9 22.6±1.9	10.1±4.0 21.2±5.0	8.7 8.7
SPW-28	Water	Jun 1987	Gr. alpha Gr. beta	8.7±1.3 12.2±5.2	10.1±4.0 9.4±3.0	8.7 8.7

Table A-3. In-house spiked samples (continued)

				Concenti	ration in p	oC i/1
L ab Code	Sample Type	Date Collected	Aņalysis	TIML Result n=3	Known Activity	Expected Precision 1s, n=3ª
SPW-29	Water	Jun 1987	Gr. alpha Gr. beta		18.9±5.0 11.8±4.0	8.7 8.7
SPM-15	Milk	Jul 1987	Sr-89 I-131 Cs-134 Cs-137	19.4±1.6 43.5±0.7 17.9±2.2 25.4±1.8	18.8±3.5 45.3±7.0 16.0±5.3 22.7±5.0	5.2 10.4 8.7 8.7
SPW-30	Water	Sep 1987	S r-89 S r-90	17.5±3.0 18.4±2.2	14.3±5.0 17.5±2.2	8.7 5.2
SPW-31	Water	Oct 1987	H-3	2053±939	2059±306	520
SPW-32	Water	Dec 1987	Gr. alpha Gr. beta		10.1±5.0 13.1±3.0	8.7 8.7
SPW-33	water	Dec 1987	Gr. alpha Gr. beta		10.1.5.0 7.9±3.0	8.7 8.7
SPW-34	Water	Dec 1987	Gr. alpha Gr. beta		5.1±3.0 7.9±3.0	8.7 8.7
SPM-16	Иilk	Jan 1988	Sr-89 Sr-90 I-131 Cs-134 Cs-137	31.7±6.0 27.8±3.5 23.2±5.0 24.2±6.0 25.1±6.0	31.8±4.7 25.5±2.7 26.4±0.5 23.8±2.3 26.5±0.8	8.7 8.7 10.4 8.7 8.7
SPM-17	Milk	Feb 1988	I-131	10.6±1.2	14.3±1.6	10.4
SPW-35	Water	Feb 1988	I-131	9.7±1.1	11 .6 ±1 .1	10.4
SPW-36	Water	Feb 1988	I-131	10.5±1.3	11.6±1.0	10.4
SPW-37	Water	Mar 1988	Sr-89 Sr-90	19.8±8.0 17.3±5.0	17.1±2.0 18.7±0.9	8.7 5.2
SPM-18	Milk	Apr 1988	I-131 Cs-134 Cs-137	26.7±5.0 30.2±5.0 26.2±5.0	33.2±2.3 31.3±2.1 29.9±1.4	10.4 8.7 8.7

Table A-3. In-house spiked samples (continued)

				Concentr	ration in p	Ci/1
Lab Code	Sample Type	Date Collected	Analysis	TIML Result n=3	Known Activity	Expected Precision 1s, n=3ª
SPW-38	Water	Apr 1988	I-131	14.2±5.0	17.1±1.1	10.4
SPW-39	Water	Apr 1988	H-3	4176±500	4439±31	724
SPW-40	Water	Apr 1988	Co-60 Cs-134 Cs-137	26.1±4.0 29.2±4.5 26.2±4.0	23.7±0.5 25.4±2.6 26.6±2.3	8.7 8.7 8.7
SPW-41	Water	Jun 1988	Gr. alpha Gr. beta	13.1±5.0 20.1±5.0	12.3±0.4 22.6±1.0	8.7 8.7
SPS-42	Milk	Jul 1988	Sr-89 Sr-90 I-131 Cs-137	15.1±1.6 18.0±0.6 88.4±4.9 22.7±0.8	16.4±5.0 18.3±5.0 86.6±8.0 20.8±6.0	8.7 8.7 10.4 8.7
SPW-43	Water	Sep 1988	Sr-89 Sr-90	48.5±3.3 10.9±1.0	50.8±8.0 11.4±3.5	8.7 5.2
SPW-44	Water	Oct 1988	Co-60 Cs-134 Cs-137	20.9±3.2 38.7±1.6 19.0±2.4	21.4±3.5 38.0±6.0 21.0±3.5	8.7 8.7 8.7
SPW-45	Water	Oct 1988	I-131	22.2±0.6	23.3±3.5	10.4
SPW-46	Water	Oct 1988	H-3	4109±43	4153±500	724
SPS-46	Milk	Oct 1988	I-131 Cs-134 Cs-137	59.8±0.9 49.6±1.8 25.8±4.6	60.6±9.0 48.6±7.5 24.7±4.0	10.4 8.7 8.7
SPW 47	Water	Dec 1988	Gr. alpha Gr. beta	11.5±2.3 26.5±2.0	15.2±5.0 25.7±5.0	8.7 8.7

a n=3 unless noted otherwise. b n=2. c n=1.

Table A-4. In-house "blank" samples.

				Concentrat	ion in pCi/1
Lab Code	Sample Type	Date Collected	Analysis	Results (4.66 ₀)	Acceptance Criteria (4.66 σ)
BL-1	D.I. Water	Nov. 1985	Gross alpha	<0.1	<1
			Gross beta	<0.4	<4
BL-2	D.I. Water	Nov. 1985	Cs-137 (gamma)	<1.9	<10
BL-3	D.I. Water	Nov. 1985	Sr-89 Sr-90	<0.5 <0.6	<5 <1
			31 - 30	10.0	11
BL-5	D.I. Water	Nov. 1985	Ra-226	<0.4	<1
			Ra-228	<0.4	<1
PW-2265	D.I. Water	Apr. 1985	Gross alpha	<0.6	<1
			Gross beta	<2.2	<4
			Sr-89	<0.2	<5
			Sr-90	<0.4	<1
			I-131	<0.2	<1
			Cs-137 (gamma)	<7.4	<10
BL-6	D.I. Water	Apr. 1986	Gross alpha	<0.4	<1
BL-7	D.I. Water	Apr. 1986	Gross alpha	<0.4	<1
BL-8	D.I. Water	June 1986	Gross alpha	<0.4	<1
BL-9	D.I. Water	June 1986	Gross alpha	<0.3	<1
SPW-3185	D.I. Water	Jan 1987	Ra-226	<0.1	<1
			Ra-228	<0.9	<1
PS-3292	Milk	Jan 1987	I-131	<0.1	<1
			Cs-134	<6.2	<10
			Cs-137	<6.4	<10
PW-3554	D.I. Water	Feb 1987	H-3	<180	<300
			Gross beta	<2.6	<4
PS-3555	Milk	eb 1987	Sr-89	<0.6	<5
			Sr-90	1.9±0.4a	<1
PS-3731	Milk	Mar 1987	Cs-134	<2.2	<10
10-0101		1101 1207	Cs-137	<2.5	<10

a Low level (1 - 4 pCi/l) of Sr-90 concentration in milk is not unusual.

Table A-4. In-house "blank" samples (continued).

				Concentrati	on in pCi/1
Lab Code	Sample Type	Date Collected	Analysis	Results (4.66 ₀)	Acceptance Criteria (4.66σ)
SPS-3732	D.I. Water	Mar 1987	Sr-89 Sr-90 I-131 Co-60 Cs-134 (gamma) Cs-137 (gamma) Ra-226 Ra-228 Np-237 Th-230 Th-232 U-234 U-235 U-238	<0.9 <0.8 <0.3 <2.3 <2.2 <2.4 <0.1 <1.0 <0.04 <0.05 <0.02 <0.03 <0.03	<55 <1 <10 <10 <10 <11 <1 <0.1 <0.1 <0.1 <0
SPS-4023	Milk	May 1987	I-131	<0.1	<1
SPS-4203	D.I. Water	May 1987	Gross alpha Gross beta	<0.7 <1.7	<1 <4
SPS-4204	Milk	May 1987	Sr-89 Sr-90	<0.5 2.4±0.6 ^a	<5 <1
SPS-4390	Milk	Jun 1987	Cs-134 Cs-137	<4.7 <5.2	<10 <10
SPS-4391	D.I. Water	Jun 1987	Sr-89 Sr-90 I-121 Co-60 Cs-137 Ra-226 Ra-228	<0.4 <0.1 <0.1 <3.8 <5.7 <0.1 <0.9	<5 <1 <1 <10 <10 <11 <1
SPW-4627	D.I. Water	Aug 1987	Gross alpha Gross beta Tritium	<0.6 <1.4 <150	<1 <4
SPS-4628	Milk	Aug 1987	Sr-89 Sr-90	<0.6 2.4±0.6 ^a	<5 <1
SPS-4847	Milk	Sep 1987	Cs-134 Cs-137	<4.4 <5.3	<10 <10

a Low level (1 - 4 pCi/l) of Sr-90 concentration in milk is not unusual.

Data Reporting Conventions

1.0. All activities, except gross alpha and gross beta, are decay corrected to collection time or the end of the collection period.

2.0. Single Measurements

Each single measurement is reported as follows:

x ± s

where x = value of the measurement:

 $s = 2\sigma$ counting uncertainty (corresponding to the 95% confidence level).

In cases where the activity is found to be below the lower limit of detection L it is reported as

<L

where L = is the lower limit of detection based on 4.66 σ uncertainty for a background sample.

3.0. Duplicate Analyses

3.1. Individual results: $x_1 \pm s_1$ $x_2 \pm s_2$

Reported result: x ± s

where x = (1/2) $(x_1 + x_2)$ s = $(1/2)\sqrt{s_1^2 + s_2^2}$

3.2. Individual results: <L1

<L2

Reported result: <L

where L = 1 ower of L_1 and L_2

3.3. Individual results: $x \pm s$

<L

Reported result: $x \pm s$ if $x \ge L$;

<L otherwise

4.0. Computation of Averages and Standard Deviations

4.1 Averages and standard deviations listed in the tables are computed from all of the individual measurements over the period averaged; for example, an annual standard deviation would not be the average of quarterly standard deviations. The average \bar{x} and standard deviation(s) of a set of n numbers $x_1, x_2, \dots x_n$ are defined as follows:

$$\overline{x} = \frac{1}{n} \Sigma x$$

$$S = \sqrt{\frac{\sum (x - \overline{x})^2}{\eta - 1}}$$

- 4.2 Values below the highest lower limit of detection are not included in the average.
- 4.3 If all of the values in the averaging group are less than the highest LLD, the highest LLD is reported.
- 4.4 If all but one of the values are less than the highest LLD, the single value x and associated two sigma error is reported.
- 4.5. In rounding off, the following rules are followed:
 - 4.5.1. If the figure following those to be retained is less than 5, the figure is dropped, and the retained figures are kept unchanged. As an example, 11.443 is rounded off to 11.44.
 - 4.5.2 If the figure following those to be retained is greater than 5, the figure is dropped, and the last retained figure is raised by 1. As an example, 11.446 is rounded off to 11.45.
 - 4.5.3. If the figure following those to be retained is 5, and if there are no figures other than zeros beyond the five, the figure 5 is dropped, and the last-place figure retained is increased by one if it is an odd number or it is kept unchanged if an even number. As an example, 11.435 is rounded off to 11.44, while 11.425 is rounded off to 11.42.

Table A-4. In-house "blank" samples (continued).

				Concentratio	n in pCi/1
Lab Code	Sample Type	Date Collected	Analysis	Results (4.66)	Acceptance Criteria (4.66)
SPS-5615	Milk	Mar 1988	Cs-134	<2.4	<10
			Cs-137	<2.5	<10
			I-131	<0.3	<1
			Sr-89	<0.4	<5
			Sr-90	2.4±0.5a	<1
SPS-5650	D.I. Water	Mar 1988	Th-228	<0.3	<1
			Th-230	<0.04	<1
			Th-232	<0.05	<1
			U-234	<0.03	<1
			U-235	<0.03	<1
			U-238	<0.03	<1
			Am-241	<0.06	<1
			Cm-242	<0.01	<1
			Pu-238	<0.08	<1
			Pu-240	<0.02	<1
SPS-6090	ttilk	Jul 1988	Sr-89	<0.5	<1
			Sr-90	1.8±0.5 a	<1
			I-131	<0.4	<1
			Cs-137	<0.4	<10
SPW-6209	Water	Jul 1988	Fe-55	<0.8	<1
SPW-6292	Water	Sep 1988	Sr-89	<0.7	<1
			Sr-90	<0.7	<1
SPS-6477	Milk	Oct 1988	I-131	<0.2	<1
			Cs-134	<6.1	<10
			Cs-137	<5.9	<10
SPW-6478	Water	Oct 1988	I-131	<0.2	<1
SPW-6479	Water	Oct 1988	Co-60	<5.7	<10
			Cs-134	<3.7	<10
			Cs-137	<4.3	<10
SPW-6480	Water	Oct 1988	H+3	<170	<300
SPW-6625	Water	Dec 1988	Gr. alpha	<0.7	<1
			Gr. beta	<1.9	<4

a Low level (1 - 4 pCi/l) of Sr-90 concentration in milk is not unusual.

ATTACHMENT B

ACCEPTANCE CRITERIA FOR "SPIKED" SAMPLES

LABORATORY PRECISION: ONE STANDARD DEVIATION VALUES FOR VARIOUS ANALYSES

Analysis	Level	One Standard Deviation for Single Determination 5 pCi/liter 5% of known value	
Gamma Emitters	5 to 100 pCi/liter or kg >100 pCi/liter or kg		
Strontium-89b	5 to 50 pCi/liter or kg >50 pCi/liter or kg	5 pCi/liter 10% of known value	
Strontium-90b	2 to 30 pCi/liter or kg >30 pCi/liter of kg	3.0 pCi/liter 10% of known value	
Potassium	>0.1 g/liter or kg	5% of known value	
Gross Alpha	<20 pCi/liter >20 pCi/liter	5 pCi/liter 25% of known value	
Gross Beta	<100 pCi/liter >100 pCi/liter	5 pCi/liter 5% of known value	
Tritium	<4,000 pCi/liter >4,000 pCi/liter	ls = (pCi/liter) = 169.85 x (known).0933 10% of known value	
Radium-226, Radium-228	<0.1 pCi/liter	15% of known value	
Plutonium	0.1 pCi/liter, gram, or sample	10% of known value	
Iodine-131, Iodine-129b	<55 pCi/liter >55 pCi/liter	6 pCi/liter 10% of known value	
Uranium-238, Nickel-63 ^b , Technetium-99 ^b	<35 pCi/liter >35 pCi/liter	6 pCi/liter 15% of known value	
Iron-55 ^b	50 to 100 pCi/liter	10 pCi/liter 10% of known value	

a From EPA publication, "Environmental Radioactivity Laboratory Intercomparison Studies Program, Fiscal Year 1981-1982, EPA-600/4-81-004.

TIML limit.

Table A-4. In-house "blank" samples (continued).

L ab C ode		ample ype	Date Collected		Concentrati	on in pCi/l
				Analysis	Results (4.66σ)	Acceptance Criteria (4.66 o)
SPS-4848	D.I.	Water	Sep 1987	I-131	<0.2	<1
SPW-4849	D.I.	Water	Sep 1987	C o-60 C s-134 C s-137 S r-89 S r-90	<4.1 <4.8 <4.0 <0.7 <0.7	<10 <10 <10 <5 <1
SPW-4850	D.I.	Water	Sep 1987	Th-228 Th-232 U-234 U-235 U-238 Am-241 Cm-242 Ra-226 Ra-228	<0.04 <0.8 <0.03 <0.02 <0.06 <0.04 <0.1	<1 <1 <1 <1 <1 <1 <1 <1 <2
SPW-4859	D.I.	Water	Oct 1987	Fe-55	<0.5	<1
SPS-5348		Milk	Dec 1987	C s-134 C s-137	<2.3 <2.5	<10 <10
SPW-5384	D.I.	Water	Dec 1987	Co-60 Cs-134 Cs-137 I-131 Ra-226 Ra-228 Sr-89 Sr-90	<2.8 <2.6 <2.8 <0.2 <0.1 <1.2 <0.5 <0.4	<10 <10 <10 <1 <1 <1 <2 <1 <1
SPW-5385	D.I.	Water	Nov 1987	Gr. alpha Gr. beta Fe-55	<0.4 <2.2 <0.3	<1 <4 <1
SPS-5386		Milk	Jan 1988	I-131	<0.1	<1
SPW-5448	"Dead"	Water	Jan 1988	H-3	<177	<300

ADDENDUM TO APPENDIX A

The following is an explanation of the reasons why certain samples were outside the control limit specified by the Environmental Protection Agency for the Interlaboratory Comparison Program starting January 1987.

Lab Code	Analysis	TIML Result	EPA Control Limit	Explanation
STM-504	Sr-89 Sr-90	57.0±4.3 32.0±1.0	60.3-77.7 32.4-37.6	Milk had high fat content which made analyses difficult. Addition of errors to TIML result would put values within EPA control limits. EPA also had the same problem in analyzing its own sample.
STW-511	Ra-228	8.1±1.4	4.6-8.0	TIML results are usually within EPA control limits. Analysis of the next sample was within EPA control limits. No further action is planned.
STW-516	Cr-51	6J.3±17.5	61.3-78.7	Results in the past have been within EPA control limits and TIML will monitor the situation in the future.
STF-524	К	1010.7±158.5	1123.5-1336.5	Error in transferrance of data. Correct data was 1105±33. Results in the past have been within the limits and TIML will monitor the situation in the future.
STW-532	I-131	9.0±2.0	6.2-8.8	Sample recounted after 12 days. The average result was 8.8±1.7 (within EPA control limits). The sample was recounted in order to check the decay. Results in the past have been within the limits and TIML will continue to monitor the situation in the future.
STW-534	Co-60	63.3±1.3	41.3-58.7	High level of Co-60 was due to contamination of beaker. Beaker was discarded upon discovery of contamination and sample was recounted. Recount results were 53.2±3.6 and 50.9±2.4.

Appendix B

Data Reporting Conventions

Appendix C

Maximum Permissible Concentrations
of Radioactivity in Air and Water
Above Background in Unrestricted Areas

Table C-1. Maximum permissible concentrations of radioactivity in air and water above natural background in unrestricted areas.

Ai	r		W	ater
Gross alpha	3	pCi/m ³	Strontium-89	3,000 pCi/1
Gross beta	100	pCi/m ³	Strontium-90	300 pCi/1
Iodine-131b	0.14	pCi/m ³	Cesium-137	20,000 pCi/1
			Barium-140	20,000 pCi/1
			Iodine-131	300 pCi/1
			Potassium-40C	3,000 pCi/1
			Gross alpha	30 pCi/1
			Gross beta	100 pCi/1
			Tritium	3 x 106 pCi/1

a Taken from Code of Federal Regulations Title 10, Part 20, Table II and appropriate footnotes. Concentrations may be averaged over a period not greater

b than one year. From 10 CFR 20 but adjusted by a factor of 700 to reduce the dose resulting

c from the air-grass-cow-milk-child pathway. A natural radionuclide.

TRANSMITTAL MANIFEST

NORTHERN STATES POWER COMPANY

NUCLEAR GENERATION DEPARTMENT

PRAIRIE ISLAND NUCLEAR GENERATING PLANT

1988 Annual Radiological Environmental Monitoring Report

Manifest Date: April 12, 1989

USNRC	3
DCD	
Regional Admin - III	
Resident Inspector	
L R Eliason	1
P H Kamman	1
Monticello Plant Manager	2
Prairie Island Plant Manager	1 2 1
ERAD Dept.	1
Attn: Records Clerk	
Media Services	1
NSP (Wisconsin)	6
NRS File	1
NSS File	1
MDH	1 1 1
Attn: Commissioner of Health	
MPCA	1
Attn: J W Ferman	
ANI Library	1
Shaw Pittman Potts & Trowbridge	1
G Charnoff	
Safety Audit Committee	9
D M Musolf	
K J Albrecht	
C W Giesler	
D E Gilberts	
F W Hartley	
H S Isbin	
J A Thie	
F P Tierney	
E L Watzl	
File (Manifest Only)	