

ENVIRONMENTAL RADIATION MONITORING PLAN FOR

Pōhakuloa Training Area
Hawaii

FINAL

Revision 1

Submitted to:
Nuclear Regulatory Commission
Office of Nuclear Material Safety and Safeguards
Washington, D.C.

Prepared by:
U.S. Army Corps of Engineers
Baltimore District, Engineering Division
Environmental and Munitions Design Center
Baltimore, MD

Prepared for:
U.S. Department of Army
Installation Management Command
Fort Sam Houston, TX

03 February 2012

List of Acronyms and Abbreviations

ARS	Archive Search Report
BAX	Battle Area Complex
Cabrera	Cabrera Services, Inc.
CFR	Code of Federal Regulations
cm	centimeter
COC	chain of custody
DI	de-ionized
DOD	Department of Defense
DOT	U.S. Department of Transportation
DQI	data quality indicator
DQO	data quality objective
DU	depleted uranium
ERMP	Environmental Radiation Monitoring Plan
GWS	gamma walkover survey
HARNG	Hawaii Army National Guard
ICP-MS	inductively coupled plasma-mass spectroscopy
ID	identification
IDW	investigation derived waste
IMCOM	U.S. Army Installation Management Command
kg	kilogram
km	kilometer
km ²	square kilometer
L	liter
LCS	laboratory control sample
LOR	letter of receipt
m	meter
MCL	maximum contaminant level
MDC	minimum detectable activity
mg/m ³	milligram per cubic meter
MIDAS	Munitions Item Disposition Action System
mL	milliliter
mm	millimeter
mrem	millirem
MS/MSD	matrix spike/matrix spike duplicate
MW	monitoring well
NA	not applicable
NELAC	National Environmental Laboratory Accreditation Conference
NRC	U.S. Nuclear Regulatory Commission
pCi/g	picocurie per gram
pCi/L	picocurie per liter
PETN	pentaerythritol tetranitrate

pH	Definition: a measure of the acidity or basicity of an aqueous solution
PID	photoionization detector
PPE	personal protective equipment
PTA	Pōhakuloa Training Area
QA/QC	quality assurance/quality control
RCA	radiological control area
RSO	radiation safety officer
RSSO	Radiation Safety Staff Officer
SB	Schofield Barracks
SOW	scope of work
TMDL	total maximum daily load
TSA	technical systems audit
μCi/mL	microcurie per milliliter
μm	micrometer
μR/h	microroentgen per hour
USACE	U.S. Army Corps of Engineers
USEPA	U.S. Environmental Protection Agency
UXO	unexploded ordnance

TABLE OF CONTENTS

1.0	INTRODUCTION	1
1.1	Objective and Scope.....	1
2.0	PROJECT BACKGROUND	2
2.1	Site Description.....	3
2.2	History of Licensed Activities	3
2.3	Nature and Extent of Radiological Contamination	5
2.3.1	Scoping Survey	5
2.3.2	Characterization and Aerial Surveys	6
2.3.3	Range 11T Construction Support Surveys.....	7
3.0	ERMP STRATEGY AND PLAN.....	8
3.1	ERMP Goals and Rationale	8
3.2	Data Quality Objectives	8
3.3	Radiation Monitoring Strategy and Plans	9
3.3.1	Groundwater	11
3.3.2	Surface Water.....	13
3.3.3	Soil	14
3.3.4	Sediment	15
3.3.5	Air	15
3.3.6	Biota	16

TABLE OF CONTENTS, cont.

4.0	PROJECT ORGANIZATION AND MANAGEMENT.....	18
4.1	Responsible Organizations.....	18
4.1.1	Nuclear Regulatory Commission.....	18
4.1.2	Department of the Army	18
4.2	Lines of Authority.....	18
5.0	FIELD PROGRAM	19
5.1	Sampling Protocol.....	19
5.1.1	Pre-Mobilization Activities.....	19
5.1.2	Groundwater Sample Collection.....	19
5.1.3	Surface Water Sample Collection.....	21
5.1.4	Soil/Sediment Sample Collection	21
5.2	Sample Handling and Management	22
5.2.1	Sample Containers	22
5.2.2	Quality Control Samples.....	23
5.2.3	Sample Identification	23
5.2.4	Sample Custody	24
5.3	Field Measurements	24
5.3.1	Field Parameters.....	24
5.3.2	Equipment Calibration and Quality Control	25
5.3.3	Equipment Maintenance and Decontamination	26
5.4	Waste Management.....	27

TABLE OF CONTENTS, cont.

5.5 Record Keeping.....	27
6.0 SAMPLE ANALYTICAL REQUIREMENTS	29
6.1 Objective	29
6.2 Analytes of Concern.....	29
6.3 Analytical Laboratory Methods	29
6.4 Laboratory Method Analytical Uncertainty	30
6.5 Evaluation of Laboratory Results	30
6.6 Laboratory Quality Control.....	31
7.0 PROGRAMMATIC QUALITY ASSURANCE	34
7.1 Planned Audits and Assessments.....	35
7.1.1 Field Sampling Technical Systems Audits	35
7.1.2 Field Analytical Laboratory Technical Systems Audits	35
7.1.3 Fixed Laboratory Technical Systems Audits.....	35
7.1.4 Data Validation Technical Systems Audit.....	36
7.1.5 Laboratory Data Package Technical Systems Audit.....	36
7.1.6 Management Systems Reviews.....	37
7.2 Corrective Actions	37
7.2.1 Corrective Actions for Sampling Activities.....	37
7.2.2 Corrective Action during Laboratory Activities	38
7.2.3 Corrective Action during Data Validation.....	38
7.2.4 Corrective Action during Data Package Review	38

TABLE OF CONTENTS, cont.

7.2.5	General Corrective Actions.....	39
7.3	Quality Assurance Management Reports.....	39
8.0	ENVIRONMENTAL RADIATION MONITORING REPORT.....	40
9.0	REFERENCES	42

LIST OF TABLES

Table 1:	ERMP Monitoring Plans and Associated Screening Levels.....	10
Table 2:	Alpha Spectrometry MDCs and ICP-MS Detection Sensitivities	30
Table 3:	Laboratory Volumetric Quality Control Samples.....	32

LIST OF FIGURES

Figure 1:	Radiation Control Area Based on Firing Characteristics	5
Figure 2:	Sample Analytical Result Evaluation Decision Tree	31

LIST OF APPENDICES

Appendix A: ERMP Plates

Appendix B: RESRAD Groundwater Modeling Results

1.0 INTRODUCTION

This plan details the environmental radiation monitoring plan (ERMP) for the depleted uranium (DU) radiation control areas (RCAs) at Pōhakuloa Training Area (PTA). The ERMP for Schofield Barracks (SB) is a separate document. The ERMP is being conducted in accordance with the terms and conditions of the U.S. Army Installation Management Command (IMCOM) Nuclear Regulatory Commission (NRC) license application.

Depleted uranium (DU) has been found or suspected to be at a number of Army ranges across the United States as a result of test firing the Davy Crockett weapon system. This weapon system, fielded from 1962 through 1968, consisted of a portable recoilless rifle capable of launching a small yield nuclear warhead that was intended as a last-ditch effort against masses of invading Soviet troops in the event of war in Germany. It used a DU-bearing spotting round (20mm spotting M101) and an inert training warhead that was fired, in accordance with Atomic Energy Commission license conditions, at a number of locations across the country including Hawaii.

Only the spotting rounds were fired in Hawaii; the dummy warhead was not fired in Hawaii (USACE, 2007). An Army shipping manifest showed that for the Hawaiian sites, 714 spotting rounds, containing about 299 pounds of DU, were sent to Hawaii between 1962 and 1968. Evidence of spotting round use has been confirmed at PTA and SB.

Section 1.0 of this plan states the purpose and scope of this ERMP. Section 0 provides an overview of the site and its history related to operations involving DU-bearing Davy Crockett M101 spotting rounds that were fired.

The ERMP objectives, strategy, and associated action levels for the environmental media of concern are detailed in Section 3.0. The project organization and the roles and responsibilities of organizations associated with this program are defined in Section 4.0. The field program is presented in Section 5.0 and includes procedures associated with sample collection and management, field measurements, equipment preparation and decontamination, waste management, and recordkeeping. Laboratory sample analytical requirements are specified in Section 6.0 and programmatic quality assurance is specified in Section 7.0. Section 8.0 provides requirements for ERMP reporting. References used in this document are noted in Section 9.0.

1.1 Objective and Scope

The objective of this ERMP is to define the strategy and associated procedures for sampling environmental media within and surrounding the RCAs at PTA and to provide the basis for determining whether NRC-licensed M101 DU is migrating out of the RCAs.

The scope of this plan is limited to the PTA RCAs and their immediate environs and to sampling media to determine the presence or absence of DU. Sampled media will be analyzed at an off site laboratory to determine isotopic uranium concentrations. Uranium isotopic activity ratios will be used to establish whether or not DU is present in sampled media.

2.0 PROJECT BACKGROUND

This section provides an overview of the site (Section 2.1), followed by a summary of NRC-licensed activities (Section 2.2). A brief summary of the environmental sampling program conducted in support of the scoping and characterization surveys and ERMP is presented in Section 2.3.

2.1 Site Description

The Pōhakuloa Training Area is a large training area on the island of Hawaii that was acquired by the United States from the State of Hawaii and private landowners. The facility was and is still used by the U.S. Army Hawaii, the U.S. Marine Corps, and the Hawaii Army National Guard (HARNG). The PTA is located on the island of Hawaii between Mauna Loa, Mauna Kea and the Hualalai Volcanic Mountains. It extends up the lower slopes of Mauna Kea to approximately 6,800 feet in elevation and to about 9,000 feet on Mauna Loa. The training area is about midway between Hilo, on the east coast, and the Army landing site at Kawaihae Harbor. The area is the largest DOD installation in Hawaii.

The PTA consists of 108,863 acres, of which 24,048 are leased by the Army from the State of Hawaii. The remaining land is ceded and includes the impact and range areas and a portion of the west maneuver area. The leased areas include the northern maneuver areas and the support complex.

As the largest training area in Hawaii, Pōhakuloa can be used to accomplish nearly all of the varying types of training required by the military forces. PTA has a 51,000 acre impact area which is over 10 times the size of the one at Schofield Barracks. There are approximately 32,000 acres free of recent lava flows and are considered fully usable for large maneuver exercises. This area is more than twice the 14,000 acres of similar training land on all of Oahu. The impact area is surrounded on the north by several ranges and designated firing points for artillery. A support area of 600 acres containing logistic and administrative facilities plus quarters for approximately 2,000 troops are located to the north at the base of Mauna Kea. The Cantonment area was constructed in April 1955 from World War II prefabricated Quonset huts. The airfield was constructed in 1956.

The general vicinity and the potential Davy Crockett ranges on the installation are shown on Plate Numbers 1 and 2 respectively in Appendix A.

2.2 History of Licensed Activities

The Davy Crockett Weapon System consisted of two types of recoilless rifle weapons. One of those, the Davy Crockett Light Weapon M28, fired a 20-millimeter (mm) Spotting M101

cartridge. This cartridge consisted of a body manufactured from a D-38 uranium alloy (^{238}U Depleted Uranium). According to the Munitions Item Disposition Action System (MIDAS) database¹, the M101 cartridge contained approximately 6.7 ounces of DU. Based on Army shipping manifests, 714 spotting rounds were potentially fired in Hawaii. Radiological surveys and visual inspection have identified evidence of M101 spotting round use at PTA and SB. The precise number of spotting rounds fired at either location is unknown.

The U.S. Army Corps of Engineers (USACE) developed Archive Search Reports (ASR) for each installation or training center at which the M101 spotting rounds were fired. The M101 spotting rounds were fired into designated range areas that consisted of an impact area centered within a target zone. The target zone (impact area) covered approximately one square kilometer. The target zone was surrounded by a larger "Surface Danger Area." A firing position was located approximately 1,000 meters from the edge of the target zone. Each of the M101 DU-affected installations identified by the ASR project contains at least one Davy Crockett range. The individual Davy Crockett impact areas are generally situated within the larger installation active training range complex and in some cases are situated within an existing unexploded ordnance (UXO) hazard area. Each of the individual Davy Crockett impact areas, delineated as the target zone and as described in the installation ASRs, will be considered a Radiation Control Area (RCA). The RCA is initially defined as the "Target Zone" shown in Figure 1.

Plate Number 2 in Appendix A identifies the initially defined RCA boundaries for PTA based on the USACE ASR. The ASR identified four possible and likely firing points for training use of the M28 weapons system. Based on the location of the firing points, the location of known or suspected historical target locations, and the characteristics of the weapon system (Figure 1), the outer boundaries of the RCAs can be estimated by the limits of the "hatched" areas on the Plate.

¹ <https://midas.dac.army.mil/default.aspx> (username and password required for access)




Figure 1: Radiation Control Area Based on Firing Characteristics

2.3 Nature and Extent of Radiological Contamination

The impact areas at PTA where DU-bearing M101 spotting rounds have been fired have been the subject of extensive radiological investigations. The currently defined RCAs are based on these surveys and historical information summarized in the ASR and are shown in Plate Number 4. The results of radiological surveys, summarized in the subsections below, indicate that DU is confined to the range and is not being transported in measureable quantities beyond the RCAs.

2.3.1 Scoping Survey

Cabrera 2008a documents the results of scoping surveys performed at PTA. This investigation consisted of site visits, surface soil and debris sampling and analysis, and radiological scanning. Laboratory analyses of the samples included alpha spectrometry for uranium isotopes. Results of the scoping survey are summarized as follows:

- The visual and scanning surveys identified no distinct surface areas with yellow, oxidized DU metal fragments.
- The visual and scanning surveys did identify non-oxidized metal fragments, partial spotting round bodies, and Davy Crockett system components on Range 11T consistent with DU and the Davy Crockett weapons.

- Pistons associated with the Davy Crockett system were identified on Ranges 11T, 10, 14, and 17
- During the scoping survey, nine soil samples were collected around the perimeter of PTA at locations areas where sediment had or may have collected from past runoff/erosion. These analysis of these soil samples showed no indication of DU.

The nine soils samples collected were analyzed for isotopic uranium by alpha spectrometry. Results ranged from 0.19 pCi/g to 0.68 pCi/g of uranium. All of the results are consistent with naturally occurring concentrations of uranium. None of the results indicate DU, based on isotopic ratios.

2.3.2 Characterization and Aerial Surveys

Cabrera 2009 documents the results of characterization surveys performed at PTA. Aerial gamma surveys and gamma walkover surveys (GWS) surveys were performed over a total of 936 and 50 acres, respectively. Surveys were concentrated in areas that were the most likely to contain Davy Crockett spotting rounds and debris from the weapons system by focusing on the range of the weapon (one to two kilometers), as well as identification of targets from the era of the weapons system. Only limited visual and radiological indications of any additional Davy Crockett spotting rounds and system debris were located and identified during these aerial and ground based surveys.

Visual reconnaissance was conducted surrounding the perimeter of the range to identify locations of additional pistons and on the range to identify other Davy Crockett system components such as back plate assemblies, spotting round bodies, aluminum tail sections, etc, associated with the weapons system. Visual surveys confirmed the presence of additional pistons, but only on those ranges which had been previously identified as locations for Davy Crockett systems use. Aerial reconnaissance identified Davy Crockett back plate assemblies on Ranges 10/11T; ground-based survey crews were directed to these locations to perform biased surveys. No on-range visual identification of Davy Crockett system components was made at any other locations on the ranges. The back plate assemblies are highly visible from the air at the low altitude flown by the helicopter, as are the tail fin assemblies when they are located on the surface of the lava.

Ground based GWS located and identified two DU metal fragments, one essentially intact spotting round body with no tail fin assembly containing the PETN (pentaerythritol tetranitrate) explosive pellet and phosphorus spotting charge, and one aluminum tail fin with the DU spotting round body still attached. The DU fragments were located in a crack in the lava. GWS surveys identified four locations on the upper area of interest on Range 11T near the proposed battle area complex (BAX) construction footprint. The spotting round body containing the energetic

material was later removed from the surface of the range and placed in an adjacent location protected from both the weather and surface disturbances from range activities.

Twenty surface soil samples were collected in areas of GWS surveys and aerial surveys on both the range and BAX construction areas. These samples were analyzed by alpha spectrometry. Isotopic ratios indicated only naturally occurring uranium with no indications of depleted uranium.

Aerial gamma surveys identified some areas of increased gamma fluence, but these locations were further investigated and analyzed and the increases were determined to be from variations in naturally occurring radioactive materials. The number of DU spotting round bodies, aluminum fin assemblies and DU fragments are much fewer than would be expected given the total number of pistons which were identified. This fact, and in comparison to the number of DU fragments, and portions of the Davy Crockett spotting rounds found at Schofield Barracks, suggests that some type of range clearance may have occurred at PTA.

2.3.3 Range 11T Construction Support Surveys

In 2010 Cabrera Services, Inc. (Cabrera) performed various radiological surveys to support BAX construction activities on PTA Range 11T for the purpose of personnel and environmental protection (documented in Cabrera, 2011). Construction work activities included improving and expanding access roads, earthmoving and grading, and construction of military targets. These surveys and radiation protection controls were performed under Cabrera's broad scope NRC radioactive materials license with NRC's concurrence.

A part of the radiological support, Cabrera collected and analyzed daily air samples during intrusive activities. All air sample results were less than 1×10^{-14} $\mu\text{Ci/mL}$ alpha. This is less than 20 percent of the NRC effluent discharge limit from 10 CFR 20 Appendix B, Table 2, Column 1. In addition, analysis of the air samples did not confirm even trace amounts of DU.

Survey activities also included the performance of exposure rate measurements, gamma walkover surveys, and radiological contamination surveys. The results of all these surveys were consistent with background levels of radiation and there were no positive indications of DU.

3.0 ERMP STRATEGY AND PLAN

In this section, the ERMP strategy and plans are presented. The overall goals of the program are presented (Section 3.1), followed by the presentation of the data quality objectives (DQOs) (Section 3.2). For each environmental medium, the rationale and basis for sampling is presented, including action criteria and associated procedures if the action criteria are exceeded (Section □).

3.1 ERMP Goals and Rationale

The overall goals of the ERMP at PTA are to provide:

- A historical and current perspective of contaminant levels in various media
- An indication of the magnitude and extent of any DU release or migration from past operations
- A timely indication of DU contaminant release and migration.

Environmental monitoring activities are necessary at PTA to ensure that potential DU within the RCAs does not pose a threat to human health and the environment through inadvertent or unanticipated release or migration. These monitoring activities include the surveillance of all credible transport pathways; the selection of suitable surveillance locations; and the application of appropriate sampling methods, techniques, and analyses. To achieve this goal, the program has been designed to meet the applicable requirements of applicable Federal and State regulations, including NRC regulations and requirements.

Because the radioactive material is isolated within the RCAs and institutional controls are in place to prevent and control access to the area, exposure is not likely to occur. However, migration of this material through groundwater, rainwater runoff, soil, air, and biota is possible, but very unlikely. The PTA ERMP was developed to provide direct surveillance of the most probable migration route through periodic sampling and analysis of radioactive constituents. The following sections present the DQOs for this ERMP and discuss the rationale for the selection of the probable migration routes, sampling locations and frequencies, and action levels and associated steps to be taken if the action levels are exceeded.

3.2 Data Quality Objectives

The DQO process is a scientific data collection planning process designed to ensure that the type, quality, and quantity of data collected are appropriate for environmental decision-making. It consists of seven prescribed steps outlined in “Data Quality Objectives Process for Hazardous Waste Site Investigations” (U.S. Environmental Protection Agency [EPA] 2000).

DQOs define the purpose of the data collection effort, clarify what the data should represent to satisfy this purpose, and specify the performance requirements for the quality of information to be obtained from the data. These outputs then are used in the final step of the DQO process to develop a data collection design that meets all requirements and constraints.

The DQO process for this ERMP applies to the RCAs at PTA and consists of the following elements corresponding to steps in the DQO process:

- The primary objective for environmental sample collection at PTA is to provide data of known and sufficient quality to determine if conditions have changed since the previous sampling events. The data will help define the nature and extent (horizontal and vertical) of DU contaminant migration if it occurs (DQO Step 1 - State the Problem).
- The environmental sampling and field measurements will provide analytical data sufficient to determine if DU contamination in the RCAs is migrating to places outside the RCAs. The data will be used to support the development and selection of appropriate corrective actions if required (DQO Step 2 - Identify the Decision).
- ERMP data from previous and current sampling events at PTA, along with data from the scoping and characterization surveys and other related studies, will provide additional inputs to meet the objectives (DQO Step 3 - Identify Inputs to the Decision).
- The boundaries of the RCAs are depicted in Plate Number 4. The Study area is defined as areas adjacent to and surrounding the RCAs to determine potential migration from the RCAs (DQO Step 4- Define the Study Boundaries).
- Uranium isotopic ratios observed at PTA ERMP sampling locations will be evaluated to determine the absence or presence of DU and the extent of contamination migration at PTA (DQO Step 5 - Develop a Decision Rule).
- The sample analysis and validation will be performed in general accordance with the procedures contained in the ERMP (DQO Step 6 - Specify Limits on Decision Errors).
- Soil will be sampled quarterly to provide sufficient data concerning contaminant concentrations and potential migration. Sampling results will be used to determine if changes have occurred in contaminant trends or if sampling of additional media is appropriate. The results of trending analyses will be used to appropriately adjust sampling frequencies, in coordination with NRC (DQO Step 7 - Optimize the Design for Obtaining Data).

3.3 Radiation Monitoring Strategy and Plans

In this section, the rationale and plans for monitoring environmental media (i.e., groundwater, surface water, sediment, soil, air, and biota) are presented. Table 1 summarizes the ERMP, including planned monitoring activities by environmental medium and associated action criteria.

Table 1: ERMP Monitoring Plans and Associated Screening Levels

Medium	Plan	DU Action Criteria	Actions
Groundwater	<i>Frequency:</i> routine sampling not planned <i>Monitoring:</i> Routine groundwater sampling is not planned.	1) Confirmed presence of DU outside the boundaries of the RCAs as determined by the following:	1) Confirmed presence of DU outside RCAs: If the $^{238}\text{U} : ^{234}\text{U}$ activity ratio exceeds three at any location in any media, the License RSO will notify the NRC within 30 calendar days of receipt of analytical sampling results. Additional confirmatory sampling will be performed as soon as practicable. Further actions may be defined based on the results of confirmatory sampling including modification of sampling frequency and/or increasing the number of sample locations. Such actions will be coordinated with the NRC. 2) Suspected presence of DU outside RCAs: If trending analyses indicate an increase in $^{238}\text{U} : ^{234}\text{U}$ activity ratio and/or an increase in $^{238}\text{U} : ^{235}\text{U}$ mass ratio, the License RSO will conduct an independent review of the results. Based on the results of this review, the scope of the ERMP may be modified. Any proposed modifications will be coordinated with the NRC prior to implementation.
Surface Water	<i>Frequency:</i> routine sampling not planned <i>Monitoring:</i> Routine surface water sampling is not planned.	$^{238}\text{U} : ^{234}\text{U}$ activity ratio from alpha spectrometry analysis equal to or greater than three.	
Soil/Sediment	<i>Frequency:</i> quarterly soil sampling <i>Monitoring:</i> This plan includes quarterly soil sampling along roadway points of egress from the RCAs.	2) Suspected presence of DU outside the boundaries of the RCAs as determined by statistically based trending analysis indicative of: an increase in $^{238}\text{U} : ^{234}\text{U}$ activity ratio based on alpha spectrometry analysis	
Air	<i>Frequency:</i> routine sampling not planned <i>Monitoring:</i> Routine air sampling is not planned as part of this ERMP.	an increase in $^{238}\text{U} : ^{235}\text{U}$ mass ratio based on inductively coupled plasma-mass spectroscopy	

Background sampling will not be performed as part of this ERMP. DU is not naturally occurring. In addition, the collection of representative and defensible reference, background, or up-gradient samples can be challenging, especially in an active range situation where access to sites can be problematic. Consequently, project decisions will be based on the presence of DU as determined by uranium isotopic ratios, not by comparisons to background. Should DU be detected in any of the sampling, the presumption will be that the DU originated from the RCAs.

Alpha spectrometry minimum detectable concentrations will be determined for each of the three uranium isotopes, ^{234}U , ^{235}U , and ^{238}U . Inductively coupled plasma-mass spectroscopy (ICP-MS) will be used for those samples that require more precise isotope ratio determinations.

3.3.1 Groundwater

Groundwater will not be routinely collected as part of the ERMP. Supporting information is provided in Section 3.3.1.1.

3.3.1.1 Rationale for Exclusion of Routine Groundwater Sampling

The RESRAD Version 6.5 radioactivity modeling code developed by the U.S. Department of Energy was used to model the transport of site DU to the underlying aquifer. Depth to groundwater beneath the RCAs is greater than 1,001 feet below ground surface (bgs) based on exploratory drilling (U.S. Army and USACE, 2004). The transport characteristics of uranium, depth to groundwater, and other site specific factors prevent the DU contamination at the site (which is only present in the near-surface) from entering the aquifer beneath PTA for tens of thousands of years.

The modeling approach included the use of RESRAD default transport parameters and site-specific data. Results indicate that uranium does not reach the groundwater until sometime around 51,000 years. RESRAD predicted the maximum concentration of DU in the aquifer at 55,000 years to be 1×10^{-9} pCi/L, which is 11 orders of magnitude less than the 10 CFR 20 Appendix B effluent discharge limit of 300 pCi/L. The RESRAD output files are included at Appendix B to this Plan. Site-specific and model-sensitive parameters are presented below (all other parameters used were RESRAD defaults).

- Uranium isotopic activity concentrations:
 - $^{234}\text{U} = 1.23 \text{ pCi/g}$
 - $^{235}\text{U} = 0.09 \text{ pCi/g}$
 - $^{238}\text{U} = 6.60 \text{ pCi/g}$
- Depth to groundwater (unsaturated zone thickness) = 305 meters
- Annual precipitation = 0.406 meters
- Uranium $K_d = 50 \text{ mL/g}$ (RESRAD default)

- Contamination depth (thickness of contaminated zone) = 0.457 meters
- Area of contaminated zone = 10,000 square meters

Uranium Activity Concentrations and Physical Distribution

DU contamination was modeled based on data collected during site characterization and historical information. The total quantity of DU on the range was conservatively assumed to be 299 pounds, the mass of all DU that was shipped to Hawaii as part of the Davey Crockett weapons system. This is conservative because it assumes that all spotting rounds were fired at PTA; it is known that a portion of these rounds were fired at SB (Cabrera, 2008a). The 299 pounds of DU was assumed to be evenly distributed over an area of 10,000 square meters to a depth of 0.457 meters (18 inches).

Depth to Groundwater

The depth from the bottom of the contaminated zone to the top of groundwater is one of the most sensitive transport parameters in this model. The time it takes the uranium to reach the aquifer is proportional to this parameter. Groundwater at PTA is estimated at greater than 1,000 ft below the soil surface. A test hole, Pōhakuloa test hole T-20, was drilled to a depth of 1,001 ft below ground surface; no groundwater was encountered in this test hole (U.S. Army and USACE, 2004). Considering that the actual depth to groundwater is not known, 305 meters (1,001 feet) was used in the model for the purpose of conservatism.

Annual Precipitation

Precipitation water penetrates the subsurface soil system and is added to the groundwater flow system, through the process of deep percolation. It is this process that serves as the primary transport mechanism for near-surface DU to flow down towards, and ultimately into, the groundwater. The average annual precipitation at PTA is reported as 10 to 16 inches (USAG-HI, 2011). For the purpose of conservatism the upper bound of the estimated precipitation, 16 inches (0.406 meters), was entered into the model.

Uranium Distribution Coefficient (K_d)

The distribution coefficient, K_d , is the ratio of the mass of solute species adsorbed or precipitated on the solids per unit of dry mass of the soil to the solute concentration in the liquids. The distribution coefficient represents the partition of the solute in the soil matrix and soil water, assuming that equilibrium conditions exist between the soil and solution phases. The transfer of radionuclides from the liquid to the solid phase or vice versa may be controlled by mechanisms such as adsorption and precipitation, depending on the radionuclides. The RESRAD model is very

sensitive to this parameter because it dictates how much uranium is “available” for transport to groundwater through percolation.

Lower K_d values predict greater uranium mobility from soil into the groundwater, reducing the time it takes uranium to enter the aquifer. EPA has reported that the lowest observed K_d for uranium in neutral conditions to be 63 mL/g (USEPA, 2006). The RESRAD default value for uranium K_d is 50 mL/g. This value was used in the DU model and is considered reasonably conservative.

3.3.1.2 General Groundwater Information

Groundwater describes any water that is located beneath the ground surface in soil pore spaces and fractures in subsurface rock. This water is stored in an aquifer—which is defined as a porous substrate, typically an underground layer of permeable rock or unconsolidated material (e.g., sand, gravel, silt, or clay)—and may either flow naturally to the surface or be extracted using pumps or wells (USAG-HI, 2011). Rainfall is the primary source of groundwater recharge on Hawai‘i Island; additionally this island has the highest recharge rate among the Hawaiian Islands (U.S. Army and USACE, 2008a). Rainfall, fog drip and occasional frost are the main sources of water for the biological resources found on PTA.

PTA experiences an average rainfall of 10 to 16 inches annually (USAG-HI, 2011). As there is no water supply at PTA, all water must be trucked approximately 40 miles. Neither Lake Waiau nor springs which occur in Pōhakuloa Gulch are used to supply potable water to PTA.

Groundwater at PTA is estimated at greater than 1,000 ft below the soil surface. Data to evaluate groundwater at PTA is very limited at this time. The majority of PTA lies within the Northwest Mauna Loa aquifer sector, which has an estimated sustainable yield of 30 million gallons per day (HDLNR, 2008).

Based on regional hydrogeological information, it is believed that the groundwater beneath PTA occurs primarily as deep basal water within older Pleistocene age basalts (U.S. Army and USACE, 2004). Exploratory well drilling was conducted in March 1965 by the DLNR near the PTA Cantonment Area. A test hole, Pōhakuloa test hole T-20 located half mile west of Mauna Kea State Park at an elevation of 6,375 feet mean sea level (msl), was drilled to a depth of 1,001 ft below ground surface (bgs); no groundwater was encountered in this test hole (U.S. Army and USACE, 2004).

3.3.2 Surface Water

Surface water will not be routinely collected as part of the ERMP. Supporting information is provided in Section 3.3.2.1.

3.3.2.1 Rationale for Exclusion of Surface Water Sampling

Surface water will not be sampled at PTA because surface water is not present due to the porous nature of the lava covered ground surface. No surface streams, lakes or other bodies of water are within the boundaries of PTA; and no perennial streams are within 15 miles (USAG-HI, 2011).

3.3.2.2 General Surface Water Information

Surface water is generally defined as waters in a river, lake, stream or estuary. Surface water is naturally replenished by precipitation and lost through natural processes such as discharge to oceans, evaporation, and subsurface seepage. The total quantity of water in any surface water system and proportions of water lost are dependent on precipitation in its watershed, storage capacity, soil permeability, runoff characteristics of land in the watershed, timing of the precipitation, and evaporation rates. No surface streams, lakes or other bodies of water are within the boundaries of PTA; and no perennial streams are within 15 miles (24.1 km) of PTA. Lake Waiau, which is located near the summit of Mauna Kea, is approximately 8 miles (12.9 km) from the installation, and is the nearest known surface water body. However, at least seven intermittent streams drain surface water off the southwestern flank of Mauna Kea and lie within the same drainage area as PTA. Popo's Gulch is the closest stream to PTA boundaries. Popo's Gulch converges with Auwaiakeakua Gulch to drain surface water toward the Waikoloa community to the west of PTA. There are three intermittent streams located within two miles (3.2 km) of the Cantonment Area (Waikahalulu Gulch, Pōhakuloa Gulch, and an unnamed gulch, which collect runoff from the southern flank of Mauna Kea) (U.S. Army and USACE, 2008a). One perennial stream runs downstream of PTA, the Waikoloa Stream, which heads towards the Kohala Mountains, runs north parallel to State Highway 19, and discharges into Kawaihae Bay through the Waiulaula Gulch (USAG-HI, 2011). According to the 303(d) List of Impaired Waters in Hawai'i, Kawaihae harbor is identified as an impaired body of water due to turbidity, and is assigned low priority for development of total maximum daily loads (TMDLs) (HDOH, 2004).

3.3.3 Soil

Soil samples will be collected quarterly at six locations along roadway points of egress from the RCAs to ensure human activities do not transport DU beyond the RCAs. Plate Number 5 identifies the six soil sample collection locations.

3.3.3.1 Rationale for and Scope of Soil Sampling

Soil ingestion can be a significant environmental pathway with regard to dose estimates. It is possible, but unlikely, that human activities on the range in the RCAs could act as a potential

transport mechanism for DU. Soil samples will be collected annually at the locations identified on Plate Number 5.

If alpha spectrometry results at any location indicate a $^{238}\text{U} : ^{234}\text{U}$ activity ratio exceeding three, IMCOM will notify the NRC within 30 calendar days of receipt of analytical sampling results. Additional sampling will be performed as soon as practicable. Further actions may be defined based on the results of confirmatory sampling including modification of sampling frequency and/or increasing the number of sample locations. Such actions will be coordinated with the NRC.

If trending analyses indicate an increase in $^{238}\text{U} : ^{234}\text{U}$ activity ratio based on alpha spectrometry analyses and/or an increase in $^{238}\text{U} : ^{235}\text{U}$ mass ratio based on ICP-MS, the License RSO will conduct an independent review of the results. Based on the results of this review, the scope of the ERMP may be modified. Any proposed modifications will be coordinated with the NRC prior to implementation.

3.3.4 Sediment

Sediment will not be routinely collected as part of the ERMP. Supporting information is provided in Section 3.3.4.1.

3.3.4.1 Rationale for Exclusion of Sediment Sampling

As discussed in Section 3.3.2.2, no surface streams, lakes or other bodies of water are within the boundaries of PTA; and no perennial streams are within 15 miles. Thus, no sedimentary material is available for sampling.

3.3.5 Air

Air samples will not be routinely collected as part of the ERMP. Supporting information is provided in Section 3.3.5.1. Samples will be collected and analyzed, however, if prescribed burns occur to remove vegetative cover on the range in accordance with the Radiation Safety Plan. The License RSO may apply to the NRC for relief from this requirement when sufficient data are accumulated that demonstrate DU is not transported in the air during range burns.

3.3.5.1 Rationale for Exclusion of Air Sampling

Extensive survey activities have occurred at both PTA and SB and substantially more DU has been found at SB. Air sampling performed during prescribed burning in a DU-contaminated area and across larger portions of the RCA at SB did not indicate the presence of DU (Cabrera, 2008b). In addition, air sampling and analysis during 2010 BAX construction at PTA Range did not confirm even trace amounts of DU (Cabrera, 2011).

M101 spotting rounds were not designed as hard target “penetrators” and were not fired into armored vehicles or targets; therefore, no significant quantities of aerosolized, highly transportable forms of DU were created by the original impact. Typically the DU round fragmented or remained mostly intact on impact. Since then, DU fragments may have undergone oxidation (depending on soil conditions), high-explosive impact from subsequent range firing, mechanical shearing and grinding by heavy tactical equipment, and prescribed or accidental range fires. DU particulates may have been created and transported by wind, explosive impact, or flame.

In order to produce particles with an activity median aerodynamic diameter (AMAD) less than 5 μm , M101 rounds must be physically acted upon, impacted or heated to temperatures over uranium’s melting point of 700-1,000 degrees Celsius (Army Environmental Policy Institute (AEPI), 1995). The type of activities that could potentially produce DU particles in the 5- μm AMAD range are: 1) use of heavy equipment on former M101 ranges could, through mechanical grinding of M101 rounds; 2) kinetic impacts between munitions and M101 rounds; and 3) incidental range fires or prescribed burns by range personnel to control vegetation. Since the DOD has prohibited the firing of high-explosive munitions into the same areas as DU since 2004 (DOD, 2004) and will restrict access to the RCAs under the NRC license, any remaining transportable DU (AMAD < 5 μm) resulting from impacts or maneuver training is assumed to be no longer present in significant quantity. The most likely identified scenario involving the production of transportable forms of DU is the occurrence of brush fires in RCAs, whether prescribed or naturally occurring.

3.3.6 Biota

Since DU is poorly adsorbed, rapidly excreted, and does not bioaccumulate, samples will not be routinely collected of plants, vegetables, wild or farm meat, fish, or milk for DU analysis as part of the ERMP. However, in the event that sampling of other environmental media indicates significant evidence of a substantial release of DU from the RCAs, the Army will include plant, animal, and aquatic creature tissue sampling in any subsequent investigations that are undertaken.

3.3.6.1 Rationale for Exclusion of Biota Sampling

DU spotting rounds fired into the RCAs during the 1960s may have been subject to artillery bombardment (until recently), maneuvers involving heavy tracked military vehicles, precipitation, and wind and water erosion. The original insoluble DU may have been oxidized into more soluble forms. DU may have been taken up from soil by plant roots, and airborne DU may have been deposited by precipitation or wind onto plant leaves and fruit. Wild animals may

eat DU-containing food or drink DU-containing water. At some installations, farm animals such as cattle or sheep graze on military lands (though typically not in active ranges).

Human receptors include on or off post recreational hunters or fishers, consumers of meat or milk from farm animals grazing on post, and consumers of vegetables or fruit off post that is contaminated by water or airborne DU.

Though these mechanisms for a complete exposure pathway seem to be in place, the actual potential for uranium uptake, transport through, and bioaccumulation in the human food chain is small to nonexistent. Uptake by plants is typically low (USEPA, 2006a) and restricted to adsorption in the outer root membranes (ATSDR, 2011). No significant translocation of uranium from soil to aboveground parts of plants has been observed; uranium levels in higher trophic levels of aquatic organisms decline due to low assimilation efficiency; bioconcentration in fish is low and may represent water extraction or accumulation on gill surfaces (ATSDR, 2011). Since lower-level organisms tend to excrete both soluble and insoluble uranium species quickly, uranium is not effectively transported through the food chain and does not bioaccumulate (USEPA, 2006a). Army sampling of deer that were exposed to DU at Jefferson Proving Ground found no DU (SAIC, 2006).

4.0 PROJECT ORGANIZATION AND MANAGEMENT

4.1 Responsible Organizations

The key organizations supporting the environmental monitoring program include the NRC and the Department of the Army.

4.1.1 Nuclear Regulatory Commission

The NRC's primary mission is to protect the public health and safety and the environment from the effects of radiation from nuclear reactors, materials, and waste facilities. The NRC also regulates these nuclear materials and facilities to promote the common defense and security. The NRC approves and oversees the implementation of IMCOM's source material license and will ensure that the terms and conditions of the license are being implemented, including implementation of the ERMP.

4.1.2 Department of the Army

For the purpose of the ERMP, the IMCOM Commander, IMCOM Radiation Safety Staff Officer (RSSO) (the License RSO), US Army Garrison (USAG) Hawaii commander, and the garrison RSO have specific responsibilities that are described in the USAG Hawaii Radiation Safety Plans.

4.2 Lines of Authority

In general, communication to the NRC should go through the garrison RSO to the IMCOM RSSO and then to the NRC. The POC for IMCOM RSSO and the garrison RSO is as follows:

IMCOM Radiation Safety Staff Officer
US Army Installation Management Command
ATTN: IMSO/301
Building 2261
2405 Gun Shed Road
Fort Sam Houston, TX 78234-1223

Garrison Radiation Safety Officer
USAG-HI Safety Office
1554 Lyman Road (Bldg 3004)
Schofield Barracks, HI 96857

5.0 FIELD PROGRAM

This section details procedures associated with the field program. In particular, it describes protocols for sampling, sample handling and management, field measurements, equipment decontamination, and waste management.

5.1 Sampling Protocol

Procedures associated with planning and conducting sampling of the RCA environs are defined in this section. These procedures include pre-mobilization activities and environmental media sampling, field measurements, equipment decontamination, and waste management.

5.1.1 Pre-Mobilization Activities

The IMCOM RSSO will notify and coordinate with the garrison RSO 60 days prior to the sampling date to ensure that support will be onsite at the time of sampling. At this time, orders for supplies and instruments will be made. In addition, the arrangements with the analytical laboratory will be completed to support analysis of samples.

5.1.2 Groundwater Sample Collection

Groundwater sample collection and analysis is not planned as part of the ERMP. However, in the unlikely event ERMP scope is modified in the future to include groundwater sample collection, procedures for sample collection and analysis are included in this ERMP.

Standard operating procedures for groundwater sampling are enumerated (note that procedures will be modified, as appropriate, for sample collection from irrigation wells and municipal/domestic wells):

1. An exposure rate measurement will be performed and recorded approximately one meter above the ground surface in the immediate vicinity of the wellhead to document pre-sampling conditions.
2. The purging of wells will be accomplished using a submersible pump. Upon opening each well, the well cover and wellhead will be inspected for damage, and organic vapors will be monitored using a photoionization detector (PID). The static water level then will be determined using a water level indicator probe. Immediately after the water level measurement, the pump intake will be installed approximately 1 foot below the top of the water surface. Each well will be purged at a rate no greater than the recharge rate of the aquifer. The water level should be monitored during purging to ensure that drawdown is not occurring. The field parameters of hydrogen ion concentration (pH), temperature, conductivity, and turbidity will be monitored and recorded during purging using a Horiba

U-10 Water Quality Meter or equivalent. Purging will be complete after the indicator parameters have stabilized within the following ranges over three consecutive readings:

- pH $\pm$ 0.2
 - Temperature $\pm$ 1° C
 - Conductivity $\pm$ 10 percent.
3. The sampler will don new nitrile rubber gloves or similar gloves.
 4. Samples will be collected using a new hand bailer tied with new colorless twine for each sample. Care will be taken when lowering the bailer into the well to prevent unnecessary aeration or contamination of the sample.
 5. A total quantity of 1 U.S. gallon of water will be collected.
 6. A portion of the first bailer full of water will be placed into a clean beaker or other suitable container, and an evaluation of temperature, pH, and conductivity will be conducted and recorded.
 7. Sample information will be recorded.
 8. The sample will not be filtered in the field.
 9. If required by the laboratory, an appropriate quantity of hydrochloric acid will be added to the sample to ensure pH is less than or equal to 2. This can be accomplished by using sample containers provided by the lab that are pre-loaded with appropriate quantities of acid.
 10. The sample will be wiped clean so that a label and security seal may be placed on it. The sample then will be placed into a sealed Ziploc bag prior to insertion into a cooler for shipment.
- Additional forms may be used to record additional well information (e.g., well depth, purging data).
11. An exposure rate measurement will be performed and recorded approximately one meter above the ground surface in the immediate vicinity of the wellhead to document post-sampling conditions.

5.1.3 Surface Water Sample Collection

Surface water sample collection and analysis is not planned as part of the ERMP. However, if a significant release of DU off the range is identified through soil sampling, surface water sampling may be implemented. As such, procedures for sample collection and analysis are included in this ERMP.

Standard operating procedures for surface water sampling are enumerated:

1. An exposure rate measurement will be performed and recorded approximately one meter above the sample collection location to document pre-sampling conditions.
2. Samples will be collected in new sample containers using the grab method. Sample containers will be positioned pointing upstream and below the surface of the water.
3. A sample quantity of 1 U.S. gallon of water will be collected.
4. Water samples will not be filtered in the field.
5. If required by the laboratory, an appropriate quantity of hydrochloric acid will be added to the sample to ensure pH is less than or equal to 2. This can be accomplished by using sample containers provided by the lab that are pre-loaded with appropriate quantities of acid.
6. The sample will be wiped clean so that a label and security seal may be placed on it. The sample then will be placed into a sealed Ziploc bag before being put into a cooler for shipment.
7. An exposure rate measurement will be performed and recorded approximately one meter above the sample collection location to document post-sampling conditions.

5.1.4 Soil/Sediment Sample Collection

Soil samples will be collected at six locations along roadway points of egress from the RCAs. Plate Number 5 identifies the six soil sample collection locations.

Sediment samples will not be collected because there is no sediment present at PTA.

Standard operating procedures for soil sampling are enumerated below:

1. An exposure rate measurement will be performed and recorded approximately one meter above the sample collection location to document pre-sampling conditions (if not performed during surface water sampling).

2. The sampler will don clean nitrile rubber or similar gloves.
3. Samples will be collected using a new or properly cleaned scoop, trowel, or other suitable tool. Samples will be placed in a glass sample jar.
4. The sample will be wiped clean so that a label and security seal may be placed on it. The sample will then be placed into a sealed Ziploc® bag before being put into a cooler with ice.
5. An exposure rate measurement will be performed and recorded approximately one meter above the sample collection location to document post-sampling conditions.

5.2 Sample Handling and Management

Because samples collected are in support of NRC license commitments, chain of custody (COC) procedures will be followed. Samples will be secured from unauthorized access during the period of sampling. Prior to shipment of samples to the analytical laboratory, a properly completed COC Record will be placed in each shipping container. Survey personnel will maintain a copy of the COC Record for verification of sample transport. Water samples should reach the analytical laboratory no later than four days from the time of sampling. To ensure that this schedule is met and that the laboratory has time to filter and preserve the samples if necessary, water samples should be collected on the first day of the sampling trip and shipped the following day. It is not necessary to ship the water, sediments, and soils together.

Sample analysis of all environmental samples will be performed through the analytical laboratory. Samples will be managed and analyzed in accordance with the established protocols and procedures of the analytical laboratory.

5.2.1 Sample Containers

The analytical laboratory will provide sample containers and labels prior to the sampling event. Sample bags, labels, and coolers will be shipped to the following address:

Schofield Barracks Military Reservation
Garrison Radiation Safety Officer
USAG-HI Safety Office
1554 Lyman Road (Bldg 3004)
Schofield Barracks, HI 96857

5.2.2 *Quality Control Samples*

Quality control (QC) samples will be collected to achieve data quality objectives. These samples include matrix spike/matrix spike duplicate (MS/MSD), field duplicate, and field replicate samples.

MS/MSD samples will be collected to evaluate the accuracy and precision of the analysis and the matrix effect of the sample on the analytical methodology. A MS/MSD sample will be collected for every 20 samples of similar matrix received at the laboratory (minimum of one MS/MSD per sampling event). MS/MSD samples do not release the laboratory from its own QC requirements for laboratory control samples (LCSs).

A field duplicate sample is a second sample collected at the same location as the original sample. Duplicate samples are collected simultaneously or in immediate succession, using identical recovery techniques, and are treated in an identical manner during storage, transportation, and analysis. The sample containers are assigned an identification number in the field so that they cannot be identified (blind duplicate) as duplicate samples by laboratory personnel performing the analysis. Specific locations are designated for collection of field duplicate samples prior to the beginning of sample collection. Field duplicates will be collected at a ratio of 1 per 10 investigative samples collected.

A field replicate sample, also called a split, is a single sample divided into two equal parts for analysis. The sample containers are assigned an identification number in the field so that they cannot be identified as replicate samples by laboratory personnel performing the analysis. Specific locations are designated for collection of field replicate samples prior to the beginning of sample collection. Replicate sample results are used to assess precision.

5.2.3 *Sample Identification*

All sample containers will have the following information listed on the label:

- Unique sample identification
- Date and time of sample collection
- Source of sample (including name, location, and sample type)
- Designation of MS/MSD
- Preservative used
- Analyses required
- Name of collector(s).

5.2.4 Sample Custody

Procedures to ensure the custody and integrity of the samples begin at the time of sampling and continue through transport, sample receipt, preparation, analysis and storage, data generation and reporting, and sample disposal. Records concerning the custody and condition of the samples are maintained in field and laboratory records.

Chain of custody records will be maintained maintain for all field and field QC samples. A sample is defined as being under a person's custody if any of the following conditions exist: (1) it is in his/her possession, (2) it is in his/her view, after being in his/her possession, (3) it was in his/her possession and he/she locked it up, or (4) it is in a designated secure area.

All sample containers will be sealed in a manner that will prevent or allow for detection of tampering if it occurs. Furthermore, each sample will be uniquely identified, labeled, and documented in the field at the time of collection.

After samples reach the laboratory, they will be checked against information reported on the COC forms for anomalies. The condition, temperature, and appropriate preservation of the samples will be checked and documented on the COC form. The occurrence of any anomalies in the received samples and decisions regarding the potentially affected samples will be documented in laboratory records.

The laboratory will confirm sample receipt and login information through the transmission of a letter of receipt (LOR) to the garrison RSO (or designee). Within 24 hours of sample receipt, the laboratory shall send a facsimile or e-mail a copy of the completed COC form, related login information, and a report specifying the condition of the samples upon receipt.

5.3 Field Measurements

Procedures associated with field measurements are described in this section. Related equipment operation and maintenance procedures are identified.

5.3.1 Field Parameters

Request for instrumentation to support the sampling program, including field measurements, will be made no later than 30 days prior to the scheduled departure date. Radiation detection instrumentation, sampling tools, and pH, temperature, and conductivity instruments either will be rented or obtained on PTA. Specific field measurements for groundwater, surface water, and radiation doses are described in the following paragraphs.

5.3.1.1 Groundwater

When collecting the groundwater sample, the field parameters of pH, temperature, conductivity, and turbidity will be monitored and recorded during purging of groundwater wells using a Horiba U-10 Water Quality Meter or equivalent. Well purging will be complete after the indicator parameters have stabilized within the following ranges over three consecutive readings:

- pH ± 0.2
- Temperature $\pm 1^{\circ}\text{C}$
- Conductivity ± 10 percent.

Measurements of static water level will be taken prior to purging and sampling and upon completion of sampling using an electronic water level indicator. The groundwater level will be measured to the nearest 0.01 ft and from a marked survey datum on the rim of the riser. The water level measurements will be recorded on the monitor well static water level form. Wells that are dry will be noted as such. Groundwater levels will be measured in all wells to be sampled in as short a period as practical. The electronic water level indicator will be decontaminated between each monitoring well measurement.

5.3.1.2 Gamma Radiation Measurements

Radiation exposure rate measurements will be taken at 1 m above the sample location and recorded. Measurements will be performed with a portable radiation survey instrument that is sensitive to gamma radiation, reads in units of exposure rate, and is appropriately calibrated. The instrument should be held 1 m above the sampling location.

Any comments and notations that may be necessary for interpretation of the results should be recorded.

5.3.2 Equipment Calibration and Quality Control

Upon receipt of instruments, appropriate instrument QC checks will be conducted to ensure proper operation prior to departure.

Radiation detection instrumentation will be checked for response against a radiation check source. This check source also should be shipped to the survey site for instrument verification onsite. The radiation check source used need not be a calibrated source because instrument response is the parameter being evaluated. The check will be performed daily or as needed to ensure accurate and precise readings.

Water quality instruments also should be verified using the manufacturer's procedures. These instruments will be calibrated daily per the manufacturer's guidelines. More frequent calibration

may be necessary if field personnel suspect that the initial calibration may have been affected by external factors (e.g., temperature or humidity). Field measurements to be performed include water level measurement, pH, conductivity, temperature, and turbidity. All equipment to be used during the field sampling will be examined to certify that it is in operating condition. This examination will include checking the manufacturer's operating manual and instructions for each instrument to ensure that all maintenance requirements are being observed.

In the event that an internally calibrated field instrument fails to meet calibration/checkout procedures, a HOLD tag will be attached, the instrument will be returned to the supplier or manufacturer, and a backup instrument will be used in its place. Project personnel responsible for calibrating and operating field instruments will receive training in the proper use of each instrument. The satisfactory operating condition of equipment and instrumentation used onsite will be verified before each piece of equipment is shipped to PTA.

5.3.3 Equipment Maintenance and Decontamination

Decontamination operations will be conducted to reduce the potential for cross-contamination from sampling equipment that will be reused. Bailers, twine, nitrile rubber gloves, and other such disposable items will not be reused but will be disposed of properly. All reusable field equipment will be decontaminated by using potable or DI water (transported to each sampling location) before sampling activities begin, between sampling activities, and after sampling activities are completed at each site. The use of DI water will be required in the decontamination process of sampling equipment that comes into direct contact with analytical samples.

Equipment decontamination for sampling activities will include rinsing the following equipment with DI water after sampling and measurements are completed at each sample location:

- Electronic water level indicators
- Probe for the water quality meter (Horiba Model U-1).

The scoops or trowels used for soil sampling will need to be decontaminated in the following manner:

- Potable water rinse
- Scrubbed in an Alconox® detergent and potable water bath
- Potable water rinse
- DI water rinse.

All rinse water will be collected in a purge water collection vessel for proper disposal. In addition, field personnel will prevent the equipment from coming into contact with potentially

contaminating substances, such as tape, oil, engine exhaust, corroded surfaces, and dirt by wrapping tools or equipment with aluminum foil when necessary.

Decontamination operations will be conducted to reduce the potential for cross-contamination from sampling equipment and machinery.

5.4 Waste Management

Waste management (e.g., purged groundwater, equipment decontamination liquids, and disposable personal protective clothing) will be addressed on a site-by-site basis. Waste may be classified as non-investigative waste or investigation-derived waste (IDW). It is anticipated that waste generated during the sampling will be acceptable for disposal in standard sanitary dumpster. Determination of standard sanitary disposal acceptability will be based on analytical sample results and/or instrument scans of materials and equipment.

Non-investigative waste, such as litter and household garbage, will be collected on an as needed basis at each sample location in a clean and orderly manner. This waste will be containerized and transported to a PTA-designated collection bin. Acceptable containers will be sealed boxes or plastic garbage bags.

IDW generated during groundwater sampling includes purged groundwater, equipment decontamination liquids, and disposable personal protective clothing. Purged groundwater and equipment decontamination liquids will be containerized in 55-gallon drums or other suitable container. Mixing of the fluids is permissible. The containers will be labeled and transported to a secure staging area designated by PTA. In no instance will a drum containing IDW be left unattended at an unsecured location. Disposable personal protective equipment (PPE) will be placed in plastic bags and disposed of in a site dumpster. PPE will be scanned for radiological contamination prior to disposal.

After field activities are completed, a representative sample of the wastewater will be collected for analysis. The sample will be a composite composed of liquid from each container of liquid IDW. Based on the results of the analysis, an appropriate disposal option will be selected. If the water meets the discharge limits, it will be released to the ground surface and any associated sampling equipment will be disposed of or stored as radiologically clean. If water analyses indicate that levels exceed discharge limits, the water will be transported and disposed of offsite.

5.5 Record Keeping

Field records will be maintained to a sufficient level of detail to re-create all sampling and measurement activities. The requirements listed in this section apply to all measuring and sampling activities. Requirements specific to individual activities are listed in the section that

addresses each activity. The information will be recorded with indelible ink in a permanently bound notebook with sequentially numbered pages. These records will be archived in an easily accessible form and made available to the U.S. Army upon request.

The following information will be recorded for all field activities: (1) location, (2) date and time, (3) identity of people performing the activity, and (4) weather conditions. The following information will be recorded for field measurements: (1) the numerical value and units of each measurement and (2) the identity of and calibration results for each field instrument.

The following additional information will be recorded for all sampling activities: (1) sample type and sampling method, (2) the identity of each sample and depth(s), where applicable, from which it was collected, (3) the amount of each sample, (4) sample description (e.g., color, odor, clarity), (5) identification of sampling devices, and (6) identification of conditions that might affect the representativeness of a sample (e.g., refueling operations, damaged casing).

The results of a sampling event completed in support of the ERMP will be documented and provided to IMCOM. The report will include, but not necessarily be limited to, planned and actual sampling events, analytical and field results, data quality assessment results, and completed forms. A draft and a final report on the sampling event will be prepared.

6.0 SAMPLE ANALYTICAL REQUIREMENTS

6.1 Objective

The analytical objective is to determine if DU residue from Davy Crockett spotting rounds has environmentally migrated from the RCAs within munitions impact areas on Army training ranges and whether the RCAs need to be enlarged.

6.2 Analytes of Concern

Natural and DU contain the uranium isotopes ^{234}U , ^{235}U , and ^{238}U . Therefore, all samples will be analyzed for ^{234}U , ^{235}U , and ^{238}U concentrations.

Isotopic uranium activity and mass ratios will be used to determine the absence or presence of DU. The expected ^{234}U to ^{238}U activity ratio in natural uranium under secular equilibrium is 1; however, because of weathering and other geochemical and physical phenomena (e.g., alpha recoil) secular equilibrium can be disturbed (Ivanovich and Harmon, 1992; Florida Institute of Phosphate Research (FIPR) 1991; Fujikawa et al. 2011; Luo et al., undated; Wong et al., 1999). The ^{235}U to ^{238}U atom ratio is much less variable and is useful for determining the presence of DU in a given sample. Therefore, ICP-MS will be used to determine ^{235}U to ^{238}U atom ratio, when needed. See Section 6.5 for more information about when ICP-MS analysis will be performed.

For natural uranium, the $^{235}\text{U}/^{238}\text{U}$ atom ratio is 0.007253 based on the best estimates for the ^{235}U and ^{238}U values from a single terrestrial source (International Union of Pure and Applied Chemistry (IUPAC), 2011).

For DU expected from M101 spotting rounds, the $^{235}\text{U}/^{238}\text{U}$ atom ratio is 0.00203 (AEPI, 1995). The values in the AEPI report are mass fractions; they have been converted to atom fractions for easy comparison to the natural uranium value.

6.3 Analytical Laboratory Methods

Alpha spectrometry, with appropriate preparation techniques for different environmental matrices, is the primary method chosen for this project. The secondary method used to more precisely determine the isotopic uranium composition is inductively coupled plasma mass spectroscopy (ICP-MS).

In Table 2, the alpha spectrometry minimum detectable concentrations (MDCs) for the radioisotopes ^{234}U , ^{235}U , and ^{238}U are provided for each sample type.

For all soil samples, the entire sample should be processed as received. Unwanted material should be removed (e.g., vegetation, large rocks, and large chunks of metal). The entire remaining sample must be dried, ground to uniform particle size (recommend ball mill or grinder), and sieved through an ASTM U.S. Series Number 40 sieve (425- μ m particle size or smaller) prior to subsampling for analysis. The subsample will proceed through the analytical procedure for total dissolution prior to extraction chromatography.

6.4 Laboratory Method Analytical Uncertainty

All reported alpha spectrometry values will include the measurement uncertainty coverage factor of 2. Uncertainties should be propagated by the laboratory whenever possible and a detailed discussion should be available for review (National Environmental Laboratory Accreditation Conference (NELAC), 2003).

6.5 Evaluation of Laboratory Results

Figure 2 illustrates the decision process for evaluation of the analytical results. As an additional test, if no sample result for each sample type exceeds an action criteria, the sample with the highest ^{238}U result will be analyzed for the ^{235}U : ^{238}U ratio analysis by ICP-MS.

Table 2: Alpha Spectrometry MDCs and ICP-MS Detection Sensitivities

Uranium Isotope	Sample Type and Activity Concentrations					
	Soil/Sediment MDC		Water MDC		Air MDC	
	Alpha Spectrometry (pCi/g)	ICP-MS (mg/kg)	Alpha Spectrometry (pCi/L)	ICP-MS (mg/L)	Alpha Spectrometry (μ Ci/mL)	ICP-MS (mg/kg)
^{234}U	0.5	0.01	1	0.01	5×10^{-15}	0.01
^{235}U	0.5	0.01	1	0.01	6×10^{-15}	0.01
^{238}U	0.5	0.05	1	0.05	6×10^{-15}	0.05

6.6 Laboratory Quality Control

Table 3 presents the quality control that applies to volumetric samples (i.e., groundwater, surface water, soil, and sediment samples) analyzed for radiochemistry at low concentrations using the following analytical methods: Total Dissolution (Soils/Sediments), Extraction Chromatography, Alpha Spectrometry.




Figure 2: Sample Analytical Result Evaluation Decision Tree

Table 3: Laboratory Volumetric Quality Control Samples

QC Sample	Frequency/Number	Method/SOP QC Acceptance Limits	Corrective Action by Analyst	Data Quality Indicator (DQI)	Measurement Performance Criteria
Instrument Energy Calibration	Weekly	Instrument Statistical established limits	Failure should preclude use of instruments for sample analysis. Depending on the degree of failure, samples may be re-prepared/re-analyzed, or data qualified (hold times must also be considered). Contact client for discussion.	Instrument Calibration Verification (Performance Checks)	Instrument Statistical established limits
Instrument Counting Efficiency	Monthly	Instrument Statistical established limits	Failure should preclude use of instruments for sample analysis. Depending on the degree of failure, samples may be re-prepared/re-analyzed, or data qualified (hold times must also be considered). Contact client for discussion.	Instrument Calibration Verification (Performance Checks)	Instrument Statistical established limits
Instrument Reference Background	Monthly	Instrument Statistical established limits	Failure should preclude use of instruments for sample analysis. Depending on the degree of failure, samples may be re-prepared/re-analyzed, or data qualified (hold times must also be considered). Contact client for discussion.	Instrument Calibration Verification (Performance Checks)	Instrument Statistical established limits
Method blank	1 per batch or 20 samples	Less than the Minimum Detectable Concentration (MDC)	Qualify if analyte is not detected in samples. If detected, identify source of contamination before further analysis and qualify only if necessary.	Laboratory Accuracy/bias Representativeness	Less than the Minimum Detectable Concentration (MDC)
Tracer	Add tracers to all samples	Uranium-232 30 percent to 100 percent	Depending on the degree of failure, samples may be re-prepared/re-analyzed, or data qualified (hold times must also be considered). Contact client for discussion.	Accuracy/bias	Uranium-232 40 percent-100 percent

QC Sample	Frequency/Number	Method/SOP QC Acceptance Limits	Corrective Action by Analyst	Data Quality Indicator (DQI)	Measurement Performance Criteria
Laboratory Control Sample (LCS)	1 per batch or 20 samples	Uranium-238 70 percent to 130 percent	If LCS is outside control limits, contact client. Depending on degree of failure (and hold time status), re-prepare/re-analyze affected batch of samples. Qualify only if necessary.	Accuracy/bias	Uranium-238 70 percent to 130 percent
Matrix Spike (MS)	1 per batch or 20 samples	Uranium-238 70 percent to 130 percent	If the recovery falls outside the designated range and the LCS for that analyte is in control, matrix-related problems will be noted and the data will be qualified.	Accuracy/bias Matrix effects	Uranium-238 70 percent to 130 percent
Matrix Spike Duplicate (MSD)	1 per batch or 20 samples	≤30 percent RPD Uranium-238 70 percent to 130 percent	If Dup is outside control and the LCS is in control, discuss with client for possible re-analysis or data qualification.	Precision	≤30 percent RPD. Uranium-238 70 percent to 130 percent

7.0 PROGRAMMATIC QUALITY ASSURANCE

This section describes ERMP assessment and oversight functions to ensure that planned program activities are implemented as described in the ERMP and that reports are provided to apprise management of the program status and any quality issues that arise during implementation. The programmatic QA processes described will ensure that the resultant data quality is adequate for its intended use and that appropriate responses are in place to address nonconformances and deviations from the project site-specific ERMP QA requirements.

The project teams should choose assessments that identify activities with the most influence on data quality and provide information about potential problems and mistakes. Due to the fact that fixed analytical laboratories adhere to quality system requirements dictated by accrediting bodies, sampling error is generally thought to contribute the majority of the measurement error associated with project data.

Therefore, all field data generation and sample collection operations should include a thorough onsite field sampling technical systems audits (TSA) at the start of field sampling activities so that effective corrective action measures can be implemented to mitigate the extent and impact of identified sampling nonconformances.

Deviations from the ERMP can also be identified by project personnel without the benefit of formal, scheduled assessments. This section also addresses those situations and describes the process by which the need for corrective action is documented, reported, and implemented and its effectiveness assessed.

After project-specific data requirements are determined, analytical laboratories are selected which can meet these requirements. When the project-specific data requirements identify procedures or criteria that a laboratory cannot meet, subcontract laboratories are acquired. The primary analytical laboratory is responsible for acquiring and oversight of contract laboratories. An SOP should be developed and implemented that describes procedures for acquiring and monitoring analytical contract laboratories to include at a minimum:

- a. Development of a Contract Statement of Work (SOW)
- b. Quality System Documentation review
- c. Performance Evaluation Samples
- d. Pre-contract award and monitoring site visits
- e. Complete laboratory assessment and site visit
- f. Use of Ad Hoc contracts for specific analyses.

7.1 Planned Audits and Assessments

This section of the ERMP also identifies the number, frequency, and types of planned assessment activities that should be performed for each project that makes up the overall program.

Assessments should be conducted periodically throughout each project by entities internal and/or external to the project to ensure that usable data are generated.

Appropriately scheduled assessments allow management to implement corrective action measures in a timely manner, thereby minimizing the impact of nonconformance on achieving the overall program quality objectives. The project-specific quality objectives dictate the type, frequency, and extent of the assessments that should be performed.

All ERMP documents, Quality Assurance Manuals, pertinent SOPs, as well as QA Assessment SOPs, should be gathered by the QA personnel that will be performing each assessment.

Checklists should be developed from these documents ensuring all pertinent quality aspects are audited and/or assessed. These checklists should then be attached to the audit or assessment reports.

7.1.1 *Field Sampling Technical Systems Audits*

Field Sampling TSA – A thorough onsite audit will be conducted, during which sampling design, equipment, instrumentation, supplies, personnel, training, sampling procedures, chain-of-custody, sample handling and tracking, data reporting, data handling and management, data tracking and control, and data verification procedures are examined for conformance with the ERMP.

The Garrison RSO, or designee, will perform at least one field sampling TSA. As there may only be a limited window for sampling, the TSA for each sampling media will be performed as early in the sampling effort as conceivably possible so that effective corrective action measures can be implemented to mitigate the extent and impact of any identified nonconformances.

7.1.2 *Field Analytical Laboratory Technical Systems Audits*

If there is a field analytical laboratory, the garrison RSO, or designee, in conjunction with the field sampling TSAs, will perform at least one field analytical TSA on that portion of the project as well.

7.1.3 *Fixed Laboratory Technical Systems Audits*

Fixed Laboratory TSA – A thorough audit of a fixed laboratory during which the facility, equipment, instrumentation, supplies, personnel, training, analytical methods/procedures, laboratory procedures, sample handling and tracking, data reporting, data handling and

management, data tracking and control, and data verification procedures are checked for conformance with the ERMP.

Complete Fixed Laboratory TSAs are performed as a part of the analytical laboratory's accreditation/certification/registration process.

Method audits are a subset of Fixed Laboratory TSAs and consist of the auditor observing the analyst while he/she performs the analytical method on actual real-world samples to ensure the analytical laboratory SOPs are followed as required by this ERMP. These types of audits are also covered under the laboratory's Quality System requirements and are required to be performed annually. Method audits are not required for this program if the laboratory can demonstrate that a method audit for the specific analytical procedure used in support of each project conducted under the ERMP has been performed within the last year. If a method audit was not performed within the last year in support of a project, the respective laboratory QA Manager or designee will conduct a method audit for that analysis at least once during that project.

7.1.4 Data Validation Technical Systems Audit

Data Validation TSA – A thorough review of the complete Data Validation Report, including a review of the associated analytical data package deliverables (tabulated and raw data) to ensure that all required analytical data package deliverables and Data Validation Report components were provided and contain the specified information.

The Data Validation TSA also ensures that the data validation criteria specified in the ERMP were met, and the method- and laboratory-specific QC acceptance criteria specified in the ERMP were met and were appropriate for achieving the program measurement performance criteria. The Data Validation TSA also evaluates whether the project-specific measurement performance criteria and data validation criteria were appropriate for meeting the specified project goals and whether analytical measurement performance usability issues affected this achievement.

The License RSO, or designee, will perform Data Validation TSAs for data that receives third party data validation.

7.1.5 Laboratory Data Package Technical Systems Audit

Laboratory Data Package TSA – This is a type of Data Validation TSA that is limited to a review of the complete analytical data package deliverable generated by the field and/or fixed laboratory or organization to ensure that all required deliverables (tabulated and raw data) are provided and contain all the information required to reproduce all reported results. The Data Package TSA also ensures that the data verification procedures specified in the ERMP were used by the

laboratory/organization producing the analytical data package deliverable. The Data Package TSA ensures that the method- and laboratory-specific QC acceptance criteria specified in the ERMP were met and were appropriate for achieving the program and project-specific measurement performance criteria.

The Analytical Laboratory Consultant, or designee, will perform Data Package TSAs for each data package generated by the analytical laboratory.

7.1.6 Management Systems Reviews

Management System Reviews – A review of an organization or organizational subset to determine if the management structure, policies, and procedures are sufficient to ensure that an effective quality system is in place to support the generation of usable program data.

The appropriate approval authority for this program reserves the right to conduct external Management Systems Reviews of any part of the study before the start of, during, or after completion of this program.

7.2 Corrective Actions

7.2.1 Corrective Actions for Sampling Activities

The mechanism for triggering a corrective action in the field will be QC checks, internal or external Field Sampling TSA findings, and/or any other informal means of identifying system nonconformances. Personnel performing sampling and in-field measurements will verbally notify the garrison RSO of any nonconformances and the resolution will be documented in the field logbook. The sampling personnel, or designee, will also provide written reports of the results of Field Sampling and Field Analytical TSAs to the garrison RSO and the IMCOM RSSO to ensure implementation of corrective actions and for inclusion in the final reports.

If appropriate, the garrison RSO will ensure that no additional work that is dependent on the nonconforming activity is performed until the corrective actions are completed. The field personnel will be responsible for ensuring that corrective actions are initiated and documented properly. Corrective action for field measurements may include:

- Repeating the measurement to check the error
- Checking for all proper adjustments for ambient conditions such as temperature
- Checking the batteries
- Recalibrating
- Checking the calibration
- Replacing the instrument or measurement devices

- Stopping work (if necessary)

7.2.2 Corrective Action during Laboratory Activities

The mechanism for triggering a corrective action in the laboratory will be QC checks, internal or external Fixed Laboratory TSA findings, and/or any other informal means of identifying system nonconformances. Corrective action decisions during chemical analyses for each project will be based on the potential for the situation to impact the quality of the data. Any issue that directly impacts project data quality objectives will be reported immediately to the Analytical Chemistry Laboratory Quality Assurance Manager and the garrison RSO. The garrison RSO, or designee, will provide written reports of the results of Fixed Laboratory TSAs to the appropriate Laboratory QA Manager and the IMCOM RSSO. When nonconformances are identified, the appropriate Laboratory QA Manager, or designee, will respond in writing with the corrective actions taken to address the nonconformances. The analytical laboratory will have a corrective action SOP that describes how corrective actions are implemented and documented within the analytical laboratory. Contract laboratories are required to adhere to corrective action procedures in accordance with their accreditation/certification standard and the contract SOW. Corrective actions may include:

- Re-analyzing the samples, if holding time criteria permits
- Resampling and analyzing
- Evaluating and amending sampling procedures
- Evaluating and amending analytical procedures
- Accepting data and acknowledging the level of uncertainty

7.2.3 Corrective Action during Data Validation

The mechanism for triggering a corrective action during data validation will be QC checks, internal or external Data Validation TSA findings, and/or any other informal means of identifying system nonconformances. The garrison RSO, or designee, will provide written reports of the results of Data Validation TSAs to the IMCOM Laboratory Consultant to ensure implementation of corrective actions and for inclusion in QA Reports to Management. When nonconformances are identified during the data validation, the Laboratory Consultant will respond in writing identifying the corrective actions taken to address the nonconformances.

7.2.4 Corrective Action during Data Package Review

The mechanism for triggering a corrective action during the data package review will be QC checks, internal or external Data Package TSA findings, and/or any other informal means of identifying system nonconformances. The Analytical Laboratory Consultant will provide written reports of the results of Data Package TSAs to the garrison RSO the appropriate supervisor of the

analytical laboratory generating the data package to ensure implementation of corrective actions. When nonconformances are identified the appropriate supervisor of the analytical laboratory will respond in writing identifying the corrective actions taken to address the nonconformances.

The content and format of corrective action responses shall be tailored to suit the project quality objectives. In certain situations, a letter documenting specific procedural changes may be a sufficient corrective action response. Appropriate procedural changes can include, but are not limited to, additional staff training, revision of SOPs, and rescheduling of field and analytical activities to ensure holding times are met. Corrective actions that require immediate implementation to ensure that project quality objectives are met may require work to cease until those corrective actions are implemented and their effectiveness verified.

7.2.5 General Corrective Actions

Corrective actions shall also be initiated whenever project personnel identify field sampling and/or analytical problems that could potentially affect data quality and/or usability. Such incidents should be documented and resolved using the procedures and personnel for planned assessments described above.

7.3 Quality Assurance Management Reports

QA Management Reports ensure that management and stakeholders are updated on the project status and the results of all QA assessments. Efficient communication of project status and problems allows management to implement timely and effective corrective actions so that project quality objectives can be met.

The contents of QA Management Reports are discussed in the corrective action section above. Assessment checklists, requests for corrective actions letters, and copies of all corrective action response letters shall be included as attachments to the QA Management Reports.

All QA Management Reports will be included in the Final Project Report.

8.0 ENVIRONMENTAL RADIATION MONITORING REPORT

The License RSO will prepare a detailed Environmental Radiation Monitoring Report that includes recommendations when the sampling is completed. The report should contain an executive summary. The report body should present a discussion of the data collected, any deviation from the sampling plan, an interpretation of the data, conclusions and recommendations. Supporting documentation should be provided in various appendices.

Prior to any release of information regarding the Environmental Radiation Monitoring Plans or Environmental Radiation Reports, consideration should be given to the public engagement and risk communications. Potential outcomes of poor risk communication have many deleterious effects, including adverse mission outcomes.

Risk communication is a combination of information tools and communication processes that can help address the perceptions and outrage associated with a real or perceived risk. This combination of tools and processes helps demonstrate a greater respect for public perceptions, emotions, and information needs, and helps build the trust necessary to discuss issues productively and/or, when needed, deliver bad news.

Although good risk communication cannot guarantee mission success, poor communication efforts will surely derail it and adversely impact trust in Army medicine. The result will be an increased need for personnel, resources and time to repeatedly explain and in some cases, defend the Army's decisions associated with the issue.

An example report format is provided below.

- Executive Summary
 - Purpose
 - Summary
 - Conclusions
 - Recommendations
- Report Body
 - Introduction
 - Authority
 - Purpose
 - General – A brief summary of location, history, and environmental setting
 - Media Sections – Include a section for each media
 - Media Data Uncertainty, and Data Gaps

- Media Results
 - Media Results Interpretation
 - Media Summary
 - Media Conclusions
 - Media Recommendations
- Summary
- Conclusions
- Recommendations
- Appendices
 - References
 - Other supporting documentation as necessary
 - Figures

9.0 REFERENCES

- (ATSDR, 2011) *Draft Toxicological Profile for Uranium, Division of Toxicology and Environmental Medicine, Applied Toxicology Branch, Atlanta, Georgia, May 2011.*
- (Cabrera, 2008a) *Technical Memorandum, Depleted Uranium Scoping Investigations for Makua Military Reservation, Pohakuloa Training Area, Schofield Barracks Impact Area Islands of Oahu and Hawaii, Cabrera Services, Inc., April 2008.*
- (Cabrera, 2008b) *Final Technical Memorandum, Schofield Barracks Firing Range Monitoring of Air Quality During Burning of Vegetation, Wahiawa, Hawaii, Cabrera Services, Inc., April 2008.*
- (Cabrera, 2009) *Final Technical Memorandum for Pohakuloa Training Area (PTA) Aerial Surveys, The Big Island (Hawaii), Hawaii, Cabrera Services, Inc., July 2009.*
- (Cabrera, 2011) *Characterization and Removal of DU munitions Debris for Pohakuloa Training Area, Draft, Cabrera Services, Inc., August 2011.*
- (DOD, 2004) *DOD Directive 4715.11, Environmental and Explosives Safety Management on Operational Ranges Within the United States, May 10, 2004.*
- (FIPR, 1991) *Radiochemistry of Uranium-Series Isotopes in Groundwater, Publication No. 05-022-092, May 1991.*
- (Fujikawa et al., 2011) *Variation in U Isotopic Ratios in Japanese Soil and Water Samples, IRPA-10 Presentation. Available at: <http://www.irpa.net/irpa10/cdrom/00809.pdf> (Accessed 1 Sep 2011).*
- (HDLNR, 2008) *Groundwater Hydrologic Units Map of Hawai'i, Hawai'i Department of Land and Natural Resources, Honolulu, Hawai'i, October, 28 2008.*
- (HDOH, 2004) *Final 2004 List of Impaired Waters in Hawai'i Prepared under Clean Water Act §303(d), Hawai'i Department of Health, 2004. Internet Web site: <http://hawaii.gov/health/environmental/envplanning/wqm/303dpcfinal.pdf>*
- (Ivanovich and Harmon, 1992) *Uranium-series Disequilibrium: Applications to Earth, Marine, and Environmental Sciences, 2nd Edition, Oxford Science Publications, Clarendon Press, Oxford, 1992.*

(Luo et al., undated) *Decay-series disequilibrium study of in-situ, long-term radionuclide transport in water-rock systems*, <http://www.osti.gov/em52/eprints/RADIONUCLIDE.PDF> (Accessed 1 Sep 2011).

(SAIC, 2006) *Deer Tissue Sampling Results, Depleted Uranium Impact Area Site Characterization, Jefferson Proving Ground, Indiana, 2006*.

(USACE, 2007) *Archive Search Report On the Use of Cartridge, 20mm Spotting M101 for Davy Crockett Light Weapon M28, Islands of Oahu and Hawaii*, U.S. Army Corps of Engineers, St. Louis District, May 2007.

(USAEPI, 1995) *Health and Environmental Consequences of Depleted Uranium use in the Army: Technical Report*, US Army Environmental Policy Institute June 1995.

(USAG-HI, 2011) *Draft Programmatic Environmental Impact Statement, Modernization of Training Infrastructure and Construction and Operation of an Infantry Platoon Battle Area (IPBA) at Pōhakuloa Training Area (PTA), Hawai‘i*, U.S. Army Garrison-Hawai‘i, October 2011.

(U.S. Army and USACE, 2004) *Final Environmental Impact Statement, Transformation of the 2nd Brigade, 25th Infantry Division (Light) to a Stryker Brigade Combat Team in Hawai‘i*, US Army and US Army Corp of Engineers, Honolulu Engineer District, May 2004.

(U.S. Army and USACE, 2008a) *Final Environmental Impact Statement, Permanent Stationing of the 2/25th Stryker Brigade Combat Team*, US Army and US Army Environmental Command, February 2008.

(USEPA, 2000) *Data Quality Objectives Process for Hazardous Waste Site Investigations, EPA/600/R-00/007*, U.S. Environmental Protection Agency, Office of Environmental Information, January 2000.

(USEPA, 2006) *Depleted Uranium Technical Brief, EPA 402-R-06-011*, U.S. Environmental Protection Agency, December 2006.

(USEPA, 2006a) *Guidance on Systematic Planning Using the Data Quality Objectives Process, EPA/240/B-06/001*, U.S. Environmental Protection Agency, Office of Environmental Information, Washington, DC., February 2006.

(Wong et al., 1999) *Isotopic Uranium Activity Ratios in California Groundwater*, Journal AWWA, 91(4), 171-185, 1 April 1999.

APPENDIX A

PTA ERMP PLATES

APPENDIX A – PTA ERMP PLATES

NAME

ERMP PLATES


Plate 1- Vicinity Map

Plate 2 - Potential Davy Crockett Ranges

Plate 3- M101 Target Zone/Impact Areas

Plate 4- RCA's

Plate 5 – Soil Sampling Locations






Legend

 POHAKULOA TRAINING AREA (APPROXIMATE)




	U. S. ARMY CORPS OF ENGINEERS ST. LOUIS DISTRICT
POHAKULOA TRAINING AREA HILO, HAWAII ISLAND OF HAWAII VICINITY MAP	
PROJ. DATE: JAN 2011	
PLATE NO. 1	






Legend

- POHAKULOA TRAINING AREA
- TARGET ZONES/IMPACT AREAS




	U. S. ARMY CORPS OF ENGINEERS ST. LOUIS DISTRICT	
	POHAKULOA TRAINING AREA HILO, HAWAII	
M101 TARGET ZONES/IMPACT AREAS		
PROJ. DATE: JAN 2011	PLATE NO. 3	



RCAs at Pohakuloa Training Area, Island of Hawaii

Plate Number 4 – PTA RCAs



Soil Sampling location: Soil00X


Plate Number 5 – PTA ERMP Sampling Locations

APPENDIX B

PTA ERMP RESRAD REPORTS

GRAPH OF ^{238}U CONCENTRATION

CONCENTRATION: U-238, Well Water



RESRAD SUMMARY REPORT

Table of Contents

Part I: Mixture Sums and Single Radionuclide Guidelines

Dose Conversion Factor (and Related) Parameter Summary ...	2
Site-Specific Parameter Summary	5
Summary of Pathway Selections	10
Contaminated Zone and Total Dose Summary	11
Total Dose Components	
Time = 0.000E+00	12
Time = 1.000E+00	13
Time = 3.000E+00	14
Time = 1.000E+01	15
Time = 3.000E+01	16
Time = 1.000E+02	17
Time = 3.000E+02	18
Time = 1.000E+04	19
Time = 5.500E+04	20
Dose/Source Ratios Summed Over All Pathways	21
Single Radionuclide Soil Guidelines	21
Dose Per Nuclide Summed Over All Pathways	22
Soil Concentration Per Nuclide	23

Dose Conversion Factor (and Related) Parameter Summary

Dose Library: FGR 12 & FGR 11

Menu	Parameter	Current Value#	Base Case*	Parameter Name
A-1	DCF's for external ground radiation, (mrem/yr)/(pCi/g)			
A-1	Ac-227 (Source: FGR 12)	4.951E-04	4.951E-04	DCF1(1)
A-1	At-218 (Source: FGR 12)	5.847E-03	5.847E-03	DCF1(2)
A-1	Bi-210 (Source: FGR 12)	3.606E-03	3.606E-03	DCF1(3)
A-1	Bi-211 (Source: FGR 12)	2.559E-01	2.559E-01	DCF1(4)
A-1	Bi-214 (Source: FGR 12)	9.808E+00	9.808E+00	DCF1(5)
A-1	Fr-223 (Source: FGR 12)	1.980E-01	1.980E-01	DCF1(6)
A-1	Pa-231 (Source: FGR 12)	1.906E-01	1.906E-01	DCF1(7)
A-1	Pa-234 (Source: FGR 12)	1.155E+01	1.155E+01	DCF1(8)
A-1	Pa-234m (Source: FGR 12)	8.967E-02	8.967E-02	DCF1(9)
A-1	Pb-210 (Source: FGR 12)	2.447E-03	2.447E-03	DCF1(10)
A-1	Pb-211 (Source: FGR 12)	3.064E-01	3.064E-01	DCF1(11)
A-1	Pb-214 (Source: FGR 12)	1.341E+00	1.341E+00	DCF1(12)
A-1	Po-210 (Source: FGR 12)	5.231E-05	5.231E-05	DCF1(13)
A-1	Po-211 (Source: FGR 12)	4.764E-02	4.764E-02	DCF1(14)
A-1	Po-214 (Source: FGR 12)	5.138E-04	5.138E-04	DCF1(15)
A-1	Po-215 (Source: FGR 12)	1.016E-03	1.016E-03	DCF1(16)
A-1	Po-218 (Source: FGR 12)	5.642E-05	5.642E-05	DCF1(17)
A-1	Ra-223 (Source: FGR 12)	6.034E-01	6.034E-01	DCF1(18)
A-1	Ra-226 (Source: FGR 12)	3.176E-02	3.176E-02	DCF1(19)
A-1	Rn-219 (Source: FGR 12)	3.083E-01	3.083E-01	DCF1(20)
A-1	Rn-222 (Source: FGR 12)	2.354E-03	2.354E-03	DCF1(21)
A-1	Th-227 (Source: FGR 12)	5.212E-01	5.212E-01	DCF1(22)
A-1	Th-230 (Source: FGR 12)	1.209E-03	1.209E-03	DCF1(23)
A-1	Th-231 (Source: FGR 12)	3.643E-02	3.643E-02	DCF1(24)
A-1	Th-234 (Source: FGR 12)	2.410E-02	2.410E-02	DCF1(25)
A-1	Tl-207 (Source: FGR 12)	1.980E-02	1.980E-02	DCF1(26)
A-1	Tl-210 (Source: no data)	0.000E+00	-2.000E+00	DCF1(27)
A-1	U-234 (Source: FGR 12)	4.017E-04	4.017E-04	DCF1(28)
A-1	U-235 (Source: FGR 12)	7.211E-01	7.211E-01	DCF1(29)
A-1	U-238 (Source: FGR 12)	1.031E-04	1.031E-04	DCF1(30)
B-1	Dose conversion factors for inhalation, mrem/pCi:			
B-1	Ac-227+D	6.724E+00	6.700E+00	DCF2(1)
B-1	Pa-231	1.280E+00	1.280E+00	DCF2(2)
B-1	Pb-210+D	2.320E-02	1.360E-02	DCF2(3)
B-1	Ra-226+D	8.594E-03	8.580E-03	DCF2(4)
B-1	Th-230	3.260E-01	3.260E-01	DCF2(5)
B-1	U-234	1.320E-01	1.320E-01	DCF2(6)
B-1	U-235+D	1.230E-01	1.230E-01	DCF2(7)
B-1	U-238	1.180E-01	1.180E-01	DCF2(8)
B-1	U-238+D	1.180E-01	1.180E-01	DCF2(9)
D-1	Dose conversion factors for ingestion, mrem/pCi:			
D-1	Ac-227+D	1.480E-02	1.410E-02	DCF3(1)
D-1	Pa-231	1.060E-02	1.060E-02	DCF3(2)
D-1	Pb-210+D	7.276E-03	5.370E-03	DCF3(3)
D-1	Ra-226+D	1.321E-03	1.320E-03	DCF3(4)
D-1	Th-230	5.480E-04	5.480E-04	DCF3(5)
D-1	U-234	2.830E-04	2.830E-04	DCF3(6)
D-1	U-235+D	2.673E-04	2.660E-04	DCF3(7)

Dose Conversion Factor (and Related) Parameter Summary (continued)

Dose Library: FGR 12 & FGR 11

Menu	Parameter	Current Value#	Base Case*	Parameter Name
D-1	U-238	2.550E-04	2.550E-04	DCF3(8)
D-1	U-238+D	2.687E-04	2.550E-04	DCF3(9)
D-34	Food transfer factors:			
D-34	Ac-227+D , plant/soil concentration ratio, dimensionless	2.500E-03	2.500E-03	RTF(1,1)
D-34	Ac-227+D , beef/livestock-intake ratio, (pCi/kg)/(pCi/d)	2.000E-05	2.000E-05	RTF(1,2)
D-34	Ac-227+D , milk/livestock-intake ratio, (pCi/L)/(pCi/d)	2.000E-05	2.000E-05	RTF(1,3)
D-34				
D-34	Pa-231 , plant/soil concentration ratio, dimensionless	1.000E-02	1.000E-02	RTF(2,1)
D-34	Pa-231 , beef/livestock-intake ratio, (pCi/kg)/(pCi/d)	5.000E-03	5.000E-03	RTF(2,2)
D-34	Pa-231 , milk/livestock-intake ratio, (pCi/L)/(pCi/d)	5.000E-06	5.000E-06	RTF(2,3)
D-34				
D-34	Pb-210+D , plant/soil concentration ratio, dimensionless	1.000E-02	1.000E-02	RTF(3,1)
D-34	Pb-210+D , beef/livestock-intake ratio, (pCi/kg)/(pCi/d)	8.000E-04	8.000E-04	RTF(3,2)
D-34	Pb-210+D , milk/livestock-intake ratio, (pCi/L)/(pCi/d)	3.000E-04	3.000E-04	RTF(3,3)
D-34				
D-34	Ra-226+D , plant/soil concentration ratio, dimensionless	4.000E-02	4.000E-02	RTF(4,1)
D-34	Ra-226+D , beef/livestock-intake ratio, (pCi/kg)/(pCi/d)	1.000E-03	1.000E-03	RTF(4,2)
D-34	Ra-226+D , milk/livestock-intake ratio, (pCi/L)/(pCi/d)	1.000E-03	1.000E-03	RTF(4,3)
D-34				
D-34	Th-230 , plant/soil concentration ratio, dimensionless	1.000E-03	1.000E-03	RTF(5,1)
D-34	Th-230 , beef/livestock-intake ratio, (pCi/kg)/(pCi/d)	1.000E-04	1.000E-04	RTF(5,2)
D-34	Th-230 , milk/livestock-intake ratio, (pCi/L)/(pCi/d)	5.000E-06	5.000E-06	RTF(5,3)
D-34				
D-34	U-234 , plant/soil concentration ratio, dimensionless	2.500E-03	2.500E-03	RTF(6,1)
D-34	U-234 , beef/livestock-intake ratio, (pCi/kg)/(pCi/d)	3.400E-04	3.400E-04	RTF(6,2)
D-34	U-234 , milk/livestock-intake ratio, (pCi/L)/(pCi/d)	6.000E-04	6.000E-04	RTF(6,3)
D-34				
D-34	U-235+D , plant/soil concentration ratio, dimensionless	2.500E-03	2.500E-03	RTF(7,1)
D-34	U-235+D , beef/livestock-intake ratio, (pCi/kg)/(pCi/d)	3.400E-04	3.400E-04	RTF(7,2)
D-34	U-235+D , milk/livestock-intake ratio, (pCi/L)/(pCi/d)	6.000E-04	6.000E-04	RTF(7,3)
D-34				
D-34	U-238 , plant/soil concentration ratio, dimensionless	2.500E-03	2.500E-03	RTF(8,1)
D-34	U-238 , beef/livestock-intake ratio, (pCi/kg)/(pCi/d)	3.400E-04	3.400E-04	RTF(8,2)
D-34	U-238 , milk/livestock-intake ratio, (pCi/L)/(pCi/d)	6.000E-04	6.000E-04	RTF(8,3)
D-34				
D-34	U-238+D , plant/soil concentration ratio, dimensionless	2.500E-03	2.500E-03	RTF(9,1)
D-34	U-238+D , beef/livestock-intake ratio, (pCi/kg)/(pCi/d)	3.400E-04	3.400E-04	RTF(9,2)
D-34	U-238+D , milk/livestock-intake ratio, (pCi/L)/(pCi/d)	6.000E-04	6.000E-04	RTF(9,3)
D-5	Bioaccumulation factors, fresh water, L/kg:			
D-5	Ac-227+D , fish	1.500E+01	1.500E+01	BIOFAC(1,1)
D-5	Ac-227+D , crustacea and mollusks	1.000E+03	1.000E+03	BIOFAC(1,2)
D-5				
D-5	Pa-231 , fish	1.000E+01	1.000E+01	BIOFAC(2,1)
D-5	Pa-231 , crustacea and mollusks	1.100E+02	1.100E+02	BIOFAC(2,2)
D-5				
D-5	Pb-210+D , fish	3.000E+02	3.000E+02	BIOFAC(3,1)
D-5	Pb-210+D , crustacea and mollusks	1.000E+02	1.000E+02	BIOFAC(3,2)
D-5				

Dose Conversion Factor (and Related) Parameter Summary (continued)

Dose Library: FGR 12 & FGR 11

Menu	Parameter	Current Value#	Base Case*	Parameter Name
D-5	Ra-226+D , fish	5.000E+01	5.000E+01	BIOFAC(4,1)
D-5	Ra-226+D , crustacea and mollusks	2.500E+02	2.500E+02	BIOFAC(4,2)
D-5				
D-5	Th-230 , fish	1.000E+02	1.000E+02	BIOFAC(5,1)
D-5	Th-230 , crustacea and mollusks	5.000E+02	5.000E+02	BIOFAC(5,2)
D-5				
D-5	U-234 , fish	1.000E+01	1.000E+01	BIOFAC(6,1)
D-5	U-234 , crustacea and mollusks	6.000E+01	6.000E+01	BIOFAC(6,2)
D-5				
D-5	U-235+D , fish	1.000E+01	1.000E+01	BIOFAC(7,1)
D-5	U-235+D , crustacea and mollusks	6.000E+01	6.000E+01	BIOFAC(7,2)
D-5				
D-5	U-238 , fish	1.000E+01	1.000E+01	BIOFAC(8,1)
D-5	U-238 , crustacea and mollusks	6.000E+01	6.000E+01	BIOFAC(8,2)
D-5				
D-5	U-238+D , fish	1.000E+01	1.000E+01	BIOFAC(9,1)
D-5	U-238+D , crustacea and mollusks	6.000E+01	6.000E+01	BIOFAC(9,2)

#For DCF1(xxx) only, factors are for infinite depth & area. See ETFG table in Ground Pathway of Detailed Report.

*Base Case means Default.Lib w/o Associate Nuclide contributions.

Site-Specific Parameter Summary

Menu	Parameter	User Input	Default	Used by RESRAD (If different from user input)	Parameter Name
R011	Area of contaminated zone (m**2)	1.000E+04	1.000E+04	---	AREA
R011	Thickness of contaminated zone (m)	4.572E-01	2.000E+00	---	THICK0
R011	Fraction of contamination that is submerged	0.000E+00	0.000E+00	---	SUBMFRACT
R011	Length parallel to aquifer flow (m)	1.000E+02	1.000E+02	---	LCZPAQ
R011	Basic radiation dose limit (mrem/yr)	2.500E+01	3.000E+01	---	BRDL
R011	Time since placement of material (yr)	0.000E+00	0.000E+00	---	TI
R011	Times for calculations (yr)	1.000E+00	1.000E+00	---	T(2)
R011	Times for calculations (yr)	3.000E+00	3.000E+00	---	T(3)
R011	Times for calculations (yr)	1.000E+01	1.000E+01	---	T(4)
R011	Times for calculations (yr)	3.000E+01	3.000E+01	---	T(5)
R011	Times for calculations (yr)	1.000E+02	1.000E+02	---	T(6)
R011	Times for calculations (yr)	3.000E+02	3.000E+02	---	T(7)
R011	Times for calculations (yr)	1.000E+04	1.000E+03	---	T(8)
R011	Times for calculations (yr)	5.500E+04	0.000E+00	---	T(9)
R011	Times for calculations (yr)	not used	0.000E+00	---	T(10)
R012	Initial principal radionuclide (pCi/g): U-234	1.230E+00	0.000E+00	---	S1(6)
R012	Initial principal radionuclide (pCi/g): U-235	9.000E-02	0.000E+00	---	S1(7)
R012	Initial principal radionuclide (pCi/g): U-238	6.600E+00	0.000E+00	---	S1(8)
R012	Concentration in groundwater (pCi/L): U-234	not used	0.000E+00	---	W1(6)
R012	Concentration in groundwater (pCi/L): U-235	not used	0.000E+00	---	W1(7)
R012	Concentration in groundwater (pCi/L): U-238	not used	0.000E+00	---	W1(8)
R013	Cover depth (m)	0.000E+00	0.000E+00	---	COVER0
R013	Density of cover material (g/cm**3)	not used	1.500E+00	---	DENSCV
R013	Cover depth erosion rate (m/yr)	not used	1.000E-03	---	VCV
R013	Density of contaminated zone (g/cm**3)	1.500E+00	1.500E+00	---	DENSCZ
R013	Contaminated zone erosion rate (m/yr)	0.000E+00	1.000E-03	---	VCZ
R013	Contaminated zone total porosity	4.000E-01	4.000E-01	---	TPCZ
R013	Contaminated zone field capacity	2.000E-01	2.000E-01	---	FCCZ
R013	Contaminated zone hydraulic conductivity (m/yr)	1.000E+01	1.000E+01	---	HCCZ
R013	Contaminated zone b parameter	5.300E+00	5.300E+00	---	BCZ
R013	Average annual wind speed (m/sec)	2.000E+00	2.000E+00	---	WIND
R013	Humidity in air (g/m**3)	not used	8.000E+00	---	HUMID
R013	Evapotranspiration coefficient	5.000E-01	5.000E-01	---	EVAPTR
R013	Precipitation (m/yr)	4.060E-01	1.000E+00	---	PRECIP
R013	Irrigation (m/yr)	2.000E-01	2.000E-01	---	RI
R013	Irrigation mode	overhead	overhead	---	IDITCH
R013	Runoff coefficient	2.000E-01	2.000E-01	---	RUNOFF
R013	Watershed area for nearby stream or pond (m**2)	1.000E+06	1.000E+06	---	WAREA
R013	Accuracy for water/soil computations	1.000E-03	1.000E-03	---	EPS
R014	Density of saturated zone (g/cm**3)	1.500E+00	1.500E+00	---	DENSAQ
R014	Saturated zone total porosity	4.000E-01	4.000E-01	---	TPSZ
R014	Saturated zone effective porosity	2.000E-01	2.000E-01	---	EPSZ
R014	Saturated zone field capacity	2.000E-01	2.000E-01	---	FCSZ
R014	Saturated zone hydraulic conductivity (m/yr)	1.000E+02	1.000E+02	---	HCSZ
R014	Saturated zone hydraulic gradient	2.000E-02	2.000E-02	---	HGWT
R014	Saturated zone b parameter	5.300E+00	5.300E+00	---	BSZ
R014	Water table drop rate (m/yr)	1.000E-03	1.000E-03	---	VWT
R014	Well pump intake depth (m below water table)	1.000E+01	1.000E+01	---	DWIBWT

Site-Specific Parameter Summary (continued)

Menu	Parameter	User Input	Default	Used by RESRAD (If different from user input)	Parameter Name
R014	Model: Nondispersion (ND) or Mass-Balance (MB)	MB	ND	---	MODEL
R014	Well pumping rate (cm**3/yr)	not used	2.500E+02	---	UW
R015	Number of unsaturated zone strata	1	1	---	NS
R015	Unsat. zone 1, thickness (m)	3.050E+02	4.000E+00	---	H(1)
R015	Unsat. zone 1, soil density (g/cm**3)	1.500E+00	1.500E+00	---	DENSUZ(1)
R015	Unsat. zone 1, total porosity	4.000E-01	4.000E-01	---	TPUZ(1)
R015	Unsat. zone 1, effective porosity	2.000E-01	2.000E-01	---	EPUZ(1)
R015	Unsat. zone 1, field capacity	2.000E-01	2.000E-01	---	FCUZ(1)
R015	Unsat. zone 1, soil-specific b parameter	5.300E+00	5.300E+00	---	BUZ(1)
R015	Unsat. zone 1, hydraulic conductivity (m/yr)	1.000E+01	1.000E+01	---	HCUZ(1)
R016	Distribution coefficients for U-234				
R016	Contaminated zone (cm**3/g)	5.000E+01	5.000E+01	---	DCNUCC(6)
R016	Unsaturated zone 1 (cm**3/g)	5.000E+01	5.000E+01	---	DCNUCU(6,1)
R016	Saturated zone (cm**3/g)	5.000E+01	5.000E+01	---	DCNUCS(6)
R016	Leach rate (/yr)	0.000E+00	0.000E+00	7.621E-03	ALEACH(6)
R016	Solubility constant	0.000E+00	0.000E+00	not used	SOLUBK(6)
R016	Distribution coefficients for U-235				
R016	Contaminated zone (cm**3/g)	5.000E+01	5.000E+01	---	DCNUCC(7)
R016	Unsaturated zone 1 (cm**3/g)	5.000E+01	5.000E+01	---	DCNUCU(7,1)
R016	Saturated zone (cm**3/g)	5.000E+01	5.000E+01	---	DCNUCS(7)
R016	Leach rate (/yr)	0.000E+00	0.000E+00	7.621E-03	ALEACH(7)
R016	Solubility constant	0.000E+00	0.000E+00	not used	SOLUBK(7)
R016	Distribution coefficients for U-238				
R016	Contaminated zone (cm**3/g)	5.000E+01	5.000E+01	---	DCNUCC(8)
R016	Unsaturated zone 1 (cm**3/g)	5.000E+01	5.000E+01	---	DCNUCU(8,1)
R016	Saturated zone (cm**3/g)	5.000E+01	5.000E+01	---	DCNUCS(8)
R016	Leach rate (/yr)	0.000E+00	0.000E+00	7.621E-03	ALEACH(8)
R016	Solubility constant	0.000E+00	0.000E+00	not used	SOLUBK(8)
R016	Distribution coefficients for daughter Ac-227				
R016	Contaminated zone (cm**3/g)	2.000E+01	2.000E+01	---	DCNUCC(1)
R016	Unsaturated zone 1 (cm**3/g)	2.000E+01	2.000E+01	---	DCNUCU(1,1)
R016	Saturated zone (cm**3/g)	2.000E+01	2.000E+01	---	DCNUCS(1)
R016	Leach rate (/yr)	0.000E+00	0.000E+00	1.894E-02	ALEACH(1)
R016	Solubility constant	0.000E+00	0.000E+00	not used	SOLUBK(1)
R016	Distribution coefficients for daughter Pa-231				
R016	Contaminated zone (cm**3/g)	5.000E+01	5.000E+01	---	DCNUCC(2)
R016	Unsaturated zone 1 (cm**3/g)	5.000E+01	5.000E+01	---	DCNUCU(2,1)
R016	Saturated zone (cm**3/g)	5.000E+01	5.000E+01	---	DCNUCS(2)
R016	Leach rate (/yr)	0.000E+00	0.000E+00	7.621E-03	ALEACH(2)
R016	Solubility constant	0.000E+00	0.000E+00	not used	SOLUBK(2)

Site-Specific Parameter Summary (continued)

Menu	Parameter	User Input	Default	Used by RESRAD (If different from user input)	Parameter Name
R016	Distribution coefficients for daughter Pb-210				
R016	Contaminated zone (cm**3/g)	1.000E+02	1.000E+02	---	DCNUCC(3)
R016	Unsaturated zone 1 (cm**3/g)	1.000E+02	1.000E+02	---	DCNUCU(3,1)
R016	Saturated zone (cm**3/g)	1.000E+02	1.000E+02	---	DCNUCS(3)
R016	Leach rate (/yr)	0.000E+00	0.000E+00	3.818E-03	ALEACH(3)
R016	Solubility constant	0.000E+00	0.000E+00	not used	SOLUBK(3)
R016	Distribution coefficients for daughter Ra-226				
R016	Contaminated zone (cm**3/g)	7.000E+01	7.000E+01	---	DCNUCC(4)
R016	Unsaturated zone 1 (cm**3/g)	7.000E+01	7.000E+01	---	DCNUCU(4,1)
R016	Saturated zone (cm**3/g)	7.000E+01	7.000E+01	---	DCNUCS(4)
R016	Leach rate (/yr)	0.000E+00	0.000E+00	5.450E-03	ALEACH(4)
R016	Solubility constant	0.000E+00	0.000E+00	not used	SOLUBK(4)
R016	Distribution coefficients for daughter Th-230				
R016	Contaminated zone (cm**3/g)	6.000E+04	6.000E+04	---	DCNUCC(5)
R016	Unsaturated zone 1 (cm**3/g)	6.000E+04	6.000E+04	---	DCNUCU(5,1)
R016	Saturated zone (cm**3/g)	6.000E+04	6.000E+04	---	DCNUCS(5)
R016	Leach rate (/yr)	0.000E+00	0.000E+00	6.377E-06	ALEACH(5)
R016	Solubility constant	0.000E+00	0.000E+00	not used	SOLUBK(5)
R017	Inhalation rate (m**3/yr)	not used	8.400E+03	---	INHALR
R017	Mass loading for inhalation (g/m**3)	not used	1.000E-04	---	MLINH
R017	Exposure duration	3.000E+01	3.000E+01	---	ED
R017	Shielding factor, inhalation	not used	4.000E-01	---	SHF3
R017	Shielding factor, external gamma	not used	7.000E-01	---	SHF1
R017	Fraction of time spent indoors	not used	5.000E-01	---	FIND
R017	Fraction of time spent outdoors (on site)	not used	2.500E-01	---	FOTD
R017	Shape factor flag, external gamma	not used	1.000E+00	>0 shows circular AREA.	FS
R017	Radii of shape factor array (used if FS = -1):				
R017	Outer annular radius (m), ring 1:	not used	5.000E+01	---	RAD_SHAPE(1)
R017	Outer annular radius (m), ring 2:	not used	7.071E+01	---	RAD_SHAPE(2)
R017	Outer annular radius (m), ring 3:	not used	0.000E+00	---	RAD_SHAPE(3)
R017	Outer annular radius (m), ring 4:	not used	0.000E+00	---	RAD_SHAPE(4)
R017	Outer annular radius (m), ring 5:	not used	0.000E+00	---	RAD_SHAPE(5)
R017	Outer annular radius (m), ring 6:	not used	0.000E+00	---	RAD_SHAPE(6)
R017	Outer annular radius (m), ring 7:	not used	0.000E+00	---	RAD_SHAPE(7)
R017	Outer annular radius (m), ring 8:	not used	0.000E+00	---	RAD_SHAPE(8)
R017	Outer annular radius (m), ring 9:	not used	0.000E+00	---	RAD_SHAPE(9)
R017	Outer annular radius (m), ring 10:	not used	0.000E+00	---	RAD_SHAPE(10)
R017	Outer annular radius (m), ring 11:	not used	0.000E+00	---	RAD_SHAPE(11)
R017	Outer annular radius (m), ring 12:	not used	0.000E+00	---	RAD_SHAPE(12)

Site-Specific Parameter Summary (continued)

Menu	Parameter	User Input	Default	Used by RESRAD (If different from user input)	Parameter Name
R017	Fractions of annular areas within AREA:				
R017	Ring 1	not used	1.000E+00	---	FRACA(1)
R017	Ring 2	not used	2.732E-01	---	FRACA(2)
R017	Ring 3	not used	0.000E+00	---	FRACA(3)
R017	Ring 4	not used	0.000E+00	---	FRACA(4)
R017	Ring 5	not used	0.000E+00	---	FRACA(5)
R017	Ring 6	not used	0.000E+00	---	FRACA(6)
R017	Ring 7	not used	0.000E+00	---	FRACA(7)
R017	Ring 8	not used	0.000E+00	---	FRACA(8)
R017	Ring 9	not used	0.000E+00	---	FRACA(9)
R017	Ring 10	not used	0.000E+00	---	FRACA(10)
R017	Ring 11	not used	0.000E+00	---	FRACA(11)
R017	Ring 12	not used	0.000E+00	---	FRACA(12)
R018	Fruits, vegetables and grain consumption (kg/yr)	not used	1.600E+02	---	DIET(1)
R018	Leafy vegetable consumption (kg/yr)	not used	1.400E+01	---	DIET(2)
R018	Milk consumption (L/yr)	not used	9.200E+01	---	DIET(3)
R018	Meat and poultry consumption (kg/yr)	not used	6.300E+01	---	DIET(4)
R018	Fish consumption (kg/yr)	not used	5.400E+00	---	DIET(5)
R018	Other seafood consumption (kg/yr)	not used	9.000E-01	---	DIET(6)
R018	Soil ingestion rate (g/yr)	not used	3.650E+01	---	SOIL
R018	Drinking water intake (L/yr)	5.100E+02	5.100E+02	---	DWI
R018	Contamination fraction of drinking water	1.000E+00	1.000E+00	---	FDW
R018	Contamination fraction of household water	not used	1.000E+00	---	FHHW
R018	Contamination fraction of livestock water	not used	1.000E+00	---	FLW
R018	Contamination fraction of irrigation water	not used	1.000E+00	---	FIRW
R018	Contamination fraction of aquatic food	not used	5.000E-01	---	FR9
R018	Contamination fraction of plant food	not used	-1	---	FPLANT
R018	Contamination fraction of meat	not used	-1	---	FMEAT
R018	Contamination fraction of milk	not used	-1	---	FMILK
R019	Livestock fodder intake for meat (kg/day)	not used	6.800E+01	---	LFI5
R019	Livestock fodder intake for milk (kg/day)	not used	5.500E+01	---	LFI6
R019	Livestock water intake for meat (L/day)	not used	5.000E+01	---	LWI5
R019	Livestock water intake for milk (L/day)	not used	1.600E+02	---	LWI6
R019	Livestock soil intake (kg/day)	not used	5.000E-01	---	LSI
R019	Mass loading for foliar deposition (g/m**3)	not used	1.000E-04	---	MLFD
R019	Depth of soil mixing layer (m)	not used	1.500E-01	---	DM
R019	Depth of roots (m)	not used	9.000E-01	---	DROOT
R019	Drinking water fraction from ground water	1.000E+00	1.000E+00	---	FGWDW
R019	Household water fraction from ground water	not used	1.000E+00	---	FGWHH
R019	Livestock water fraction from ground water	not used	1.000E+00	---	FGWLW
R019	Irrigation fraction from ground water	not used	1.000E+00	---	FGWIR
R19B	Wet weight crop yield for Non-Leafy (kg/m**2)	not used	7.000E-01	---	YV(1)
R19B	Wet weight crop yield for Leafy (kg/m**2)	not used	1.500E+00	---	YV(2)
R19B	Wet weight crop yield for Fodder (kg/m**2)	not used	1.100E+00	---	YV(3)
R19B	Growing Season for Non-Leafy (years)	not used	1.700E-01	---	TE(1)
R19B	Growing Season for Leafy (years)	not used	2.500E-01	---	TE(2)
R19B	Growing Season for Fodder (years)	not used	8.000E-02	---	TE(3)
R19B	Translocation Factor for Non-Leafy	not used	1.000E-01	---	TIV(1)

Site-Specific Parameter Summary (continued)

Menu	Parameter	User Input	Default	Used by RESRAD (If different from user input)	Parameter Name
R19B	Translocation Factor for Leafy	not used	1.000E+00	---	TIV(2)
R19B	Translocation Factor for Fodder	not used	1.000E+00	---	TIV(3)
R19B	Dry Foliar Interception Fraction for Non-Leafy	not used	2.500E-01	---	RDRY(1)
R19B	Dry Foliar Interception Fraction for Leafy	not used	2.500E-01	---	RDRY(2)
R19B	Dry Foliar Interception Fraction for Fodder	not used	2.500E-01	---	RDRY(3)
R19B	Wet Foliar Interception Fraction for Non-Leafy	not used	2.500E-01	---	RWET(1)
R19B	Wet Foliar Interception Fraction for Leafy	not used	2.500E-01	---	RWET(2)
R19B	Wet Foliar Interception Fraction for Fodder	not used	2.500E-01	---	RWET(3)
R19B	Weathering Removal Constant for Vegetation	not used	2.000E+01	---	WLAM
C14	C-12 concentration in water (g/cm**3)	not used	2.000E-05	---	C12WTR
C14	C-12 concentration in contaminated soil (g/g)	not used	3.000E-02	---	C12CZ
C14	Fraction of vegetation carbon from soil	not used	2.000E-02	---	CSOIL
C14	Fraction of vegetation carbon from air	not used	9.800E-01	---	CAIR
C14	C-14 evasion layer thickness in soil (m)	not used	3.000E-01	---	DMC
C14	C-14 evasion flux rate from soil (1/sec)	not used	7.000E-07	---	EVSNI
C14	C-12 evasion flux rate from soil (1/sec)	not used	1.000E-10	---	REVSNI
C14	Fraction of grain in beef cattle feed	not used	8.000E-01	---	AVFG4
C14	Fraction of grain in milk cow feed	not used	2.000E-01	---	AVFG5
STOR	Storage times of contaminated foodstuffs (days):				
STOR	Fruits, non-leafy vegetables, and grain	1.400E+01	1.400E+01	---	STOR_T(1)
STOR	Leafy vegetables	1.000E+00	1.000E+00	---	STOR_T(2)
STOR	Milk	1.000E+00	1.000E+00	---	STOR_T(3)
STOR	Meat and poultry	2.000E+01	2.000E+01	---	STOR_T(4)
STOR	Fish	7.000E+00	7.000E+00	---	STOR_T(5)
STOR	Crustacea and mollusks	7.000E+00	7.000E+00	---	STOR_T(6)
STOR	Well water	1.000E+00	1.000E+00	---	STOR_T(7)
STOR	Surface water	1.000E+00	1.000E+00	---	STOR_T(8)
STOR	Livestock fodder	4.500E+01	4.500E+01	---	STOR_T(9)
R021	Thickness of building foundation (m)	not used	1.500E-01	---	FLOOR1
R021	Bulk density of building foundation (g/cm**3)	not used	2.400E+00	---	DENSEFL
R021	Total porosity of the cover material	not used	4.000E-01	---	TPCV
R021	Total porosity of the building foundation	not used	1.000E-01	---	TPFL
R021	Volumetric water content of the cover material	not used	5.000E-02	---	PH2OCV
R021	Volumetric water content of the foundation	not used	3.000E-02	---	PH2OFL
R021	Diffusion coefficient for radon gas (m/sec):				
R021	in cover material	not used	2.000E-06	---	DIFCV
R021	in foundation material	not used	3.000E-07	---	DIFFL
R021	in contaminated zone soil	not used	2.000E-06	---	DIFCZ
R021	Radon vertical dimension of mixing (m)	not used	2.000E+00	---	HMIX
R021	Average building air exchange rate (1/hr)	not used	5.000E-01	---	REXG
R021	Height of the building (room) (m)	not used	2.500E+00	---	HRM
R021	Building interior area factor	not used	0.000E+00	---	FAI
R021	Building depth below ground surface (m)	not used	-1.000E+00	---	DMFL
R021	Emanating power of Rn-222 gas	not used	2.500E-01	---	EMANA(1)
R021	Emanating power of Rn-220 gas	not used	1.500E-01	---	EMANA(2)
TITL	Number of graphical time points	32	---	---	NPTS
TITL	Maximum number of integration points for dose	17	---	---	LYMAX
TITL	Maximum number of integration points for risk	257	---	---	KYMAX

Summary of Pathway Selections

Pathway	User Selection
1 -- external gamma	suppressed
2 -- inhalation (w/o radon)	suppressed
3 -- plant ingestion	suppressed
4 -- meat ingestion	suppressed
5 -- milk ingestion	suppressed
6 -- aquatic foods	suppressed
7 -- drinking water	active
8 -- soil ingestion	suppressed
9 -- radon	suppressed
Find peak pathway doses	suppressed

Total Dose TDOSE(t), mrem/yr

Basic Radiation Dose Limit = 2.500E+01 mrem/yr

Total Mixture Sum $M(t)$ = Fraction of Basic Dose Limit Received at Time (t)

[illegible]

Maximum TDOSE(t): 1.107E-01 mrem/yr at t = 5.500E+04 years

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 0.000E+00 years

Water Independent Pathways (Inhalation excludes radon)

Radio- Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
U-234	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
U-235	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
U-238	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 0.000E+00 years

Water Dependent Pathways

Radio- Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
U-234	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
U-235	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
U-238	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000

*Sum of all water independent and dependent pathways.

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
 As mrem/yr and Fraction of Total Dose At t = 1.000E+00 years

Water Independent Pathways (Inhalation excludes radon)

Radio- Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
U-234	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
U-235	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
U-238	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
 As mrem/yr and Fraction of Total Dose At t = 1.000E+00 years

Water Dependent Pathways

Radio- Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
U-234	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
U-235	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
U-238	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000

*Sum of all water independent and dependent pathways.

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)

As mrem/yr and Fraction of Total Dose At t = 3.000E+00 years

Water Independent Pathways (Inhalation excludes radon)

Radio- Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
U-234	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
U-235	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
U-238	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)

As mrem/yr and Fraction of Total Dose At t = 3.000E+00 years

Water Dependent Pathways

Radio- Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
U-234	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
U-235	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
U-238	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000

*Sum of all water independent and dependent pathways.

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
 As mrem/yr and Fraction of Total Dose At t = 1.000E+01 years

Water Independent Pathways (Inhalation excludes radon)

Radio- Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
U-234	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
U-235	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
U-238	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
 As mrem/yr and Fraction of Total Dose At t = 1.000E+01 years

Water Dependent Pathways

Radio- Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
U-234	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
U-235	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
U-238	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000

*Sum of all water independent and dependent pathways.

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
 As mrem/yr and Fraction of Total Dose At t = 3.000E+01 years

Water Independent Pathways (Inhalation excludes radon)

Radio- Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
U-234	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
U-235	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
U-238	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
 As mrem/yr and Fraction of Total Dose At t = 3.000E+01 years

Water Dependent Pathways

Radio- Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
U-234	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
U-235	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
U-238	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000

*Sum of all water independent and dependent pathways.

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+02 years

Water Independent Pathways (Inhalation excludes radon)

Radio- Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
U-234	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
U-235	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
U-238	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+02 years

Water Dependent Pathways

Radio- Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
U-234	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
U-235	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
U-238	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000

*Sum of all water independent and dependent pathways.

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 3.000E+02 years

Water Independent Pathways (Inhalation excludes radon)

Radio- Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
U-234	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
U-235	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
U-238	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 3.000E+02 years

Water Dependent Pathways

Radio- Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
U-234	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
U-235	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
U-238	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000

*Sum of all water independent and dependent pathways.

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+04 years

Water Independent Pathways (Inhalation excludes radon)

Radio- Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
U-234	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
U-235	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
U-238	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
As mrem/yr and Fraction of Total Dose At t = 1.000E+04 years

Water Dependent Pathways

Radio- Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
U-234	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
U-235	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
U-238	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000

*Sum of all water independent and dependent pathways.

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
 As mrem/yr and Fraction of Total Dose At t = 5.500E+04 years

Water Independent Pathways (Inhalation excludes radon)

Radio- Nuclide	Ground		Inhalation		Radon		Plant		Meat		Milk		Soil	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
U-234	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
U-235	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
U-238	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000
Total	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000

Total Dose Contributions TDOSE(i,p,t) for Individual Radionuclides (i) and Pathways (p)
 As mrem/yr and Fraction of Total Dose At t = 5.500E+04 years

Water Dependent Pathways

Radio- Nuclide	Water		Fish		Radon		Plant		Meat		Milk		All Pathways*	
	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.	mrem/yr	fract.
U-234	5.558E-02	0.5019	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	5.558E-02	0.5019
U-235	1.755E-10	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.755E-10	0.0000
U-238	5.516E-02	0.4981	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	5.516E-02	0.4981
Total	1.107E-01	1.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	0.000E+00	0.0000	1.107E-01	1.0000

*Sum of all water independent and dependent pathways.

Dose/Source Ratios Summed Over All Pathways
 Parent and Progeny Principal Radionuclide Contributions Indicated

Parent (i)	Product (j)	Thread Fraction	DSR(j,t) At Time in Years (mrem/yr)/(pCi/g)								
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04
U-234	U-234	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	2.083E-11
U-234	Th-230	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	4.503E-06
U-234	Ra-226+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	9.299E-03
U-234	Pb-210+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	3.589E-02
U-234	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	4.519E-02
U-235+D	U-235+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	2.299E-11
U-235+D	Pa-231	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	6.270E-10
U-235+D	Ac-227+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	1.301E-09
U-235+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	1.951E-09
U-238	U-238	5.400E-05	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	1.184E-15
U-238+D	U-238+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	2.311E-11
U-238+D	U-234	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	3.514E-12
U-238+D	Th-230	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	8.560E-07
U-238+D	Ra-226+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	1.720E-03
U-238+D	Pb-210+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	6.637E-03
U-238+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	8.358E-03

The DSR includes contributions from associated (half-life ≤ 180 days) daughters.

Single Radionuclide Soil Guidelines G(i,t) in pCi/g
 Basic Radiation Dose Limit = 2.500E+01 mrem/yr

Nuclide (i)	t = 0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04
U-234	*6.247E+09	*6.247E+09	*6.247E+09	*6.247E+09	*6.247E+09	*6.247E+09	*6.247E+09	*6.247E+09	5.532E+02
U-235	*2.161E+06	*2.161E+06	*2.161E+06	*2.161E+06	*2.161E+06	*2.161E+06	*2.161E+06	*2.161E+06	*2.161E+06
U-238	*3.361E+05	*3.361E+05	*3.361E+05	*3.361E+05	*3.361E+05	*3.361E+05	*3.361E+05	*3.361E+05	2.991E+03

*At specific activity limit

Summed Dose/Source Ratios DSR(i,t) in (mrem/yr)/(pCi/g)
 and Single Radionuclide Soil Guidelines G(i,t) in pCi/g
 at t_{min} = time of minimum single radionuclide soil guideline
 and at t_{max} = time of maximum total dose = 5.500E+04 years

Nuclide (i)	Initial (pCi/g)	t _{min} (years)	DSR(i,t _{min})	G(i,t _{min}) (pCi/g)	DSR(i,t _{max})	G(i,t _{max}) (pCi/g)
U-234	1.230E+00	5.500E+04	4.519E-02	5.532E+02	4.519E-02	5.532E+02
U-235	9.000E-02	5.500E+04	1.951E-09	*2.161E+06	1.951E-09	*2.161E+06
U-238	6.600E+00	5.500E+04	8.358E-03	2.991E+03	8.358E-03	2.991E+03

*At specific activity limit

Individual Nuclide Dose Summed Over All Pathways
 Parent Nuclide and Branch Fraction Indicated

Nuclide (j)	Parent (i)	THF(i)	DOSE(j,t), mrem/yr									
			t= 0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04	
U-234	U-234	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	2.562E-11
U-234	U-238	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	2.319E-11
U-234	ΣDOSE(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	4.881E-11
Th-230	U-234	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	5.539E-06
Th-230	U-238	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	5.650E-06
Th-230	ΣDOSE(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	1.119E-05
Ra-226	U-234	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	1.144E-02
Ra-226	U-238	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	1.135E-02
Ra-226	ΣDOSE(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	2.279E-02
Pb-210	U-234	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	4.414E-02
Pb-210	U-238	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	4.380E-02
Pb-210	ΣDOSE(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	8.795E-02
U-235	U-235	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	2.070E-12
Pa-231	U-235	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	5.643E-11
Ac-227	U-235	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	1.170E-10
U-238	U-238	5.400E-05	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	7.817E-15
U-238	U-238	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	1.525E-10
U-238	ΣDOSE(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	1.525E-10

THF(i) is the thread fraction of the parent nuclide.

Individual Nuclide Soil Concentration
 Parent Nuclide and Branch Fraction Indicated

Nuclide (j)	Parent (i)	THF(i)	S(j,t), pCi/g									
			t=	0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04
U-234	U-234	1.000E+00		1.230E+00	1.221E+00	1.202E+00	1.140E+00	9.785E-01	5.738E-01	1.249E-01	9.524E-34	0.000E+00
U-234	U-238	9.999E-01		0.000E+00	1.857E-05	5.486E-05	1.734E-04	4.466E-04	8.730E-04	5.702E-04	1.469E-34	0.000E+00
U-234	ΣS(j):			1.230E+00	1.221E+00	1.202E+00	1.140E+00	9.790E-01	5.747E-01	1.255E-01	1.099E-33	0.000E+00
Th-230	U-234	1.000E+00		0.000E+00	1.103E-05	3.284E-05	1.066E-04	2.968E-04	7.741E-04	1.301E-03	1.248E-03	6.246E-04
Th-230	U-238	9.999E-01		0.000E+00	8.378E-11	7.464E-10	8.005E-09	6.516E-08	5.148E-07	1.927E-06	2.495E-06	1.249E-06
Th-230	ΣS(j):			0.000E+00	1.103E-05	3.284E-05	1.066E-04	2.969E-04	7.746E-04	1.303E-03	1.250E-03	6.258E-04
Ra-226	U-234	1.000E+00		0.000E+00	2.388E-09	2.130E-08	2.293E-07	1.888E-06	1.549E-05	6.333E-05	9.212E-05	4.611E-05
Ra-226	U-238	9.999E-01		0.000E+00	1.210E-14	3.232E-13	1.154E-11	2.804E-10	7.241E-09	7.406E-08	1.842E-07	9.222E-08
Ra-226	ΣS(j):			0.000E+00	2.388E-09	2.130E-08	2.293E-07	1.888E-06	1.550E-05	6.340E-05	9.231E-05	4.620E-05
Pb-210	U-234	1.000E+00		0.000E+00	2.455E-11	6.471E-10	2.206E-08	4.739E-07	8.650E-06	5.108E-05	8.208E-05	4.108E-05
Pb-210	U-238	9.999E-01		0.000E+00	9.344E-17	7.402E-15	8.458E-13	5.527E-11	3.427E-09	5.590E-08	1.641E-07	8.217E-08
Pb-210	ΣS(j):			0.000E+00	2.455E-11	6.471E-10	2.206E-08	4.740E-07	8.653E-06	5.114E-05	8.224E-05	4.117E-05
U-235	U-235	1.000E+00		9.000E-02	8.932E-02	8.797E-02	8.340E-02	7.161E-02	4.200E-02	9.147E-03	7.169E-35	0.000E+00
Pa-231	U-235	1.000E+00		0.000E+00	1.890E-06	5.583E-06	1.764E-05	4.544E-05	8.877E-05	5.788E-05	1.367E-35	0.000E+00
Ac-227	U-235	1.000E+00		0.000E+00	2.965E-08	2.555E-07	2.445E-06	1.472E-05	5.053E-05	3.941E-05	1.006E-35	0.000E+00
U-238	U-238	5.400E-05		3.564E-04	3.537E-04	3.483E-04	3.302E-04	2.836E-04	1.663E-04	3.622E-05	2.839E-37	0.000E+00
U-238	U-238	9.999E-01		6.600E+00	6.550E+00	6.450E+00	6.115E+00	5.251E+00	3.080E+00	6.707E-01	5.257E-33	0.000E+00
U-238	ΣS(j):			6.600E+00	6.550E+00	6.451E+00	6.116E+00	5.251E+00	3.080E+00	6.708E-01	5.257E-33	0.000E+00

THF(i) is the thread fraction of the parent nuclide.

RESRAD.EXE execution time = 1.79 seconds

RESRAD DETAILED REPORT

Table of Contents

Part II: Source Terms, Factors, and Parameters for Individual Pathways

Source Factors for Ingrowth and Decay	
Radioactivity Only	3
Combined Radioactivity and Leaching	3
Ground Pathway	
Source Term Parameters	4
Time Dependence of Source Geometry	4
Occupancy, Cover/Depth, and Area Factors	6
Dose Conversion and Environmental Transport Factors .	8
Dose/Source Ratios	9
Inhalation Pathway (radon excluded)	
Dose/Source Ratios	10
Pathway Factors	10
Dose Conversion and Environmental Transport Factors .	11
Radon Pathway	
Flux and Parameters	12
Concentration and Parameters	13
Working Levels	14
Dose/Source Ratios	15
Groundwater and Surface Water Pathway Segments	
Transport Time Parameters for Unsaturated Zone Strata	17
Dilution Factor Parameters for Mass-Balance (MB) Mode	18..
Primary Parameters Used to Calculate Ratios	18
Water/Soil Concentration Ratios	19

Table of Contents (cont.)

Part II: Source Terms, Factors, and Parameters for Individual Pathways

Food Pathways

Storage Times for Contaminated Foodstuffs	20
Storage Time Ingrowth and Decay Factors	20
Storage Correction Factors	
Drinking Water	21
Irrigation Water	21
Livestock Water	23
Plants	24
Livestock Fodder	25
Meat and Milk	26
Fish and Crustacea	27
Area and Depth Factors	28
Dose Conversion and Environmental Transport Factors	
Plant	30
Meat	32
Milk	35
Fish	38
Drinking Water	38
Dose/Source Ratios	
Plant	39
Plant Total	43
Meat	44
Meat Total	49
Milk	50
Milk Total	55
Fish	56
Drinking Water	57
Concentration Ratios	
Plant/Air and Plant/Water	58
Plant/Soil	58
Meat/Fodder, Fodder/Air, Fodder/Water	61
Fodder/Soil	62
Meat/Soil	64
Milk/Soil	66

Soil Ingestion Pathway

Dose/Source Ratios.....	68
Dose Conversion and Environmental Transport Factors .	68

Source Factors for Ingrowth and Decay
 Radioactivity Factors Only
 Parent and Progeny Principal Radionuclide Contributions Indicated

Parent (i)	Product (j)	Thread Fraction	ID(j,t) = THF(j)*S1(j,t)/S1(i,0) At Time in Years									
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04	
U-234	U-234	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	9.999E-01	9.997E-01	9.991E-01	9.720E-01	8.556E-01	
U-234	Th-230	1.000E+00	0.000E+00	9.002E-06	2.701E-05	9.001E-05	2.700E-04	8.997E-04	2.696E-03	8.486E-02	3.593E-01	
U-234	Ra-226+D	1.000E+00	0.000E+00	1.950E-09	1.754E-08	1.947E-07	1.747E-06	1.921E-05	1.679E-04	6.618E-02	3.486E-01	
U-234	Pb-210+D	1.000E+00	0.000E+00	2.004E-11	5.328E-10	1.870E-08	4.373E-07	1.068E-05	1.363E-04	6.592E-02	3.485E-01	
U-235+D	U-235+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	9.999E-01	
U-235+D	Pa-231	1.000E+00	0.000E+00	2.116E-05	6.347E-05	2.116E-04	6.345E-04	2.114E-03	6.327E-03	1.907E-01	6.877E-01	
U-235+D	Ac-227+D	1.000E+00	0.000E+00	3.332E-07	2.937E-06	3.037E-05	2.258E-04	1.477E-03	5.667E-03	1.902E-01	6.874E-01	
U-238	U-238	5.400E-05	5.400E-05	5.400E-05	5.400E-05	5.400E-05	5.400E-05	5.400E-05	5.400E-05	5.400E-05	5.400E-05	
U-238+D	U-238+D	9.999E-01	9.999E-01	9.999E-01	9.999E-01	9.999E-01	9.999E-01	9.999E-01	9.999E-01	9.999E-01	9.999E-01	
U-238+D	U-234	9.999E-01	0.000E+00	2.835E-06	8.504E-06	2.835E-05	8.504E-05	2.834E-04	8.501E-04	2.795E-02	1.444E-01	
U-238+D	Th-230	9.999E-01	0.000E+00	1.276E-11	1.148E-10	1.276E-09	1.148E-08	1.275E-07	1.147E-06	1.227E-03	3.123E-02	
U-238+D	Ra-226+D	9.999E-01	0.000E+00	1.842E-15	4.973E-14	1.840E-12	4.958E-11	1.822E-09	4.813E-08	7.937E-04	2.895E-02	
U-238+D	Pb-210+D	9.999E-01	0.000E+00	1.423E-17	1.138E-15	1.346E-13	9.704E-12	8.486E-10	3.570E-08	7.877E-04	2.892E-02	

Source Factors for Ingrowth and Decay
 Combined Radioactivity and Leaching Factors
 Parent and Progeny Principal Radionuclide Contributions Indicated

Parent (i)	Product (j)	Thread Fraction	SF(j,t) = THF(j)*S1(j,t)/S1(i,0) At Time in Years								
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04
U-234	U-234	1.000E+00	1.000E+00	9.924E-01	9.774E-01	9.266E-01	7.955E-01	4.665E-01	1.015E-01	7.743E-34	0.000E+00
U-234	Th-230	1.000E+00	0.000E+00	8.968E-06	2.670E-05	8.667E-05	2.413E-04	6.293E-04	1.058E-03	1.014E-03	5.078E-04
U-234	Ra-226+D	1.000E+00	0.000E+00	1.941E-09	1.731E-08	1.864E-07	1.535E-06	1.259E-05	5.149E-05	7.490E-05	3.749E-05
U-234	Pb-210+D	1.000E+00	0.000E+00	1.996E-11	5.261E-10	1.793E-08	3.853E-07	7.032E-06	4.153E-05	6.673E-05	3.340E-05
U-235+D	U-235+D	1.000E+00	1.000E+00	9.924E-01	9.774E-01	9.266E-01	7.956E-01	4.667E-01	1.016E-01	7.966E-34	0.000E+00
U-235+D	Pa-231	1.000E+00	0.000E+00	2.100E-05	6.204E-05	1.960E-04	5.049E-04	9.864E-04	6.431E-04	1.519E-34	0.000E+00
U-235+D	Ac-227+D	1.000E+00	0.000E+00	3.295E-07	2.839E-06	2.716E-05	1.636E-04	5.614E-04	4.379E-04	1.118E-34	0.000E+00
U-238	U-238	5.400E-05	5.400E-05	5.359E-05	5.278E-05	5.004E-05	4.296E-05	2.520E-05	5.488E-06	4.301E-38	0.000E+00
U-238+D	U-238+D	9.999E-01	9.999E-01	9.924E-01	9.773E-01	9.266E-01	7.956E-01	4.666E-01	1.016E-01	7.965E-34	0.000E+00
U-238+D	U-234	9.999E-01	0.000E+00	2.813E-06	8.312E-06	2.627E-05	6.766E-05	1.323E-04	8.640E-05	2.226E-35	0.000E+00
U-238+D	Th-230	9.999E-01	0.000E+00	1.269E-11	1.131E-10	1.213E-09	9.873E-09	7.800E-08	2.919E-07	3.781E-07	1.893E-07
U-238+D	Ra-226+D	9.999E-01	0.000E+00	1.833E-15	4.897E-14	1.748E-12	4.249E-11	1.097E-09	1.122E-08	2.791E-08	1.397E-08
U-238+D	Pb-210+D	9.999E-01	0.000E+00	1.416E-17	1.122E-15	1.282E-13	8.374E-12	5.192E-10	8.470E-09	2.487E-08	1.245E-08

The effect of volatilization was also considered when computing the source factors for H-3 and C-14.

Cover Erosion rate (vcv):	0.001000 m/yr	
Contaminated Zone Erosion rate (vcz):	0.000000 m/yr	
Water Table Drop rate (vwt):	0.001000 m/yr	
Precipitation rate (Pr):	0.406000 m/yr	
Cover Removal Time (Tc):	0.000E+00 yr	
Overhead irrigation rate (Irr):	0.200 m/yr	Runoff coefficient (Cr): 0.200
Evapotranspiration coeff. (Ce):	0.500	Infiltration rate (In): 0.262 m/yr
Bulk soil density (rhob):	1.500 g/cm**3	Effective porosity (pe): 0.000

Radio-nuclide (i)	Distribution Coefficient Kd(i), cm ³ /g	Leaching Ratio q(i)
Ac-227	2.000000E+01	1.010E-02
Pa-231	5.000000E+01	4.064E-03
Pb-210	1.000000E+02	2.036E-03
Ra-226	7.000000E+01	2.906E-03
Th-230	6.000000E+04	3.401E-06
U-234	5.000000E+01	4.064E-03
U-235	5.000000E+01	4.064E-03
U-238	5.000000E+01	4.064E-03

Time Dependence of Source Geometry

Time Dependence of Cover Depth [Cd(i,t)]

[illegible]

[illegible]

[illegible]

Dose Conversion and Environmental Transport Factors for the Ground Pathway (p=1)

Nuclide (i)	DCF(i,1)*	ETFG(i,t) At Time in Years (dimensionless)									
		t=	0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04
Ac-227	4.951E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
At-218	5.847E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
Bi-210	3.606E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
Bi-211	2.559E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
Bi-214	9.808E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
Fr-223	1.980E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
Pa-231	1.906E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
Pa-234	1.155E+01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
Pa-234m	8.967E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
Pb-210	2.447E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
Pb-211	3.064E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
Pb-214	1.341E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
Po-210	5.231E-05	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
Po-211	4.764E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
Po-214	5.138E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
Po-215	1.016E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
Po-218	5.642E-05	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
Ra-223	6.034E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
Ra-226	3.176E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
Rn-219	3.083E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
Rn-222	2.354E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
Th-227	5.212E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
Th-230	1.209E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
Th-231	3.643E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
Th-234	2.410E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
Tl-207	1.980E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
Tl-210	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	4.017E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235	7.211E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238	1.031E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00

* - Units are (mrem/yr)/(pCi/g) at infinite depth and area. Multiplication by ETFG(i,t) converts to site conditions.

Dose/Source Ratios for External Radiation from the Ground (p=1)
 Parent and Progeny Principal Radionuclide Contributions Indicated

Parent (i)	Product (j)	Thread Fraction	DSR(j,1,t) At Time in Years (mrem/yr)/(pCi/g)									
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04	
U-234	U-234	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-234	Th-230	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-234	Ra-226+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-234	Pb-210+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-234	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-235+D	U-235+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-235+D	Pa-231	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-235+D	Ac-227+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-235+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-238	U-238	5.400E-05	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-238+D	U-238+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-238+D	U-234	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-238+D	Th-230	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-238+D	Ra-226+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-238+D	Pb-210+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-238+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	

The DSR includes contributions from associated (half-life ≤ 180 days) daughters.

Dose/Source Ratios for Inhalation Pathway, Excluding Radon (p=2)
Parent and Progeny Principal Radionuclide Contributions Indicated

[illegible]

The DSR includes contributions from associated (half-life ≤ 180 days) daughters.

Pathway Factors for the Inhalation Pathway (radon excluded)

Area (A):	1.0000E+04 m**2	Occupancy Factor (FO2):	4.5000E-01
Area Factor (FA2):	1.6930E-01	Annual Air Intake (FI2):	8.4000E+03 m**3/yr
Cover Depth [Cd(0)]:	0.0000E+00 m	Mass Loading (ASR2):	1.0000E-04 g/m**3
Contaminated Zone Thickness [T(0)]:	4.5720E-01 m	FA2 * FO2 * FI2 * ASR2:	6.3995E-02 g/yr

[illegible]

Dose Conversion and Environmental Transport Factors for the Inhalation Pathway, Excluding Radon (p=2)

Parent (i)	Product (j)	DCF(j,2)*	ETF(j,2,t) At Time in Years (g/yr)									
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04	
U-234	U-234	1.320E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-234	Th-230	3.260E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-234	Ra-226+D	8.594E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-234	Pb-210+D	2.320E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-235+D	U-235+D	1.230E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-235+D	Pa-231	1.280E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-235+D	Ac-227+D	6.724E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-238	U-238	1.180E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-238+D	U-238+D	1.180E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-238+D	U-234	1.320E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-238+D	Th-230	3.260E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-238+D	Ra-226+D	8.594E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-238+D	Pb-210+D	2.320E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	

* - The dose conversion factor units are mrem/pCi.

Parameters Used for Calculating Indoor and Outdoor Radon Flux

	*Floor	Cover	Contaminated
	Material	Material	Zone
Radon Diffusion Coefficient (m**2/s)	3.000E-07	2.000E-06	2.000E-06
Total Porosity	1.000E-01	4.000E-01	4.000E-01
Volumetric Water Content	3.000E-02	5.000E-02	3.061E-01
Bulk Density (g/cm**3)	2.400E+00	1.500E+00	1.500E+00
Rn-222 Emanation Coefficient	2.500E-01	2.500E-01	2.500E-01
Initial Thickness (m)	1.500E-01	0.000E+00	4.572E-01

Building Depth Below Ground Surface *(DMFL): -1.000E+00 (m)
Negative DMFL shows building depth adjusted (if necessary) for no penetration
of contaminated zone. Actual values used *(DMFLACT), m:

```
t=          0.0000E+00  1.0000E+00  3.0000E+00  1.0000E+01  3.0000E+01  1.0000E+02  3.0000E+02  1.0000E+04  5.5000E+04
DMFLACT= 0.0000E+00  0.0000E+00  0.0000E+00  0.0000E+00  0.0000E+00  0.0000E+00  0.0000E+00  0.0000E+00  0.0000E+00
Building indoor area factor *(FAI):          0.000E+00
FAI <= 0.0 shows calculated time-dependent value based on amount of wall area
extending into the contaminated zone. Actual values used *(FAIACT):
```

[illegible]

* - Parameters are used only for indoor radon flux

Time Dependence of Outdoor Radon Flux [FLUXO(i,t)]

[illegible]

Time Dependence of Indoor Radon Flux [FLUXI(i,t)]

[illegible]

Parameters Used for Calculating Indoor and Outdoor Radon Concentration

Radon Vertical Dimension of Mixing (HMIX):	2.000E+00	(m)
Average Annual Wind Speed (WIND):	2.000E+00	(m/sec)
Building Room Height (HRM):	2.500E+00	(m)
Building Air Exchange Rate (REXG):	5.000E-01	(1/hr)

Time Dependence of Outdoor Radon Concentration [CRNO(i,t)]

[illegible]

Time Dependence of Indoor Radon Concentration [HCONC(i,r)]

[illegible]

Outdoor Working Levels of Radon [WLOTD(i,t)]

Nuclide		WLOTD(i,t) (WL)								
(i)	t=	0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04
U-234		0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00
U-238		0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00

Indoor Working Levels of Radon [WLIND(i,t)]

Nuclide		WLIND(i,t) (WL)								
(i)	t=	0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04
U-234		0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00
U-238		0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00	0.0000E+00

Fraction of Time Spent Outdoors (FOTD): 2.500E-01
Fraction of Time Spent Indoors (FIND): 5.000E-01

Subpathway: Outdoor and Indoor Radon Flux

Progeny Principal Radionuclide Contribution

[illegible]

The DSR includes contributions from associated (half-life ≤ 180 days) daughters.

Subpathway: Indoor Radon from Water Usage

Progeny Principal Radionuclide Contribution

[illegible]

Dose/Source Ratios for Radon Pathway (p=9)

Subpathway: Indoor Radon from Water Usage

Parent and Progeny Principal Radionuclide Contributions Indicated

Parent (i)	Product (j)	Thread Fraction	DSRRNW(j,t) At Time in Years (mrem/yr)/(pCi/g)									
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04	
U-238+D	U-238+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-238+D	U-234	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-238+D	Th-230	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-238+D	Ra-226+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-238+D	Pb-210+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-238+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	

The DSR includes contributions from associated (half-life ≤ 180 days) daughters.

Transport Time Parameters for Unsaturated Zone Stratum No. 1

Stratum thickness [h(1)]: 305.000000 m

Bulk soil material density [rhob(1)]: 1.500000 g/cm**3

Effective porosity [peuz(1)]: 0.200000

Hydraulic conductivity [Khuz(1)]: 10.000000 m/yr

Total porosity [ptuz(1)]: 0.400000

Soil specific b parameter [buz(1)]: 5.300000

Saturation ratio [sruz(1)]: 0.765151

Radio-nuclide	Distribution Coefficient	Retardation Factor	Transport Time
(i)	Kduz(i,1), cm**3/g	Rduz(i,1)	Dtuz(i,1), yr
Ac-227	2.0000E+01	9.9020E+01	1.7613E+04
Pa-231	5.0000E+01	2.4605E+02	4.3766E+04
Pb-210	1.0000E+02	4.9110E+02	8.7354E+04
Ra-226	7.0000E+01	3.4407E+02	6.1201E+04
Th-230	6.0000E+04	2.9406E+05	5.2306E+07
U-234	5.0000E+01	2.4605E+02	4.3766E+04
U-235	5.0000E+01	2.4605E+02	4.3766E+04
U-238	5.0000E+01	2.4605E+02	4.3766E+04

Transport Time Parameters for Unsaturated Zone created by the Falling Water Table

Water table drop rate [vwt]: 0.001000 m/yr

Bulk soil material density [rhobaq]: 1.500000 g/cm**3

Effective porosity [peaq]: 0.200000

Hydraulic conductivity [Khaq]: 100.000000 m/yr

Total porosity [ptaq]: 0.400000

Soil specific b parameter [baq]: 5.300000

Saturation ratio [sruaq]: 0.645978

Radio-nuclide	Distribution Coefficient	Retardation Factor	Minimum Transport Time
(i)	Kdaq(i), cm**3/g	Rduaq(i)	Dtuag(i), yr
Ac-227	2.0000E+01	1.1710E+02	1.0777E+03
Pa-231	5.0000E+01	2.9126E+02	7.3269E+03
Pb-210	1.0000E+02	5.8151E+02	3.5045E+04
Ra-226	7.0000E+01	4.0736E+02	1.5355E+04
Th-230	6.0000E+04	3.4831E+05	Infinite
U-234	5.0000E+01	2.9126E+02	7.3269E+03
U-235	5.0000E+01	2.9126E+02	7.3269E+03
U-238	5.0000E+01	2.9126E+02	7.3269E+03

Dilution Factor Parameters for Mass-Balance (MB) Model

Contaminated zone area (A): 10000.000000 m**2
 Infiltration rate (In): 0.262400 m/yr
 Evapotranspiration coefficient (Ce): 0.500000
 Runoff coefficient (Cr): 0.200000
 Precipitation rate (Pr): 0.406000 m/yr
 Overhead irrigation rate (Irr): 0.200000 m/yr
 Well pumping rate (Uw): 250.000000 m**3/yr
 Dilution factor (fi, all i): 1.000000

Primary Parameters Used for Calculating Water/Soil
 Concentration Ratios for Groundwater Pathway Segment

Model used: Mass-Balance (MB)

Bulk soil density in contaminated zone (rhob): 1.500 g/cm**3

Radio- nuclide (i)	Dilution Factor	Retardation Factor	Breakthrough Time		Rise Time
	f(i)	Rdcz(i)	Chain year	Single Nuclide Dt(i), yr	dt(i), yr
Ac-227	1.000E+00	9.902E+01	1.869E+04	1.869E+04	0.000E+00
Pa-231	1.000E+00	2.460E+02	5.109E+04	5.109E+04	0.000E+00
Pb-210	1.000E+00	4.911E+02	5.109E+04	1.224E+05	0.000E+00
Ra-226	1.000E+00	3.441E+02	5.109E+04	7.656E+04	0.000E+00
Th-230	1.000E+00	2.941E+05	5.109E+04	Infinite	0.000E+00
U-234	1.000E+00	2.460E+02	5.109E+04	5.109E+04	0.000E+00
U-235	1.000E+00	2.460E+02	5.109E+04	5.109E+04	0.000E+00
U-238	1.000E+00	2.460E+02	5.109E+04	5.109E+04	0.000E+00

Water/Soil Concentration Ratios [WSR(j,1,t)] for Groundwater Pathway Segment

[illegible]

Water/Soil Concentration Ratios [WSR(j,2,t)] for Surface Water Pathway Segment

Watershed Area (Aw) = 1.0000E+06 m**2

Contaminated Zone Area (A) = 1.0000E+04 m**2

Dilution Factor (f') = 1.0000E-02

Soil Density (rhob) = 1.5000E+00 kg/m**3

[illegible]

Storage Times For Contaminated Foodstuffs

k	Food Item	STOR_T(k), days
1	non-leafy plants	14.
2	leafy plants	1.
3	milk	1.
4	meat	20.
5	fish	7.
6	crustacea	7.
7	well water	1.
8	surface water	1.
9	livestock fodder	45.

Storage Time Ingrowth and Decay Factors

Storage Time for k'th Foodstuff: $t = \text{STOR_T}(k)$, days

Parent (i)	Product (j)	Thread Fraction	STOR_ID(i,j,t) = CONCE(i,j,t)/CONCE(i,i,0)									
			t = 1.400E+01	1.000E+00	1.000E+00	2.000E+01	7.000E+00	7.000E+00	1.000E+00	1.000E+00	4.500E+01	
Ac-227	Ac-227	1.000E+00	9.988E-01	9.999E-01	9.999E-01	9.983E-01	9.994E-01	9.994E-01	9.999E-01	9.999E-01	9.961E-01	
Pa-231	Pa-231	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
Pa-231	Ac-227	1.000E+00	1.219E-03	8.716E-05	8.716E-05	1.742E-03	6.099E-04	6.099E-04	8.716E-05	8.716E-05	3.915E-03	
Pb-210	Pb-210	1.000E+00	9.988E-01	9.999E-01	9.999E-01	9.983E-01	9.994E-01	9.994E-01	9.999E-01	9.999E-01	9.962E-01	
Ra-226	Ra-226	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	9.999E-01	
Ra-226	Pb-210	1.000E+00	1.191E-03	8.510E-05	8.510E-05	1.701E-03	5.955E-04	5.955E-04	8.510E-05	8.510E-05	3.822E-03	
Th-230	Th-230	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
Th-230	Ra-226	1.000E+00	1.661E-05	1.186E-06	1.186E-06	2.372E-05	8.303E-06	8.303E-06	1.186E-06	1.186E-06	5.337E-05	
Th-230	Pb-210	1.000E+00	9.888E-09	5.047E-11	5.047E-11	2.018E-08	2.472E-09	2.472E-09	5.047E-11	5.047E-11	1.021E-07	
U-234	U-234	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-234	Th-230	1.000E+00	3.450E-07	2.465E-08	2.465E-08	4.929E-07	1.725E-07	1.725E-07	2.465E-08	2.465E-08	1.109E-06	
U-234	Ra-226	1.000E+00	2.865E-12	1.462E-14	1.462E-14	5.846E-12	7.162E-13	7.162E-13	1.462E-14	1.462E-14	2.960E-11	
U-234	Pb-210	1.000E+00	1.137E-15	4.146E-19	4.146E-19	3.315E-15	1.422E-16	1.422E-16	4.146E-19	4.146E-19	3.774E-14	
U-235	U-235	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-235	Pa-231	1.000E+00	8.110E-07	5.793E-08	5.793E-08	1.159E-06	4.055E-07	4.055E-07	5.793E-08	5.793E-08	2.607E-06	
U-235	Ac-227	1.000E+00	4.946E-10	2.524E-12	2.524E-12	1.009E-09	1.237E-10	1.237E-10	2.524E-12	2.524E-12	5.105E-09	
U-238	U-238	5.400E-05	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-238	U-238	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-238	U-234	1.000E+00	1.087E-07	7.762E-09	7.762E-09	1.552E-07	5.433E-08	5.433E-08	7.762E-09	7.762E-09	3.493E-07	
U-238	Th-230	1.000E+00	1.875E-14	9.565E-17	9.565E-17	3.826E-14	4.687E-15	4.687E-15	9.565E-17	9.565E-17	1.937E-13	
U-238	Ra-226	1.000E+00	1.038E-19	3.782E-23	3.782E-23	3.025E-19	1.297E-20	1.297E-20	3.782E-23	3.782E-23	3.446E-18	
U-238	Pb-210	1.000E+00	3.090E-23	8.045E-28	8.045E-28	1.287E-22	1.931E-24	1.931E-24	8.045E-28	8.045E-28	3.296E-21	

CONCE(i,j,t)/CONCE(i,i,0) is the concentration ratio of Product(j) at time t to Parent(i) at start of storage time.

Storage Time Correction Factors
Drinking Water from Well and/or Surface
Harvest Time = t - 2.74E-03 yr; Consumption Time = t yr

Parent (i)	Product (j)	Thread Fraction	CFWW(j,t,1) # At Time in Years								
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04
U-234	U-234	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-234	Th-230	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-234	Ra-226+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-234	Pb-210+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-235+D	U-235+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-235+D	Pa-231	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-235+D	Ac-227+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238	U-238	5.400E-05	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	U-238+D	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	U-234	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	Th-230	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	Ra-226+D	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	Pb-210+D	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00

#Correction factor = (concentration in media at consumption time)/(concentration at harvest time).

Storage Time Correction Factors
Irrigation Water for Nonleafy Plants from Well and/or Surface
Harvest Time = t - 4.11E-02 yr; Consumption Time = t - 3.83E-02 yr

Parent (i)	Product (j)	Thread Fraction	CFWW(j,t,2) # At Time in Years									
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04	
U-234	U-234	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-234	Th-230	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-234	Ra-226+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-234	Pb-210+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-235+D	U-235+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-235+D	Pa-231	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-235+D	Ac-227+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-238	U-238	5.400E-05	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-238+D	U-238+D	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-238+D	U-234	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-238+D	Th-230	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-238+D	Ra-226+D	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-238+D	Pb-210+D	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	

#Correction factor = (concentration in media at consumption time)/(concentration at harvest time).

Storage Time Correction Factors

Irrigation Water for Leafy Plants from Well and/or Surface

Harvest Time = t - 5.48E-03 yr; Consumption Time = t - 2.74E-03 yr

Parent (i)	Product (j)	Thread Fraction	CFWW(j,t,3)# At Time in Years								
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04
U-234	U-234	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-234	Th-230	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-234	Ra-226+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-234	Pb-210+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-235+D	U-235+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-235+D	Pa-231	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-235+D	Ac-227+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238	U-238	5.400E-05	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	U-238+D	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	U-234	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	Th-230	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	Ra-226+D	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	Pb-210+D	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00

#Correction factor = (concentration in media at consumption time)/(concentration at harvest time).

Storage Time Correction Factors

Irrigation Water for Livestock (Milk) Fodder from Well and/or Surface

Harvest Time = t - 1.29E-01 yr; Consumption Time = t - 1.26E-01 yr

Parent (i)	Product (j)	Thread Fraction	CFWW(j,t,5)# At Time in Years									
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04	
U-234	U-234	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-234	Th-230	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-234	Ra-226+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-234	Pb-210+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-235+D	U-235+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-235+D	Pa-231	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-235+D	Ac-227+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-238	U-238	5.400E-05	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-238+D	U-238+D	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-238+D	U-234	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-238+D	Th-230	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-238+D	Ra-226+D	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-238+D	Pb-210+D	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	

#Correction factor = (concentration in media at consumption time)/(concentration at harvest time).

Storage Time Correction Factors
 Irrigation Water for Livestock (Meat) Fodder from Well and/or Surface
 Harvest Time = t - 1.81E-01 yr; Consumption Time = t - 1.78E-01 yr

Parent (i)	Product (j)	Thread Fraction	CFWW(j,t,7)# At Time in Years								
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04
U-234	U-234	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-234	Th-230	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-234	Ra-226+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-234	Pb-210+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-235+D	U-235+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-235+D	Pa-231	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-235+D	Ac-227+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238	U-238	5.400E-05	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	U-238+D	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	U-234	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	Th-230	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	Ra-226+D	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	Pb-210+D	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00

#Correction factor = (concentration in media at consumption time)/(concentration at harvest time).

Storage Time Correction Factors
 Livestock (Milk) Water from Well and/or Surface
 Harvest Time = t - 5.48E-03 yr; Consumption Time = t - 2.74E-03 yr

Parent (i)	Product (j)	Thread Fraction	CFWW(j,t,4)# At Time in Years								
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04
U-234	U-234	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-234	Th-230	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-234	Ra-226+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-234	Pb-210+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-235+D	U-235+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-235+D	Pa-231	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-235+D	Ac-227+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238	U-238	5.400E-05	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	U-238+D	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	U-234	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	Th-230	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	Ra-226+D	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	Pb-210+D	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00

#Correction factor = (concentration in media at consumption time)/(concentration at harvest time).

Storage Time Correction Factors
 Livestock (Meat) Water from Well and/or Surface
 Harvest Time = t - 5.75E-02 yr; Consumption Time = t - 5.48E-02 yr

Parent (i)	Product (j)	Thread Fraction	CFWW(j,t,6) # At Time in Years								
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04
U-234	U-234	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-234	Th-230	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-234	Ra-226+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-234	Pb-210+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-235+D	U-235+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-235+D	Pa-231	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-235+D	Ac-227+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238	U-238	5.400E-05	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	U-238+D	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	U-234	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	Th-230	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	Ra-226+D	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	Pb-210+D	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00

#Correction factor = (concentration in media at consumption time)/(concentration at harvest time).

Storage Time Correction Factors for Nonleafy Plants
 Harvest Time = t - 3.83E-02 yr; Consumption Time = t yr

Parent (i)	Product (j)	Thread Fraction	CF3(j,1,t)# At Time in Years									
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04	
U-234	U-234	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-234	Th-230	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-234	Ra-226+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-234	Pb-210+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.001E+00	
U-235+D	U-235+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-235+D	Pa-231	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-235+D	Ac-227+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	9.996E-01	
U-238	U-238	5.400E-05	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-238+D	U-238+D	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-238+D	U-234	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-238+D	Th-230	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-238+D	Ra-226+D	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-238+D	Pb-210+D	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.001E+00	

#Correction factor = (concentration in media at consumption time)/(concentration at harvest time).

Storage Time Correction Factors for Leafy Plants
 Harvest Time = t - 2.74E-03 yr; Consumption Time = t yr

Parent (i)	Product (j)	Thread Fraction	CF3(j,2,t)# At Time in Years								
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04
U-234	U-234	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-234	Th-230	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-234	Ra-226+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-234	Pb-210+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-235+D	U-235+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-235+D	Pa-231	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-235+D	Ac-227+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238	U-238	5.400E-05	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	U-238+D	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	U-234	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	Th-230	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	Ra-226+D	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	Pb-210+D	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00

#Correction factor = (concentration in media at consumption time)/(concentration at harvest time).

Storage Time Correction Factors for Livestock (Meat) Fodder
 Harvest Time = t - 1.78E-01 yr; Consumption Time = t - 5.48E-02 yr

Parent (i)	Product (j)	Thread Fraction	CFLF(j,1,t)# At Time in Years								
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04
U-234	U-234	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-234	Th-230	1.000E+00	1.000E+00	1.371E+00	1.107E+00	1.030E+00	1.009E+00	1.002E+00	1.000E+00	1.000E+00	1.000E+00
U-234	Ra-226+D	1.000E+00	1.000E+00	1.009E+00	1.002E+00	1.001E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	9.999E-01
U-234	Pb-210+D	1.000E+00	1.000E+00	2.808E+00	1.531E+00	1.158E+00	1.057E+00	1.024E+00	1.015E+00	1.013E+00	1.002E+00
U-235+D	U-235+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-235+D	Pa-231	1.000E+00	1.000E+00	1.038E+00	1.011E+00	1.003E+00	1.001E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-235+D	Ac-227+D	1.000E+00	1.000E+00	2.226E+00	1.360E+00	1.111E+00	1.044E+00	1.024E+00	1.019E+00	1.017E+00	9.987E-01
U-238	U-238	5.400E-05	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	U-238+D	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	U-234	9.999E-01	1.000E+00	1.150E+00	1.044E+00	1.013E+00	1.004E+00	1.001E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	Th-230	9.999E-01	1.000E+00	1.799E+00	1.220E+00	1.061E+00	1.019E+00	1.005E+00	1.001E+00	1.000E+00	1.000E+00
U-238+D	Ra-226+D	9.999E-01	1.000E+00	1.016E+00	1.004E+00	1.001E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	9.999E-01
U-238+D	Pb-210+D	9.999E-01	1.000E+00	3.416E+00	1.706E+00	1.208E+00	1.074E+00	1.028E+00	1.016E+00	1.013E+00	1.002E+00

#Correction factor = (concentration in media at consumption time)/(concentration at harvest time).

Storage Time Correction Factors for Livestock (Milk) Fodder
 Harvest Time = t - 1.26E-01 yr; Consumption Time = t - 2.74E-03 yr

Parent (i)	Product (j)	Thread Fraction	CFLF(j,2,t)# At Time in Years								
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04
U-234	U-234	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-234	Th-230	1.000E+00	1.000E+00	1.349E+00	1.105E+00	1.030E+00	1.009E+00	1.002E+00	1.000E+00	1.000E+00	1.000E+00
U-234	Ra-226+D	1.000E+00	1.000E+00	1.008E+00	1.002E+00	1.001E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	9.999E-01
U-234	Pb-210+D	1.000E+00	1.000E+00	2.701E+00	1.521E+00	1.157E+00	1.057E+00	1.024E+00	1.015E+00	1.013E+00	1.002E+00
U-235+D	U-235+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-235+D	Pa-231	1.000E+00	1.000E+00	1.035E+00	1.011E+00	1.003E+00	1.001E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-235+D	Ac-227+D	1.000E+00	1.000E+00	2.153E+00	1.354E+00	1.110E+00	1.044E+00	1.024E+00	1.019E+00	1.017E+00	9.987E-01
U-238	U-238	5.400E-05	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	U-238+D	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	U-234	9.999E-01	1.000E+00	1.141E+00	1.043E+00	1.012E+00	1.004E+00	1.001E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	Th-230	9.999E-01	1.000E+00	1.749E+00	1.216E+00	1.061E+00	1.019E+00	1.005E+00	1.001E+00	1.000E+00	1.000E+00
U-238+D	Ra-226+D	9.999E-01	1.000E+00	1.015E+00	1.004E+00	1.001E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	9.999E-01
U-238+D	Pb-210+D	9.999E-01	1.000E+00	3.272E+00	1.693E+00	1.207E+00	1.074E+00	1.028E+00	1.016E+00	1.013E+00	1.002E+00

#Correction factor = (concentration in media at consumption time)/(concentration at harvest time).

Storage Time Correction Factors for Meat
 Harvest Time = t - 5.48E-02 yr; Consumption Time = t yr

Parent (i)	Product (j)	Thread Fraction	CF45(j,1,t)# At Time in Years								
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04
U-234	U-234	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-234	Th-230	1.000E+00	1.000E+00	1.213E+00	1.068E+00	1.020E+00	1.006E+00	1.001E+00	1.000E+00	1.000E+00	1.000E+00
U-234	Ra-226+D	1.000E+00	1.000E+00	1.004E+00	1.001E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-234	Pb-210+D	1.000E+00	1.000E+00	1.297E+00	1.129E+00	1.045E+00	1.017E+00	1.007E+00	1.004E+00	1.004E+00	1.001E+00
U-235+D	U-235+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-235+D	Pa-231	1.000E+00	1.000E+00	1.003E+00	1.001E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-235+D	Ac-227+D	1.000E+00	1.000E+00	3.791E+01	1.430E+01	5.478E+00	2.925E+00	2.098E+00	1.918E+00	1.849E+00	1.292E+00
U-238	U-238	5.400E-05	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	U-238+D	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	U-234	9.999E-01	1.000E+00	1.058E+00	1.019E+00	1.006E+00	1.002E+00	1.001E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	Th-230	9.999E-01	1.000E+00	1.433E+00	1.138E+00	1.040E+00	1.013E+00	1.003E+00	1.001E+00	1.000E+00	1.000E+00
U-238+D	Ra-226+D	9.999E-01	1.000E+00	1.008E+00	1.002E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	Pb-210+D	9.999E-01	1.000E+00	1.352E+00	1.161E+00	1.058E+00	1.022E+00	1.008E+00	1.005E+00	1.004E+00	1.001E+00

#Correction factor = (concentration in media at consumption time)/(concentration at harvest time).

Storage Time Correction Factors for Milk
 Harvest Time = t - 2.74E-03 yr; Consumption Time = t yr

Parent (i)	Product (j)	Thread Fraction	CF45(j,2,t)# At Time in Years								
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04
U-234	U-234	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-234	Th-230	1.000E+00	1.000E+00	1.351E+00	1.117E+00	1.034E+00	1.011E+00	1.002E+00	1.000E+00	1.000E+00	1.000E+00
U-234	Ra-226+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-234	Pb-210+D	1.000E+00	1.000E+00	1.037E+00	1.016E+00	1.006E+00	1.002E+00	1.001E+00	1.001E+00	1.001E+00	1.000E+00
U-235+D	U-235+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-235+D	Pa-231	1.000E+00	1.000E+00	1.249E+00	1.081E+00	1.024E+00	1.008E+00	1.002E+00	1.001E+00	1.000E+00	1.000E+00
U-235+D	Ac-227+D	1.000E+00	1.000E+00	1.002E+00	1.001E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	9.999E-01
U-238	U-238	5.400E-05	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	U-238+D	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	U-234	9.999E-01	1.000E+00	1.003E+00	1.001E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	Th-230	9.999E-01	1.000E+00	1.698E+00	1.233E+00	1.069E+00	1.022E+00	1.005E+00	1.001E+00	1.000E+00	1.000E+00
U-238+D	Ra-226+D	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00
U-238+D	Pb-210+D	9.999E-01	1.000E+00	1.045E+00	1.020E+00	1.007E+00	1.003E+00	1.001E+00	1.001E+00	1.001E+00	1.000E+00

#Correction factor = (concentration in media at consumption time)/(concentration at harvest time).

Storage Time Correction Factors for Fish & Crustacea
 Harvest Time = t - 1.92E-02 yr; Consumption Time = t yr

Parent (i)	Product (j)	Thread Fraction	CFF(j,1,t) # At Time in Years									
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04	
U-234	U-234	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-234	Th-230	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-234	Ra-226+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-234	Pb-210+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	9.995E-01	
U-235+D	U-235+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-235+D	Pa-231	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-235+D	Ac-227+D	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	9.997E-01	
U-238	U-238	5.400E-05	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-238+D	U-238+D	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-238+D	U-234	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-238+D	Th-230	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-238+D	Ra-226+D	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	
U-238+D	Pb-210+D	9.999E-01	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	1.000E+00	9.995E-01	

#Correction factor = (concentration in media at consumption time)/(concentration at harvest time).

[illegible]

Area and Depth Factors for Plant (p=3), Meat (p=4), and Milk (p=5) Pathways
 Ditch Irrigation (q=3)

Area Factor for Plant Foods [FA(3)] = 0.50

Nuclide (i)	t=	Depth Factor FD(i,3,t) (dimensionless)								
		0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04
Ac-227		1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00
Pa-231		1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00
Pb-210		1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00
Ra-226		1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00
Th-230		1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00
U-234		1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00
U-235		1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00
U-238		1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00	1.0000E+00

Area and Depth Factors for Plant (p=3), Meat (p=4), and Milk (p=5) Pathways
 Overhead Irrigation (q=4)

Area Factor for Plant Foods [FA(3)] = 0.50

The Depth Factor Value

FD(i,p,q,t) = 1.0000E+00

is applicable for all radionuclides(i) and times(t).

Area and Depth Factors for Meat (p=4) and Milk (p=5) Pathways
 Transfer from Livestock Water (q=5) and Soil (q=6) Intake

Area Factor for Meat and Milk [FA(p),p=4,5] = 0.50

The livestock water subpathway (q=5) and livestock soil intake subpathway (q=6)
 occur only for the meat (p=4) and milk (p=5) pathways.

Dose Conversion and Environmental Transport Factors for the Plant Food Pathway (p=3)

Subpathway: Root Uptake from Contaminated Soil (q=1)

Parent (i)	Product (j)	DCF(j,3)*	ETF(j,3,1,t) At Time in Years (g/yr)								
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04
U-234	U-234	2.830E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Th-230	5.480E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Ra-226+D	1.321E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Pb-210+D	7.276E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	U-235+D	2.673E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Pa-231	1.060E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Ac-227+D	1.480E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238	U-238	2.550E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-238+D	2.687E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-234	2.830E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Th-230	5.480E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Ra-226+D	1.321E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Pb-210+D	7.276E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00

* - The dose conversion factor units are mrem/pCi.

Dose Conversion and Environmental Transport Factors for the Plant Food Pathway (p=3)

Subpathway: Foliar Uptake from Contaminated Dust (q=2)

Parent (i)	Product (j)	DCF(j,3)*	ETF(j,3,2,t) At Time in Years (g/yr)								
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04
U-234	U-234	2.830E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Th-230	5.480E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Ra-226+D	1.321E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Pb-210+D	7.276E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	U-235+D	2.673E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Pa-231	1.060E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Ac-227+D	1.480E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238	U-238	2.550E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-238+D	2.687E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-234	2.830E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Th-230	5.480E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Ra-226+D	1.321E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Pb-210+D	7.276E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00

* - The dose conversion factor units are mrem/pCi.

Dose Conversion and Environmental Transport Factors for the Plant Food Pathway (p=3)
 Subpathway: Ditch Irrigation (q=3)

Parent (i)	Product (j)	DCF(j,3)*	ETF(j,3,3,t) * SF(j,t) At Time in Years (g/yr)									
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04	
U-234	U-234	2.830E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Th-230	5.480E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Ra-226+D	1.321E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Pb-210+D	7.276E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	U-235+D	2.673E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Pa-231	1.060E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Ac-227+D	1.480E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238	U-238	2.550E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-238+D	2.687E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-234	2.830E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Th-230	5.480E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Ra-226+D	1.321E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Pb-210+D	7.276E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00

* - The dose conversion factor units are mrem/pCi.

Dose Conversion and Environmental Transport Factors for the Plant Food Pathway (p=3)
 Subpathway: Overhead Irrigation (q=4)

Parent (i)	Product (j)	DCF(j,3)*	ETF(j,3,4,t) * SF(j,t) At Time in Years (g/yr)									
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04	
U-234	U-234	2.830E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Th-230	5.480E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Ra-226+D	1.321E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Pb-210+D	7.276E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	U-235+D	2.673E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Pa-231	1.060E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Ac-227+D	1.480E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238	U-238	2.550E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-238+D	2.687E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-234	2.830E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Th-230	5.480E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Ra-226+D	1.321E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Pb-210+D	7.276E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00

* - The dose conversion factor units are mrem/pCi.

Dose Conversion and Environmental Transport Factors for the Meat Pathway (p=4)
Subpathway: Fodder Root Uptake from Contaminated Soil (q=1)

Parent (i)	Product (j)	DCF(j,4)*	ETF(j,4,1,t) At Time in Years (g/yr)									
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04	
U-234	U-234	2.830E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Th-230	5.480E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Ra-226+D	1.321E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Pb-210+D	7.276E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	U-235+D	2.673E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Pa-231	1.060E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Ac-227+D	1.480E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238	U-238	2.550E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-238+D	2.687E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-234	2.830E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Th-230	5.480E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Ra-226+D	1.321E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Pb-210+D	7.276E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00

* - The dose conversion factor units are mrem/pCi.

Dose Conversion and Environmental Transport Factors for the Meat Pathway (p=4)
Subpathway: Fodder Foliar Uptake from Contaminated Dust (q=2)

Parent (i)	Product (j)	DCF(j,4)*	ETF(j,4,2,t) At Time in Years (g/yr)									
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04	
U-234	U-234	2.830E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Th-230	5.480E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Ra-226+D	1.321E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Pb-210+D	7.276E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	U-235+D	2.673E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Pa-231	1.060E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Ac-227+D	1.480E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238	U-238	2.550E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-238+D	2.687E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-234	2.830E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Th-230	5.480E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Ra-226+D	1.321E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Pb-210+D	7.276E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00

* - The dose conversion factor units are mrem/pCi.

Dose Conversion and Environmental Transport Factors for the Meat Pathway (p=4)
 Subpathway: Ditch Irrigation (q=3)

Parent (i)	Product (j)	DCF(j,4)*	ETF(j,4,3,t) * SF(j,t) At Time in Years (g/yr)								
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04
U-234	U-234	2.830E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Th-230	5.480E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Ra-226+D	1.321E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Pb-210+D	7.276E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	U-235+D	2.673E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Pa-231	1.060E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Ac-227+D	1.480E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238	U-238	2.550E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-238+D	2.687E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-234	2.830E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Th-230	5.480E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Ra-226+D	1.321E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Pb-210+D	7.276E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00

* - The dose conversion factor units are mrem/pCi.

Dose Conversion and Environmental Transport Factors for the Meat Pathway (p=4)
 Subpathway: Overhead Irrigation (q=4)

Parent (i)	Product (j)	DCF(j,4)*	ETF(j,4,4,t) * SF(j,t) At Time in Years (g/yr)								
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04
U-234	U-234	2.830E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Th-230	5.480E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Ra-226+D	1.321E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Pb-210+D	7.276E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	U-235+D	2.673E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Pa-231	1.060E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Ac-227+D	1.480E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238	U-238	2.550E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-238+D	2.687E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-234	2.830E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Th-230	5.480E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Ra-226+D	1.321E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Pb-210+D	7.276E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00

* - The dose conversion factor units are mrem/pCi.

Dose Conversion and Environmental Transport Factors for the Meat Pathway (p=4)
 Subpathway: Livestock Water (q=5)

Parent (i)	Product (j)	DCF(j,4)*	ETF(j,4,5,t) * SF(j,t) At Time in Years (g/yr)									
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04	
U-234	U-234	2.830E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Th-230	5.480E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Ra-226+D	1.321E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Pb-210+D	7.276E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	U-235+D	2.673E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Pa-231	1.060E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Ac-227+D	1.480E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238	U-238	2.550E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-238+D	2.687E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-234	2.830E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Th-230	5.480E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Ra-226+D	1.321E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Pb-210+D	7.276E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00

* - The dose conversion factor units are mrem/pCi.

Dose Conversion and Environmental Transport Factors for the Milk Pathway (p=5)
 Subpathway: Fodder Root Uptake from Contaminated Soil (q=1)

Parent (i)	Product (j)	DCF(j,5)*	ETF(j,5,1,t) At Time in Years (g/yr)									
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04	
U-234	U-234	2.830E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Th-230	5.480E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Ra-226+D	1.321E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Pb-210+D	7.276E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	U-235+D	2.673E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Pa-231	1.060E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Ac-227+D	1.480E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238	U-238	2.550E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-238+D	2.687E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-234	2.830E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Th-230	5.480E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Ra-226+D	1.321E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Pb-210+D	7.276E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00

* - The dose conversion factor units are mrem/pCi.

Dose Conversion and Environmental Transport Factors for the Milk Pathway (p=5)
 Subpathway: Fodder Foliar Uptake from Contaminated Dust (q=2)

Parent (i)	Product (j)	DCF(j,5)*	ETF(j,5,2,t) At Time in Years (g/yr)									
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04	
U-234	U-234	2.830E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Th-230	5.480E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Ra-226+D	1.321E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Pb-210+D	7.276E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	U-235+D	2.673E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Pa-231	1.060E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Ac-227+D	1.480E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238	U-238	2.550E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-238+D	2.687E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-234	2.830E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Th-230	5.480E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Ra-226+D	1.321E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Pb-210+D	7.276E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00

* - The dose conversion factor units are mrem/pCi.

Dose Conversion and Environmental Transport Factors for the Milk Pathway (p=5)

Subpathway: Ditch Irrigation (q=3)

Parent (i)	Product (j)	DCF(j,5)*	ETF(j,5,3,t) * SF(j,t) At Time in Years (g/yr)									
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04	
U-234	U-234	2.830E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Th-230	5.480E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Ra-226+D	1.321E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Pb-210+D	7.276E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	U-235+D	2.673E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Pa-231	1.060E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Ac-227+D	1.480E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238	U-238	2.550E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-238+D	2.687E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-234	2.830E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Th-230	5.480E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Ra-226+D	1.321E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Pb-210+D	7.276E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00

* - The dose conversion factor units are mrem/pCi.

Dose Conversion and Environmental Transport Factors for the Milk Pathway (p=5)

Subpathway: Overhead Irrigation (q=4)

Parent (i)	Product (j)	DCF(j,5)*	ETF(j,5,4,t) * SF(j,t) At Time in Years (g/yr)									
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04	
U-234	U-234	2.830E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Th-230	5.480E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Ra-226+D	1.321E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Pb-210+D	7.276E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	U-235+D	2.673E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Pa-231	1.060E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Ac-227+D	1.480E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238	U-238	2.550E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-238+D	2.687E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-234	2.830E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Th-230	5.480E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Ra-226+D	1.321E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Pb-210+D	7.276E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00

* - The dose conversion factor units are mrem/pCi.

Dose Conversion and Environmental Transport Factors for the Milk Pathway (p=5)
 Subpathway: Livestock Water (q=5)

Parent (i)	Product (j)	DCF(j,5)*	ETF(j,5,5,t) * SF(j,t) At Time in Years (g/yr)									
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04	
U-234	U-234	2.830E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Th-230	5.480E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Ra-226+D	1.321E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Pb-210+D	7.276E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	U-235+D	2.673E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Pa-231	1.060E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Ac-227+D	1.480E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238	U-238	2.550E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-238+D	2.687E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-234	2.830E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Th-230	5.480E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Ra-226+D	1.321E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Pb-210+D	7.276E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00

* - The dose conversion factor units are mrem/pCi.

Dose Conversion and Environmental Transport Factors for the Fish Pathway (p=6)

Parent (i)	Product (j)	DCF(j,6)*	ETF(j,6,t) * SF(j,t) At Time in Years (g/yr)								
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04
U-234	U-234	2.830E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Th-230	5.480E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Ra-226+D	1.321E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Pb-210+D	7.276E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	U-235+D	2.673E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Pa-231	1.060E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Ac-227+D	1.480E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238	U-238	2.550E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-238+D	2.687E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-234	2.830E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Th-230	5.480E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Ra-226+D	1.321E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Pb-210+D	7.276E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00

* - The dose conversion factor units are mrem/pCi.

Dose Conversion and Environmental Transport Factors for the Drinking Water Pathway (p=7)

Parent (i)	Product (j)	DCF(j,7)*	ETF(j,7,t) * SF(j,t) At Time in Years (g/yr)								
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04
U-234	U-234	2.830E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	7.384E-08
U-234	Th-230	5.480E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	8.217E-03
U-234	Ra-226+D	1.321E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	7.040E+00
U-234	Pb-210+D	7.276E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	4.932E+00
U-235+D	U-235+D	2.673E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	8.629E-08
U-235+D	Pa-231	1.060E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	5.934E-08
U-235+D	Ac-227+D	1.480E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	8.818E-08
U-238	U-238	2.550E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	4.660E-12
U-238+D	U-238+D	2.687E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	8.629E-08
U-238+D	U-234	2.830E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	1.246E-08
U-238+D	Th-230	5.480E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	1.562E-03
U-238+D	Ra-226+D	1.321E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	1.302E+00
U-238+D	Pb-210+D	7.276E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	9.121E-01

* - The dose conversion factor units are mrem/pCi.

Dose/Source Ratios for Internal Radiation from Ingestion of Plant Foods (p=3)

Subpathway: Root Uptake from Contaminated Soil (q=1)

Parent and Progeny Principal Radionuclide Contributions Indicated

Parent (i)	Product (j)	Thread Fraction	DSR(j,3,1t) At Time in Years (mrem/yr)/(pCi/g)									
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04	
U-234	U-234	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Th-230	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Ra-226+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Pb-210+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	U-235+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Pa-231	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Ac-227+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238	U-238	5.400E-05	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-238+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-234	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Th-230	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Ra-226+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Pb-210+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00

The DSR includes contributions from associated (half-life ≤ 180 days) daughters.

Dose/Source Ratios for Internal Radiation from Ingestion of Plant Foods (p=3)

Subpathway: Foliar Uptake from Contaminated Dust (q=2)

Parent and Progeny Principal Radionuclide Contributions Indicated

Parent (i)	Product (j)	Thread Fraction	DSR(j,3,2t) At Time in Years (mrem/yr)/(pCi/g)									
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04	
U-234	U-234	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Th-230	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Ra-226+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Pb-210+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	U-235+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Pa-231	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Ac-227+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238	U-238	5.400E-05	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-238+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-234	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Th-230	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Ra-226+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Pb-210+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00

The DSR includes contributions from associated (half-life ≤ 180 days) daughters.

Dose/Source Ratios for Internal Radiation from Ingestion of Plant Foods (p=3)

Subpathway: Ditch Irrigation (q=3)

Parent and Progeny Principal Radionuclide Contributions Indicated

Parent (i)	Product (j)	Thread Fraction	DSR(j,3,3t) At Time in Years (mrem/yr)/(pCi/g)									
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04	
U-234	U-234	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Th-230	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Ra-226+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Pb-210+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	U-235+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Pa-231	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Ac-227+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238	U-238	5.400E-05	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-238+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-234	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Th-230	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Ra-226+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Pb-210+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00

The DSR includes contributions from associated (half-life ≤ 180 days) daughters.

Dose/Source Ratios for Internal Radiation from Ingestion of Plant Foods (p=3)

Subpathway: Overhead Irrigation (q=4)

Parent and Progeny Principal Radionuclide Contributions Indicated

Parent (i)	Product (j)	Thread Fraction	DSR(j,3,4t) At Time in Years (mrem/yr)/(pCi/g)									
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04	
U-234	U-234	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Th-230	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Ra-226+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Pb-210+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	U-235+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Pa-231	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Ac-227+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238	U-238	5.400E-05	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-238+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-234	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Th-230	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Ra-226+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Pb-210+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00

The DSR includes contributions from associated (half-life ≤ 180 days) daughters.

Dose/Source Ratios for Internal Radiation from Ingestion of Plant Foods (p=3)

Total for All Subpathways

Parent and Progeny Principal Radionuclide Contributions Indicated

Parent (i)	Product (j)	Thread Fraction	DSR(j,3,t) At Time in Years (mrem/yr)/(pCi/g)									
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04	
U-234	U-234	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Th-230	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Ra-226+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Pb-210+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	U-235+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Pa-231	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Ac-227+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238	U-238	5.400E-05	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-238+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-234	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Th-230	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Ra-226+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Pb-210+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00

The DSR includes contributions from associated (half-life ≤ 180 days) daughters.

Dose/Source Ratios for Internal Radiation from Ingestion of Meat (p=4)
Subpathway: Fodder Root Uptake from Contaminated Soil (q=1)
Parent and Progeny Principal Radionuclide Contributions Indicated

Parent (i)	Product (j)	Thread Fraction	DSR(j,4,1t) At Time in Years (mrem/yr)/(pCi/g)									
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04	
U-234	U-234	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Th-230	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Ra-226+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Pb-210+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	U-235+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Pa-231	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Ac-227+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238	U-238	5.400E-05	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-238+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-234	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Th-230	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Ra-226+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Pb-210+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00

The DSR includes contributions from associated (half-life ≤ 180 days) daughters.

Dose/Source Ratios for Internal Radiation from Ingestion of Meat (p=4)

Subpathway: Fodder Foliar Uptake from Contaminated Dust (q=2)

Parent and Progeny Principal Radionuclide Contributions Indicated

Parent (i)	Product (j)	Thread Fraction	DSR(j,4,2t) At Time in Years (mrem/yr)/(pCi/g)								
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04
U-234	U-234	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Th-230	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Ra-226+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Pb-210+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	U-235+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Pa-231	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Ac-227+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238	U-238	5.400E-05	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-238+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-234	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Th-230	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Ra-226+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Pb-210+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00

The DSR includes contributions from associated (half-life ≤ 180 days) daughters.

Dose/Source Ratios for Internal Radiation from Ingestion of Meat (p=4)

Subpathway: Ditch Irrigation (q=3)

Parent and Progeny Principal Radionuclide Contributions Indicated

Parent (i)	Product (j)	Thread Fraction	DSR(j,4,3t) At Time in Years (mrem/yr)/(pCi/g)									
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04	
U-234	U-234	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Th-230	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Ra-226+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Pb-210+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	U-235+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Pa-231	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Ac-227+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238	U-238	5.400E-05	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-238+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-234	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Th-230	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Ra-226+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Pb-210+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00

The DSR includes contributions from associated (half-life ≤ 180 days) daughters.

Dose/Source Ratios for Internal Radiation from Ingestion of Meat (p=4)

Subpathway: Overhead Irrigation (q=4)

Parent and Progeny Principal Radionuclide Contributions Indicated

Parent (i)	Product (j)	Thread Fraction	DSR(j,4,4t) At Time in Years (mrem/yr)/(pCi/g)									
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04	
U-234	U-234	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Th-230	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Ra-226+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Pb-210+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	U-235+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Pa-231	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Ac-227+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238	U-238	5.400E-05	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-238+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-234	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Th-230	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Ra-226+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Pb-210+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00

The DSR includes contributions from associated (half-life ≤ 180 days) daughters.

Dose/Source Ratios for Internal Radiation from Ingestion of Meat (p=4)

Subpathway: Livestock Water (q=5)

Parent and Progeny Principal Radionuclide Contributions Indicated

Parent (i)	Product (j)	Thread Fraction	DSR(j,4,5t) At Time in Years (mrem/yr)/(pCi/g)									
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04	
U-234	U-234	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Th-230	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Ra-226+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Pb-210+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	U-235+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Pa-231	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Ac-227+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238	U-238	5.400E-05	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-238+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-234	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Th-230	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Ra-226+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Pb-210+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00

The DSR includes contributions from associated (half-life ≤ 180 days) daughters.

Dose/Source Ratios for Internal Radiation from Ingestion of Meat (p=4)

Total for All Subpathways

Parent and Progeny Principal Radionuclide Contributions Indicated

Parent (i)	Product (j)	Thread Fraction	DSR(j,4,t) At Time in Years (mrem/yr)/(pCi/g)									
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04	
U-234	U-234	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Th-230	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Ra-226+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Pb-210+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	U-235+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Pa-231	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Ac-227+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238	U-238	5.400E-05	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-238+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-234	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Th-230	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Ra-226+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Pb-210+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00

The DSR includes contributions from associated (half-life ≤ 180 days) daughters.

Dose/Source Ratios for Internal Radiation from Ingestion of Milk (p=5)

Subpathway: Fodder Root Uptake from Contaminated Soil (q=1)

Parent and Progeny Principal Radionuclide Contributions Indicated

Parent (i)	Product (j)	Thread Fraction	DSR(j,5,1t) At Time in Years (mrem/yr)/(pCi/g)									
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04	
U-234	U-234	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Th-230	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Ra-226+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Pb-210+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	U-235+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Pa-231	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Ac-227+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238	U-238	5.400E-05	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-238+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-234	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Th-230	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Ra-226+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Pb-210+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00

The DSR includes contributions from associated (half-life ≤ 180 days) daughters.

Dose/Source Ratios for Internal Radiation from Ingestion of Milk (p=5)

Subpathway: Fodder Foliar Uptake from Contaminated Dust (q=2)

Parent and Progeny Principal Radionuclide Contributions Indicated

Parent (i)	Product (j)	Thread Fraction	DSR(j,5,2t) At Time in Years (mrem/yr)/(pCi/g)									
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04	
U-234	U-234	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-234	Th-230	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-234	Ra-226+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-234	Pb-210+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-234	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-235+D	U-235+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-235+D	Pa-231	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-235+D	Ac-227+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-235+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-238	U-238	5.400E-05	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-238+D	U-238+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-238+D	U-234	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-238+D	Th-230	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-238+D	Ra-226+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-238+D	Pb-210+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-238+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	

The DSR includes contributions from associated (half-life ≤ 180 days) daughters.

Dose/Source Ratios for Internal Radiation from Ingestion of Milk (p=5)

Subpathway: Ditch Irrigation (q=3)

Parent and Progeny Principal Radionuclide Contributions Indicated

Parent (i)	Product (j)	Thread Fraction	DSR(j,5,3t) At Time in Years (mrem/yr)/(pCi/g)									
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04	
U-234	U-234	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Th-230	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Ra-226+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Pb-210+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	U-235+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Pa-231	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Ac-227+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238	U-238	5.400E-05	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-238+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-234	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Th-230	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Ra-226+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Pb-210+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00

The DSR includes contributions from associated (half-life ≤ 180 days) daughters.

Dose/Source Ratios for Internal Radiation from Ingestion of Milk (p=5)

Subpathway: Overhead Irrigation (q=4)

Parent and Progeny Principal Radionuclide Contributions Indicated

Parent (i)	Product (j)	Thread Fraction	DSR(j,5,4t) At Time in Years (mrem/yr)/(pCi/g)								
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04
U-234	U-234	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Th-230	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Ra-226+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Pb-210+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	U-235+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Pa-231	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Ac-227+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238	U-238	5.400E-05	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-238+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-234	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Th-230	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Ra-226+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Pb-210+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00

The DSR includes contributions from associated (half-life ≤ 180 days) daughters.

Dose/Source Ratios for Internal Radiation from Ingestion of Milk (p=5)

Subpathway: Livestock Water (q=5).

Parent and Progeny Principal Radionuclide Contributions Indicated

Parent (i)	Product (j)	Thread Fraction	DSR(j,5,5t) At Time in Years (mrem/yr)/(pCi/g)									
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04	
U-234	U-234	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Th-230	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Ra-226+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Pb-210+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	U-235+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Pa-231	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Ac-227+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238	U-238	5.400E-05	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-238+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-234	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Th-230	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Ra-226+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Pb-210+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00

The DSR includes contributions from associated (half-life ≤ 180 days) daughters.

Dose/Source Ratios for Internal Radiation from Ingestion of Milk (p=5)

Total for All Subpathways

Parent and Progeny Principal Radionuclide Contributions Indicated

Parent (i)	Product (j)	Thread Fraction	DSR(j,5,t) At Time in Years (mrem/yr)/(pCi/g)								
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04
U-234	U-234	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Th-230	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Ra-226+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Pb-210+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	U-235+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Pa-231	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Ac-227+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238	U-238	5.400E-05	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-238+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-234	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Th-230	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Ra-226+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Pb-210+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00

The DSR includes contributions from associated (half-life ≤ 180 days) daughters.

Dose/Source Ratios for Internal Radiation from the Ingestion of Fish (p=6)
 Parent and Progeny Principal Radionuclide Contributions Indicated

Parent (i)	Product (j)	Thread Fraction	DSR(j,6,t) At Time in Years (mrem/yr)/(pCi/g)									
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04	
U-234	U-234	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Th-230	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Ra-226+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	Pb-210+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	U-235+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Pa-231	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	Ac-227+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-235+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238	U-238	5.400E-05	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-238+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	U-234	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Th-230	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Ra-226+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	Pb-210+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-238+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00

The DSR includes contributions from associated (half-life ≤ 180 days) daughters.

Dose/Source Ratios for Internal Radiation from the Ingestion of Drinking Water (p=7)

Parent and Progeny Principal Radionuclide Contributions Indicated

Parent (i)	Product (j)	Thread Fraction	DSR(j,7,t) At Time in Years (mrem/yr)/(pCi/g)								
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04
U-234	U-234	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	2.083E-11
U-234	Th-230	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	4.503E-06
U-234	Ra-226+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	9.299E-03
U-234	Pb-210+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	3.589E-02
U-234	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	4.519E-02
U-235+D	U-235+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	2.299E-11
U-235+D	Pa-231	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	6.270E-10
U-235+D	Ac-227+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	1.301E-09
U-235+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	1.951E-09
U-238	U-238	5.400E-05	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	1.184E-15
U-238+D	U-238+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	2.311E-11
U-238+D	U-234	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	3.514E-12
U-238+D	Th-230	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	8.560E-07
U-238+D	Ra-226+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	1.720E-03
U-238+D	Pb-210+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	6.637E-03
U-238+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	8.358E-03

The DSR includes contributions from associated (half-life ≤ 180 days) daughters.

Plant/Air and Plant/Water Concentration Ratios

Mass loading [ASR(3)]: 1.000E-04 g/m²*3

Area Factor for Mass Loading [FA(2)]: 1.693E-01

Nuclide (i)	FAR(i,3,2,1) m ² /3/g	FAR(i,3,2,2) m ² /3/g	FWR(i,3,3,1) L/g	FWR(i,3,3,2) L/g	FWR(i,3,4,1) L/g	FWR(i,3,4,2) L/g
Ac-227	5.4545E-02	2.6156E-01	2.8288E-07	4.1568E-07	3.4522E-04	1.6554E-03
Pa-231	5.4545E-02	2.6156E-01	1.1326E-06	1.6651E-06	3.4522E-04	1.6554E-03
Pb-210	5.4545E-02	2.6156E-01	1.1330E-06	1.6659E-06	3.4522E-04	1.6554E-03
Ra-226	5.4545E-02	2.6156E-01	4.5312E-06	6.6622E-06	3.4522E-04	1.6554E-03
Th-230	5.4545E-02	2.6156E-01	1.1216E-07	1.6824E-07	3.4522E-04	1.6554E-03
U-234	5.4545E-02	2.6156E-01	2.8315E-07	4.1628E-07	3.4522E-04	1.6554E-03
U-235	5.4545E-02	2.6156E-01	2.8315E-07	4.1628E-07	3.4522E-04	1.6554E-03
U-238	5.4545E-02	2.6156E-01	2.8315E-07	4.1628E-07	3.4522E-04	1.6554E-03

FAR(i,p,q,k) is the plant/air concentration ratio for airborne contaminated dust,
 and FWR(i,p,q,k) is the plant/water concentration ratio. See groundwater displays
 for water/soil concentration ratios.

Plant/Soil Concentration Ratios, FSR(i,3,q,k,t)

Root Uptake (q=1) and Foliar Dust Deposition (q=2)

Nonleafy (k=1) and/or Leafy (k=2) Vegetables

Nuclide(i)				
Parent	Product	FSR(i,3,1,k)	FSR(i,3,2,1)	FSR(i,3,2,2)
U-234	U-234	2.5000E-03	9.2345E-07	4.4282E-06
U-234	Th-230	1.0000E-03	9.2345E-07	4.4282E-06
U-234	Ra-226+D	4.0000E-02	9.2345E-07	4.4282E-06
U-234	Pb-210+D	1.0000E-02	9.2345E-07	4.4282E-06
U-235+D	U-235+D	2.5000E-03	9.2345E-07	4.4282E-06
U-235+D	Pa-231	1.0000E-02	9.2345E-07	4.4282E-06
U-235+D	Ac-227+D	2.5000E-03	9.2345E-07	4.4282E-06
U-238	U-238	2.5000E-03	9.2345E-07	4.4282E-06
U-238+D	U-238+D	2.5000E-03	9.2345E-07	4.4282E-06
U-238+D	U-234	2.5000E-03	9.2345E-07	4.4282E-06
U-238+D	Th-230	1.0000E-03	9.2345E-07	4.4282E-06
U-238+D	Ra-226+D	4.0000E-02	9.2345E-07	4.4282E-06
U-238+D	Pb-210+D	1.0000E-02	9.2345E-07	4.4282E-06

Plant/Soil Concentration Ratio, FSR(j,3,q,k,t)

Ditch Irrigation (q=3)

[illegible]

Plant/Soil Concentration Ratio, FSR(j,3,q,k,t)

Overhead Irrigation (g=4) and Nonleafy Vegetables (k=1)

[illegible]

Overhead Irrigation (q=4) and Leafy Vegetables (k=2)

[illegible]

Meat/Fodder, Milk/Fodder, Fodder/Air and Fodder/Water Concentration Ratios

FI(4,q): 68.0 kg/day FI(5,q): 55.0 kg/day q=1,2,3,4
FI(4,q): 50.0 L/day FI(5,q): 160.0 L/day q=5
FI(4,q): 0.5 kg/day FI(5,q):

Nuclide (i)	FQR(i,4) d/kg	FQR(i,5) d/kg	FAR(i,3,2,3) m**3/g	FWR(i,3,3,3) L/g	FWR(i,3,4,3) L/g
Ac-227	2.0000E-05	2.0000E-05	2.8659E-01	1.3323E-07	1.8139E-03
Pa-231	5.0000E-03	5.0000E-06	2.8659E-01	5.3317E-07	1.8139E-03
Pb-210	8.0000E-04	3.0000E-04	2.8659E-01	5.3323E-07	1.8139E-03
Ra-226	1.0000E-03	1.0000E-03	2.8659E-01	2.1328E-06	1.8139E-03
Th-230	1.0000E-04	5.0000E-06	2.8659E-01	5.6081E-08	1.8139E-03
U-234	3.4000E-04	6.0000E-04	2.8659E-01	1.3329E-07	1.8139E-03
U-235	3.4000E-04	6.0000E-04	2.8659E-01	1.3329E-07	1.8139E-03
U-238	3.4000E-04	6.0000E-04	2.8659E-01	1.3329E-07	1.8139E-03

FI(p,q) are the fodder (q=1,2,3,4), livestock water (q=5) and soil (q=6) intake rates;
FQR(i,p) are the transfer coefficients from contaminated fodder of livestock
water to meat (p=4) or milk (p=5). FAR(i,3,2,3) are the fodder/air
concentration ratios, and FWR(i,3,3,3) and FWR(i,3,4,3) are the fodder/
water concentration ratios for ditch and overhead irrigation, respectively.

Nuclide (i)		QSR(i,p,1)	QSR(i,p,2)
Parent	Product		
U-234	U-234	2.5000E-03	4.8520E-06
U-234	Th-230	1.0000E-03	4.8520E-06
U-234	Ra-226+D	4.0000E-02	4.8520E-06
U-234	Pb-210+D	1.0000E-02	4.8520E-06
U-235+D	U-235+D	2.5000E-03	4.8520E-06
U-235+D	Pa-231	1.0000E-02	4.8520E-06
U-235+D	Ac-227+D	2.5000E-03	4.8520E-06
U-238	U-238	2.5000E-03	4.8520E-06
U-238+D	U-238+D	2.5000E-03	4.8520E-06
U-238+D	U-234	2.5000E-03	4.8520E-06
U-238+D	Th-230	1.0000E-03	4.8520E-06
U-238+D	Ra-226+D	4.0000E-02	4.8520E-06
U-238+D	Pb-210+D	1.0000E-02	4.8520E-06

Fodder/Soil Concentration Ratio, QSR(j,p,q,t), for Meat and Milk Pathways
Ditch Irrigation (q=3)

[illegible]

[illegible][illegible]

Meat/Soil Concentration Ratio, FSR(j,4,q,t)
Overhead Irrigation (α=4)

[illegible]

Meat/Soil Concentration Ratio, FSR(j,4,q,t)
Livestock Water (q=5)

[illegible]

Milk/Soil Concentration Ratios, FSR(i,5,q,t)

Root Uptake ($q=1$) and Foliar Dust Deposition ($q=2$)

Nuclide (i)		Nuclide (j)	
Parent	Product	FSR(i,5,1)	FSR(i,5,2)
U-234	U-234	4.1910E-05	1.6012E-07
U-234	Th-230	0.0000E+00	0.0000E+00
U-234	Ra-226+D	0.0000E+00	0.0000E+00
U-234	Pb-210+D	0.0000E+00	0.0000E+00
U-235+D	U-235+D	4.1910E-05	1.6012E-07
U-235+D	Pa-231	0.0000E+00	0.0000E+00
U-235+D	Ac-227+D	0.0000E+00	0.0000E+00
U-238	U-238	2.2631E-09	8.6462E-12
U-238+D	U-238+D	4.1908E-05	1.6011E-07
U-238+D	U-234	0.0000E+00	0.0000E+00
U-238+D	Th-230	0.0000E+00	0.0000E+00
U-238+D	Ra-226+D	0.0000E+00	0.0000E+00
U-238+D	Pb-210+D	0.0000E+00	0.0000E+00

Milk/Soil Concentration Ratio, FSR(j,5,q,t)

Ditch Irrigation ($q=3$)[illegible]

Milk/Soil Concentration Ratio, FSR(j,5,q,t)
Overhead Irrigation (q=4)

[illegible]

Milk/Soil Concentration Ratio, FSR(j,5,q,t)
Livestock Water (q=5)

[illegible]

Dose/Source Ratios for Soil Ingestion Pathway (p=8)
 Parent and Progeny Principal Radionuclide Contributions Indicated

Parent (i)	Product (j)	Thread Fraction	DSR(j,8,t) At Time in Years (mrem/yr)/(pCi/g)									
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04	
U-234	U-234	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-234	Th-230	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-234	Ra-226+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-234	Pb-210+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-234	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-235+D	U-235+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-235+D	Pa-231	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-235+D	Ac-227+D	1.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-235+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-238	U-238	5.400E-05	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-238+D	U-238+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-238+D	U-234	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-238+D	Th-230	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-238+D	Ra-226+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-238+D	Pb-210+D	9.999E-01	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-238+D	ΣDSR(j)		0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	

The DSR includes contributions from associated (half-life ≤ 180 days) daughters.

Dose Conversion and Environmental Transport Factors for the Soil Ingestion Pathway (p=8)

Parent (i)	Product (j)	DCF(j,8)*	ETF(j,8,t) At Time in Years (g/yr)									
			0.000E+00	1.000E+00	3.000E+00	1.000E+01	3.000E+01	1.000E+02	3.000E+02	1.000E+04	5.500E+04	
U-234	U-234	2.830E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-234	Th-230	5.480E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-234	Ra-226+D	1.321E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-234	Pb-210+D	7.276E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-235+D	U-235+D	2.673E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-235+D	Pa-231	1.060E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-235+D	Ac-227+D	1.480E-02	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-238	U-238	2.550E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-238+D	U-238+D	2.687E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-238+D	U-234	2.830E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-238+D	Th-230	5.480E-04	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-238+D	Ra-226+D	1.321E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	
U-238+D	Pb-210+D	7.276E-03	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	

* - The dose conversion factor units are mrem/pCi.

RESRAD CONCENTRATION REPORT

Table of Contents

Part IV: Concentration of Radionuclides

Concentration of radionuclides in different media

Time= 0.000E+00	2
Time= 1.000E+00	3
Time= 3.000E+00	4
Time= 1.000E+01	5
Time= 3.000E+01	6
Time= 1.000E+02	7
Time= 3.000E+02	8
Time= 1.000E+04	9
Time= 5.500E+04	10

Concentration of radionuclides in environmental media
 at t = 0.000E+00 years

	Contaminat- ed Zone	Surface Soil*	Air Par- ticulate	Well Water	Surface Water
Radio- Nuclide	pCi/g	pCi/g	pCi/m**3	pCi/L	pCi/L
Ac-227	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
Pa-231	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
Pb-210	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
Ra-226	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
Th-230	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	1.230E+00	1.230E+00	2.082E-05	0.000E+00	0.000E+00
U-235	9.000E-02	9.000E-02	1.524E-06	0.000E+00	0.000E+00
U-238	6.600E+00	6.600E+00	1.117E-04	0.000E+00	0.000E+00

*The Surface Soil is the top layer of soil within the user specified mixing zone/depth.

Concentrations in the media occurring in pathways that are suppressed are calculated using the current input parameters, i.e. using parameters appearing in the input screen when the pathways are active.

Concentration of radionuclides in foodstuff media
 at t = 0.000E+00 years*

	Drinking Water	Nonleafy Vegetable	Leafy Vegetable	Fodder Meat	Fodder Milk	Meat	Milk	Fish	Crustacea
Radio- Nuclide	pCi/L	pCi/kg	pCi/kg	pCi/kg	pCi/kg	pCi/kg	pCi/L	pCi/kg	pCi/kg
Ac-227	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
Pa-231	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
Pb-210	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
Ra-226	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
Th-230	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00	0.000E+00
U-234	0.000E+00	1.563E+00	1.568E+00	1.568E+00	1.568E+00	2.454E-01	4.207E-01	0.000E+00	0.000E+00
U-235	0.000E+00	1.144E-01	1.147E-01	1.147E-01	1.147E-01	1.795E-02	3.079E-02	0.000E+00	0.000E+00
U-238	0.000E+00	8.388E+00	8.411E+00	8.414E+00	8.414E+00	1.317E+00	2.258E+00	0.000E+00	0.000E+00

*Concentrations are at consumption time and include radioactive decay and ingrowth during storage time.
 For livestock fodder, consumption time is t minus meat or milk storage time.

Concentrations in the media occurring in pathways that are suppressed are calculated using the current input parameters, i.e. using parameters appearing in the input screen when the pathways are active.

Concentration of radionuclides in environmental media
 at t = 1.000E+00 years

Radio- Nuclide	Contaminat- ed Zone	Surface Soil*	Air Par- ticulate	Well Water	Surface Water
	pCi/g	pCi/g	pCi/m**3	pCi/L	pCi/L
Ac-227	2.965E-08	2.965E-08	5.020E-13	0.000E+00	0.000E+00
Pa-231	1.890E-06	1.890E-06	3.199E-11	0.000E+00	0.000E+00
Pb-210	2.455E-11	2.455E-11	4.156E-16	0.000E+00	0.000E+00
Ra-226	2.388E-09	2.388E-09	4.042E-14	0.000E+00	0.000E+00
Th-230	1.103E-05	1.103E-05	1.867E-10	0.000E+00	0.000E+00
U-234	1.221E+00	1.221E+00	2.067E-05	0.000E+00	0.000E+00
U-235	8.932E-02	8.932E-02	1.512E-06	0.000E+00	0.000E+00
U-238	6.550E+00	6.550E+00	1.109E-04	0.000E+00	0.000E+00

*The Surface Soil is the top layer of soil within the user specified mixing zone/depth.

Concentrations in the media occurring in pathways that are suppressed are calculated using the current input parameters, i.e. using parameters appearing in the input screen when the pathways are active.

Concentration of radionuclides in foodstuff media
 at t = 1.000E+00 years*

Radio- Nuclide	Drinking Water	Nonleafy Vegetable	Leafy Vegetable	Fodder Meat	Fodder Milk	Meat	Milk	Fish	Crustacea
	pCi/L	pCi/kg	pCi/kg	pCi/kg	pCi/kg	pCi/kg	pCi/L	pCi/kg	pCi/kg
Ac-227	0.000E+00	4.617E-08	3.842E-08	5.709E-08	6.234E-08	1.300E-08	3.642E-10	0.000E+00	0.000E+00
Pa-231	0.000E+00	9.329E-06	9.589E-06	8.207E-06	8.704E-06	7.279E-06	8.875E-09	0.000E+00	0.000E+00
Pb-210	0.000E+00	1.645E-10	1.279E-10	1.952E-10	2.255E-10	2.455E-11	7.650E-12	0.000E+00	0.000E+00
Ra-226	0.000E+00	4.498E-08	4.827E-08	3.312E-08	3.740E-08	3.334E-09	3.245E-09	0.000E+00	0.000E+00
Th-230	0.000E+00	5.935E-06	5.675E-06	6.382E-06	6.674E-06	6.849E-07	3.963E-08	0.000E+00	0.000E+00
U-234	0.000E+00	1.552E+00	1.556E+00	1.558E+00	1.558E+00	2.436E-01	4.176E-01	0.000E+00	0.000E+00
U-235	0.000E+00	1.135E-01	1.138E-01	1.140E-01	1.140E-01	1.783E-02	3.056E-02	0.000E+00	0.000E+00
U-238	0.000E+00	8.327E+00	8.348E+00	8.361E+00	8.358E+00	1.307E+00	2.241E+00	0.000E+00	0.000E+00

*Concentrations are at consumption time and include radioactive decay and ingrowth during storage time.

For livestock fodder, consumption time is t minus meat or milk storage time.

Concentrations in the media occurring in pathways that are suppressed are calculated using the current input parameters, i.e. using parameters appearing in the input screen when the pathways are active.

Concentration of radionuclides in environmental media
 at t = 3.000E+00 years

	Contaminat- ed Zone	Surface Soil*	Air Par- ticulate	Well Water	Surface Water
Radio- Nuclide	pCi/g	pCi/g	pCi/m**3	pCi/L	pCi/L
Ac-227	2.555E-07	2.555E-07	4.325E-12	0.000E+00	0.000E+00
Pa-231	5.583E-06	5.583E-06	9.453E-11	0.000E+00	0.000E+00
Pb-210	6.471E-10	6.471E-10	1.096E-14	0.000E+00	0.000E+00
Ra-226	2.130E-08	2.130E-08	3.605E-13	0.000E+00	0.000E+00
Th-230	3.284E-05	3.284E-05	5.560E-10	0.000E+00	0.000E+00
U-234	1.202E+00	1.202E+00	2.035E-05	0.000E+00	0.000E+00
U-235	8.797E-02	8.797E-02	1.489E-06	0.000E+00	0.000E+00
U-238	6.451E+00	6.451E+00	1.092E-04	0.000E+00	0.000E+00

*The Surface Soil is the top layer of soil within the user specified mixing zone/depth.

Concentrations in the media occurring in pathways that are suppressed are calculated using the current input parameters, i.e. using parameters appearing in the input screen when the pathways are active.

Concentration of radionuclides in foodstuff media
 at t = 3.000E+00 years*

	Drinking Water	Nonleafy Vegetable	Leafy Vegetable	Fodder Meat	Fodder Milk	Meat	Milk	Fish	Crustacea
Radio- Nuclide	pCi/L	pCi/kg	pCi/kg	pCi/kg	pCi/kg	pCi/kg	pCi/L	pCi/kg	pCi/kg
Ac-227	0.000E+00	3.506E-07	3.275E-07	3.935E-07	4.057E-07	4.290E-08	2.998E-09	0.000E+00	0.000E+00
Pa-231	0.000E+00	2.811E-05	2.837E-05	2.704E-05	2.752E-05	2.292E-05	2.326E-08	0.000E+00	0.000E+00
Pb-210	0.000E+00	3.664E-09	3.318E-09	4.201E-09	4.407E-09	5.347E-10	1.723E-10	0.000E+00	0.000E+00
Ra-226	0.000E+00	4.221E-07	4.321E-07	3.842E-07	3.984E-07	3.643E-08	3.254E-08	0.000E+00	0.000E+00
Th-230	0.000E+00	1.703E-05	1.685E-05	1.756E-05	1.784E-05	1.850E-06	9.707E-08	0.000E+00	0.000E+00
U-234	0.000E+00	1.528E+00	1.532E+00	1.535E+00	1.534E+00	2.399E-01	4.113E-01	0.000E+00	0.000E+00
U-235	0.000E+00	1.118E-01	1.121E-01	1.123E-01	1.123E-01	1.756E-02	3.009E-02	0.000E+00	0.000E+00
U-238	0.000E+00	8.201E+00	8.221E+00	8.235E+00	8.232E+00	1.287E+00	2.207E+00	0.000E+00	0.000E+00

*Concentrations are at consumption time and include radioactive decay and ingrowth during storage time.
 For livestock fodder, consumption time is t minus meat or milk storage time.

Concentrations in the media occurring in pathways that are suppressed are calculated using the current input parameters, i.e. using parameters appearing in the input screen when the pathways are active.

Concentration of radionuclides in environmental media
 at t = 1.000E+01 years

	Contaminat- ed Zone	Surface Soil*	Air Par- ticulate	Well Water	Surface Water
Radio- Nuclide	pCi/g	pCi/g	pCi/m**3	pCi/L	pCi/L
Ac-227	2.445E-06	2.445E-06	4.139E-11	0.000E+00	0.000E+00
Pa-231	1.764E-05	1.764E-05	2.987E-10	0.000E+00	0.000E+00
Pb-210	2.206E-08	2.206E-08	3.734E-13	0.000E+00	0.000E+00
Ra-226	2.293E-07	2.293E-07	3.882E-12	0.000E+00	0.000E+00
Th-230	1.066E-04	1.066E-04	1.805E-09	0.000E+00	0.000E+00
U-234	1.140E+00	1.140E+00	1.930E-05	0.000E+00	0.000E+00
U-235	8.340E-02	8.340E-02	1.412E-06	0.000E+00	0.000E+00
U-238	6.116E+00	6.116E+00	1.035E-04	0.000E+00	0.000E+00

*The Surface Soil is the top layer of soil within the user specified mixing zone/depth.

Concentrations in the media occurring in pathways that are suppressed are calculated using the current input parameters, i.e. using parameters appearing in the input screen when the pathways are active.

Concentration of radionuclides in foodstuff media
 at t = 1.000E+01 years*

	Drinking Water	Nonleafy Vegetable	Leafy Vegetable	Fodder Meat	Fodder Milk	Meat	Milk	Fish	Crustacea
Radio- Nuclide	pCi/L	pCi/kg	pCi/kg	pCi/kg	pCi/kg	pCi/kg	pCi/L	pCi/kg	pCi/kg
Ac-227	0.000E+00	3.191E-06	3.121E-06	3.352E-06	3.382E-06	1.576E-07	2.816E-08	0.000E+00	0.000E+00
Pa-231	0.000E+00	8.941E-05	8.969E-05	8.851E-05	8.895E-05	7.400E-05	7.021E-08	0.000E+00	0.000E+00
Pb-210	0.000E+00	1.162E-07	1.124E-07	1.233E-07	1.251E-07	1.608E-08	5.401E-09	0.000E+00	0.000E+00
Ra-226	0.000E+00	4.626E-06	4.658E-06	4.502E-06	4.549E-06	4.197E-07	3.648E-07	0.000E+00	0.000E+00
Th-230	0.000E+00	5.455E-05	5.465E-05	5.535E-05	5.562E-05	5.791E-06	2.914E-07	0.000E+00	0.000E+00
U-234	0.000E+00	1.449E+00	1.453E+00	1.455E+00	1.455E+00	2.275E-01	3.900E-01	0.000E+00	0.000E+00
U-235	0.000E+00	1.060E-01	1.063E-01	1.065E-01	1.064E-01	1.664E-02	2.853E-02	0.000E+00	0.000E+00
U-238	0.000E+00	7.775E+00	7.794E+00	7.807E+00	7.804E+00	1.221E+00	2.092E+00	0.000E+00	0.000E+00

*Concentrations are at consumption time and include radioactive decay and ingrowth during storage time.
 For livestock fodder, consumption time is t minus meat or milk storage time.

Concentrations in the media occurring in pathways that are suppressed are calculated using the current input parameters, i.e. using parameters appearing in the input screen when the pathways are active.

Concentration of radionuclides in environmental media
 at t = 3.000E+01 years

	Contaminat- ed Zone	Surface Soil*	Air Par- ticulate	Well Water	Surface Water
Radio- Nuclide	pCi/g	pCi/g	pCi/m**3	pCi/L	pCi/L
Ac-227	1.472E-05	1.472E-05	2.493E-10	0.000E+00	0.000E+00
Pa-231	4.544E-05	4.544E-05	7.692E-10	0.000E+00	0.000E+00
Pb-210	4.740E-07	4.740E-07	8.025E-12	0.000E+00	0.000E+00
Ra-226	1.888E-06	1.888E-06	3.196E-11	0.000E+00	0.000E+00
Th-230	2.969E-04	2.969E-04	5.027E-09	0.000E+00	0.000E+00
U-234	9.790E-01	9.790E-01	1.657E-05	0.000E+00	0.000E+00
U-235	7.161E-02	7.161E-02	1.212E-06	0.000E+00	0.000E+00
U-238	5.251E+00	5.251E+00	8.890E-05	0.000E+00	0.000E+00

*The Surface Soil is the top layer of soil within the user specified mixing zone/depth.

Concentrations in the media occurring in pathways that are suppressed are calculated using the current input parameters, i.e. using parameters appearing in the input screen when the pathways are active.

Concentration of radionuclides in foodstuff media
 at t = 3.000E+01 years*

	Drinking Water	Nonleafy Vegetable	Leafy Vegetable	Fodder Meat	Fodder Milk	Meat	Milk	Fish	Crustacea
Radio- Nuclide	pCi/L	pCi/kg	pCi/kg	pCi/kg	pCi/kg	pCi/kg	pCi/L	pCi/kg	pCi/kg
Ac-227	0.000E+00	1.894E-05	1.878E-05	1.944E-05	1.949E-05	5.069E-07	1.687E-07	0.000E+00	0.000E+00
Pa-231	0.000E+00	2.307E-04	2.310E-04	2.302E-04	2.305E-04	1.917E-04	1.784E-07	0.000E+00	0.000E+00
Pb-210	0.000E+00	2.443E-06	2.412E-06	2.508E-06	2.520E-06	3.306E-07	1.129E-07	0.000E+00	0.000E+00
Ra-226	0.000E+00	3.828E-05	3.837E-05	3.796E-05	3.808E-05	3.522E-06	3.038E-06	0.000E+00	0.000E+00
Th-230	0.000E+00	1.514E-04	1.522E-04	1.529E-04	1.531E-04	1.596E-05	7.926E-07	0.000E+00	0.000E+00
U-234	0.000E+00	1.245E+00	1.248E+00	1.250E+00	1.249E+00	1.954E-01	3.349E-01	0.000E+00	0.000E+00
U-235	0.000E+00	9.103E-02	9.126E-02	9.141E-02	9.137E-02	1.429E-02	2.450E-02	0.000E+00	0.000E+00
U-238	0.000E+00	6.676E+00	6.692E+00	6.703E+00	6.701E+00	1.048E+00	1.796E+00	0.000E+00	0.000E+00

*Concentrations are at consumption time and include radioactive decay and ingrowth during storage time.
 For livestock fodder, consumption time is t minus meat or milk storage time.

Concentrations in the media occurring in pathways that are suppressed are calculated using the current input parameters, i.e. using parameters appearing in the input screen when the pathways are active.

Concentration of radionuclides in environmental media

at t = 1.000E+02 years

Radio-	Contaminat-	Surface	Air Par-	Well	Surface
Nuclide	ted Zone	Soil*	ticulate	Water	Water
	pCi/g	pCi/g	pCi/m**3	pCi/L	pCi/L
Ac-227	5.053E-05	5.053E-05	8.554E-10	0.000E+00	0.000E+00
Pa-231	8.877E-05	8.877E-05	1.503E-09	0.000E+00	0.000E+00
Pb-210	8.653E-06	8.653E-06	1.465E-10	0.000E+00	0.000E+00
Ra-226	1.550E-05	1.550E-05	2.624E-10	0.000E+00	0.000E+00
Th-230	7.746E-04	7.746E-04	1.311E-08	0.000E+00	0.000E+00
U-234	5.747E-01	5.747E-01	9.730E-06	0.000E+00	0.000E+00
U-235	4.200E-02	4.200E-02	7.111E-07	0.000E+00	0.000E+00
U-238	3.080E+00	3.080E+00	5.214E-05	0.000E+00	0.000E+00

*The Surface Soil is the top layer of soil within the user specified mixing zone/depth.

Concentrations in the media occurring in pathways that are suppressed are calculated using the current input parameters, i.e. using parameters appearing in the input screen when the pathways are active.

Concentration of radionuclides in foodstuff media

at t = 1.000E+02 years*

Radio-	Drinking	Nonleafy	Leafy	Fodder	Fodder	Meat	Milk	Fish	Crustacea
Nuclide	Water	Vegetable	Vegetable	Meat	Milk				
	pCi/L	pCi/kg	pCi/kg	pCi/kg	pCi/kg	pCi/kg	pCi/L	pCi/kg	pCi/kg
Ac-227	0.000E+00	6.467E-05	6.443E-05	6.587E-05	6.589E-05	1.247E-06	5.777E-07	0.000E+00	0.000E+00
Pa-231	0.000E+00	4.510E-04	4.514E-04	4.513E-04	4.514E-04	3.754E-04	3.469E-07	0.000E+00	0.000E+00
Pb-210	0.000E+00	4.425E-05	4.402E-05	4.487E-05	4.492E-05	5.938E-06	2.041E-06	0.000E+00	0.000E+00
Ra-226	0.000E+00	3.147E-04	3.149E-04	3.141E-04	3.143E-04	2.910E-05	2.504E-05	0.000E+00	0.000E+00
Th-230	0.000E+00	3.944E-04	3.969E-04	3.976E-04	3.977E-04	4.147E-05	2.051E-06	0.000E+00	0.000E+00
U-234	0.000E+00	7.306E-01	7.325E-01	7.337E-01	7.334E-01	1.147E-01	1.966E-01	0.000E+00	0.000E+00
U-235	0.000E+00	5.340E-02	5.353E-02	5.362E-02	5.360E-02	8.383E-03	1.437E-02	0.000E+00	0.000E+00
U-238	0.000E+00	3.916E+00	3.925E+00	3.932E+00	3.930E+00	6.147E-01	1.054E+00	0.000E+00	0.000E+00

*Concentrations are at consumption time and include radioactive decay and ingrowth during storage time.

For livestock fodder, consumption time is t minus meat or milk storage time.

Concentrations in the media occurring in pathways that are suppressed are calculated using the current input parameters, i.e. using parameters appearing in the input screen when the pathways are active.

Concentration of radionuclides in environmental media
 at t = 3.000E+02 years

	Contaminat- ed Zone	Surface Soil*	Air Par- ticulate	Well Water	Surface Water
Radio- Nuclide	pCi/g	pCi/g	pCi/m**3	pCi/L	pCi/L
Ac-227	3.941E-05	3.941E-05	6.672E-10	0.000E+00	0.000E+00
Pa-231	5.788E-05	5.788E-05	9.798E-10	0.000E+00	0.000E+00
Pb-210	5.114E-05	5.114E-05	8.658E-10	0.000E+00	0.000E+00
Ra-226	6.340E-05	6.340E-05	1.073E-09	0.000E+00	0.000E+00
Th-230	1.303E-03	1.303E-03	2.205E-08	0.000E+00	0.000E+00
U-234	1.255E-01	1.255E-01	2.124E-06	0.000E+00	0.000E+00
U-235	9.147E-03	9.147E-03	1.549E-07	0.000E+00	0.000E+00
U-238	6.708E-01	6.708E-01	1.136E-05	0.000E+00	0.000E+00

*The Surface Soil is the top layer of soil within the user specified mixing zone/depth.

Concentrations in the media occurring in pathways that are suppressed are calculated using the current input parameters, i.e. using parameters appearing in the input screen when the pathways are active.

Concentration of radionuclides in foodstuff media
 at t = 3.000E+02 years*

	Drinking Water	Nonleafy Vegetable	Leafy Vegetable	Fodder Meat	Fodder Milk	Meat	Milk	Fish	Crustacea
Radio- Nuclide	pCi/L	pCi/kg	pCi/kg	pCi/kg	pCi/kg	pCi/kg	pCi/L	pCi/kg	pCi/kg
Ac-227	0.000E+00	5.039E-05	5.025E-05	5.123E-05	5.122E-05	8.895E-07	4.504E-07	0.000E+00	0.000E+00
Pa-231	0.000E+00	2.941E-04	2.943E-04	2.945E-04	2.945E-04	2.449E-04	2.259E-07	0.000E+00	0.000E+00
Pb-210	0.000E+00	2.610E-04	2.601E-04	2.638E-04	2.638E-04	3.494E-05	1.203E-05	0.000E+00	0.000E+00
Ra-226	0.000E+00	1.288E-03	1.289E-03	1.288E-03	1.288E-03	1.193E-04	1.025E-04	0.000E+00	0.000E+00
Th-230	0.000E+00	6.630E-04	6.675E-04	6.682E-04	6.682E-04	6.969E-05	3.441E-06	0.000E+00	0.000E+00
U-234	0.000E+00	1.595E-01	1.599E-01	1.602E-01	1.601E-01	2.504E-02	4.293E-02	0.000E+00	0.000E+00
U-235	0.000E+00	1.163E-02	1.166E-02	1.168E-02	1.167E-02	1.826E-03	3.129E-03	0.000E+00	0.000E+00
U-238	0.000E+00	8.528E-01	8.549E-01	8.563E-01	8.560E-01	1.339E-01	2.295E-01	0.000E+00	0.000E+00

*Concentrations are at consumption time and include radioactive decay and ingrowth during storage time.
 For livestock fodder, consumption time is t minus meat or milk storage time.

Concentrations in the media occurring in pathways that are suppressed are calculated using the current input parameters, i.e. using parameters appearing in the input screen when the pathways are active.

Concentration of radionuclides in environmental media
 at t = 1.000E+04 years

	Contaminat- ted Zone	Surface Soil*	Air Par- ticulate	Well Water	Surface Water
Radio- Nuclide	pCi/g	pCi/g	pCi/m**3	pCi/L	pCi/L
Ac-227	1.006E-35	1.006E-35	1.704E-40	0.000E+00	0.000E+00
Pa-231	1.367E-35	1.367E-35	2.315E-40	0.000E+00	0.000E+00
Pb-210	8.224E-05	8.224E-05	1.392E-09	0.000E+00	0.000E+00
Ra-226	9.231E-05	9.231E-05	1.563E-09	0.000E+00	0.000E+00
Th-230	1.250E-03	1.250E-03	2.117E-08	0.000E+00	0.000E+00
U-234	1.099E-33	1.099E-33	1.861E-38	0.000E+00	0.000E+00
U-235	7.169E-35	7.169E-35	1.214E-39	0.000E+00	0.000E+00
U-238	5.257E-33	5.257E-33	8.901E-38	0.000E+00	0.000E+00

*The Surface Soil is the top layer of soil within the user specified mixing zone/depth.

Concentrations in the media occurring in pathways that are suppressed are calculated using the current input parameters, i.e. using parameters appearing in the input screen when the pathways are active.

Concentration of radionuclides in foodstuff media
 at t = 1.000E+04 years*

	Drinking Water	Nonleafy Vegetable	Leafy Vegetable	Fodder Meat	Fodder Milk	Meat	Milk	Fish	Crustacea
Radio- Nuclide	pCi/L	pCi/kg	pCi/kg	pCi/kg	pCi/kg	pCi/kg	pCi/L	pCi/kg	pCi/kg
Ac-227	0.000E+00	1.286E-35	1.283E-35	1.307E-35	1.307E-35	2.190E-37	1.150E-37	0.000E+00	0.000E+00
Pa-231	0.000E+00	6.948E-35	6.951E-35	6.961E-35	6.958E-35	5.786E-35	5.332E-38	0.000E+00	0.000E+00
Pb-210	0.000E+00	4.196E-04	4.183E-04	4.238E-04	4.238E-04	5.615E-05	1.934E-05	0.000E+00	0.000E+00
Ra-226	0.000E+00	1.876E-03	1.876E-03	1.876E-03	1.876E-03	1.737E-04	1.493E-04	0.000E+00	0.000E+00
Th-230	0.000E+00	6.363E-04	6.407E-04	6.412E-04	6.412E-04	6.687E-05	3.302E-06	0.000E+00	0.000E+00
U-234	0.000E+00	1.398E-33	1.401E-33	1.403E-33	1.403E-33	2.194E-34	3.761E-34	0.000E+00	0.000E+00
U-235	0.000E+00	9.114E-35	9.137E-35	9.152E-35	9.148E-35	1.431E-35	2.453E-35	0.000E+00	0.000E+00
U-238	0.000E+00	6.684E-33	6.700E-33	6.711E-33	6.709E-33	1.049E-33	1.799E-33	0.000E+00	0.000E+00

*Concentrations are at consumption time and include radioactive decay and ingrowth during storage time.
 For livestock fodder, consumption time is t minus meat or milk storage time.

Concentrations in the media occurring in pathways that are suppressed are calculated using the current input parameters, i.e. using parameters appearing in the input screen when the pathways are active.

Concentration of radionuclides in environmental media
 at t = 5.500E+04 years

Radio- Nuclide	Contaminat- ed Zone	Surface Soil*	Air Par- ticulate	Well Water	Surface Water
	pCi/g	pCi/g	pCi/m**3	pCi/L	pCi/L
Ac-227	0.000E+00	0.000E+00	0.000E+00	1.556E-11	1.556E-13
Pa-231	0.000E+00	0.000E+00	0.000E+00	1.047E-11	1.047E-13
Pb-210	4.117E-05	4.117E-05	6.969E-10	2.370E-02	2.370E-04
Ra-226	4.620E-05	4.620E-05	7.822E-10	3.383E-02	3.383E-04
Th-230	6.258E-04	6.258E-04	1.060E-08	4.003E-05	4.003E-07
U-234	0.000E+00	0.000E+00	0.000E+00	3.393E-10	3.393E-12
U-235	0.000E+00	0.000E+00	0.000E+00	1.523E-11	1.523E-13
U-238	0.000E+00	0.000E+00	0.000E+00	1.117E-09	1.117E-11

*The Surface Soil is the top layer of soil within the user specified mixing zone/depth.

Concentrations in the media occurring in pathways that are suppressed are calculated using the current input parameters, i.e. using parameters appearing in the input screen when the pathways are active.

Concentration of radionuclides in foodstuff media
 at t = 5.500E+04 years*

Radio- Nuclide	Drinking Water	Nonleafy Vegetable	Leafy Vegetable	Fodder Meat	Fodder Milk	Meat	Milk	Fish	Crustacea
	pCi/L	pCi/kg	pCi/kg	pCi/kg	pCi/kg	pCi/kg	pCi/L	pCi/kg	pCi/kg
Ac-227	1.556E-11	5.376E-12	2.577E-11	2.823E-11	2.822E-11	6.969E-14	8.083E-14	2.334E-12	1.555E-10
Pa-231	1.047E-11	3.628E-12	1.735E-11	1.903E-11	1.902E-11	9.088E-12	1.361E-14	1.047E-12	1.152E-11
Pb-210	2.370E-02	8.423E-03	3.948E-02	4.328E-02	4.328E-02	3.323E-03	1.858E-03	7.106E-02	2.373E-02
Ra-226	3.383E-02	1.277E-02	5.717E-02	6.237E-02	6.237E-02	5.956E-03	8.866E-03	1.691E-02	8.457E-02
Th-230	4.003E-05	3.323E-04	3.870E-04	3.936E-04	3.936E-04	3.417E-05	1.705E-06	4.003E-05	2.002E-04
U-234	3.393E-10	1.173E-10	5.618E-10	6.163E-10	6.161E-10	2.002E-11	5.290E-11	3.393E-11	2.036E-10
U-235	1.523E-11	5.263E-12	2.522E-11	2.766E-11	2.765E-11	8.985E-13	2.374E-12	1.523E-12	9.138E-12
U-238	1.117E-09	3.860E-10	1.849E-09	2.029E-09	2.028E-09	6.589E-11	1.741E-10	1.117E-10	6.701E-10

*Concentrations are at consumption time and include radioactive decay and ingrowth during storage time.
 For livestock fodder, consumption time is t minus meat or milk storage time.

Concentrations in the media occurring in pathways that are suppressed are calculated using the current input parameters, i.e. using parameters appearing in the input screen when the pathways are active.